Biological and physical evidence for extreme seasonality in central Permian Pangea

Cindy V. Looy^a, Stephanie L. Ranks^a, Dan S. Chaney^b, Sophie Sanchez^c, Sébastien Steyer^d, Roger M. H. Smith^{ef}, Christian A. Sidor^g, Timothy S. Myers^h, Oumarou Ideⁱ, Neil J. Tabor^h

^a*Department of Integrative Biology and Museum of Paleontology, University of California, Berkeley, Berkeley CA 94720, USA*

^b*Department of Paleobiology, NMNH Smithsonian Institution, Washington, D.C. 20560, USA*

^c*Department of Physiology and Developmental Biology, Evolutionary Biology Centre, Uppsala University, Norbyvägen 18A, 75236 Uppsala, Sweden*

^d*Centre de Recherches en Paléobiodiversité et Paléoenvironnements (UMR 7207 CNRS-MNHN-UPMC), Muséum national d'Histoire naturelle, CP 38, 8 rue Buffon, F-75008 Paris, France*

^e*Karoo Palaeontology, Iziko: South African Museum, Cape Town 8000, South Africa*

^f*Evolutionary Studies Institute, University of Witwatersrand, 1 Jan Smuts Avenue, Johannesburg 2000, South Africa*

^g*Burke Museum and Department of Biology, University of Washington, Seattle, Washington 98195, USA*

^h*Huffington Department of Earth Sciences, Southern Methodist University, Dallas TX 75275-0395*

ⁱ*Institut de Recherches en Sciences Humaines, Niamey, Niger Republic*

Corresponding author: Cindy V. Looy

Address: 1005 Valley Life Sciences Bldg #3140, Berkeley CA 94720-3140

Phone: ++1-510-642-1607

E-mail: looy@berkeley.edu

Abstract

Climate models indicate increased desertification in the continental interior of Pangea during the Permian, which would have affected the composition of the flora and fauna. We present a multi-proxy paleoenvironmental reconstruction of a terrestrial ecosystem in central Pangea of Lopingian age. The reconstruction is based on biological and physical data from the Moradi Formation, located in the Tim Mersoi Basin, northern Niger. Paleosols and sedimentological evidence indicate that the prevailing climate was semi-arid to very arid with marked intervals of high water availability. Carbon stable isotope data from organic matter and paleosols suggest that both the soil productivity and actual evapotranspiration were very low, corresponding to arid conditions. Histological analysis of pareiasaur bones shows evidence of active metabolism and reveals distinct growth marks. These interruptions of bone formation are indicative of growth rhythms, and are considered as markers for contrasting seasonality or episodic climate events. The macrofossil floras have low diversity and represent gymnosperm-dominated woodlands. Most notable are ovuliferous dwarf shoots of voltzian conifers, and a 25-m long tree trunk with irregularly positioned branch scars. The combined biological and physical evidence suggests that the Moradi Formation was deposited under a generally arid climate with recurring periods of water abundance, allowing for a well-established ground water-dependent ecosystem. With respect to its environment, this system is comparable with modern ecosystems such as the southern African Namib Desert and the Lake Eyre Basin in Australia, which are discussed as modern analogues.

Keywords

Niger, Permian, Central Pangea, climate, desert, paleobiology

1. Introduction

Lithological and phytogeographic data show that the region currently forming the interior of Africa experienced increased aridification from the Late Carboniferous-early Permian to the middle to late Permian (Gibbs et al., 2002; Rees et al., 2002). Climate models estimate that desert-like conditions prevailed in central Pangea during the Lopingian, with most recent models suggesting summertime high temperatures around 40-50° C (Fluteau et al., 2001; Peyser and Poulsen, 2008; Kiehl and Shields, 2005). The change from seasonally wet Cisuralian climate to desert-like Lopingian climate must have considerably altered both the composition of the flora and fauna in the local communities, and also their potential pathways for dispersal (Sidor et al., 2005). How exactly the biota responded is difficult to assess, however, as arid climates typically do not create the best conditions for plant preservation (Gastaldo and Demko, 2010), leaving an unsatisfactory gap in our knowledge of central Pangean ecosystems.

Fortunately, rare fossil vertebrate and plant assemblages have, in the past ten years, been discovered from the Upper Permian in northern Niger (Damiani et al., 2006; Sidor et al., 2003, 2005; Smiley et al., 2008; Steyer et al., 2006; Tsuji et al., 2013; Turner et al., 2015). During Lopingian times, deposition of the Moradi Formation of the Tim Mersoï sub-basin, Iullemeden Basin, occurred approximately 10° S of the paleoequator in central Pangea. Studies of vertebrate fossils from the Moradi Formation have suggested a community structure and taxonomic composition similar to those described from Morocco (e.g. Germain, 2010; Jalil and Dutuit, 1996; Jalil and Janvier, 2005; Sidor, 2013; Steyer and Jalil, 2009). They are, however, distinctly different from any of the better understood southern Pangean assemblages (e.g. South Africa, Tanzania, Zambia, and Malawi), which were dominated by therapsids (e.g. Angielczyk et al., 2013; Sidor et al., 2005, 2013). The faunas from Niger and Morocco both are dominated by temnospondyls, pareiasaurs, and captorhinids. The two herbivorous reptiles described from the Moradi Formation of Niger, the captorhinid *Moradisaurus* and the pareiasaur *Bunostegos*, are known only from

the formation (de Ricqlès and Taquet, 1982; Sidor et al., 2003; Taquet, 1969; Tsuji et al., 2013; Turner et al., 2015). Similarly, both of the temnospondyls described, *Nigerpeton* and *Saharastega*, represent relicts, or holdovers of lineages represented in faunas thought to have gone extinct near the Pennsylvanian-Permian boundary elsewhere (Damiani et al., 2006; Sidor et al., 2005, 2013; Steyer et al., 2006). The only terrestrial predator discovered in these faunas is represented by fragmentary gorgonopsid remains, not sufficient for a generic identification (Smiley et al., 2008). Due to the lack of any obvious physical barrier between the faunas of central and southern Pangea Sidor et al. (2005) hypothesized that climate was the factor likely responsible for the biogeographic isolation and endemism of the Moradi assemblage.

These climatic interpretations are corroborated by sedimentary, morphological, and mineralogical data from paleosol profiles in the Moradi Formation, which display evidence of prevailing arid conditions at the time of development. (Tabor et al., 2011; Smith et al., 2015). The paleosols and associated sedimentary strata are characteristic of those formed under a range of conditions from well to poorly drained, and represent stable floodplains of ephemeral, anabranching fluvial depositional systems (Tabor et al., 2011). These soil types are indicative of semi-arid areas with relatively shallow water tables, and include those formed on plains surrounding ephemeral lakes or playa environments. Modern analogues of these soil types can be found in the Namib Desert (Watson, 1992) of southwestern Africa and the Lake Eyre Basin, central and eastern Australia (Tabor et al., 2011; Fielding and Alexander, 2001; Tooth and Nanson, 2000). Both fossil and extant landscapes represent Heinrich Walter's zonobiome 3, or hot and arid subtropical deserts with very little rain (Breckle, 2002).

In this paper new biological and physical evidence for extreme seasonality in the Moradi Formation are presented, using a wide array of ecological proxies. The paleosol and sedimentological data are described briefly, and the stable carbon isotope compositions of paleosol calcite and co-existing organic matter are presented and discussed in terms of their relevance to interpretations of paleoclimate, biological productivity and water availability for floral and faunal communities across these paleolandscapes. Data

from tetrapod bone histology is presented for evidence of seasonality, including cyclical periodic cessation of growth. The Moradi gymnosperm-dominated fossil floras, including a large tree trunk, are described for the first time, and compared to modern analogs in order to constrain the environmental circumstances under which these plants grew. Finally, the implications of the combined data on the interpretation of climate for Lopingian central Pangea are discussed.

2. Material and methods

2.1. Geological setting

The Carboniferous-Permian stratigraphy and structural evolution of the Tim Mersoï sub-basin of the Iullemeden basin in north-central Niger was recently reviewed in Tabor et al. (2011). At the time of active Permian deposition, the basin was moving northward from ~30 to ~10° S (Scotese et al., 1999), and remained at least 2000 km from the nearest open ocean. The Moradi Formation is the terminal lithostratigraphic unit of the Izegouandane Group and, in the study area, comprises ~100 m of dominantly red mudstones with secondary fluvial channel sandstones and conglomerates as well as rare coarse breccias (Fig. 1). A Lopingian age of deposition for Izegouandane Group strata is primarily based upon biostratigraphic correlations of fossil vertebrates in the Moradi Formation to contemporaneous strata in Russia and South Africa (Sidor et al., 2005; Lucas, 2004; Taquet, 1969).

The specimens described in this paper were collected from several localities located approximately 20 km west of Arlit, Agadez Department, northern Niger (see o’Keefe et al., 2005 or Damiani et al., 2006 for a map of the study area). The localities are within 1 km of each other and all fall within the upper half of the Moradi Formation. Detailed locality data are available to qualified researchers at the Musée National du Niger in Niamey (MNN) or by contacting Christian A. Sidor. The fossil material described here is housed in the paleontological collections at the MNN.

2.2. Sedimentary environment and taphonomy of plant fossil localities

The upper Moradi succession is made up of three distinctive associations of sedimentary facies, which were interpreted as having accumulated in discrete sub-environments of the Moradi alluvial plain. These are (1) high sinuosity channel fill and point bar, (2) loessic floodplain and incised distributary channels, and (3) end-point playa. The stratigraphic position and sedimentological log of the fossil localities recovered by our team in the 2003 (S) and 2006 (T) expeditions is shown in Figure 1. The plant fossils from locality T9 occur within the loessic floodplain facies in a 0.25m-thick bed of pale red massively-bedded calcareous siltstone (“siltite”) with olive mottling along vertically orientated, downward branching root molds. This facies directly overlies a 3 m-thick conglomeratic sandstone displaying gently dipping ($\pm 20^\circ$) beds on its upper surface, which are interpreted as lateral accretion units of a large low-angle point bar. The fossiliferous “siltite” is immediately succeeded by a sequence of carbonate mudrock breccia and thinly-bedded micritic limestone interpreted as the accumulated deposits of a semi-permanent playa lake that most likely formed at the end-point of an ephemeral distributary channel (Smith et al., 2009, 2015). For a paleogeographic map and more information on the study area see Smith et al., 2015.

The leaf fossils occur as three-dimensional molds scattered at random orientations throughout the pale red siltstone, rather than concentrated along horizontal bedding planes. The lack of clearly-defined bedding planes within a bed of homogeneous unsorted silt-sized grains suggests that the leaf-bearing bed was deposited in a single episode and fairly rapidly as is common in windstorm-generated loess (Giles et al., 2013). The presence of root channels and the pervasive carbonate cement are indicative of incipient pedogenesis with relatively high water-tables in an evaporative setting, such as would be encountered on the sparsely-vegetated margins of an arid zone floodplain lake.

2.3. Fossil Flora

Fossil plants were observed and collected from a total of three localities (Fig. 1). Two (field localities T09 and T24) produced vegetative remains and one (field locality T25) produced fossil wood. Localities T09 and T24, in the upper part of the Moradi Formation, produced vegetative and fertile plant fossils in the

form of dark rust-colored carbonaceous dust in three-dimensional molds of a lighter colored fine-grained siltstone. The material was photographed using a Nikon D300 with a 60mm macro lens. Leaf fossils were found at both localities. Ovuliferous dwarf shoots of a voltzian conifer were found only at T24. Of the sixty-one dwarf shoots collected, twenty-eight were selected for morphological measurements based on well-defined outlines and completeness. Because terminology used to describe ovuliferous cones of early conifers deviates from that for modern species, terms used herein are explained (see Hernandez-Castillo et al. 2001; Rothwell et al. 2005). “Bracts” are foliar appendages of the cone (or fertile zone) axis, these are interpreted as a modified leaves. “Ovuliferous dwarf shoots” or “dwarf shoots” are lateral branches of limited growth, arising in axils of the bracts. A “sporophyll” is a potentially seed bearing appendage of an ovuliferous dwarf shoot, while a “sterile scale” is a foliar appendage.

Locality fossil forest deposit (field locality T25). One fossil tree trunk was partially excavated to a length of 25 m. The log was oriented roughly around 0° N-S with a dip of ~02° degrees. The angle of the fossil tree trunk arises from deposition at an orthogonal angle upon a point-bar deposit of a meandering stream. The length, diameter, branch base positions and shape of the exposed surface of the prone trunk were recorded. Small greenish-gray (silicified) wood samples were collected from the lower part of the trunk. These samples were thin-sectioned using standard techniques employed for petrified plant fossil material (e.g., Hass and Rowe, 1999). In the immediate area (< 0.1 km²) around the excavated log, there were other green haloes in the red sediment, as well as mounds of fragmented petrified wood. Petrified wood fragments also were found near vertebrate-producing sites S44 and S48, upon microscopic examination, no cellular structures were observed.

2.4. Paleosols and carbonate samples for geochemical analysis

The geochemical part of this study focuses on $\delta^{13}\text{C}$ values of paleosol calcite and co-existing organic matter samples from 23 different paleosol profiles distributed among three different stratigraphic sections

of the Moradi Formation. The paleosols from which these samples were collected exhibit three distinct morphologies (Tabor et al., 2011).

Paleosol type A: Type A paleosols occur in very fine to medium sandstones, and are especially common within the upper 1–2 m of thick sandstones that preserve trough cross-strata. They include vertical to horizontal calcareous tubules ranging from 1–100 mm in diameter, and ~10 mm to 3.7 m in length that are interpreted to be rhizoliths (*sensu* Klappa, 1980) or root petrifications (Tabor et al., 2011). Rhizoliths are typically composed of micritic calcite with alveolar texture on the exterior of the tubule and void-filling sparry calcite crystals on the interior of the tubule that grew inward. They rarely occur in such abundance that they dominate the volume of stratigraphic horizons, and are similar to calcified root mat horizons that have been interpreted to form just below the groundwater table (Wright, 1990). This paleosol type also rarely includes stratiform calcareous layers near the base of profiles that are interpreted to be groundwater calcrete (*sensu* Semeniuk, 1985, 1981; Fig 3F).

Paleosol type B: Type B paleosols occur in blocky red mudstones and include an abundance of discrete carbonate nodules, vertically stacked carbonate nodules (*sensu* Blodgett, 1988) and vertically and horizontally orientated carbonate tubules composed of micritic calcite. Calcareous nodules are interpreted to be stage II pedogenic nodules (*sensu* Machette, 1985), whereas calcareous tubules are interpreted as rhizoliths (Klappa, 1980). These paleosol profiles are typically noncalcareous in the upper ~15 cm of the profiles. Below that level they often include micritic calcareous cements along the exteriors of mudstone blocks; these are interpreted to be pedogenic calcans (Tabor et al., 2011).

Paleosol type C: Type C paleosol profiles occur in red mudstones that exhibit wedge-shape aggregate structure and low-angle slickenplanes. Profiles of this paleosol type are noncalcareous in the upper 15–30 cm of the profiles, but carbonate nodules and concretions ranging from a few mm to about 50 mm in

diameter and irregular carbonate tubules are common (5-20% by volume) to abundant (>20% by volume; Tabor et al., 2011).

In addition to the carbonate in the paleosol profiles discussed above, early-burial groundwater calcite and void-filling sparry calcite cements also are found in association with these paleosol profiles. Several examples of each are included in this work for comparison of stable carbon isotope compositions with micritic calcite nodules, tubules and calcans in paleosol profiles.

2.5. Stable carbon isotope geochemistry

Carbonate samples that were collected from at least 50 cm beneath the interpreted paleo-surface of paleosol profiles were drilled directly from thin-sections, matching billets, or polished slabs using a table-top drill system with 100 mm diamond bits on an x, y, z stage. Between 2 and 20 mg of carbonate-bearing powders were loaded into reaction vessels, and atmospheric gases were evacuated. Samples were then dissolved in 100% H₃PO₄ *in vacuo* at 25° C for about 16 hours to produce CO₂. CO₂ samples were cryogenically purified, and CO₂ yields were determined via mercury manometer with a precision of ±0.2 μmol. The resulting CO₂ was analyzed for carbon and oxygen isotope ratios using a Finigan MAT 252 isotope ratio mass spectrometer at Southern Methodist University.

Although fossil macroplant impressions and petrifications occur in the Moradi Formation (reported in this paper), macroscopic examples of fossilized organic plant matter were not observed. However, several examples of paleosol carbonate nodules and tubules occlude finely disseminated, organic-rich domains. These paleosol carbonate samples were pulverized, acidified in concentrated (~12 N) HCl in order to dissolve carbonate and eliminate it as a possible undesirable source of carbon during pyrolysis of organic matter. The organic-enriched residue was then rinsed with ionized water until the rinse water remained at the original pH (~5.6) after at least 10 minutes of contact with the acid-leached sample. The residues were then freeze-dried to remove absorbed water. The carbon content and δ¹³C values of the samples were

determined from CO₂ produced by closed system combustion of the organic matter following the methods of Boutton (1991).

In order to provide a context for the interpretation of Moradi Formation paleosol calcite and organic matter $\delta^{13}\text{C}$ values, paleosol carbonate and organic matter from three other Middle and Upper Permian basins are included in this work. These include the Middle and Upper Permian Quanzijie and Wutongguo formations of the Turpan Basin, Xinjiang Province, northwestern China (Yang et al., 2010). Samples from the Quanzijie Formation were collected from two different paleosol profiles in the lower half of the Formation (see Yang et al., 2010 for the specific lithostratigraphic section). Also included are calcite and co-existing organic matter stable carbon isotope compositions from Upper Permian (Changhsingian) Balfour Formation, South Africa (Gastaldo et al., 2013; Gastaldo et al., 2014), and Upper Permian Buntsandstein Formation, Italy.

Isotope values are reported in standard delta notation:

$$\delta^{13}\text{C} = (\text{R}_{\text{sample}}/\text{R}_{\text{standard}} - 1) * 1000,$$

where R is the ratio of heavy to light stable isotope present in the sample or standard, and δ values are reported relative to the Pee Dee Belemnite standard (PDB; Craig 1957) for both carbon and oxygen isotope ratios.

2.6. Bone histology

Four isolated pareiasaur bones, provisionally referred to *Bunostegos akokanensis* (humerus – MNN MOR145, radius – MNN MOR146, ulna – MNN MOR147, scapula – MNN MOR148), were sampled from the Moradi Formation at locality T22 (Tabor et al., 2011). The fossil material is preserved in three dimensions. The bones were prepared for osteohistology and embedded in resin (Lamm, 2013). The long

bones were sectioned at mid-shaft and a thin section was made just dorsally to the narrowest region of the scapula. They were observed and photographed using optical microscopes (Nikon Eclipse 80i and Leica CTR6000) with either ordinary transmitted light or polarized light.

The skeletochronological analysis (Castanet et al., 1993) that was performed consists of blind counting the number of growth marks that remain visible in the appositional deposit of compact primary tissue. A growth mark is composed of (1) a zone, corresponding to the amount of bone rapidly deposited during favorable periods of food supply; (2) an annulus, made of lamellar bone formed when environmental conditions become poorer; and in extreme cases, (3) a line of arrested growth (LAG) when the animal needs to reduce their metabolism and enter into lethargy or dormancy for survival (e.g., hibernation, aestivation; Castanet et al., 1993). LAGs are thus considered as biomarkers of contrasted seasonality. Growth marks are evidence of biological cycles interspersed throughout tetrapod bones. They have been intensively studied among extant species with regard to their age and development (Castanet, 1994; Castanet and Baez, 1991; Castanet et al., 2003). Growth mechanisms of periosteal bone are similar for all tetrapods, and thus can be assumed to be so for fossils (de Ricqlès, 1981, 1975; de Ricqlès et al., 2004). Therefore, growth marks are a proxy for the reconstruction of fossil tetrapod life history traits (e.g. Köhler et al., 2012; Sanchez et al., 2008, 2010b; Scheyer and Sander, 2009; Steyer et al., 2004) and the palaeoecological conditions influencing them (e.g. Konietzko-Meier and Klein, 2013; Sanchez and Schoch, 2013; Sanchez et al., 2010a; Steyer et al., 2004).

3. Results

3.1. Fossil flora

Systematics:

Leaves – The macrofloral elements from the two plant localities produced four morphological leaf types.

Description:

Morphotype 1: (T09, T24; 28 specimens) Isolated narrow strap-like leaves. These leaves were the only type found at the T24 locality. They are linear, straight to slightly curved, ranging between 100-125 mm in length and 2.2-3.5 mm in width (Fig. 2A). They taper slightly at their square base and have obtuse apices.

Morphotype 2: (T09; 2 specimens) Isolated linear leaves of variable size, approximately 85-133 mm long by 5-11 mm wide, with obtuse apices (Fig. 2B). The base is unknown. The leaves have prominent striations running parallel to the leaf margin (see inset). These striations could represent veins or stomatal rows.

Morphotype 3: (T09; 2 specimens) Shoot fragments with leaves. The leaves are short and narrow, linear, 5-22 mm long and 0.5-1.2 mm wide, with obtuse apices (Fig. 2C-D). They arise from the axis at angles between 26-75 degrees.

Morphotype 4: (T09; 4 specimens) Shoot fragments with obovate leaves. Leaves are 9-21 mm long and 2.2-3.6 mm wide, with obtuse tips and have slightly to greatly tapered bases that attach at angles of 12-50 degrees (Fig. 2 E-F).

Ovuliferous dwarf shoots – (T24; 61 specimens) The dwarf shoots have at least ten partially fused scales that are flattened and bilaterally symmetrical, longer than they are wide (Fig. 3). The total length of the dwarf shoots varies from 8-21 mm and width from 9-17 mm. Each dwarf shoot bears two types of foliar elements, interpreted as sterile scales and sporophylls. The sterile scales are narrow-triangular in shape with acuminate apices and are positioned in at least two planes. They have fused bases but separate in their upper 2-7 mm, with widths of 0.2-0.7 mm immediately distal to the point of separation of the scales. The sporophylls seem to be interspersed among the sterile ones. The more centrally positioned sporophylls are oblong to triangular with a rounded apex, shorter and fleshier than the sterile scales, and

about 3.9-6.6 mm long and about 0.7-1.8 mm wide. The position of the ovule attachment area could not be discerned. The scale-bearing, fused part of the ovuliferous dwarf shoot is much wider than high *en face* with a broad, rounded base (Fig. 3C-E). The basal part is long and stalk-like, with a length ranging from 8.5-12 mm and diameter/width between 0.5-3.3 mm. It is not clear if the stalk-like base was attached to the bract. It was possible to determine the original shape of the ovuliferous dwarf shoots by the impressions left on the matrix. In one specimen, in which both part and counterpart are preserved, the fused scale-bearing part of the dwarf shoot displays a thickness of 4 mm (Fig. 3F).

Log – (T25) The exposed straight basipetal part of the Moradi tree measured 58 cm in diameter (Fig. 4). During excavation, more than 24.9 m of the trunk was exposed. As the base of the trunk is not preserved, all length measurements are from the basipetal end of the log as preserved. The basipetal end is not flared. Flaring of the trunk would suggest that the tree base was within < 0.5 m. Thus, the true base of tree, were it to have been preserved, was more than a meter below the preserved base. The distal-most exposed end of the trunk had a width of 20 cm, and continued into unexcavated strata. Based on the size of the branches that had broken off the trunk, the distal end of the preserved tree is probably not more than 1 m beyond the end of the excavation. Therefore, whether the true based was preserved or not, the length of the trunk in life was greater than the fossilized specimen, as exposed in excavation. Scars of irregularly spaced branches are present along most of the length, but are less common and not as prominent in the basal 9 m. Above 9 m the branch scars gradually increase in size. Between 9 and 11 m the elliptical branch scars range from 4 x 3 cm to 10 x 8 cm (l x w). Above 11 m, larger branch scars occur, ranging from 10 x 19 cm to 22 x 15 cm, with smaller branch scars increasingly interspersed above 19 m. At the very top of the exposed portion there is a branch scar as large as the width of the tapered trunk (10 x 10 cm), indicating that it had bifurcated at that point.

Thin sections revealed poor anatomical preservation (Poole 2013, pers. comm.), making identification of the wood problematic. No evidence for growth rings or growth intervals were observed, this however is

not to say definitively that this specimen did not have these features *in vivo*. Isolated wood fragments from other vertebrate-rich localities were identified as *Agathoxylon* (Bamford personal communication), a wood morphology characteristic of coniferophytes, cordaitaleans, and conifers (Philippe and Bamford, 2008). The cross section of the basal part of the trunk is massive and shows no evidence of a large central pith. The absence of a central pith excludes the cordaitaleans and suggests that the fossil trunk is likely a conifer. The observed irregular branching pattern is characteristic of orthotropically branching voltzian conifers (Rothwell et al., 2005).

3.2. Stable carbon isotope geochemistry

Paleosol Calcite - Paleosol carbonate samples from the Moradi Formation (n = 69) range from 19 - 100% calcite (Table 1). Among the samples analyzed, calcans and stage II carbonate nodules have lower, whereas stage III carbonate layers and rhizoliths have higher wt. % calcite (Table 2). Micrite $\delta^{13}\text{C}$ values (n = 66) range from -8.2 to 3.3‰ (mean = -3.0 ± 1.6 ‰). Spar and microspar cement $\delta^{13}\text{C}$ values (n = 3) range from -4.6 to -3.4‰ (mean = -4.2 ± 0.7 ‰).

In general, micrite from BkIII horizons and rhizoliths record the most positive $\delta^{13}\text{C}$ values, whereas BKII nodules record the most negative $\delta^{13}\text{C}$ and values. Calcans and early-burial groundwater cements record intermediate $\delta^{13}\text{C}$ values. This pattern of isotope distribution among the carbonate textures is also mimicked by the pattern of isotope distribution among paleosol carbonates with respect to their host lithology. That is, calcite from sandstone profiles generally records more positive $\delta^{13}\text{C}$ values, calcite from mudstone profiles records more negative $\delta^{13}\text{C}$ values, and calcite from siltstones has intermediate $\delta^{13}\text{C}$ values.

As mentioned, other Middle and Upper Permian paleosol carbonate samples were analyzed in order to place the Moradi Formation calcite isotope values in a broader context. Quanzijie Formation micrite samples (Middle Permian, NW China; n = 4) record $\delta^{13}\text{C}$ values between -5.2 and -3.9‰ (mean = -

4.6‰). Wutonggou Formation micrite samples (Upper Permian, NW China; n = 2) record $\delta^{13}\text{C}$ values between -13.5 and -10.4. Balfour Formation micrite samples (Upper Permian, South Africa; n = 10) record $\delta^{13}\text{C}$ values between -17.1 and -5.3‰ (mean = -11.6‰). Buntsandstein micrites (Upper Permian, northern Italy; n = 9) record $\delta^{13}\text{C}$ values between -3.2 and -5.9‰ (mean = -4.6‰). Thus, micrite $\delta^{13}\text{C}$ values from the Moradi Formation paleosol carbonates are more positive than Middle and Upper Permian samples from other sites.

Organic Matter - Residues (n = 9) from acid-reaction of carbonate nodules, rhizoliths, and calcans in the Moradi Fm. range from 0.053 – 1.187 wt. % carbon (Table 2). $\delta^{13}\text{C}$ values of CO_2 produced via pyrolysis of organic matter in these residues range from -20.8 to -24.5‰. The difference between corresponding calcite and organic matter carbon isotope values, or $\Delta^{13}\text{C}_{\text{CC-OM}}$, ranges from 18.2 to 24.4‰.

Middle Permian Quanzijie Formation residues (n = 2) have organic matter $\delta^{13}\text{C}$ values that range from -19.8 to -20.3‰, whereas $\Delta^{13}\text{C}_{\text{CC-OM}}$ values range from 15.2 to 15.6‰. Upper Permian Balfour Formation residues (n = 5) range from -25.3 to -26.6‰, whereas $\Delta^{13}\text{C}_{\text{CC-OM}}$ values range from 3.3 to 20.4‰. Upper Permian Buntsandstein and Wutonggou formations carbonate samples were not available for production of organic matter-rich residues. However, coalified branch axes and fossil leaves in close stratigraphic association with paleosol calcite samples are available for determination of organic matter $\delta^{13}\text{C}$ values. Fossil wood samples from the Buntsandstein Formation range from -21.6 to -22.6‰, resulting in $\Delta^{13}\text{C}_{\text{CC-OM}}$ values from 16.3 to 18.7‰. Organic matter $\delta^{13}\text{C}$ values from the Wutonggou Formation range from -21.5 to -24.7‰, resulting in $\Delta^{13}\text{C}_{\text{CC-OM}}$ values from 11.1 to 11.2‰. Therefore, although $\delta^{13}\text{C}$ values of organic matter from the Moradi Formation are similar to those from contemporaneous sites, Moradi Formation $\Delta^{13}\text{C}_{\text{CC-OM}}$ values are larger than contemporaneous sites because of their generally more positive calcite $\delta^{13}\text{C}$ values.

3.3. Bone histology

Scapula (MNN MOR148) - Although only half of the scapula is available for sectioning, the bone tissue is very well preserved (Fig. 5A-B). As the result of membrane ossification (Francillon-Vieillot et al., 1990), the scapula is composed of three layers (Fig. 5 A): 1) a slightly compact basal layer of bone; 2) a cancellous core; and 3) an outer compact layer built out by appositional cyclical deposition. In the third layer, five clear LAGs are visible that are regularly spaced out from each other (Fig. 5B). The vascularization is very dense and represented by a longitudinal mesh of primary osteons. Although the collagen matrix is mainly fibrous, some patches of pseudo-lamellar bone are also scarcely distributed over the cortex. The cancellous core exhibits a large amount of radial trabeculae, covered with a secondary lamellar deposit. The basal layer is pierced by numerous, small and simple canals and large primary osteons.

Radius (MNN MOR146) – The radius is a long bone composed of a cortex surrounding a medullary cavity that is secondarily filled with trabeculae (Fig. 5D). These trabeculae form part of the spongiosa. As a result of remodeling (resorption and redeposition), the spongiosa extends to the inner region of the cortex (Fig. 5D). It comprises numerous secondary lamellar trabeculae that are oriented in all directions. The compact cortical bone represents one third of the section area. It is made of fibrous and pseudo-lamellar bone (Fig. 5E, H). In the crest region, some radial extrinsic fibers cut through the bone matrix (Fig. 5E). The bundles of fibers are associated with a dense radial vascular mesh of primary osteons. In the other part of the cortex, the vascularization is composed of longitudinal primary osteons, organized in concentric rows (Fig. 5H). The *pareiasaur* radius shows a pattern of three LAGs visible in a few places of the compact cortex (Fig. 5G). The distance between these LAGs is constant over the entire thickness of the compacta.

Ulna (MNN MOR147) - The mid-shaft histology of *Bunostegos* ulna exhibits a very densely vascularized cortex. The vascular mesh is made of longitudinal and circular vascular canals and osteons. Based on this great cortical porosity, the spongiosa occupies almost the entire section area. The central part of the

medullary cavity remains free of trabeculae. Because of recrystallization, the bone tissue is too badly preserved to identify any LAG or analyze the bone matrix organization.

Humerus (MNN MOR145) - The surface of the bone has been very much weathered, as a result the compact layer of cortical bone has completely disappeared. The preserved spongiosa has been greatly remodelled and post-mortem recrystallized, and does not preserve to the primary tissue or LAGs. The spongiosa is composed of lamellar and pseudo-lamellar trabeculae, which are relatively loose and evenly distributed in the central region of the section, but denser and slightly organized in concentric rows in the rest of the section.

4. Discussion

This study combines lithological indicators with biological and geochemical proxies of paleoclimate in order to establish a robust reconstruction of terrestrial paleoenvironments. Due to the broad nature of the datasets presented here, they provide a unique insight into climate during Moradi Formation deposition and the extent of seasonality over this region of Pangea.

4.1. Fossil flora

Log - In the area around the excavated log, there were green haloes in the red sediment, as well as mounds of fragmented petrified wood. Though not excavated, each of these occurrences likely represented the presence of a section of a petrified fossil log. Based on the conglomeratic nature of the sediment in which the excavated log was entombed, these logs, taken in toto, may represent a log jam. This means that the 25 m long, possible conifer could have been transported downstream from a wetter upland region. Wood samples from the fossil tree did not show evidence of a seasonal or episodic climate. This could be a preservation artifact, but given its size it is also likely that the root system was able to tap into ground water as its permanent water source.

Taxonomic affiliation - The ovuliferous dwarf shoots of the Moradi conifers have a morphology that is comparable to members of the voltzian Voltziales (or voltzians), a monophyletic group of Paleozoic conifers that includes members of the Majonicaceae (Rothwell et al., 2005). In a phylogenetic analysis by Rothwell et al. (2005) the Permian voltzians were the sister group to the lebachiod walchians, which comprise the other major clade of late-Paleozoic conifer taxa. The ovuliferous cones or fertile zones in these two groups are compound, with bract and dwarf shoots helically arranged along the main axis. The bilaterally symmetrical walchian dwarf shoots consist of more than ten sterile scales with one to three interspersed sporophylls. In contrast to ovuliferous dwarf shoots of walchians, in Late Paleozoic voltzians the basal part of the dwarf-shoot is stalk-like, the vegetative scales and sporophylls are fused at the base and form a flattened structure of distinctly bilateral organization. The vegetative scales and sporophylls are either arranged side by side along the width of the fused structure, or are arranged in two to three planes. In the latter case the sporophylls are always positioned on the adaxial side of the dwarf shoot, with some of the sterile scales forming the abaxial row(s).

The gross morphology of the Moradi dwarf shoots is similar those of members of the Majonicaceae. They have a long stalk, fused bases of vegetative scales and sporophylls, and several vegetative scales positioned on the abaxial side. The only thing reminiscent of walchian dwarf-shoot morphology, is the higher number of scales on the Moradi shoots compared to most voltzian conifer (five or less). This, however, is not a synapomorphy shared by all voltzians, as also the undisputed voltzian *Dolomitia cittertia* had many sterile scales on the abaxial side of its ovuliferous dwarf-shoots. Therefore, despite their poor conservation, they can be confidently assigned to the voltzian Voltziales. Most voltzian Voltziales have an irregular orthotropic branching pattern and medium-sized, bifacial ovate to linear leaves (Clement-Westerhof, 1987; Schweitzer, 1963, 1996; Rothwell et al., 2005). So it is quite possible that some of the Moradi leaves were produced by the same plan that produced the ovuliferous dwarf shoots.

Floral biogeographic implications - Before the discovery of the Moradi floras, Permian voltzian conifers were mainly known from Euramerican basins. The oldest known voltzians are found in latest early Permian to early middle Permian deposits from the Midland Basin, north-central Texas, United States (e.g., DiMichele et al., 2001, 2004; Looy 2007, Looy and Stevenson, 2014). Recovered from a channel-form silty deposit, there are in the sub and superjacent strata weakly developed Inceptisols (Tabor and Montañez, 2004; DiMichele et al., 2006; Montañez et al., 2007) and primary gypsum (DiMichele et al., 2001, 2008), suggesting growth on short-lived flood plains in a generally strongly seasonally dry climate. Until now, Lopingian voltzian conifers were exclusively known from seasonally wet plant communities from Europe, particularly from the Zechstein Group of Western Europe and the Val Gardena Sandstone of the Southern Alps (Clement-Westerhof, 1987; Schweitzer, 1986, 1996). The Moradi dwarf shoots show that voltzian conifers were not restricted to arid environments in the equatorial tropics and northern paratropical parts, but instead had a range extending to similar latitudes in the southern hemisphere.

4.2. Stable carbon isotope geochemistry

Soil CO₂ concentrations (C_s; see appendix) estimates from the Moradi Formation (Niger) are the lowest of all Middle and Upper Permian sites; 500 to 1300 ppmV (mean = 800 ±300; Table 3, Appendix). C_s estimates from the Buntsandstein Formation (Italy) range from 1200 to 2800 ppmV (mean = 1800 ±700). The Upper Permian Balfour Fm. samples that conform to 2-component soil CO₂ mixing exhibit a large range of C_s estimates from 800 to 6000 ppmV. The Middle Permian Quanzijie Formation samples provide generally higher and uniform C_s estimates, 6300 ±2600 ppmV.

Differences in soil PCO₂ across a landscape are principally related to differences in soil productivity, which is strongly controlled by the amount of moisture in the soil that is available for floral and microbial metabolic processes (Myers et al., 2012; Tabor et al., 2013; Brook et al., 1983). In this regard, C_s estimates presented herein suggest that the Lopingian soils of the Moradi Formation were among the least productive studied to date. Such low productivity may have been related to more arid conditions, and

increased moisture stress of the Permian soil floras in Niger, compared with other Lopingian paleogeographic sites.

The following equation, modified from Brook and others (1983), relates the concentration of mean growing-season soil CO₂ to actual evapotranspiration:

$$\text{Log } \frac{C_S}{10^6} = \text{Log } \frac{C_A}{10^6} + 2.09 * (1 - e^{-0.00172 * AET})$$

C_S and C_A are as defined in Eqn 1. AET is actual evapotranspiration in units of mm/yr., which is the quantity of water that is removed from a surface due to the processes of evaporation and transpiration. If all other factors are held constant, AET will be less in arid climates than in humid climates, because there is less precipitation delivered to the soil to be used in evapotranspiration. Actual evapotranspiration in naturally occurring well-drained soils is almost always less than mean annual precipitation, because some moisture can be lost to through-flow (deep percolation or groundwater) and soil runoff. Therefore, AET estimates effectively provide *minimum* estimates of mean annual precipitation.

Moradi Formation AET estimates are the lowest among all study sites; 110 ±50 mm/yr (Table 2). This is followed, in order of increasing AET estimates, by Buntsandstein (240 ±70 mm/yr), Balfour (320 ±290 mm/yr) and Quanzijie (540 ±110 mm/yr) formations. Thus AET estimates based upon δ¹³C values of co-existing calcite and organic matter indicate that the paleogeographic location of the Moradi Fm. was the most arid of Permian sites studied.

In spite of the evidence for aridity garnered from this stable carbon isotope geochemical analysis, the fauna associated with the Moradi formation strata includes large amphibians. This would seem to contradict the geochemistry results. Furthermore, lithological evidence such as shrink-swell features in

paleosol profiles, lateral accretion surfaces in fluvial channels, and non-marine limestones in end-point playa deposits indicate that water was abundant at least intermittently.

Several explanations can account for these different paleo-environmental signals. One possibility is that the paleosols, from which no plant fossils have been recovered, are time equivalent but lateral to the fluvial sediments in the section. This would mean the regional climate was arid with local rivers, and included poorly-drained areas with shallow water tables (cf. DiMichele et al., 2006). Another explanation is that the fluvial sediments represent the wetter times during fluctuating successions of wetter and drier phases. The Moradi sediments were deposited in the continental interior of Pangea, and the Aïr mountains located to the east of the Tim Mersoï Basin would have received additional rainfall due to elevation, but maybe not the massive amounts required to move all the conglomeratic material and other sediments into the lowland. The Moradi flora and fauna could be representative of wetter intervals, and the paleosols that developed during the wetter phases could have either been reworked by the streams or strongly overprinted by the dry phase paleosols.

4.3. Bone histology

The studied pareiasaur bones showed a fibrous matrix and a great density of their vascular network (Fig. 3B, F, and H). This indicates that these individuals had relatively fast bone deposition and active metabolisms. This is in accordance with previous studies, which suggest South-African pareiasaurs experienced rapid growth and short development periods (de Ricqlès, 1978a, b; Canoville and Chinsamy-Turan, 2011).

Although these fossils show evidence of active metabolism, they also exhibit growth marks which are reliable indicators of growth rhythms (Castanet et al., 1993). These growth marks end up as lines of arrested growth (LAGs). In extant tetrapods, LAGs are generally laid down annually (e.g. Castanet et al., 1996; Castanet, 1994). The five LAGs counted on the scapula therefore suggest this individual was at

least five years old. The radius must have belonged to an individual older than three years. Since the distance between the LAGs remains constantly large over the period of bone deposition (0.92-1.17 mm between two LAGs at mid-shaft in the radius, Fig. 5G; 1.07-1.71 mm close to the narrowest region of the scapula, Fig. 5B), it would seem that the corresponding individuals died while still in development (the LAG interdistance usually decreases with acquisition of sexual maturity; Castanet et al., 2003). This hypothesis regarding the ontogeny of these individuals should be nevertheless tested within the framework of a complete growth-series analysis.

The presence of LAGs in specimens tentatively referred to *Bunostegos* probably results from seasonal or episodic changes in the local climate. The LAG patterns are observed in various individuals, thereby suggesting that these changes must have been contrasted enough to induce successive periods of fast growth and lethargy within this taxon. The regularity of the LAG patterns would evoke a relatively constant succession of seasons. As pareiasaurs have sometimes been considered amphibious (e.g. Krilloff et al., 2008), it is possible that an annual fluctuation in water supply could be one of the major reasons for the cyclical growth patterns inferred here from observations of bone histology.

In each bone where LAG patterns were visible (Fig. 5B, G), bone deposition stopped in the early part of a zone (i.e., when active growth restarted) after the period of dormancy. The time of death is therefore suspected to have occurred during a period when environmental conditions were favorable, i.e., when water was once again available. These observations agree with the taphonomic interpretation of flood events resulting in the accumulation of multiple pareiasaur skeletons.

4.4. The Moradi environment and modern analogues – In places the loessic floodplain silts of the upper Moradi Formation contain discrete shallow roughly circular depressions measuring some 30-50m in diameter filled with a 4-5m thick succession of thinly-bedded lacustrine limestone (Facies C, Fig. 1). At the base of each sequence is a relatively thick and distinctive chaotic limestone “biscuit” breccia filling

deep elongate troughs. The biscuits are platy clasts of reworked calcareous mudstone that appear to have been ripped up from the desiccated mudflats around the lake margins and rapidly dumped into channels cut into the lake bed. Overlying the basal breccia is a succession of thinly bedded and finely laminated carbonate mudstones that display 5-7 stacked depositional cycles. Each 0.5 m cycle comprises climbing ripple cross-lamination with the topmost ripple forms preserved, grading into horizontal laminated lime rich mudstone and terminating in a claystone veneered paleosurface commonly showing desiccation cracks filled with pale red aeolian silt stone and the preferential preservation of trackways. Such sequences are common in playa lake settings and reflect episodic sediment laden floodwaters entering a ponded water body. Interestingly, the flood sequences are overlain by finely laminated mudstone terminating in massive micrite with wrinkle marks and pustular algal-matted patches with desiccation cracks. The superposition of these deep water lacustrine facies on their marginal facies indicates that upper Moradi playa lakes enlarged and deepened over time. A very similar association of edgewise limestone biscuit breccias overlain by turbidite sequences recording decelerating flow, as well as in-situ reed stems, has been recorded in Pleistocene end-point playa deposits of the Namib desert (Smith and Mason, 1998). The classic modern example of this environment is in the hyperarid central Namib Desert where the Sossus river terminates in a dune-dammed playa lake at Sossusvlei.

Paleosol and biotic data from the Moradi Formation give seemingly contradictory results about paleoclimate. Although the paleosol types found in the Moradi (Protosols (similar to Aridisols, Inceptisols, and Entisols) Calcisols, and Vertisols indicate that the prevailing climate was semi- to very arid with pronounced seasonality, there is also sedimentological evidence that these same areas experienced intervals of great water abundance (Tabor et al., 2011; Smith et al., 2015; this paper). Possible present-day analogues for conditions interpreted from the Niger Permian paleosol data, are found in the deserts that are among the driest in the world (Goudie, 2010; Nanson et al., 1992): the Lake Eyre Basin, Central Australia and the Namib Desert, southwestern Africa. Heinrich Walter (Walter, 1984) classified the present global climate zones in zonobiomes (regions defined by particular climates and

ecologies) and orobiomes (biomes located in mountain regions). Both the Namib Desert and Lake Eyre Basin are classified as zonobiome 3: hot, arid subtropical deserts (Breckle, 2002).

The Lake Eyre Basin of central Australia is a barren wasteland of salt plains and sand dunes most years. About twice in every hundred years, the landscape is transformed as heavy rains in the northeast fill the tributary rivers and flood the basin, creating an inland sea (McMahon et al., 2008; Serventy, 1985). While this flooded state occurs infrequently, there are more frequent periods of short small-scale flooding, which fill dry creek beds and temporarily bring life to the floodplains surrounding the lake (Bonython and Mason, 1953). Despite the extremely variable environmental conditions, life thrives in the Lake Eyre Basin. The plants there have morphological and physiological features that permit them to survive prolonged periods of drought, such as persistent seed banks and root systems that penetrate both wide and deep into the soil (Bowman, 2002). Shady groves of large eucalyptus trees provide shelter for large mammals and reptiles, and amphibians (Cockerham, 2004; Serventy, 1985). Such communities can weather the inevitable periods of drought by tapping into deeper water tables, despite being highly saline and saturated in carbonates (Gregory, 1906).

The central and inland regions of the Namib Desert lack the fog belts that somewhat mitigate the dry conditions in the coastal regions. Away from the shore, the greatest amount of vegetation exists in interdune areas, with the greatest diversity in the east (Seely and Ward, 1989). The Kuiseb River course winds through the dunes and escarpments, creating linear ground-water dependent oases that support pockets of desert life. Acacia trees form woodland regions that provide habitats for populations of a wide variety of animals (Seely and Ward, 1989). As evidenced by both the Lake Eyre Basin and the Namib Desert, despite being surrounded by a generally arid climate, even small areas with increased water availability can sustain a wide range of flora and fauna, including large vertebrates and trees. Thus, finding fossil amphibians in the Moradi Formation, i.e. non-amniotic vertebrates that need water at

least for their reproduction, is not so strange when the dynamics revealed by modern analogues are taken into consideration.

5. Conclusions

Description Moradi ecosystem - In sum, we have organisms that lack an amniotic egg and therefore depend on water for reproduction (temnospondyl amphibians), paleosols indicating semi-arid conditions (Tabor et al., 2011), carbon stable isotope data from paleosols indicating arid climatic conditions, and evidence of tall, robust plant life. All of these facts combined suggest that the Permian Moradi Formation was deposited under a generally arid climate that experienced cyclical intermittent periods of water abundance, a place of extreme seasonality or episodicity, supporting what was probably a well-established ground water dependent ecosystem. During the drier, probably longer intervals, we suggest that the Moradi ecosystem was a large arid and relatively flat plain with a sparse plant cover adapted to shallow ground water resources and temporary water ponds in which rare temnospondyl amphibians survived. During the shorter but intense wet intervals, this ecosystem was cut and drained by temporary meandering systems occasionally filled by locally catastrophic flash floods (Smith et al., 2009, 2015).

Acknowledgements

We are indebted to the Nigerien Ministry of Higher Education, Research, and Technology, the National Museum of Niger, and the University of Niamey for supporting our research. We thank T. Smiley, S. Thomas, A. Dindine, A. Maga and B. Tahirou for assistance in the field. N.J. Tabor thanks Francesco Massari (Padova University) for introducing and leading him through the late Permian strata of Italy. S. Streyer and S. Sanchez acknowledge M. Lemoine (MNHN, Paris) for making the bone thin sections. C. Looy thanks I. Poole for making the wood thin sections. W.A. DiMichele and S. Finnegan are thanked for proof reading the draft manuscript. Two anonymous reviewers provided thoughtful and critical reviews, which greatly improved the paper. This project was supported by the National Science Foundation grants EAR 0617250 to N.J. Tabor and EAR-0617718 to C.A. Sidor. S. Sanchez was funded by ERC Advanced

Investigator Grant 233111 (to P. E. Ahlberg), and S.L. Ranks work was funded by the Hellman Foundation (to C.V. Looy). This is University of California Museum of Paleontology Contribution No. xxx.

References

- Angielczyk, K.D., Steyer, J.S., Sidor, C.A., Smith, R.M.H., Whatley, R.L., Tolan, S., 2013. Permian and Triassic dicynodont (Therapsida: Anomodontia) faunas of the Luangwa Basin, Zambia: taxonomic update and implications for dicynodont biogeography and biostratigraphy, in: Kammerer, C.F., Angielczyk, K.D., Fröbisch, J. (Eds.), The early evolutionary history of Synapsida. Springer, Dordrecht, pp. 93–138
- Blodgett, R.H., 1988. Calcareous paleosols in the Triassic Dolores Formation, southwestern Colorado. Geol. S. Am. S. 216, 103–121.
- Bonython, C.W., Mason, B., 1953. The filling and drying of Lake Eyre. Geogr. J. 119, 321–330.
- Boutton, T.W., 1991. Stable carbon isotope ratios of natural materials: II. Atmospheric, terrestrial, marine and freshwater environments, in: Coleman, D.C., Fry, B. (Eds.), Carbon Isotope Techniques. Academic Press, San Diego, pp. 155–170.
- Bowman, D.M.J.S., 2002. The Australian summer monsoon: a biogeographic perspective. Aust. Geogr. Stud. 40, 261–277.
- Breckle, S.-W., 2002. Walter's vegetation of the earth: the ecological systems of the geo-biosphere, Springer, Berlin.

665 Brook, G.A., Folkoff, M.E., Box, E.O., 1983. A world model of soil carbon dioxide. *Earth Surf. Proc.*
666 *Land.* 8, 79–88.

667

668 Canoville, A., Chinsamy-Turan, A., 2011. Growth patterns and palaeoecology of pareiasaurs (parareptilia,
669 pareiasauridae) inferred from long bone histology and microanatomy. *J. Vertebr. Paleontol.* 31 (Suppl. 3),
670 83.

671

672 Castanet, J., 1994. Age estimation and longevity in Reptiles. *Gerontology* 40, 174–192.

673

674 Castanet, J., Baez, M., 1991. Adaptation and evolution in *Gallotia* lizards from the Canary Islands: age,
675 growth, maturity and longevity. *Amphibia-Reptilia* 12, 81–102.

676

677 Castanet, J., Francillon-Vieillot, H., Bruce, R.C., 1996. Age estimation in desmognathine salamanders
678 assessed by skeletochronology. *Herpetologica* 52, 160–171.

679

680 Castanet, J., Francillon-Vieillot, H., Meunier, F.-J., de Ricqlès, A., 1993. Bone and individual aging, in:
681 Hall, B.K. (Ed.), *Bone Vol. 7: Bone Growth-B.* CRC Press, Boca Raton, pp. 245–283.

682

683 Castanet, J., Francillon-Vieillot, H., de Ricqlès, A., 2003. The skeletal histology of the Amphibia, in:
684 Heatwole, H., Davies, M. (Eds.), *Amphibian Biology, Vol. 5: Osteology.* Surrey Beatty & Sons, Chipping
685 Norton, pp. 1598–1683.

686

687 Clement-Westerhof, J.A., 1987. Aspects of Permian palaeobotany and palynology. VII. The
688 Majonicaceae, a new family of Late Permian conifers. *Rev. Palaeobot. Palynol.* 52, 375–402.

689

690 Cockerham, S.T., 2004. Irrigation and planting density affect river red gum growth. *Calif. Agr.* 58, 40–43.

691

692 Craig, H., 1957, Isotopic standards for carbon and oxygen and correction factors for mass-spectrometric
693 analysis of carbon dioxide: *Geochim. Cosmochim. Acta* 12, 133–149.

694

695 Damiani, R.O., Sidor, C.A., Steyer, J.-S., Smith, R.M.H., O’Keefe, F.R., Larsson, H.C.E., Maga, A., Ide,
696 O., 2006. The vertebrate fauna of the Upper Permian of Niger. V. The primitive temnospondyl
697 *Saharastega moradiensis*. *J. Vertebr. Paleontol.* 26, 559–572.

698

699 de Ricqlès, A., 1981. Recherches paléohistologiques sur les os longs des tétrapodes. VI—Les
700 Stégocéphales. *Ann. Paléontol.* 67, 141–157.

701

702 de Ricqlès, A., 1978a. Recherches paléohistologiques sur les os longs des tétrapodes. VII—Sur la
703 classification, la signification fonctionnelle et l’histoire des tissus osseux des tétrapodes (Troisième partie:
704 Evolution: considérations phylogénétiques). *Ann. Paléontol.* 64, 85–111.

705

706 de Ricqlès, A., 1978b. Recherches paléohistologiques sur les os longs des tétrapodes. VII—Sur la
707 classification, la signification fonctionnelle et l’histoire des tissus osseux des tétrapodes (Troisième partie:
708 Les problèmes du déterminisme des types de tissus osseux). *Ann. Paléontol.* 64, 153–184.

709

710 de Ricqlès, A., 1975. Recherches paléohistologiques sur les os longs des tétrapodes. VII—Sur la
711 classification, la signification fonctionnelle et l’histoire des tissus osseux des tétrapodes (Première partie:
712 Structures). *Ann. Paléontol.* 61, 49–129.

713

714 de Ricqlès, A., Castanet, J., Francillon-Vieillot, H., 2004. The 'message' of bone tissue in
715 paleoherpetology. *Ital. J. Zool.* 71 S2, 3–12.

716

717 de Ricqlès, A., Taquet, P., 1982. La faune de vertébrés du Permien Supérieur du Niger.—I. Le
 718 captorhinomorphe *Moradisaurus grandis* (Reptilia, Cotylosauria). Ann. Paléontol. 68, 33–106.
 719
 720 DiMichele, W.A., Hook, R.W., Nelson, W.J., Chaney, D.S., 2004 An unusual Middle Permian flora from
 721 the Blaine Formation (Pease River Group, Leonardian-Guadalupian Series) of King County, west Texas.
 722 J. Paleontol. 78, 765–782.
 723
 724 DiMichele, W.A., Mamay, S.H., Chaney, D.S., Hook, R.W., 2001. An Early Permian flora with Late
 725 Permian and Mesozoic affinities from north-central Texas. J. Paleontol. 75, 449–460.
 726
 727 DiMichele, W.A., Tabor, N.J., Chaney, D.S., Nelson, W.J., 2006. From wetlands to wet spots:
 728 environmental tracking and the fate of Carboniferous elements in Early Permian tropical floras, in: Greb,
 729 S.F., DiMichele, W.A. (Eds.), Wetlands through time. Geol. S. Am. S. 399, 223–248.
 730
 731 DiMichele, W.A., Kerp, H., Tabor, N.J., Looy, C.V., 2008. The so-called “Paleophytic–Mesophytic”
 732 transition in equatorial Pangea—Multiple biomes and vegetational tracking of climate change through
 733 geological time. Palaeogeogr. Palaeoclimatol. Palaeoecol. 268, 152–163.
 734
 735
 736 Fielding, C.R., Alexander, J., 2001. Fossil trees in ancient fluvial channel deposits: evidence of seasonal
 737 and longer-term climatic stability. Palaeogeogr. Palaeoclimatol. Palaeoecol. 170, 59–80.
 738
 739 Fluteau, F., Besse, J., Broutin, J. Ramstein, G. 2001. The Late Permian climate. What can be inferred
 740 from climate modelling concerning Pangea scenarios and Hercynian range altitude? Palaeogeogr.
 741 Palaeoclimatol. Palaeoecol. 167, 39–71.
 742

743 Francillon-Vieillot, H., de Buffrénil, V., Castanet, J., Géraudie, J., Meunier, F.-J., Sire, J.-Y., Zylberberg,
 744 L., de Ricqlès, A., 1990. Microstructure and mineralization of vertebrate skeletal tissues, in: Carter, J.G.
 745 (Ed.), *Skeletal biomineralization: patterns, processes and evolutionary trends*. American Geophysical
 746 Union, San Francisco, 175–234.
 747
 748 Gastaldo, R.A., Demko, T.M., 2010. The relationship between continental landscape evolution and the
 749 plant-fossil record: long term hydrologic controls on preservation, in: Allison, P.A., Bottjer, D.J. (Eds.),
 750 *Taphonomy. Process and Bias Through Time*. *Top. Geobiol.* 32, 250–285.
 751
 752 Gastaldo, R.A., Pludow, B.A., Neveling, J., 2013, Mud aggregates from the Katberg Formation, South
 753 Africa: Additional evidence for Early Triassic degradational landscapes. *J. Sediment. Res.* 83, 531–540.
 754
 755 Gastaldo, R.A., Knight, C., Neveling, J., Tabor, N., 2014. Late Permian Paleosols from Wapadsberg Pass,
 756 South Africa: Implications for Changhsingian Climate. *Geo. Soc. Am. Bull.* 126, 665–679.
 757
 758 Germain, D., 2010. The Moroccan diplocaulid: the last lepospondyl, the single one on Gondwana. *Hist.*
 759 *Biol.* 22, 4–39.
 760
 761 Gibbs, M.T., Rees, P.M., Kutzbach, J.E., Ziegler, A.M., Behling, P.J., Rowley, D.B., 2002. Simulations
 762 of Permian climate and comparisons with climate sensitive sediments. *J. Geol.* 110, 33–55.
 763
 764 Giles, M.J., Soreghan, M.J., Benison, K.C., Soreghan, G.S., Hasiotis, S.T., 2013. Lakes, loess and
 765 paleosols in the Permian Wellington Formation of Oklahoma, USA: Implications for paleoclimate and
 766 paleogeography of the mid-continent. *J. Sedimentary Res.* 83, 825–846.
 767

768 Goudie, A., 2010. Namib sand sea: large dunes in an ancient desert, in: Migon, P. (Ed),
 769 Geomorphological Landscapes of the World. Springer, Netherlands, 163–169.
 770
 771 Gregory, J.W., 1906. The dead heart of Australia: A journey around Lake Eyre in the summer of 1901 -
 772 1902, with some account of the Lake Eyre basin and the flowing wells of central Australia. John Murray,
 773 London.
 774
 775 Hass, H., Rowe, N.P., 1999. Thin sections and wafering, in: Jones, T.P., Rowe, N.P. (Eds.), Fossil plants
 776 and spores: modern techniques. The Geological Society, London, 76–81.
 777
 778 Hernandez-Castillo, G.R., Rothwell, G.W., Mapes, G., 2001. Thucydiaceae fam. nov., with a review and
 779 reevaluation of Paleozoic walchian conifers. *Int. J. Plant. Sci.* 162, 1155–1185.
 780
 781 Jalil, N.E., Dutuit, J.M., 1996. Permian captorhinid reptiles from the Argana Formation, Morocco.
 782 *Palaeontology* 39, 907–918.
 783
 784 Jalil, N.E., Janvier, P., 2005. Les pareiasaures (Amniota, Parareptilia) du Permien supérieur du Bassin
 785 d'Argana, Maroc. *Geodiversitas* 27, 35–132.
 786
 787 Kiehl, J.T., Shields, C.A., 2005. Climate simulation of the latest Permian: Implications for mass
 788 extinction. *Geology* 33, 757–760.
 789
 790 Klappa, C.F., 1980. Rhizoliths in terrestrial carbonates: classification, recognition, genesis and
 791 significance. *Sedimentology* 27, 613–629.
 792

793 Köhler, M., Marín-Moratalla, N., Jordana, X., Aanes, R., 2012. Seasonal bone growth and physiology in
 794 endotherms shed light on dinosaur physiology. *Nature* 487, 358–361.
 795
 796 Konietzko-Meier, D., Klein, N., 2013. Unique growth pattern of *Metoposaurus diagnosticus*
 797 *krasiejowensis* (Amphibia, Temnospondyli) from the Upper Triassic of Krasiejów, Poland. *Palaeogeogr.*
 798 *Palaeoclimatol. Palaeoecol.* 370, 145–157.
 799
 800 Kriloff, A., Germain, D., Canoville, P., Vincent, P., Sache, M., Laurin, M., 2008. Evolution of bone
 801 microanatomy of the tetrapod tibia and its use in palaeobiological inference. *J. Evolution. Biol.* 21, 807–
 802 826.
 803
 804 Lamm, E.-T., 2013. Preparation and sectioning of specimens, in: Padian, K., Lamm, E.-T. (Eds.), *Bone*
 805 *histology of fossil tetrapods*. University of California Press, Berkeley, 55-160.
 806
 807 Looy, C.V., 2007. Extending the range of derived Late Paleozoic conifers: *Lebowskia* gen. nov.
 808 (Majonicaceae). *Int. J. Plant Sci.* 168, 957–972.
 809
 810 Looy, C.V., Stevenson, R.A., 2014. Earliest occurrence of autorotating seeds in conifers: the Permian
 811 (Kungurian-Roadian) *Manifera talaris* gen. et sp. nov. *Int. J. Plant Sci.* 175, 841–854.
 812
 813 Lucas, S.G., 2004. A global hiatus in the Middle Permian tetrapod fossil record. *Stratigraphy* 1, 47–64.
 814
 815 Machette, M.N., 1985. Calcic soils of the southwestern United States, in: Wiede, D.L. (Ed.), *Soils and*
 816 *Quaternary Geology of the Southwestern United States*, Geological Society of America Special Paper
 817 203, 1-22.
 818

819 McMahon, T.A., Murphy, R.E., Peel, M.C., Costelloe, J.F., Chiew, F.H.S., 2008. Understanding the
820 surface hydrology of the Lake Eyre Basin: Part 1—Rainfall. *J. Arid Environ.* 72, 1853–1868.
821

822 Montañez, I.P., Tabor, N.J., Niemeier, D., DiMichele, W.A., Frank, T.D., Fielding, C.R., Isbell, J.L., 2007.
823 CO₂-forced climate and vegetation instability during Late Paleozoic deglaciation. *Science* 315, 87–91.
824

825 Myers, T.S., Tabor, N.J., Jacobs, L.L., Mateus, O., 2012, Estimating soil pCO₂ using paleosol carbonates:
826 implications for the relationship between primary productivity and faunal richness in ancient terrestrial
827 ecosystems. *Paleobiology* 38, 585–604.
828

829 Nanson, G.C., Price, D.M., Short, S.A., 1992. Wetting and drying of Australia over the past 300 ka.
830 *Geology* 20, 791–794.
831

832 O'Keefe, F.R., Sidor, C.A., Larsson, H.C.E., Maga, A., Ide, O., 2005. The vertebrate fauna of the Upper
833 Permian of Niger—III, Morphology and ontogeny of the hindlimb of *Moradisaurus grandis* (Reptilia,
834 Captorhinidae). *J. Vertebr. Paleontol.* 25, 309–319.
835

836 Peyser, C.E., Poulsen, C.J., 2008. Controls on Permo-Carboniferous precipitation over tropical Pangaea: a
837 GCM sensitivity study. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 268, 181–192.
838

839 Philippe, M., Bamford, M.K., 2008. A key to morphogenera used for Mesozoic conifer-like woods. *Rev.*
840 *Palaeobot. Palyno.* 148, 184–207.
841

842 Rees, P.M., Ziegler, A.M., Gibbs, M.T., Kutzbach, J.E., Behling, P.J., Rowley, D.B., 2002. Permian
843 phytogeographic patterns and climate data/model comparisons. *J. Geol.* 110, 1–31.
844

845 Rothwell GW, Mapes, G., Hernandez-Castillo, G.R., 2005 *Hanskerpia* gen. nov. and phylogenetic
846 relationships among the most ancient conifers (Voltziales). *Taxon* 54, 733–750.

847

848 Sanchez, S., Schoch, R.R., 2013. Bone histology reveals a high environmental and metabolic plasticity as
849 a successful evolutionary strategy in a long-lived homeostatic Triassic temnospondyl. *Evol. Biol.* 40,
850 627–647.

851

852 Sanchez, S., Klembara, J., Castanet, J., Steyer, J.-S., 2008. Salamander-like development in a
853 seymouriamorph revealed by palaeohistology. *Biol. Letters* 4, 411–414.

854

855 Sanchez, S., Schoch, R.R., de Ricqlès, A., Steyer, J.-S., 2010a. Palaeoecological and palaeoenvironmental
856 influences revealed by long-bone palaeohistology: the example of the Permian branchiosaurid *Apateon*,
857 in: Vecoli, M., Clement, G., Meyer Berthaud, B. (Eds.), *The terrestrialization process: modelling complex*
858 *interactions at the biosphere-geosphere Interface*. *Geol. Soc. Spec. Publ.* 339, 139–149.

859

860 Sanchez, S., de Ricqlès, A., Schoch, R.R., Steyer J.-S., 2010b. Developmental plasticity of limb bone
861 microstructural organization in *Apateon*: histological evidence of paedomorphic conditions in
862 branchiosaurs. *Evol. Dev.* 12, 315–328.

863

864 Scheyer, T.M., Sander, P.M., 2009. Bone microstructures and mode of skeletogenesis in osteoderms of
865 three pareiasaur taxa from the Permian of South Africa. *J. Evolution. Biol.* 22, 1153–1162.

866

867 Schweitzer, H.J., 1963. Der weibliche Zapfen von *Pseudovoltzia liebeana* and seine Bedeutung für die
868 Phylogenie der Koniferen. *Palaeontogr. Abt. B* 113, 1–29.

869

870 Schweitzer, H.J., 1986. The land flora of the English and German Zechstein sequences, in: Harwood,
 871 G.M, Smith. D.B. (Eds.), The English Zechstein and related topics. Geological Society Special
 872 Publication 22. Geological Society-Blackwell Scientific, Oxford, 31–54.
 873
 874 Schweitzer, H.J., 1996 *Voltzia hexagona* (Bischoff) Geinitz aus dem mittleren Perm Westdeutschlands.
 875 Palaeontogr. Abt. B 293, 1–22.
 876
 877 Scotese, C.R., Boucot, A.J., McKerrow, W.S., 1999. Gondwanan palaeogeography and
 878 palaeoclimatology. J. Afr. Earth Sci. 28, 99–114.
 879
 880 Seely M.K., Ward, J.D., 1989. Ecological aspects of dunes in the central Namib Desert. Dunes '89
 881 excursion 2A field guide, Desert Ecological Research Unit, Gobabeb.
 882
 883 Semeniuk, V., 1981. Calcrete in Quaternary coastal dunes in southwestern Australia: a capillary-rise
 884 phenomenon associated with plants. J. Sediment. Petrol. 51, 47–68.
 885
 886 Semeniuk, V., 1985. Distribution of calcrete in Holocene coastal sands in relationship to climate,
 887 southwestern Australia. J. Sediment. Petrol. 55, 86–95.
 888
 889 Serventy, V., 1985. The desert sea. The miracle of Lake Eyre in flood. MacMillan Company of Australia,
 890 Melbourne.
 891
 892 Sidor, C.A., 2013. The vertebrate fauna of the Upper Permian of Niger—VIII. *Nigerpeton ricqlesi*
 893 (Temnospondyli: Cochleosauridae) and tetrapod biogeographic provinces. C. R. Palevol. 12, 463–472.
 894

895 Sidor, C.A., Blackburn, D.C., Gado, B., 2003. The vertebrate fauna of the Upper Permian of Niger—II.
896 Preliminary description of a new pareiasaur. *Palaeontol. Afr.* 39, 42–52.
897
898 Sidor, C.A., O'Keefe, F.R., Damiani, R., Steyer, J.S., Smith, R.M.H., Larson, H.C.E., Sereno, P.C., Ide,
899 O., Maga, A., 2005. Permian tetrapods from the Sahara show climate-controlled endemism in Pangea.
900 *Nature* 434, 886–889.
901
902 Sidor, C.A., Vilhena, D.A., Angielczyk, K.D., Huttenlocker, A.K., Nesbitt, S.J., Peacock, B.R., Steyer,
903 J.S., Smith, R.M.H., Tsuji, L.A.. 2013. Provincialization of terrestrial faunas following the end-Permian
904 mass extinction. *P. Natl. Acad. Sci. USA* 110, 8129–8133.
905
906 Smiley, T.M., Sidor, C.A., Maga, A., Ide, O., 2008. The vertebrate fauna of the Upper Permian of Niger.
907 —VI. First evidence of a gorgonopsian therapsid. *J. Vertebr. Paleontol.* 28, 543–547.
908
909 Smith, R.M.H., Mason, T.R., 1998. Sedimentary Environments and Trace Fossils of Tertiary Oasis
910 Deposits in the Central Namib Desert, Namibia. *Palaios* 13, 547–559.
911
912 Smith, R.M.H., Sidor, C.A., Tabor, N.J., Steyer, J.S., Chaney, D.S., 2009. Vertebrate taphonomy and
913 ichnology of a Permian 'wet desert' in central Pangea. *Palaeontol. Africana* 44, 179–183.
914
915 Smith, R.M.H., Sidor, C.A., Tabor, N.J., Steyer, J.S., 2015. Sedimentology and vertebrate taphonomy of
916 the Moradi Formation of northern Niger: A Permian wet desert in the tropics of Pangaea. *Palaeogeogr.*
917 *Palaeoclimatol. Palaeoecol.* 440, 128–141.
918
919 Steyer, J.-S., Jalil, N.E., 2009. First evidence of a temnospondyl in the Late Permian of the Argana Basin,
920 Morocco. *Spec. Pap. Palaeontol.* 81, 155–160.

921

922 Steyer, J.-S., Damiani, R., Sidor, C.A., O’Keefe, F.R., Larsson, H.C.E., Maga, A., Ide, O., 2006. The

923 vertebrate fauna of the Upper Permian of Niger. —IV. *Nigerpeton ricqlèsi* (Temnospondyli:

924 Cochleosauridae), and the edopoid colonization of Gondwana. J. Vertebr. Paleontol. 26, 18–28.

925

926 Steyer, J.-S., Laurin, M., Castanet, J., de Ricqlès, A., 2004. First histological and skeletochronological

927 data on temnospondyl growth: palaeoecological and palaeoclimatological implications. Palaeogeogr.

928 Palaeoclimatol. Palaeoecol. 206, 193–201.

929

930 Tabor, N.J., Montañez, I.P., 2004. Permo-Pennsylvanian alluvial paleosols (north-central Texas): High-

931 resolution proxy records of the evolution of early Pangean paleoclimate. Sedimentology 51, 851–884.

932

933 Tabor, N.J., Smith, R.M.H., Steyer, S.J., Sidor, C.A., Poulsen, C.J., 2011. The Permian Moradi Formation

934 of northern Niger: Paleosol morphology, petrography and mineralogy. Palaeogeogr. Palaeoclimatol.

935 Palaeoecol. 299, 200–213.

936

937 Tabor, N.J., Myers, T.S., Gulbranson, E., Rasmussen, C., Sheldon, N.D., 2013. Light stable isotope

938 composition of modern calcareous soil profiles in California: Implications for CO₂ reconstructions from

939 calcareous paleosols. Soc. Econ. Paleontol. Mineral. Spec. Publ. 104, 17–34.

940

941 Taquet, P., 1969. Première découverte en Afrique d’un reptile captorhinomorphe (Cotylosaurien). C. R.

942 Acad. Sci. Ser. D 268, 779–781.

943

944 Tooth, S., Nanson, G.C., 2000. The role of vegetation in the Fm. of anabranching channels in an

945 ephemeral river, northern plains, arid central Australia. Hydrol. Process. 14, 3099–3117.

946

947 Tsuji, L.A., Sidor, C.A., Steyer, J.-S., Smith, R.M.H., Tabor, N.J., Ide, O., 2013. The vertebrate fauna of
 948 the Upper Permian of Niger—VII. Cranial anatomy and relationships of *Bunostegos akokanensis*
 949 (Pareiasauria). J. Vertebr. Paleontol. 33, 747–763.

950

951 Turner, M.L., Tsuji, L.A., Ide, O., Sidor, C.A., 2015. The vertebrate fauna of the upper Permian of
 952 Niger—IX. The appendicular skeleton of *Bunostegos akokanensis* (Parareptilia: Pareiasauria), Journal of
 953 Vertebrate Paleontology. <http://dx.doi.org/10.1080/02724634.2014.994746>

954

955 Walter, H., 1984. Vegetation und Klimazonen. Grundriß der globalen Ökologie. Verlag E. Ulmer,
 956 Stuttgart.

957

958 Watson, A., 1992. Desert soils, in: Martini, I., Chesworth, W. (Eds.), Weathering, Soils and Paleosols.
 959 Elsevier, Amsterdam, 225–260.

960

961 Wright, V.P., 1990. A micromorphological classification of fossil and recent calcic and petrocalcic
 962 microstructures, in: Douglas, L.A. (Ed.), Soil Micromorphology: A Basic and Applied Science. Elsevier,
 963 Amsterdam, 401–407.

964

965 Yang, W., Feng, Qi., Liu, Y.Q., Tabor, N., Lin, J.Y., Thomas, S., Miggins, D., 2010. Depositional
 966 environments and cyclo- and chronostratigraphy of uppermost Carboniferous–Lower Triassic fluvial–
 967 lacustrine deposits. Global Planet. Change 73, 15–113.

968

Figure Captions

Fig. 1. Stratigraphy, depositional facies and fossils of the Late Permian upper Moradi Formation in Northern Niger. A= high sinuosity, gravelly sand bed channel fill; B= loessic floodplain with anastomosed distributary channels; C= end-point lake with exposed carbonate mudflats; D= gravel-bed braided channel fill. **Add info on new part of the figure.**

Fig. 2. Fossil leaves from the Moradi Formation found at localities T09 and T24. Scale bars equal 5 mm. (A) MNN MOR139, narrow strap-like leaves of Morphotype 1. (B) MNN MOR140, wide strap-like leaves of Morphotype 2. Lengthwise striations from thick veins can be seen in the inset. (C) MNN MOR141 and (D) MNN MOR142, needle-like leaves of Morphotype 3. (E) MNN MOR143 and (F) MNN MOR144 are impressions of the obovate leaves of Morphotype 4.

Fig. 3. Ovuliferous conifer dwarf shoots. Scale bars equal 5 mm. (A) MNN MOR114 shows a distinct lobed central scale. (B) MNN MOR115 shows that the scales come in a variety of thicknesses on each individual shoot. (C) MNN MOR116 displays the remarkable number of scales an individual shoot can possess. (D) MNN MOR118 is paired with its counterpart (E) to form (F), which is a rare example of how the full dwarf shoot would have looked in cross-section after burial, with a thickness of at least 4 mm (not accounting for desiccation). (G) MNN MOR44 shows a prominent lobed scale on the leftmost side. (H) MNN MOR117, (I) MNN MOR119, and (J) MNN MOR120 are examples of complete dwarf shoots, including stalks. (K) MNN MOR121 shows the point of attachment for two dwarf shoots joined at the base. It is likely that attached dwarf shoots were arranged in groups.

Fig. 4. Diagram of the Moradi tree trunk fossil. Scale bar in the lower left-hand corner of each inset represents 0.5 m. Red circles along the tree trunk diagram show position and size of branch scars. At the base of the trunk, from 0-1.5 m, the remains were badly damaged; likewise from 13.45-14.40 m, the

remains were broken and crumbled. From 23-24 m, the trunk was mostly buried. Likewise, past 25 m the trunk was not excavated.

Fig. 5. Histology of isolated pareiasaur bones from the Moradi Formation, Upper Permian of Niger. (A) Thin section, made from a scapula (MNN MOR148; ordinary transmitted light), displaying three layers: the basal layer (bl), the cancellous core (cc) and the outer layer (ol). (B) Detail of the region framed in A (polarized light). The red arrows show the regular pattern of lines of arrested growth (LAGs). (C) Radius (MNN MOR146) as preserved. The black arrow indicates the location where the thin section was made at mid-shaft. (D) Mid-shaft thin section of radius showing the compact cortical bone (c) and the spongiosa (s) (ordinary transmitted light). The frames, displayed from top to bottom, correspond to the images E-H. (E) Detail of the compact cortex (polarized light) showing a region of rearrangement with radial extrinsic fibers oriented in the same axis as the vascular mesh (vc). The primary bone tissue shows patches of pseudo-lamellar bone (plb). (F) Detail of the compact cortex displaying the organization of primary osteons (po) and the fibrous primary bone matrix (fb) in the same region as E (polarized light). (G) Detail of another region of compact cortical bone showing a regular LAG pattern (ordinary transmitted light). LAGs are highlighted by the red arrows. (H) Fibrous bone matrix (fb) pierced by circumferential rows of primary osteons (po) (polarized light).

Table 1. Sample information for carbonates analyzed from the Moradi, Quanzijie, Wutonggou, Balfour, Buntsandstein Formations.

Table 2. Sample information and stable carbon isotope values for calcite ($\Delta^{13}\text{C}_{\text{CC}}$) and organic matter ($\delta^{13}\text{C}_{\text{OM}}$) as well as estimates of Soil PCO_2 (in ppmV) and actual evapotranspiration (AET) from the Moradi, Quanzijie, Wutonggou, Balfour, Buntsandstein Formations.

Appendix

CO₂ mixing in soils - Soils may be characterized by mixing of one, two, or three components of soil CO₂, which can result in significantly different $\delta^{13}\text{C}$ values of soil CO₂ and the resulting pedogenic calcite (Tabor et al., 2013; Sheldon and Tabor, 2009; Feng and Yapp, 2009; Tabor and Yapp, 2005; Yapp, 2002; Hsieh and Yapp, 1999). The three CO₂ components are derived from (1) oxidation of *in situ* organic matter, (2) tropospheric CO₂, and (3) dissolution of pre-existing carbon-bearing minerals such as calcite in the soil.. None of the paleosols from the Moradi Formation or other sites in China, South Africa and Italy are stratigraphically associated with marine or lacustrine limestone strata. Therefore, the following discussion considers only 1- and 2-component mixing of soil CO₂ as viable explanations for carbon isotope data collected from Middle and Upper Permian paleosols.

Soils characterized by one component of CO₂ form under a chemically closed or semi-closed system (e.g. Tabor et al., 2013; Tabor et al., 2007; Gluyas, 1984; Sheldon and Tabor, 2009) that arises in settings where gaseous diffusion between the soil and the troposphere is limited, usually as a result of poorly drained, waterlogged conditions within the soil profile. Diffusion from the troposphere (including CO₂) to the wet soil is limited, and aqueous O₂ in the soil is quickly consumed by the oxidation of organic matter (Feng and Hsieh, 1998). Under these conditions, both bicarbonate and calcite will have $\delta^{13}\text{C}$ values that are equivalent to that of the oxidizing organic matter in the closed-system soil. Soil profiles forming under semi-closed conditions will result in calcite $\delta^{13}\text{C}$ values that are less than 14.6‰ more positive than coexisting organic matter (at 20° C), which reflects the diffusive 4.4‰ enrichment in $\delta^{13}\text{C}$ in the biologically-derived CO₂ (Cerling et al., 1991), as well as an additional 10.2‰ carbon isotope enrichment from gaseous CO₂ to calcite due to carbon isotope fractionation between carbonate species (at mildly alkaline pH; Romanek et al., 1992) in open system 2-component CO₂ mixing.

1046 For example, both samples ($n = 2$) from the Upper Permian Wutonggou Formation and over half ($n = 3$)
 1047 of the samples from the Upper Permian Balfour Formation have $\delta^{13}\text{C}_{\text{m(CC)}}$ values that are less than 14.6‰
 1048 more positive than co-existing $\delta^{13}\text{C}_{\text{OM}}$ values, and therefore do not permit meaningful estimates of Soil
 1049 CO_2 concentrations (C_{S} ; Table 2). The paleosols of the Balfour Formation exhibit drab matrix colors, little
 1050 to no soil structure, as well as pyrite and gypsum crystals and are interpreted to have been formed in
 1051 swampy bottomlands (Gastaldo et al., 2009; 2014; Tabor et al., 2007; Retallack et al., 2003). Upper
 1052 Permian Wutonggou paleosols have drab matrix colors, iron and manganese concretions, and are
 1053 interpreted to have formed on a poorly-drained lake plain (Yang et al., 2007; Thomas et al., 2010). The
 1054 results from these five samples were therefore interpreted to reflect calcite crystallization in a closed or
 1055 semi-closed system dominated by one component of CO_2 derived from oxidation of in-situ organic
 1056 matter, and are not considered further. All other Middle and Upper Permian samples have $\delta^{13}\text{C}_{\text{m(CC)}}$ and
 1057 $\delta^{13}\text{C}_{\text{OM}}$ values consistent with formation in a soil characterized by mixing of two components of CO_2 , and
 1058 therefore may provide information about variations in soil PCO_2 (C_{S}) among those sites.
 1059
 1060 Soils characterized by two-component soil CO_2 mixing assumes that the only two sources contributing to
 1061 soil CO_2 are CO_2 from the open atmosphere (tropospheric CO_2) and CO_2 from *in situ* oxidation of
 1062 biological carbon in the soil. Steady-state solutions to the one-dimensional Fickian diffusion equation
 1063 yield reasonable representations of $\delta^{13}\text{C}$ values of soil CO_2 with depth in modern soils characterized by
 1064 mixing of two sources of CO_2 (Cerling, 1984). In these models, PCO_2 and $\delta^{13}\text{C}$ values of Earth's
 1065 atmosphere are the upper boundary conditions, and there is a depth-dependent CO_2 production term that
 1066 describes the oxidation of organic carbon in the soil. Above a characteristic CO_2 production depth within
 1067 the soil, both the concentration and $\delta^{13}\text{C}$ of soil CO_2 will progressively approach values of the Earth's
 1068 atmosphere. The characteristic production depth in carbonate-forming soils varies, but tends to be at least
 1069 300-500 mm beneath the soil surface (Breecker et al., 2009; Cerling and Quade, 1993; Quade et al.,
 1070 1989). It is important to note that calcite samples from all paleosols considered herein were collected
 1071 from at least 500 mm beneath the interpreted surface of the paleosol profile, and therefore calcite $\delta^{13}\text{C}$

values should reflect the maximum contribution of CO₂ from oxidation of organic matter, and provide maximum soil PCO₂ (C_S) estimates. Thus, CO₂ in soil gas can be generally regarded as a mixture of two isotopically distinct end-members in these soils: atmospheric CO₂ and CO₂ derived from *in-situ* oxidation of soil organic matter. Calcite δ¹³C values that are ≥14.6‰ more positive than co-existing oxidizing organic matter reflect proportionately larger contributions from isotopically lighter atmospheric CO₂ to the soil atmosphere.

Permian Soil CO₂ estimates - Although its potential has been largely ignored to date (however, see Yapp and Poths, 1996; Sheldon and Tabor, 2009; Myers et al., 2012), paleosol calcite δ¹³C values may be tremendously useful as a proxy of soil PCO₂ and biological productivity for times prior to the evolution of C₄ plants and their rise to dominance in some ecosystems (pre-Miocene time). The following equation relates measured calcite and organic matter δ¹³C values to soil PCO₂ estimates:

$$C_{S(cc)} = C_{A(cc)} * \frac{\delta^{13}C_{A(cc)} - \delta^{13}C_{O(cc)}}{\delta^{13}C_{m(cc)} - \delta^{13}C_{O(cc)}}.$$

The subscript “(cc)” indicates calcite. δ¹³C_{m(cc)} is the measured δ¹³C value of pedogenic calcite (δ¹³C_{CC} in Table 2). The δ¹³C values subscripted with “A” and “O” are the δ¹³C values of calcite formed solely in equilibrium with CO₂ derived from the atmosphere and from oxidation of soil organic matter, respectively. C_A refers to the concentration of CO₂ gas in the soil if the only contribution to soil CO₂ were from the atmosphere. C_S is the actual concentration of CO₂ that was present in the soil. Both C_A and C_S are in units of ppmV. The value for (δ¹³C_{A(cc)} - δ¹³C_{O(cc)}) is assumed to be +16‰ (Yapp and Poths, 1991, 1992, 1993, 1996), because it represents the approximate difference between the estimated δ¹³C values of pre-industrial atmospheric CO₂ (-6.5‰) and recent C₃ continental biota (-27‰), after the latter value is adjusted for the diffusive 4.4 ‰ enrichment in δ¹³C in the biologically-derived CO₂ of the soil (Cerling et

1096 al., 1991). Note that other ($\delta^{13}C_{A(cc)} - \delta^{13}C_{O(cc)}$) values have been suggested (e.g. $\sim 13.7\text{‰}$, Arens et al.,
 1097 2000), but differences in this quantity result in relatively minor differences in estimated $p\text{CO}_2$ values
 1098 (Sheldon and Tabor, 2009; Tabor et al., 2004; Yapp and Poths, 1991).
 1099
 1100 $\delta^{13}C_{O(cc)}$ values may be estimated from $\delta^{13}C$ values of co-existing organic matter within the paleosol
 1101 profiles (Montañez et al., 2007; Tabor et al., 2004; Yapp and Poths, 1996; $\delta^{13}C_{OM}$ in Table 2). $\delta^{13}C_{O(cc)}$
 1102 estimates are 14.6‰ heavier than $\delta^{13}C$ values of organic matter, which reflects the diffusive 4.4‰
 1103 enrichment in $\delta^{13}C$ in the biologically-derived CO_2 (Cerling et al., 1991), as well as an additional 10.2‰
 1104 carbon isotope enrichment from gaseous CO_2 to calcite due to carbon isotope fractionation between
 1105 carbonate species (at mildly alkaline pH and 20°C ; Romanek et al., 1992). If $\delta^{13}C_{O(cc)}$ estimates are more
 1106 positive than $\delta^{13}C_{m(cc)}$ values from the same paleosol, C_s estimates will have a negative value. As
 1107 discussed earlier, negative C_s values have no physical meaning, and rather reflect that either (1) calcite
 1108 crystallization occurred in a chemically closed or semi-closed system or (2) the organic matter and calcite
 1109 $\delta^{13}C$ values do not share a common pedogenic origin.
 1110
 1111 C_A values are not known with certainty for Middle and Late Permian time. Estimates for Middle and Late
 1112 Permian paleoatmospheric PCO_2 range from $\sim\text{PAL}$ (Present Atmospheric Level; $\sim 300\text{ ppmV}$) to $\sim 17 \times$
 1113 PAL (e.g. Berner, 2005; Royer et al., 2004; Retallack, 2002). However, it is not necessary to know C_A
 1114 explicitly. This reflects the fact that the troposphere is a well-mixed system (e.g. Cerling, 1984), so for
 1115 any given interval of time in earth history, C_A will be effectively equivalent at all places upon Earth's
 1116 surface. Therefore, a comparative analysis of paleosol calcite and organic matter from
 1117 penecontemporaneous sites should delineate differences in C_s , as related to variations in biologically-
 1118 produced CO_2 , since C_A would have been equivalent among all sites considered. With respect to these
 1119 aforementioned considerations, it was assumed that all paleosol samples (1) are penecontemporaneous
 1120 (with respect to significant temporal changes in C_A values) and (2) formed in the presence of a constant

1121 atmospheric PCO_2 (C_A) value of 300 ppmV. Note that this assumed C_A value will result in relatively low
1122 C_S estimates, and that higher C_A values will produce correspondingly higher estimates of C_S (Sheldon and
1123 Tabor, 2009; Tabor et al., 2013).
1124

1125 **Citations for the Appendix**

- 1126 Arens, N.C., Jahren, A.H., Amundson, R., 2000. Can C3 plants faithfully record the carbon isotopic
1127 composition of atmospheric carbon dioxide? *Paleobiology* 26, 134–164.
- 1128 Berner, R.A., 2005. GEOCARBSULF: A combined model for phanerozoic atmospheric O₂. *Geochimica*
1129 *et Cosmochimica. Acta* 70, 5653–5664.
- 1130 Breecker, D.O., Sharp, Z.D., McFadden, L.D., 2009. Seasonal bias in the formation and stable isotopic of
1131 pedogenic carbonate in modern soils from central New Mexico, USA. *Geological Society of*
1132 *America Bulletin* 121, 630–640.
- 1133 Cerling, T.E., 1984. The stable isotopic composition of modern soil carbonate and its relationship to
1134 climate. *Earth and Planetary Science Letters* 71, 229–240.
- 1135 Cerling, T.E., Quade, J., 1993. Stable carbon and oxygen isotopes in soil carbonates, in: Swart, P.K.,
1136 Lohmann, K.C., McKenzie, J., Savin, S. (Eds.), *Climate Change in Continental Isotopic Records*.
1137 Washington (DC). *Geophysical Monograph Series* 78, American Geophysical Union, 217–231.
- 1138 Cerling, T.E., Solomon, D.K., Quade, J., Bowman, J.R., 1991, On the isotopic composition of carbon in
1139 soil carbon dioxide. *Geochimica et Cosmochimica. Acta* 55, 3403–3405.
- 1140 Feng, J., and Hsieh, Y.P., 1998. Wetlands and aquatic processes; sulfate reduction in freshwater wetland
1141 soils and the effects of sulfate and substrate loading. *Journal of Environmental Quality* 27, 968–972.
- 1142 Feng, W., Yapp, C.J., 2009. Paleoenvironmental implications of concentration and ¹³C/¹²C ratios of
1143 Fe(CO₃)OH in goethite from a mid-latitude Cenomanian laterite in southwestern Minnesota.
1144 *Geochimica et Cosmochimica. Acta* 73, 2559–2580.
- 1145 Gastaldo, R.A., Neveling, J., Clark, C.K., Newbury, S.S., 2009. The terrestrial Permian-Triassic boundary
1146 event bed is a non-event. *Geology* 37, 199–202.
- 1147 Gastaldo, R.A., Knight, C., Neveling, J., Tabor, N., 2014. Late Permian Paleosols from Wapadsberg Pass,
1148 South Africa: Implications for Changhsingian Climate. *Geo. Soc. Am. Bull.* 126, 665–679.
- 1149 Gluyas, J.G., 1984. Early carbonate diagenesis within Phanerozoic shales and sandstones of the NW
1150 European shelf. *Clay Mineralogy* 19, 309–321.

1151 Hsieh, J.C.C., Yapp, C.J., 1999. Stable carbon isotope budget of CO₂ in a wet, modern soil as inferred
 1152 from Fe(CO₃)OH in pedogenic goethite: Possible role of calcite dissolution. *Geochimica et*
 1153 *Cosmochimica Acta* 63, 767–783.

1154 Montañez, I.P., Tabor, N.J., Niemeier, D., DiMichele, W.A., Frank, T.D., Fielding, C.R., Isbell, J.L., 2007.
 1155 CO₂-forced climate and vegetation instability during Late Paleozoic deglaciation. *Science* 315, 87–
 1156 91.

1157 Myers, T.S., Tabor, N.J., Jacobs, L.L., Mateus, O., 2012, Estimating soil pCO₂ using paleosol carbonates:
 1158 implications for the relationship between primary productivity and faunal richness in ancient
 1159 terrestrial ecosystems. *Paleobiology* 38, 585–604.

1160 Quade, J., Cerling, T.E., Bowman, J.R., 1989, Development of Asian monsoon revealed by marked
 1161 ecological shift during the latest Miocene in northern Pakistan: *Nature* 342, 162–66.

1162 Retallack, G.J., 2002. Carbon dioxide and climate over the past 300 million years, in: *Understanding*
 1163 *climate change: proxies, chronology and ocean-atmosphere interactions*, Gröcke, D.R., Kucera, M.
 1164 (Eds.), *Philosophical Transactions of the Royal Society of London* 360, 659–674.

1165 Retallack, G.J., Smith, R.M.H., Ward, P.D., 2003, Vertebrate extinction across the Permian-Triassic
 1166 boundary in the Karoo Basin of South Africa. *Geological Society of America Bulletin* 115, 1133–
 1167 1152.

1168 Romanek, C.S., Grossman, E.L., Morse, J.W., 1992, Carbon isotopic fractionation in synthetic aragonite
 1169 and calcite: effects of temperature and precipitation rate. *Geochimica et Cosmochimica. Acta* 56,
 1170 419–430.

1171 Royer, D.L., Berner, R.A., Montañez, I.P., Tabor, N.J., Beerling, D.J., 2004. CO₂ as a primary driver of
 1172 Phanerozoic climate change. *GSA Today* 14, 4–10.

1173 Sheldon, N.D., Tabor, N.J., 2009. Quantitative Paleoenvironmental and Paleoclimatic Reconstruction
 1174 using paleosols. *Earth Science Reviews* 95, 1–52.

1175 Tabor, N.J., Yapp, C.J., 2005. Incremental vacuum dehydration-decarbonation experiments on a natural
1176 gibbsite (-Al(OH)_3): CO_2 abundance and ^{13}C values. *Geochimica et Cosmochimica. Acta* 69, 519–
1177 527.

1178 Tabor, N.J., Yapp, C.J., Montañez, I.P., 2004. Goethite, Calcite, and Organic Matter from Permian and
1179 Triassic Soils: Carbon Isotopes and CO_2 Concentrations. *Geochimica et Cosmochimica. Acta* 68,
1180 1503–1517.

1181 Tabor, N.J., Montanez, I.P., Steiner, M., Schwindt, D., 2007. The ^{13}C values of Permo-Triassic carbonates
1182 from South Africa reflect a stinking, sulfurous swamp, not atmospheric conditions.
1183 *Palaeogeography, Palaeoclimatology, Palaeoecology* 252, 370–381.

1184 Tabor, N.J., Myers, T.S., Gulbranson, E., Rasmussen, C., Sheldon, N.D., 2013. Light stable isotope
1185 composition of modern calcareous soil profiles in California: Implications for CO_2 reconstructions
1186 from calcareous paleosols. *SEPM Special Publication* 104, 17–34.

1187 Thomas, S.G., Tabor, N.J., Yang, W., 2010. Morphology, mineralogy and elemental chemistry of
1188 paleosols across the Permo-Triassic boundary in strata from the Turpan basin, Bodga Shan, NW
1189 China: revised and accepted by *Palaeogeography, Palaeoclimatology, Palaeoecology* 308, 41–64.

1190 Yang, W., Liu, Y., Feng, Q., Lin, J., Zhou, D., Wang, D., 2007. Sedimentary evidence of Early–Late
1191 Permian mid-latitude continental climate variability, southern Bogda Mountains, NW China.
1192 *Palaeogeography, Palaeoclimatology, Palaeoecology* 252, 239–258.

1193 Yapp, C.J., 2002. Erratum to Crayton J. Yapp (2001), “Mixing of CO_2 in surficial environments as
1194 recorded by the concentration and $\delta^{13}\text{C}$ values of the $\text{Fe}(\text{CO}_3)\text{OH}$ component in goethite,” *GCA*,
1195 Vol. 65, 4115–4130. *Geochimica et Cosmochimica Acta* 66, 1497.

1196 Yapp, C.J., Poths, H., 1991. $^{13}\text{C}/^{12}\text{C}$ ratios of the Fe(III) carbonate component in natural goethites, in:
1197 Taylor Jr, H.P., O’Neil, J.R., Kaplan, I.R. (Eds.), *In Stable Isotope Geochemistry: A Tribute to*
1198 Samuel Epstein, *Geochemical Society Special Publication* 3, 257–270.

1199 Yapp, C.J., Poths, H., 1992. Ancient atmospheric CO_2 pressures inferred from natural goethites. *Nature*
1200 355, 342–344.

- 1201 Yapp, C.J., Poths, H., 1993. The carbon isotope geochemistry of goethite (α -FeOOH) in ironstone of the
1202 Upper Ordovician Neda Formation, Wisconsin, USA: Implications for early Paleozoic continental
1203 environments. *Geochimica et Cosmochimica Acta* 57, 2599–2611.
- 1204 Yapp, C.J., Poths, H., 1996. Carbon isotopes in continental weathering environments and variations in
1205 ancient atmospheric CO₂ pressure. *Earth and Planetary Science Letters* 137, 71–82.

FACIES

quartz pebble and cobble conglomerate		trough cross-bedding
silcrete clast and nodule conglomerate		horizontal lamination
carbonate mudrock breccia		root moulds
thickly-bedded massive siltstone		invertebrate burrows
thinly-bedded calcareous mudstone		vertebrate fossil
ripple cross-lamination		tetrapod trackways
calcite-filled shrinkage cracks and nodules		palaeocurrent direction
desiccation cracks		leaf fossil
lateral-accretion units		fossil wood
		unconformity

figure 1

figure 1

figure 2

figure 3

figure 4

figure 5

Table 1 calcite isotope data

TABLES

Sample	Section	Position in Stratigraphy	Level (m)	Paleosol Type	Host Sediment	Description	Texture	Wt % Calcite	$\delta^{13}\text{C}$	$\delta^{18}\text{O}$
LMOR1	1	Lower	N/A	A	sandstone	rhizolith	micrite	24.2	-1.9	-6.9
LMOR2	1	Lower	N/A	B	mudstone	Calcans	micrite	52.1	-2.9	-7
LMOR4	1	Upper	N/A	B	mudstone	Calcans	micrite	64.2	-2.9	-7
LMOR8	1	Upper	N/A	B	mudstone	Calcans	micrite	23.2	-3.2	-6.1
UMOR1A	2	Upper	15	B	mudstone	BKII nodules	micrite	78.2	-3.2	-5.8
UMOR1A	2	Upper	15	B	mudstone	BKII nodules	micrite	81.3	-3.3	-5.7
UMOR9A	2	Upper	28	A	mud-ball conglomerate	rhizolith	micrite	88.6	-4.9	-4.3
UMOR9B	2	Upper	28	A	mud-ball conglomerate	rhizolith	micrite	50.2	-5.5	-6.1
UMOR10A	2	Upper	28	A	mud-ball conglomerate	rhizolith	micrite	89.7	-5.5	-5.2
UMOR11A	2	Upper	29	A	sandstone	rhizolith	micrite	93.1	-4	-5.1
UMOR12A	2	Upper	32	A	sandstone	calcrete	micrite	90.6	-3.7	-2.1
UMOR12B	2	Upper	32	A	sandstone	calcrete	micrite	90.9	-4.9	-3.1
2MOR2	3	middle	50	B	mudstone	Calcans	micrite	60.2	-3	-7.9
2MOR6	3	middle	50.5	B	mudstone	Calcans	micrite	56.3	-3.5	-6.4
2MOR6	3	middle	50.5	B	mudstone	Calcans	micrite	56.5	-2.6	-6.3
2MOR8	3	middle	5.2	B	mudstone	BKII nodules	micrite	37.5	-5.8	-8
2MOR9	3	middle	11.7	B	mudstone	BKII nodules	micrite	47.2	-2.6	-7.4
2MOR9A	3	middle	11.7	B	mudstone	BKII nodules	sparite	53.9	-3.4	-7.1
2MOR9B	3	middle	11.7	B	mudstone	BKII nodules	micrite	59.7	-2.6	-7.4
2MOR11	3	middle	27	B	mudstone	BKII nodules	micrite	57.1	-5.2	-3.8
2MOR12	3	middle	40	A	sandstone	Rhizolith	micrite	33.6	-3	-7.9
MOR15A	3	upper	90	A	sandstone	Rhizolith	micrite	100.1	-3.8	-4.2
2MOR15A	3	upper	90	A	sandstone	Rhizolith	micrite	94.5	-3.6	-4.1
2MOR17A	3	upper	98.5	A	sandstone	Rhizolith	micrite	87.5	-3.5	-3.8
2MOR18	3	upper	101	A	mudstone	BKII nodules	micrite	42.7	-1.8	-8.4
2MOR18	3	upper	101	A	mudstone	BKII nodules	micrite	42.7	-1.9	-8.4
MW6	4	middle	N/A	A	sandstone	rhizolith	micrite	89.1	1	1.1
MW6A	4	middle	N/A	A	sandstone	rhizolith	micrite	88.7	-3.8	-4.4
MW7A	4	middle	N/A	A	sandstone	rhizolith	micrite	87.8	-4	-3.6
MW7cc1	4	middle	N/A	A	sandstone	rhizolith	sparite	85.5	-4.6	-6
MW7cc2	4	middle	N/A	A	sandstone	rhizolith	sparite	82.9	-4.6	-6
MW8	4	middle	N/A	A	sandstone	rhizolith	micrite	63.0	-3.3	-4.1
MW8A	4	middle	N/A	A	sandstone	rhizolith	micrite	65.7	-3	-7.2
MW9	4	middle	N/A	A	sandstone	rhizolith	micrite	65.8	-3.2	-4.3

MW9A	4	middle	N/A	A	sandstone	rhizolith	micrite	82.2	-3.4	-3.6
MW9B	4	middle	N/A	A	sandstone	rhizolith	micrite	80.0	-3	-4.2
MW9C	4	middle	N/A	A	sandstone	rhizolith	micrite	51.3	-2.8	-7.1
MW10a	4	middle	N/A	A	sandstone	rhizolith	micrite	99.0	-3.8	-4.2
MW10A	4	middle	N/A	A	sandstone	rhizolith	micrite	99.0	-3.6	-3.7
MW10A	4	middle	N/A	A	sandstone	rhizolith	micrite	99.0	-3.6	-3.8
MW10B	4	middle	N/A	A	sandstone	rhizolith	micrite	89.8	-3.7	-4.8
MW10C	4	middle	N/A	A	sandstone	rhizolith	micrite	85.2	-3.9	-4.8
MW10C	4	middle	N/A	A	sandstone	rhizolith	micrite	63.2	-3.9	-4.4
				N/A		groundwater				
MW12a	4	middle	32		sandstone	concretion	micrite	55.0	-3.7	-4.4
MW13A	4	middle	34.1	A	sandstone	rhizolith	micrite	90.4	-3.4	-3.9
MW15A	4	middle	40.2	A	mudstone	BKII nodules	micrite	100.7	-3.6	-3.6
MW15B	4	middle	40.2	A	mudstone	BKII nodules	micrite	95.2	-3.4	-4
MW15C	4	middle	40.2	A	mudstone	BKII nodules	micrite	94.9	-3.3	-3.7
MW15D	4	middle	40.2	A	mudstone	BKII nodules	micrite	89.8	-2.9	-3.4
MW16A	4	middle	47	A	sandstone	rhizolith	micrite	88.9	-2.4	-4.9
MW17	4	middle	50.6	A	siltstone	rhizolith	micrite	51.9	-2.3	-4.3
MW24	4	middle	64.7	C	mudstone	BKII nodule	micrite	47.3	-1.8	-7.8
MW24	4	middle	64.7	C	Mudstone	BKII nodule	Micrite	19.1	-1.8	-7.8
MW27A	4	upper	75	A	sandstone	rhizolith	micrite	89.4	-4.1	-6.9
MW27A	4	upper	75	A	sandstone	rhizolith	micrite	90.0	-3.6	-4.9
MW27B	4	upper	75	A	sandstone	rhizolith	micrite	47.5	-2.6	-6
MW27	4	upper	75	A	sandstone	rhizolith	micrite	91.2	-3.8	-6
MW28	4	upper	78	A	sandstone	rhizolith	micrite	87.5	-3.7	-6
MW29	4	upper	81	B	mudstone	BKII nodules	micrite	34.2	-2.2	-5.2
				N/A		groundwater				
MW31	4	upper	84.8		sandstone	concretion	micrite	39.5	-0.8	-6
MW32a	4	upper	84.8	A	sandstone	rhizolith	micrite	56.7	-2.5	-4.8
MW32B	4	upper	84.8	A	sandstone	rhizolith	micrite	63.2	-0.5	-0.9
MW33	4	upper	86	B	mudstone	calcans	micrite	31.5	0.5	0.5
MW34	4	upper	94.8	A	sandstone	rhizolith	micrite	67.3	1	2.3
MW35	4	upper	95.5	B	mudstone	BKIII horizon	micrite	75.3	-1.8	-2.3
NW36	4	upper	96.3	B	mudstone	BKII nodules	micrite	71.0	-2.8	-5.3
MW37	4	upper	100.5	B	mudstone	BKIII horizon	micrite	68.5	1.5	3.3
MT7	5	upper	7.5	C	mudstone	BKII nodule	micrite	78.3	-8.2	-10.2
MT8	5	upper	2	C	mudstone	BKII nodule	micrite	55.3	-0.5	-3.7

China – Middle Permian Quanzijie formation										
QF2a	-	-	-	-	mudstone	rhizolith	micrite	82.1	-3.9	-9.4
QF2b	-	-	-	-	mudstone	rhizolith	micrite	71.7	-4.5	-9.9
QF10a	-	-	-	-	mudstone	BKII nodule	micrite	84.3	-5.2	-9.4
QF10b	-	-	-	-	mudstone	BKII nodule	micrite	77.9	-4.9	-9.9
China – Upper Permian Wutonggou formation										
WY-A1	-				mudstone	BKII nodule	micrite		-10.4	-4.1
TR3-13	-				mudstone	BKII nodule	micrite		-13.5	-8.0
South Africa – Upper Permian Balfour formation										
23606-9a					mudstone	BKII nodule	micrite	35.1	-10.7	-21.5
23606-9b					mudstone	BKII nodule	micrite	35	-9.3	-21.6
21606-8a					mudstone	BKII nodule	micrite	43	-14.4	-14.8
21606-8b					mudstone	BKII nodule	micrite	39	-14.0	-14.3
23606-23a					mudstone	BKII nodule	micrite	46.3	-5.5	-21.3
23606-23b					mudstone	BKII nodule	micrite	47.2	-5.3	-21.0
21606-6a					mudstone	BKII nodule	micrite	42.5	-9.1	-22.2
21606-6b					mudstone	BKII nodule	micrite	50	-15.4	-21.7
21606-1a					mudstone	BKII nodule	micrite	38.7	-15.4	-22.0
21606-1b					mudstone	BKII nodule	micrite	24.7	-17.1	-21.7
Italy – Upper Permian Buntsandstein formation										
JE1a					mudstone	BKII nodule	micrite	*	-4.5	-1.5
JE1b					mudstone	BKII nodule	micrite	*	-4.5	-0.9
JE8a					mudstone	BKII nodule	micrite	*	-5.4	-3.7
JE8b					mudstone	BKII nodule	micrite	*	-5.9	-4.9
JE10					mudstone	BKII nodule	micrite	*	-5.3	-2.8
JE11					mudstone	BKII nodule	micrite	*	-5.0	-2.6
BU3a1					mudstone	BKII nodule	micrite	*	-3.2	-8.9
BU3a2					mudstone	rhizolith	micrite	*	-3.5	-8.9
BU3a3					mudstone	BKII nodule	micrite	*	-3.7	-9.4

*Samples were analyzed by automated common acid-bath on-line system at University of California Davis. This method does not include measurements of wt. % calcite.

table 2				Wt □□□					
Sample	Position in Stratigraphy	Paleosol Type	Description	□ carbon	$\delta^{13}\text{C}_{\text{OM}}$	$\delta^{13}\text{C}_{\text{CC}}^1$	$\Delta^{13}\text{C}_{\text{CC-OM}}$	Soil PCO_2	AET (mm/yr)
Niger – Middle/Upper Permian Moradi formation									
LMOR1	Lower	A	rhizolith	0.170	-24.3	-1.9	22.4	600	73
MW7	Middle	A	rhizolith	0.329	-23.6	0.8	24.4	500	49
MW18	Middle	A	rhizolith	0.417	-22.7	-3.5	19.2	1000	148
MW19	Middle	A	rhizolith	1.187	-21.9	-2.9	19.0	1100	163
LMOR8	Upper	B	BKII nodule	0.214	-24.7	-3.2	21.5	700	95
UMOR1	Upper	B	rhizolith	0.117	-25.0	-3.3	21.7	700	95
UMOR11	Upper	B	Calcans	0.053	-24.5	-4	20.5	800	114
MW27B	Upper	C	BKII nodule	0.091	-20.8	-2.6	18.2	1300	190
MW37	Upper	B	Calcans	0.098	-24.4	-2.1	22.3	600	73
China – Middle Permian Quanzijie formation									
QF2	-	-	rhizolith	0.620	-19.8	-4.2	15.6	4800	465
QF10	-	-	rhizolith	0.547	-20.3	-5.1	15.2	8000	622
China – Upper Permian Wutonggou formation									
TD83	-	-	BKII nodule		-21.5	-10.4	11.1	N/A	N/A
TR-H2	-	-	BKII nodule		-24.7	-13.5	11.2	N/A	N/A
South Africa – Upper Permian Balfour formation									
23606-9	-	-	Calcans	0.150	-26.1	-10.7	15.4	6000	529
21606-8	-	-	rhizolith	0.097	-25.8	-14.2	11.6	N/A	N/A
23606-23	-	-	rhizolith	0.088	-25.9	-5.4	20.4	800	114
23606-6	-	-	rhizolith	0.071	-26.6	-15.4	11.2	N/A	N/A
21606-1	-	-	rhizolith	0.040	-25.3	-22.0	3.3	N/A	N/A
Italy – Upper Permian Buntsandstein formation									
JE1 ²	-	-	Coaly wood	- ²	-22.6	-4.5	18.1	1400	203
JE9 ³	-	-	Coaly wood	- ²	-21.6	-5.3	16.3	2800	335
JE11	-	-	Coaly wood	- ²	-22.3	-5.0	17.3	1800	248
BU4 ⁴	-	-	Coaly wood	- ²	-22.4	-3.7	18.7	1200	177

¹ $\delta^{13}\text{C}_{\text{cc}}$ values reported in this table are the average of all calcite $\delta^{13}\text{C}$ values collected for the sample (see Table 2).

² $\delta^{13}\text{C}^{\text{OM}}$ values for these samples were determined from fossilized organic matter in close stratigraphic association with paleosol calcite samples. These organic matter samples were analyzed with a Carlo-Erba continuous flow Elemental Analyzer interfaced with a Fisons Optima isotope ratio mass spectrometer. This method does not provide an estimate of wt. % organic carbon.

³ $\delta^{13}\text{C}_{\text{cc}}$ values are average of JE-1 calcite samples in Table 2.

⁴ $\delta^{13}\text{C}_{\text{cc}}$ values are average of JE-8 calcite samples in Table 2.

⁵ $\delta^{13}\text{C}_{\text{cc}}$ values are average of BU3 calcite samples in Table 2.