**Redox-dependent distribution of early macro-organisms: Evidence from the terminal
Ediacaran Khatyspyt Formation in Arctic Siberia**

Huan Cui ^{a,b,*}, Dmitriy V. Grazhdankin ^{c,d}, Shuhai Xiao ^e, Sara Peek ^{b,1}, Vladimir I. Rogov ^c,
Natalia V. Bykova ^e, Natalie Sievers ^{b,2}, Xiao-Ming Liu ^f, Alan J. Kaufman ^b

^a Department of Geoscience and NASA Astrobiology Institute, University of Wisconsin–
Madison, Madison, WI 53706, USA

^b Department of Geology and Earth System Science Interdisciplinary Center, University of
Maryland, College Park, MD 20742, USA

^c Trofimuk Institute of Petroleum Geology and Geophysics, Siberian Branch Russian Academy
of Sciences, Novosibirsk 630090, Russia

^d Department of Geology and Geophysics, Novosibirsk State University, Novosibirsk 630090,
Russia

^e Department of Geosciences, Virginia Tech, Blacksburg, VA 24061, USA

^f Department of Geological Sciences, University of North Carolina, Chapel Hill, NC 27599, USA

* Corresponding author: Huan.Cui@Wisc.EDU (H. Cui)

¹ Current address: United States Geological Survey, Menlo Park, CA 94025, USA

² Current address: School of Earth, Energy & Environmental Sciences, Stanford University, CA
94305, USA

Abstract

The Ediacaran Period witnessed the first appearance of macroscopic animal life in
Earth's history. However, the biogeochemical context for the stratigraphic occurrence of early

metazoans is largely uncertain, in part due to the dearth of integrated paleobiological and chemostratigraphic datasets. In this study, a comprehensive geochemical analysis was conducted on the fossiliferous Khatyspyt Formation in Arctic Siberia, in order to gain insights into the Ediacaran paleoenvironments. This study was designed to specifically address the relationship between paleoredox conditions and Ediacaran fossil occurrences in the Khatyspyt Formation. Our data reveal a dramatic shift in pyrite sulfur isotope compositions ($\delta^{34}\text{S}_{\text{pyrite}}$) from ca. -20% to ca. 50% , and this shift is intriguingly associated with the first occurrence of Ediacara-type macrofossils at the studied section, suggesting a possible link between seawater redox conditions and distribution of early macroscopic organisms. Based on multiple lines of sedimentological and geochemical evidence, we propose that the development of oceanic euxinia — which may be widespread in the continental margins due to enhanced oxidative weathering in the terminal Ediacaran Period — may have locally prohibited the colonization of Ediacara-type organisms and resulted in low $\delta^{34}\text{S}_{\text{pyrite}}$ values in the lower Khatyspyt Formation. In the middle and upper Khatyspyt Formation, progressive secular transition from euxinic to non-euxinic and more habitable conditions may have allowed for the colonization of Ediacara-type and other macro-organisms.

Keywords: Chemostratigraphy; Redox condition; Early metazoan distribution; Carbon and sulfur cycles; Ocean stratification; Ediacaran Period

1. Introduction

The Ediacaran Period (ca. 635–541 Ma) holds the answers to key questions related to the origins of the modern Earth system. In particular, Ediacaran strata contain the planet’s first unambiguous evidence of macroscopic metazoans and assemblages of “Ediacara-type” fossils

([Xiao and Laflamme, 2009](#)). Our present understanding of the origin of animals depends critically on the ability to interpret fossil impressions in siliciclastic sediments made by the Ediacara-type macro-organisms ([Gehling, 1999](#); [Narbonne, 2005](#); [Fedonkin et al., 2007](#)) and to document their spatial and temporal distribution ([Grazhdankin, 2011, 2014](#); [Narbonne et al., 2014](#)). However, siliciclastic rocks, particularly coarse-grained sandstones, offer limited opportunities for biogeochemical reconstructions of the deep-time record. Fortunately, Ediacara-type macrofossils can also be preserved in carbonate rocks, including the Khatyspyt Formation in Siberia ([Fedonkin, 1990](#); [Nagovitsin et al., 2015](#); [Rogov et al., 2015](#)) and the Dengying Formation in South China ([Ding and Chen, 1981](#); [Sun, 1986](#); [Xiao et al., 2005](#); [Chen et al., 2014](#)). Chemostratigraphic investigation of these fossiliferous carbonate successions can provide critical geochemical data that complement our understanding from siliciclastic successions.

Ediacaran animals and perhaps Ediacara-type macro-organisms are believed to be oxygen-breathing life forms ([Cloud, 1968](#); [Cloud, 1976](#); [Xiao, 2014](#)), therefore a putative rise in atmospheric oxygen during the Ediacaran Period ([Derry et al., 1992](#); [Kaufman et al., 1993](#)) may have dictated their evolutionary trajectories and their environmental distribution. Compilations of redox-sensitive proxies at broad timescale suggest a general pattern of rising atmospheric oxygen levels during the late Proterozoic ([Kah et al., 2004](#); [Canfield et al., 2007](#); [Kump, 2008](#); [Kah and Bartley, 2011](#); [Shields-Zhou and Och, 2011](#); [Lyons et al., 2014](#); [Planavsky et al., 2014](#); [Liu et al., 2016](#)), although oceanic anoxia has also been argued to remain persistent in many parts of the ocean even in the Ediacaran Period ([Canfield et al., 2008](#); [Sperling et al., 2015c](#); [Reinhard et al., 2016](#); [Sahoo et al., 2016](#)). In addition, individual case studies on integrative chemostratigraphy and biostratigraphy of the terminal Ediacaran strata, including the Blueflower Formation in NW Canada ([Johnston et al., 2013](#); [Macdonald et al., 2013](#); [Sperling et al., 2015a](#)),

the Nama Group in Namibia ([Hall et al., 2013](#); [Darroch et al., 2015](#); [Wood et al., 2015](#)), and the Dengying Formation in South China ([Duda et al., 2014](#); [Cui et al., 2016a](#)) reveal dynamic redox histories in these depositional basins, suggesting a complex relationship between the emergence of macrometazoans and the putative Ediacaran oxygenation. To further test the various hypotheses about the relationship between the rise of animals and atmospheric oxygen levels, we carried out an integrative investigation of the terminal Ediacaran Khatyspyt Formation in northern Siberia, which contains a moderate diversity of Ediacara-type macrofossils and abundant carbonate rocks for chemostratigraphic analysis ([Knoll et al., 1995](#); [Pelechaty et al., 1996a](#)). The goal of this study is to assess the effect of redox conditions on the distribution of early macro-organisms — particularly Ediacara-type macro-organisms — in the Khatyspyt Formation, using carbon, oxygen, sulfur, and strontium isotopes, as well as trace element concentrations.

2. Geological background

2.1. Lithostratigraphy

The fossiliferous Khatyspyt Formation is well exposed along the right-hand side tributaries of the Olenek River (i.e., the Khorbusuonka and Kersyuke rivers) that drains the Olenek Uplift in the northeastern part of the Siberian platform, Republic of Sakha, Russia (**Fig. 1, 2A**) ([Nagovitsin et al., 2015](#)). Sedimentological observations from the studied Khatyspyt interval (0–130 m in Fig. 3) suggest an overall shelf marine environment with relatively deeper depth compared with the Maastakh Formation and the overlying Turkut Formation. The lowermost Khatyspyt Formation is dominated by thick packages of intraclastic limestone occurring as channelized bodies up to 7 m in thickness, with abundant tilted angular intraclasts (**Fig. 2B**) and

soft-sediment deformation structures (**Fig. 2C**), and are laterally persistent over tens to hundreds of meters. The majority of this formation consists of finely laminated medium-bedded limestone with occasional small cross beddings (**Fig. 2D**). The succession also includes intervals of alternating thin layers of limestone and shale (**Fig. 2E**), abundant carbonaceous compressions preserved on the bedding planes (**Fig. 2F**), packages of thin-bedded (**Fig. 2G**) and thick-bedded limestone (**Fig. 2H**). The latter often show wavy bedding (**Fig. 2J**). Finely laminated limestones (**Fig. 2I**) sometimes contain abundant accumulations of *Nenoxites* fossils, which were initially regarded as meniscate trace fossils ([Sokolov and Fedonkin, 1984](#); [Rogov et al., 2012](#); [Rogov et al., 2013a](#); [Rogov et al., 2013b](#); [Rogov et al., 2015](#)), but have alternatively been interpreted as enigmatic tubular body fossils similar to *Shaanxilithes* that has been reported from the terminal Ediacaran strata in South China and India ([Hua et al., 2004](#); [Shen et al., 2007](#); [Meyer et al., 2012](#); [Brasier et al., 2013](#); [Gámez Vintaned and Zhuravlev, 2013](#); [Mángano and Buatois, 2014](#); [Tarhan et al., 2014](#)). The uppermost Khatyspyt Formation is characterized by soft-sediment slump and deformation features (**Fig. 2K**) and gutter cast structures (**Fig. 2L**). Sedimentological observations suggest deposition within a starved intracratonic rift developed in a marine ramp setting, with intraclastic limestone representing debris flows deposited adjacent to inferred syn-sedimentary faults ([Knoll et al., 1995](#); [Pelechaty et al., 1996a](#); [Nagovitsin et al., 2015](#)).

The Khatyspyt Formation overlies pebble-bearing trough and tabular cross-bedded channelized sandstone, microbial dolostone and finely laminated shale of the Maastakh Formation, and is overlain by microbial dolostone, cross-bedded oolitic and pisolitic dolostone, microbialite bioherms of the Turkut Formation (**Fig. 3**). These three formations constitute the Khorbosuonka Group, which is bounded by regional unconformities ([Sokolov and Fedonkin, 1984](#); [Knoll et al., 1995](#); [Pelechaty et al., 1996a](#); [Nagovitsin et al., 2015](#)).

2.2. Geochronological constraints

Available geochronometric constraints for the Khorbusuonka Group and the overlying Kessyusa Group are currently based on two ages. A diatreme that penetrates the Turkut and the lowermost Syhargalakh formations in a section along the Khorbusuonka River yield a U–Pb zircon age of 543.9 ± 0.24 Ma ([Bowring et al., 1993](#); [Rogov et al., 2015](#)). This age provides a minimum age constraint on the Khatyspyt Formation and a maximum age constraint on the first appearance of the trace fossil *Treptichnus pedom* in the overlying strata of the uppermost Syhargalakh Formation. A U–Pb zircon age of 529.56 ± 0.24 Ma has recently been reported for a volcanic tuff in the Mattaia Formation (Kessyusa Group) higher in the sequence ([Kaufman et al., 2012](#)) (**Fig.3**). These two ages are consistent with an inferred terminal Ediacaran age for the studied Khatyspyt Formation.

2.3. Biostratigraphy

The Khatyspyt Formation is exceptionally fossiliferous, with multiple taphonomic windows into the Earth’s earliest macroscopic communities. The fossil assemblages contain soft-bodied organisms (Avalon-type biota) with moldic preservation through authigenic carbonate cementation, carbonaceous compression macrofossils (Miaohe-type biota) ([Grazhdankin et al., 2008](#); [Nagovitsin et al., 2015](#)), and the enigmatic fossil *Nenoxites* ([Rogov et al., 2012](#)). The fossils are unevenly distributed in the Khatyspyt Formation. Apart from sporadic occurrences of *Nenoxites*, rare low-relief discoidal structures ([microbial colonies sensu Grazhdankin and Gerdes, 2007](#)), and a single small individual resembling *Khatyspytia grandis*, the lower part of the formation (i.e., the interval of 0–67.7 m from the base) is essentially unfossiliferous. The interval from 15 m to 40 m was not sampled because of poor exposure, resulting in an incomplete

lithostratigraphic column of the lower Khatyspyt Formation. Regardless, a cursory inspection of this interval suggests that it is non-fossiliferous, although future investigation of better exposures is needed to verify this observation.

Diverse Avalon-type fossils, including the globally distributed rangeomorph taxon *Charnia masoni*, “*Hiemalora*”-type holdfast structures ([sensu Hofmann et al., 2008](#)), and the arboreomorph *Khatyspytia grandis* showing similarities with *Charniodiscus procerus* from Newfoundland ([sensu Xiao and Laflamme, 2009](#)), appear suddenly in finely laminated limestones of the Khatyspyt Formation (at ~42 m). There is no visible evidence of any significant change in depositional rate or facies at this stratigraphic level. Avalon-type fossils are found throughout the rest of the Khatyspyt Formation with little variation in taxonomic composition. *Nenoxites* fossils also become increasingly abundant in the upper part of the Khatyspyt Formation. Abundant and diverse carbonaceous compression macrofossils, including some Miaohu-type taxa ([Xiao et al., 2002](#)), first appear in the mid-Khatyspyt Formation. The appearance of carbonaceous compressions is not accompanied by notable isotopic perturbation, thus may be related to taphonomic factors ([Grazhdankin et al., 2008](#)).

The paleontological characteristics of the Khatyspyt Formation is not particularly indicative of specific Ediacaran stratigraphic age. Although macrofossil assemblages consisting of palaeopascichnids, rangeomorph, erniettomorph, and/or arboreomorph frondose taxa in the absence of many other iconic Ediacaran forms usually (but not always) suggest a late Ediacaran age ([Kolesnikov et al., 2015](#); [Boag et al., 2016](#); [Darroch et al., 2016](#)). However, it should also be taken with caution that ecological control of Ediacaran organisms may have imposed serious limitation in the use of biostratigraphic correlation ([Grazhdankin, 2004](#); [Gehling and Droser, 2013](#); [Grazhdankin, 2014](#)). Supposing the *Nenoxites* ichnofabric ([sensu Rogov et al., 2012](#)) and

the body fossil *Shaanxilithes* refer to approximately the same structure, then the latter could be used as a reliable biostratigraphic marker for global correlation of terminal Ediacaran strata immediately preceding the oldest small skeletal fossils ([Shen et al., 2007](#); [Zhuravlev et al., 2009](#); [Meyer et al., 2012](#); [Tarhan et al., 2014](#); [Rogov et al., 2015](#); [Darroch et al., 2016](#); [Xiao et al., 2016](#)). Indeed, the lowest stratigraphic occurrence of the small skeletal fossil *Cambrotubulus decurvatus* in the northwestern slope of the Olenek Uplift is found to be 1.4 m above the base of the Turkut Formation that conformably overlies the Khatyspyt Formation.

3. Samples and analytical methods

The samples were collected from the Khatyspyt Formation at the 0601 (GPS: 71° 08' 28.80" N, 123° 52' 23.97" E) and 0605 (GPS: 71° 12' 17.20" N, 123° 39' 35.43" E) sections. The 0601 section is of particular interest because it has yielded an exceptionally preserved assemblage of diverse Ediacara-type macrofossils, representing the first appearance of these fossils in the Khatyspyt Formation (**Fig. 3**). Using regionally consistent occurrence of thick-bedded limestones and volcanic tuffs as marker beds, we correlated the 0605 section with a covered interval (90–106 m) at the 0601 section in order to achieve more complete chemostratigraphic sampling (**Figs. 4, 5**). The sampled interval in this study covers the lowest occurrence of Ediacara-type macrofossils, and is composed of thin-bedded carbonates deposited in normal marine environments.

A total of 176 samples were systematically analyzed to obtain a complete suite of chemostratigraphic data, including carbonate carbon ($\delta^{13}\text{C}_{\text{carb}}$) and oxygen isotopes ($\delta^{18}\text{O}_{\text{carb}}$), strontium isotopes ($^{87}\text{Sr}/^{86}\text{Sr}$), organic carbon isotopes ($\delta^{13}\text{C}_{\text{org}}$), sulfur isotopes of total sulfur in acidified residuals ($\delta^{34}\text{S}_{\text{TS}}$, comprising dominantly pyrite S with trace amount of organic S),

carbonate-associated sulfate sulfur isotopes ($\delta^{34}\text{S}_{\text{CAS}}$), and major and trace element concentrations. Sample preparation work, including sample cutting, crushing, acidification, leaching, and polishing, was conducted at the Trofimuk Institute of Petroleum Geology and Geophysics, Siberian Branch of the Russian Academy of Sciences, Novosibirsk. Elemental analyses were conducted in the Carnegie Institution of Washington. All the isotope analyses were conducted in the Department of Geology, University of Maryland by using standard methods ([e.g., McFadden et al., 2008](#); [Cui, 2015](#); [Cui et al., 2015](#); [Cao et al., 2016](#); [Cui et al., 2016a](#); [Cui et al., 2016b](#); [Cui et al., under review](#)), which are briefly outlined below.

3.1. Carbon and oxygen isotope analysis

Rock samples were cut and polished for detailed petrographic observation and micro-drilling in order to obtain powders from the least-altered, least-recrystallized, and purest phases for carbonate carbon and oxygen isotope analysis. The powders were measured with a Multicarb inlet device in-line with an Elementar Isoprime continuous-flow isotope ratio mass spectrometer, and precision for both isotopes was routinely better than 0.1‰.

3.2. Organic carbon and pyrite sulfur isotope analyses

The organic carbon, total sulfur isotope compositions were measured by combustion of the decalcified residuals to CO_2 or SO_2 with a Eurovector elemental analyzer in-line with a second Elementar Isoprime isotope ratio mass spectrometer. Around 15 g of bulk crushed sample was acidified with 3 M HCl. Acidified residues were washed with ultra-pure Milli-Q (18 M Ω) water, centrifuged, decanted, and dried. The residues were packed into folded tin cups with V_2O_5 for combustion to CO_2 or SO_2 in a Eurovector elemental analyzer in-line with a second Elementar Isoprime isotope ratio mass spectrometer, which measured isotope abundances.

Uncertainties for carbon and sulfur isotope measurements determined by multiple analyses of standard materials during analytical sessions are better than 0.1‰ and 0.3‰, respectively.

3.3. Sulfate sulfur isotope analysis

Approximately 100 g of bulk carbonate powders were used for extraction of carbonate-associated sulfate (CAS). To minimize the contamination of soluble non-CAS sulfate ([Marengo et al., 2008](#); [Wotte et al., 2012](#); [Peng et al., 2014](#); [Schobben et al., 2015](#)), bulk powders were leached by 10% NaCl until no sulfate could be tested from the leachate, and then washed with Milli-Q water for at least 3 times prior to acidification of the leached powders with 3 M HCl to release CAS. Both non-CAS and CAS precipitates were collected as BaSO₄ three days after BaCl₂ was added to the solution for $\delta^{34}\text{S}_{\text{leachate}}$ and $\delta^{34}\text{S}_{\text{CAS}}$ analyses, respectively. The BaSO₄ precipitates were packed into folded tin cups with V₂O₅ for combustion to SO₂ in a Eurovector elemental analyzer in-line with a second Elementar Isoprime isotope ratio mass spectrometer. Uncertainties for sulfur isotope measurements determined by multiple analyses of standard materials during analytical sessions are better than 0.3‰. Sulfur isotope fractionation between sulfide and sulfate are calculated as $\Delta\delta^{34}\text{S} \approx \delta^{34}\text{S}_{\text{CAS}} - \delta^{34}\text{S}_{\text{TS}}$, assuming the sulfur species in the acidified results of our pyrite-rich samples are dominated by pyrite (i.e., $\delta^{34}\text{S}_{\text{TS}} \approx \delta^{34}\text{S}_{\text{pyrite}}$).

3.4. Strontium isotope analyses

For strontium isotope analysis, only limestone samples were selected for extraction and measurement. Micro-drilled powders (ca. 10 mg) were leached three times in 0.2 M ammonium acetate (pH ~ 8.2) to remove exchangeable Sr from non-carbonate minerals, and then rinsed three times with Milli-Q water. The leached powder was centrifuged, decanted, and acidified with doubly distilled 0.5 M acetic acid overnight to release strontium from the carbonate crystal

structure. The supernatant was centrifuged to remove insoluble residues and then decanted, dried, and subsequently dissolved in 200 μl of 3 M HNO_3 . Strontium separation was carried out via cation exchange using a small polyethylene column containing ~ 1 cm of Eichrom Sr specific resin. The column was rinsed with 400 μl of 3 M HNO_3 before the dissolved sample was loaded onto the column. After loading, the sample was sequentially eluted with 200 μl of 3 M HNO_3 , 600 μl of 7 M HNO_3 , and 100 μl of 3 M HNO_3 to remove the Ca, Rb and REE fractions; the Sr fraction adsorbs strongly to the resin in an acidic environment. The Sr fraction was removed by elution with ~ 800 μl of 0.05 M HNO_3 and the resultant eluate collected and dried. Approximately 200–300 ng of the dried sample was transferred onto a degassed and pre-baked (~ 4.2 A under high vacuum) high purity Re filament with 0.7 μl of Ta_2O_5 activator. The prepared filaments were measured using the VG Sector 54 thermal ionization mass spectrometer (TIMS) facility of the University of Maryland Geochemistry Laboratories. Filaments were transferred to a sample carousel, heated under vacuum ($\sim 10^{-7}$ to 10^{-8} atm) to a temperature between 1450 $^\circ\text{C}$ and 1650 $^\circ\text{C}$, and analyzed when a stable signal (>1.0 V) was detected on the mass 88 ion beam. Approximately 100 $^{87}\text{Sr}/^{86}\text{Sr}$ ratios were collected for each sample. Final data have been corrected for fractionation using the standard value $^{87}\text{Sr}/^{86}\text{Sr} = 0.1194$. The fraction of ^{87}Sr resulting from in situ decay from ^{87}Rb was removed by measurement of rubidium abundance at mass 85. Repeated analysis of the NBS SRM987 standard yields an average value of $^{87}\text{Sr}/^{86}\text{Sr} = 0.710245 \pm 0.000011$ (2σ) during the analytical window.

3.5. Elemental analyses

Based on petrographic observations, carbonate samples with low detritus contents were selected for major and trace element analysis. Aliquots of micro-drilled carbonate powders were

dissolved in 0.4 M HNO₃, centrifuged, and then analyzed for elemental concentrations. The resulting solutions were analyzed on a Thermo Scientific iCAP-Q ICP-MS (Inductively Coupled Plasma – Mass Spectrometry) at the Carnegie Institution of Washington. Precision of these analyses as determined by repeated measurements of a house standard carbonate was < 5% (2σ) for major elements with high concentrations and < 10% (2σ) for the REEs. Cerium anomaly values are calculated by using the formula $Ce/Ce^* = [Ce]_{PAAS} / ([Pr]^2_{PAAS} / [Nd]_{PAAS})$ in order to further constrain the redox conditions of the depositional environment (e.g., [Lawrence et al., 2006](#); [Ling et al., 2013](#); [Cui et al., 2015](#); [Cui et al., 2016a](#)).

4. Geochemical results

4.1. $\delta^{13}C_{carb}$ and $\delta^{18}O_{carb}$ data

Chemostratigraphic profiles of the Khatyspyt Formation reveal strong fluctuations in carbon and sulfur isotope compositions (**Figs. 4, 5**). The percent carbonate values of most limestone samples are high (approaching 100%) except the shaley interval in the 0605 section (**Fig. 4A**). Carbonate carbon isotope ($\delta^{13}C_{carb}$) data show a positive excursion (up to ca. 5‰) in the intraclastic limestone interval of the 0601B section, and then decrease to lower values (**Fig. 4B**). The lower 45 meters of the measured section generally reveals a $\delta^{13}C_{carb}$ negative excursion and is succeeded by more positive values up section. Carbonate oxygen isotopes ($\delta^{18}O_{carb}$) mostly remain – 5‰, except a slight positive excursion to ca. 0‰ in the lower 45 meters (**Fig. 4C**).

4.2. $\delta^{13}C_{org}$ and TOC data

Total organic carbon content data in the Khatyspyt Formation (**Fig. 4F**) are generally high ($0.28\% \pm 0.4$, 1σ) compared with time-equivalent Gaojiashan Member in South China ($0.08\% \pm 0.15$, 1σ). Organic carbon isotope values ($\delta^{13}C_{org}$) are mostly between -30‰ and -40‰ (**Fig. 4D**), providing a relatively large total carbon isotope fractionation ($\Delta\delta^{13}C$) (**Fig. 4E**) compared with some older Ediacaran successions ([McFadden et al., 2008](#); [Sansjofre et al., 2011](#); [Cui et al., 2015](#)) or the Phanerozoic record ([Hayes et al., 1999](#); [Oehlert et al., 2012](#); [Oehlert and Swart, 2014](#)).

4.3. $\delta^{34}S_{py}$ and $\delta^{34}S_{CAS}$ data

A dramatic increase in pyrite sulfur isotopic compositions ($\delta^{34}S_{py}$) from ca. -20‰ to as high as ca. 50‰ occurs in the lower Khatyspyt Formation in the 0601 section (**Fig. 5B**). Such a large magnitude of $\delta^{34}S_{py}$ excursion has never been reported in the terminal Ediacaran Period (see equivalent chemostratigraphic records in [Fike and Grotzinger, 2008](#); [Ries et al., 2009](#); [Cui et al., 2016a](#)) (**Fig. 7**). Two stages could be defined to describe this dramatic $\delta^{34}S_{py}$ excursion. The lower 45 meters (Stage 1), sampled at section 0601B and lower part of section 0601C, is characterized by negative $\delta^{34}S_{py}$ values and an increase from -20‰ to ca. 0‰ . The next 85 meters of the measured Khatyspyt Formation (Stage 2) is characterized by positive $\delta^{34}S_{py}$ values with an average value of $31.7\text{‰} \pm 10.1\text{‰}$ (1σ) and maximum value as high as 55‰ . In contrast with this dramatic variation in $\delta^{34}S_{py}$, sulfur isotope values ($\delta^{34}S_{CAS}$) of carbonate-associated sulfate (CAS) are less variable, with an average value of $+40.7\text{‰} \pm 6.5\text{‰}$ (1σ). Assuming the sulfur species in the acidified residuals of our samples are dominated by pyrite (e.g., [McFadden et al., 2008](#); [Zhelezinskaia et al., 2014](#); [Cui et al., 2016a](#)), calculated sulfur isotope differences

($\Delta\delta^{34}\text{S}$) between sulfate and pyrite are large in the lower 45 meters (up to 50‰), but decline to lower and even negative values up section (**Fig. 5C**). Coincident with the large positive excursion in $\delta^{34}\text{S}_{\text{py}}$ is a dramatic decrease in CAS concentration (**Fig. 5D**). CAS concentration data in the lower 50 meters are remarkably high (up to almost 2000 ppm), but drop to much lower values (ca. 100 ppm) in the upper section. Total sulfur concentration (TS) data are generally below 0.2%, with higher values in the shaley interval of the 0605 section (**Fig. 5E**). Intriguingly, the first appearance datum (FAD) of soft-bodied Ediacara-type macro-organisms coincides with the negative-to-positive shifts in both $\delta^{13}\text{C}_{\text{carb}}$ and $\delta^{34}\text{S}_{\text{py}}$ in the lower Khatyspyt Formation (**Figs. 4, 5**).

4.4. $^{87}\text{Sr}/^{86}\text{Sr}$ and elemental concentration data

$^{87}\text{Sr}/^{86}\text{Sr}$ values in the Khatyspyt Formation (**Fig. 4H**) are ca. 0.7080 ± 0.0001 , which are consistent with published data from the same section ([Vishnevskaya et al., 2013](#)). Rb/Sr and Mn/Fe concentration ratios of the samples analyzed for $^{87}\text{Sr}/^{86}\text{Sr}$ are all consistently low, which are mostly <0.01 and <0.1 , respectively. Ce/Ce* data (**Fig. 4G**) remain steady values of 0.84 ± 0.06 (1σ) throughout the section.

5. Discussion

5.1. Diagenetic evaluation

Multiple lines of evidence (**Fig. 6**) suggest that the measured samples from the Khatyspyt Formation are well preserved, and experienced little diagenetic alteration. In contrast with the proposed diagenetic alteration trend ([Arthur, 2009](#); [Knauth and Kennedy, 2009](#); [Derry, 2010](#); [Oehlert and Swart, 2014](#)), which typically shows a positive correlation between $\delta^{13}\text{C}_{\text{carb}}$ and

$\delta^{18}\text{O}_{\text{carb}}$ resulting from meteoric water alteration and organic carbon oxidation during fluid percolation, the Khatyspyt samples shows a broadly negative correlation between $\delta^{13}\text{C}_{\text{carb}}$ and $\delta^{18}\text{O}_{\text{carb}}$ (**Fig. 6A**), suggesting little influence by meteoric water or deep burial.

It has been proposed that sulfur isotopic compositions of carbonate-associated sulfate ($\delta^{34}\text{S}_{\text{CAS}}$) could possibly be contaminated by pyrite oxidation ([Marenco et al., 2008](#); [Wotte et al., 2012](#); [Theiling and Coleman, 2015](#)) or secondary atmospheric sulfate ([Peng et al., 2014](#)). If true, cross plot between CAS content and $\delta^{34}\text{S}_{\text{CAS}}$ should follow a mixing line. However, [CAS]– $\delta^{34}\text{S}_{\text{CAS}}$ cross plot reveal little correlation (**Fig. 6F**), suggesting minimum contamination in the measured $\delta^{34}\text{S}_{\text{CAS}}$ values. Another line of evidence supporting the fidelity of our $\delta^{34}\text{S}_{\text{CAS}}$ data comes from the general consistency with comparable $\delta^{34}\text{S}$ values (ca. 40‰) of bedded anhydrite from the terminal Ediacaran Ara Group in Oman ([Fike and Grotzinger, 2008](#); [Fike and Grotzinger, 2010](#)), suggesting a global seawater signal of ca. 40‰. The consistent $\delta^{34}\text{S}_{\text{CAS}}$ values (ca. 40‰) independent of [CAS] abundances in the Khatyspyt Formation are also consistent with previous studies ([Lyons et al., 2004](#); [Gill et al., 2008](#); [Cui et al., 2016a](#)), demonstrating that CAS can track seawater sulfate compositions. We also plot concentration data against isotopic data (e.g., $\delta^{13}\text{C}_{\text{org}}$ –TOC, $\delta^{34}\text{S}_{\text{TS}}$ –TS) in order to test whether any diagenetic processes could preferentially change the isotopic compositions. These cross-plots show little correlation (**Figs. 6B, C**) as would be expected from diagenetic alteration.

In addition to $\delta^{34}\text{S}_{\text{TS}}$ and $\delta^{34}\text{S}_{\text{CAS}}$ data, sulfur isotope compositions of non-CAS from leachate solutions ($\delta^{34}\text{S}_{\text{leachate}}$) were also analyzed (**Fig. 6G**). The $\delta^{34}\text{S}_{\text{leachate}}$ data are mostly higher than $\delta^{34}\text{S}_{\text{TS}}$ in the lower 45 m of the section. In contrast, $\delta^{34}\text{S}_{\text{leachate}}$ values are mostly lower than the $\delta^{34}\text{S}_{\text{TS}}$ in the upper Khatyspyt Formation where superheavy pyrite is common. The sources of sulfate in the leachate solution could be diverse, including pyrite oxidation

([Marenco et al., 2008](#); [Wotte et al., 2012](#); [Theiling and Coleman, 2015](#)), secondary atmospheric sulfate ([Peng et al., 2014](#)), or loosely bounded secondary sulfate derived from diagenetic fluids. Thus, $\delta^{34}\text{S}_{\text{leachate}}$ values may be controlled by various geological factors at different stratigraphic intervals. Regardless, it is notable that all $\delta^{34}\text{S}_{\text{CAS}}$ data are consistently higher than $\delta^{34}\text{S}_{\text{leachate}}$ data, demonstrating the importance of leaching pretreatment before CAS extraction.

In summary, sedimentological observations and geochemical data suggest that the limestones in the studied Khatyspyt Formation are well preserved, and provide reliable geochemical signals for chemostratigraphic interpretation.

5.2. Redox constraints for the depositional basin

The dominance of carbonate lithology and the presence of Ediacara-type fossils in the Khatyspyt Formation provide important geobiological links between early macro-organisms and their marine environments ([Xiao et al., 2013](#)). However, because the most widely used tools for redox reconstruction of marine sediments—including iron speciation analysis (e.g., [Canfield et al., 2007](#); [Canfield et al., 2008](#); [Poulton and Canfield, 2011](#)) and redox-sensitive trace element analysis (e.g., [Scott et al., 2008](#); [Sahoo et al., 2012](#); [Li et al., 2015b](#); [Sahoo et al., 2016](#))—are best calibrated for shale lithofacies (but see [Clarkson et al., 2014](#); [Wood et al., 2015](#)), constraints on the redox conditions of the Khatyspyt Formation need to be sought elsewhere.

Given the intimate link between cerium abundance in carbonates and seawater/porewater redox conditions (e.g., [Zhou et al., 2012](#); [Ling et al., 2013](#)), cerium anomaly $\text{Ce}/\text{Ce}^* = [\text{Ce}]_{\text{PAAS}} / ([\text{Pr}]_{\text{PAAS}}^2 / [\text{Nd}]_{\text{PAAS}})$ has been explored to constrain the redox conditions of the depositional environment. Cerium exists in two redox states in marine environments, namely Ce(III) and Ce(IV). Trivalent Ce can be oxidized into tetravalent Ce in oxic environments ([Tanaka et al.,](#)

[2010](#)), and then scavenged by Mn-oxides and hydroxides and removed from the seawater. Thus, in oxygenated seawaters, tetravalent Ce is preferentially removed from the system, leaving the seawaters with a Ce negative anomaly in REE patterns. In contrast, trivalent Ce is more soluble in reduced environments, resulting in relatively high Ce/Ce* values ([Bau et al., 1996](#)). Therefore, Ce/Ce* ratios of carbonates may help to trace redox changes in seawater/porewater if the samples have not yet experienced significant diagenetic alteration. Here in the Khatyspyt Formation, Ce/Ce* values of well-preserved limestone samples are consistently high throughout the section, with an average value of 0.84 ± 0.06 (1σ), suggesting possible anoxic conditions.

Supporting evidence for an overall anoxic condition of the Khatyspyt Formation also comes from sedimentological observations in the field, where thinly bedded limestone are mostly laminated and remarkable rich in organic matter, including bitumen. Total organic carbon (TOC) abundances in most of the carbonate samples are high (0.28% in average) compared with the correlative Gaojiashan Member (0.08% in average), and $\delta^{13}\text{C}_{\text{org}}$ values range from -24.1‰ to -38.9‰ and average at -33.2‰ , suggesting the possible presence of chemoautotrophic or methanotrophic communities at the oxic-anoxic interface in the water column or sediments ([Hayes, 1993](#); [Brocks et al., 2005](#); [Johnston et al., 2009](#); [Jiang et al., 2012](#); [Houghton et al., 2014](#); [Guo et al., 2015](#)).

Sulfur isotope data and sulfur abundance data have also been used to constrain the redox conditions of the Khatyspyt Formation. In the lower 45 meters of the studied interval, negative $\delta^{34}\text{S}_{\text{py}}$ values (as low as -20‰) and remarkably high sulfate contents ([CAS] up to 1969 ppm) suggest strong euxinic conditions and syngenetic pyrite formation in the water column (e.g., [Canfield and Teske, 1996](#); [Li et al., 2010](#); [Cui et al., 2016a](#); [Kah et al., 2016](#)); whereas in the upper 85 meters of the measured section, relatively lower [CAS] concentrations and anomalously

high $\delta^{34}\text{S}_{\text{py}}$ values suggest non-euxinic conditions in the water column and authigenic pyrite precipitation within marine sediments in an environment with relatively low sulfate concentration (e.g., [Ries et al., 2009](#); [Fike et al., 2015](#); [Wu et al., 2016](#)), although the marine sulfate reservoir was large enough to be isotopically buffered, as suggested by the relatively invariant $\delta^{34}\text{S}_{\text{CAS}}$ values.

Collectively, integrated sedimentological and geochemical data suggest that the Khatyspyt Formation records a transition from euxinic conditions in the lower 45 meters to non-euxinic conditions in the upper 85 meters.

5.3. Stratigraphic correlation at a global scale

In order to investigate the effect of redox conditions on early metazoans at a global scale, four upper Ediacaran successions are considered for integrated stratigraphic correlation: the Ara Group in Oman, the Nama Group in Namibia, the Gaojiashan/Shibantan Member in South China, and the Khatyspyt Formation (**Fig. 7**). The late Ediacaran age of these sections is supported by biostratigraphic data and corroborated by available U–Pb zircon dates from volcanic tuffs ([Bowring et al., 1993](#); [Grotzinger et al., 1995](#); [Amthor et al., 2003](#); [Condon et al., 2005](#); [Halverson, 2006](#)). Reliable chemostratigraphic data in high stratigraphic resolution are all available for these four successions ([Fike and Grotzinger, 2008](#); [Ries et al., 2009](#); [Cui et al., 2016a](#)).

5.3.1. Chemostratigraphic $\delta^{13}\text{C}$ correlation

Integrated sedimentological, stratigraphic, and geochemical studies of carbonate successions of terminal Ediacaran records demonstrate the utility of carbon isotope

chemostratigraphy for intra- and inter-basinal correlations. The late Ediacaran $\delta^{13}\text{C}_{\text{carb}}$ record of northeastern Siberia reflects global variations seen elsewhere, featuring (in ascending order) a strong $\delta^{13}\text{C}_{\text{carb}}$ positive shift to values near 6‰ in the Maastakh Formation and basal Khatyspyt Formation, an intermediate interval of relatively little isotopic change with a monotonic decrease in $\delta^{13}\text{C}_{\text{carb}}$ from 2‰ to near 0‰ throughout most of the Khatyspyt Formation and Turkut formations, and a negative excursion to ca. – 4‰ just beneath the Ediacaran–Cambrian boundary ([Knoll et al., 1995](#); [Pelechaty et al., 1996a](#); [Pelechaty et al., 1996b](#)). Similar $\delta^{13}\text{C}_{\text{carb}}$ positive excursions have also been revealed in the broadly equivalent Nama Group of southern Namibia ([Kaufman et al., 1991](#); [Saylor et al., 1998](#)), the Ara Group in Oman ([Fike and Grotzinger, 2008](#)), and the Dengying Formation in South China ([Jiang et al., 2007](#); [Zhou and Xiao, 2007](#); [Wang et al., 2014](#); [Cui et al., 2016a](#)).

5.3.2. Chemostratigraphic $^{87}\text{Sr}/^{86}\text{Sr}$ correlation

Given the much longer residence time of strontium in the ocean than the seawater mixing time, strontium isotopes ($^{87}\text{Sr}/^{86}\text{Sr}$) have been widely used as a tool for inter-basinal stratigraphic correlation (e.g., [Halverson et al., 2007](#); [Cui et al., 2015](#)). Terminal Ediacaran limestones typically have $^{87}\text{Sr}/^{86}\text{Sr}$ ratios ca. 0.7084, including the fossiliferous Shibantan Member of the Dengying Formation in South China ([Jiang et al., 2007](#)), the Blueflower Formation in NW Canada ([Narbonne et al., 1994](#)), the Nama and Witvlei groups in Namibia and South Africa ([Kaufman et al., 1993](#)), the Tamengo Formation of Corumba Group in Brazil ([Gaucher et al., 2003](#); [Boggiani et al., 2010](#)), and the Itapucumi Group in NW Paraguay ([Warren et al., 2011](#)). However, $^{87}\text{Sr}/^{86}\text{Sr}$ values of the Khatyspyt Formation are consistently ca. 0.7080 ([Vishnevskaya et al., 2013 and this study](#)), raising questions about a $^{87}\text{Sr}/^{86}\text{Sr}$ chemostratigraphic correlation between the Khatyspyt Formation and other terminal Ediacaran successions worldwide.

Two possibilities may account for the relatively lower $^{87}\text{Sr}/^{86}\text{Sr}$ values of the Khatyspyt Formation compared with other roughly equivalent successions mentioned above. Largely due to inappropriate lithology (i.e., dolostone, instead of limestone) in the other sections, the 0.7084 values are mainly based on limited analyses of sections in low stratigraphic resolutions, thus may not represent a complete evolution of the $^{87}\text{Sr}/^{86}\text{Sr}$ during that time. The overall 0.7080 values of the Khatyspyt Formation thus may reflect primary seawater signals of a time interval that has not yet been captured in other terminal Ediacaran data sets before. Alternatively, it is also possible that the lower $^{87}\text{Sr}/^{86}\text{Sr}$ values in the Khatyspyt Formation may reflect a secondary signal overprinted by enhanced hydrothermal input (e.g., [Frei et al., 2011](#)). We cannot resolve these alternative interpretations with the available data.

5.3.3. Chemostratigraphic $\delta^{34}\text{S}$ correlation

Chemostratigraphic profiles of paired sulfur isotopes ($\delta^{34}\text{S}_{\text{sulfate}}$ and $\delta^{34}\text{S}_{\text{py}}$) reported from upper Ediacaran Gaojiashan Member in South China ([Cui et al., 2016a](#)), Nama Group in Namibia ([Ries et al., 2009](#)), and Ara Group in Oman ([Fike et al., 2006](#); [Fike and Grotzinger, 2008](#)) reveal both similarities and differences when compared to sulfur isotope profiles of the Khatyspyt Formation (**Figs. 7, 8**). All these sections consistently show a large positive shift in $\delta^{34}\text{S}_{\text{py}}$; however, the absolute values and the magnitude of this positive shift are different among the four successions, which may reflect a global perturbation in S cycling overprinted by local signals (e.g., [Loyd et al., 2013](#); [Wood et al., 2015](#)). The sulfate sulfur isotopes are generally steady at around 40‰ in Ara, Gaojiashan, and Khatyspyt units ([Fike and Grotzinger, 2010](#); [Cui et al., 2016a](#)). However, the Nama Group shows lower $\delta^{34}\text{S}_{\text{CAS}}$ values with large variations (27.3‰ $\pm$ 10.3‰, 1 σ) ([Ries et al., 2009](#)). It should be noted that $\delta^{34}\text{S}_{\text{sulfate}}$ in Oman was measured from bedded anhydrite, while $\delta^{34}\text{S}_{\text{sulfate}}$ data in the other three successions were measured from

extracted carbonate-associated sulfate (CAS). Thus, the $\delta^{34}\text{S}_{\text{sulfate}}$ profile in Oman is probably more reliable given the possible contamination of CAS during chemical weathering in the field ([Peng et al., 2014](#)) or lab extraction ([Marenco et al., 2008](#)). Another notable feature is the occurrence of superheavy pyrite ($\delta^{34}\text{S}_{\text{py}} > \delta^{34}\text{S}_{\text{CAS}}$) in both Nama Group and the Khatyspyt Formation (**Figs. 7, 8**). If we accept the global $\delta^{34}\text{S}_{\text{sulfate}}$ signal is ca. 40‰ as preserved in evaporates, then the comparatively lower $\delta^{34}\text{S}_{\text{CAS}}$ values [ca. 27.3‰ $\pm$ 10.3‰ (1 σ)] in the Nama Group and the anomalously high $\delta^{34}\text{S}_{\text{py}}$ values in both Nama Group and the Khatyspyt Formation need to be explained (see the next section).

The inconsistent absolute values in sulfur isotopes among different localities may result from potential alteration of $\delta^{34}\text{S}_{\text{CAS}}$ during post-deposition (e.g., [Peng et al., 2014](#)), or locally expressed $\delta^{34}\text{S}_{\text{pyrite}}$ signals due to largely heterogeneous sulfur reservoirs in the late Ediacaran ocean (cf. [Loyd et al., 2012](#)). Nevertheless, the overall increasing trend in $\delta^{34}\text{S}_{\text{pyrite}}$ and decreasing trend in $\Delta\delta^{34}\text{S}$ are consistent among the sections, suggesting an overall redox transition from euxinic conditions to a more ventilated ocean. It should be noted that these euxinic water masses may have largely been constrained within shelf basin environments ([Lyons et al., 2014](#); [Cui et al., 2015](#)) instead of an open ocean phenomenon. They may have been synchronized by a global enhancement of sulfate flux into the ocean shelf environment during an oxidation event.

5.4. Controlling factors for the chemostratigraphic $\delta^{34}\text{S}$ excursion

Multiple factors, for example, ventilation effect of bioturbation (cf. [Canfield and Farquhar, 2009](#)), sealing effect of microbial mat (e.g., [Bouougri and Porada, 2007](#); [Cai et al., 2010](#); [Grotzinger and Al-Rawahi, 2014](#)), or change in sedimentation rate (cf. [Schiffbauer et al.,](#)

[2014](#)) of marine sediments, could have potentially caused variation of $\delta^{34}\text{S}_{\text{pyrite}}$ compositions in the studied Khatyspyt Formation. Here we briefly discuss these factors below.

5.4.1. Ventilation effect of bioturbation

It has been proposed that enhanced ventilation of marine sediments and oxidation of authigenic pyrite could be triggered by bioturbation, which consequently would increase the sulfate concentration in the seawater ([Canfield and Farquhar, 2009](#)). The activity of bioturbation is generally low during the late Ediacaran Period ([Tarhan and Droser, 2014](#); [Tarhan et al., 2015](#)). Here we tentatively evaluate the intensity of bioturbation via the abundance of the enigmatic *Nenoxites* fossils, though its origin still remains debatable ([Rogov et al., 2012](#); [Brasier et al., 2013](#); [Gámez Vintaned and Zhuravlev, 2013](#); [Rogov et al., 2013a](#); [Rogov et al., 2013b](#); [Mángano and Buatois, 2014](#); [Budd and Jensen, 2015](#); [Rogov et al., 2015](#)). If *Nenoxites* fossils indeed result from bioturbation as proposed by [Rogov et al. \(2012\)](#), then it is revealed that bioturbation appears both below and above the euxnic interval, and becomes more abundant starting from ca. 90 m of the studied section (**Figs. 4, 5**). However, these intervals with inferred evidence of bioturbation all yield lower [CAS] concentration values (**Fig. 5D**) instead of higher sulfate concentration predicted by [Canfield and Farquhar \(2009\)](#). Thus, we preclude the possibility of bioturbation as a major controlling factor for the chemostratigraphic $\delta^{34}\text{S}$ excursion.

5.4.2. Sealing effect of microbial mat

The occurrence of microbial mats could also potentially influence the $\delta^{34}\text{S}$ composition of pyrite in marine sediments. Below non-bioturbated and microbially sealed sediments where the water–sediment interface represents a significant diffusion barrier, pyrite may have higher $\delta^{34}\text{S}$ composition. This mechanism may have played a role in the time-equivalent Gaojiashan

Member in South China, where chemostratigraphic higher $\delta^{34}\text{S}_{\text{pyrite}}$ values roughly correspond to the limestone interval with abundant microbial mats (Cui et al., 2016a). However, in the studied Khatyspyt Formation, no clear evidence of microbial mats has been found throughout the section. Thus, this possibility cannot readily explain the observed chemostratigraphic $\delta^{34}\text{S}$ excursion either.

5.4.3. Change in sedimentation rate

Sedimentation rate of marine sediments has been proposed to be a controlling factor in S cycles within the pore water system (Schiffbauer et al., 2014). Higher sedimentation rate may decrease the sulfate diffusion between seawater and pore water, thereby increase the $\delta^{34}\text{S}$ composition of authigenic pyrite. However, in the studied Khatyspyt Formation, the lithological facies seem to be largely constant with no clear depositional change or discontinuity (Figs. 2, 3). Moreover, negative $\delta^{34}\text{S}$ excursion covers both channelized intraclastic limestone (presumably with relatively higher sedimentation rate) and finely laminated limestone (presumably with relatively lower sedimentation rate). Therefore, the scenario of changing sedimentation rate could not readily explain the $\delta^{34}\text{S}$ variation in the Khatyspyt Formation.

Taken together, based on the consideration above, it is unlikely that variation in the intensity of bioturbation, abundance of microbial mats, or sedimentation rate could well explain the large $\delta^{34}\text{S}$ excursion revealed in this study. In light of similar patterns of $\delta^{34}\text{S}$ profiles from roughly equivalent successions in Oman and South China (Figs. 7, 9), it appears that this event is a global phenomenon and directly associated with the formation of distinct superheavy pyrite (see the next section). Thus, the interpretation of the $\delta^{34}\text{S}$ profiles should not only explain the

overall positive shift in $\delta^{34}\text{S}_{\text{pyrite}}$, but also reconcile the remarkably ^{34}S -enriched pyrite in the upper section.

5.5. Biogeochemical origin of the superheavy pyrite

Paired sulfur isotopes of pyrite and sulfate from the Khatyspyt Formation are strongly decoupled. In the measured section, CAS sulfur isotope values are consistently around $40.7\text{‰} \pm 6.5\text{‰}$ (1σ), which are similar to $\delta^{34}\text{S}_{\text{CAS}}$ data from well-preserved limestone samples in the broadly equivalent Gaojiahsan Member of the Dengying Formation in South China (Cui et al., 2016a) and bedded anhydrite in the Ara Group in Oman (Fike and Grotzinger, 2008) (Figs. 5, 7). However, sulfur isotope compositions of total sulfur ($\delta^{34}\text{S}_{\text{TS}}$) in the acidified residuals of the Khatyspyt Formation reveal a large positive shift (Fig. 5B). Assuming sulfur in acidified residuals of these samples is dominantly pyrite (e.g., McFadden et al., 2008; Zhelezinskaia et al., 2014; Cui et al., 2015), calculated sulfur isotope differences between sulfate and sulfide $\Delta\delta^{34}\text{S} = \delta^{34}\text{S}_{\text{CAS}} - \delta^{34}\text{S}_{\text{TS}}$ are large in the lower Khatyspyt Formation (ca. 40–50‰), transitioning to negative values in the middle section and then to ca. 20–30‰ in the upper section (Fig. 5C). It should be noted, however, that the lack of negative $\Delta\delta^{34}\text{S}$ values in the upper Khatyspyt Formation may reflect the limited number of $\delta^{34}\text{S}_{\text{CAS}}$ analyses. If the global sulfate $\delta^{34}\text{S}$ compositions during the terminal Ediacaran Period are 40‰ as recorded in bedded anhydrite in Oman (Fike and Grotzinger, 2008; Fike and Grotzinger, 2010), a number of samples in the upper 50 meters of the Khatyspyt Formation would give negative $\Delta\delta^{34}\text{S}$ values.

5.5.1. Published models for superheavy pyrite

Superheavy pyrite and reverse fractionation ($\delta^{34}\text{S}_{\text{CAS}} < \delta^{34}\text{S}_{\text{py}}$) have also been reported from the time-equivalent Nama Group in Namibia (Ries et al., 2009) and basal Ediacaran cap

carbonates in northwestern China ([Shen et al., 2008](#)). Traditional sulfur isotope models have difficulty explaining such phenomenon. Some workers have attributed the reverse fractionation to decoupling of the surface- and bottom-water sulfur reservoirs through ocean stratification ([Shen et al., 2008](#); [Kah et al., 2016](#)). This leads to Rayleigh distillation and the formation of ^{34}S -enriched bottom waters, from which sulfate for pyrite precipitation is sourced. Similarly, CAS formed in surface waters could have lower $\delta^{34}\text{S}$ values resulting from input from oxidative weathering of terrestrial pyrite (e.g., [Shen et al., 2008](#)) or oxidation of deeper euxinic water mass (e.g., [Thompson and Kah, 2012](#); [Kah et al., 2016](#)). However, this mechanism would not work in the Khatyspyt Formation, which shows cross beddings and intraclasts (**Fig. 2**) and is interpreted to be deposited in a generally shallow environment in a marine ramp setting.

Experimental studies of microbial cultures have shown that the degree of sulfur isotope fractionation accompanying bacterial sulfate reduction (BSR) in seawater approaches to zero when $[\text{SO}_4^{2-}]$ is less than 200 μM ([Habicht et al., 2002](#)). This may partly explain the large ^{34}S enrichment of pyrite observed in the Khatyspyt Formation and may reflect low seawater $[\text{SO}_4^{2-}]$ in the Proterozoic time. Additionally, the inferred non-euxinic conditions in the upper Khatyspyt Formation means that pyrite formation likely occurred within porewaters, and sulfate availability for BSR may have been limited by diffusion, leading to relatively small sulfur isotope fractionations. Despite this, low seawater sulfate concentration or availability by itself does not adequately explain the anomalous superheavy pyrite values, because microbial culture studies show that sulfur isotope fractionation during BSR approaches zero but does not invert as seawater sulfate declines. Thus, additional mechanism is required.

Abiotic oxidation of dissolved sulfide during sediment reworking has recently been invoked to explain the reversed S isotope fractionation or positive S isotope variations ([Aller,](#)

2014; Fike et al., 2015). Experimental studies reveals that abiotic oxidation of dissolved sulfide could result in a fractionation of $\Delta\delta^{34}\text{S}_{\text{reactant-product}} = \delta^{34}\text{S}_{\text{sulfide}} - \delta^{34}\text{S}_{\text{sulfate}} = 5.2 \pm 1.4\text{‰}$ (Fry et al., 1988). During abiotic oxidation of sulfide, ^{32}S in sulfide is preferentially oxidized, causing the residual sulfide reservoir to be more enriched in ^{34}S . This mechanism has been proposed to explain the reversed S isotope fractionation in the terminal Ediacaran Nama Group (Ries et al., 2009) and positive $\delta^{34}\text{S}_{\text{py}}$ values in both modern marine sediments (Gao et al., 2013; Fike et al., 2015) and Triassic strata (Thomazo et al., 2016). However, the limited fractionation (ca. 5‰) associated with abiotic oxidation of sulfide does not fully explain the large $\delta^{34}\text{S}_{\text{py}}$ shift in the Khatyspyt Formation. For example, the $\delta^{34}\text{S}_{\text{py}}$ offset between shallow (as high as 20‰) and deep (as low as -35‰) drill cores of modern marine sediments reaches ca. 55‰ (Fike et al., 2015), which would require almost complete oxidation of sulfide and thus exceedingly low pyrite contents in the sediments. Similarly, the $\delta^{34}\text{S}_{\text{py}}$ variations in Triassic strata show a large positive excursion from ca. -20‰ to 40‰ (Thomazo et al., 2016), which is also too large to be adequately explained by abiotic oxidation of sulfide.

Bacterial sulfide oxidation may also contribute to the ^{34}S -enrichment of residual sulfide. In experiments with *Thiobacillus concretivorus* grown on H_2S as the sulfur source, enrichment of ^{32}S has been detected in the product sulfate. Lab results show that sulfur isotope fractionation between initial sulfide and the sulfate product during biological oxidation of H_2S by *Thiobacillus concretivorus* ($\Delta\delta^{34}\text{S}_{\text{reactant-product}} = \delta^{34}\text{S}_{\text{sulfide}} - \delta^{34}\text{S}_{\text{sulfate}}$) can be as much as 9‰ (Kaplan and Rafter, 1958) and up to 18‰ (Kaplan and Rittenberg, 1964). This magnitude of fractionation has not yet been confirmed in natural environment, but its potential role in generating superheavy pyrite should be further investigated in future research.

5.5.2. A synthetic model

In light of increasingly higher TOC content and possible anoxia conditions in the upper Khatyspyt section, it is likely that the large $\delta^{34}\text{S}_{\text{pyrite}}$ positive shift may reflect enhanced reaction rates of microbial sulfate reduction (MSR) and pyrite formation. Environmentally controlled experiments suggest that MSR-related fractionation is related to sulfate reduction rates that are dependent on the availability of organic substrates as electron donors ([Canfield et al., 2010](#); [Leavitt et al., 2013](#); [Wing and Halevy, 2014](#); [Gomes and Hurtgen, 2015](#)), and the degree of S isotope fractionation is inversely proportional to the MSR rate ([Harrison and Thode, 1958](#); [Kaplan and Rittenberg, 1964](#)). This mechanism has been well demonstrated in modern marine sediments, particularly in the sulfate–methane transition zone where sufficient methane flux could significantly accelerate the reaction rates of microbial sulfate reduction ([Borowski et al., 1996](#); [Borowski et al., 2000](#); [Jørgensen et al., 2004](#); [Borowski et al., 2013](#); [Lin et al., 2016a](#); [Lin et al., 2016b](#)).

The occurrences of superheavy pyrite are not limited to the Ediacaran System; in fact, they have also been reported from Tonian-Cryogenian ([Xu et al., 1990](#); [Li et al., 1999](#); [Liu et al., 2006](#); [Guilbaud et al., 2015](#)), Cambrian ([Gill et al., 2011](#)), and Ordovician strata ([Chen et al., 2013](#); [Sim et al., 2015](#); [Kah et al., 2016](#)). However, superheavy pyrite is remarkably common in the Ediacaran System ([Canfield, 2001](#); [Prokoph et al., 2008](#); [Canfield and Farquhar, 2012](#)) (**Figs. 8, 9**), possibly indicating unusual sulfur cycling and redox conditions in a strongly stratified ocean at the dawn of metazoan diversification ([Logan et al., 1995](#)). In the context of a progressively oxidized atmosphere ([Shields-Zhou and Och, 2011](#); [Canfield, 2014](#)), enhanced oxidative weathering could have significantly enhanced the continental weathering flux, bring nutrients and sulfate into the terminal Ediacaran ocean, promoting primary production and water

column bacterial sulfate reduction ([Horton, 2015](#); [Cui et al., 2016a](#); [Cui et al., 2016b](#)). Such a scenario is also consistent with the multiple S isotope study ([Wu et al., 2014](#)) and progressive increase in the global seawater $^{87}\text{Sr}/^{86}\text{Sr}$ ratios during the late Neoproterozoic ([Halverson et al., 2007](#); [Shields, 2007](#); [McArthur et al., 2012](#); [Xiao et al., 2016](#)). Consistent with this view, largely based on paired S analysis of the time-equivalent Nafun and Ara groups, it was proposed that an increase in pyrite burial, most likely driven by enhanced primary production and sequestration of organic carbon, led to the large paired S isotope anomaly (named Ara anomaly) in the terminal Ediacaran Period ([Fike and Grotzinger, 2008](#)), which is notably coupled with earlier reports of elevated oxidation of the atmosphere ([Kaufman et al., 2007](#)), enhanced organic carbon burial ([Hayes et al., 1999](#)), and widespread phosphorite deposition ([Cook and Shergold, 1984](#); [Papineau, 2010](#); [Muscente et al., 2015](#); [Cui et al., 2016b](#)).

5.6. Oceanic redox conditions and fossil distribution

Integrated litho-, bio- and chemo-stratigraphy allows us to uncover possible link between the distribution of early macro-organisms and their paleo-environmental context. The Khatyspyt Formation reveals an intriguing relationship between the first local appearance of Ediacara-type macrofossils and a positive shift in both carbonate carbon and pyrite sulfur isotopes at ca. 42 m in stratigraphy (**Fig. 5**). The occurrence of Ediacaran fossils in the non-euxinic intervals in the Khatyspyt Formation the Gaojiashan Member indicates that this overall isotopic-paleontological coupling relationship may reflect a biological response of early macro-organisms to more habitable redox conditions with less hydrogen sulfide in the water column (**Fig. 10**).

The poisonous nature of dissolved hydrogen sulfide in the water column would have been a source of stress for Ediacaran complex eukaryotes ([Anbar and Knoll, 2002](#)), and would have

been a major control on where communities would have been able to establish themselves in the Ediacaran oceans (**Fig. 10A**). As euxinic conditions in the lower Khatyspyt Formation are replaced by non-euxinic and more habitable conditions in the upper Khatyspyt Formation, macrofossils particularly Ediacara-type fossils become more abundant in the upper 85 meters of the studied interval (**Fig. 10B**). Although the atmospheric and oceanic oxygen contents remained below modern levels until the early Paleozoic ([Kah and Bartley, 2011](#); [Chen et al., 2015](#)), they may have already crossed the threshold to support animal metabolisms and activities ([Knoll and Sperling, 2014](#); [Mills and Canfield, 2014](#); [Mills et al., 2014](#); [Planavsky et al., 2014](#); [Sperling et al., 2015b](#)). Ediacara-type organisms preserved in the Khatyspyt Formation may have lived in non-euxinic (weakly oxic and perhaps ferruginous) conditions. Indeed, iron speciation studies from time-equivalent strata in NW Canada ([Johnston et al., 2013](#); [Sperling et al., 2015a](#)) and Namibia ([Wood et al., 2015](#)) also support the view that early Ediacaran animals may have lived in dynamic redox environments fluctuating between ferruginous and oxic conditions.

Oceanic euxinia may have been widespread in Mesoproterozoic oceans ([Canfield, 1998](#); [Poulton et al., 2004](#); [Scott et al., 2008](#); [Gilleaudeau and Kah, 2013](#)) and may have been common in Ediacaran continental margins due to enhanced oxidative weathering (e.g., [Li et al., 2010](#); [Wang et al., 2012](#); [Li et al., 2015a](#); [Li et al., 2015b](#); [Och et al., 2015](#); [Sahoo et al., 2016](#)).

Whether early macro-organisms evolved the capability to detoxify water-column sulfide still remains unknown. Data from this study, however, indicate that the local appearance of Ediacara-type macrofossils in the Khatyspyt Formation does not support the suggestion that early metazoans were capable of tolerating euxinic conditions. If ocean euxinia had a lethal effect on early metazoans, more work is needed to constrain the effect of spatiotemporal variation of

euxinic water masses on the distribution of early metazoans (e.g., [Li et al., 2010](#); [Cui et al., 2015](#); [Li et al., 2015b](#); [Och et al., 2015](#)).

6. Conclusions

Based on integrated litho-, bio-, and chemo-stratigraphy of the terminal Ediacaran Khatyspyt Formation in Arctic Siberia, a large positive $\delta^{34}\text{S}_{\text{py}}$ excursion with a magnitude of ca. 70‰ is reported here for the first time in this time interval. This shift of $\delta^{34}\text{S}_{\text{py}}$ values from –20‰ to 50‰ is closely coupled with the local appearance of Ediacara-type macrofossils in the studied section, suggesting an intriguing geobiological response of early metazoans to dynamic redox conditions. Based on multiple lines of sedimentological and geochemical evidence, we proposed that euxinic conditions in the continental margin may have excluded Ediacara-type organisms and early metazoans from the lower Khatyspyt Formation. The progressive transition from euxinic to non-euxinic conditions allowed the colonization of early macro-organisms in the upper Khatyspyt Formation.

Acknowledgments

We thank Konstantin Nagovitsin and Boris Kochnev for their assistance in the field; Mike Evans, Rebecca Ohly, Courtney Ray, Rebecca E. Plummer and Yongbo Peng for lab assistance in the UMD Paleoclimate CoLaboratory; and Richard Walker, Igor Puchtel, Jingao Liu and Katherine Bermingham for the guidance on strontium isotope analysis in the UMD TIMS Laboratory. We also thank James Farquhar, Roberta Rudnick, and Xianguo Lang for helpful discussion in the course of this project. This paper benefits from constructive review from Linda Kah and an anonymous reviewer.

This research is funded by the NASA Exobiology grant (NNX12AR91G to AJK and NNX15AL27G to SX), the NSF Sedimentary Geology and Paleontology grant (EAR0844270 to AJK; EAR1528553 to SX), the Carnegie Institution of Washington Postdoctoral Fellowship to XML. Part of the stratigraphic, sedimentological and paleoecological studies of the Khatyspyt Formation was supported by the Russian Science Foundation (grant 14–17–00409 to DVG) and the Russian National Geographic Committee for Research and Exploration (grants 8227–07, 8637–09, 9031–11 to DVG). HC acknowledges the NASA Astrobiology Institute for support.

Supplementary materials

Table S1: Chemostratigraphic data of the terminal Ediacaran Khatyspyt Formation at the 0601 and 0605 sections, Arctic Siberia, Russia.

Figure S1: Integrated chemostratigraphic profiles of the Khatyspyt Formation at the 0601 and 0605 sections, Arctic Siberia, Russia.

Figure captions

Figure 1: Map showing measured sections of the Khatyspyt Formation (including the studied 0601 and 0605 sections marked as red dots) along the Khorbusuonka River in the Olenek Uplift, northeastern Siberia, Republic of Sakha, Russia.

Figure 2. Sedimentary features of the Khatyspyt Formation. **(A)** Outcrop of the Khatyspyt Formation and stratigraphic position of the studied interval; **(B)** Intraclastic limestone suggesting

a local slope break in a ramp setting; (C) Dolomitized intraclastic limestones with soft-sediment deformation structures suggesting a local slope break in a ramp setting; (D) Cross-bedded limestone in a mid-ramp setting; (E) Alternating limestones and shales in an outer ramp setting; (F) Carbonaceous compressions preserved on a bedding plane of finely laminated limestones; (G) Thin-bedded limestones in a mid-ramp setting; (H) Alternating thin-bedded and thick-bedded limestone characteristic of a distal mid ramp to proximal outer ramp zone (pencil for scale); (I) Thin-bedded limestone with debated ichnofabric in an outer to mid-ramp setting; (J) Wavy-bedded limestone in a mid-ramp setting; (K) Soft-sediment deformation structures in the interval of limestone-shale alternations suggesting a local slope break in a ramp setting; (L) Thin-bedded limestone gutter cast in an inner ramp setting. The scales in I, J and K are around 21cm, 7 cm, and 10 cm in length, respectively. Please refer to Fig. 4 for stratigraphic position of each photograph.

Figure 3. Integrated litho-, bio and chrono-stratigraphy of the Khorbusuonka and lowermost Kessyusa groups in Arctic Siberia. Controversial *Nenoxites/Shaanxilithes*-type structure occurs in the basal Khatyspyt Formation, which may represent the oldest evidence of bioturbation (Rogov et al., 2012), or enigmatic ribbon-shaped body fossils (Shen et al., 2007; Meyer et al., 2012; Tarhan et al., 2014). Typical Ediacara-type fossils were found in the middle Khatyspyt Formation (Grazhdankin et al., 2008). Small skeletal fossils were found in the overlying Turkut Formation and Syhargalakh formations (Nagovitsin et al., 2015). The Turkut Formation is penetrated by diatremes that yielded a U–Pb zircon date of 543.9 ± 0.3 Ma (Bowring et al., 1993). The small shelly fossil *Anabarites trisulcatus* was found in limestone clasts in the tuff breccia from the diatreme. The limestone clasts are inferred to have derived from the Turkut Formation, although the exact stratigraphic horizon of *Anabarites trisulcatus* has not yet been located in the

Turkut Formation. A new U–Pb zircon date of 529.56 ± 0.24 Ma has recently been reported for a volcanic tuff in the Mattaia Formation (Kessyusa Group) higher in the sequence ([Kaufman et al., 2012](#)). Both ages are measured by TIMS (Thermal Ionization Mass Spectrometer).

Figure 4. Chemostratigraphic profiles of the terminal Ediacaran Khatyspyt Formation at the 0601 and 0605 sections, Arctic Siberia. Measured geochemical profiles include carbonate percentage (carbonate%), carbonate carbon ($\delta^{13}\text{C}_{\text{carb}}$, ‰ V-PDB) and oxygen isotopes ($\delta^{18}\text{O}_{\text{carb}}$, ‰ V-PDB), organic carbon isotopes ($\delta^{13}\text{C}_{\text{org}}$, ‰ V-PDB), carbon isotope fractionations ($\Delta\delta^{13}\text{C}_{\text{carb-org}}$), total organic carbon content (TOC, %), Ce anomaly (Ce/Ce^*) calculated using the formula $\text{Ce}/\text{Ce}^* = [\text{Ce}]_{\text{PAAS}} / ([\text{Pr}]_{\text{PAAS}}^2 / [\text{Nd}]_{\text{PAAS}})$ ([Ling et al., 2013](#)), and strontium isotope ratios ($^{87}\text{Sr}/^{86}\text{Sr}$). The pink background color represents interpreted euxinic conditions. Red fitted lines represent five-point running average. Poor exposure of the interval from 15 m to 40 m resulted in a sampling gap and an incomplete lithological column in the lower Khatyspyt Formation, although a limited scope of field observation recovered no fossils in this unsampled interval. Refer to Fig. 4 for stratigraphic position of the studied interval.

Figure 5. Chemostratigraphic profiles of the terminal Ediacaran Khatyspyt Formation at the 0601 and 0605 sections, Arctic Siberia. Measured geochemical profiles include carbonate carbon isotopes ($\delta^{13}\text{C}_{\text{carb}}$, ‰ V-PDB), total sulfur isotopes in acidified residuals ($\delta^{34}\text{S}_{\text{TS}}$, ‰ V-CDT; including pyrite and trace amount of organic S), carbonate-associated sulfate sulfur isotopes ($\delta^{34}\text{S}_{\text{CAS}}$, ‰ V-CDT), sulfur isotope fractionations ($\Delta\delta^{34}\text{S}_{\text{CAS-pyrite}}$) assuming the $\delta^{34}\text{S}_{\text{TS}}$ signals are dominantly derived from pyrite, total sulfur content (TS%, dominated by pyrite with trace amount of organic S), and carbonate-associated sulfate concentration ([CAS] in ppm). The pink background color represents euxinic conditions. Fitted lines represent five-point running average for $\delta^{13}\text{C}_{\text{carb}}$ and $\delta^{34}\text{S}_{\text{TS}}$ data, and two-point running average for $\delta^{34}\text{S}_{\text{CAS}}$, $\Delta\delta^{34}\text{S}$ and [CAS] data.

Poor exposure of the interval from 15 m to 40 m resulted in a sampling gap and an incomplete lithological column in the lower Khatyspyt Formation, although a limited scope of field observation recovered no fossils in this unsampled interval. Please refer to Fig. 4 for stratigraphic position of the studied interval.

Figure 6. (A–F) Cross plots of $\delta^{13}\text{C}_{\text{carb}}-\delta^{18}\text{O}_{\text{carb}}$, $\delta^{13}\text{C}_{\text{org}}-\text{TOC}\%$, $\delta^{34}\text{S}_{\text{TS}}-\text{TS}\%$, $\text{TOC}\%-\text{TS}\%$, $\delta^{34}\text{S}_{\text{TS}}-\delta^{34}\text{S}_{\text{CAS}}$, $\delta^{34}\text{S}_{\text{CAS}}-[\text{CAS}]$. (G) Sulfur isotope data of total sulfur ($\delta^{34}\text{S}_{\text{TS}}$, red), carbonate-associated sulfate ($\delta^{34}\text{S}_{\text{CAS}}$, blue), and leachate solutions ($\delta^{34}\text{S}_{\text{leachate}}$, empty circles) collected during leaching. Note that the $\delta^{34}\text{S}_{\text{leachate}}$ data are systematically higher than $\delta^{34}\text{S}_{\text{TS}}$ in the lower 45 m, while are generally lower than the $\delta^{34}\text{S}_{\text{TS}}$ data measured from acidified residuals dominated by superheavy pyrite in the upper section. All $\delta^{34}\text{S}_{\text{CAS}}$ data are higher than $\delta^{34}\text{S}_{\text{leachate}}$ data, demonstrating the significance of leaching pretreatment before CAS extraction. All the data are measured from the studied interval in the Khatyspyt Formation at the 0601 and 0605 sections, Arctic Siberia.

Figure 7. Global comparison of carbonate C and O isotope data ($\delta^{13}\text{C}_{\text{carb}}$, $\delta^{18}\text{O}_{\text{carb}}$), paired S isotope data ($\delta^{34}\text{S}_{\text{SO}_4}$, $\delta^{34}\text{S}_{\text{py}}$), and sulfur isotope fractionation ($\Delta\delta^{34}\text{S}$) data in terminal Ediacaran successions. Data source: Siberia (this study), South China ([Cui et al., 2016a](#)), Oman ([Fike et al., 2006](#); [Fike and Grotzinger, 2008](#)), and Namibia ([Ries et al., 2009](#)). Note that Oman $\delta^{34}\text{S}_{\text{SO}_4}$ data were measured from bedded anhydrite, whereas $\delta^{34}\text{S}_{\text{SO}_4}$ data in Siberia, South China and Namibia were measured from extracted carbonate-associated sulfate (CAS). Fitted lines represent five-point running average for most of the data, and two-point running average for $\delta^{34}\text{S}_{\text{CAS}}$ and $\Delta\delta^{34}\text{S}$ data in Gaojiashan and Siberia due to relatively smaller data set. The pink background color represents putative euxinic conditions. Biomineralizing animals have been found from sections in Oman ([Amthor et al., 2003](#)), Namibia ([Germs, 1972](#)) and South China

([Bengtson and Zhao, 1992](#)). Frondose Ediacara fossils have been found from sections in Siberia ([Grazhdankin et al., 2008](#)) and Namibia ([Grotzinger et al., 1995](#)). Similar fossils have also been found in the Shibantan Member in the Yangtze Gorges area ([Ding and Chen, 1981](#); [Sun, 1986](#); [Chen et al., 2014](#)), which is generally correlated with the Gaojiashan Member in southern Shaanxi Province of South China. Note that the enigmatic fossil *Nenoxites/Shaanxilithes* also appears to be restricted to non-euxinic intervals in both the Gaojiashan Member ([Shen et al., 2007](#); [Meyer et al., 2012](#)) and the Khatyspyt Formation ([Rogov et al., 2012](#)).

Figure 8. Histograms showing paired sulfur isotope data ($\delta^{34}\text{S}_{\text{pyrite}}$ in red, $\delta^{34}\text{S}_{\text{sulfate}}$ in blue) from the Ara Group in Oman ([Fike and Grotzinger, 2008](#)), the Gaojiashan Member in South China ([Cui et al., 2016a](#)), the Khatyspyt Formation in Arctic Siberia (this study), and the Nama Group in Namibia ([Ries et al., 2009](#)). Note that the $\delta^{34}\text{S}_{\text{pyrite}}$ data in the Khatyspyt Formation and the Nama Group are anomalously high compared with the other two sections.

Figure 9. Evaporite, carbonate-associated sulfate (CAS), and pyrite sulfur isotope data through Earth history. Paired $\delta^{34}\text{S}$ data are compiled from the literature [after ([Canfield and Farquhar, 2009](#); [Och and Shields-Zhou, 2012](#); [Sahoo et al., 2012](#); [Cui et al., 2016a](#))]. The small panels are frequency distribution of paired sulfur isotopes of the Phanerozoic (541–0 Ma), Ediacaran (635–541 Ma), Tonian–Cryogenian (1000–635 Ma), Mesoproterozoic (1600–1000 Ma), Paleoproterozoic (2500–1600 Ma), and Archean (4000–2500 Ma). Note the relatively heavier $\delta^{34}\text{S}$ data in both sulfate and pyrite during the Ediacaran Period.

Figure 10. Biogeochemical model for the Khatyspyt Formation. (A) Redox conditions in the lower studied interval (5–45 m) of the Khatyspyt Formation are characterized by redox stratification with oxic conditions on the surface ocean and euxinic conditions in depth. Sulfide re-oxidation and enhanced microbial sulfate reduction lead to more negative $\delta^{13}\text{C}_{\text{carb}}$ and $\delta^{34}\text{S}_{\text{pyrite}}$

774 values. **(B)** Redox conditions in the upper studied interval (45–130 m) of the Khatyspyt
 775 Formation are characterized by non-euxinic conditions with abundant Ediacara-type macrofossils.
 776 Ferruginous or oxic conditions drive authigenic pyrite formed within marine sediments, therefore
 777 resulting in superheavy pyrite due to limited sulfate concentration in porewaters. **(C)** Simplified
 778 mechanisms for S isotope cycles of the Khatyspyt Formation. **(D)** Simplified mechanisms for C
 779 isotope cycles of the Khatyspyt Formation. MSR = microbial sulfate reduction.

780 **References**

- 781 Aller, R.C., 2014. Sedimentary diagenesis, depositional environments, and benthic fluxes.
 782 Treatise on Geochemistry 8, 293–334.
- 783 Amthor, J.E., Grotzinger, J.P., Schröder, S., Bowring, S.A., Ramezani, J., Martin, M.W., Matter,
 784 A., 2003. Extinction of *Cloudina* and *Namacalathus* at the Precambrian–Cambrian boundary in
 785 Oman. *Geology* 31, 431–434.
- 786 Anbar, A.D., Knoll, A., 2002. Proterozoic ocean chemistry and evolution: a bioinorganic bridge?
 787 *Science* 297, 1137–1142.
- 788 Arthur, M.A., 2009. Carbonate rocks deconstructed. *Nature* 460, 698–699.
- 789 Bau, M., Koschinsky, A., Dulski, P., Hein, J.R., 1996. Comparison of the partitioning behaviours
 790 of yttrium, rare earth elements, and titanium between hydrogenetic marine ferromanganese crusts
 791 and seawater. *Geochim. Cosmochim. Acta* 60, 1709–1725.
- 792 Bengtson, S., Zhao, Y., 1992. Predatorial borings in late Precambrian mineralized exoskeletons.
 793 *Science* 257, 367–369.
- 794 Boag, T.H., Darroch, S.A., Laflamme, M., 2016. Ediacaran distributions in space and time:
 795 testing assemblage concepts of earliest macroscopic body fossils. *Paleobiology*, in press.
- 796 Boggiani, P.C., Gaucher, C., Sial, A.N., Babinski, M., Simon, C.M., Riccomini, C., Ferreira,
 797 V.P., Fairchild, T.R., 2010. Chemostratigraphy of the Tamengo Formation (Corumbá Group,
 798 Brazil): A contribution to the calibration of the Ediacaran carbon-isotope curve. *Precambrian Res.*
 799 182, 382–401.
- 800 Borowski, W.S., Paull, C.K., Ussler, W., 1996. Marine pore-water sulfate profiles indicate in situ
 801 methane flux from underlying gas hydrate. *Geology* 24, 655–658.

802 Borowski, W.S., Hoehler, T.M., Alperin, M.J., Rodriguez, N.M., Paull, C.K., 2000. Significance
803 of anaerobic methane oxidation in methane-rich sediments overlying the Blake Ridge gas
804 hydrates, Proceedings of the ocean drilling program, scientific results, pp. 87–99.

805 Borowski, W.S., Rodriguez, N.M., Paull, C.K., Ussler, W., 2013. Are ³⁴S-enriched authigenic
806 sulfide minerals a proxy for elevated methane flux and gas hydrates in the geologic record? Mar.
807 Pet. Geol. 43, 381–395.

808 Bouougri, E., Porada, H., 2007. Mat-related features from the Terminal Ediacaran Nudaus
809 Formation, Nama Group, Namibia, in: Schieber, J., Bose, P.K., Eriksson, P.G., Banerjee, S.,
810 Sarkar, S., Altermann, W., Catuneanu, O. (Eds.), Atlas of Microbial Mat Features Preserved
811 within the Siliciclastic Rock Record. Elsevier, pp. 214–221.

812 Bowring, S.A., Grotzinger, J.P., Isachsen, C.E., Knoll, A.H., Pelechaty, S.M., Kolosov, P., 1993.
813 Calibrating rates of Early Cambrian evolution. Science 261, 1293–1298.

814 Brasier, M.D., McIlroy, D., Liu, A.G., Antcliffe, J.B., Menon, L.R., 2013. The oldest evidence of
815 bioturbation on Earth: Comment. Geology 41, e289.

816 Brocks, J.J., Love, G.D., Summons, R.E., Knoll, A.H., Logan, G.A., Bowden, S.A., 2005.
817 Biomarker evidence for green and purple sulphur bacteria in a stratified Palaeoproterozoic sea.
818 Nature 437, 866–870.

819 Budd, G.E., Jensen, S., 2015. The origin of the animals and a ‘Savannah’ hypothesis for early
820 bilaterian evolution. Biological Reviews, in press.

821 Cai, Y., Hua, H., Xiao, S., Schiffbauer, J.D., Li, P., 2010. Biostratinomy of the late Ediacaran
822 pyritized Gaojiashan Lagerstätte from southern Shaanxi, South China: importance of event
823 deposits. Palaios 25, 487–506.

824 Canfield, D.E., Teske, A., 1996. Late Proterozoic rise in atmospheric oxygen concentration
825 inferred from phylogenetic and sulphur-isotope studies. Nature 382, 127–132.

826 Canfield, D.E., 1998. A new model for Proterozoic ocean chemistry. Nature 396, 450–453.

827 Canfield, D.E., 2001. Biogeochemistry of Sulfur Isotopes, in: Valley, J.W., Cole, D. (Eds.),
828 Stable Isotope Geochemistry, Reviews in Mineralogy and Geochemistry. Mineralogical Society
829 of America, pp. 607–636.

830 Canfield, D.E., Poulton, S.W., Narbonne, G.M., 2007. Late-Neoproterozoic deep-ocean
831 oxygenation and the rise of animal life. Science 315, 92–95.

832 Canfield, D.E., Poulton, S.W., Knoll, A.H., Narbonne, G.M., Ross, G., Goldberg, T., Strauss, H.,
833 2008. Ferruginous conditions dominated later Neoproterozoic deep-water chemistry. Science 321,
834 949–952.

835 Canfield, D.E., Farquhar, J., 2009. Animal evolution, bioturbation, and the sulfate concentration
836 of the oceans. Proc. Natl Acad. Sci. USA 106, 8123–8127.

- 837 Canfield, D.E., Farquhar, J., Zerkle, A.L., 2010. High isotope fractionations during sulfate
838 reduction in a low-sulfate euxinic ocean analog. *Geology* 38, 415–418.
- 839 Canfield, D.E., Farquhar, J., 2012. The global sulfur cycle, in: Knoll, A.H., Canfield, D.E.,
840 Konhauser, K.O. (Eds.), *Fundamentals of Geobiology*. John Wiley & Sons, Ltd, Oxford, UK, pp.
841 49–64.
- 842 Canfield, D.E., 2014. Proterozoic Atmospheric Oxygen, in: Holland, H.D., Turekian, K.K. (Eds.),
843 *Treatise on Geochemistry* (2nd Edition). Elsevier, Oxford, pp. 197–216.
- 844 Cao, H., Kaufman, A.J., Shan, X., Cui, H., Zhang, G., 2016. Sulfur isotope constraints on marine
845 transgression in the lacustrine upper cretaceous Songliao Basin, Northeastern China. *Palaeogeogr.*
846 *Palaeoclimatol. Palaeoecol.* 451, 152–163.
- 847 Chen, D., Wang, J., Racki, G., Li, H., Wang, C., Ma, X., Whalen, M.T., 2013. Large sulphur
848 isotopic perturbations and oceanic changes during the Frasnian–Famennian transition of the Late
849 Devonian. *J. Geol. Soc. London* 170, 465–476.
- 850 Chen, X., Ling, H.-F., Vance, D., Shields-Zhou, G.A., Zhu, M., Poulton, S.W., Och, L.M., Jiang,
851 S.-Y., Li, D., Cremonese, L., Archer, C., 2015. Rise to modern levels of ocean oxygenation
852 coincided with the Cambrian radiation of animals. *Nat. Commun.* 6, 7142.
- 853 Chen, Z., Zhou, C., Xiao, S., Wang, W., Guan, C., Hua, H., Yuan, X., 2014. New Ediacara
854 fossils preserved in marine limestone and their ecological implications. *Sci. Rep.* 4, 4180.
- 855 Clarkson, M., Poulton, S., Guilbaud, R., Wood, R., 2014. Assessing the utility of Fe/Al and Fe-
856 speciation to record water column redox conditions in carbonate-rich sediments. *Chem. Geol.*
857 382, 111–122.
- 858 Cloud, P., 1976. Beginnings of biospheric evolution and their biogeochemical consequences.
859 *Paleobiology* 2, 351–387.
- 860 Cloud, P.E., 1968. Atmospheric and hydrospheric evolution on the primitive Earth. *Science* 160,
861 729–736.
- 862 Condon, D., Zhu, M., Bowring, S., Wang, W., Yang, A., Jin, Y., 2005. U–Pb ages from the
863 neoproterozoic Doushantuo Formation, China. *Science* 308, 95–98.
- 864 Cook, P.J., Shergold, J.H., 1984. Phosphorus, phosphorites and skeletal evolution at the
865 Precambrian–Cambrian boundary. *Nature* 308, 231–236.
- 866 Cui, H., 2015. Authigenesis, biomineralization, and carbon–sulfur cycling in the Ediacaran ocean
867 (Ph.D. dissertation), Department of Geology. University of Maryland, College Park.
- 868 Cui, H., Kaufman, A.J., Xiao, S., Zhu, M., Zhou, C., Liu, X.-M., 2015. Redox architecture of an
869 Ediacaran ocean margin: Integrated chemostratigraphic ($\delta^{13}\text{C}$ – $\delta^{34}\text{S}$ – $^{87}\text{Sr}/^{86}\text{Sr}$ –Ce/Ce*)
870 correlation of the Doushantuo Formation, South China. *Chem. Geol.* 405, 48–62.

871 Cui, H., Kaufman, A.J., Xiao, S., Peek, S., Cao, H., Min, X., Cai, Y., Siegel, Z., Liu, X.M., Peng,
872 Y., Schiffbauer, J.D., Martin, A.J., 2016a. Environmental context for the terminal Ediacaran
873 biomineralization of animals. *Geobiology* 14, 344–363.

874 Cui, H., Xiao, S., Zhou, C., Peng, Y., Kaufman, A.J., Plummer, R.E., 2016b. Phosphogenesis
875 associated with the Shuram Excursion: Petrographic and geochemical observations from the
876 Ediacaran Doushantuo Formation of South China. *Sediment. Geol.* 341, 134–146.

877 Cui, H., Alan, K.J., Xiao, S., Zhou, C., under review. Methane-derived authigenic carbonate
878 linked to the Shuram Excursion: Petrographic and geochemical evidence for an early diagenesis
879 origin from the uppermost Doushantuo Formation in South China. *Chem. Geol.*

880 Darroch, S.A.F., Sperling, E.A., Boag, T.H., Racicot, R.A., Mason, S.J., Morgan, A.S., Tweedt,
881 S., Myrow, P., Johnston, D.T., Erwin, D.H., Laflamme, M., 2015. Biotic replacement and mass
882 extinction of the Ediacara biota. *P. Roy. Soc. Lond. B Bio.* 282, 20151003.

883 Darroch, S.A.F., Boag, T.H., Racicot, R.A., Tweedt, S., Mason, S.J., Erwin, D.H., Laflamme, M.,
884 2016. A mixed Ediacaran-metazoan assemblage from the Zaris Sub-basin, Namibia. *Palaeogeogr.*
885 *Palaeoclimatol. Palaeoecol.* 459, 198–208.

886 Derry, L.A., Kaufman, A.J., Jacobsen, S.B., 1992. Sedimentary cycling and environmental
887 change in the Late Proterozoic: evidence from stable and radiogenic isotopes. *Geochim.*
888 *Cosmochim. Acta* 56, 1317–1329.

889 Derry, L.A., 2010. A burial diagenesis origin for the Ediacaran Shuram–Wonoka carbon isotope
890 anomaly. *Earth Planet. Sci. Lett.* 294, 152–162.

891 Ding, Q., Chen, Y., 1981. Discovery of soft metazoan from the Sinian System along eastern
892 Yangtze Gorge, Hubei. *Journal of the Wuhan College of Geology* 2, 53–57, in Chinese with
893 English abstract.

894 Duda, J.-P., Blumenberg, M., Thiel, V., Simon, K., Zhu, M., Reitner, J., 2014. Geobiology of a
895 palaeoecosystem with Ediacara-type fossils: The Shibantan Member (Dengying Formation,
896 South China). *Precambrian Res.* 255, 48–62.

897 Fedonkin, M.A., 1990. Systematic description of Vendian metazoa, in: Sokolov, B.S.,
898 Iwanowski, A.B. (Eds.), *The Vendian System: Vol. 1 Paleontology*. Springer-Verlag Berlin
899 Heidelberg, Berlin, Germany, pp. 71–120.

900 Fedonkin, M.A., Gehling, J.G., Grey, K., Narbonne, G.M., Vickers-Rich, P., 2007. The rise of
901 animals: evolution and diversification of the kingdom Animalia. John Hopkins University Press,
902 Baltimore, Maryland, USA.

903 Fike, D.A., Grotzinger, J.P., Pratt, L.M., Summons, R.E., 2006. Oxidation of the Ediacaran
904 ocean. *Nature* 444, 744–747.

905 Fike, D.A., Grotzinger, J.P., 2008. A paired sulfate–pyrite $\delta^{34}\text{S}$ approach to understanding the
906 evolution of the Ediacaran–Cambrian sulfur cycle. *Geochem. Cosmochim. Acta* 72, 2636–2648.

907 Fike, D.A., Grotzinger, J.P., 2010. A $\delta^{34}\text{S}_{\text{SO}_4}$ approach to reconstructing biogenic pyrite burial in
 908 carbonate–evaporite basins: An example from the Ara Group, Sultanate of Oman. *Geology* 38,
 909 371–374.

910 Fike, D.A., Bradley, A.S., Rose, C.V., 2015. Rethinking the ancient sulfur cycle. *Annu. Rev.*
 911 *Earth Planet. Sci.* 43, 593–622.

912 Frei, R., Gaucher, C., Døssing, L.N., Sial, A.N., 2011. Chromium isotopes in carbonates — A
 913 tracer for climate change and for reconstructing the redox state of ancient seawater. *Earth Planet.*
 914 *Sci. Lett.* 312, 114–125.

915 Fry, B., Ruf, W., Gest, H., Hayes, J., 1988. Sulfur isotope effects associated with oxidation of
 916 sulfide by O_2 in aqueous solution. *Chem. Geol.* 73, 205–210.

917 Gámez Vintaned, J.A., Zhuravlev, A.Y., 2013. The oldest evidence of bioturbation on Earth:
 918 Comment. *Geology* 41, e299.

919 Gao, J., Fike, D.A., Aller, R.C., 2013. Enriched pyrite $\delta^{34}\text{S}$ signals in modern tropical deltaic
 920 muds, 2013 AGU Fall Meeting. American Geophysical Union, San Francisco, p. 0352.

921 Gaucher, C., Boggiani, P., Sprechmann, P., Sial, A., Fairchild, T., 2003. Integrated correlation of
 922 the Vendian to Cambrian Arroyo del Soldado and Corumbá Groups (Uruguay and Brazil):
 923 palaeogeographic, palaeoclimatic and palaeobiologic implications. *Precambrian Res.* 120, 241–
 924 278.

925 Gehling, J.G., 1999. Microbial mats in terminal Proterozoic siliciclastics: Ediacaran death masks.
 926 *Palaaios* 14, 40–57.

927 Gehling, J.G., Droser, M.L., 2013. How well do fossil assemblages of the Ediacara Biota tell
 928 time? *Geology* 41, 447–450.

929 Germs, G.J., 1972. New shelly fossils from Nama Group, south west Africa. *Am. J. Sci.* 272,
 930 752–761.

931 Gill, B.C., Lyons, T.W., Frank, T.D., 2008. Behavior of carbonate-associated sulfate during
 932 meteoric diagenesis and implications for the sulfur isotope paleoproxy. *Geochim. Cosmochim.*
 933 *Acta* 72, 4699–4711.

934 Gill, B.C., Lyons, T.W., Young, S.A., Kump, L.R., Knoll, A.H., Saltzman, M.R., 2011.
 935 Geochemical evidence for widespread euxinia in the Later Cambrian ocean. *Nature* 469, 80–83.

936 Gilleaudeau, G.J., Kah, L.C., 2013. Oceanic molybdenum drawdown by epeiric sea expansion in
 937 the Mesoproterozoic. *Chem. Geol.* 356, 21–37.

938 Gomes, M.L., Hurtgen, M.T., 2015. Sulfur isotope fractionation in modern euxinic systems:
 939 Implications for paleoenvironmental reconstructions of paired sulfate–sulfide isotope records.
 940 *Geochim. Cosmochim. Acta* 157, 39–55.

- 941 Grazhdankin, D., 2004. Patterns of distribution in the Ediacaran biotas: facies versus
942 biogeography and evolution. *Paleobiology* 30, 203–221.
- 943 Grazhdankin, D., Gerdes, G., 2007. Ediacaran microbial colonies. *Lethaia* 40, 201–210.
- 944 Grazhdankin, D.V., Balthasar, U., Nagovitsin, K.E., Kochnev, B.B., 2008. Carbonate-hosted
945 Avalon-type fossils in arctic Siberia. *Geology* 36, 803–806.
- 946 Grazhdankin, D.V., 2011. Ediacaran Biota, in: Reitner, J., Thiel, V. (Eds.), *Encyclopedia of*
947 *Geobiology*. Springer, Netherlands, pp. 342–348.
- 948 Grazhdankin, D.V., 2014. Patterns of evolution of the Ediacaran soft-bodied biota. *J. Paleontol.*
949 88, 269–283.
- 950 Grotzinger, J., Al-Rawahi, Z., 2014. Depositional facies and platform architecture of
951 microbialite-dominated carbonate reservoirs, Ediacaran–Cambrian Ara Group, Sultanate of
952 Oman. *AAPG Bull.* 98, 1453–1494.
- 953 Grotzinger, J.P., Bowring, B.Z., Saylor, B.Z., Kaufman, A.J., 1995. Biostratigraphic and
954 geochronologic constraints on early animal evolution. *Science* 270, 598–604.
- 955 Guilbaud, R., Poulton, S.W., Butterfield, N.J., Zhu, M., Shields-Zhou, G.A., 2015. A global
956 transition to ferruginous conditions in the early Neoproterozoic oceans. *Nat. Geosci.* 8, 466–470.
- 957 Guo, H., Du, Y., Kah, L.C., Hu, C., Huang, J., Huang, H., Yu, W., Song, H., 2015. Sulfur isotope
958 composition of carbonate-associated sulfate from the Mesoproterozoic Jixian Group, North
959 China: Implications for the marine sulfur cycle. *Precambrian Res.* 266, 319–336.
- 960 Habicht, K.S., Gade, M., Thamdrup, B., Berg, P., Canfield, D.E., 2002. Calibration of sulfate
961 levels in the Archean ocean. *Science* 298, 2372–2374.
- 962 Hall, M., Kaufman, A.J., Vickers-Rich, P., Ivantsov, A., Trusler, P., Linnemann, U., Hofmann,
963 M., Elliott, D., Cui, H., Fedonkin, M., Hoffmann, K.-H., Wilson, S.A., Schneider, G., Smith, J.,
964 2013. Stratigraphy, palaeontology and geochemistry of the late Neoproterozoic Aar Member,
965 southwest Namibia: Reflecting environmental controls on Ediacara fossil preservation during the
966 terminal Proterozoic in African Gondwana. *Precambrian Res.* 238, 214–232.
- 967 Halverson, G.P., 2006. A Neoproterozoic chronology, in: Xiao, S., Kaufman, A.J. (Eds.),
968 *Neoproterozoic Geobiology and Paleobiology*. Springer, Berlin, Germany, pp. 231–271.
- 969 Halverson, G.P., Dudás, F.Ö., Maloof, A.C., Bowring, S.A., 2007. Evolution of the $^{87}\text{Sr}/^{86}\text{Sr}$
970 composition of Neoproterozoic seawater. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 256, 103–129.
- 971 Harrison, A., Thode, H., 1958. Mechanism of the bacterial reduction of sulfate from isotope
972 fractionation studies. *Trans. Faraday Soc.* 54, 84–92.
- 973 Hayes, J.M., 1993. Factors controlling ^{13}C contents of sedimentary organic compounds:
974 Principles and evidence. *Mar. Geol.* 113, 111–125.

- 975 Hayes, J.M., Strauss, H., Kaufman, A.J., 1999. The abundance of ^{13}C in marine organic matter
976 and isotopic fractionation in the global biogeochemical cycle of carbon during the past 800 Ma.
977 Chem. Geol. 161, 103–125.
- 978 Hofmann, H., O'Brien, S., King, A., 2008. Ediacaran biota on bonavista peninsula,
979 Newfoundland, Canada. J. Paleontol. 82, 1–36.
- 980 Horton, F., 2015. Did phosphorus derived from the weathering of large igneous provinces
981 fertilize the Neoproterozoic ocean? Geochem. Geophys. Geosy. 16, 1723–1738.
- 982 Houghton, J., Fike, D., Druschel, G., Orphan, V., Hoehler, T.M., Des Marais, D.J., 2014. Spatial
983 variability in photosynthetic and heterotrophic activity drives localized $\delta^{13}\text{C}_{\text{org}}$ fluctuations and
984 carbonate precipitation in hypersaline microbial mats. Geobiology 12, 557–574.
- 985 Hua, H., Chen, Z., Zhang, L.-y., 2004. *Shaanxilithes* from lower Taozichong Formation,
986 Guizhou Province and its geological and paleobiological significance. Journal of Stratigraphy 28,
987 265–269, in Chinese.
- 988 Jiang, G., Kaufman, A.J., Christie-Blick, N., Zhang, S., Wu, H., 2007. Carbon isotope variability
989 across the Ediacaran Yangtze platform in South China: Implications for a large surface-to-deep
990 ocean $\delta^{13}\text{C}$ gradient. Earth Planet. Sci. Lett. 261, 303–320.
- 991 Jiang, G., Wang, X., Shi, X., Xiao, S., Zhang, S., Dong, J., 2012. The origin of decoupled
992 carbonate and organic carbon isotope signatures in the early Cambrian (ca. 542–520Ma) Yangtze
993 platform. Earth Planet. Sci. Lett. 317–318, 96–110.
- 994 Johnston, D.T., Wolfe-Simon, F., Pearson, A., Knoll, A.H., 2009. Anoxygenic photosynthesis
995 modulated Proterozoic oxygen and sustained Earth's middle age. Proc. Natl Acad. Sci. USA 106,
996 16925–16929.
- 997 Johnston, D.T., Poulton, S.W., Tosca, N.J., O'Brien, T., Halverson, G.P., Schrag, D.P.,
998 Macdonald, F.A., 2013. Searching for an oxygenation event in the fossiliferous Ediacaran of
999 northwestern Canada. Chem. Geol. 362, 273–286.
- 1000 Jørgensen, B.B., Böttcher, M.E., Lüschen, H., Neretin, L.N., Volkov, I.I., 2004. Anaerobic
1001 methane oxidation and a deep H_2S sink generate isotopically heavy sulfides in Black Sea
1002 sediments 1. Geochim. Cosmochim. Acta 68, 2095–2118.
- 1003 Kah, L.C., Lyons, T.W., Frank, T.D., 2004. Low marine sulphate and protracted oxygenation of
1004 the Proterozoic biosphere. Nature 431, 834–838.
- 1005 Kah, L.C., Bartley, J.K., 2011. Protracted oxygenation of the Proterozoic biosphere. Int. Geol.
1006 Rev. 53, 1424–1442.
- 1007 Kah, L.C., Thompson, C.K., Henderson, M.A., Zhan, R., 2016. Behavior of marine sulfur in the
1008 Ordovician. Palaeogeogr. Palaeoclimatol. Palaeoecol. 458, 133–153.

- 1009 Kaplan, I.R., Rafter, T.A., 1958. Fractionation of stable isotopes of sulfur by Thiobacilli. *Science*
1010 127, 517–518.
- 1011 Kaplan, I.R., Rittenberg, S.C., 1964. Microbiological fractionation of sulphur isotopes.
1012 *Microbiology* 34, 195–212.
- 1013 Kaufman, A.J., Hayes, J.M., Knoll, A.H., Germs, G.J.B., 1991. Isotopic compositions of
1014 carbonates and organic carbon from upper Proterozoic successions in Namibia: stratigraphic
1015 variation and the effects of diagenesis and metamorphism. *Precambrian Res.* 49, 301–327.
- 1016 Kaufman, A.J., Jacobsen, S.B., Knoll, A.H., 1993. The Vendian record of Sr and C isotopic
1017 variations in seawater: implications for tectonics and paleoclimate. *Earth Planet. Sci. Lett.* 120,
1018 409–430.
- 1019 Kaufman, A.J., Corsetti, F.A., Varni, M.A., 2007. The effect of rising atmospheric oxygen on
1020 carbon and sulfur isotope anomalies in the Neoproterozoic Johnnie Formation, Death Valley,
1021 USA. *Chem. Geol.* 237, 47–63.
- 1022 Kaufman, A.J., Peek, S., Aaron, J., Cui, H., Grazhdankin, D., Rogov, V., Xiao, S., Buchwaldt, R.,
1023 Bowring, S., 2012. A shorter fuse for the Cambrian Explosion?, 2012 GSA Annual Meeting. The
1024 Geological Society of America, Charlotte, North Carolina, USA.
- 1025 Knauth, L.P., Kennedy, M.J., 2009. The late Precambrian greening of the Earth. *Nature* 460,
1026 728–732.
- 1027 Knoll, A.H., Grotzinger, J.P., Kaufman, A.J., Kolosov, P., 1995. Integrated approaches to
1028 terminal Proterozoic stratigraphy: an example from the Olenek Uplift, northeastern Siberia.
1029 *Precambrian Res.* 73, 251–270.
- 1030 Knoll, A.H., Sperling, E.A., 2014. Oxygen and animals in Earth history. *Proc. Natl Acad. Sci.*
1031 USA 111, 3907–3908.
- 1032 Kolesnikov, A.V., Marusin, V.V., Nagovitsin, K.E., Maslov, A.V., Grazhdankin, D.V., 2015.
1033 Ediacaran biota in the aftermath of the Kotlinian Crisis: Asha Group of the South Urals.
1034 *Precambrian Res.* 263, 59–78.
- 1035 Kump, L.R., 2008. The rise of atmospheric oxygen. *Nature* 451, 277–278.
- 1036 Lawrence, M.G., Greig, A., Collerson, K.D., Kamber, B.S., 2006. Rare Earth Element and
1037 Yttrium variability in South East Queensland waterways. *Aquatic Geochemistry* 12, 39–72.
- 1038 Leavitt, W.D., Bradley, A.S., Halevy, I., Johnston, D.T., 2013. Influence of sulfate reduction
1039 rates on the Phanerozoic sulfur isotope record. *Proc. Natl Acad. Sci. USA* 110, 11244–11249.
- 1040 Li, C., Love, G.D., Lyons, T.W., Fike, D.A., Sessions, A.L., Chu, X., 2010. A stratified redox
1041 model for the Ediacaran ocean. *Science* 328, 80–83.

- 1042 Li, C., Cheng, M., Algeo, T.J., Xie, S., 2015a. A theoretical prediction of chemical zonation in
1043 early oceans (>520 Ma). *Sci. China Earth Sci.* 11, 1901–1909.
- 1044 Li, C., Planavsky, N.J., Shi, W., Zhang, Z., Zhou, C., Cheng, M., Tarhan, L.G., Luo, G., Xie, S.,
1045 2015b. Ediacaran marine redox heterogeneity and early animal ecosystems. *Sci. Rep.* 5, 17097.
- 1046 Li, R., Chen, J., Zhang, S., Lei, J., Shen, Y., Chen, X., 1999. Spatial and temporal variations in
1047 carbon and sulfur isotopic compositions of Sinian sedimentary rocks in the Yangtze platform,
1048 South China. *Precambrian Res.* 97, 59–75.
- 1049 Lin, Q., Wang, J., Algeo, T.J., Sun, F., Lin, R., 2016a. Enhanced framboidal pyrite formation
1050 related to anaerobic oxidation of methane in the sulfate-methane transition zone of the northern
1051 South China Sea. *Mar. Geol.* 379, 100–108.
- 1052 Lin, Z., Sun, X., Peckmann, J., Lu, Y., Xu, L., Strauss, H., Zhou, H., Gong, J., Lu, H., Teichert,
1053 B.M.A., 2016b. How sulfate-driven anaerobic oxidation of methane affects the sulfur isotopic
1054 composition of pyrite: A SIMS study from the South China Sea. *Chem. Geol.* 440, 26–41.
- 1055 Ling, H.-F., Chen, X., Li, D., Wang, D., Shields-Zhou, G.A., Zhu, M., 2013. Cerium anomaly
1056 variations in Ediacaran–earliest Cambrian carbonates from the Yangtze Gorges area, South
1057 China: Implications for oxygenation of coeval shallow seawater. *Precambrian Res.* 225, 110–127.
- 1058 Liu, T.-b., Maynard, J.B., Alten, J., 2006. Superheavy S isotopes from glacier-associated
1059 sediments of the Neoproterozoic of south China: Oceanic anoxia or sulfate limitation? *Geol. Soc.
1060 Am. Mem.* 198, 205–222.
- 1061 Liu, X.-M., Kah, L.C., Knoll, A.H., Cui, H., Kaufman, A.J., Shahar, A., Hazen, R.M., 2016.
1062 Tracing Earth's O₂ evolution using Zn/Fe ratios in marine carbonates. *Geochem. Persp. Lett.* 2,
1063 24–34.
- 1064 Logan, G.A., Hayes, J., Hieshima, G.B., Summons, R.E., 1995. Terminal Proterozoic
1065 reorganization of biogeochemical cycles. *Nature* 376, 53–56.
- 1066 Loyd, S.J., Marenco, P.J., Hagadorn, J.W., Lyons, T.W., Kaufman, A.J., Sour-Tovar, F., Corsetti,
1067 F.A., 2012. Sustained low marine sulfate concentrations from the Neoproterozoic to the
1068 Cambrian: Insights from carbonates of northwestern Mexico and eastern California. *Earth Planet.
1069 Sci. Lett.* 339–340, 79–94.
- 1070 Loyd, S.J., Marenco, P.J., Hagadorn, J.W., Lyons, T.W., Kaufman, A.J., Sour-Tovar, F., Corsetti,
1071 F.A., 2013. Local $\delta^{34}\text{S}$ variability in ~580Ma carbonates of northwestern Mexico and the
1072 Neoproterozoic marine sulfate reservoir. *Precambrian Res.* 224, 551–569.
- 1073 Lyons, T.W., Walter, L.M., Gellatly, A.M., Martini, A.M., Blake, R.E., 2004. Sites of anomalous
1074 organic remineralization in the carbonate sediments of South Florida, USA: the sulfur cycle and
1075 carbonate-associated sulfate, in: Amend, J.P., Edwards, K.J., Lyons, T.W. (Eds.), *Sulfur
1076 Biogeochemistry—Past and Present: Geological Society of America Special Paper* 379.
1077 Geological Society of America, Boulder, Colorado, USA, pp. 161–176.

- 1078 Lyons, T.W., Reinhard, C.T., Planavsky, N.J., 2014. The rise of oxygen in Earth's early ocean
1079 and atmosphere. *Nature* 506, 307–315.
- 1080 Macdonald, F.A., Strauss, J.V., Sperling, E.A., Halverson, G.P., Narbonne, G.M., Johnston, D.T.,
1081 Kunzmann, M., Schrag, D.P., Higgins, J.A., 2013. The stratigraphic relationship between the
1082 Shuram carbon isotope excursion, the oxygenation of Neoproterozoic oceans, and the first
1083 appearance of the Ediacara biota and bilaterian trace fossils in northwestern Canada. *Chem. Geol.*
1084 362, 250–272.
- 1085 Mángano, M.G., Buatois, L.A., 2014. Decoupling of body-plan diversification and ecological
1086 structuring during the Ediacaran–Cambrian transition: evolutionary and geobiological feedbacks.
1087 *P. Roy. Soc. Lond. B Bio.* 281, 20140038.
- 1088 Marengo, P.J., Corsetti, F.A., Hammond, D.E., Kaufman, A.J., Bottjer, D.J., 2008. Oxidation of
1089 pyrite during extraction of carbonate associated sulfate. *Chem. Geol.* 247, 124–132.
- 1090 McArthur, J.M., Howarth, R.J., Shields, G.A., 2012. Strontium Isotope Stratigraphy, in:
1091 Gradstein, F.M., Ogg, J.G., Schmitz, M.D., Ogg, G.M. (Eds.), *The Geologic Time Scale*.
1092 Elsevier, Boston, pp. 127–144.
- 1093 McFadden, K.A., Huang, J., Chu, X., Jiang, G., Kaufman, A.J., Zhou, C., Yuan, X., Xiao, S.,
1094 2008. Pulsed oxidation and biological evolution in the Ediacaran Doushantuo Formation. *Proc.*
1095 *Natl Acad. Sci. USA* 105, 3197–3202.
- 1096 Meyer, M., Schiffbauer, J.D., Xiao, S., Cai, Y., Hua, H., 2012. Taphonomy of the upper
1097 Ediacaran enigmatic ribbonlike fossil *Shaanxilithes*. *Palaios* 27, 354–372.
- 1098 Mills, D.B., Canfield, D.E., 2014. Oxygen and animal evolution: Did a rise of atmospheric
1099 oxygen “trigger” the origin of animals? *BioEssays* 36, 1145–1155.
- 1100 Mills, D.B., Ward, L.M., Jones, C., Sweeten, B., Forth, M., Treusch, A.H., Canfield, D.E., 2014.
1101 Oxygen requirements of the earliest animals. *Proc. Natl Acad. Sci. USA* 111, 4168–4172.
- 1102 Muscente, A.D., Hawkins, A.D., Xiao, S., 2015. Fossil preservation through phosphatization and
1103 silicification in the Ediacaran Doushantuo Formation (South China): a comparative synthesis.
1104 *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 434, 46–62.
- 1105 Nagovitsin, K.E., Rogov, V.I., Marusin, V.V., Karlova, G.A., Kolesnikov, A.V., Bykova, N.V.,
1106 Grazhdankin, D.V., 2015. Revised Neoproterozoic and Terreneuvian stratigraphy of the Lena-
1107 Anabar Basin and north-western slope of the Olenek Uplift, Siberian Platform. *Precambrian Res.*
1108 270, 226–245.
- 1109 Narbonne, G.M., Kaufman, A.J., Knoll, A.H., 1994. Integrated chemostratigraphy and
1110 biostratigraphy of the Windermere Supergroup, northwestern Canada: Implications for
1111 Neoproterozoic correlations and the early evolution of animals. *Geol. Soc. Am. Bull.* 106, 1281–
1112 1292.

- 1113 Narbonne, G.M., 2005. The Ediacara Biota: Neoproterozoic origin of animals and their
1114 ecosystems. *Annu. Rev. Earth Planet. Sci.* 33, 421–442.
- 1115 Narbonne, G.M., Laflamme, M., Trusler, P.W., Dalrymple, R.W., Greentree, C., 2014. Deep-
1116 water Ediacaran fossils from northwestern Canada: taphonomy, ecology, and evolution. *J.*
1117 *Paleontol.* 88, 207–223.
- 1118 Och, L.M., Shields-Zhou, G.A., 2012. The Neoproterozoic oxygenation event: Environmental
1119 perturbations and biogeochemical cycling. *Earth-Sci. Rev.* 110, 26–57.
- 1120 Och, L.M., Cremonese, L., Shields-Zhou, G.A., Poulton, S.W., Struck, U., Ling, H., Li, D., Chen,
1121 X., Manning, C., Thirlwall, M., Strauss, H., Zhu, M., 2015. Palaeoceanographic controls on
1122 spatial redox distribution over the Yangtze Platform during the Ediacaran–Cambrian transition.
1123 *Sedimentology* 63, 378–410.
- 1124 Oehlert, A.M., Lamb-Wozniak, K.A., Devlin, Q.B., Mackenzie, G.J., Reijmer, J.J.G., Swart,
1125 P.K., 2012. The stable carbon isotopic composition of organic material in platform derived
1126 sediments: implications for reconstructing the global carbon cycle. *Sedimentology* 59, 319–335.
- 1127 Oehlert, A.M., Swart, P.K., 2014. Interpreting carbonate and organic carbon isotope covariance
1128 in the sedimentary record. *Nat. Commun.* 5, 4672.
- 1129 Papineau, D., 2010. Global biogeochemical changes at both ends of the Proterozoic: Insights
1130 from phosphorites. *Astrobiology* 10, 165–181.
- 1131 Pelechaty, S.M., Grotzinger, J.P., Kashirtsev, V.A., Zhernovsky, V.P., 1996a.
1132 Chemostratigraphic and sequence stratigraphic constraints on Vendian–Cambrian basin
1133 dynamics, northeast Siberian craton. *J. Geol.* 104, 543–563.
- 1134 Pelechaty, S.M., Kaufman, A.J., Grotzinger, J.P., 1996b. Evaluation of $\delta^{13}\text{C}$ chemostratigraphy
1135 for intrabasinal correlation: Vendian strata of northeast Siberia. *Geol. Soc. Am. Bull.* 108, 992–
1136 1003.
- 1137 Peng, Y., Bao, H., Pratt, L.M., Kaufman, A.J., Jiang, G., Boyd, D., Wang, Q., Zhou, C., Yuan,
1138 X., Xiao, S., Loyd, S., 2014. Widespread contamination of carbonate-associated sulfate by
1139 present-day secondary atmospheric sulfate: Evidence from triple oxygen isotopes. *Geology* 42,
1140 815–818.
- 1141 Planavsky, N.J., Reinhard, C.T., Wang, X., Thomson, D., McGoldrick, P., Rainbird, R.H.,
1142 Johnson, T., Fischer, W.W., Lyons, T.W., 2014. Low Mid-Proterozoic atmospheric oxygen
1143 levels and the delayed rise of animals. *Science* 346, 635–638.
- 1144 Poulton, S.W., Fralick, P.W., Canfield, D.E., 2004. The transition to a sulphidic ocean ~1.84
1145 billion years ago. *Nature* 431, 173–177.
- 1146 Poulton, S.W., Canfield, D.E., 2011. Ferruginous conditions: A dominant feature of the ocean
1147 through Earth's history. *Elements* 7, 107–112.

- 1148 Prokoph, A., Shields, G.A., Veizer, J., 2008. Compilation and time-series analysis of a marine
1149 carbonate $\delta^{18}\text{O}$, $\delta^{13}\text{C}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{34}\text{S}$ database through Earth history. *Earth-Sci. Rev.* 87, 113–
1150 133.
- 1151 Reinhard, C.T., Planavsky, N.J., Olson, S.L., Lyons, T.W., Erwin, D.H., 2016. Earth's oxygen
1152 cycle and the evolution of animal life. *Proc. Natl Acad. Sci. USA* 113, 8933–8938.
- 1153 Ries, J.B., Fike, D.A., Pratt, L.M., Lyons, T.W., Grotzinger, J.P., 2009. Superheavy pyrite
1154 ($\delta^{34}\text{S}_{\text{pyr}} > \delta^{34}\text{S}_{\text{CAS}}$) in the terminal Proterozoic Nama Group, southern Namibia: A consequence of
1155 low seawater sulfate at the dawn of animal life. *Geology* 37, 743–746.
- 1156 Rogov, V.I., Marusin, V., Bykova, N.V., Goy, Y., Nagovitsin, K.E., Kochnev, B.B., Karlova, G.,
1157 Grazhdankin, D.V., 2012. The oldest evidence of bioturbation on Earth. *Geology* 40, 395–398.
- 1158 Rogov, V.I., Marusin, V., Bykova, N., Goy, Y., Nagovitsin, K., Kochnev, B., Karlova, G.,
1159 Grazhdankin, D.V., 2013a. The oldest evidence of bioturbation on Earth: Reply. *Geology* 41,
1160 e290.
- 1161 Rogov, V.I., Marusin, V., Bykova, N.V., Goy, Y., Nagovitsin, K.E., Kochnev, B.B., Karlova, G.,
1162 Grazhdankin, D.V., 2013b. The oldest evidence of bioturbation on Earth: Reply. *Geology* 41,
1163 e300.
- 1164 Rogov, V.I., Karlova, G.A., Marusin, V.V., Kochnev, B.B., Nagovitsin, K.E., Grazhdankin, D.V.,
1165 2015. Duration of the first biozone in the Siberian hypostratotype of the Vendian. *Russ. Geol.*
1166 *Geophys.* 56, 573–583.
- 1167 Sahoo, S.K., Planavsky, N.J., Kendall, B., Wang, X., Shi, X., Scott, C., Anbar, A.D., Lyons,
1168 T.W., Jiang, G., 2012. Ocean oxygenation in the wake of the Marinoan glaciation. *Nature* 489,
1169 546–549.
- 1170 Sahoo, S.K., Planavsky, N.J., Jiang, G., Kendall, B., Owens, J.D., Wang, X., Shi, X., Anbar,
1171 A.D., Lyons, T.W., 2016. Oceanic oxygenation events in the anoxic Ediacaran ocean.
1172 *Geobiology*, In press.
- 1173 Sansjofre, P., Ader, M., Trindade, R., Elie, M., Lyons, J., Cartigny, P., Nogueira, A., 2011. A
1174 carbon isotope challenge to the snowball Earth. *Nature* 478, 93–96.
- 1175 Saylor, B.Z., Kaufman, A.J., Grotzinger, J.P., Urban, F., 1998. A composite reference section for
1176 terminal Proterozoic strata of southern Namibia. *J. Sediment. Res.* 68, 1223–1235.
- 1177 Schiffbauer, J.D., Xiao, S., Cai, Y., Wallace, A.F., Hua, H., Hunter, J., Xu, H., Peng, Y.,
1178 Kaufman, A.J., 2014. A unifying model for Neoproterozoic–Palaeozoic exceptional fossil
1179 preservation through pyritization and carbonaceous compression. *Nat. Commun.* 5, 5754.
- 1180 Schobben, M., Stebbins, A., Ghaderi, A., Strauss, H., Korn, D., Korte, C., 2015. Flourishing
1181 ocean drives the end-Permian marine mass extinction. *Proc. Natl Acad. Sci. USA* 112, 10298–
1182 10303.

- 1183 Scott, C., Lyons, T.W., Bekker, A., Shen, Y., Poulton, S.W., Chu, X., Anbar, A.D., 2008.
1184 Tracing the stepwise oxygenation of the Proterozoic ocean. *Nature* 452, 456–459.
- 1185 Shen, B., Xiao, S., Dong, L., Chuanming, Z., Liu, J., 2007. Problematic macrofossils from
1186 Ediacaran successions in the North China and Chaidam blocks: Implications for their
1187 evolutionary roots and biostratigraphic significance. *J. Paleontol.* 81, 1396–1411.
- 1188 Shen, B., Xiao, S., Kaufman, A.J., Bao, H., Zhou, C., Wang, H., 2008. Stratification and mixing
1189 of a post-glacial Neoproterozoic ocean: Evidence from carbon and sulfur isotopes in a cap
1190 dolostone from northwest China. *Earth Planet. Sci. Lett.* 265, 209–228.
- 1191 Shields-Zhou, G., Och, L., 2011. The case for a Neoproterozoic oxygenation event: geochemical
1192 evidence and biological consequences. *GSA Today* 21, 4–11.
- 1193 Shields, G., 2007. A normalised seawater strontium isotope curve: possible implications for
1194 Neoproterozoic–Cambrian weathering rates and the further oxygenation of the Earth. *eEarth* 2,
1195 35–42.
- 1196 Sim, M.S., Ono, S., Hurtgen, M.T., 2015. Sulfur isotope evidence for low and fluctuating sulfate
1197 levels in the Late Devonian ocean and the potential link with the mass extinction event. *Earth*
1198 *Planet. Sci. Lett.* 419, 52–62.
- 1199 Sokolov, B., Fedonkin, M., 1984. The Vendian as the terminal system of the Precambrian.
1200 *Episodes* 7, 12–19.
- 1201 Sperling, E.A., Carbone, C., Strauss, J.V., Johnston, D.T., Narbonne, G.M., Macdonald, F.A.,
1202 2015a. Oxygen, facies, and secular controls on the appearance of Cryogenian and Ediacaran
1203 body and trace fossils in the Mackenzie Mountains of northwestern Canada. *Geol. Soc. Am. Bull.*
1204 128, 558–575.
- 1205 Sperling, E.A., Knoll, A.H., Girguis, P.R., 2015b. The ecological physiology of Earth's second
1206 oxygen revolution. *Annu. Rev. Ecol. Evol. Syst.* 46, 215–235.
- 1207 Sperling, E.A., Wolock, C.J., Morgan, A.S., Gill, B.C., Kunzmann, M., Halverson, G.P.,
1208 Macdonald, F.A., Knoll, A.H., Johnston, D.T., 2015c. Statistical analysis of iron geochemical
1209 data suggests limited late Proterozoic oxygenation. *Nature* 523, 451–454.
- 1210 Sun, W., 1986. Late Precambrian pennatulids (sea pens) from the eastern Yangtze Gorge, China:
1211 *Paracharnia* gen. nov. *Precambrian Res.* 31, 361–375.
- 1212 Tanaka, K., Tani, Y., Takahashi, Y., Tanimizu, M., Suzuki, Y., Kozai, N., Ohnuki, T., 2010. A
1213 specific Ce oxidation process during sorption of rare earth elements on biogenic Mn oxide
1214 produced by *Acremonium* sp. strain KR21-2. *Geochim. Cosmochim. Acta* 74, 5463–5477.
- 1215 Tarhan, L.G., Droser, M.L., 2014. Widespread delayed mixing in early to middle Cambrian
1216 marine shelfal settings. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 399, 310–322.

- 1217 Tarhan, L.G., Hughes, N.C., Myrow, P.M., Bhargava, O.N., Ahluwalia, A.D., Kudryavtsev, A.B.,
1218 2014. Precambrian–Cambrian boundary interval occurrence and form of the enigmatic tubular
1219 body fossil *Shaanxilithes ningqiangensis* from the Lesser Himalaya of India. *Palaeontology* 57,
1220 283–298.
- 1221 Tarhan, L.G., Droser, M.L., Planavsky, N.J., Johnston, D.T., 2015. Protracted development of
1222 bioturbation through the early Palaeozoic Era. *Nat. Geosci.* 8, 865–869.
- 1223 Theiling, B.P., Coleman, M., 2015. Refining the extraction methodology of carbonate associated
1224 sulfate: Evidence from synthetic and natural carbonate samples. *Chem. Geol.* 411, 36–48.
- 1225 Thomazo, C., Vennin, E., Brayard, A., Bour, I., Mathieu, O., Elmeknassi, S., Olivier, N.,
1226 Escarguel, G., Bylund, K.G., Jenks, J., Stephen, D.A., Fara, E., 2016. A diagenetic control on the
1227 Early Triassic Smithian–Spathian carbon isotopic excursions recorded in the marine settings of
1228 the Thaynes Group (Utah, USA). *Geobiology* 14, 220–236.
- 1229 Thompson, C.K., Kah, L.C., 2012. Sulfur isotope evidence for widespread euxinia and a
1230 fluctuating oxycline in Early to Middle Ordovician greenhouse oceans. *Palaeogeogr.*
1231 *Palaeoclimatol. Palaeoecol.* 313–314, 189–214.
- 1232 Vishnevskaya, I., Kochnev, B., Letnikova, E., Kiseleva, V.Y., Pisareva, N., 2013. Sr isotope
1233 signatures in the Vendian Khorbusuonka group of the Olenek uplift (Northeastern Siberian
1234 Platform). *Dokl. Earth Sci.* 449, 298–302.
- 1235 Wang, L., Shi, X., Jiang, G., 2012. Pyrite morphology and redox fluctuations recorded in the
1236 Ediacaran Doushantuo Formation. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 333–334, 218–227.
- 1237 Wang, X., Shi, X., Jiang, G., Tang, D., 2014. Organic carbon isotope gradient and ocean
1238 stratification across the late Ediacaran–Early Cambrian Yangtze Platform. *Sci. China Earth Sci.*
1239 57, 919–929.
- 1240 Warren, L.V., Fairchild, T.R., Gaucher, C., Boggiani, P.C., Poire, D.G., Anelli, L.E., Inchausti,
1241 J.C., 2011. *Corumbella* and *in situ Cloudina* in association with thrombolites in the Ediacaran
1242 Itapucumi Group, Paraguay. *Terra Nova* 23, 382–389.
- 1243 Wing, B.A., Halevy, I., 2014. Intracellular metabolite levels shape sulfur isotope fractionation
1244 during microbial sulfate respiration. *Proc. Natl Acad. Sci. USA* 111, 18116–18125.
- 1245 Wood, R.A., Poulton, S.W., Prave, A.R., Hoffmann, K.H., Clarkson, M.O., Guilbaud, R., Lyne,
1246 J.W., Tostevin, R., Bowyer, F., Penny, A.M., Curtis, A., Kasemann, S.A., 2015. Dynamic redox
1247 conditions control late Ediacaran metazoan ecosystems in the Nama Group, Namibia.
1248 *Precambrian Res.* 261, 252–271.
- 1249 Wotte, T., Shields-Zhou, G.A., Strauss, H., 2012. Carbonate-associated sulfate: Experimental
1250 comparisons of common extraction methods and recommendations toward a standard analytical
1251 protocol. *Chem. Geol.* 326–327, 132–144.

- 1252 Wu, C., Zhang, Z., Xiao, J., Fu, Y., Shao, S., Zheng, C., Yao, J., Xiao, C., 2016. Nanhuan
1253 manganese deposits within restricted basins of the southeastern Yangtze Platform, China:
1254 Constraints from geological and geochemical evidence. *Ore Geology Reviews* 75, 76–99.
- 1255 Wu, N., Farquhar, J., Strauss, H., 2014. $\delta^{34}\text{S}$ and $\Delta^{33}\text{S}$ records of Paleozoic seawater sulfate
1256 based on the analysis of carbonate associated sulfate. *Earth Planet. Sci. Lett.* 399, 44–51.
- 1257 Xiao, S., Yuan, X., Steiner, M., Knoll, A.H., 2002. Macroscopic carbonaceous compressions in a
1258 terminal Proterozoic shale: A systematic reassessment of the Miaohu Biota, South China. *J.*
1259 *Paleontol.* 76, 347–376.
- 1260 Xiao, S., Shen, B., Zhou, C., Xie, G., Yuan, X., 2005. A uniquely preserved Ediacaran fossil
1261 with direct evidence for a quilted bodyplan. *Proceedings of the National Academy of Sciences of*
1262 *the United States of America* 102, 10227–10232.
- 1263 Xiao, S., Laflamme, M., 2009. On the eve of animal radiation: phylogeny, ecology and evolution
1264 of the Ediacara biota. *Trends Ecol. Evol.* 24, 31–40.
- 1265 Xiao, S., Droser, M., Gehling, J.G., Hughes, I.V., Wan, B., Chen, Z., Yuan, X., 2013. Affirming
1266 life aquatic for the Ediacara biota in China and Australia. *Geology* 41, 1095–1098.
- 1267 Xiao, S., 2014. Oxygen and early animal evolution, in: Holland, H.D., Turekian, K.K. (Eds.),
1268 *Treatise on Geochemistry, Second Edition, vol. 6 (The Atmosphere - History)*. Elsevier, Oxford,
1269 pp. 231–250.
- 1270 Xiao, S., Narbonne, G.M., Zhou, C., Laflamme, M., Grazhdankin, D.V., Moczyłowska-Vidal,
1271 M., Cui, H., 2016. Toward an Ediacaran Time Scale: Problems, protocols, and prospects.
1272 Episodes, in press.
- 1273 Xu, X., Huang, H., Liu, B., 1990. Manganese deposits of the Proterozoic Datangpo Formation,
1274 South China: genesis and palaeogeography, in: Parnell, J., Ye, L.J., Chen, C.M. (Eds.),
1275 *Sediment-Hosted Mineral Deposits: Proceedings of a Symposium*. Blackwell Publishing
1276 Ltd., Oxford, UK, Beijing, People's Republic of China, 30 July–4 August 1988, pp. 39–50.
- 1277 Zhelezinskaia, I., Kaufman, A.J., Farquhar, J., Cliff, J., 2014. Large sulfur isotope fractionations
1278 associated with Neoproterozoic microbial sulfate reduction. *Science* 346, 742–744.
- 1279 Zhou, C., Xiao, S., 2007. Ediacaran $\delta^{13}\text{C}$ chemostratigraphy of South China. *Chem. Geol.* 237,
1280 89–108.
- 1281 Zhou, C., Jiang, S., Xiao, S., Chen, Z., Yuan, X., 2012. Rare earth elements and carbon isotope
1282 geochemistry of the Doushantuo Formation in South China: Implication for middle Ediacaran
1283 shallow marine redox conditions. *Chin. Sci. Bull.* 57, 1998–2006.
- 1284 Zhuravlev, A.Y., Gámez Vintaned, J.A., Ivantsov, A.Y., 2009. First finds of problematic
1285 Ediacaran fossil *Gaojiashania* in Siberia and its origin. *Geol. Mag.* 146, 775–780.
- 1286

Figure 1

A Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Oman Figure 7

Namibia

South China

Siberia

Figure 9

lower $[\text{SO}_4]_{\text{SW}}$

non-euxinic

authigenic pyrite within sediments (^{34}S -enriched)
enhanced organic C recycling ($^{13}\text{C}_{\text{org}}$ -depleted)

B Khatyspyt Fm (40–130 m)

MSR within sediments

sulfide oxidation

oxic

higher $[\text{SO}_4]_{\text{SW}}$

sulfide

sulfate

sulfate reduction

euxinic

syngenetic pyrite in water column ($^{34}\text{S}_{\text{py}}$ -depleted)
MSR-derived alkalinity ($^{13}\text{C}_{\text{carb}}$ -depleted)

A Khatyspyt Fm (5–40 m)

C

$\delta^{13}\text{C}_{\text{org}}$ negative anomaly

$\delta^{13}\text{C}_{\text{carb}}$ positive anomaly $+6\text{‰}$

MSR-derived alkalinity
organic C recycling

enhanced organic
carbon burial

–10 –5 0 +5 +10

$\delta^{13}\text{C}_{\text{carb}}$ (‰, PDB)

–40 –35 –30 –25 –20

$\delta^{13}\text{C}_{\text{org}}$ (‰, PDB)

D

superheavy pyrite

terminal Ediacaran seawater 40‰

terminal Ediacaran pyrite

ca. 10‰

sulfide oxidation

MSR within sediments
enhanced sulfate reduction rate

–20 –10 0 +10 +20 +30 +40 +50 +60

$\delta^{34}\text{S}$ (‰, CDT)

Figure 10