Submitted to Palaeogeography, Palaeoclimatology, Palaeoecology on April 4th, 2016,
revised and resubmitted on October 9th, 2016

Abstract

We present new, monthly-resolved Sr/Ca-based sea-surface temperature (SST) records from two species of massive coral, *Orbicella faveolata* and *Siderastrea siderea*, from the Dry Tortugas National Park, FL (DTNP). We combine these new records with published data from three additional *S. siderea* coral colonies to generate a 278-year long multi-species stacked Sr/Ca-SST record from DTNP. The composite record of mean annual Sr/Ca-SST at DTNP shows pronounced decadal-scale variability with a range of 1 to 2 °C. Notable cool intervals in the Sr/Ca-derived SST lasting about a decade centered at ~1845, ~1935, and ~1965 are associated with reduced summer Sr/Ca-SST (monthly maxima < 29 °C), and imply a reduction in the spatial extent of the Atlantic Warm Pool (AWP). There is significant coherence between the composite DTNP Sr/Ca-SST record and the Atlantic Multidecadal Oscillation (AMO) index, with the AMO lagging Sr/Ca-SST at DTNP by 9 years. Low frequency variability in the Gulf Stream surface transport, which originates near DTNP, may provide a link for the lagged relationship between multidecadal variability at DTNP and the AMO.

Introduction

Understanding the drivers of inter-annual to decadal climate variability during the late Holocene is critical in making accurate predictions of future climate change.

Instrumental observations of climatic parameters such as sea-surface temperature (SST) aid in understanding the natural climate system, but limited spatial and temporal coverage of instrumental records leaves large gaps in our knowledge of climate history (Deser et al., 2010). Proxy records developed from geologic archives, such as corals, help fill these data gaps and provide high-resolution records of local and regional paleoceanographic variability. Previous studies have established that the ratio of Strontium to Calcium (Sr/Ca) of coral skeletal aragonite varies primarily as a function of water temperature at the time of coral growth (e.g., Swart et al., 2002; Corregge, 2006; Smith et al., 2006; Maupin et al., 2008; DeLong et al., 2011; Flannery and Poore, 2013; DeLong et al., 2014).

In this study, we present two new Sr/Ca-based SST records from two widespread Atlantic/Caribbean species of massive scleractinian corals, *Siderastraea siderea* and *Orbicella faveolata* (formerly *Montastraea faveolata*), from the Dry Tortugas National Park (DTNP). We then integrate these new records with previously published coral Sr/Ca-SST records from DTNP to provide an updated estimate of SST variability at this locality for the last two centuries.

DTNP consists of a shallow platform with several small shoals located at the western end of the Florida Keys, approximately 100 km west of Key West, Florida (Figure 1, inset). DTNP lies along the northern boundary of the Florida Straits, where Gulf of Mexico and Caribbean waters flow into the North Atlantic Ocean via the Florida Current. This forms the beginning of the Gulf Stream, the primary surface return component of northern Atlantic Meridional Overturning Circulation (AMOC). Patterns in SST variability at DTNP are positively correlated with broader-scale SST patterns in the Gulf of Mexico (GOM,) and along the Gulf Stream ($r \geq 0.7$, $p < 0.01$) (Figure 1). Given

its sensitivity to Gulf Stream variability, coral-based SST reconstructions at DTNP can provide insights into past changes in surface circulation in the western Atlantic Ocean.

DTNP is also situated within the Atlantic Warm Pool (AWP), a feature defined by surface waters ≥ 28.5 °C that develops seasonally in the region (Wang and Enfield, 2001; Wang et al., 2006). The AWP first develops during the boreal spring and expands through the GOM, Caribbean Sea and western tropical North Atlantic from July to October (Wang et al., 2006; Wang et al., 2008). Multidecadal variability in the size of the AWP correlates with variations in rainfall in the Caribbean, formation and intensification of North Atlantic hurricanes, and variations in atmospheric transport of moisture to the North American continent (Wang et al., 2006; Wang et al., 2008; Richey et al., 2009; DeLong et al., 2014).

Multidecadal-scale variations in SST of the North Atlantic Ocean have been identified in the instrumental record (Schlesinger and Ramankutty, 1994) and are commonly referred to as the Atlantic Multidecadal Oscillation (AMO; Kerr, 2005). An AMO index calculated from detrended SST anomalies averaged over the North Atlantic Ocean from 0–70 °N has shown that the AMO oscillates between positive (warm) and negative (cold) phases (Enfield et al., 2001) with a periodicity of 65–80 years (Schlesinger and Ramankutty, 1994; Kerr, 2000; Enfield et al., 2001). Shifts in the AMO phase have been linked to changes in precipitation (Enfield et al., 2001) and frequency of landfalls of major hurricanes in North America (Goldenberg et al., 2001). However, the short duration of instrumental SST records has only allowed for the direct documentation of about 1½ AMO cycles, with the early portion of the instrumental record based on limited observations (Alexander et al., 2014). Because of these limitations, the actual

periodicity and persistence of the AMO before the 20th century remains uncertain (see Kilbourne et al., 2014). The forcing for the AMO is unknown, but several studies indicate that the decadal-scale changes in North Atlantic SST involve fluctuations due to internal variations in oceanic heat transport, specifically AMOC (Delworth and Mann, 2000; Collins and Sinha, 2003; Sutton and Hodson, 2003; Knight et al, 2005; Buckley and Marshall, 2016).

In this study we present two new coral Sr/Ca-SST records from DTNP with well-constrained chronologies and approximately monthly sampling resolution. These new records are combined with published records of Sr/Ca-SST from DTNP (DeLong et al., 2014) to generate a stacked Sr/Ca-SST record that incorporates five distinct coral colonies and two different species (*Siderastrea siderea* and *Orbicella faveolata*). This new record is situated in a region that is sensitive to the AMO and AMOC variations, and, therefore, provides insight into the link between the two systems.

2. Methods

2.1 Coral Collection

The United States Geological Survey (USGS) collected the coral cores used in this study in the area of Pulaski Shoal at the northern end of DTNP in 2008 (*S. siderea* colonies A* and F1*; *O. faveolata* colony B3) and 2012 (*S. siderea* colony CG2) (Figure 1, inset). An additional *S. siderea* colony (93DRYSS2*) was collected from DTNP in 1993. Details of field operations are provided in Maupin et al. (2008), DeLong et al. (2011), Hickey et al. (2013), Flannery and Poore (2013), and Weinzierl et al. (2016).

2.2 Chronology

All coral cores were slabbed along the major vertical growth axis and underwent a nondestructive cleaning treatment using a Branson Sonifer 450 sonication device. X-radiographs of each coral slab taken at the United States Geological Survey Coastal and Marine Science Center in St. Petersburg, Florida, USA, revealed annual couplets of high- and low- density bands. Previous work in the Florida Keys (Hudson et al., 1976) on *Orbicella* (formerly *Montastraea*) spp. demonstrated that each couplet of high- and low- density bands equates to an annual growth band, with the high-density bands forming in late summer or early fall (the time with warmest SSTs). An age model for *O. faveolata* colony B3 was developed by counting annual high-density bands identified in the x-radiographs backwards from the year of coral collection, 2008, and cross-checked by overlapping the raw Sr/Ca records onto the x-radiographs as described in Flannery and Poore (2013). This method produced an age model for colony B3 that extended back to 1893 (Figures S1 and S2).

Siderastrea siderea grows slower than *O. faveolata* (DeLong et al., 2011). X-radiographs of the *S. siderea* CG2 coral slabs show annual density bands, but they are less defined than *O. faveolata* bands and therefore difficult to distinguish. A preliminary age model was constructed for *S. siderea* CG2 using the annual density bands, and the final age model was constructed using the geochemistry record by counting the annual Sr/Ca cycles backwards from 2012 (the year of coral collection). This procedure produced an age model for CG2 that ranged from 1837–2012. The chronology was cross-checked and verified by overlapping the geochemistry onto the x-radiograph (Figure S3), following the procedure of Flannery and Poore, 2013.

Scanning electron microscopy (SEM) analysis (performed at the University of South Florida on a Hitachi SEM) on select intervals (particularly intervals with notable Sr/Ca excursions, Figures 2, S1, and S2) from coral *O. faveolata* colony B3 and *S. siderea* colony CG2 revealed no evidence of diagenetic alteration (Figures S3 and S4) that may affect the Sr/Ca records stored within. Some secondary precipitation of aragonite needles appears in the pore spaces between thecal walls (Figures S3 and S4), but proper sampling techniques avoided these areas and only removed material from pristine thecal walls.

2.3 Sr/Ca Determinations

Subsampling of both coral species was done using a tri-axial micromill. Preliminary analysis of the *O. faveolata* colony B3 x-radiographs (Flannery and Poore, 2013; Figure S1) suggested an average linear extension rate of $\sim 8.5 \pm 1.9 \text{ mm yr}^{-1}$ (1σ , $n = 115$) throughout the entire coral, with the lowest linear extension rate of 4.5 mm yr^{-1} occurring in 1969. In order to achieve the highest sampling resolution while still obtaining enough powder for analysis, we removed skeletal samples of *O. faveolata* B3 along a continuous thecal wall with a 0.5 mm drill-bit and 0.5 mm forward increment. Each density band couplet (representing one year of coral growth) yielded 15–20 skeletal powder samples, resulting in greater than monthly coral Sr/Ca resolution.

The annual banding on the *S. siderea* colony CG2 x-radiographs indicated an average linear extension rate of $\sim 5 \text{ mm yr}^{-1}$. To ensure that we captured the full annual cycle, we divided 5 mm by 12 (the average number of samples needed to capture ~monthly resolution) to obtain a forward increment of 0.417 mm.

Between 88 and 237 μg of subsampled skeletal powder was weighed out for each sample from both *O. faveolata* colony B3 and *S. siderea* colony CG2 and was acidified using ultrapure 2% HNO_3 in order to obtain sample solutions with a target Ca concentration of 20 ppm. A PerkinElmer 7300 dual view inductively coupled plasma-optical emission spectrometer (ICP-OES) was used to make the Sr/Ca determinations. Sr/Ca was corrected for instrumental drift using the internal gravimetric standard (IGS) method devised by Schrag (1999). The average corrected IGS precision for Sr/Ca was 0.01 mmol mol^{-1} (1σ , $n=303$) prior to August, 2013 (when a new IGS standard was created) and 0.008 post-2013 (1σ , $n=464$). A second in-house standard (homogenized powder from a *Porites lutea* coral specimen) analyzed for Sr/Ca to test for any potential matrix effects had an average corrected precision of 0.02 mmol mol^{-1} (1σ , $n=1104$). These precisions are comparable with the long-term ICP-OES analytical precision for this instrument. In order to assess how our Sr/Ca determinations compare to those made in other laboratories, we also analyzed the JCp-1 standard used in an inter-laboratory comparison study of Sr/Ca (Hathorne et al., 2013). The average Sr/Ca value determined for the JCp-1 standard on our ICP-OES was $8.823 \text{ mmol mol}^{-1} \pm 0.019$ ($n = 50$, 2σ), which is within analytical error of the mean ($8.831 \text{ mmol mol}^{-1}$) reported in Hathorne et al. (2013).

Measured Sr/Ca data were converted from depth to time using the default settings in Analyseries 2.0.4.2 (Paillard et al., 1996) for *O. faveolata* B3 (1893–2008) and *S. siderea* CG2 (1837–2012) (Figure 2).

2.4 *In situ* SST Data

185 National Oceanic and Atmospheric Administration (NOAA) National Data Buoy
186 Center (NDBC, <http://www.ndbc.noaa.gov>; NOAA, 2015) provides twice-hourly SST
187 data from DTNP at two stations: DRYF1 (1992–2005 CE) and PLSF1 (2008–2010 CE)
188 (Figure 1); however, the buoy data contain temporal gaps due to equipment failure, which
189 limits the data available for the present study to only 6 (1995, 1996, 1998, 2001, 2003,
190 2005) complete years.

191 USGS personnel have been monitoring SST at Pulaski Shoal from 2009 to present
192 using Onset® HOBO® Water Temp Pro V2 (U22-001) temperature loggers placed at 5
193 m water depth, adjacent to the coral colonies used in this study. The logger monthly
194 temperature records show that SST at DTNP has a 12 °C seasonal range, with summers
195 exceeding 31 °C and winters falling to 19 °C (Figure S5; Kuffner, 2015). Several gridded
196 SST data product sets are also available for the DTNP area. We compared the
197 HADISST1 (Rayner et al., 2003) data set with a 1° x 1° box centered on 82 °W and 24 °N
198 (1870–2013) to the *in situ* temperature loggers (2010–2014). The HADISST1 data set for
199 the DTNP region does not capture the annual winter SST minima recorded by the
200 corals/*in situ* temperature loggers. The average winter minimum *in situ* logger
201 temperature is lower than the HADISST1 data by 2.23 °C (21.66 °C and 23.89 °C,
202 respectively) (Figure 2B), over the complete record, and is 1.35 °C lower than the
203 HADISST1 dataset over the common age interval (2010–2013). Conversely, the average
204 summer maximum *in situ* logger temperature is on average 0.62° C warmer than the
205 HADISST1 when all data is used (29.74 °C and 30.36 °C, respectively) (Figure2B) and
206 0.17 °C cooler than the HADISST1 data over the common age interval (2010–2013).

When interpreting the summer and winter Sr/Ca-SST extremes, we use the values generated from the *in situ* temperature logger (Figure 2B).

2.5 SST Estimates

The Sr/Ca records were converted to SST (Figures 2 and 3) using previously-published calibration equations specific to each colony and species (*O. faveolata*, Flannery and Poore, 2013; *S. siderea*, LH SST calibrated equation, Table 4, DeLong et al., 2014). The equation for *O. faveolata* is $\text{Sr/Ca} = -0.0392 \times \text{SST} + 10.205$ (Flannery and Poore, 2013) and the equation for *S. siderea* is $\text{Sr/Ca} = -0.043 \times \text{SST} + 10.11$ (LH SST calibrated equation, Table 4, DeLong et al., 2014). DeLong et al. (2011, 2014) produced their calibration equation using *Siderastrea siderea* colony A collected from DTNP in August 2008. The DRYF1 and PLSF1 combined data sets were used for the calibration of Sr/Ca to SST in Flannery and Poore (2013) and DeLong et al. (2011; 2014). Mean annual Sr/Ca-SST was determined by averaging all Sr/Ca-SST for a given calendar year (Figure 2).

2.6 Statistical Analysis

The Sr/Ca-SST records produced in this study using *O. faveolata* and *S. siderea* were combined with coral-based (*S. siderea*) Sr/Ca-SST records from DeLong et al. (2014) into a stacked Sr/Ca-SST record spanning 1733–2012 using a Monte Carlo routine modified from the algorithm used in Thirumalai et al. (2016). We incorporated analytical and calibration uncertainty (*S. siderea*, DeLong et al. 2014; *O. faveolata*, Flannery and Poore, 2013) in a Monte Carlo framework (n=5000) while converting the Sr/Ca records to SST, prior to stacking them. A median SST stack along with corresponding 32nd and

68th percentile error bounds was produced. Frequency analysis using the multi-taper method (number of tapers = 3) was performed using KSpectra version 3.4.3 with a red noise AR1 process to test for confidence at the 90% level. All data produced in this study is available in Flannery et al. (2016).

3. Results and Discussion

3.1 New Sr/Ca-SST estimates from DTNP

The Sr/Ca and derived SST records for *O. faveolata* colony B3 and *S. siderea* colony CG2 are shown in Figure 2A. Raw Sr/Ca between the two coral species is offset such that Sr/Ca from *O. faveolata* colony B3 is, on average, 0.2 mmol mol⁻¹ higher than the Sr/Ca from *S. siderea* colony CG2. This offset reflects different partitioning of Sr during calcification in the two species, a “vital effect” noted previously by DeLong et al. (2011). Other studies stress caution when interpreting Sr/Ca-based SST relationships (Alpert et al., 2016) due to vital effects, but we circumvent this issue by independently calibrating each coral colony to local C-MAN buoy SST. In the case of *S. siderea* at DTNP, we found that the calibration of CG2 to local C-MAN buoy SST resulted in a Sr/Ca-SST calibration within error of the LH equation established by DeLong et al. (2014), and therefore we apply the DeLong et al. (2014) LH equation to all *S. siderea* Sr/Ca data in this study.

The average summer SST maximum for the *O. faveolata* colony B3 record is 29.4 °C (±1.3 °C), and 29.1 (±1.3 °C) for the *S. siderea* colony CG2 record, which is consistent with available *in situ* instrumental measurements (1997–2003; 2009–2015; see Figure S5) and the gridded HADISST from DTNP (Figure 2B). Sr/Ca-SST estimates that reach or

exceed 31 °C are also realistic as the observed maximum daily SST at DTNP was > 31 °C in 1998, 2000, 2011 and 2014 according to the *in situ* instrumental temperature logger measurements. In each record there are three years in which Sr/Ca-SST estimates exceed 32 °C. While it is unlikely that monthly mean SST exceeded 32 °C, these values are analytically sound and within the calibration uncertainty of observed monthly maximum SST at DTNP. Sr/Ca-SST records from both *O. faveolata* colony B3 and *S. siderea* colony CG2 exhibit decadal intervals of reduced summer SST during the 1930s and 1960s; an interval of reduced summer SST maxima is also present in *S. siderea* colony CG2 around 1840 (Figure 2A).

The average winter monthly SST minima for the *O. faveolata* colony B3 and *S. siderea* colony CG2 is 21.4 °C (± 1.6 °C) and 20.4 °C (± 1.3 °C), respectively; the coldest years occur in 1939 and 1973, coinciding with El Niño events recorded for those years (Poore et al., 2001). Comparison of *in situ* observations at DTNP with HADISST1 show the SST gridded data products capture summer maxima at DTNP fairly well, but underestimate winter SST minima by ~3 °C (DeLong et al., 2011; Flannery et al., 2013). Overall, the *S. siderea* monthly winter SST minima are 1 °C cooler than the *O. faveolata* B3 estimates (Figure 2B). It is possible that this results from calibration uncertainties, or systematic bias in seasonal growth rates between the two species. When performing additional treatment of the data, we did not remove any anomalously “high” or “low” Sr/Ca-SST values.

Where the *O. faveolata* colony B3 and *S. siderea* colony CG2 Sr/Ca-SST records overlap, the average offset in their mean annual Sr/Ca-SST estimates is 0.37 °C, but with divergences of up to ± 3 °C in mean annual SST. One such interval of divergence occurs

1966–1974, when both coral colonies record the coolest decade in the 20th century– but *O. faveolata* colony B3 Sr/Ca-SST averages 1.7 °C cooler than the *S. siderea* colony CG2 Sr/Ca-SST. Conversely, during the interval 1991–1995 *O. faveolata* colony B3 shows mean annual Sr/Ca-SSTs that are on average 2.3 °C warmer than those of *S. siderea* CG2. We hypothesize that slowed or halted calcification of *S. siderea* due to thermal stress (warm or cold) may result in *S. siderea*-based Sr/Ca-SST reconstructions failing to record seasonal temperature extremes during anomalously cold/warm intervals. For example, during 1991–1995 *O. faveolata* B3 records the warmest mean annual SST (28.04 °C) for any 5-year interval in the 20th century, and summer maxima exceeded 30 °C for each of those years in the *O. faveolata* colony B3 record. During that warm interval (as recorded *O. faveolata* colony B3), linear extension rates in the *S. siderea* CG2 are anomalously low (Figure S6), approaching the low threshold (2.1 mm yr⁻¹) for recording reliable mean annual Sr/Ca-SST established by DeLong et al. (2016). We interpret the most recent part (1990–2011) of the CG2 record with caution, as the average extension rate falls to 3.9 mm yr⁻¹, effectively reducing the sampling resolution from 12 to 9 samples per year. Despite discrepancies in interannual variability observed in the *O. faveolata* colony B3 and *S. siderea* colony CG2 Sr/Ca-SST records, similar low-frequency (decadal to multidecadal) variability suggests that the low-frequency trends represent a regional signal.

3.2 Intercolony Comparison at DTNP

With the addition of the new records produced in this study, there are now mean annual Sr/Ca-SST time series from five individual colonies of two different species at

DTNP (*O. faveolata* B3 and *S. siderea* colonies CG2, A, F1, and 93DRYSS2)(Figure 3A). Comparing inter-annual variability between the five different colonies we only find significant correlation between the GC2 and A2 *S. siderea* colonies ($r=0.34$, $p<0.01$), which overlap from 1837-1955. This suggests that there is a lot of noise in the mean annual SST estimates from any given colony calcifying at DTNP. All of the corals examined in this study were obtained from a relatively small geographic area and from similar water depths; therefore, we do not think the differences are related to hydrographic variability. It is likely that changes in growth rate related to thermal stress (either warm or cold), predation, nutrient availability, or other ecological factors may be the cause of the heterogeneity in the SST signal among these coral colonies at DTNP. For example, extension rates of *S. siderea* CG2 are anomalously low between 1990 and 2008, resulting in lower resolution sampling (as low as ~5 samples per annual cycle in 1995) compared with the older part of the record. During this interval, we see a large divergence between the B3 and CG2 SST records. Therefore, we suspect mean annual SSTs after 1990 in the *S. siderea* CG2 record (stippled section in Figure 3A) have been biased by undersampling the annual cycle. Despite these discrepancies between individual colonies, there is coherency among the suite of DTNP cores at the decadal to multidecadal scale. This underscores the value of composite records, that have the ability to average out these highly localized and ecological overprints on the regional signal.

3.3 A three-century multi-species composite SST reconstruction from DTNP

We developed a new, stacked Sr/Ca-SST reconstruction for DTNP by combining our Sr/Ca-SST estimates from *O. faveolata* colony B3 and *S. siderea* colony CG2 with

published Sr/Ca-SST from *S. siderea* colonies A (A1 and A2), F1, and 93DRYSS2 in the DTNP from DeLong et al. (2014). The stacked Sr/Ca-SST record (Figure 3B) shows that mean annual Sr/Ca-SST at DTNP ranged from a minimum of 23.74 °C in 1965 to a maximum of 27.63 °C in 2011 (3.89 °C range) during the last 278 years. In contrast with the reconstruction of DeLong et al. (2014), the updated DTNP Sr/Ca-SST reconstruction does not show a secular warming trend over the last century, but does show a pronounced 0.8 °C multidecadal warming trend from 1970–2012. This is consistent with the findings of Kuffner et al. (2015), which show that the 20th century SST warming trend in the Florida Keys can be attributed almost entirely to 0.8 °C of warming from 1975–2007. The magnitude of this warming is double the 0.4 °C warming observed over the same time interval in the western Atlantic Ocean coral-based reconstruction of Tierney et al. (2015).

The oldest section of the DTNP stacked record coincides with the end of the Little Ice Age (LIA, 1400–1850 CE). Several studies in the Caribbean/GOM region demonstrate that SSTs in the AWP were generally cooler during the LIA, but also variable, with distinct cool intervals alternating with more moderate conditions (Richey et al., 2009). Several such decadal-scale cooling events occur in the DTNP Sr/Ca-SST reconstruction centered around 1740, 1790, 1820 and 1845. Aside from the cooling centered at 1965, these four intervals are the coolest decadal excursions in the three-century reconstruction, and likely represent a regional-scale cooling event associated with the end of the LIA. These four decadal cooling events correspond with four of the coolest decades observed in the 450-year western Atlantic coral SST reconstruction (Tierney et al., 2015), however there is no secular warming trend from the end of the LIA to the 20th century in the DTNP SST reconstruction.

The cool interval at DTNP around 1935 coincides with cool SST anomalies in the southwestern GOM (del Monte-Luna et al., 2015). This regional cooling also corresponds with a major drought (the Dust Bowl) centered over the central Great Plains (USA). The Dust Bowl was a complicated event partly related to human activities, but also a major drought interval that lasted for about 9 years (Cook et al., 2008). Several studies (e.g., Schubert et al., 2004; Sutton and Hodson, 2005) suggest that decadal-scale variations in tropical Atlantic Ocean mean annual SSTs influence precipitation variability over the North American mid-continent and elsewhere along the North Atlantic margin. During the Dust Bowl (1932–1938), warm tropical Atlantic SST anomalies correspond with lower amounts of precipitation over continental North America, leading to dry conditions (Schubert et al., 2004). The warm, tropical SSTs observed during this interval are thought to result in a cyclonic circulation anomaly in the lower troposphere that inhibits transport of moisture northward from the GOM into the central United States (Schubert et al., 2004).

Observational and modeling data show that the precipitation deficit during the peak of the Dust Bowl drought (1932–1938) occurred primarily in the spring and summer (Schubert et al., 2004). As noted above, SSTs in the southwestern GOM (del Monte-Luna et al., 2015) and the Dry Tortugas area were cool during the Dust Bowl interval. Moreover, our study shows that lower mean annual SST in the Dry Tortugas area at this time was driven by lower summer (warm season) temperatures (Figure 2B).

The most prominent cool event in the DTNP Sr/Ca-SST record occurs during the decade centered around 1965, and coincides with a previously-documented period of cooling in the North Atlantic Ocean (Hodson et al., 2014). A cooler North Atlantic during

the 1960s to the 1970s is evident in a number North Atlantic SST data sets (e.g., Baines and Folland, 2007; Thompson et al., 2010) as well as North American tree-ring based temperature reconstructions (Jacoby and D'Arrigo, 1989)(for further discussion of the 1960s cooling see section 3.5).

3.4 Atlantic Warm Pool Variability

During the LIA, proxy records throughout the Caribbean and GOM region including the Cariaco Basin, Jamaica, Puerto Rico, and the northern GOM suggest that mean annual SST was between 1 °C and 3 °C lower than modern values (Richey et al., 2009). Based on the wide geographic distribution of low mean annual SST, Richey et al. (2009) suggest that the spatial extent and intensity of the AWP was reduced during the LIA. Because the AWP is defined by SSTs at or exceeding 28.5 °C, summer SSTs lower than 28.5 °C may indicate reduced or no expansion of the AWP into the Florida Straits.

The sub-annually resolved coral records from DTNP support the interpretation that a reduced or less intense AWP was associated with cool events (i.e., around 1845, 1930s and 1960s) in DTNP over the last few hundred years. With the exception of these cool intervals, the monthly maximum summer Sr/Ca-SST estimates from DTNP corals *S. siderea* colony CG2 and *O. faveolata* colony B3 (Figure 2) are consistently at or above 29 °C for the last 174 years. In contrast, during the major cool intervals the monthly summer maxima SST estimates are about 2° C lower (at or below 28 °C), a pattern that is particularly obvious in *O. faveolata* colony B3. The lower summer SSTs identified in the DTNP record in the 1930s and 1960s coincide with lower SSTs anomalies in the southwestern GOM (del Monte-Luna et al., 2015), which supports our conclusion that the

AWP was weakly-developed during those times. Similarly, the lower monthly summer maxima Sr/Ca-SST at DTNP during the major cool event in *S. siderea* CG2 centered around 1845 indicates the AWP did not extend to DTNP at that time, consistent with reduced AWP area during the latter portion of the LIA.

Additionally, an abrupt shift to cooler temperature occurs in the Sr/Ca-SST composite reconstruction in 2000. Monthly summer Sr/Ca-SST maxima associated with the possible cool event from 2000–2010 do not show a consistent pattern. Monthly summer maxima Sr/Ca-SST in *S. siderea* CG2 are at or above 29 °C during the first part of the interval (2000–2006) but below 28 °C for the second part of the interval (2006–2010). This reduction in mean annual Sr/Ca-SST between 2006 and 2010 may reflect a sampling bias towards winter months due to a reduction in linear extension rates during this time (see Figure S6).

3.5 Decadal-Scale Variability and the AMO

Comparison of the DTNP Sr/Ca-SST reconstruction with the AMO index of Enfield et al. (2001) reveals similar multidecadal structure in the data sets during the 20th Century. The maximum correlation between the two records occurs when the AMO index is lagged by 9 years behind the DTNP SST record ($r = 0.28$, $p < 0.001$) (Figure 4B). The strongest example of coherence between the AMO and DTNP SST is the cooling trend in the stacked Sr/Ca-SST reconstruction from 1948 to 1965 (Figure 4B), and subsequent warming into the 1990's. We performed a 10-year running correlation between the 9-year lagged AMO index (Enfield et al., 2001) and the stacked Sr/Ca-SST reconstruction (this study) from 1856–2008 (the period of overlap between the records). Although the AMO

index and the coral Sr/Ca-SST reconstruction are significantly correlated throughout the interval of overlap, several decadal intervals of the record yield higher positive correlations ($r > 0.7$, $p < 0.001$, Figure 4A), with the interval centered on 1970 displaying the highest correlation ($r > 0.8$, $p < 0.001$). The divergence of the AMO index and the coral Sr/Ca-SST record after 1990 may be the result of anthropogenic overprints on the climate signal. We note that the oldest portion of the AMO index is derived from the Kaplan et al., (1998) data set, which has lower spatial and temporal resolution and may not adequately reflect AMO variability.

We performed frequency analysis (MTM, $n =$ three tapers) of our stacked Sr/Ca-SST reconstruction to test for the persistence of significant multidecadal oscillations in SST in the Florida Straits. Our analyses (Figure 4C) revealed significant periodicities (90% confidence level) of 28–30 years and 80–90 years. The multispecies composite SST reconstruction presented in this study is one of few annually resolved SST proxy records in the North Atlantic that extends back in time beyond the 19th Century, and can address the question of whether the AMO persisted beyond the observational period. The coherence between the coral-based SST reconstruction and AMO index suggests that SST in the Florida Straits is sensitive to the AMO on multidecadal timescales, and the frequency analysis reveals that this variability persists for the past three centuries. Significant multi-decadal oscillations were also found in a coral growth-rate based SST reconstruction from the Bahamas over the same time interval (Saenger et al., 2009).

We also note that significant variability with a period of ~30 years has been found in several additional proxy records sensitive to the AMO. MTM analysis of a Mg/Ca-SST record from the Pigmy Basin, northern GOM (Richey et al., 2007) revealed a significant

33-to 34-year cycle, attributed to the influence of the AMO (Poore et al., 2009). MTM analysis of the Gray et al. (2004) Northern Hemisphere tree-ring reconstruction shows a significant cycle of ~40 years, which was also ascribed to AMO variability (Poore et al., 2009).

During the cooling event of the 1960s, Hodson et al. (2014) found a lag between oceanic cooling in the northern extension of the Gulf Stream and a negative AMO (see Hodson et al. 2014, Figure 5). Rather than homogeneous cooling throughout the North Atlantic, the northern extension of the Gulf Stream cools first, followed by the SubPolar Gyre, and then the tropical Atlantic. The AMO index and cooling in the Sub Polar Gyre and tropical Atlantic lag the northern Gulf Stream cooling by ~ 6–9 years (Hodson et al., 2014; Knudsen et al., 2014). Our results, along with the observations of Hodson et al. (2014), support the interpretation (e.g., Delworth and Mann, 2000; Collins and Sinha, 2003; Sutton and Hodson, 2003; Knight et al., 2005) that the AMO is linked to changes in Atlantic meridional overturning circulation (AMOC). Increased transport of Caribbean waters by the Florida Current and then Gulf Stream into the North Atlantic Ocean leads to warming in the Northern Atlantic, but it takes about 9 years for the warming to propagate throughout the entire basin and to be recorded in the AMO index. Decreased AMOC leads to a reversal of this process. Our results do not address the cause of the change in AMOC but do indicate that the AMO and AMOC are closely linked.

4. Summary and Conclusions

We developed two new monthly-resolved coral Sr/Ca-based SST records from an *Orbicella faveolata* (1893–2008) colony and a *Siderastrea siderea* colony (1837–2012) from Dry Tortugas National Park in the Florida Straits. We combined these new records with three previously-published *S. siderea* Sr/Ca-SST records from DTNP to develop a 278 year-long multi-species, annually resolved composite SST reconstruction.

The early part of the reconstruction (1733-1850) coincides with the end of the Little Ice Age, and exhibits 3 of the 4 coolest decadal excursions in the record. However, the mean SST estimate from that interval during the LIA is not significantly different from the late 20th Century SST mean. The most prominent cooling event in the 20th Century is a decade centered around 1965. This corresponds to a basin-wide cooling in the North Atlantic and cool phase of the AMO. During the Dust Bowl (1932-1938), an interval of persistent drought in the central United States, the DTNP record indicates a modest cooling in mean annual SST. The Dust Bowl cooling is distinct from other decadal cooling events because it is dominated by a reduction of summer maximum SST by 2 °C relative to the climatic mean, observed in both *O. faveolata* and *S. siderea*.

The multidecadal trends in SST at DTNP are significantly correlated with the AMO index, with the AMO lagging the DTNP SST record by 9 years. The covariance between SST at DTNP and the AMO index suggests a link between low frequency SST variability in the Florida Straits and the greater North Atlantic. Spectral analysis of the DTNP SST reconstruction over the last three centuries highlights significant periodicity in the 80–90 year range, suggesting that the AMO pattern persists beyond the instrumental record.

Acknowledgements

We thank Don Hickey, Chris Reich, B.J. Reynolds, and Jordan Sanford from the United States Geological Survey St. Petersburg Coastal and Marine Science Center, the M/V *Fort Jefferson* crew, and Dry Tortugas National Park personnel for field assistance during this study. Daniel Umberger, Adis Muslic, Tara Roeder, and Kelsey Roberts from the USGS St. Petersburg Coastal and Marine Science Center provided assistance with sample preparation. Tony Greco provided scanning electron microscope assistance. We thank Christopher Maupin from Texas A&M University and Lauren Toth from the USGS St. Petersburg Coastal and Marine Science Center for discussions that improved the manuscript. We appreciate helpful reviews from Brad Rosenheim and an anonymous reviewer. This research was funded jointly by the USGS Climate and Land Use Research & Development Program and the Coastal and Marine Geology Program. The study was conducted under scientific research permits DTNP-2008-SCI-0015 and DTNP-2012-SCI-0001. Any use of trade names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

References

- Alexander M. A., K. H. Kilbourne and J. A. Nye (2014), Climate variability during warm and cold phases of the Atlantic Multidecadal Oscillation (AMO) 1871–2008. *J. Mar. Syst.*, 133, 14-26, doi:10.1016/j.jmarsys.2013.07.017
- Alpert, A. E., A. L. Cohen, D. W. Oppo, T. M. DeCarlo, J. M. Gove, and C. W. Young (2016), Comparison of equatorial Pacific sea surface temperature variability and trends with Sr/Ca records from multiple corals, *Paleoceanography*, 31, doi:10.1002/2015PA002897.

506 Baines, P. G., and C. K. Folland (2007), Evidence for a rapid global climate shift across
 507 the late 1960s, *Journal of Climate*, 20(12), 2721-2744, doi.org/10.1175/JCLI4177.1
 508 Buckley, M. W. and J. Marshall (2016), Observations, inferences, and mechanisms of the
 509 Atlantic Meridional Overturning Circulation: A review, *Rev. Geophys.*, 54,
 510 doi:10.1002/2015RG000493
 511 Cook, B. I., R. L. Miller and R. Seager (2008), Dust and sea surface temperature forcing
 512 of the 1930's 'Dust Bowl' drought. *Geophysical Research Letters*, (35), L08710, doi:
 513 10.1029/2008GL033486
 514 Collins, M., and B. Sinha (2003), Predictability of decadal variations in the thermohaline
 515 circulation and climate, *Geophysical Research Letters*, 30(6),
 516 doi:10.1029/2002GL016504
 517 Correge, T. (2006), Sea surface temperature and salinity reconstruction from coral
 518 geochemical tracers, *Palaeogeography Palaeoclimatology Palaeoecology*, 232(2-4),
 519 408-428, doi:10.1016/j.palaeo.2005.10.014
 520 del Monte-Luna, P., H. Villalobos, and F. Arreguin-Sanchez (2015), Variability of sea
 521 surface temperature in the southwestern Gulf of Mexico, *Continental Shelf Research*,
 522 102, 73-79, doi:10.1016/j.csr.2015.04.017
 523 DeLong, K. L., J. A. Flannery, C. R. Maupin, R. Z. Poore, and T. M. Quinn (2011), A
 524 coral Sr/Ca calibration and replication study of two massive corals from the Gulf of
 525 Mexico, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 307, 117–128,
 526 doi:10.1016/j.palaeo.2011.05.005.
 527 DeLong, K. L., J. A. Flannery, R. Z. Poore, T. M. Quinn, C. R. Maupin, K. Lin, and C. C.

528 Shen (2014), A reconstruction of sea surface temperature variability in the southeastern
 529 Gulf of Mexico from 1734–2008 CE using cross-dated Sr/Ca records from the coral
 530 *Siderastrea siderea*, *Paleoceanography*, doi:10.1002/2013PA002524.

531 DeLong, K. L., C. R. Maupin, J. A. Flannery, T. M. Quinn, and C. C. Shen (2016),
 532 Refining temperature reconstructions with the Atlantic coral *Siderastrea siderea*,
 533 *Palaeogeography, Palaeoclimatology, Palaeoecology*,
 534 doi:10.1016/j.palaeo.2016.08.028.

535 Delworth, T. L., and M. E. Mann (2000), Observed and simulated multidecadal
 536 variability in the Northern Hemisphere, *Climate Dynamics*, 16(9), 661–676, doi:
 537 10.1007/s003820000075

538 Deser, C., A. S. Phillips, and M. A. Alexander (2010), Twentieth century tropical sea
 539 surface temperature trends revisited, *Geophysical Research Letters*, 37, 6,
 540 doi:10.1029/2010GL043321

541 Enfield, D. B., A. M. Mestas-Núñez, and P. J. Trimble (2001), The Atlantic multidecadal
 542 oscillation and its relation to rainfall and river flows in the continental U.S., *Geophys.*
 543 *Res. Lett.*, 28, 2077–2080, doi:10.1029/2000GL012745

544 Flannery, J. A., and R. Z. Poore (2013), Sr/Ca proxy sea-surface temperature
 545 reconstructions from modern and Holocene *Montastraea faveolata* specimens from the
 546 Dry Tortugas National Park, in *Understanding and Predicting Change in the Coastal*
 547 *Ecosystems of the Northern Gulf of Mexico, Journal of Coastal Research*, edited by
 548 J.C. Brock, J.A. Barras and S.J. Williams, pp. 20–31, Coconut Creek, Florida,
 549 doi:10.2112/SI63-003.1.

Flannery, J. A., J. N. Richey, K. Thirumalai, R. Z. Poore, and K. L. DeLong (2016),
Multi-species coral Sr/Ca based sea-surface temperature reconstruction data
using *Orbicella faveolata* and *Siderastrea siderea* from Dry Tortugas National Park,
Florida: U.S. Geological Survey data release, <http://dx.doi.org/10.5066/F7CC0XS9>.

Goldenberg, S. B., C. W. Landsea, A. M. Mestas-Nunez, and W. M. Gray (2001), The
recent increase in Atlantic hurricane activity: Causes and implications, *Science*,
293(5529), 474-479, doi: 10.1126/science.1060040

Gray, S. T., L. J. Graumlich, J. L. Betancourt, and G. T. Pederson (2004), A tree-ring
based reconstruction of the Atlantic Multidecadal Oscillation since 1567 A.D.
Geophysical Research Letters, 31:L12205, doi:10.1029/2004GL019932.

Hathorne, E. C., A. Gagnon, T. Felis, J. Adkins, R. Asami, W. Boer, N. Caillon, D. Case,
K. M. Cobb, E. Douville, P. deMenocal, A. Eisenhauer, D. Garbe-Schönberg, W.
Geibert, S. Goldstein, K. Huguen, M. Inoue, H. Kawahata, M. Kölling, F. L. Cornec, B.
K. Linsley, H. V. McGregor, P. Montagna, I. S. Nurhati, T. M. Quinn, J. Raddatz, H.
Rebaubier, L. Robinson, A. Sadekov, R. Sherrell, D. Sinclair, A. W. Tudhope, G. Wei,
H. Wong, H.C. Wu, and C. You (2013), Interlaboratory study for coral Sr/Ca and other
element/Ca ratio measurements, *Geochem. Geophys. Geosyst.*, 14, 3730–3750,
doi:10.1002/ggge.20230.

Hickey, D., C. D. Reich, K. L. DeLong, R. Z. Poore, and J. C. Brock (2013), Holocene
core logs and site methods for modern reef and head-coral cores: Dry Tortugas
National Park, Florida, U.S. Geological Survey Open-File Report 2012–1095, 27 pp,
Dept. of the Interior, U.S. Geological Survey, Washington DC.,
<http://pubs.usgs.gov/of/2012/1095>.

573 Hodson, D. L. R., J. I. Robson, and R. T. Sutton (2014), An Anatomy of the Cooling of
 574 the North Atlantic Ocean in the 1960s and 1970s, *Journal of Climate*, 27(21), 8229-
 575 8243, doi.org/10.1175/JCLI-D-14-00301.1
 576 Hudson, J. H., E. A. Shinn, R. B. Halley, and B. Lidz (1976), Sclerochronology: A tool
 577 for interpreting past environments, *Geology*, v. 4, p. 361-364, doi:10.1130/0091-
 578 7613(1976)
 579 Jacoby, G. C. and R. D. D'Arrigo (1989), Reconstructed northern hemisphere annual
 580 temperature since 1671 based on high-latitude tree-ring data from North America,
 581 *Climatic Change* 14:39-59, doi:10.1007/BF00140174
 582 Kaplan, A., M. A. Cane, Y. Kushnir, A. C. Clement, M. B. Blumenthal, and B.
 583 Rajagopalan (1998), Analyses of global sea surface temperature 1856-1991, *Journal of*
 584 *Geophysical Research-Oceans*, 103(C9), 18567-18589, doi:10.1029/97JC01736
 585 Kerr, R. A. (2000), A North Atlantic climate pacemaker for the centuries, *Science*,
 586 288(5473), 1984-1986 doi: 10.1126/science.288.5473.1984
 587 Kerr, R. A. (2005), Climate change - Atlantic climate pacemaker for millennia past,
 588 decades hence?, *Science*, 309(5731), 41-+, doi: 10.1126/science.309.5731.41
 589 Kilbourne K. H., M. A. Alexander and J. A. Nye (2014), A low latitude paleoclimate
 590 perspective on Atlantic multidecadal variability. *J. Marine Syst.*, 133, 4-13.
 591 doi:10.1016/j.jmarsys.2013.09.004
 592 Knight, J. R., R. J. Allan, C. K. Folland, M. Vellinga, and M. E. Mann (2005), A
 593 signature of persistent natural thermohaline circulation cycles in observed climate,
 594 *Geophysical Research Letters*, 32(20), doi:10.1029/2005GL024233

595 Knudsen, M. F., B. H. Jacobsen, M.-S. Seidenkrantz, and J. Olsen (2014), Evidence for
 596 external forcing of the Atlantic Multidecadal Oscillation since termination of the Little
 597 Ice Age, *Nature Communications*, 5, doi:10.1038/ncomms4323
 598 Kuffner, I. B. (2015), Underwater temperature on off-shore coral reefs of the Florida
 599 Keys, U.S.A., 2009 to 2015, U.S. Geological Survey data release, doi:
 600 10.5066/F71C1TZK
 601 Kuffner, I. B., B. H. Lidz, J. H. Hudson, and J. S. Anderson (2015), A Century of Ocean
 602 Warming on Florida Keys Coral Reefs: Historic In Situ Observations, *Estuaries and*
 603 *Coasts*, 38(3), 1085-1096, doi:10.1007/s12237-014-9875-5
 604 Maupin, C. R., T. M. Quinn, and R. B. Halley (2008), Extracting a climate signal from
 605 the skeletal geochemistry of the Caribbean coral *Siderastrea siderea*, *Geochemistry,*
 606 *Geophysics, Geosystems*, doi:10.1029/2008GC002106.
 607 NOAA (2015), National Data Buoy Center. URL: <http://www.ndbc.noaa.gov>; accessed
 608 on January 4, 2015.
 609 Paillard, D., L. Labeyrie, and P. Yiou (1996), Macintosh program performs time-series
 610 analysis, *Eos Trans. AGU*, 77, 379, doi:10.1029/96EO00259.
 611 Poore, R. Z., J. Darling, H. J. Dowsett, and L. Wright (2001), Variations in River Flow to
 612 the Gulf of Mexico: Implications for Paleoenvironmental Studies of Gulf of Mexico
 613 Marine Sediments, *USGS Bulletin* 2187, <http://pubs.usgs.gov/bulletin/b2187/title.html>
 614 Poore, R. Z., K. L. DeLong, J. N. Richey, and T. M. Quinn (2009), Evidence of
 615 multidecadal climate variability and the Atlantic Multidecadal Oscillation from a Gulf
 616 of Mexico sea-surface temperature-proxy record, *Geo-Marine Letters*, 29(6), 477-484,
 617 doi:10.1007/s00367-009-0154-6

618 Rayner, N. A., D. E. Parker, E. B. Horton, C. K. Folland, L. V. Alexander, D. P. Rowell,
 619 E. C. Kent, and A. Kaplan (2003), Global analyses of sea surface temperature, sea ice,
 620 and night marine air temperature since the late nineteenth century *J. Geophys. Res.* Vol.
 621 108, doi:10.1029/2002JD002670

622 Richey, J. N., R. Z. Poore, B. P. Flower, and T. Quinn (2007), 1400 yr multiproxy record
 623 of climate variability from the northern Gulf of Mexico, *Geology*, 35(5), 423-426,
 624 doi:10.1130/G23507A.1

625 Richey, J. N., R. Z. Poore, B. P. Flower, T. M. Quinn, and D. J. Hollander (2009),
 626 Regionally coherent Little Ice Age cooling in the Atlantic Warm Pool, *Geophysical*
 627 *Research Letters*, 36, doi:10.1029/2009GL040445

628 Schlesinger, M. E., and N. Ramankutty (1994), An oscillation in the global climate
 629 system of period 65–70 years, *Nature*, 367(6465), 723-726, doi:10.1038/367723a0

630 Schrag, D. P. (1999), Rapid analysis of high-precision Sr/Ca ratios in corals and other
 631 marine carbonates, *Paleoceanography*, 14, 97-102, doi:10.1029/1998PA900025.

632 Smith, S. V., R. W. Buddemeier, R. C. Redalje, and J. E. Houck (1979), Strontium-
 633 calcium thermometry in coral skeletons, *Science*, 204, 404-406,
 634 doi:10.1126.science.204.4391.404.

635 Smith, J. M., T. M. Quinn, K. P. Helmle, and R. B. Halley (2006), Reproducibility of
 636 geochemical and climatic signals in the Atlantic coral *Montastraea faveolata*.
 637 *Paleoceanography*, 21, 10.1029/2005PA001187

638 Sutton, R. T., and D. L. R. Hodson (2003), Influence of the ocean on North Atlantic
 639 climate variability 1871-1999, *Journal of Climate*, 16(20), 3296-3313,
 640 doi.org/10.1175/1520-0442(2003)016<3296:IOTOON>2.0.CO;2

641 Swart, P. K., H. Elderfield, and M. J. Greaves (2002), A high-resolution calibration of
 642 Sr/Ca thermometry using the Caribbean coral *Montastraea annularis*. *Geochemistry*
 643 *Geophysics Geosystems* 3, 8402, doi: 10.1029/2002GC000306
 644 Thirumalai, K., T. M. Quinn, and G. Marino (2016), Constraining past seawater $\delta^{18}\text{O}$ and
 645 temperature records developed from foraminiferal
 646 geochemistry, *Paleoceanography*, 31, doi:[10.1002/2016PA002970](https://doi.org/10.1002/2016PA002970).
 647
 648 Thompson, D. W. J., J. M. Wallace, J. J. Kennedy, and P. D. Jones (2010), An abrupt
 649 drop in Northern Hemisphere sea surface temperature around 1970, *Nature*, 467
 650 (7314), 444-447, doi:10.1038/nature09394
 651 Tierney, J. E., N. J. Abram, K. J. Anchukaitis, M. N. Evans, C. Giry, K. H. Kilbourne, C.
 652 P. Saenger, H. C. Wu, and J. Zinke (2015), Tropical sea surface temperatures for the
 653 past four centuries reconstructed from coral archives, *Paleoceanography*, 30(3), 226-
 654 252, doi:[10.1002/2014PA002717](https://doi.org/10.1002/2014PA002717)
 655 Wang, C. Z., and D. B. Enfield (2001), The tropical Western Hemisphere warm
 656 pool, *Geophysical Research Letters*, 28(8), 1635-1638, doi: 10.1029/2000GL011763
 657 Wang, C., D. B. Enfield, S.-k. Lee, and C. W. Landsea (2006), Influences of the Atlantic
 658 warm pool on western hemisphere summer rainfall and Atlantic hurricanes, *Journal of*
 659 *Climate*, 19(12), 3011-3028, <http://dx.doi.org/10.1175/JCLI3770.1>
 660 Wang, C. Z., S. K. Lee, and D. B. Enfield (2008), Atlantic Warm Pool acting as a link
 661 between Atlantic Multidecadal Oscillation and Atlantic tropical cyclone activity,
 662 *Geochemistry Geophysics Geosystems*, 9, doi: 10.1029/2007GC001809

Weinzierl, M.S., C. D. Reich, T. D. Hickey, L. A. Bartlett, and I. B. Kuffner (2016),
Collection methods and descriptions of coral cores extracted from massive corals in
Dry Tortugas National Park, Florida, USA, U.S. Geological Survey Open-File Report
2016–xxxx, xx pp, Dept. of the Interior, U.S. Geological Survey, Washington DC.,
<http://pubs.usgs.gov/of/2016/xxxx>

Figure Captions

Figure 1. Map of detrended, mean annual SST from the HADISST1 data set centered on
the Dry Tortugas National Park (82 °W, 24 °N) correlated over 1870-2013 with grid
points all across the ocean (shown from 0 °–90 °N). Correlation coefficients range from -
0.7 to 0.7 (hatching represents 99% significance confidence interval). Inset: Map of the
Dry Tortugas National Park, Florida. The locations of coral cores are indicated on the
map by yellow stars (colonies with a * indicate previously-published data used in
conjunction with coral cores collected in this study), and National Buoy Data Center
(NBDC) stations DRYF1 and PLSF1 that monitor SST are represented by red circles. A
core was collected from the living *Orbicella faveolata* coral colony B3 at Pulaski Shoal
in August 2008 (along with *Siderastrea siderea* colony A used for comparison purposes
in this study). The core from the living *Siderastrea siderea* colony CG2 was collected in
March 2012 from Pulaski Shoal. Pulaski Shoal is approximately 7 km from NBDC buoy
DRYF1.

Figure 2. (A) Sub-annual (~monthly) Sr/Ca determinations (mmol mol^{-1}) (light pink) and mean-annual Sr/Ca values (dark pink) for *Siderastrea siderea* CG2 (1837–2012). Corresponding Sr/Ca-SST ($^{\circ}\text{C}$) values were scaled to the Sr/Ca axis using the calibration equation $[\text{Sr/Ca} = -0.43 \cdot \text{SST} + 10.11]$ from DeLong et al. (2014) and are plotted on the second y-axis. Sub-annual (~monthly) Sr/Ca determinations (mmol mol^{-1}) (light purple) and mean-annual Sr/Ca values (dark purple) for *Orbicella faveolata* B3 (1893–2008). Corresponding Sr/Ca-SST ($^{\circ}\text{C}$) values were scaled to the Sr/Ca axis using the calibration equation $[\text{Sr/Ca} = -0.0392 \cdot \text{SST} + 10.205]$ from Flannery and Poore (2013) and are plotted on the second y-axis. Sr/Ca axes are reversed so that warmer values are up. (B) Summer maximum and winter minimum Sr/Ca-SST for *S. siderea* colony CG2 (pink) and *O. faveolata* colony B3 (purple). Small dashed lines indicate the average summer and winter SSTs (29.74°C and 23.89°C , respectively) from the HADISST dataset for the Dry Tortugas (1870–2013). Large dashed lines indicate the average summer and winter SSTs (30.36°C and 21.66°C , respectively) from the *in situ* temperature logger present at DTNP (2010–2014).

Figure 3. (A) Mean-annual Sr/Ca-SST ($^{\circ}\text{C}$) values for *Siderastrea siderea* records CG2 (light pink), A1 (light gray), A2 (gray), F1 (gray), and 93DRYSS2 (dark gray) and *Orbicella faveolata* B3 (light purple). Each mean-annual Sr/Ca-SST record was smoothed using a 3-point running mean (CG2 dark pink, A1 dark gray, A2 dark gray, F1 dark gray, 93DRYSS2 dark gray, and B3 dark purple). The dark-pink stippled area in the CG2 record (1990–2012) indicates an interval where the Sr/Ca-SST values significantly deviate from the trends in the other records. (B) The stacked mean annual Sr/Ca-SST ($^{\circ}\text{C}$,

black line) created by combining the mean annual Sr/Ca-SST data from all 5 coral colonies (*O. faveolata* B3, and *S. siderea* colonies CG2, A, F1, and 93DRYSS2) using a mean-based stacking scheme with 5000 Monte Carlo simulations. Shaded intervals represent $\pm 68\%$ error boundaries. (C) The length of time covered by each individual colony. Colors are consistent with those in part A.

Figure 4. A) 10-year running correlation between the stacked Sr/Ca-SST record and the AMO index of detrended SST anomalies (Enfield et al., 2001). The shaded box represents the interval with the highest correlation ($r > 0.7$ and $p < 0.01$). B) The stacked mean-annual Sr/Ca-SST ($^{\circ}\text{C}$) reconstruction (black line, shaded intervals represent $\pm 68\%$ error boundaries plotted alongside the AMO index of detrended SST anomalies [$^{\circ}\text{C}$]; blue curve, [Enfield et al., 2001]). The records are plotted on top of each other where the best correlation ($r = 0.28$,) occurs (the AMO record lags the Sr/Ca-SST record by about 9 years). C) Plot of MTM spectral analysis performed on the stacked Sr/Ca-SST reconstruction (black line) with a red-noise AR1 process (red line, 90% confidence interval). Significant peaks occur at 80–90 years, 28–30 years and 2–8 years.

726

727

729

730