Thermal equation of state and stability of (Mg_{0.06}Fe_{0.94})O

June K. Wicks^{a,c}, Jennifer M. Jackson^a, Wolfgang Sturhahn^a, Kirill K.
Zhuravlev^b, Sergey N. Tkachev^b, Vitali B. Prakapenka^b

^a*Division of Geological and Planetary Sciences, California Institute of Technology,
Pasadena, California, USA*

^b*Center for Advanced Radiation Sources, University of Chicago, Chicago, IL 60637, USA*

^c*Now at Department of Geosciences, Princeton University, Princeton, NJ, USA*

Abstract

We present the pressure-volume-temperature (P – V – T) equation of state of polycrystalline (Mg_{0.06}Fe_{0.94})O (Mw94) determined from laser-heated x-ray diffraction experiments up to 122 GPa and 2100 K, conditions approaching those of the deep mantle. We conducted two sets of experiments, one with an *in situ* Fe metal oxygen fugacity buffer and one without such a buffer. The internal pressure markers used in these experiments were $B2$ -NaCl and hcp -Fe in the buffered experiment and $B2$ -NaCl in the unbuffered experiment. In the sampled P – T range of the high temperature part of this study, only the $B1$ structure of Mw94 was observed, indicating that the addition of Mg to FeO stabilizes the $B1$ phase with respect to the $B8$ phase at these conditions. Both datasets were fit to a Birch-Murnaghan and Mie-Grüneisen-Debye thermal equation of state using a new open-source fitting routine, also presented here. Analysis of these data sets using the same internal pressure marker shows that the P – V – T data of Mw94 obtained in the unbuffered experiment are well explained by the equation of state parameters determined from the buffered data set. We have also compared the thermal equation of state of Mw94 with that of wüstite and conclude that Mw94 has measurably distinct thermoelastic properties compared with those of wüstite. We use the results obtained in the buffered experiment to

Email address: jwicks@princeton.edu (June K. Wicks)

determine the density and bulk sound velocity of Mw94 at the base of the mantle and compare these values to geophysical observations of ultralow-velocity zones.

Keywords: equations of state, core-mantle boundary, (Mg,Fe)O

1. Introduction

1.1. Motivation

Iron-poor members of the periclase-wüstite (MgO-FeO) solid solution may comprise up to 20% of the Earth's lower mantle. According to the "pyrolite" model, the chemistry of this phase contains 10–20 % FeO (Ringwood, 1975). The simplest assumption is that the lower mantle is homogeneous, yet seismic studies reveal discontinuities and lateral heterogeneity in the deep mantle that suggest lateral chemical inhomogeneities (Garnero et al., 2007). Chemical analyses of magma associated with hotspot volcanism from mantle plumes reportedly sourced from the lower mantle also give evidence for chemical heterogeneity, promising a more complex lower mantle (Mukhopadhyay, 2012). To understand the role of (Mg,Fe)O in these variations, the thermodynamic and elasticity systematics of ferropericlase, $(\text{Mg}_{1-x}\text{Fe}_x)\text{O}$ with $x < 0.5$, has been extensively studied as a function of composition, pressure, and temperature. However, more iron-rich compositions of (Mg,Fe)O have not been explored in the same detail.

Ultralow-velocity zones (ULVZs) at the core-mantle boundary give insight into the chemical heterogeneity of the lowermost mantle. As their name implies, ULVZs are characterized by very low seismic wave velocities (10–30%) (e.g. Garnero and Helmberger, 1998; Helmberger et al., 2000; Rost et al., 2006; Sun et al., 2013). Requiring concomitant density increase and sound velocity decrease with respect to the surrounding mantle, ULVZs are best explained by Fe-enrichment. Recent considerations of a crystallizing primordial magma ocean show that enrichment of Fe in the lowermost mantle is possible, preserved as

25 either iron-rich solids or residual melt (Labrosse et al., 2007; Lee et al., 2010;
26 Nomura et al., 2011).

27 Incorporation of 40% Fe into magnesium silicate post-perovskite has been
28 found to reduce shear wave velocities to 33% lower than that of the 1D Preliminary
29 Reference Earth Model (PREM) (Mao et al., 2006; Dziewonski and Anderson,
30 1981). An iron-rich oxide, namely $(\text{Mg}_{0.16}\text{Fe}_{0.84})\text{O}$, also has low shear wave
31 velocities—50% lower than that of PREM (Wicks et al., 2010). Partial melting
32 has also been suggested as a plausible mechanism (Williams et al., 1998; Lay
33 et al., 2004; Labrosse et al., 2007).

34 Dynamic models have been used to explore the stability and topology of
35 ULVZs to differentiate the observable characteristics of various candidate as-
36 semblages. Hernlund and Tackley (2007) found that the amount of partial melt
37 required to decrease the sound velocities of a phase assemblage to match ULVZs
38 would percolate and pool at the base of the mantle, a result that could be pre-
39 vented if the ULVZ were stirred (Hernlund and Jellinek, 2010). Dynamic models
40 exploring solid ULVZs show that ULVZ shape can be correlated to chemical den-
41 sity anomaly, which in turn can be used to explore the tradeoff between density
42 and sound velocities of $(\text{Mg,Fe})\text{O}$ and bridgmanite (Bower et al., 2011; Sun
43 et al., 2013).

44 We are motivated, therefore, to measure the pressure-volume-temperature
45 equation of state of iron-rich $(\text{Mg,Fe})\text{O}$, so that the density of iron-rich ULVZs
46 can be properly modeled, and the relationship between sound velocity and den-
47 sity can further constrain the composition of ULVZs.

48 *1.2. Previous experimental studies*

49 Previous pressure-volume-temperature (P – V – T) studies have shown that
50 the MgO – FeO solid solution is complicated by the existence of phase transitions,
51 a spin transition, and defect clustering as a function of FeO component (e.g.
52 Lin et al., 2003; Fei et al., 2007a; Lin et al., 2005; Speziale et al., 2005; Kantor
53 et al., 2009; Mao et al., 2011). The MgO endmember is known to be cubic
54 beyond the pressure and temperature conditions of the Earth’s mantle (Duffy

55 and Ahrens, 1993). The FeO endmember is thought to be cubic at the pressures
 56 and temperatures of the interior of the Earth, but at lower temperatures is found
 57 to transform to a rhombohedral structure at moderate pressures (e.g. Shu et al.,
 58 1998) and to the *B8* NiAs structure at higher pressures (e.g. Fei and Mao, 1994;
 59 Fischer et al., 2011b).

60 At 300 K, iron-rich (Mg,Fe)O undergoes a cubic to rhombohedral phase
 61 transition at 8-40 GPa, with a phase transition pressure sensitive to both com-
 62 position and hydrostaticity. Studies of the bulk modulus at 0 GPa as a function
 63 of composition show differing trends due to sample stoichiometry. Studies of
 64 non-stoichiometric, iron-rich (Mg,Fe)O show that both K_S and K_T decreases as
 65 a function of Fe concentration (Jacobsen et al., 2002; Richet et al., 1989), where
 66 K_S is determined from measurements of volume, composition, and of V_P and
 67 V_S using ultrasonic interferometry, and K_T is determined in a P - V compres-
 68 sion study. The trend is opposite for stoichiometric samples, where ultrasonic
 69 interferometry studies for iron-poor samples display a positive trend of K_S with
 70 increasing iron content (Jacobsen et al., 2002). In the iron endmember, K_T does
 71 indeed depend on stoichiometry, with $\text{Fe}_{0.99}\text{O}$ being much less compressible than
 72 $\text{Fe}_{<0.98}\text{O}$ (Zhang, 2000).

73 Thermal expansion, $\alpha = \frac{1}{V}(\frac{\partial V}{\partial T})_P$ has been shown to be insensitive to Fe
 74 content for Mg-containing ferropericlase (Zhang and Kostak, 2002; Westrenen
 75 et al., 2005; Komabayashi et al., 2010; Mao et al., 2011). Yet, it appears to
 76 be sensitive to defect concentration. In the Fe end member, α is about 30%
 77 larger for $\text{Fe}_{0.942}\text{O}$ than $\text{Fe}_{0.987}\text{O}$ at pressures and temperatures up to 5.4 GPa
 78 and 1073 K (Zhang and Zhao, 2005). In this study, we aim to determine the
 79 P - V - T equation of state ($\text{Mg}_{0.06}\text{Fe}_{0.94}\text{O}$) at conditions approaching those of
 80 the deep mantle to constrain the thermoelasticity of iron-rich (Mg,Fe)O and to
 81 see if these aforementioned trends apply to iron-rich (Mg,Fe)O.

82 1.3. Experimental considerations

83 In this paper we discuss two types of experiments, "buffered" and "un-
 84 buffered". In previous high- PT experiments on wüstite, an *in situ* Fe metal

oxygen buffer has been used (Fischer et al., 2011b; Seagle et al., 2008; Ozawa et al., 2010). The motivation of using such a buffer has been to decrease the vacancy population of wüstite at high pressures and temperatures, as some studies indicate that the thermoelastic properties of wüstite vary as a function of vacancy concentration (Zhang, 2000; Zhang and Zhao, 2005).

2. Materials and Methods

Polycrystalline ($\text{Mg}_{0.06}^{57}\text{Fe}_{0.94}\text{O}$) was synthesized from ^{57}Fe (95% enriched, *Isoplex*) and MgO powders in a 1 atm gas-mixing furnace at 1673 K for two runs of 20 hours each. A H_2/CO_2 gas mixture was used to control the oxygen fugacity of the synthesis to just above that of the iron-wüstite fugacity buffer ($\log f_{\text{O}_2} = -9$). Sample composition was measured using a JEOL-JXA-8200 electron microprobe, reporting a composition of ($\text{Mg}_{0.058(1)}\text{Fe}_{0.942(1)}\text{O}$), where the number in parentheses is the error on the last digit. A conventional Mössbauer spectrum (Figure 1, inset) caps the ferric content at 5% of the total iron, thus capping the vacancy concentration at 5%.

Two experiments were conducted in this study. For the unbuffered experiment, a symmetric diamond anvil cell with 300 μm -culet diamonds was prepared. ($\text{Mg}_{0.06}\text{Fe}_{0.94}\text{O}$), hereafter referred to as Mw94, was lightly ground with NaCl powder (1:1 by volume) in an agate mortar under ethanol, allowed to dry, then pressed into a pellet. This mixture was loaded between two thin NaCl plates ($<10 \mu\text{m}$), and the remaining space in the rhenium gasket sample chamber was filled with neon using the COMPRES/GSECARS gas-loading system (Rivers et al., 2008). To keep the NaCl dehydrated, the pellet was consistently stored in a desiccator when not in use, and the loaded diamond anvil was placed under vacuum for about one hour and purged with argon before being put into the gas-loading system.

We conducted a second experiment with an *in situ* Fe metal oxygen buffer for comparison. In this second, buffered, high pressure experiment, 250 μm -culet diamonds were used. Otherwise, the preparation differed in that the sample

114 pellet (Mw94 and NaCl) also included a Fe metal (1:1 Mw94:Fe by weight) as
 115 an *in situ* oxygen buffer and pressure marker. Thus, both experiments contained
 116 an intimate mixture of Mw94 and NaCl in the sample pellet, so that NaCl may
 117 be used as a common pressure marker to compare the results of the buffered
 118 and unbuffered experiments.

119 The high temperature powder X-ray diffraction (XRD) experiments were
 120 conducted at the 13-ID-D beamline (GeoSoilEnviroCars) at the Advanced Photon
 121 Source, Argonne National Laboratory. Using an incident X-ray beam of $\lambda =$
 122 $0.3344 \text{ \AA}$ and focus spot size of $4 \text{ }\mu\text{m} \times 4 \text{ }\mu\text{m}$, angle-dispersive X-ray diffrac-
 123 tion patterns were recorded onto a MAR165 CCD detector and subsequently
 124 integrated using Fit2D (Hammersley et al., 1996). CeO_2 was used to calibrate
 125 the sample to detector distance at 1 bar. Samples were compressed to about 35
 126 GPa at room temperature before heating. High temperatures were achieved *in*
 127 *situ* by laser heating from both sides by $1.064 \text{ }\mu\text{m}$ Yb fiber lasers with 'flat top'
 128 profiles (Prakapenka et al., 2008) and temperatures determined spectroradio-
 129 metrically (e.g. Heinz and Jeanloz, 1987; Shen et al., 2001) using the gray body
 130 approximation over the 600-800 nm range of thermal emission. Temperatures
 131 were measured during the collection of the diffraction patterns. In the buffered
 132 experiment, only the upstream temperatures were used due to technical difficul-
 133 ties with the downstream temperature determinations. The diffraction peaks
 134 were relatively sharp compared to ambient temperature spectra, indicating that
 135 temperature gradients are likely small. Example XRD patterns are shown in
 136 Figure 1.

137 The 2θ angles corresponding to lattice reflections of $(\text{Mg}_{0.06}\text{Fe}_{0.94})\text{O}$, NaCl,
 138 Fe, and Ne were determined by fitting the patterns with Voigt peaks using Igor
 139 Pro (WaveMetrics, Lake Oswego, OR, USA). In the multi-peak fitting routine in
 140 Igor Pro, XRD pattern backgrounds were fit locally with a cubic function. Unit
 141 cell volumes were determined using unweighted linear regression using the *Unit*
 142 *Cell* refinement software package (Holland and Redfern, 1997), which assumes
 143 a minimum uncertainty of 0.005 degrees on each lattice reflection, with errors
 144 on the unit cell weighted by goodness of fit.

Figure 1: Example integrated XRD patterns showing peak identifications for $B2$ -NaCl, hcp -Fe, and Ne. Pressures were determined using the equation of state of $B2$ -NaCl (Fei et al., 2007b) for the unbuffered dataset (top two patterns), and hcp -Fe (Dewaele et al., 2006; Murphy et al., 2011) for the buffered dataset (bottom three patterns). $(Mg_{0.06}Fe_{0.94})O$ is rhombohedral at room temperature (Mw_R , $R\bar{3}m$) and cubic at high temperature (Mw_C , $Fm\bar{3}m$). Unidentified peaks are labeled with *. Arrows indicate location of expected B8 peaks at these conditions (Ozawa et al., 2011). Inset: A conventional Mössbauer spectrum of Mw94 before heating at 1 bar (Wicks, 2013), with two example fits (no Fe^{3+} , green, and 5% Fe^{3+} , red.)

Volumes of cubic Mw94 were refined using at least three of the following four reflections: 111, 200, 220, and 331. *hcp*-Fe volumes were refined using at least 6 of the 8 following reflections: 100, 200, 101, 102, 110, 103, 200, and 112. *B2*-NaCl volumes were refined using at least 5 of the 6 following reflections: 100, 110, 111, 200, 210, and 211. Neon volumes were determined from at least one of the two reflections: 111, 200. Rhombohedral Mw94 volumes were refined from at least 3 of the 5 following reflections: 003, 101, 102, 104, and 110.

The *B2*-NaCl thermal equation of state of Fei et al. (2007b) was used to determine pressure in the unbuffered experiment, and to compare that dataset to the buffered experiment, because each contained an intimate mixture of NaCl and Mw94. Pressures in the buffered experiment were determined using the unit cell volume of *hcp*-Fe. We used the equation of state of *hcp*-Fe given by Dewaele et al. (2006), and the thermal pressure given by Murphy et al. (2011). The difference between the two pressure scales is small, with a resulting pressure increase ranging from 0.01 to 0.4 GPa for the Murphy et al. (2011) values.

The greatest source of error in these measurements comes from the temperature determination. Past studies have analyzed these sources of error in great detail for experiments conducted in the laser-heated diamond-anvil cell (e.g. Heinz and Jeanloz, 1987; Kavner and Jeanloz, 1998; Panero and Jeanloz, 2001; Shim et al., 2000; Shen et al., 2001; Benedetti and Loubeyre, 2004; Noguchi et al., 2013). The errors depend heavily on the individual experiments, with cited temperature uncertainties ranging from a few percent to 10 percent. Considering the measured range of standard deviations from multiple measurements for each XRD pattern collected in our study (between 1 and 104 K) and previous reports suggesting an accuracy of such measurements to be between about 100 and 200 K (Shen et al., 2001; Sturhahn and Jackson, 2007; Noguchi et al., 2013), we report a temperature uncertainty of 100 K.

The error in pressure is a function of the uncertainty in both the volume and temperature of the *in situ* pressure calibrant (*hcp*-Fe or *B2*-NaCl, in this case). However, in determining the correct error assignments for the equation of state fitting, we made adjustments to ensure that errors were not double-counted.

176 Calculated volumes and pressures of the buffered and unbuffered datasets are
177 presented in Tables D.8 and D.9, respectively.

178 3. Results

179 3.1. Phase identification

180 In the P - T range of study, we identified cubic Mw94 at high temperature,
181 and rhombohedral or a mixture of rhombohedral and cubic Mw94 at room
182 temperature. We interpret the mixture to be a result of incomplete back-
183 transformation of the cubic to rhombohedral phase on quench. We plot our
184 phase identifications in Figure 2. In the diffraction patterns of the buffered
185 experiment, all peaks could be assigned to known phases in the sample cham-
186 ber (Mw94, NaCl, Fe, Ne). In the unbuffered experiment, there were a few
187 weak peaks that could not be assigned to known phases of Mw94, NaCl, or
188 Fe. We will return to this topic later in our discussion of stoichiometry. Our
189 cold compression points are shown in gray, bracketing the room temperature
190 cubic-rhombohedral transition between 13 and 24 GPa.

191 As we did not identify the $B8$ structure in our P - T range of study, we ob-
192 serve that the addition of Mg to FeO stabilizes the $B1$ phase with respect to the
193 $B8$ phase at these conditions. At 300 K, our data bracket the $B1$ to rhombohe-
194 dral transition indicated for FeO, but our data coverage at room temperature
195 is not high enough to make comparisons. At high temperature, our observation
196 of only the cubic structure is inconsistent with those of Kondo et al. (2004)
197 and Ohta et al. (2014). Kondo et al. (2004) reports an expansion of the stabil-
198 ity field of rhombohedrally-structured (Mg,Fe)O up to 2000 K at 120 GPa for
199 $(\text{Mg}_{0.05}\text{Fe}_{0.95})\text{O}$, $(\text{Mg}_{0.1}\text{Fe}_{0.9})\text{O}$ and $(\text{Mg}_{0.2}\text{Fe}_{0.8})\text{O}$ sandwiched between corun-
200 dum disks, regardless of composition. Ohta et al. (2014) reported a similar
201 rhombohedral phase boundary for $(\text{Mg}_{0.2}\text{Fe}_{0.8})\text{O}$ prepared with silica glass as a
202 pressure medium. The use of a neon pressure medium in our experiments could
203 be one explanation for the discrepancy.

Figure 2: Phase identification of $(\text{Mg}_{0.06}\text{Fe}_{0.94})\text{O}$ in P - T space. $(\text{Mg}_{0.06}\text{Fe}_{0.94})\text{O}$ is cubic (unbuffered $\square$, buffered $\blacksquare$) at high temperature and rhombohedral (unbuffered $\diamond$, buffered $\blacklozenge$) at room temperature. Before-heating points from the buffered experiment are shown below 25 GPa in gray ($\blacksquare, \blacklozenge$). Light gray lines and text: the latest phase diagram of wüstite (Fei and Mao, 1994; Fischer et al., 2011a,b; Kondo et al., 2004; Ozawa et al., 2010, 2011). Pressures shown here in the buffered experiment used *hcp*-Fe as a pressure marker, while pressures in the unbuffered experiment used *B2*-NaCl. Black dashed line: *B1*-rhombohedral phase boundary for iron (Mg,Fe)O of varying composition (Kondo et al., 2004). Error bars shown represent standard deviation of individual temperature measurements.

3.2. Thermal equation of state fitting using MINUTI

The buffered and unbuffered P - V - T data of Mw94 were fit using an open-source thermal equation of state fitting routine (seos) within the MINeral physics UTility software package, MINUTI version 1.1.0 (Sturhahn, 2015). We describe the thermal equation of state and parameter fitting here.

The free energy or Helmholtz energy is written as the sum of a term describ-

ing the elastic deformation F_{el} and a term describing the lattice vibrations

$$F(V, T) = F_{el}(V) + F_{th}(V, T) - F_{th}(V, T_0) = F_{el}(V) - k_B T \ln Z_{th}(V, T) + k_B T_0 \ln Z_{th}(V, T_0) \quad (1)$$

The partition function Z gives the contribution of lattice vibrations and is expressed by the phonon density of states

$$\ln Z_{th}(V, T) = - \int \ln(2 \sinh \frac{\omega}{2k_B T}) D(\omega, V) d\omega \quad (2)$$

The elastic part of (1) can be selected to be of type Birch-Murnaghan, Vinet, or logarithmic. In this study, we used a Birch-Murnaghan type, which is given by

$$F_{el}(V) = \frac{9}{2} K_0 V_0 f^2 \{1 + f (K'_0 - 4)\} \quad (3)$$

with $f = \frac{1}{2} \left\{ \left(\frac{V_0}{V} \right)^{2/3} - 1 \right\}$

In these relations, V_0 is the volume at zero pressure, K_0 is the isothermal bulk modulus at zero pressure, and K'_0 is the pressure derivative of the isothermal bulk modulus at zero pressure. All values are assumed at a reference temperature T_0 (300 K, in this case). Therefore at the reference temperature the EOS is given by the elastic part only.

The seos program uses the Debye approximation for the phonon density of states which features one volume-dependent parameter, the Debye temperature Θ . For $\omega \leq k_B \Theta$

$$D(\omega) = \frac{9}{k_B \Theta} \left(\frac{\omega}{k_B \Theta} \right)^2 \quad (4)$$

With this approximation, the free energy of the Debye phonon gas is given by

$$F_{th}(V, T) = 3k_B T \ln \left(2 \sinh \frac{\Theta}{2T} \right) - k_B T I\left(\frac{\Theta}{T}\right) - \frac{3}{8} k_B \Theta \quad (5)$$

with

$$I(t) = \frac{3}{t^3} \int_0^t \frac{x^3 dx}{e^x - 1} \quad (6)$$

227 The thermal pressure now takes the Mie-Grüneisen-Debye form

$$p_{th}(V, T) = - \left(\frac{\partial F_{th}}{\partial V} \right)_T = \frac{\gamma}{V} \left(3k_B T I \left(\frac{\Theta}{T} \right) + \frac{9}{8} k_B \Theta \right) \quad , \quad (7)$$

228 where $\gamma = -\partial \ln \Theta / \partial \ln V$ is the Debye-Grüneisen parameter which is expressed
229 by the seos program using a scaling law

$$\gamma(V) = \gamma_0 \left(\frac{V}{V_0} \right)^q \quad , \quad (8)$$

230 where γ_0 , q , and V_0 are inputs to the program. The expression for the Debye
231 temperature follows via integration of $\gamma = -\partial \ln \Theta / \partial \ln V$ to result in

$$\Theta(V) = \Theta_0 \exp \left[\frac{\gamma_0 - \gamma(V)}{q} \right] \quad , \quad (9)$$

232 where Θ_0 is an input to the seos program. For the data sets presented here, q
233 was fixed to 0.5. The Debye temperature, Θ_0 , was fixed to 426 K, calculated
234 from the measured Debye velocity of the starting material according to the
235 following relation:

$$\Theta_0 = \frac{h}{k} \left(\frac{3N_A \rho}{4\pi\mu} \right)^{1/3} V_D \quad (10)$$

236 where h is Planck's constant, k is the Boltzmann constant, N_A is Avogadro's
237 Number, μ is the reduced mass, $\rho = 5.94 \pm 0.02$ g/cc, and $V_D = 3.075 \pm 0.014$
238 km/s (Wicks, 2013). In this experiment, the equation of state fitting is not
239 sensitive to the Debye temperature, as the temperatures in this experiment are
240 much greater than those of the calculated Debye temperature.

241 For a given pressure p and temperature T , the volume is calculated by solving
242 the thermal EOS

$$p = - \frac{\partial F_{el}}{\partial V} + p_{th}(V, T) - p_{th}(V, T_0) \quad . \quad (11)$$

243 Input data are specified as sets $\{p_i, V_i, T_i, \delta p_i, \delta V_i, \delta T_i\}$ consisting of pressure,
244 volume, temperature, and their uncertainties. The EOS is fitted to these data by
245 variation of parameters by minimizing the normalized mean-square deviation.
246 In seos, some number of the fit parameters may have priors with uncertainties.

247 The weights are determined by uncertainties of the data (see Appendix A for
248 details).

249 If data and priors constrain the fit parameters reasonably well a solution with
250 minimal reduced χ^2 is obtained. These optimal parameters have errors related
251 to data variation and prior uncertainties. The errors are estimated as variances
252 (square errors) by Equation A.3. We report the fit-parameter correlation matrix,
253 $\sigma_{jj'}$, for each fitting result in the Appendix. The volume V is calculated based
254 on the EOS, Eq. 3, for various pressures and temperatures as specified in
255 the standard input file. If fit parameters are defined, the temperature-reduced
256 volumes V_{ri} are calculated from the data $\{p_i, V_i, T_i\}$ as

$$V_{ri} = V_i - V(p_i, T_i) + V(p_i, T_0) \quad , \quad (12)$$

257 where T_0 is the specified reference temperature. A plot of $V(p_i, T_0)$ and V_{ri}
258 versus pressure provides an easier visual assessment of deviations between data
259 and model (see examples in Section 3). The equation of state parameters and
260 their associated uncertainties are described in detail in Appendix B.

261 Other features, such as F-f plots are also determined from seos (Sturhahn,
262 2015), however are not reported in this paper. The seos executable also features
263 a spin crossover form of an equation of state that determines spin contribu-
264 tions (Chen et al., 2012).

265 With the best-fit parameters obtained from the thermal equation of state
266 fitting and the fit-parameter correlation matrix, the following parameters can be
267 calculated at a specified temperature over a specified pressure range: isother-
268 mal bulk modulus K_T , adiabatic seismic velocity V_ϕ , and thermal expansion
269 coefficient α , with uncertainties reflecting the reported correlations of the fitted
270 parameters.

271 3.3. Dataset with in-situ Fe metal buffer

272 The equation of state parameters of both cubic and rhombohedral (Mg_{0.06}Fe_{0.94})O
273 of the buffered dataset, 38–122 GPa, were fit using pressures given by the equa-
274 tion of state of *hcp*-Fe (Dewaele et al., 2006; Murphy et al., 2011). Equation of

state parameters are shown in Table 1.

Table 1: Equation of state parameters of cubic and rhombohedrally-structured (Mg_{0.06}Fe_{0.94})O for the buffered dataset using *hcp*-Fe as an internal pressure marker. Room temperature parameters correspond to a Birch-Murnaghan equation of state, while thermal parameters correspond to the Mie-Grüneisen equation of state. *Denotes numbers that were fixed in the fitting.

	<i>B1</i> -Mw94	<i>r</i> -Mw94
V_0 ($\text{\AA}^3/\text{atom}$)	9.92 ± 0.04	9.97 ± 0.11
K_{0T} (GPa)	197.4 ± 6.5	184 ± 16
K'_{0T}	2.79 ± 0.09	2.98 ± 0.25
θ_0 (K)	426*	
γ_0	1.72 ± 0.08	
q	0.50*	
reduced χ^2	1.07 ± 0.13	2.84 ± 0.51

Fitting with a Birch-Murnaghan equation of state yields $V_0 = 9.92 \pm 0.04 \text{ \AA}^3/\text{atom}$, $K_{0T} = 197.4 \pm 6.5$ GPa, and $K'_{0T} = 2.79 \pm 0.09$. Using the high temperature Mie-Grüneisen formalism, the fitted γ_0 was 1.72 ± 0.08 . The rhombohedral Mw94 phase was also fit to a Birch-Murnaghan EOS, yielding the parameters $V_0 = 9.97 \pm 0.11 \text{ \AA}^3/\text{atom}$, $K_{0T} = 184 \pm 16$ GPa, and $K'_{0T} = 2.98 \pm 0.25$.

Figure 3 shows the results from this buffered experiment, showing volume as a function of pressure and temperature. Also shown are the volumes measured as we compressed the cell before heating. We did not include them in the fit, as they were data collected without annealing. Near room-pressure, the steeper $\partial V/\partial P$, *i.e.* lower K_{0T} , is more consistent with a (magnesian)wüstite containing 2-5% vacancies (Zhang, 2000; Jacobsen et al., 2002). Projection of the high temperature data to a reference temperature of 300 K and associated residuals allows for visualization of goodness-of-fit (Equation 12, Figure 5a).

As the Mw94 sample pellet in the buffered experiment also contained a mixture of NaCl, this buffered dataset was also fit using pressures given by the equation of state of *B2*-NaCl (Table 2). When using pressures given by the equation of state of NaCl (Fei et al., 2007b), the fitted $V_0 = 9.92 \pm 0.03 \text{ \AA}^3/\text{atom}$, $K_{0T} = 199.8 \pm 5.3$ GPa, $K'_{0T} = 2.67 \pm 0.07$, and $\gamma_0 = 1.17 \pm 0.06$.

Figure 3: P - V - T data and isotherms of 1200 K, 1600 K, and 2000 K of $B1$ -structured $(\text{Mg}_{0.06}\text{Fe}_{0.94})\text{O}$ in the buffered experiment (Table 1). Pressures were determined using the equation of state of Fe (Dewaele et al., 2006; Murphy et al., 2011). Gray diamonds show rhombohedral $(\text{Mg}_{0.06}\text{Fe}_{0.94})\text{O}$ at room temperature, and the black line is the 300 K fit to these data, reduced $\chi^2 = 1.07$ (Equation 12). Open circles: pre-heating data points at pressures determined by the equation of state of $B1$ -NaCl (JCPDS 5-0628). Fitting residuals and data points projected to reference temperature of 300 K are shown in Figure 5a.

3.4. Dataset with no buffer

The equation of state of the unbuffered dataset (30–70 GPa) was fit using pressures given by the equation of state of $B2$ -NaCl (Table 2). Fitting to a Birch-Murnaghan EOS yields $V_0 = 9.75 \pm 0.04 \text{ Å}^3/\text{atom}$, $K_{0T} = 237 \pm 10 \text{ GPa}$, $K'_{0T} = 2.26 \pm 0.18$, and $\gamma_0 = 1.34 \pm 0.11$.

Consideration of the fitted parameters of the unbuffered dataset at face value

Table 2: Equation of state parameters of cubic (Mg_{0.06}Fe_{0.94})O for the buffered and unbuffered datasets using *B2*-NaCl as an internal pressure marker. Room temperature parameters correspond to a Birch-Murnaghan equation of state, while thermal parameters correspond to the Mie-Grüneisen equation of state. *Denotes numbers that were fixed in the fitting. ^aA prior of 190±15 GPa was placed on K_{0T} .

	buffered	unbuffered	unbuffered (prior)
V_0 ($\text{\AA}^3/\text{atom}$)	9.92 ± 0.03	9.75 ± 0.04	9.87 ± 0.03
K_{0T} (GPa)	199.8 ± 5.3	237 ± 10	202 ± 6^a
K'_{0T}	2.67 ± 0.07	2.26 ± 0.18	2.89 ± 0.13
θ_0 (K)	426*	426*	426*
γ_0	1.17 ± 0.06	1.34 ± 0.11	1.39 ± 0.11
q	0.50*	0.50*	0.50*
reduced χ^2	0.61 ± 0.10	0.20 ± 0.05	0.23 ± 0.06

may lead to unphysical conclusions about the equation of state of (Mg,Fe)O, often a concern of fitting a dataset with limited pressure-temperature range. Initial volume was not fixed in these fits due to the uncertainty of defect concentration, although the starting material measured volume gives a lower bound of 9.9 $\text{\AA}^3/\text{atom}$. Expected bulk moduli of iron-rich (Mg,Fe)O are closer to 180 GPa (Jacobsen et al., 2002), while the buffered dataset fit produced a value around 190 GPa. Such factors can be used to inform the fitting routine in the form of a prior. When fitting the unbuffered dataset with a prior placed on K_{0T} of 190±15 GPa, the resulting EOS parameters are $V_0 = 9.87 \pm 0.03 \text{ \AA}^3/\text{atom}$, $K_{0T} = 202 \pm 6$ GPa, $K'_{0T} = 2.89 \pm 0.13$, and $\gamma_0 = 1.39 \pm 0.11$. This fit results in a slight increase in the reduced χ^2 (0.23 compared to 0.20), a result of the penalty in using a prior (equation A.1). Given the limited pressure-temperature range of the unbuffered study compared to the buffered study, the equation of state fitted using such a prior on K_{0T} (Table 2) is favored. Caution should be used in extrapolating this EOS beyond the PT conditions studied here.

Figure 4 shows the results from the unbuffered experiment, showing volume change of (Mg_{0.06}Fe_{0.94})O as a function of pressure and temperature, using NaCl as a pressure marker. A visual comparison of the two fits (with and without a prior on K_{0T}) can be seen in Figure 6 showing that in the region of data

Figure 4: P - V - T data and isotherms at 1200 K, 1400 K, and 1700 K of $B1$ -structured $(\text{Mg}_{0.06}\text{Fe}_{0.94})\text{O}$ in the unbuffered experiment. Equation of state parameters corresponding to these curves are those using a prior of $K_{0T}=190(15)$ GPa in Table 2. The 300 K isotherm for this EOS is also shown up to 30 GPa. Pressures were determined using the equation of state of $B2$ -NaCl (Fei et al., 2007b) (reduced $\chi^2 = 0.23$). Fitting residuals and data points projected to reference temperature of 300 K are shown in Figure 5b.

coverage, the two fits agree with each other very well.

4. Discussion

4.1. Phase diagram of iron-rich $(\text{Mg},\text{Fe})\text{O}$

The phase identifications presented in Figure 2 are consistent with a previous result in that there is no observable $B8$ -structured $(\text{Mg}_{0.06}\text{Fe}_{0.94})\text{O}$ in the pressure and temperature range studied (Kondo et al., 2004). Our study, however,

disagrees with previous reports of the location of the rhombohedral-cubic phase boundary of iron-rich (Mg,Fe)O at high temperature. Where Kondo et al. (2004) find a transition of rhombohedral to cubic (Mg_{0.1}Fe_{0.9})O and (Mg_{0.05}Fe_{0.95})O at 1100 to 1500 K in the pressure range of 70 to 100 GPa, we find no evidence of rhombohedral (Mg_{0.06}Fe_{0.94})O at the high temperatures explored in our study (Figure 2).

In comparison to the phase diagram of FeO, we show that substitution of Mg into FeO expands the stability field of the rhombohedral phase with respect to the *B8* phase.

Decomposition of the (Mg,Fe)O solid solution at high P – T conditions has been reported (Dubrovinsky et al., 2000). More recently, (Mg_{0.05}Fe_{0.95})O was found to decompose into *B8* FeO and (Mg_{0.06}Fe_{0.94})O in a diffraction pattern at 123 GPa, after heating up to 2800 K. This sample was also recovered to 1 bar for ATEM measurements and shown to have these two distinct phases (Ohta et al., 2014). Such compelling evidence for a high P - T experiment on Mw95 at pressures around 70 GPa was not presented, although Mw95 was suggested to have decomposed (Ohta et al., 2014). The starting composition we investigated in this study is one of their suggested reaction products, Mw94. Nevertheless, we searched our diffraction patterns for such evidence and do not see end member FeO in any of our diffraction patterns. It is possible that Mw94 does decompose, but at higher P - T conditions than what we sampled or that (Mg,Fe)O forms a complete solid solution up to at least 94% FeO.

4.2. Equations of state for iron-rich (Mg,Fe)O

The isothermal bulk modulus K_{0T} resulting from the equation of state fitting to the high- PT data of the buffered and unbuffered data sets is at the upper end of that predicted for nearly stoichiometric (Mg,Fe)O (Jacobsen et al., 2002). Previous work shows that the bulk modulus of (Mg,Fe)O decreases with increasing Fe content, from ~160 GPa for the MgO end member to 150 GPa for Fe_{0.95}O, likely due to increasing defect content. Conversely, stoichiometric examples show the opposite trend— bulk modulus increases with increasing Fe

content, up to 180 GPa for stoichiometric FeO (Jacobsen et al., 2002). All of our fits fall on the high side of this range of bulk moduli, but are admittedly not very well constrained by a compression study such as this one, with no annealed data at low pressure.

In Figure 6, we compare the data and thermal equations of state determined for the buffered and unbuffered experiments using their common internal pressure marker, the *B2*-NaCl pressure scale (Fei et al., 2007b). We do not discern an effect of buffering the composition on the equation of state of the Mw94 sample investigated here.

As a more quantitative measure, two datasets can be compared by using one dataset’s results as a prior on the other. When fitting the unbuffered dataset, using buffered values and associated error bars as priors, the resulting reduced chi-squared is 0.27 ± 0.06 , not worse than that of the fit without said priors. As a result, the two datasets are not distinguishable in their overlapping P-T range. The values reported in Table 2 show that some of the fitted parameters are consistent within error bars, with the exception of K_{0T} and K'_{0T} . A discrepancy in these two parameters, in turn, is not surprising, given the limited compression range of the unbuffered dataset, especially in the low pressure range where K_{0T} and K'_{0T} are most tightly constrained.

In Figure 7, error ellipses showing the tradeoff between equation of state parameters K'_{0T} versus γ_0 and V_0 versus K_{0T} demonstrate the overlap in thermal EOS parameters between buffered and unbuffered datasets when using the same internal pressure marker (*B2*-NaCl).

Another assessment is the level of compatibility in the thermal EOS parameters between the buffered Mw94 experiment (this study) and that of wüstite (Fischer et al., 2011b). As both experiments were conducted with an *in situ* Fe metal buffer/pressure marker, any measurable differences should be due to the incorporation of Mg into wüstite. When evaluating the buffered ($\text{Mg}_{0.06}\text{Fe}_{0.94}\text{O}$) dataset, using the FeO equation of state values from Fischer et al. (2011b) as priors ($\gamma = 1.42(5)$, $K_{0T} = 149(1)$, $K'_{0T} = 3.60(4)$), the resulting reduced chi-squared increased from 1.07 to 1.74 ± 0.16 , indicating that the two datasets are

386 not compatible.

387 From the equation of state parameters reported in Table 1, the thermal ex-
388 pansion coefficient α at specified output temperatures can be computed (Eq.
389 B.5) with uncertainties (Eqs. A.5, B.7). Figure 8 shows the thermal expansion
390 coefficient α (Equation B.5) determined from the buffered dataset calculated at
391 1500 K and 3800 K over the pressure range 0–140 GPa. We chose the value
392 of 3800 K, because the melting temperature of wüstite at core-mantle bound-
393 ary conditions is suggested to be about 3900 K (Fischer and Campbell, 2010).
394 Also shown are the thermal expansion coefficients calculated for FeO which was
395 measured up to 156 GPa and 3100 K (Fischer et al., 2011b) and for MgO which
396 was measured up to 196 GPa and 3700 K (Tange et al., 2009, Fit2-Vinet). It
397 is obvious from Figure 8 that the thermal equation of state of buffered wüstite
398 (Fischer et al., 2011a) and buffered Mw94 using *hcp*-Fe as a pressure marker in
399 each study are significantly distinct, as we discussed earlier. With this consid-
400 eration, together with the Mw94 fit quality (e.g., Figure 5a) and the parameter
401 correlations (Appendix C), we conclude that Mw94 has measurably distinct
402 thermoelastic properties compared with those of wüstite.

403 4.3. Sample Stoichiometry

404 The collected Mössbauer spectrum of the starting material (Figure 1, inset)
405 was fit using CONUSS (Sturhahn, 2000). A fit of a single Fe^{2+} site with an
406 isomer shift of 1.03 ± 0.04 mm/s (with respect to bcc iron) and a quadrupole
407 splitting of 1.21 ± 0.03 mm/s (distribution of 0.43 ± 0.03 mm/s) best fit the data
408 ($\chi^2 = 0.99$), but does not capture the asymmetry of the doublet. The addition
409 of a Fe^{3+} site with an isomer shift of 0.4 mm/s and a quadrupole splitting of
410 0.3 mm/s better (McCammon and Price, 1985) describes the spectrum visually
411 with resulting hyperfine parameters of the ferrous site within uncertainties of
412 the above-mentioned parameters ($\chi^2 = 0.95$).

413 The discrepancy between un-annealed low pressure data (0 to about 13 GPa,
414 Figure 3) and annealed high pressure measurements in the buffered experiment
415 suggests a physical difference between cold-compressed and annealed iron-rich

(Mg,Fe)O. As mentioned, the steeper slope is more consistent with a sample containing 2-5% vacancies (Zhang, 2000; Jacobsen et al., 2002). It has been proposed that vacancy concentrations in non-stoichiometric FeO-bearing samples are reduced with the exsolution of (Fe,Mg)Fe₂O₄ at high pressure and temperature, ensuring the stoichiometry of the (Mg,Fe)O phase regardless of oxygen fugacity (Zhang and Zhao, 2005; McCammon et al., 1998). It has also been proposed in the FeO end member that this exsolution occurs at the pressure of the magnetite to *h*-Fe₃O₄ transition (Fei, 1996).

It may be possible that this reaction proceeded at high temperatures in the unbuffered experiment, as the small, unidentified peaks may be attributed to a high pressure oxide phase such as *h*-Fe₃O₄, MgFe₂O₄ or Fe₄O₅ (Dubrovinsky et al., 2003; Lavina et al., 2011; Andrault and Bolfan-Casanova, 2001), Appendix E. The resulting consequences for sample chemistry are unclear. It could be possible that an exsolved oxide is either more, less, or equally iron-rich than the remaining material, leaving behind a sample that is less, more, or equally iron-rich to the starting composition. Similarly, we assume that the buffered experiment is achieved by the addition of Fe⁰ for every Fe³⁺, assured by the lack of excess peaks. In the end member scenario, the assumption of 5% vacancies would create a sample enriched in iron in comparison to the unbuffered experiment. As mentioned previously, the buffered and unbuffered datasets were statistically similar to each other, but inconsistent with FeO, so we conclude that resulting chemical variation between the two experiments must be small.

4.4. Geophysical Implications

When extrapolated to the conditions near the core-mantle boundary, at 135 GPa and 3800 K (e.g. Tackley, 2012), our buffered equation of state predicts a density of $\rho = 8.32 \pm 0.04$ g/cm³, bulk modulus $K_T = 423 \pm 8$ GPa, bulk sound velocity $v_\phi = 7.14 \pm 0.07$ km/s, and thermal expansion $\alpha = 2.01 \pm 0.01$ (10⁻⁵ K⁻¹). Figure 9 shows the calculated density and bulk sound velocities at 3800 K of (Mg_{0.06}Fe_{0.94})O, determined from the fitted thermal equation of state. Covariance among parameters is considered in the determination of uncertainty

(see Appendix C). Also shown are density and bulk sound velocity of MgO (red dotted line, Tange et al. (2009), Fit2-Vinet), wüstite (red dashed line, Fischer et al. (2011b)), bridgmanite (blue dotted line, (Tange et al., 2012, Vinet Fit)), calcium silicate perovskite (green dotted line, (Noguchi et al., 2013, Model 1)), and the bulk mantle, represented by PREM (Dziewonski and Anderson, 1981). Also shown at 135 GPa is the range of constrained density from seismic studies of ultralow-velocity zones (Rost et al., 2006), which may contain iron-rich (Mg,Fe)O.

Iron-rich (Mg,Fe)O in solid form is denser than the silicate-dominated lower mantle (MgSiO_3), but not as dense as the liquid Fe-alloy outer core (Figure 9). Our constrained value of α can thus be used in geodynamic models to compute the thermal buoyancy of iron-rich (Mg,Fe)O, for example to model the dynamics of ULVZs (e.g. Bower et al., 2011). The bulk sound velocity of Mw94 is lower than both the lowermost mantle and the core.

Using geodynamic modeling, Bower et al. (2011) showed that chemical density anomalies of 2–8% produced ULVZs with a variety of reasonable ULVZ morphologies. In the future, these characteristic morphologies could be used to constrain composition with sufficient 2-d and 3-d seismic imaging. Other studies cite ULVZs of around 10% density anomaly (e.g. Rost et al., 2006). If we assumed that a ULVZ was formed by entrainment of iron-rich material (such as Mw94) with ambient mantle, a 2–10% density anomaly could be explained by 4–20% volume fraction, respectively. To match ULVZ densities, a smaller(larger) volume fraction would be required for mixing (Mg,Fe)O with a greater(smaller) Fe content.

The relative density of iron-rich (Mg,Fe)O compared to its surroundings may also be useful when interpreting seismic signatures on the core side of the core-mantle boundary, which could be a reservoir of solid precipitates on the top of the core (Buffett et al., 2000). These precipitates can aid as a diffusion barrier between the core and the mantle, and thus far display seismic signatures that have been difficult to distinguish from similar features on the mantle side.

476 5. Summary

477 We have determined the P - V - T equation of state of $(\text{Mg}_{0.06}\text{Fe}_{0.94})\text{O}$ in
478 two distinct sets of experiments: one with an Fe oxygen fugacity buffer and one
479 without Fe. Considering the covariance of the fitted equation of state parameters
480 using the seos module in the MINUTI software package, we have determined the
481 level of compatibility between the buffered and unbuffered data sets of Mw94,
482 as well as between Mw94 and wüstite, using the same pressure indicators. We
483 find that the addition of small amounts of Mg causes measurable affects on the
484 thermoelastic properties of wüstite.

485 We see a clear difference in the location of the phase boundary between
486 cubic and rhombohedral iron-rich $(\text{Mg,Fe})\text{O}$ at high pressure and temperature,
487 compared with previous studies. Unlike Kondo et al. (2004), we do not find
488 rhombohedrally-structured iron-rich $(\text{Mg,Fe})\text{O}$ in our high temperature mea-
489 surements. In addition, we also do not find the $B8$ structure in our high tem-
490 perature measurements, indicating that Mg stabilizes the cubic phase upon
491 substitution into FeO (Fischer et al., 2011b).

492 The calculated bulk sound velocity and density from this study can be used
493 to constrain geophysical models of ULVZs that consider an assemblage of phases
494 that include iron-rich $(\text{Mg,Fe})\text{O}$.

495 Acknowledgements

496 We thank D. Zhang and C.A. Murphy for assistance with the measure-
497 ments, A.S. Wolf and D.B. Bower for discussions, and NSF-EAR 0711542 and
498 CSEDI EAR-0855815 for financial support. We thank E.E. Alp for conducting
499 the conventional Mössbauer spectroscopy measurements at Argonne National
500 Laboratory. This work was performed at GeoSoilEnviroCARS (Sector 13),
501 Advanced Photon Source (APS), Argonne National Laboratory. GeoSoilEnviroCARS
502 is supported by the National Science Foundation - Earth Sciences (EAR-1128799)
503 and Department of Energy - Geosciences (DE-FG02-94ER14466). Use of the
504 Advanced Photon Source was supported by the U.S. Department of Energy,

505 Office of Science, Office of Basic Energy Sciences, under contract No. DE-
506 AC02-06CH11357. The gas-loading system at GSECARS and conventional
507 Mössbauer spectroscopy laboratory are supported in part by COMPRES un-
508 der NSF Cooperative Agreement EAR 06-49658. Sample synthesis was carried
509 out in the petrological facilities at Caltech. Microprobe analyses were carried
510 out at the Caltech GPS Division Analytical Facility (funded in part by the
511 MRSEC Program of the NSF under DMR-0080065). Finally, we would like to
512 acknowledge two anonymous reviewers, whose feedback helped to improve the
513 manuscript.

Figure 5: Fitting residuals of $(\text{Mg}_{0.06}\text{Fe}_{0.94})\text{O}$ and projected data at 300 K. A) buffered dataset (reduced $\chi^2 = 1.07$). B) unbuffered dataset fit with a prior on $K_{0T} = 190(15)$ GPa (reduced $\chi^2 = 0.23$).

Figure 6: Overlay of data and equations of state of $(\text{Mg}_{0.06}\text{Fe}_{0.94})\text{O}$ in both experiments, with isotherms corresponding to 1100 K, 1500 K, and 1900 K. The pressure of the buffered (circles, solid lines) and unbuffered (squares, dashed and dotted lines) datasets were determined by the equation of state of their common internal pressure marker, $B2\text{-NaCl}$ (Fei et al., 2007b, Table 2). *dotted line denotes the fit of the unbuffered dataset with a prior on K_{0T} .

Figure 7: Error ellipses of equation of state parameters of unbuffered (red, using a prior on K_0) and buffered (blue) dataset using $B2\text{-NaCl}$ as a pressure marker (Table 2), demonstrating tradeoff of parameters K'_{0T} versus γ_0 and V_0 versus K_{0T} , respectively. 1 and 2- σ standard deviations are shown with solid and dashed lines, respectively.

Figure 8: Linear thermal expansion coefficient (α , Equation B.5) calculated at 1500 K and 3800 K as a function of pressure for $(\text{Mg}_{0.06}\text{Fe}_{0.94})\text{O}$ with an *in situ* Fe buffer (blue). Error bars correspond to uncertainties calculated in seos (see Appendix). Also shown are same values calculated for FeO (Fischer et al., 2011b, black) and MgO (Tange et al., 2009, red). Dashed lines indicate regions where the values are extrapolated beyond the P - T range of study.

Figure 9: Density and bulk sound velocity of $(\text{Mg}_{0.06}\text{Fe}_{0.94})\text{O}$ calculated at 3800 K as a function of pressure and depth. Also shown: MgO, (red dotted line, Tange et al., 2009), FeO (red dashed line, Fischer et al., 2011b), magnesium silicate bridgmanite (blue dotted line, Tange et al., 2012, Vinet Fit), calcium silicate perovskite, (green dotted line, Noguchi et al., 2013, Model 1), PREM (black solid line, Dziewonski and Anderson, 1981). Also shown at 135 GPa: range of constrained density from seismic studies of ultralow-velocity zones (e.g. Rost et al., 2006).

514 Andrault, D., Bolfan-Casanova, N., 2001. High-pressure phase transformations
 515 in the MgFe_2O_4 and Fe_2O_3 - MgSiO_3 systems. *Phys. Chem. Min.* 28, 211–217.

516 Benedetti, L.R., Loubeyre, P., 2004. Temperature gradients, wavelength-
 517 dependent emissivity, and accuracy of high and very-high temperatures mea-
 518 sured in the laser-heated diamond cell. *High Press. Res.* 24, 423–445.

519 Bower, D.J., Wicks, J.K., Gurnis, M., Jackson, J.M., 2011. A geodynamic and
 520 mineral physics model of a solid-state ultralow-velocity zone. *Earth Planet.*
 521 *Sci. Lett.* 303, 193–202.

522 Buffett, B.A., Garnero, E.J., Jeanloz, R., 2000. Sediments at the Top of Earth’s
 523 Core. *Science* 290, 1338–1342.

524 Chen, B., Jackson, J.M., Sturhahn, W., Zhang, D., Zhao, J., Wicks, J.K.,
 525 Murphy, C.A., 2012. Spin crossover equation of state and sound velocities of
 526 $(\text{Mg}_{0.65}\text{Fe}_{0.35})\text{O}$ ferropericlasite to 140 GPa. *J. Geophys. Res.* 117.

527 Dewaele, A., Datchi, F., Loubeyre, P., Mezouar, M., 2008. High-pressure–
 528 high-temperature equations of state of neon and diamond. *Phys. Rev. B* 77,
 529 094106.

530 Dewaele, A., Loubeyre, P., Occelli, F., Mezouar, M., Dorogokupets, P.I.,
 531 Torrent, M., 2006. Quasihydrostatic Equation of State of Iron above 2 Mbar.
 532 *Phys. Rev. Lett.* 97, 215504.

533 Dubrovinsky, L., Dubrovinskaia, N., Saxena, S.K., Annersten, H., Hålenius,
 534 E., Harryson, H., Tutti, F., Rekhi, S., Le Bihan, T., 2000. Stability of
 535 Ferropericlasite in the Lower Mantle. *Science* 289, 430–432.

536 Dubrovinsky, L.S., Dubrovinskaia, N., McCammon, C., Rozenber, G.K., Ahuja,
 537 R., Osorio-Guillen, J.M., Dmitriev, V., Weber, H.P., Le Bihan, T., Johansson,
 538 B., 2003. The structure of the metallic high-pressure Fe_3O_4 polymorph: ex-
 539 perimental and theoretical study. *J. Phys.: Condens. Matter* 15, 7697–7706.

540 Duffy, T.S., Ahrens, T.K., 1993. Thermal expansion of mantle and core mate-
541 rials at very high pressures. *Geophys. Res. Lett.* 20, 1103–1106.

542 Dziewonski, A.M., Anderson, D.L., 1981. Preliminary reference Earth model.
543 *Phys. Earth Planet. Inter.* 25, 297–356.

544 Fei, Y., 1996. Crystal chemistry of FeO at high pressure and temperature. The
545 *Geochemical Society, Special Publication Number 5*, 243–254.

546 Fei, Y., Zhang, L., Corgne, A., Watson, H., A., R., Meng, Y., Prakapenka,
547 V., 2007a. Phase transition and metallization of FeO at high pressures and
548 temperatures. *Geophys. Res. Lett.* 34, L17307.

549 Fei, Y., Mao, H.k., 1994. In Situ Determination of the NiAs Phase of FeO at
550 High Pressure and Temperature. *Science* 266, 1678–1680.

551 Fei, Y., Ricolleau, A., Frank, M., Mibe, K., Shen, G., Prakapenka, V., 2007b.
552 Toward an internally consistent pressure scale. *Proc. Natl. Acad. of Sci.* 104,
553 9182–9186.

554 Fischer, R.A., Campbell, A.J., 2010. High-pressure melting of wüstite. *Am.*
555 *Mineral.* 95, 1473–1477.

556 Fischer, R.A., Campbell, A.J., Lord, O.T., Shofner, G.A., Dera, P., Prakapenka,
557 V.B., 2011a. Phase transition and metallization of FeO at high pressures and
558 temperatures. *Geophys. Res. Lett.* 38, L24301.

559 Fischer, R.A., Campbell, A.J., Shofner, G.A., Lord, O.T., Dera, P., Prakapenka,
560 V.B., 2011b. Equation of state and phase diagram of FeO. *Earth Planet. Sci.*
561 *Lett.* 304, 496–502.

562 Garnero, E.J., Helmberger, D.V., 1998. Further structural constraints and un-
563 certainties of a thin laterally varying ultralow-velocity layer at the base of the
564 mantle. *J. Geophys. Res.* 103, 12,495–12,509.

565 Garnero, E.J., Lay, T., McNamara, A., 2007. Implications of lower-mantle
 566 structural heterogeneity for existence and nature of whole-mantle plumes.
 567 GSA Special Papers 430, 79–101.

568 Hammersley, A.O., Svensson, S.O., Hanfland, M., Fitch, A.N., Hausermann,
 569 D., 1996. Two-dimensional detector software: from real detector to idealized
 570 image or two-theta scan. High Press. Res. 14, 235–248.

571 Heinz, D.L., Jeanloz, R., 1987. Temperature measurements in the laser-heated
 572 diamond cell. Geophysical Monograph Series 39, 113–127.

573 Helmberger, D.V., Ni, S., Wen, L., Ritsema, J., 2000. Seismic evidence for
 574 ultralow-velocity zones beneath Africa and eastern Atlantic. J. Geophys. Res.
 575 105, 23,865–23,878.

576 Hernlund, J., Jellinek, A.M., 2010. Dynamics and structure of a stirred partially
 577 molten ultralow-velocity zone. Earth Planet. Sci. Lett. 296, 1–8.

578 Hernlund, J., Tackley, P., 2007. Some dynamical consequences of partial melting
 579 in Earth’s deep mantle. Phys. Earth Planet. Inter. 162, 149–163.

580 Holland, T.J.B., Redfern, S.A.T., 1997. Unit cell refinement from powder
 581 diffraction data: the use of regression diagnostics. Mineral. Mag. 61, 65–77.

582 Jacobsen, S.D., Reichmann, H.J., Spetzler, H., Mackwell, S.J., Smyth, J.R.,
 583 Angel, R.J., McCammon, C.A., 2002. Structure and elasticity of single-crystal
 584 (Mg,Fe)O and a new method of generating shear waves for gigahertz ultrasonic
 585 interferometry. J. Geophys. Res. 107, 5867–5871.

586 Kantor, I., Dubrovinsky, L., McCammon, C., Steinle-Neumann, G., Kantor, A.,
 587 Skorodumova, N., Pascarelli, S., Aquilanti, G., 2009. Short-range order and
 588 Fe clustering in $\text{Mg}_{1-x}\text{Fe}_x\text{O}$ under high pressure. Phys. Rev. B 80, 014204.

589 Kavner, A., Jeanloz, R., 1998. High-pressure melting curve of platinum. J.
 590 Appl. Phys. 83, 7553–7559.

591 Komabayashi, T., Hirose, K., Nagaya, Y., Sugimura, E., Ohishi, Y., 2010. High-
592 temperature compression of ferropericlase and the effect of temperature on
593 iron spin transition. *Earth Planet. Sci. Lett.* 297, 691–699.

594 Kondo, T., Ohtani, E., Hirao, N., Yagi, T., Kikegawa, T., 2004. Phase transi-
595 tions of (Mg,Fe)O at megabar pressures. *Phys. Earth Planet. Inter.* 143-144,
596 201–213.

597 Labrosse, S., Hernlund, J., Coltice, N., 2007. A crystallizing dense magma ocean
598 at the base of the Earth’s mantle. *Nature* 450, 866–869.

599 Lavina, B., Dera, P., Kim, E., Men, Y., Downs, R., Weck, P.F., Sutton, S.R.,
600 Zhao, Y., 2011. Discovery of the recoverable high-pressure iron oxide Fe₄O₅.
601 *Proc. Natl. Acad. of Sci.* 108, 17281–17285.

602 Lay, T., Garnero, E.J., Williams, Q., 2004. Partial melting in a thermo-chemical
603 boundary layer at the base of the mantle. *Phys. Earth Planet. Inter.* 146, 441–
604 467.

605 Lee, C.T.A., Luffi, P., Hoink, T., Li, J., Dasgupta, R., Hernlund, J., 2010.
606 Upside-down differentiation and generation of a ‘primordial’ lower mantle.
607 *Nature* 463, 30–U102.

608 Lin, J.F., Heinz, D.L., Mao, H., Hemley, R.J., Devine, J.M., Li, J., Shen, G.,
609 2003. Stability of magnesiowüstite in Earth’s lower mantle. *Proc. Natl. Acad.*
610 *of Sci.* 100, 4405–4408.

611 Lin, J.F., Struzhkin, V., Jacobsen, S., Hu, M., Chow, P., Kung, J., Liu, H.,
612 Mao, H., Hemley, R., 2005. Spin transition of iron in magnesiowüstite in the
613 Earth’s lower mantle. *Nature* 436, 377–380.

614 Mao, W.L., Lin, J.F., Liu, J., Prakapenka, V.B., 2011. Thermal equation of state
615 of lower-mantle ferropericlase across the spin crossover. *J. Geophys. Res.* 38,
616 L23308.

617 Mao, W.L., Mao, H., Sturhahn, W., Zhao, J., Prakapenka, V.B., Meng, Y., Shu,
618 J., Fei, Y., Hemley, R.J., 2006. Iron-Rich Post-Perovskite and the Origin of
619 Ultralow-Velocity Zones. *Science* 312, 564–565.

620 McCammon, C., Peyronneau, J., Poirier, J.P., 1998. Low ferric iron content
621 of (Mg,Fe)O at high pressures and temperatures. *Geophys. Res. Lett.* 25,
622 1589–1592.

623 McCammon, C.A., Price, D.C., 1985. Mössbauer spectra of $\text{Fe}_{x\text{O}}$ ($x \leq 0.95$). *Phys.*
624 *Chem. Min.* 11, 250–254.

625 Mukhopadhyay, S., 2012. Early differentiation and volatile accretion recorded
626 in deep-mantle neon and xenon. *Nature* 486, 101–104.

627 Murphy, C.A., Jackson, J.M., Sturhahn, W., Chen, B., 2011. Melting and
628 thermal pressure of hcp-Fe from the phonon density of states. *Phys. Earth*
629 *Planet. Inter.* 188, 114–120.

630 Noguchi, M., Komabayashi, T., Hirose, K., Ohishi, Y., 2013. High-temperature
631 compression experiments of CaSiO_3 perovskite to lowermost mantle condi-
632 tions and its thermal equation of state. *Phys. Chem. Min.* 40, 81–91.

633 Nomura, R., Ozawa, H., Tateno, S., Hirose, K., Hernlund, J., Muto, S., Ishii,
634 H., Hiraoka, N., 2011. Spin crossover and iron-rich silicate melt in the Earth's
635 deep mantle. *Nature* 473, 199–203.

636 Ohta, K., Fujino, K., Kuwayama, Y., Kondo, T., Shimizu, K., Ohishi, Y., 2014.
637 Highly conductive iron-rich (Mg,Fe)O magnesiowüstite and its stability in the
638 Earth's lower mantle. *J. Geophys. Res.* 119, 4656–4665.

639 Ozawa, H., Hirose, K., Ohta, K., Ishii, H., Hiraoka, N., Ohishi, Y., Seto, Y.,
640 2011. Spin crossover, structural change, and metallization in NiAs-type FeO
641 at high pressure. *Phys. Rev. B* 84, 134417.

642 Ozawa, H., Hirose, K., Tateno, S., Sata, N., Ohishi, Y., 2010. Phase transition
643 boundary between B1 and B8 structures of FeO up to 210 GPa. *Phys. Earth*
644 *Planet. Inter.* 179, 1157–163.

645 Panero, W.R., Jeanloz, R., 2001. Temperature gradients in the laser-heated
646 diamond anvil cell. *J. Geophys. Res.* 106, 6493–6498.

647 Prakapenka, V.B., Kuba, A., Kuznetsov, A., Kaskin, A., Shkurikhin, O., Dera,
648 P., Rivers, M.L., Sutton, S.R., 2008. Advanced flat top laser heating system
649 for high pressure research at GSECARS: application to the melting behavior
650 of germanium. *High Press. Res.* 28, 225–235.

651 Richet, P., Mao, H.k., Bell, P.M., 1989. Bulk Moduli of Magnesio-wüstites From
652 Static Compression Measurements. *J. Geophys. Res.* 94, 3037–3045.

653 Ringwood, A.E., 1975. Bulk Moduli of Magnesio-wüstites From Static
654 Compression Measurements. McGraw-Hill, New York.

655 Rivers, M., Prakapenka, V.B., Kubo, A., Pullins, C., Holl, C.M., Jacobsen, S.D.,
656 2008. The COMPRES/GSECARS gas-loading system for diamond anvil cells
657 at the Advanced Photon Source. *High Press. Res.* 28, 273–292.

658 Rost, S., Garnero, E.J., Williams, Q., 2006. Fine-scale ultralow-velocity zone
659 structure from high-frequency seismic array data. *J. Geophys. Res.* 111,
660 B09310.

661 Seagle, C., Heinz, D., Campbell, A., Prakapenka, V., Wanless, S., 2008. Melting
662 and thermal expansion in the Fe–FeO system at high pressure. *Earth Planet.*
663 *Sci. Lett.* 265, 655–665.

664 Shen, G., Rivers, M.L., Wang, Y., Sutton, S.R., 2001. Laser heated diamond
665 cell system at the Advanced Photon Source for *in situ* x-ray measurements
666 at high pressure and temperature. *Rev. Sci. Instrum.* 72, 1273–1282.

667 Shim, S., Duffy, T.S., Shen, G., 2000. The stability and P–V–T equation of
668 state of CaSiO₃ perovskite in the Earth’s lower mantle. *J. Geophys. Res.* 105,
669 25,955–25,968.

670 Shu, J., Mao, H.k., Hu, J., Fei, Y., Hemley, R.J., 1998. Single-Crystal X-ray
671 Diffraction of Wüstite to 30 GPa Hydrostatic Pressure. *N. Jb. Miner. Abh.*
672 172, 309–323.

673 Speziale, S., Milner, A., Lee, V.E., Clark, S.M., Pasternak, M.P., Jeanloz, R.,
 674 2005. Iron spin transition in Earth's mantle. *Proc. Natl. Acad. of Sci.* 102,
 675 17918–17922.

676 Sturhahn, W., 2000. CONUSS and PHOENIX: Evaluation of nuclear resonant
 677 scattering data. *Hyperfine Interactions* 125, 149–172.

678 Sturhahn, W., 2015. www.nrixs.com.

679 Sturhahn, W., Jackson, J.M., 2007. Geophysical applications of nuclear resonant
 680 spectroscopy. *GSA Special Papers* 421, 157.

681 Sun, D., Helmberger, D.V., Jackson, J.M., Clayton, R.W., Bower, D.J., 2013.
 682 Rolling hills on the core–mantle boundary. *Earth Planet. Sci. Lett.* 361, 333–
 683 342.

684 Tackley, P.J., 2012. Dynamics and evolution of the deep mantle resulting from
 685 thermal, chemical, phase and melting effects. *Earth-Science Reviews* 110,
 686 1–25.

687 Tange, Y., Kuwayama, Y., Irifune, T., Funakoshi, K.i., Ohishi, Y., 2012. P - V - T
 688 equation of state of MgSiO_3 perovskite based on the MgO pressure scale: A
 689 comprehensive reference for mineralogy of the lower mantle. *J. Geophys. Res.*
 690 117, B06201.

691 Tange, Y., Nishihara, Y., Tsuchiya, T., 2009. Unified analyses for P - V - T equa-
 692 tion of state of MgO : A solution for pressure-scale problems in high P - T
 693 experiments. *J. Geophys. Res.* 114, B03208.

694 Westrenen, W.v., Li, J., Fei, Y., Frank, M.R., Hellwig, H., Komabayashi, T.,
 695 Mibe, K., Minarik, W.G., Orman, J.A.V., Watson, H.C., Funakoshi, K.i.,
 696 Schmidt, M.W., 2005. Thermoelastic properties of $(\text{Mg}_{0.64}\text{Fe}_{0.36})\text{O}$ ferroperi-
 697 clase based on in situ X-ray diffraction to 26.7 GPa and 2173 K. *Phys. Earth*
 698 *Planet. Inter.* 151, 163–176.

- 699 Wicks, J.K., 2013. Sound Velocities and Equation of State of iron-rich (Mg,Fe)O.
700 Ph.D. thesis. California Institute of Technology.
- 701 Wicks, J.K., Jackson, J.M., Sturhahn, W., 2010. Very low sound velocities
702 in iron-rich (Mg,Fe)O: Implications for the core-mantle boundary region.
703 Geophys. Res. Lett. 37, L15304.
- 704 Williams, Q., Revenaugh, J., Garnero, E., 1998. A Correlation Between Ultra-
705 Low Basal Velocities in the Mantle and Hot Spots. Science 281, 546.
- 706 Zhang, J., 2000. Effect of pressure on the thermal expansion of MgO up to 8.2
707 GPa. Phys. Chem. Min. 27, 145–148.
- 708 Zhang, J., Kostak, P., 2002. Thermal equation of state of magnesiowüstite
709 (Mg_{0.6}Fe_{0.4})O. Phys. Earth Planet. Inter. 129, 301–311.
- 710 Zhang, J., Zhao, Y., 2005. Effects of defect and pressure on the thermal expan-
711 sivity of Fe_XO. Phys. Chem. Min. 32, 241–247.

712 **Appendix A. Parameter fitting**

713 Here we describe the details of the thermal equation of state parameter fit-
714 ting. These details are also provided in the complete manual to MINUTI, pro-
715 vided by Sturhahn (2015). The input data for the seos executable in MINUTI
716 are specified as N sets $\{p_i, V_i, T_i, \delta p_i, \delta V_i, \delta T_i\}$ consisting of pressure, volume,
717 temperature, and their uncertainties. The EOS is fitted to these data by varia-
718 tion of n parameters $\{x_j\}$ by minimizing the normalized mean-square deviation
719 (method of weighted least squares), with the reduced χ^2 described by

$$\chi^2 = \frac{1}{(N + p - n)} \left\{ \sum_{i=1}^N w_i (V_i - V(p_i, T_i, \{x_j\}))^2 + \sum_{k=1}^p \frac{(x_k - X_k)^2}{\delta^2 X_k} \right\} . \quad (\text{A.1})$$

Some number p of the fit parameters may have priors X_k with uncertainties δX_k . The weights w_i are determined by uncertainties of the data via

$$w_i^{-1} = \delta^2 V_i + \left(\frac{\partial V}{\partial p} \right)_T^2 \delta^2 p_i + \left(\frac{\partial V}{\partial T} \right)_p^2 \delta^2 T_i + \delta^2 V_i + V_i^2 \left(\frac{\delta^2 p_i}{K_i^2} + \alpha_i^2 \delta^2 T_i \right) , \quad (\text{A.2})$$

where K_i and α_i are bulk modulus and thermal expansion calculated from the EOS at $\{p_i, T_i\}$, respectively. If data and priors constrain the fit parameters reasonably well a solution with minimal χ^2 is obtained. These optimal parameters have errors related to data variation and prior uncertainties. The errors are estimated as variances (square errors) by

$$\delta x_j = \sqrt{\sigma_{jj}} \quad \text{with} \quad \sigma_{jj'} = \chi_{min}^2 \left[\sum_{i=1}^N w_i \frac{\partial V(p_i, T_i)}{\partial x_l} \frac{\partial V(p_i, T_i)}{\partial x_{l'}} + \delta_{ll'} \frac{1}{\delta^2 X_l} \right]_{jj'}^{-1} , \quad (\text{A.3})$$

where $\sigma_{jj'}$ is the covariance matrix and $[\dots]^{-1}$ denotes matrix inversion. The derivatives are calculated with the optimal parameters. The fit-parameter correlation matrix is given by

$$c_{jj'} = \frac{\sigma_{jj'}}{\sqrt{\sigma_{jj}\sigma_{j'j'}}} . \quad (\text{A.4})$$

The variance of a function F of the fit parameters can be calculated as

$$\delta^2 F(\{x_j\}) = \sum_{jj'=1}^n \sigma_{jj'} \frac{\partial F}{\partial x_j} \frac{\partial F}{\partial x_{j'}} \quad (\text{A.5})$$

The fit-parameter correlation matrix for each fitting result is reported in Appendix C.

Appendix B. EOS parameters and their uncertainties

The isothermal bulk modulus is calculated by direct differentiation of the EOS

$$K(p, T) = - \left(\frac{\partial p}{\partial \ln V} \right)_T . \quad (\text{B.1})$$

736 The seismic velocity is calculated via

$$v_\phi(p, T) = \sqrt{(1 + \alpha \gamma T) (K/\rho_0) (V/V_0)} \quad , \quad (\text{B.2})$$

737 where α is the thermal expansion coefficient, γ is the is the Debye-Grüneisen
 738 parameter, ρ_0 is the reference density. If fit parameters are defined the variances
 739 are calculated using (A.5) with the derivatives

$$\frac{1}{K} \frac{\partial K}{\partial x_j} = \frac{1}{V} \frac{\partial V}{\partial x_j} + \frac{K}{V} \frac{\partial}{\partial p} \left(\frac{\partial V}{\partial x_j} \right) \quad (\text{B.3})$$

740 and

$$\frac{1}{v_\phi} \frac{\partial v_\phi}{\partial x_j} = \frac{1}{V} \frac{\partial V}{\partial x_j} + \frac{K}{2V} \frac{\partial}{\partial p} \left(\frac{\partial V}{\partial x_j} \right) \quad . \quad (\text{B.4})$$

741 Thermal expansion is calculated according to

$$\alpha = \frac{1}{V} \left(\frac{\partial V}{\partial T} \right)_p = -\frac{1}{V} \left(\frac{\partial S}{\partial p} \right)_T \quad . \quad (\text{B.5})$$

742 The thermal contribution to the entropy S is given by

$$S(V, T) = -3k_B \ln \left(2 \sinh \frac{\Theta}{2T} \right) + 4k_B I \left(\frac{\Theta}{T} \right) + k_B \frac{3\Theta}{2T} \quad , \quad (\text{B.6})$$

743 where Θ is the Debye temperature and I was defined in Eq. (6). If fit
 744 parameters are defined the variance of α is calculated using Eq. A.5 with the
 745 derivative

$$\frac{\partial \alpha}{\partial x_j} = -\frac{\alpha}{V} \frac{\partial V}{\partial x_j} + \frac{1}{V} \frac{\partial}{\partial p} \left(\frac{\partial S}{\partial x_j} \right) \quad . \quad (\text{B.7})$$

746 **Appendix C. Correlation matrices**

747 In the fitting procedure of seos, the correlation matrices are provided for all
748 fitted parameters (Equation A.4). The resulting correlation matrices from the
749 fitting of the four different data sets from the two experiments (buffered and
750 unbuffered) are reported here.

Table C.3: $B1-Mw94$ in the buffered experiment using Fe as a pressure marker (Table 1).

	γ_0	V_0	K_{0T}	K'_{0T}
γ_0	1.000	-0.416	-0.001	0.112
V_0	-0.416	1.000	-0.897	0.799
K_{0T}	-0.001	-0.897	1.000	-0.973
K'_{0T}	0.112	0.799	-0.973	1.000

Table C.4: $B1-Mw94$ in the buffered experiment using NaCl as a pressure marker (Table 2).

	γ_0	V_0	K_{0T}	K'_{0T}
γ_0	1.000	-0.458	0.045	0.050
V_0	-0.458	1.000	-0.896	0.807
K_{0T}	0.045	-0.896	1.000	-0.974
K'_{0T}	0.050	0.807	-0.974	1.000

Table C.5: Rhombohedrally-structured Mw94 ($r-Mw94$) in the buffered experiment at 300 K, using Fe as a pressure marker (Table 1).

	V_0	K_{0T}	K'_{0T}
V_0	1.000	-0.909	0.558
K_{0T}	-0.909	1.000	-0.843
K'_{0T}	0.558	-0.843	1.000

Table C.6: *B1–Mw94* in the unbuffered experiment using NaCl as a pressure marker (Table 2).

	γ_0	V_0	K_{0T}	K'_{0T}
γ_0	1.000	-0.337	-0.162	0.244
V_0	-0.337	1.000	-0.866	0.798
K_{0T}	-0.162	-0.866	1.000	-0.987
K'_{0T}	0.244	0.798	-0.987	1.000

Table C.7: *B1–Mw94* in the unbuffered experiment using NaCl as a pressure marker, with $K_{0T} = 190 \pm 15$ GPa as a prior (Table 2).

	γ_0	V_0	K_{0T}	K'_{0T}
γ_0	1.000	-0.602	-0.083	0.299
V_0	-0.602	1.000	-0.724	0.496
K_{0T}	-0.083	-0.724	1.000	-0.945
K'_{0T}	0.299	0.496	-0.945	1.000

Table D.8: Pressure-volume-temperature data for the buffered experiment. (Mg_{0.06}Fe_{0.94})O volumes were refined using at least three of the following four reflections: 111, 200, 220, and 311. *hcp*-Fe volumes were calculated using at least 6 of the following 8 reflections: 100, 200, 101, 102, 110, 103, 200, and 112. *B2*-NaCl volumes were refined using at least 5 of the 6 following reflections: 100, 110, 111, 200, 210, and 211. Ne volumes were calculated from at least one of the two reflections: 111, 200. ^aPressure was determined by the equation of state of *hcp*-Fe from Dewaele et al. (2006) and Murphy et al. (2011). ^bPressure was determined from the equation of state of *B2*-NaCl (Fei et al., 2007b). ^cPressure was determined from the equation of state of Ne (Dewaele et al., 2008). ^dNo error bar is given due to too few reflections. Temperature uncertainties in this table reflect the standard deviation (note that these values were not used in the EOS fitting; see text for details).

P_{Fe}^a (GPa)	P_{NaCl}^b (GPa)	P_{Ne}^c (GPa)	T (K)	V_{Mw94e} (Å ³)	V_{Fe} (Å ³)	V_{NaCl} (Å ³)	V_{Ne}^d (Å ³)
116.5(7)	111.7(5)	116.1	1046(37)	6.935(26)	16.50(2)	20.13(2)	25.52
118.2(7)	113.1(5)	118.6	1290(28)	6.943(19)	16.50(2)	20.11(3)	25.48
118.9(7)	113.5(4)	118.4	1406(35)	6.952(20)	16.51(2)	20.12(2)	25.55
120.1(8)	114.7(5)	119.7	1611(52)	6.954(19)	16.53(2)	20.11(3)	25.56
122.0(7)	115.7(4)	121.7	1912(31)	6.950(21)	16.55(2)	20.14(2)	25.59
102.7(4)	98.9(3)	102.5	1143(7)	7.170(14)	16.90(1)	20.86(2)	26.48
103.1(4)	99.2(3)	102.3	1223(6)	7.174(16)	16.90(1)	20.87(2)	26.54
104.0(4)	99.7(3)	103.6	1397(14)	7.191(16)	16.92(1)	20.89(2)	26.54
105.5(4)	100.9(4)	104.6	1640(104)	7.188(14)	16.94(1)	20.89(2)	26.60
106.3(4)	101.4(3)	105.6	1744(40)	7.191(17)	16.94(1)	20.89(2)	26.59
107.8(4)	102.3(3)	106.5	2003(46)	7.197(18)	16.97(1)	20.90(2)	26.67
95.5(4)	89.6(1.7)	95.5	1237(4)	7.257(28)	17.13(1)	21.48(11)	27.07
96.4(4)	90.8(1.3)	95.9	1416(3)	7.286(13)	17.16(1)	21.45(9)	27.15

97.3(3)	91.5(1.5)	96.6	1606(19)	7.296(15)	17.18(1)	21.47(10)	27.20
98.7(3)	92.1(1.5)	97.3	1832(21)	7.343(12)	17.19(1)	21.50(10)	27.29
90.9(2)	87.1(3)	91.7	1279(38)	7.403(25)	17.29(1)	21.66(2)	27.40
91.3(2)	87.4(3)	91.8	1396(6)	7.432(25)	17.31(1)	21.68(2)	27.47
92.3(2)	88.3(3)	92.8	1578(36)	7.428(33)	17.33(1)	21.68(2)	27.50
94.3(2)	89.5(3)	93.6	1891(34)	7.449(18)	17.36(1)	21.69(2)	27.65
95.1(2)	90.1(3)	93.6	2036(36)	7.451(16)	17.37(1)	21.70(2)	27.74
94.9(2)	90.3(3)	94.0	2109(81)	7.429(13)	17.40(1)	21.71(2)	27.76
85.4(4)	82.4(5)	86.4	1208(1)	7.520(27)	17.45(1)	21.98(4)	27.82
85.5(4)	82.4(5)	86.3	1287(0)	7.520(27)	17.48(2)	22.00(3)	27.89
85.8(3)	83.0(4)	86.5	1407(37)	7.524(25)	17.50(1)	22.00(3)	27.95
86.4(4)	83.9(3)	87.1	1559(14)	7.539(29)	17.52(1)	21.98(2)	28.00
77.7(5)	75.8(4)	79.6	1106(9)	7.647(17)	17.70(2)	22.44(3)	28.40
78.4(5)	76.3(3)	80.2	1248(4)	7.661(20)	17.71(2)	22.45(3)	28.44
80.2(4)	77.5(4)	81.6	1510(16)	7.663(16)	17.73(2)	22.45(3)	28.49
81.0(5)	78.2(3)	82.5	1694(58)	7.664(14)	17.75(2)	22.47(3)	28.54
76.6(3)	73.2(3)	77.6	1215(5)	7.704(9)	17.77(1)	22.69(2)	28.68
77.5(4)	74.3(3)	78.2	1351(6)	7.714(8)	17.78(1)	22.65(2)	28.72
78.4(3)	74.9(3)	79.7	1516(26)	7.721(9)	17.79(1)	22.67(2)	28.70
79.2(3)	75.5(3)	80.0	1628(24)	7.724(8)	17.80(1)	22.66(3)	28.75
80.1(3)	76.3(3)	81.4	1811(63)	7.737(7)	17.83(1)	22.67(3)	28.75
81.9(4)	77.4(3)	82.4	2048(12)	7.744(10)	17.84(1)	22.66(2)	28.83
73.4(3)	69.8(3)	73.5	1253(1)	7.799(21)	17.90(1)	23.00(3)	29.14
74.0(3)	70.0(4)	73.7	1362(9)	7.805(22)	17.91(1)	23.02(3)	29.22
74.9(3)	70.4(4)	74.9	1506(9)	7.807(25)	17.92(1)	23.05(4)	29.20
76.1(3)	71.3(3)	75.8	1706(16)	7.809(19)	17.94(1)	23.04(3)	29.27
77.9(2)	72.3(3)	76.5	1960(43)	7.813(16)	17.97(1)	23.06(2)	29.40
77.4(2)	72.5(3)	76.5	1922(40)	7.814(13)	17.97(1)	23.03(3)	29.38
61.0(5)	59.1(7)	63.2	1242(4)	8.047(20)	18.41(2)	24.02(7)	30.40
61.5(4)	59.5(5)	63.9	1340(25)	8.049(18)	18.42(2)	24.03(6)	30.40

62.5(4)	60.4(4)	65.0	1527(6)	8.057(18)	18.45(2)	24.01(4)	30.44
64.0(3)	61.2(4)	66.2	1761(12)	8.061(15)	18.47(1)	24.04(4)	30.51
65.2(2)	62.0(3)	67.7	1959(31)	8.065(17)	18.50(1)	24.06(4)	30.50
54.5(6)	52.6(4)	56.3	1223(4)	8.214(11)	18.70(3)	24.74(5)	31.38
55.3(6)	53.1(5)	57.0	1327(0)	8.216(9)	18.71(3)	24.74(6)	31.40
57.0(5)	54.3(6)	58.2	1570(2)	8.220(8)	18.72(3)	24.72(7)	31.48
57.7(5)	54.5(5)	59.3	1704(16)	8.227(6)	18.74(2)	24.76(6)	31.46
58.6(4)	55.2(5)	60.0	1830(4)	8.229(5)	18.75(2)	24.75(6)	31.49
59.3(4)	55.4(6)	60.3	1918(7)	8.233(15)	18.76(2)	24.77(7)	31.55
52.3(4)	49.5(3)	52.4	1523(8)	8.326(11)	18.94(2)	25.29(4)	32.40
51.2(4)	48.7(3)	51.5	1319(1)	8.316(11)	18.91(2)	25.27(4)	32.30
53.9(3)	50.2(2)	53.6	1740(52)	8.337(10)	18.95(2)	25.31(3)	32.47
44.3(6)	41.5(5)	45.3	1336(25)	8.562(26)	19.28(4)	26.30(8)	33.53
43.7(8)	41.6(5)	45.2	1354(4)	8.565(22)	19.32(5)	26.30(7)	33.59
44.4(8)	41.9(4)	45.4	1451(28)	8.561(17)	19.33(4)	26.32(6)	33.69
46.4(5)	42.8(4)	46.6	1672(42)	8.558(12)	19.32(3)	26.32(7)	33.77
47.4(5)	43.4(4)	47.3	1822(49)	8.560(13)	19.34(3)	26.33(6)	33.85
48.1(5)	43.6(3)	47.8	1902(110)	8.566(14)	19.34(3)	26.34(5)	33.85
38.7(4)	37.0(4)	38.8	1366(16)	8.693(7)	19.63(2)	27.07(7)	35.12
39.2(4)	36.9(2)	38.9	1431(18)	8.694(7)	19.63(3)	27.12(4)	35.22
39.8(4)	37.4(3)	39.3	1521(33)	8.703(7)	19.64(3)	27.11(6)	35.28
42.0(4)	38.3(3)	40.9	1794(8)	8.719(7)	19.65(3)	27.15(6)	35.36
42.7(3)	38.8(4)	42.0	1906(90)	8.722(20)	19.66(2)	27.13(7)	35.27

Table D.9: Pressure-volume-temperature data for the unbuffered experiment. (Mg_{0.06}Fe_{0.94})O volumes were refined using the three reflections: 111, 200, and 220. B2-NaCl volumes were refined using at least 3 of the 4 following reflections: 100, 110, 111, and 200. Ne volumes were calculated from at least one of the two reflections: 111, 200. ^aPressure was determined from the equation of state of B2-NaCl (Fei et al., 2007b). ^bPressure was determined from the equation of state of Ne (Dewaele et al., 2008). ^cNo error bar is given due to too few reflections. Temperature uncertainties in this table reflect the standard deviation (note that these values were not used in the EOS fitting; see text for details).

P_{NaCl}^a (GPa)	P_{Ne}^b (GPa)	T (K)	V_{Mw94} ($\text{\AA}^3$)	V_{NaCl} ($\text{\AA}^3$)	V_{Ne}^c ($\text{\AA}^3$)
71.5(4)	73.1	1227(11)	7.77(2)	22.84(3)	29.17
71.7(6)	73.2	1312(14)	7.78(2)	22.85(5)	29.23
72.0(5)	73.5	1442(28)	7.79(2)	22.88(4)	29.30
71.4(4)	70.2	1208(23)	7.77(2)	22.85(3)	28.77
71.7(4)	72.8	1317(27)	7.78(2)	22.86(4)	29.19
71.9(4)	72.9	1413(26)	7.78(2)	22.88(4)	29.26
69.5(5)	71.0	1192(44)	7.86(3)	23.00(5)	29.38
70.0(3)	70.6	1306(62)	7.85(2)	23.01(3)	29.52
70.5(3)	70.9	1433(19)	7.86(1)	23.01(3)	29.61
69.9(3)	68.1	1173(15)	7.83(2)	22.96(3)	29.00
70.1(3)	70.3	1279(36)	7.84(2)	22.98(2)	29.45
70.5(4)	70.5	1396(49)	7.85(2)	23.00(3)	29.52
67.4(7)	69.3	1226(28)	7.89(6)	23.21(7)	29.61
67.7(6)	69.1	1293(17)	7.89(5)	23.21(6)	29.69
68.0(5)	69.0	1403(20)	7.90(5)	23.22(5)	29.80
67.4(5)	68.9	1180(23)	7.88(5)	23.18(5)	29.61
67.8(5)	69.3	1306(7)	7.89(5)	23.20(5)	29.68
67.9(6)	69.3	1351(15)	7.89(5)	23.21(5)	29.72

68.1(5)	69.3	1420(16)	7.90(4)	23.22(5)	29.78
63.3(2)	65.8	1216(7)	7.96(4)	23.58(2)	30.04
63.6(2)	63.8	1295(2)	7.97(4)	23.59(2)	30.38
64.1(3)	64.9	1412(17)	7.97(3)	23.60(3)	30.34
63.6(1)	65.7	1212(10)	7.95(3)	23.55(1)	30.04
63.8(2)	66.2	1289(11)	7.96(3)	23.57(2)	30.05
64.1(3)	65.6	1430(13)	7.97(3)	23.60(3)	30.27
60.5(4)	61.0	1215(27)	8.05(5)	23.86(4)	30.68
61.1(1)	62.1	1335(14)	8.05(4)	23.86(1)	30.64
61.4(3)	59.7	1424(15)	8.05(3)	23.87(3)	31.08
62.1(7)	62.6	1446(23)	8.03(1)	23.81(7)	30.69
57.6(2)	59.0	1203(24)	8.13(4)	24.16(3)	30.95
58.0(3)	57.9	1317(40)	8.13(3)	24.18(3)	31.24
58.4(2)	57.5	1424(43)	8.13(3)	24.19(2)	31.42
58.3(3)	58.3	1390(20)	8.13(3)	24.18(3)	31.26
57.9(2)	59.4	1207(36)	8.11(3)	24.13(2)	30.90
58.1(2)	57.7	1308(12)	8.12(3)	24.16(2)	31.26
58.4(1)	57.8	1382(19)	8.12(3)	24.16(1)	31.33
58.8(4)	59.4	1451(37)	8.13(3)	24.16(5)	31.17
55.7(2)	56.5	1245(22)	8.18(3)	24.39(2)	31.38
56.2(4)	54.4	1337(18)	8.18(2)	24.39(4)	31.83
56.1(2)	55.1	1371(16)	8.18(2)	24.41(3)	31.75
56.1(3)	56.1	1386(32)	8.17(2)	24.41(3)	31.60
55.9(5)	55.5	1270(23)	8.15(3)	24.38(6)	31.57
56.1(2)	55.6	1346(18)	8.16(3)	24.40(2)	31.64
56.4(3)	55.6	1422(15)	8.17(2)	24.40(3)	31.73
54.0(1.3)	55.6	1224(38)	8.21(4)	24.58(16)	31.50
54.6(1.1)	55.5	1396(42)	8.22(3)	24.59(13)	31.70
55.4(1.2)	55.3	1579(16)	8.23(2)	24.59(14)	31.96
55.7(1.0)	55.9	1670(24)	8.23(1)	24.60(12)	31.98

55.8(1.0)	55.5	1731(9)	8.24(1)	24.63(11)	32.12
54.4(7)	54.9	1140(21)	8.15(8)	24.49(8)	31.51
54.7(6)	54.7	1239(6)	8.16(6)	24.51(7)	31.65
55.4(6)	54.8	1437(16)	8.18(3)	24.52(6)	31.88
55.3(6)	54.1	1483(53)	8.22(1)	24.56(6)	32.05
46.8(1.1)	48.6	1195(20)	8.39(4)	25.45(15)	32.67
47.6(1.0)	49.4	1398(34)	8.40(3)	25.46(13)	32.79
48.2(8)	48.7	1591(24)	8.41(2)	25.50(11)	33.19
48.8(8)	50.0	1760(26)	8.42(1)	25.51(10)	33.19
45.2(1.1)	48.8	1248(24)	8.42(5)	25.71(15)	32.71
45.9(1.1)	49.1	1429(15)	8.42(4)	25.72(16)	32.90
46.5(1.1)	48.8	1613(15)	8.44(3)	25.75(15)	33.21
46.8(1.0)	51.3	1705(35)	8.45(2)	25.75(14)	32.84
41.3(1.0)	44.3	1186(32)	8.55(2)	26.23(15)	33.52
42.0(8)	45.9	1387(6)	8.56(2)	26.26(13)	33.48
42.7(8)	46.2	1561(33)	8.57(2)	26.27(13)	33.69
42.8(8)	45.7	1584(12)	8.57(2)	26.27(12)	33.83
38.6(1.0)	39.1	1265(35)	8.62(4)	26.72(16)	3.27
39.1(9)	39.3	1398(37)	8.63(4)	26.72(15)	3.27
39.1(9)	39.4	1395(23)	8.63(4)	26.72(15)	3.27
39.7(8)		1562(38)	8.65(4)	26.75(14)	
39.8(8)		1613(28)	8.65(4)	26.76(14)	
38.4(7)	39.0	1165(51)	8.60(3)	26.69(12)	3.26
39.2(9)		1436(45)	8.63(3)	26.74(15)	
39.7(8)		1594(17)	8.65(4)	26.76(14)	
34.6(6)	35.5	1207(77)	8.72(1)	27.37(11)	35.75
35.0(7)	35.9	1295(68)	8.73(1)	27.36(12)	35.78
35.3(6)	36.4	1384(50)	8.76(1)	27.38(12)	35.82
35.5(6)	37.2	1475(25)	8.77(1)	27.40(11)	35.77
33.2(9)	34.1	1224(31)	8.76(2)	27.64(17)	36.20

33.6(7)	35.3	1454(41)	8.79(1)	27.74(13)	36.32
34.3(1.1)	36.4	1630(31)	8.81(1)	27.76(22)	36.35
31.9(7)	34.1	1235(36)	8.75(7)	27.90(15)	36.22
32.2(9)	33.9	1384(65)	8.78(5)	27.97(19)	36.59
32.4(8)	34.4	1526(21)	8.82(4)	28.04(17)	36.75
32.4(5)	33.9	1391(60)	8.79(4)	27.93(9)	36.62
32.8(4)	34.3	1510(31)	8.81(4)	27.95(8)	36.74

752 Appendix E. Unexplained Peaks

753 As described in the text, weak unexplained peaks were observed in our un-
 754 buffered dataset (Figure 1). It has been proposed that vacancy concentrations
 755 in non-stoichiometric FeO-bearing samples are reduced with the exsolution of
 756 (Fe,Mg)Fe₂O₄ at high pressure and temperature, according to

$$(Mg_xFe_y)O \rightarrow a(Mg_{x'}Fe_{1-x'})Fe_2O_4 + b(Mg_{x''}Fe_{y''})O \quad (E.1)$$

757 where $(x'' + y'') > (x + y)$, ensuring the stoichiometry of the (Mg,Fe)O phase
 758 regardless of oxygen fugacity (Zhang and Zhao, 2005; McCammon et al., 1998;
 759 Fei, 1996).

760 In Figure E.10, measured XRD patterns are compared to candidate oxides
 761 measured in other studies. As the chemical composition of the Mw94 at the
 762 P-T conditions sampled in this study is not known, it could be possible that the
 763 unidentified peaks belong to a phase with composition (Mg,Fe)₃O₄, but they
 764 do not match that of MgFe₂O₄ or *h*-Fe₃O₄. They are consistent, however, with
 765 peaks of Fe₄O₅. We do not see any of the remaining several peaks predicted
 766 for this phase, the most intense of which fall under ours (8 degrees), which may
 767 rule it out as a potential candidate. More dedicated studies need to be carried
 768 out to precisely determine both the composition, identity, and phase relations
 769 of exsolution products before definite claims can be made.

Figure E.10: Example integrated XRD patterns for the unbuffered dataset showing peak identifications for $(\text{Mg}_{0.06}\text{Fe}_{0.94})\text{O}$, $B2\text{-NaCl}$, and Ne. Pressures were determined using the equation of state of $B2\text{-NaCl}$ (Fei et al., 2007b). $(\text{Mg}_{0.06}\text{Fe}_{0.94})\text{O}$ is cubic at high temperature (Mw_C , $Fm\text{-}3m$) and rhombohedral at room temperature upon quench (Mw_R , $R\text{-}3m$). In this quench pattern at 300 K, a mixture of cubic and rhombohedral Mw_{94} is observed. Unidentified peaks are labeled with *. For comparison, reflections of candidate iron oxides are shown below: MgFe_2O_4 measured at 31.6 GPa (Andrault and Bolfan-Casanova, 2001), $h\text{-Fe}_3\text{O}_4$ calculated at 31 GPa (Dubrovinsky et al., 2003), and Fe_4O_5 measured at 30 GPa (Lavina et al., 2011)