Strong seismic scatterers near the core-mantle boundary north of the Pacific Anomaly

Xiaolong Ma^{a,b}, Xinlei Sun^{a,c,*}, Douglas. A. Wiens^c, Lianxing Wen^{d,e}, Andrew Nyblade^f, Sridhar Anandakrishnan^f, Rick Aster^g, Audrey Huerta^h, Terry Wilsonⁱ

^aState Key Laboratory of Isotope Geochemistry, Guangzhou Institute of

Geochemistry, CAS

^bUniversity of Chinese Academy of Sciences, Beijing 10069, China

^cDepartment of Earth and Planetary Sciences, Washington University (St. Louis)

^dDepartment of Geosciences, State University of New York at Stony Brook

^eDepartment of Earth and Space Sciences, University of Science and Technology of China

^fDepartment of Geosciences, Pennsylvania State University

^gDepartment of Geosciences, Colorado State University

^hDepartment of Geological Sciences, Central Washington University

ⁱDepartment of Geological Sciences, Ohio State University

* Corresponding author Email address: xsun@gig.ac.cn

23 **Abstract**

24 Tomographic images have shown that there are clear high-velocity heterogeneities to
25 the north of the Pacific Anomaly near the core-mantle boundary (CMB), but the
26 detailed structure and origin of these heterogeneities are poorly known. In this study,
27 we analyze PKP precursors from earthquakes in the Aleutian Islands and Kamchatka
28 Peninsula recorded by seismic arrays in Antarctica, and find that these heterogeneities
29 extend ~400 km above the CMB and are distributed between 30-45°N in latitude. The
30 scatterers show the largest P-wave velocity perturbation of 1.0-1.2% in the center
31 (160-180°E) and ~0.5% to the east and west (140-160°E, 180-200°E). ScS-S
32 differential travel-time residuals reveal similar features. We suggest that these seismic
33 scatterers are the remnants of ancient subducted slab material. The lateral variations
34 may be caused either by different slabs, or by variations in slab composition resulting
35 from their segregation process.

36 **Keywords**

37 Seismic scatterers; PKP precursor; Subducted slabs; Lateral heterogeneities; Core-
38 mantle boundary

39

40 **1. Introduction**

41 The core-mantle boundary (CMB) is one of the most significant boundary layers
42 within the Earth. This layer and its adjacent regions, especially D'', are key to
43 understanding dynamic processes within the Earth, such as the source of mantle
44 plumes, the fate of subducted slabs, and material and heat exchange between the

mantle and core (Young and Lay, 1987; Wyssession et al., 1994; Lay et al., 1998; Garnero, 2000; Garnero, 2004; McNamara and Zhong, 2005; Lay and Garnero, 2011). Previous seismological studies have found complex heterogeneities near the CMB, such as Large Low Shear Velocity Provinces (LLSVPs) (e.g., Wen 2001; Ni et al, 2003; Wang and Wen, 2007; Garnero and McNamara, 2008; He and Wen, 2009; He and Wen, 2012), Ultra-Low-Velocity zones (ULVZ) (Garnero et al., 1999; Rost et al., 2006; Garnero et al, 2007; Rost and Garnero, 2010; Yao and Wen, 2014), anisotropy (e.g., Kendal and Silver, 1996; Lay et al., 1998; Garnero et al., 2004; Long, 2009), and seismic scatterers (e.g., Cleary and Haddon, 1972; Husebye and King, 1976; Vidale and Hedlin 1998; Thomas et al., 1999; Hedlin and Shearer, 2000; Cao and Romanowicz, 2007; Vanacore et al., 2010). Although high-velocity anomaly regions are well studied and are commonly attributed to subducted slabs, many regions, because of inadequate sampling by conventional seismic phases, are poorly imaged. In addition, conventional tomographic methods often poorly resolve small-scale structures due to limited frequency content. In this case, unconventional methods, such as those specifically focusing on scattered waves, may provide additional information. For the lowermost mantle, PKP precursors are possible candidates to serve this purpose.

PKP precursors are P waves that are scattered by small-scale elastic heterogeneities in the mantle and/or topographic irregularities on the CMB (e.g., Cleary and Haddon, 1972; Haddon and Cleary, 1974; Doornbos, 1976; Doornbos, 1978; Bataille and Flatté, 1988; Cormier, 1995; Hedlin and Shearer, 2000). Because

of the geometry ray-paths of these seismic phases, the scattered P waves can precede the core phase (PKIKP) by up to 20s and are typically best recorded at a distance range between 120-142° (Fig. 1). Particular advantages of using PKP precursors are that they are sensitive to heterogeneity at a scale as small as ~10 km, and are not contaminated by the coda of other phases due to their earlier arrival times. There have been successful applications of PKP precursors studies on global scales (e.g., [Hedlin et al., 1997](#); [Cormier, 1999](#); [Margerin and Nolet, 2003b](#); [Mancinelli and Shearer, 2013](#)), as well as regional scales (e.g., [Vidale and Hedlin 1998](#); [Wen and Helmberger, 1998](#); [Niu and Wen, 2001](#); [Miller and Niu, 2008](#); [Thomas et al., 2009](#); [Frost et al., 2013](#)). In regional studies that incorporate well-sited arrays of seismographs, greater information about the anomalies, such as their boundary locations and sharpness, may be resolvable.

In this study, we focus on the CMB region to the north of the Pacific Anomaly, a region where tomographic images consistently show large-scale high-velocity anomalies ([Li and Romanowicz, 1996](#); [Grand et al, 2002](#); [Zhao, 2004](#); [Li et al., 2008](#); [Simmons et al., 2010](#)) but where high resolution details are lacking. We use PKP precursors from earthquakes in the Aleutian Islands and Kamchatka Peninsula recorded by recently deployed seismic arrays in Antarctica to locate the scatterers and to investigate their velocity variations. Our results show that the observed precursors are caused by seismic scatterers in the lowermost mantle, and the strengths of these scatterers vary according to their locations. Furthermore, we also analyze ScS-S travel time residuals sampling the same region, and find that the results show similar

variation pattern. We suggest that the lateral variations in this region may be caused by heterogeneities from different subducted slabs, or by varying chemical compositions of the slab resulting from their segregation process.

2. CMB region heterogeneities from PKP precursors

To constrain small-scale heterogeneities near the CMB beneath the Pacific region, we collected PKP precursor data recorded by Antarctic seismic arrays (GAMNIS and POLENET-ANET array) from earthquakes in the Aleutian Islands and Kamchatka Peninsula (Fig. 2). The GAMSEIS and POLENET/ANET networks consist of temporary and semi-permanent seismic stations deployed in East and West Antarctica beginning in 2008 for understanding the structure and solid-earth ice sheet interactions of the continent (Heeszel et al., 2013; Accardo et al., 2014; Anthony et al., 2015). These new seismic stations provide much improved coverage of the CMB along paths not previously sampled due to sparse global station distribution in the far-southern latitudes. Here we use 59 Antarctic seismic stations (26 in GAMSEIS and 33 in POLENET/ANET) deployed during 2009-2011, and earthquakes with magnitude greater than 5.5 (Table1) at a distance range of 134-143°. All seismograms are band-pass filtered between 0.5-2.0 Hz, and only those with low noise levels and clear precursor signals are selected. PKP precursor travel times and envelope amplitudes are then analyzed to locate the scatterers and to calculate the velocity variations within them.

Fig. 3 shows representative PKP precursor waveforms from earthquakes in three different regions (western, middle and eastern region). The relative amplitudes of

PKP precursors in the middle region (central panel) are obviously much larger than those in the other two regions, and are larger than the PKIKP phases at some distances. Investigating the larger sample of earthquakes in each region, we find that the amplitudes have no relation with the earthquake depth (Table 1). By comparing the entry and exit points of PKIKP rays at the CMB (Fig. 2) from these earthquakes, it is found that the exit points of the earthquakes from different regions are intermixed, while the entry points are well separated into different locations.

2.1 Results from PKP arrival times

To determine the exact locations of the seismic scatterers, we adopt Wen's method (Wen, 2000) that utilizes the arrival times of PKP precursors. First, we divide the region of the possible scatterer locations (20°N to 50°N, 140°E to 200°E, and from the CMB to 600 km above the CMB) into six depth ranges with separation of 100 km. In each depth range, the region is further divided into a 1.0° by 1.0° uniform grid of nodes. Then we calculate the scatterer probability and hit count for each node by comparing the predicted and observed precursor arrival times of all earthquakes. The probability at a given node is the ratio of the number of seismic rays whose PKP precursor onsets sample this node over the total number of seismic rays in this study, and the hit count is the number of PKP precursors sampling this node. Detailed information about the method can be found in Wen (2000). In general, the grids with high probability and large hit counts are most likely where the PKP precursors originate.

We calculate the hit counts and probability near the CMB region for both source side and receiver side. We find that, at the same depth, both the hit counts and probability for receiver side are smaller than those of the source side. Considering together the fact that the piercing points of PKIKP at the CMB are well separated in source side while are intermixed with each other in receiver side (Fig. 2), and the precursor waveform differences in different locations, we suggest that the precursors are most likely from the CMB region of the source side, the region to the north of the Pacific Anomaly.

Fig. 4 shows results of the scatterer probability and hit counts calculations for the three earthquake clusters at four depths in source side: the CMB, and 200 km, 300 km, 400 km above the CMB. The scattering regions are approximately distributed in 30-45°N in latitude, and 140-160°E, 160-180°E, 180-200°E in longitude, respectively (Fig. 4). Because the scatterer probability and hit counts in our study become insignificant at scattering depth shallower than 400 km above the CMB, we suggest that the scatterers are most likely confined within the lowermost 400 km of the mantle.

2.2 Results from PKP precursor amplitude

PKP precursor amplitude is determined not only by the scatterer location, but also their geometry, lateral distribution and velocity perturbation. Although PKP precursor arrival time can locate where the scatterers are, they cannot give any information about the velocity contrast between the scatterers and the ambient mantle. To solve this problem, we look further into the precursor amplitude.

We assume that the PKP precursors are due to PKP propagation through a random inhomogeneous elastic media. Based on Chernov's acoustic scattering theory (Chernov, 1960) adapted to elastic media (Haddon and Cleary, 1974; Doornbos, 1976), and assume the mantle behaves as a Poisson solid with single scattering and an exponential geostatistical autocorrelation function (Wu and Aki, 1985), the average scattered power can be written as (Hedlin et al., 1997; Hedlin and Shearer, 2000):

$$\langle |\Phi_s(\theta)|^2 \rangle = \frac{2k^4 a^3 v^2 V A^2}{\pi r^2} \frac{\frac{1}{4} [\cos(\theta) + \frac{1}{3} + \frac{2}{3} \cos^2(\theta)]^2}{[1 + 4k^2 a^2 \sin^2(\theta/2)]^2} \quad (1)$$

where V is the scattering volume, v the rms velocity perturbation, A is the incident wave amplitude, k is the wavenumber, r is the distance between the scatterer and receiver, and θ is the scattering angle.

We develop our own code to calculate the precursor energy based on equation (1). In our calculation, AK135 model (Kennett et al., 1995) is used, and both PKP AB and BC phase are considered for the precursor energy. We also include all the refraction coefficients and attenuation factors.

In this study, the scatterer volume can be calculated based on the results of section 2.1. Previous studies have been empirically suggested that 6-10 km is the general scale length for scatterers within the Earth's lower mantle (e.g., Hedlin and Shearer, 2000; Helffrich and Wood, 2001; Mancinelli and Shearer, 2013). In order to compare with previous results, and because our observed precursors can only be seen at period less than 2 seconds, we choose 8 km as the scale length in our calculation.

Using these assumptions, velocity perturbation is the only unknown and can be constrained by trial and error.

Similar to the method described in Section 2.1, we first divide the scattering volume into thinner (20 km) depth layers, with each layer further divided into $0.5^\circ \times 0.5^\circ$ grids. The area of each layer is based on the results from Section 2.1, so the scattering areas are different at different depths. Then for each grid node on each layer, we calculate its energy power based on equation (1). The energy kernel of each layer is obtained by summing up the scattering power of all the corresponding grid nodes in that layer.

Because the PKIKP phase in the seismogram may be contaminated by precursors, and it is difficult to obtain P waves with similar azimuths as PKIKP due to earthquake-station geometry, we could not extract the PKIKP source time function from observed seismograms. Instead, we use Direct Solution Methods (DSM; [Geller et al., 1995](#); [Kawai and Geller, 2006](#)) to calculate the synthetic seismograms for every station and then stack PKIKP phases to estimate the source time function for each event. In the calculation, all the attenuation and refraction coefficients are considered. The inner core attenuation is set to be 360 ([Bhattacharyya et al., 1993](#)). After convolving the energy kernels with the source time functions (in power), we calculate the square root of the energy power and normalize it to the theoretical amplitude of PKIKP.

Fig.5 shows three representative seismograms from events located in the three different regions (east, middle and west). In this figure, the seismograms are binned

over 1° distance and synthetic curves are superimposed on them. Note that the synthetic amplitudes increase gradually with distance and travel time. For the eastern and western regions (longitude 140-160°E, 180-200°E), the best fit is a model with ~0.5% bulk perturbation in P-wave velocity (Fig. 5a, c), and in the middle region (longitude 160-180°E), the best model is a scattering volume with 1.0%-1.2% P-wave velocity perturbation (Fig.5b). The amplitude variations of the scatterers in the three sub-regions suggest these regions must have different characteristics.

3. Results from S-ScS travel times

S-wave velocities can also be investigated to study the lower mantle characteristics near the scatterer locations. We searched the global data systematically for both broadband and long period tangential S waveforms. The earthquakes we selected are from 1990 to 2014. To obtain high-quality data, we only chose earthquakes with magnitude greater than 5.5 and depths greater than 100 km. The distance range of the S data is between 50-85° (Fig. 6).

After removing instrument response and transferring them to ground motion, we band-pass filtered the broad and long period data with frequency bands 0.05-2 Hz and 0.01-0.1 Hz, respectively. We carefully check the quality of those waveforms and finally select 423 waveforms.

ScS-S residual travel-times are defined as:

$$(\text{ScS-S})_{\text{residual}} = (\text{ScS} - \text{S})_{\text{observed}} - (\text{ScS-S})_{\text{predicted}} \quad (2)$$

Although differential travel times can reduce the error from uncertainties of earthquake location, origin time, and crust and upper mantle heterogeneities, it can be

biased by heterogeneities from lower mantle (Wyssession et al., 1994; Wyssession et al., 2001). We correct both S and ScS travel time anomalies above 400 km from the CMB using the 3-D tomographic model GypSuM (Simmons et al., 2010), thus reducing the heterogeneity effect from the mantle above our study region.

Fig. 7 shows the relationship between S, ScS and ScS-S residuals. Obviously ScS residuals show strong correlation with ScS-S while S residuals show weak correlation. This relationship demonstrates that those differential ScS-S residuals are mainly from ScS path in the lowermost mantle, and negative residuals indicate the anomaly is from high-velocity heterogeneities.

We estimate S velocity variations based on the following equation

$$\delta v / v = -\delta t / t \quad (3)$$

where v is the averaged velocity, δt is the corrected travel-time residuals, and t is the total travel time of ScS through the bottom 400 km of the mantle.

Fig. 8 shows the distribution of averaged shear-velocity variations on ScS reflection points at the CMB. Although not all the scattering regions are well sampled, the three regions can still be distinguished and are sketched in the figure. The S-wave velocity variations are approximately 1.0%-1.5% on the eastern and western regions, while the middle region shows a different S velocity variation of 2.0-3.0%.

A possible concern is that the travel times may be influenced by the adjacent Pacific Anomaly. Since most of the S waves travel well above the Pacific Anomaly, this is only a concern for the ScS phases. Thus we carefully checked the ScS piercing points at 400 km above the CMB, and found that all of them bypass the Pacific

Anomaly's north boundary (Fig. 6). Therefore, the ScS travel time data are not affected by the Pacific Anomaly, and they show correct trend of the heterogeneities in the same region as the PKP precursors.

4. Discussion

4.1 Comparison with previous studies

There have been various studies on PKP precursors. In terms of global PKP precursor data, single scattering theory have proposed a model of $0.5\% \pm 0.1\%$ rms velocity perturbations in a 200 km layer above the CMB (Bataille and Flatté, 1988), or 1.0% rms velocity perturbation in the whole mantle (Hedlin and Shearer, 1997). On the other hand, multiple scattering theory has given different results. For example, Margerin and Nolet (2003b) suggested a model with 0.1% P velocity perturbations in the whole mantle, similar to the 0.2% whole mantle perturbation model proposed by Mancinelli and Shearer (2013). Margerin and Nolet (2003a) claimed that the single scattering theory could probably wrongly estimate the strength of the scatterer, and the limit of the rms velocity perturbation from single scatter would be 0.5%. Later, Mancinelli and Shearer (2013) found that the single and multiple scattering methods give similar results, and in all the scattered waves, single scattered waves dominate in number (90%).

Above studies are from averaged PKP precursors in global scale, so the PKP precursor amplitudes are weak. However, anomalous PKP precursors do exist in various regions, not only in amplitude, but also in shape (e.g., Vidale and Hedlin, 1998; Wen and Helmberger, 1998; Hedlin and Shearer, 2000; Vanacore et al., 2010;

Waszek et al, 2015). These anomalies may reflect regional heterogeneities, such as in our study.

To further compare single and multiple scattering methods, we used the phonon code (Shearer and Earle, 2004; Mancinelli and Shearer, 2013), which considers all the scattered waves, to model the precursor amplitudes of our data. We confine the scattering region in the 400 km above the CMB and set the scale length to 8 km, same as that in our single scattering modeling. The result suggests that P wave velocity perturbations of 0.2% and 0.5% on the two sides and the center region respectively fit the data well. Considering that we only confine depth in the multiple scattering modeling, the difference is understandable. We deduce that a more realistic regional scatterer should reduce the difference between these two methods. In general, the single and multiple scattering theory should produce similar results: on one hand, the single scattered waves dominate the scattered waves in number; on the other hand, the intrinsic attenuation may suppress the multiple scattered waves more.

Finally, strong scattering in the lithosphere would also affect the precursor amplitude relative to the peak PKIKP (Mancinelli and Shearer, 2013), and neglecting this effect may overestimate the rms velocity perturbation of the scatterer. However, the lateral variations of the scattering strength would remain the same in our study. Because this effect is beyond the content of this paper, we did not investigate deeply on this topic.

4.2 Origin of the scatterers

Examination of mantle tomographic models (e.g., [Grand et al, 2002](#); [Zhao, 2004](#); [Li et al., 2008](#); [Simmons et al., 2010](#)) shows that the heterogeneous regions identified in this study are part of the high-velocity anomalies near the CMB, as the ScS data have suggested. Although it has been long noticed that there is a “ring” of fast anomalies surrounding the slow anomalies in the center of Pacific (e.g., [Wyssession, 1996](#); [Maruyama et al 2007](#)), the detailed structure and origin of these features still remain ambiguous.

Using P diffracted waves, Wyssession ([1996](#)) inverted global large-scale heterogeneities near the CMB region, and found that northern Pacific rim region was coincident with the Panthalasa paleo-subduction. Based on geological and seismic tomographic data, Van der Meer et al ([2012](#)) reconstructed intra-oceanic arc subduction and suggested that a series of paleo-subduction zones subducted in the mid-Panthalasa ocean. Their reconstruction results show that the locations of the subduction zones ~200 Myr ago overlap partly with the scatterer locations we find in the lowermost mantle. On the other hand, Maruyama et al ([2007](#)) reconstructed the paleo-subduction location back to 180 Myr based on the hot spot frame and current plate movements. By comparing their results to the current tomographic structures in the mantle, they speculated that the high-velocity anomaly surrounding the Pacific Anomaly near the CMB was the graveyard of Rodinia subduction (0.75-1.0 Ga), since the high-velocity region does not correspond with the locations of Mesozoic subduction zones in their study.

Our results show that the fast regions to the north of the Pacific Anomaly have lateral variations, with central region having larger high velocity and two side regions having relatively smaller high velocities. Based on the previously described studies, we also suggest that the high velocity anomalies represent subducted paleo-slabs. A related question is how to interpret the different levels of velocity heterogeneity in the three sub-regions. One explanation is that the three sub-regions are from different paleo-slabs (Oku-Niikappu, Kolyma-Omolon, and Anadyr-Koryak plate, from left to right), and these slabs may have different thermal and subduction histories, as Van der Meer et al (2012) have suggested.

Alternatively, these sub-regions could also be from a single paleo-slab, where different parts of the slab were separated during the segregation/mixing process of the slab in the CMB (Christensen and Hoffman, 1994). For example, Lee and Chen (2007) proposed that when the slab subducts into the mantle, the oceanic crust could separate from the underlying lithospheric mantle due to a weak serpentinized zone inside the slab. Because the oceanic crust transformed to high-pressure phases has negative buoyancy in the mantle, it can sink into the lower mantle (Stixrude and Lithgow-Bertelloni, 2012). Moreover, the rheological differences between the recycled crust and surrounding mantle may defer the homogenization and lead to incomplete mixing (Metcalf et al., 1995; Helffrich and Wood, 2001; Van KeKen, 2002), making it possible for the subducted slab to remain in the CMB region for a long time. Furthermore, during the homogenization, the segregation of different minerals within the crust could also result in compositional heterogeneities (Sun et al.,

2011) because of low viscosity effects (here the Pacific Anomaly may serve this purpose). Thus, different compositions from a segregation process can explain our observed lateral variations within the fast anomalies.

The relative amplitude differences between PKP precursors and PKIKP are also affected by inner core attenuation (Wen and Niu, 2002; Waszek, et al. 2015) or topography at the CMB (Doornbos, 1978; Bataille and Flatté, 1988). To test the inner core effect, we checked PKIKP turning points of the earthquakes in the three sub-regions, and found that they were highly random and inter-mixed with each other. Thus the regional inner core attenuation variations could not be the cause. We also chose different Q value from 250 to 600 (Cormier and Xu, 1998; Yu and Wen, 2006) in PKIKP synthetics, and found that the change of the relative amplitude is ~20%, which will not affect our result greatly. For topography effect, the largest PKP precursor amplitude in our study, similar to that of Wen (2000), would require topography of several kilometers at the CMB. According to Wen (2000), that is incompatible with CMB dynamics. Overall, we contend that the lower mantle heterogeneity model described here provides a good explanation for the PKP precursor observations and is consistent with ScS travel times and mantle tomographic models.

Although we have both P and S velocity variations in the study region, we did not compare them directly. The reason is that our P-wave velocity perturbations are statistical variations within a range, but S-wave velocity variations are absolute average variations in the region. Thus it is not proper to compare them directly.

Investigation of the ratio of P and S anomalies would require similar observations and processing methods for both P and S waves.

5. Conclusions

We observe clear PKP precursors at Antarctica seismic stations from earthquakes in the Aleutian Islands and the Kamchacha Penninsula. By analyzing the onsets and amplitudes of these precursors, we find that there exist seismic heterogeneities in the lowermost mantle to the north of Pacific Anomaly, between latitude 30-45°N. These heterogeneities exhibit lateral variations: the middle region (longitude 160°-180°E) has small-scale P-wave velocity perturbations of ~0.5%, while on the two sides (140°-160°E and 180°-200°E), the P-wave velocity perturbations are 1.0-1.2%. Our ScS-S travel time residuals sampling similar regions also show similar results: the middle and the two sides have S velocity variations of about 2.0-3.0% and 1.0-1.5%, respectively.

The heterogeneities may be the remnants of ancient subducted slabs. We propose that the lateral variations are caused by either different slabs, or by different compositional fragments arising from material segregation during subduction.

Although there may be discrepancies between single and multiple scattering calculation in determining the rms perturbation of scatterer, our results suggest that they are generally on the same order. Other effects such as near surface scattering may need further investigation in the future.

Acknowledgments

We would like to thank V.F. Cormier and another reviewer for their constructive comments/suggestions, which have greatly improved the quality of the manuscript. We also thank N.J. Mancinelli and P.M. Shearer for providing us the phonon code generously. X. Ma and X. Sun are supported by China National Science Foundation (No. 41274057) and the 973 project of Ministry of Science and Technology of China (2014CB845901), and China National Science Foundation (No. 41330209). The Antarctic seismic data were collected with the support of the National Science Foundation under grants ANT-0537597, ANT-0632209, and ANT-0838934. Figures are made using Generic Mapping Tools ([Wessel and Smith, 1995](#)). The facilities of IRIS Data Services, and specifically the IRIS Data Management Center, were used for access to waveforms, related metadata, and/or derived products used in this study. IRIS Data Services are funded through the Seismological Facilities for the Advancement of Geoscience and EarthScope (SAGE) Proposal of the National Science Foundation under Cooperative Agreement EAR-1261681.

References

Accardo, N.J., Wiens, D.A., Hernandez, S., Aster, R.C., Nyblade, A., Huerta, A., Anandakrishnan, S., Wilson, T., Heeszel, D.S., Dalziel, I.W.D., 2014. Upper mantle seismic anisotropy beneath the West Antarctic Rift System and surrounding region from shear wave splitting analysis. *Geophys. J. Int.* 198, 414–429.

392 Anthony, R.E., Aster, R.C., Wiens, D.A., Nybalade, A., Anandakrishnan, S., Huerta,
 393 A., Winberry, J.P., Wilson, T., Rowe, C., 2015. The seismic noise environment
 394 of Antarctica. *Seismol. Res. Lett.* 86 (1), 89-100.

395 Bataille, K., Flatté, S.M., 1988. Inhomogeneities near the core-mantle boundary
 396 inferred from short-period scattered PKP waves recorded at the global digital
 397 seismograph network. *J. Geophys. Res.* 93 (B12), 15057-15064.

398 Bhattacharyya, J., Shearer, P., Master G., 1993. Inner core attenuation from short-
 399 period PKP(BC) versus PKP(DF) waveforms. *Geophys. J. Int.* 114, 1-11.

400 Cao, A., Romanowicz, B., 2007. Locating scatterers in the mantle using array analysis
 401 of PKP precursors from an earthquake doublet. *Earth planet. Sci. Lett.* 255, 22–
 402 31.

403 Chernov, L.A., 1960. *Wave Propagation in a Random Medium.* trans. Silverman,
 404 R.A. McGraw-Hill, New York.

405 Christensen, U.R., Hofmann, W., 1994. Segregation of subducted oceanic crust in the
 406 convecting mantle. *J. Geophys. Res.* 99 (B10), 19867-19884.

407 Cleary, J.R., Haddon, R.A.W., 1972. Seismic wave scattering near the core-mantle
 408 boundary: a new interpretation of precursor to PKP. *Nature* 240, 549-551.

409 Cormier, V.F., 1995. Time-domain modelling of PKIKP precursors for constraints on
 410 the heterogeneity in the lowermost mantle. *Geophys. J. Int.* 121, 725-736.

411 Cormier, V.F., Xu, L., 1998. Seismic Attenuation of the inner core: Viscoelastic or
 412 stratigraphic? *Geophys. Res. Lett.* 25, 4019-4022.

413 Cormier, V.F., 1999. Anisotropy of heterogeneity scale lengths in the lower mantle
 414 from PKIKP precursors. *Geophys. J. Int.* 136, 373-384.

415 Doornbos, D.J., 1976. Characteristics of lower mantle inhomogeneities from scattered
 416 waves. *Geophys. J. R. astr. Soc.* 44, 447-470.

417 Doornbos, D.J., 1978. On seismic-wave scattering by a rough core-mantle boundary.
 418 *Geophys. J. R. astr. Soc.* 53, 643-662.

419 Frost, D.F., Rost, S., Selby, N.D., Stuart, G.W., 2013. Detection of a tall ridge at the
 420 core–mantle boundary from scattered PKP energy. *Geophys. J. Int.* 123, 449-470.

421 Garnero, E.J., Vidale, J.E., 1999. ScP: a probe of ultralow velocity zones at the base
 422 of the mantle. *Geophys. Res. Lett.* 26 (3), 377-380.

423 Garnero, E.J., 2000. Heterogeneity of lowermost mantle. *Annu. Rev. Earth Planet.*
 424 *Sci.* 28, 509-537.

425 Garnero, E.J., 2004. A new paradigm for Earth's core-mantle boundary. *Science* 304,
 426 834-836.

427 Garnero, E.J., Maupin V., Lay, T., Fouch, M.J., 2004. Variable azimuthal anisotropy
 428 in Earth's lowermost mantle. *Science* 306, 259–261.

429 Garnero, E.J., McNamara, A.K., 2008. Structure and dynamics of Earth's lower
 430 mantle. *Science* 320, 626-628.

431 Geller, R.J., Takeuchi, N., 1995. A new method for computing highly accurate DSM
 432 synthetic seismograms. *Geophys. J. Int.* 123, 449-470.

433 Grand, S.P., 2002. Mantle shear-wave tomography and the fate of subducted slabs.
 434 *Philos. Trans. R. Soc. London, Ser. A.* 360, 2475-2491.

435 Haddon, R.A., Cleary, R.J., 1974. Evidence for scattering of seismic PKP waves near
 436 the core-mantle boundary. *Phys. Earth Planet. Inter.* 8, 211-234.

437 Hedlin, M.A.H., Shearer, P.M., 2000. An analysis of large-scale variations in small-
 438 scale mantle heterogeneity using Global Seismographic Network recordings of
 439 precursors to PKIKP. *J. Geophys. Res.* 105 (B6), 13665-13673.

440 Hedlin, M.A.H., Shearer, P.M., Earle P.S., 1997. Seismic evidence for small-scale
 441 heterogeneity throughout the Earth's mantle. *Nature* 387, 145-150.

442 Heeszel, D.S., Wiens, D.A., Nyblade, A.A., Hansen, S.E., Kanao, M., An, M., Zhao,
 443 Y., 2013. Rayleigh wave constraints on the structure and tectonic history of the
 444 Gamburtsev Subglacial Mountains, East Antarctica. *J. Geophys. Res. Solid Earth*
 445 118, 2138-2153

446 He, Y., Wen, L., 2009. Structural features and shear-velocity structure of the "Pacific
 447 Anomaly". *J. Geophys. Res.* 114, B02309.

448 He, Y., Wen, L., 2012. Geographic boundary of the "Pacific Anomaly" and its
 449 geometry and transitional structure in the north. *J. Geophys. Res.* 117, B09308.

450 Helffrich, G., Wood, B., 2001. The Earth's mantle. *Nature* 412, 501-507.

451 Husebye, E.S., King, D.W., 1976. Precursors to PKIKP and seismic wave scattering
 452 near the mantle-core boundary. *J. Geophys. Res.* 81 (11), 1870-1882.

453 Kawai, K., Takeuchi, N., Geller, R.J., 2006. Complete synthetic seismograms up to 2
 454 Hz for transversely isotropic spherically symmetric media. *Geophys. J. Int.* 164,
 455 411-424.

456 Kendall, J.M., Silver, P.G., 1996. Constraints from seismic anisotropy on the nature
457 of the lowermost mantle. *Nature* 381, 409-412.

458 Kennett, B.L.N., Engdahl, E.R., and Buland, R., 1995. Constraints on seismic
459 velocities in the Earth from travel times. *Geophys. J. Int.* 122, 108-124.

460 Lay, T., Williams, Q., Garnero, E.J., 1998. The core-mantle boundary layer and deep
461 earth dynamics. *Nature* 392, 461–468.

462 Lay, T., Garnero, E.J., 2011. Deep mantle seismic modeling and imaging. *Annu. Rev.*
463 *Earth Planet. Sci.* 39, 91-123.

464 Lee, C.-T.A., Chen, W., 2007. Possible density segregation of subducted oceanic
465 lithosphere along a weak serpentinite layer and implications for compositional
466 stratification of the Earth's mantle. *Earth Planet. Sci. Lett.* 255, 357-366.

467 Li, X., Romanowicz, B., 1996. Global mantle shear velocity model developed using
468 nonlinear asymptotic coupling theory. *J. Geophys. Res.* 101 (B10), 22245-22272.

469 Li, C., van der Hilst, R.D., Engdahl, E.R., Burdick, S., 2008. A new global model for
470 P wave speed variations in Earth's mantle. *Geochem. Geophys. Geosyst.* 9,
471 Q05018.

472 Long, M.D., 2009. Complex anisotropy in D'' beneath the eastern Pacific from SKS–
473 SKKS splitting discrepancies. *Earth Planet. Sci. Lett.* 283, 181-189.

474 Mancinelli, N.J., Shearer, P.M., 2013. Reconciling discrepancies among estimates of
475 small-scale mantle heterogeneity from PKP precursors. *Geophys. J. Int.* 195,
476 1721-1729.

477 Margerin, L., Nolet, G., 2003a. Multiple scattering of high-frequency seismic waves
 478 in the deep Earth: Modeling and numerical examples. *J. Geophys. Res.* 108 (B5),
 479 2234.

480 Margerin, L., Nolet, G., 2003b. Multiple scattering of high-frequency seismic waves
 481 in the deep Earth: PKP precursors analysis inversion for mantle granularity. *J.*
 482 *Geophys. Res.* 108 (B11), 2514.

483 Maruyama, S., Santosh, M., Zhao, D., 2007. Superplume, supercontinent, and post-
 484 perovskite: mantle dynamics and anti-plate tectonics on the Core-Mantle
 485 Boundary. *Gondwana Res.* 11, 7-37.

486 McNamara, A.K., Zhong, S., 2005. Thermochemical structures beneath Africa and the
 487 Pacific Ocean. *Nature* 437, 1136-1139.

488 Metcalfe, G., Bina, C.R., Ottino, J.M., 1995. Kinematic considerations for mantle
 489 mixing. *Geophys. Res. Lett.* 22 (7), 743-746.

490 Miller, M.S., Niu, F., 2008. Bulldozing the core-mantle boundary: Localized seismic
 491 scatterers beneath the Caribbean Sea. *Phys. Earth Planet. Inter.* 170, 89-94.

492 Ni, S., Helmberger, D.V., 2003. Seismological constraints on the South African
 493 superplume: Could be the oldest distinct structure on earth. *Earth Planet. Sci.*
 494 *Lett.* 206, 119-131.

495 Niu, F., Wen, L., 2001. Strong seismic scatterers near the core-mantle boundary west
 496 of Mexico. *Geophys. Res. Lett.* 28 (18), 3557-3560.

497 Rost, S., Garnero, E.J., Williams, Q., 2006. Fine-scale ultralow-velocity zone
 498 structure from high-frequency seismic array data. *J. Geophys. Res.* 111, B09310.

499 Rost, S., Garnero, E.J., Stefan, W., 2010. Thin and intermittent ultralow-velocity
 500 zones. *J. Geophys. Res.* 115, B06312.

501 Shearer, P.M., Earle, P.S., 2004. The global short-period wavefield modelled with a
 502 Monte Carlo seismic phonon method. *Geophys. J. Int.* 158, 1103-1107.

503 Simmons, N.A., Forte, A.M., Boschi, L., Grand, S.P., 2010. GyPSuM: A joint
 504 tomographic model of mantle density and seismic wave speeds. *J. Geophys. Res.*
 505 115, B12310.

506 Stixrude, L., Lithgow-Bertelloni, C., 2012. Geophysics of chemical heterogeneity in
 507 the mantle. *Annu. Rev. Earth planet. Sci.* 40(1), 569–595.

508 Sun, W., Ding, X., Hu, Y., Zartman, R.E., Arculus, R.J., Kamenetsky, V.S., Chen, M.,
 509 2011. The fate of subducted oceanic crust : a mineral segregation model.
 510 *International Geology Review.* 53(8), 879-893.

511 Thomas, C., Kendall, J.M., Helffrich, G., 2009. Probing two low-velocity regions
 512 with PKP b-caustic amplitudes and scattering. *Geophys. J. Int.* 178, 503-512.

513 Thomas, C., Weber, M., Wicks, C.W., Scherbaum, F., 1999. Small scatterers in the
 514 lower mantle observed at German broadband arrays. *J. Geophys. Res.* 104(B7),
 515 15073-15088.

516 Vanacore, E., Niu, F., Ma, Y., 2010. Large angle reflection from a dipping structure
 517 recorded as a PKIKP precursor: Evidence for a low velocity zone at the core-
 518 mantle boundary beneath the Gulf of Mexico. *Earth Planet. Sci. Lett.* 293, 54-62.

519 Van der Meer, D.G., Torsvik, T.H., Spakman, W., van Hinsbergen, D.J.J., Amaru,
 520 M.L., 2012. Intra-Panthalassa Ocean subduction zones revealed by fossil arcs
 521 and mantle structure. *Nature Geosci.* 5, 215-219.

522 Van Keken, P.E., Hauri, E.H., Ballentine, C.J., 2002. Mantle mixing: the generation,
 523 preservation, and destruction of chemical heterogeneity. *Annu. Rev. Earth*
 524 *Planet. Sci.* 30, 493-525.

525 Vidale, J.E., Hedlin, M.A.H., 1998. Evidence for partial melt at the core-mantle
 526 boundary north of Tonga from the strong scattering of seismic waves. *Nature*
 527 391, 682-685.

528 Wang, Y., Wen, L., 2007. Geometry and P and S velocity structure of the “African
 529 Anomaly”. *J. Geophys. Res.* 112, B05313.

530 Waszek, L., Thomas, C., Deuss, A., 2015. PKP precursor: Implications for global
 531 scatterers. *Geophys. Res. Lett.* 42 (10), 3829-3838.

532 Wen, L., Helmberger, D., 1998. Ultra low velocity zones near the core-mantle
 533 boundary from broadband PKP precursors. *Science* 279, 1701-1703.

534 Wen, L., 2000. Intense seismic scattering near the core-mantle boundary beneath the
 535 Comoros hotspot. *Geophys. Res. Lett.* 27 (22), 3627-3630.

536 Wen, L., 2001. Seismic evidence for a rapidly varying compositional anomaly at the
 537 base of the Earth’s mantle beneath the Indian Ocean. *Earth Planet. Sci. Lett.* 194,
 538 83-95.

539 Wen, L., Niu, F., 2002. Seismic velocity and attenuation structures in the top of the
 540 Earth’s inner core. *J. Geophys. Res.* 107 (B11), 2273.

541 Wessel, P., Smith, W.H.F., 1995. New version of the generic mapping tools released.
 542 EOS. Trans. Am. Geophys. Un. 76, 329.

543 Wyssession, M.E., 1996. Large-scale structure at the core–mantle boundary from
 544 diffracted waves. *Nature* 382, 244 -248.

545 Wyssession, M.E., Bartkó L., Wilson, J.B., 1994. Mapping the lowermost mantle using
 546 core-reflected shear waves. *J. Geophys. Res.* 99 (B7), 13667-13684.

547 Wyssession, M.E., Fischer, K.M., Al-eqabi, G.I., Shore, P.J., Gurari, I., 2001. Using
 548 MOMA broadband array ScS-S data to image smaller-scale structures at the base
 549 of the mantle. *Geophys. Res. Lett.* 28 (5), 867-870.

550 Wu, R.S., Aki, K., 1985. Elastic wave scattering by a random medium and the small-
 551 scale inhomogeneities in the lithosphere. *J. Geophys. Res.* 90 (B12), 10261-
 552 10273.

553 Yao, J., Wen, L., 2014. Seismic structure and ultra-low velocity zones at the base of
 554 the Earth's mantle beneath Southeast Asia. *Phys. Earth Planet. Inter.* 233, 103-
 555 111.

556 Young, C.J., Thorne, L., 1987. The core-mantle boundary. *Ann. Rev. Earth Planet.*
 557 *Sci.* 15, 25-46.

558 Yu W., Wen L., 2006. Seismic velocity and attenuation structures in the top 400 km
 559 of the Earth's inner core along equatorial paths. *J. Geophys. Res.* 111, B07308.

560 Zhao, D., 2004. Global tomographic images of mantle plumes and subducting slabs:
 561 insight into deep Earth dynamics. *Phys. Earth Planet. Inter.* 146, 3-34.

Figure captions

Figure 1. (a) Ray-paths of PKIKP and scattered PKP waves. The PKP precursors are scattered PKP waves generated by the seismic scatterers (black dots) in the lower mantle beneath the source or receiver. The five-pointed star denotes the earthquake at the surface, and the triangle denotes the seismic station. (b) Travel time curves of four branches of PKP. Due to the unusual ray paths caused by scatterers in the lower mantle, the scattered waves can precede PKIKP by up to ~20s. The shaded region indicates the possible earlier arrival times for precursors.

Figure 2. Maps showing the distributions of the earthquakes and stations used in this study. (a) The background is the CMB shear velocity perturbation map from Grand (2002). The stars indicate earthquake locations. The circles with different colors and inverted triangles show PKIKP entry and exit points on the CMB generated by the three groups of earthquakes. Note that the earthquakes are divided into three groups (western, middle, and eastern region from left to right), and the entry points of the three groups of earthquakes at CMB are well separated on the source side while being intermixed on the receiver side. (b) Distributions of seismic stations in Antarctica. The squares and triangles denote GAMSEIS and POLENET/ANET network stations, respectively.

Figure 3. Examples of PKP precursors observed in the Antarctic arrays from earthquakes in different locations (Fig. 2b). Each trace is aligned with the hand-picked PKIKP phase ($t=0$). The hand-picked precursor onsets are marked by squares. The seismograms in (a) and (c) are from two events that occurred in western and eastern region, respectively. The seismograms in (b) are from events that occurred in the middle region. Note that the PKP precursors in (a) and (c) are weak while the precursors from the middle region show clear and comparable amplitudes to that of PKIKP waves.

Figure 4. Maps of scatterer probability and hit count for events in the eastern (a), middle (b), and western (c) region, respectively. Four depths of seismic scattering are presented: 0 km, 200 km, 300 km, and 400 km above the CMB. At each depth, the left figure is the scatterer probability map and the right is the hit count map. The red stars denote event locations. The regions with high scatterer probability and large hit count are considered to be the most likely locations from where the PKP precursor are generated.

Figure 5. Observed (black) and synthetic PKP precursor envelopes (blue) for three representative events in three regions (see Fig. 3). The seismograms are aligned with the hand-picked PKIKP phase ($t=0$). The envelopes of PKIKP (black smoothed curve)

for each event are stacked from individual seismograms, which are calculated using DSM ([Kawai and Geller, 2006](#)). (a,c) Synthetic curves are calculated for 0.5% rms velocity perturbation within 400 km above the CMB in the scattering regions shown in Fig. 4a,c. (b) Synthetic curves are the same as in (a,c) except for ~1.2% velocity perturbation in the region in Fig. 4b. The scale length of the scatterers is 8 km in our model.

Figure 6. Map of the study region with surface projection of ScS data. Yellow and cyan circles indicate ScS bouncing points at CMB from broadband and long-period data, respectively. The gray lines denote the great-circle path connecting the earthquakes (red stars) and seismic stations (blue squares). The red dashed line is the boundary of the Pacific Anomaly from [He and Wen \(2012\)](#). Note that the ScS paths bypass the Pacific Anomaly. The inset shows the ray paths of S and ScS at epicentral distances from 40° to 80°. Pink and blue lines denote the S and ScS ray-paths respectively.

Figure 7. Correlations between observed ScS-S residuals with S (a) and ScS (b) residuals after correcting travel time anomalies 400 km above the CMB using model GyPSuM10 ([Simmons et al., 2010](#)). Dashed lines are one to one correlation.

628

629 **Figure 8.** Averaged shear wave velocity perturbations at the bottom 400 km of the
630 lower mantle. The white crosses, which denote fast velocities, are plotted at ScS
631 reflection points at the CMB. The background colors are smoothed S velocity
632 perturbations based on our observations. Three regions (east, middle and west) with
633 different heterogeneity variations are delineated approximately by the three dashed
634 ellipses.

Figure-1a

Figure-1b

Figure-2

Figure-3a

Figure-3b

Figure-3c

Figure-4a

2491 km

2591 km

2691 km

2891 km

Figure-4b

2491 km

2591 km

2691 km

2891 km

Figure-4c

Figure-5a

Figure-5b

Figure-5c

Figure-6

Figure-7

Figure-8

Table 1

Events used in this study for PKP precursor analysis

Date	Time (GMT)	Latitude (deg)	Longitude (deg)	Depth (km)	Magnitude (mb)	Region
2009.01.26	19:11:47	51.9550	-171.1590	25.3000	5.6	East
2010.07.18	05:56:44	52.8760	-169.8480	14.0000	6.3	
2011.06.24	03:09:39	52.0500	-171.8360	52.0000	6.9	
2011.09.02	10:55:53	52.1710	-171.7080	32.0000	6.5	
2009.12.10	02:30:52	53.4170	152.7560	656.2000	6.1	West
2010.07.30	03:56:13	52.4980	159.8430	23.0000	6.1	
2009.03.30	12:07:28	51.5350	-178.2580	31.0000	6.0	Middle
2009.06.22	19:55:24	51.2770	-178.2010	35.0000	5.6	
2009.07.06	14:53:12	50.4350	176.9920	22.0000	6.0	
2009.10.07	05:38:37	52.1570	178.0520	138.4000	5.6	
2009.12.17	20:01:21	51.3980	179.9630	35.0000	6.2	