Daytime cold exposure and salt intake
based on nocturnal urinary sodium excretion:
A cross-sectional analysis of The HEIJO-KYO study
Short title: Cold exposure and salt intake

Keigo SAEKI M.D., Ph.D.¹, Kenji OBAYASHI M.D., Ph.D.¹,
Nobuhiro TONE², and Norio KURUMATANI M.D., Ph.D.¹

¹Department of Community Health and Epidemiology, Nara Medical University School of Medicine, Nara, Japan

²Center for Academic Industrial and Governmental Relations, Nara Medical University School of Medicine, Nara, Japan

Corresponding Author

Keigo Saeki, MD

840 Shijocho, Kashiharashi, Nara 634-8521, Japan

Department of Community Health and Epidemiology, Nara Medical University School of Medicine, Nara, Japan

E mail: saekik@naramed-u.ac.jp

Phone: +81-744-29-8841, Fax: +81-744-29-0673

Abstract

Increased cardiovascular incidence in winter is partly explained by higher blood pressure due to cold exposure. Although higher salt intake induced by cold exposure has been reported in mice, the association remains unclear in humans. To investigate the association between salt intake and cold exposure in winter, a cross-sectional study was conducted among 860 elderly subjects (mean $\pm$ standard deviation: 72.0 ± 7.1 years).

We determined ambient temperature at every 10 min according to indoor temperature measured in the subjects' home, outdoor temperature, and self-administered diary logging time spent outdoors. Salt intake was estimated by nocturnal sodium excretion rate of overnight urine collection. A 1 °C lower daytime ambient temperature was significantly associated with a higher urinary sodium excretion rate by 0.07 mmol/h in the subsequent night independent of age, sex, body weight, alcohol intake, calcium channel blocker use, diabetes, household income, estimated glomerular filtration rate, daytime physical activity ($p = 0.02$). After further adjustment for outdoor temperature and day length, the lowest tertile groups of ambient daytime temperature ($10.1 \pm 2.3^{\circ}\text{C}$) showed the nocturnal urinary sodium excretion rate was higher by 14.2% (7.62 vs. 6.54 mmol/h) compared with the highest tertile group ($19.3 \pm 1.8^{\circ}\text{C}$). Higher sodium excretion rate was associated with higher nighttime ambulatory blood pressure ($p <$

0.01) and its lower nocturnal dipping ($p < 0.01$). Significant association between higher salt intake and daytime cold exposure partly explain the mechanism of higher blood pressure in winter, and suggest that a reduction of cold exposure might be effective to decrease salt intake.

Key Words: Cold exposure, Indoor temperature, Outdoor temperature

Nocturnal urinary sodium excretion rate, Ambulatory blood pressure monitoring

Introduction

Higher mortality in winter than other seasons is worldwide phenomenon [1-4].

Increased mortality from ischemic heart diseases and stroke in winter [5-7] is partly explained by a higher blood pressure (BP) due to cold exposure. [8, 9]

As an index of cold exposure, outdoor temperature is convenient because it is usually available from meteorological office. However, it may diverse from the really-exposed temperature, because it poorly correlate with indoor temperature especially in cold climate [10]. Ambient temperatures defined as indoor temperature while the participants stayed at home and outdoor temperature while the participants were out of their home showed the strongest association with ambulatory BP than outdoor temperature and indoor temperature. [10, 11]

Higher salt intake estimated using 24 h urinary sodium excretion is associated with higher BP from within-population and cross-population analyses in the INTERSALT study [12, 13]. Systematic review from randomized controlled studies demonstrated effectiveness of a low-salt diet in decreasing BP.[14] Meta-analysis of prospective cohort studies including over 100000 participants showed that higher salt intake is significantly associated with greater incidence of stroke and total cardiovascular disease (CVD).[15]

Higher salt intake induced by cold exposure was reported in mice, [16, 17] and it may also explain the higher BP due to cold exposure. However, the influence of cold exposure on salt intake in human has not reported. The association between ambient temperature and salt intake may provide important evidence to develop a suitable environment for reduction of salt intake to prevent CVD.

To quantify the association between ambient temperature and salt intake, we conducted the cross-sectional study among 880 elderly participants. We estimated salt intake from the nocturnal urinary sodium excretion because of the high correlation between 24 h and overnight sodium excretion in an Asian population. [18], [19]

Methods

Participants

From September to April in 2010, 2011, 2012, and 2013, we recruited 880 home-dwelling males and females, aged 60 years or more for HEIJO-KYO (Housing Environments and Health Investigation among Japanese Older People in Nara, Kansai Region), a prospective community-based cohort study. The study protocol has been previously described. [20] Of all 880 participants, we excluded 20 participants with missing of indoor temperature or overnight urine collection during the colder season

(October–April), 860 were remained for analysis (Fig.1). All participants provided a written informed consent and the study protocol was approved by the Nara Medical University Ethics Review Board.

Study protocol

The participants' characteristics, including age, gender, smoking, household income, and drinking habits, and medication use were obtained using a standardized questionnaire. After the waist circumference at the level of the umbilicus in the standing position, an overnight fasting venous sample was obtained. At noon, we started measurements of indoor temperature, ambulatory BP and actigraphy for 48 h. The participants were instructed to collect overnight urine at the second night. We revisited each participant at home to retrieve instruments and collect their diaries logging bedtime, rising time, duration in bedroom, and time spent outdoors.

Nocturnal urinary sodium excretion

After discarding the last void at bedtime, participants were asked to collect urine until the first morning void. The sodium concentration was measured using an ion-selective electrode method in a commercial laboratory (SRL Co. Inc., Tokyo, Japan). The total

nocturnal urinary sodium excretion (mmol) and nocturnal urinary sodium excretion rate (mmol/h) were calculated from the total urine volume, the sodium concentration, and the duration of urine collection.

Daytime ambient temperature

Indoor temperature was measured in the living room and bedroom 60 cm above the floor. Bed temperature was measure at center of the bed 50cm from the headboard. These temperatures were measured at 10 min intervals using a Thermochron iButtons (DS1922L; Maxim Integrated, Dallas, TX, USA) with an accuracy of $\pm 0.5^{\circ}\text{C}$ from -10°C to $+65^{\circ}\text{C}$ and a 0.0624°C resolution. Outdoor temperature was also measured at 10 min intervals and was provided by the local meteorological office in Nara (latitude, 34° N). We defined the ambient temperature as indoor temperature (temperature in the living room or bedroom) while the participants were at living room or bedroom, and the outdoor temperature while the participants were out of their homes. The at-home and out-of-home periods were determined according to the participants' self-reported diaries logging the time spent outdoors and the time spent at living room and bedroom. The mean ambient temperature during the last daytime (rising–bedtime) before the nocturnal urine collection was calculated.

Ambulatory BP and physical activity

Physical activity was determined at 1 min epochs using an actigraph (Actiwatch 2; Respironics Inc., Murrysville, PA, USA) worn on the non-dominant arm. Ambulatory BP was measured using a validated device (TM-2430; A&D Co. Ltd., Tokyo, Japan) on the non-dominant arm at 30 min intervals. We calculated the mean of two days for the daytime systolic BP (blood pressure), nighttime systolic BP, and dipping [(daytime systolic BP – nighttime systolic BP)/ (daytime systolic BP) × 100].

Other measurements

Venous blood samples were analyzed using a standard clinical chemical analysis to determine the concentration of creatinine, glycated hemoglobin (HbA1c), and fasting plasma glucose. Diabetes was defined based on medical history, the current administration of anti-diabetic treatment, or fasting plasma glucose of at least 126 mg/dl and HbA1c of at least 6.5% (National Glycohemoglobin Standardization Program value). The estimated glomerular filtration rate (eGFR) was calculated by the Japanese Society of Nephrology - Chronic Kidney Disease Practice Guide formula: $eGFR \text{ (ml/min/1.73 m}^2\text{)} = 194 \times [\text{serum creatinine (mg/dl)}]^{-1.094} \times [\text{age (years)}]^{-0.287}$. The

result was multiplied by a correction factor of 0.739 for women. Data of the day length from sun rise to sunset in Nara on the measurement days were obtained from National astronomical Observatory of Japan. The habitual intake of soup and fruit were asked using self-administered questionnaire as follows: “How many cups of Miso soup and other kind of soup do you have in a week?”, “How often do you take fruit in a week?”, and “Please select your quantity of your usual fruit intake from 1) small size, 2) standard portion size, and 3) large portion size? The product of the fruit intake frequency in a week and the size (1 to 3) was calculated.

Statistical analysis

For continuous variables with a normal distribution, the mean $\pm$ standard deviation was reported. For variables not distributed normally, the median and interquartile ranges were reported. The associations of variables with the nocturnal urinary sodium excretion rate were assessed using univariate and multivariate linear regression models. Multivariate adjusted mean values among tertile groups of daytime ambient temperature were compared using ANCOVA (analysis of covariance).

We assessed associations of nocturnal urinary sodium excretion rate with daytime

ambient temperature as continuous variables (Table 2) and tertile groups (Table3).

We explored the variables as potential confounders with marginal to significant association ($p < 0.20$) in the univariate model. In the multivariate models, we simultaneously adjusted for potential confounders. To detect severe multicollinearity we calculated the variance inflation factors. All p -values were two-sided, and those <0.05 were considered statistically significant. All statistical analyses were performed using the SPSS ver. 21.0 software (SPSS, Chicago, IL, USA).

Results

The mean age among 860 participants was 72.0 ± 7.1 (mean $\pm$ SD) and 49.2% were males. Tertile groups of ambient temperature were T1 ($10.1 \pm 2.3^{\circ}\text{C}$), T2 ($15.0 \pm 1.1^{\circ}\text{C}$), and T3 ($19.3 \pm 1.8^{\circ}\text{C}$), respectively. Daytime ambient temperature was significantly associated with nighttime ambient temperature, bed temperature at night, daytime outdoor temperature, and nighttime outdoor temperature. We did not find a significant difference in household income and education status across the tertile groups (Table 1).

In the univariate linear regression analysis, the daytime ambient temperature was significantly associated with the nocturnal urinary sodium excretion rate ($\beta = -0.07$,

95% CI: -0.13 to -0.01 , $p = 0.03$). On the other hand, nocturnal urinary sodium excretion rate did not show significant association with daytime outdoor temperature ($p = 0.44$), nighttime outdoor temperature ($p = 0.36$), nighttime ambient temperature ($p = 0.30$), and bed temperature in nighttime ($p = 0.30$).

In the multivariate linear regression model, the ambient temperature showed a significant inverse association with the nocturnal urinary sodium excretion rate even after adjusting for potential confounders such as age, male, body weight, alcohol intake, CCB (calcium channel blocker), and diuretics use, household income, and daytime physical activity. ($\beta = -0.07$, 95% CI: -0.13 to -0.01 , $p = 0.02$; Table 2). In this multivariate model, we did not detect severe multicollinearity with variance inflation factors of 10 and above.

Similarly, categorical analysis using tertile groups of daytime ambient temperature showed significant and inverse association with nocturnal urinary sodium excretion rate [T1: 7.49 mmol/h, (95% CI: 7.06 – 7.92); T2: 7.01 (6.58 – 7.44); and T3: 6.65 (6.22 – 7.08); p for trend < 0.01], and total nocturnal urinary sodium excretion [T1: 62.4 mmol (58.7 – 66.0); T2: 58.5 (54.9 – 62.2); and T3: 54.3 (50.7 – 58.0); p for trend < 0.01 ; Figure 2).

In Table 3, a significant association between tertile groups of daytime ambient

temperature and nocturnal urinary sodium excretion rate stayed significant even after adjustment for all potential confounders in univariate models ($p < 0.2$) plus day length (model 2, $p = 0.006$). After further adjustment for outdoor temperature, a lower daytime ambient temperature by 9.2°C ; (T1: 10.1°C vs. T3: 19.3°C) was associated with a higher sodium excretion rate by 14.2 % (7.62 vs. 6.54 mmol/h; model 3). Adjustment for pressure natriuresis induced by nighttime systolic BP during urine collection did not attenuate the significant association ($p = 0.005$, model 4).

In stratified analysis by age groups, gender, and body weight groups, we found consistently inverse association between daytime ambient temperature and nocturnal sodium excretion rate. The higher tertile group of daytime ambient temperature is significantly associated with lower sodium excretion rate ($p = 0.048$) in younger group (≤ 72 years) and elder group (>72 years; $p = 0.042$). As for gender, the lower ambient temperature group ($\leq$ median: 15.07°C) was significantly associated with higher sodium excretion ($p = 0.025$) among female, and the lower tertile group of daytime ambient temperature was significantly associated with higher sodium excretion rate among male ($p = 0.019$). As for obesity, the lower ambient temperature group ($\leq 15.07^{\circ}\text{C}$) was significantly associated with higher sodium excretion ($p = 0.028$) among the low weight group ($\leq$ median: 57kg), and the lower tertile group of daytime ambient temperature

was associated with higher sodium excretion rate ($p = 0.019$) among the higher weight group (> 57 kg). All these associations were independent of other potential confounders such as independent of alcohol intake, CCB administration, diabetes, eGFR, income, daytime physical activity, outdoor temperature, and day length.

Of 860, 789 participants (91.7%) answered the questions about soup and fruit intake. The mean soup intake was 6.0 ± 3.6 (SD) cups in a week, and the fruit intake index was 11.45 ± 7.7 with almost normal distribution. After further adjustment for hot soup intake and fruit intake in the multivariate model in the table 2, we found consistently significant association between daytime ambient temperature and nocturnal sodium excretion rate ($p = 0.039$).

Figure 3 shows mean values of daytime systolic BP, nighttime systolic BP, and dipping by the tertile groups of the nocturnal urinary sodium excretion rate (T1: 3.6 ± 0.1 , T2: 6.4 ± 0.1 , and T3: 11.2 ± 0.1 mmol/h) adjusted for age, gender, abdominal circumference, current smoking status, alcohol intake, anti-hypertensive medication, and daytime physical activity. Nocturnal urinary sodium excretion rate was significantly associated with nighttime systolic BP (p for trend < 0.01) and dipping (p for trend < 0.01). The nighttime systolic BP in the highest tertile group of the nocturnal sodium excretion rate was significantly higher than in the lowest tertile group (119.2 vs.

112.5mmHg, $p < 0.001$). The dipping in the lowest tertile group of sodium excretion rate was significantly higher than that in the highest tertile group (16.2% vs. 11.7%, $p < 0.01$). In contrast, the nocturnal urinary sodium excretion rate did not show a significant association with daytime systolic BP (p for trend = 0.64). We did not find significant association between daytime ambient temperature and nighttime systolic BP ($p = 0.22$).

Discussion

A higher salt intake is shown to be associated with a lower ambient temperature as evidenced by the higher nocturnal urinary sodium excretion rates observed. These higher nocturnal urinary sodium excretion rates were significantly associated with the lower ambient temperature in the previous day independent of potential confounders such as age, gender, body weight, alcohol intake, CCB administration, diabetes, household income, eGFR, and daytime physical activity ($\beta = -0.07$, 95% CI: -0.13 to -0.01, $p = 0.02$).

Although Hata et al. showed higher sodium excretion rates in winter (monthly outdoor temperature: 4.8°C) than in summer (29.1°C) among 14 normotensive and 20 hypertensive men,[21] the association between temperatures which each participants exposed and urinary sodium excretion have not yet been investigated. The findings from

the present study are concordant with the experiments using rats [22] and mice. [17] Dejima et al. showed that cold exposure (7°C–9°C) for 6 h caused a higher salt intake during the 9 h following cold exposure [16], and the increased salt intake was accompanied by a higher concentration of noradrenaline in the kidney and plasma renin activity during cold exposure compared with the control group. [17] The mechanism of cold-induced salt intake is explained by the activation of the renin angiotensin system (RAS) by noradrenaline [23] and the increase of salt intake by RAS. [24] In the present study, cold exposure at during daytime cause higher sympathetic activity and RAS activation, and may cause higher salt appetite during daytime based on previous studies. In contrast, influence of cold exposure derived from indoor temperature attenuates during bed [25]. The higher salt intake during daytime may cause higher nocturnal sodium excretion at night.

Pressure natriuresis may increase nocturnal urinary excretion. [26-28] In the present study, we found significant association between ambient temperature and nocturnal urinary sodium excretion rate even after adjusting for ambulatory BP during urine collection (model 4 in Table 3).

The strengths of the current study include the measurement of ambulatory temperature and the large-scale sample size. The ambient temperature in the present analysis was

estimated from both outdoor and indoor temperatures because it allowed for a better prediction of morning BP surge and daytime ambulatory BP than crude indoor and outdoor temperatures. [10, 11] Owing to the large-scale sample size, we could assess the independent association of ambient temperature and salt intake independent of a variety of potential confounders.

However, the present study has some limitations. First, we estimated salt intake through nocturnal urine collection, which is inferior to 24 h urine collection, because it provides a higher ratio of intra- to inter-individual variance than the 24-hour urine collection (2.12 vs. 1.48). [18] However, the association between the daytime ambient temperature and the subsequent nocturnal urinary sodium excretion may be acceptable. In addition, sodium excretion in the 24 h urine collection and overnight urine collection showed a high correlation among the Asian population (correlation coefficient: 0.86–0.92). [18, 19] The second limitation of the present study is the lack of information regarding sodium loss from sweat. To address this, we assessed the physical activity measured using the actigraph. Third limitation lies in the way to estimate ambient temperature. We use outdoor temperature as exposed temperature while the participants go out of home. Therefore, we may have overestimated cold exposure when participants left their homes but remained in an indoor environment. Fourth, we lack information about the intake of

total energy and other nutrition in the present study. Finally we recruited participants with cooperation of local-resident associations and elderly-resident clubs in six areas of Nara prefecture with non-random sampling (participation rate unclear), and the generalizability of the study may be limited. However, the proportion of study participants using anti-hypertensive and anti-hyperglycemic medications was similar to that of a large-scale national survey among 3499 elderly (≥ 60 years) individuals conducted in 2010 (anti-hypertensive medication use: 40.3%; anti-diabetic treatment: 12.0%). [29]

The clinical implications of the present study can be interpreted through previous evidence regarding the influence of salt intake on BP and incidence of CVD. High salt intake can cause cardiovascular diseases (CVDs) via hypertension. Concordant with a previous study[28], we found a significant association between sodium excretion and nighttime BP. Nighttime BP has been shown to provide a superior prediction for CVD incidence compared with daytime systolic BP. [30] Salt-induced hypertension is caused by a higher extracellular volume, [31] the cellular hypertrophy of arterial smooth muscle, cardiac myocytes,[32] and endothelial dysfunction due to decreased endothelial nitric oxide.[33, 34] Furthermore, epidemiologic studies have shown a direct harmful effect of salt intake on stroke, left ventricular hypertrophy,[35, 36] the progression of

renal disease, and proteinuria [37, 38] independent of BP. The present study suggests that these harmful effects of salt intake may be associated with ambient temperature. A large scale prospective study showed that a 100 mmol increase in the 24 h sodium excretion is associated with a higher hazard ratio for the mortality of coronary heart disease, CVD, and all-cause mortality by 51%, 45%, and 26%, respectively. [39] We found that the nocturnal urinary sodium excretion rate was 1.08mmol/h lower when the ambient temperature was 9.3°C higher (model 3 in Table 3,). Based on the night-to-day ratio of sodium excretion (0.79) [28] and the mean duration of urine collection in the present study (498 ± 82.1 min), 1.08 mmol/h is equivalent to approximately 30.4 mmol sodium excretion in 24 h. This meant that cold exposure by 9.3 °C is associated with a 15.5%, 13.7%, and 7.9% higher mortalities by coronary heart disease, CVD, and all-cause mortality, respectively.

In summary, we found significant cross-sectional association between lower ambient temperature and higher salt intake independent of potential confounders from community-based 860 elderly. Although our findings suggest that a reduction of cold exposure may be effective to decrease salt intake, further well-designed prospective randomized controlled studies are required.

Author contributions

K.S and K.O designed the study and conducted data collection. N.T contributed to methods of measurement. N.K contributed to interpretation of data. K.S analyzed the data and prepared draft of this study and all authors reviewed it.

Acknowledgement

This work was supported by grants from the Department of Indoor Environmental Medicine, Nara Medical University; JSPS KAKENHI Grant Number 2586447, 15H04776, 15H04777; Mitsui Sumitomo Insurance Welfare Foundation; Meiji Yasuda, Life Foundation of Health and Welfare; Osaka Gas Group Welfare Foundation; Japan Diabetes Foundation; and Japan Science and Technology Agency. We are indebted to all participants of this study. We would also like to thank Sachiko Uemura, Naomi Takenaka, Keiko Nakajima, and Syoichi Takamiya for their valuable support during the data collection.

We would like to thank Sachiko Uemura, Keiko Nakajima, and Naomi Takenaka for their valuable support during the data collection.

REFERENCES

- [1] Healy, J. D. Excess winter mortality in Europe: a cross country analysis identifying key risk factors. *J Epidemiol Community Health*. 2003,57:784-9.
- [2] Curriero, F., Heiner, K., Samet, J., Zeger, S., Strug, L., Patz, J. Temperature and mortality in 11 cities of the eastern United States. *Am J Epidemiol*. 2002,155:80-7.
- [3] Ou, C. Q., Song, Y. F., Yang, J., Chau, P. Y., Yang, L., Chen, P. Y., et al. Excess winter mortality and cold temperatures in a subtropical city, Guangzhou, China. *PLoS One*. 2013,8:e77150.
- [4] Xu, B., Liu, H., Su, N., Kong, G., Bao, X., Li, J., et al. Association between winter season and risk of death from cardiovascular diseases: a study in more than half a million inpatients in Beijing, China. *BMC Cardiovasc. Disord*. 2013,13:93.
- [5] Turin, T. C., Kita, Y., Murakami, Y., Rumana, N., Sugihara, H., Morita, Y., et al. Higher stroke incidence in the spring season regardless of conventional risk factors: Takashima Stroke Registry, Japan, 1988-2001. *Stroke*. 2008,39:745-52.
- [6] Arntz, H. R., Willich, S. N., Schreiber, C., Brüggemann, T., Stern, R., Schultheiß, H. P. Diurnal, weekly and seasonal variation of sudden death. Population-based analysis of 24061 consecutive cases. *Eur Heart J*. 2000,21:315-20.
- [7] Wolf, K., Schneider, A., Breitner, S., von Klot, S., Meisinger, C., Cyrys, J., et al. Air temperature and the occurrence of myocardial infarction in Augsburg, Germany. *Circulation*. 2009,120:735-42.
- [8] Modesti, P. A., Morabito, M., Bertolozzi, I., Massetti, L., Panci, G., Lumachi, C., et al. Weather-related changes in 24-hour blood pressure profile: effects of age and implications for hypertension management. *Hypertension*. 2006,47:155-61.
- [9] Modesti, P. A., Morabito, M., Massetti, L., Rapi, S., Orlandini, S., Mancia, G., et al. Seasonal blood pressure changes: an independent relationship with temperature and daylight hours. *Hypertension*. 2013,61:908-14.
- [10] Saeki, K., Obayashi, K., Iwamoto, J., Tone, N., Okamoto, N., Tomioka, K., et al. Stronger association of indoor temperature than outdoor temperature with blood pressure in colder months. *J Hypertens*. 2014,32:1582-9.
- [11] Saeki, K., Obayashi, K., Iwamoto, J., Tone, N., Okamoto, N., Tomioka, K., et al. The relationship between indoor, outdoor and ambient temperatures and morning BP surges from inter-seasonally repeated measurements. *J. Hum. Hypertens*. 2014,28:482-8.
- [12] Elliott, P., Stamler, J., Nichols, R., Dyer, A. R., Stamler, R., Kesteloot, H., et al. Intersalt revisited: further analyses of 24 hour sodium excretion and blood pressure within and across populations. Intersalt Cooperative Research Group. *BMJ*. 1996,312:1249-53.

- [13] Stamler, J. The INTERSALT Study: background, methods, findings, and implications. *Am J Clin Nutr.* 1997,65:626s-42s.
- [14] Graudal, N., Hubeck-Graudal, T., Jurgens, G. Effects of low sodium diet versus high sodium diet on blood pressure, renin, aldosterone, catecholamines, cholesterol, and triglyceride. *Cochrane Database Syst Rev.* 2011,9:CD004022.
- [15] Strazzullo, P., D'Elia, L., Kandala, N. B., Cappuccio, F. P. Salt intake, stroke, and cardiovascular disease: meta-analysis of prospective studies. *BMJ.* 2009,339:b4567.
- [16] Dejima, Y., Kim, S., Kashiwazaki, H., Suzuki, T. A spontaneous increase of salt intake and change of colonic temperature in mice exposed to cold. *Appetite.* 1991,16:169-91.
- [17] Dejima, Y., Fukuda, S., Ichijoh, Y., Takasaka, K., Ohtsuka, R. Cold-induced salt intake in mice and catecholamine, renin and thermogenesis mechanism. *Appetite.* 1996,26:203-19.
- [18] Lisheng, L., Deyu, Z., Lue, J., Liao, Y., Liu, K., Stamler, J. Variability of Urinary Sodium and Potassium Excretion in North Chinese Men. *Journal of Hypertension.* 1987,5:331-5.
- [19] He, J., Klag, M. J., Whelton, P. K., Chen, J. Y., Mo, J. P., Qian, M. C., et al. Agreement between overnight and 24-hour urinary cation excretions in southern Chinese men. *Am J Epidemiol.* 1993,137:1212-20.
- [20] Obayashi, K., Saeki, K., Iwamoto, J., Okamoto, N., Tomioka, K., Nezu, S., et al. Positive effect of daylight exposure on nocturnal urinary melatonin excretion in the elderly: a cross-sectional analysis of the HEIJO-KYO study. *J Clin Endocrinol Metab.* 2012,97:4166-73.
- [21] Hata, T., Ogihara, T., Maruyama, A., Mikami, H., Nakamaru, M., Naka, T., et al. The seasonal variation of blood pressure in patients with essential hypertension. *Clin. Exp. Hypertens. A.* 1982,4:341-54.
- [22] Yoshida, T., Okuno, T., Kawabata, T., Morimoto, T. Salt consumption and body fluid balance during cold exposure in rats. *Physiol Behav.* 1994,55:163-7.
- [23] Johnson, J. A., Davis, J. O., Witty, R. T. Effects of catecholamines and renal nerve stimulation on renin release in the nonfiltering kidney. *Circ. Res.* 1971,29:646-53.
- [24] Fregly, M. J., Rowland, N. E. Role of renin-angiotensin-aldosterone system in NaCl appetite of rats. *Am. J. Physiol.* 1985,248:R1-11.
- [25] Saeki, K., Obayashi, K., Iwamoto, J., Tanaka, Y., Tanaka, N., Kubo, H., et al.. Influence of room heating on ambulatory blood pressure in winter: a randomised controlled study *J Epidemiol Community Health.* 2013,67:484-90.
- [26] Granger, J. P. Pressure natriuresis. Role of renal interstitial hydrostatic pressure. *Hypertension.* 1992,19:I9-17.
- [27] Granger, J. P., Alexander, B. T., Llinas, M. Mechanisms of pressure natriuresis. *Curr.*

- Hypertens. Rep. 2002,4:152-9.
- [28] Staessen, J. A., Birkenhager, W., Bulpitt, C. J., Fagard, R., Fletcher, A. E., Lijnen, P., et al. The relationship between blood pressure and sodium and potassium excretion during the day and at night. *J Hypertens.* 1993,11:443-7.
- [29] The National Health and Nutrition Survey Japan 2010; available at: <http://www.mhlw.go.jp/bunya/kenkou/eiyou/h22-houkoku.html> (Japanese; accessed on May 1, 2015).
- [30] Boggia, J., Li, Y., Thijs, L., Hansen, T. W., Kikuya, M., Bjorklund-Bodegard, K., et al. Prognostic accuracy of day versus night ambulatory blood pressure: a cohort study. *Lancet.* 2007,370:1219-29.
- [31] Guyton, A. C. Blood pressure control--special role of the kidneys and body fluids. *Science.* 1991,252:1813-6.
- [32] Gu, J. W., Anand, V., Shek, E. W., Moore, M. C., Brady, A. L., Kelly, W. C., et al. Sodium induces hypertrophy of cultured myocardial myoblasts and vascular smooth muscle cells. *Hypertension.* 1998,31:1083-7.
- [33] Oberleithner, H., Riethmuller, C., Schillers, H., MacGregor, G. A., de Wardener, H. E., Hausberg, M. Plasma sodium stiffens vascular endothelium and reduces nitric oxide release. *Proc. Natl. Acad. Sci. U. S. A.* 2007,104:16281-6.
- [34] Dickinson, K. M., Keogh, J. B., Clifton, P. M. Effects of a low-salt diet on flow-mediated dilatation in humans. *Am J Clin Nutr.* 2009,89:485-90.
- [35] Kupari, M., Koskinen, P., Virolainen, J. Correlates of left ventricular mass in a population sample aged 36 to 37 years. Focus on lifestyle and salt intake. *Circulation.* 1994,89:1041-50.
- [36] Schmieder, R. E., Messerli, F. H., Garavaglia, G. E., Nunez, B. D. Dietary salt intake. A determinant of cardiac involvement in essential hypertension. *Circulation.* 1988,78:951-6.
- [37] Verhave, J. C., Hillege, H. L., Burgerhof, J. G., Janssen, W. M., Gansevoort, R. T., Navis, G. J., et al. Sodium intake affects urinary albumin excretion especially in overweight subjects. *J Intern Med.* 2004,256:324-30.
- [38] Cianciaruso, B., Bellizzi, V., Minutolo, R., Tavera, A., Capuano, A., Conte, G., et al. Salt intake and renal outcome in patients with progressive renal disease. *Miner. Electrolyte Metab.* 1998,24:296-301.
- [39] Tuomilehto, J., Jousilahti, P., Rastenyte, D., Moltchanov, V., Tanskanen, A., Pietinen, P., et al. Urinary sodium excretion and cardiovascular mortality in Finland: a prospective study. *Lancet.* 2001,357:848-51.

Table1 Characteristic of 860 participants by tertile groups of daytime ambient temperature

Variables	Daytime ambient temperature† (°C)			<i>p</i> for trend
	T1 (<13.1) n=288	T2 (13.1-16.9) n=287	T3 (≥16.9) n=285	
Environment				
Ambient temperature†, mean (SD)				
Daytime	10.1 (2.3)	15.0 (1.1)	19.3 (1.8)	
Nighttime	9.0 (3.4)	12.5 (3.1)	16.4 (3.8)	<0.01
Bed temperature at night, mean (SD)	29.4 (5.2)	29.9 (4.4)	30.4 (3.8)	0.02
Outdoor temperature, °C, mean (SD)				
Daytime	5.4 (2.7)	8.7 (3.9)	13.1 (4.9)	<0.01
Nighttime	1.9 (3.1)	4.8 (3.8)	8.9 (4.9)	<0.01
Day length, min, mean (SD)	656.6 (39.6)	666.2 (46.8)	677.8 (50.6)	<0.01
Basic characteristics				
Age, mean (SD)	71.9 (7.4)	71.7 (6.8)	72.7 (7.1)	0.17
Male, n (%)	147 (51.0)	141 (49.1)	132 (46.3)	0.24
Body weight, kg, mean (SD)	57.2 (10.8)	57.6 (10.7)	57.7 (10.2)	0.59
Current Smoker, n (%)	20 (6.9)	13 (4.5)	11 (3.9)	0.06
Alcohol intake (≥30g/day)	46 (16.0)	37 (12.9)	33 (1.6)	0.07
Clinical characteristics				
Anti-hypertensive use, n (%)	117 (40.6)	136 (47.4)	125 (43.9)	0.45
ACEI/ARBs	74 (25.7)	90 (31.4)	81 (28.4)	0.48
CCB	75 (26.0)	94 (32.8)	85 (29.8)	0.31
Diuretics§	20 (6.9)	16 (5.6)	18 (6.3)	0.74
Diabetes, n (%)	32 (11.1)	34 (11.8)	36 (12.6)	0.52
eGFR, ml/min/1.73m ² , mean (SD)	73.1 (15.3)	73.4 (14.9)	70.1 (14.4)	0.02
Socio-economic status				
Household Income ‡, n (%)	114 (39.6)	115 (40.1)	116 (40.7)	0.81
Education (≥13 years), n (%)	70 (24.3)	74 (25.8)	68 (23.9)	1.00
Other covariates				
Daytime Physical activity, counts/min, mean (SD)	319.9 (117.8)	312.8 (106.8)	284.2 (110.3)	<0.01
Duration spent outdoor in daytime, min, median (IOR)	179(70,346)	182 (76,354)	201 (81,335)	0.57¶

T1–T3, Tertiles of daytime ambient temperature; SD, standard deviation; IQR, inter quartile range;; ACE, angiotensin converting enzyme inhibitor; ARBs, angiotensin receptor blockers; CCB, calcium channel blocker; eGFR, estimated glomerular filtration rate.

† Ambient temperature is indoor temperature during at-home and outdoor temperature during out-of home period.

‡ ≥4 million yen/year

¶ *p*-value was estimated after log-transformation

§ Diuretics include thiazide, loop diuretics, and spironolactone.

Table 2 Association of variables with nocturnal urinary sodium excretion rate

Variables	Univariate model		Multivariate model [‡]		Partial R ²
	$\beta^{\dagger}$	(95%CI)	Adjusted β	(95%CI)	p -value
Environment					
Ambient temperature, (per 1°C)					
Daytime	-0.07	(-0.13, -0.01)	-0.07	(-0.13, -0.01)	0.02
Nighttime	-0.03	(-0.08, 0.03)			0.30
Bed temperature at night, (per 1°C)	0.03	(-0.03, 0.08)			0.30
Outdoor temperature, (per 1°C)					
Daytime	-0.02	(-0.07, 0.03)			0.44
Nighttime	-0.02	(-0.07, 0.03)			0.36
Day length, (per 30 min)	0.02	(-0.14, 0.18)			0.82
Basic Characteristics					
Age, (per 1 year)	0.05	(0.02, 0.09)	0.08	(0.04, 0.12)	<0.01
Male	1.88	(1.39, 2.36)	0.96	(0.33, 1.58)	<0.01
Body weight, (per 1 kg)	0.09	(0.07, 0.12)	0.08	(0.05, 0.11)	<0.01
Current Smoker	0.63	(-0.51, 1.76)			0.28
Alcohol intake (≥ 30 g/day)	1.19	(0.47, 1.91)	0.10	(-0.65, 0.85)	0.80
Clinical characteristics					
Anti-hypertensive use	0.19	(-0.32, 0.69)			0.47
ACEI/ARB	-0.10	(-0.65, 0.45)			0.72
CCB	0.78	(0.24, 1.33)	0.27	(-0.27, 0.82)	0.32
Diuretics	-0.13	(-1.17, 0.91)			0.81
Diabetes	0.91	(0.14, 1.68)	0.53	(-0.21, 1.28)	0.16
eGFR, per 10 ml/min/1.73m ²	-0.13	(-0.29, 0.04)	0.03	(-0.13, 0.20)	0.68
Socio-economic status					
Household income, ≥ 4 million yen/year,	-0.45	(-0.97, 0.08)	-0.37	(-0.88, 0.14)	0.16
Education, ≥ 13 years	-0.19	(-0.77, 0.38)			0.51
Life style					
Daytime PA (per 100counts/min)	-0.16	(-0.38, 0.05)	0.17	(-0.05, 0.39)	0.14
Duration spent outdoor, (per 1 min) §	0.016	(-0.11, 0.14)			0.81

ACE, angiotensin converting enzyme inhibitor; ARBs, angiotensin receptor blockers CCB, calcium channel blocker; eGFR, estimated glomerular filtration rate; PA, physical activity.

[†] β coefficient show the difference in nocturnal urinary sodium excretion rate (mmol/h) associated with 1 unit change of each covariates.

[‡] multivariate model simultaneously include all covariate with $p < 0.2$ in univariate model.

§ Included after log-transformation due to its skewed distribution.

Table 3 Adjusted nocturnal urinary sodium excretion rate by tertiles of daytime ambient temperature

models	Nocturnal urinary sodium excretion rate (95%CI)			<i>p</i> -value <i>trend</i>	<i>p</i> -value T1 vs. T3 *
	T1 (10.1±2.3°C) n=288	T2 (15.0±1.1°C) n=287	T3 (19.3±1.8°C) n=285		
model 1 †	7.51 (7.09, 7.92)	6.96 (6.54, 7.37)	6.66 (6.25, 7.08)	0.005	0.016
model 2 ‡ (day length adjusted)	7.51 (7.09, 7.92)	6.96 (6.54, 7.37)	6.67 (6.25, 7.10)	0.006	0.019
model 3 § (outdoor temperature adjusted)	7.62 (7.14, 8.10)	6.97 (6.55, 7.38)	6.54 (6.05, 7.03)	0.006	0.019
model 4 ¶ (nighttime systolic BP adjusted)	7.59 (7.12, 8.05)	7.00 (6.60, 7.41)	6.52 (6.04, 7.00)	0.005	0.017

T, Tertile; BP, blood pressure; 95%CI, 95% confidence interval; CCB, calcium channel blocker; eGFR, estimated glomerular filtration rate.

† adjusted for all variables with $p < 0.20$ in univariate regression model (Table2) such as age, gender, body weight, alcohol intake, CCB administration, diabetes, eGFR, and household income, daytime physical activity.

‡ adjusted for covariates in model 1 plus day length (per 30min).

§ adjusted for covariates in model 2 plus outdoor temperature

¶ adjusted for covariates in model 3 plus nighttime systolic blood pressure

* after Bonferoni correction for multiple comparison

Legends of Figures

Figure 1 Flow chart of participants.

Figure 2 Nocturnal Urinary Sodium Excretion by Daytime Ambient Temperature.

Error bars show 95% confidence intervals. Each column demonstrates mean values nocturnal urinary sodium excretion rate (mmol/h), and total nocturnal urinary sodium excretion (mmol) by tertile groups of daytime ambulatory temperature.

Figure 3 Nocturnal Urinary Sodium Excretion Rate and Ambulatory Blood Pressure

Error bars show 95% confidence intervals. Each column demonstrates mean values of daytime BP, nighttime BP and Dipping by tertile groups of nocturnal sodium excretion rate. Mean values were adjusted for age, gender, abdominal circumference, current smoking, alcohol intake, antihypertensive medication, and daytime physical activity.

P-values for the mean difference between T1 and T3 were estimated by Bonferroni correction.

Figure 1 Flow chart of participants

Figure 2 Nocturnal Urinary Sodium Excretion by Daytime Ambient Temperature

Nocturnal urinary sodium excretion rate

Total nocturnal urinary sodium excretion

Daytime Ambient Temperature (°C)

Figure 3 Nocturnal Urinary Sodium Excretion Rate and Ambulatory Blood Pressure