Kangaroo rats change temperature when investigating rattlesnake predators

Hannes A. Schraft*^{1,2} & Rulon W. Clark¹

Keywords

Infrared

Predator-induced stress

Predator-prey interaction

Stress-induced hyperthermia

Thermography

Abstract

Predator presence causes acute stress in mammals. A prey animal's stress response increases its chance of survival during life-threatening situations through adaptive changes in behavior and physiology. Some components of the physiological stress response can lead to changes in body surface temperatures. Body temperature changes in prey could provide information about prey state to predators that sense heat, such as pit vipers. We determined whether wild rodents undergo a stress-induced change in body surface temperature upon detecting and investigating rattlesnake predators. We staged encounters between free-ranging Merriam's kangaroo rats (*Dipodomys merriami*) and tethered Mojave rattlesnakes (*Crotalus scutulatus*) at baited feeding stations, and recorded interactions with a thermal-imaging camera. Kangaroo rats showed a significant change in maximum head temperature, snout temperature, and hind leg temperature during interactions with rattlesnakes. This supports the hypothesis that presence of a predator induces body temperature changes in prey animals. If changes in prey heat signature are detectable by heat-sensitive rattlesnakes, rattlesnakes could use this information to evaluate prey vigilance or arousal before striking; however, more detailed information on the sensory ecology of the pit organ under field conditions is needed to evaluate this possibility.

Introduction

Prey animals have evolved behavioral and physiological responses to escape predators and cope with the risks associated with predator presence (Preisser et al., 2005; Clinchy et al., 2013). The physiological stress response increases survival during acute and transitory life-threatening situations (Cannon, 1915; Hawlena and Schmitz, 2010; Clinchy et al., 2013). Predator-induced stress in mammals is a suite of neuroendocrine processes that include physiological responses such as changes in the level of stress hormones, heat shock proteins, cardiovascular activity, breathing rate, gluconeogenesis, and activity in the gastrointestinal tract (Cannon, 1915; Wingfield and Ramenofsky, 1999; Hawlena and Schmitz, 2010).

These physiological responses can result in temperature changes in prey animals. Changes in body temperature can occur due to increased cardiovascular activity induced by increased levels of the stress hormones (catecholamines and glucocorticoids; Hawlena and Schmitz, 2010). For example, exposure to predator scent directly leads to an elevated heart rate in elk (*Cervus canadensis*; Chabot et al., 1996). In a related manner, blood is shunted away from the gastrointestinal tract and to muscles as part of a fight-or-flight response, and breathing rate increases (Carere and van Oers, 2004; Sunardi et al., 2007; Willmer et al., 2009; Hawlena and Schmitz, 2010). Concurrently, acute stress results in elevated metabolism in most endotherms (Ward et al., 1996; Chabot et al., 1996; Hawlena and Schmitz, 2010), leading to a temporary rise in core body temperature (Zethof et al., 1994; Careau et al., 2011). This phenomenon, termed ‘stress-induced hyperthermia’ has been demonstrated in a variety of taxa, including rodents, primates, and birds (Baker et al., 1976; Briese and Cabanac, 1991; Zethof et al., 1994; Carere and van Oers, 2004; Careau et al., 2011). It manifests itself over the course of minutes and is often shown in response to handling (e.g. Korhonen et al., 2000). Stress-induced hyperthermia presumably also happens during acute predator stress, but this possibility has not yet been examined in wild animals (but see discussion of Rundus et al., 2007 below).

We hypothesized that these core body temperature changes translate to changes in body surface temperature when small mammals detect and interact with their predators. We studied this in kangaroo rats as they responded to a key predator, rattlesnakes. This question has additional relevance in this system due to the heat-sensing capabilities of rattlesnakes (Goris,

2011). Rattlesnakes have evolved highly specialized infrared detectors that are used to efficiently hunt prey, even in complete darkness (Noble and Schmidt, 1937; Kardong and Smith, 2002; Cadena et al., 2013). Through facial pit organs, rattlesnakes and other crotaline snakes can detect the infrared radiation emitted by endothermic animals (Goris, 2011; Cadena et al., 2013). Depending on the magnitude of the temperature change and the sensitivity of the pit organ system, temperature changes in prey animals during predator encounters may be detectable by infrared-sensitive snakes. For infrared-sensitive predators, one potential avenue of prey evaluation is prey body temperature, and snakes may use information gained through infrared radiation to evaluate prey condition or degree of vigilance. In the only example to date, tail flagging California ground squirrels (*Otospermophilus beecheyi*) showed an increase in tail temperatures of up to 3°C when signaling at rattlesnakes, which in turn was associated with a behavioral reaction by snakes to this temperature change (Rundus et al., 2007).

In this study, we experimentally exposed free-ranging kangaroo rats (a common prey item of many rattlesnake species) to tethered rattlesnakes at baited feeding sites and quantified kangaroo rat body surface temperature over the course of their interaction. We hypothesized that the presence of a major predator would represent a stressor sufficient to cause measureable increases in body temperature.

Methods

Study site and animals

Experimental trials took place outside of Rodeo, New Mexico, USA (31.888° N 109.031° W), between June 5 and August 4, 2015. Located in the Chihuahan Desert, the 200 ha study area consists of mesquite shrubland and supports large populations of Mojave rattlesnakes (*Crotalus scutulatus*) and Merriam's kangaroo rats (*Dipodomys merriami*). We captured adult kangaroo rats using Sherman live-traps baited with sunflower seeds (N = 59 individuals, 32m/27f, weight = 52.2 g ± 15.3 SD, hind foot length = 35.6 mm ± 3.4 SD). The study period coincides with reproductive activity in this population of Merriam's kangaroo rats (pers. obs.). We marked all individuals with a numbered metal ear tag and unique Nyanzol pelage dye markings for visual

identification of individuals from a distance. Adult rattlesnakes were collected by us at night from nearby roads (N = 4) or provided to us by the Chiricahua Desert Museum (N = 2). We only used snakes large enough to prey on adult kangaroo rats (snout-vent length 525-705 mm, 122-292 g). Snakes were housed individually in plastic bins between trials and provided with water and food. All snakes were returned to their original location upon completion of the experiment. The San Diego State University Institutional Animal Care and Use Committee (APF 13-08-015C) approved all methods.

Experimental design

We used sunflower seeds to attract individual marked kangaroo rats to bait stations near their original trapping location. Capture and trials were separated by at least two days in all cases. A trial started when, upon discovering bait stations, individuals began shuttling seeds away for caching. We recorded kangaroo rats during this period for up to 10 minutes using a thermal imaging camera (see below) to measure baseline body temperature. We then used a temporary absence of the kangaroo rat to tether a live rattlesnake at the bait station. Snakes were tethered by using a combination of athletic tape and duct tape to attach small metal carabiners on two points along the body of the snake, which were in turn secured to metal stakes in the ground. We kept snake body temperature cooled in between trials by housing them in secured plastic bins with insulated icepacks; this facilitated handling during tethering. Although cooling snakes undoubtedly affected their behavior, snakes in this study were used only as stimuli to measure the potential effects of rattlesnake presence on kangaroo rat temperature. When returning to the bait station, the kangaroo rat discovered the snake and typically interacted with it intermittently for up to 10 min, at which point we terminated the trial. Figure 1 shows the trapping and trial sequence. Discovery of the snake was indicated by a freeze and stare response, followed by close approaches and jumps backward away from the snake (see Randall et al., 1995 and Clark et al., 2016 for detailed characterizations of the antisnake behavior of kangaroo rats).

In order to gain insight into whether potential temperature changes were caused by the presence of a predator or simply the appearance of a novel feature at the bait station, we performed a small number of trials with an inanimate object (empty plastic jar) instead of

rattlesnake predator. Due to logistical constraints in the field, the number of these trials was much smaller than the number of rattlesnake trials (60 rattlesnake trials vs. 17 control trials). Table 1 details the number of measurements, individuals, and trials in each trial type. Some individuals were tested multiple times. Trials on the same individual were separated by at least two full days. Trials took place between 21:00h and 05:00h, during which time free-ranging kangaroo rats and Mojave rattlesnakes tend to be naturally active during summer months.

We recorded kangaroo rats from a distance of ~2 meters throughout each trial with a FLIR T-420 thermal imaging camera (FLIR Systems, Wilsonville, OR, USA) connected to a laptop running FLIR Tools+ v5.4.1 software. The camera automatically calibrated itself intermittently against an internal thermocouple. Emissivity was assumed to be 0.95 (Speakman and Ward, 1998; Tattersall et al., 2004).

Temperature variables and data extraction

We extracted still frames from infrared videos every 4 s and measured the temperature of four locations on the kangaroo rat's body using FLIR Tools+ v5.4.1 infrared imaging software. We measured the temperature of the head, snout, tail base, and hind leg (Figure 2). For head temperature, we measured maximum temperature (eye in all cases). For hind leg measurements, an ellipse was drawn that included the maximum possible area of a kangaroo rat's hind leg, and we recorded the average temperature of the area covered by the ellipse. Because of the dynamic nature of the interaction, we were not able to record all temperatures from all still frames. Table 1 shows the total number of temperature measurements taken at each body location. We only used temperature measurements taken within 10 min of the start of each experimental period (initial bait station discovery, addition of rattlesnake or addition of plastic jar).

Statistical analysis

We used linear mixed models to compare baseline kangaroo rat body temperatures with temperatures after discovery of a rattlesnake or plastic jar (Bolker et al., 2009). We fitted separate models with each measured body part (head, snout, tail base, hind leg) as the response variable and experimental condition (baseline, rattlesnake, control) as fixed effects. We included random

intercepts for snake ID, trial ID, and kangaroo rat ID (some kangaroo rats were the subjects of multiple trials, see above). We included trial ID as a random intercept because kangaroo rats had slightly different baseline temperatures in every trial. Circadian variation can explain differences in body temperature of about 2°C (Willmer et al., 2009), and body surface temperature tracks ambient temperature to an extent (e.g. Hill et al., 1980; Shine and Sun, 2003; Willmer et al. 2009). We therefore included the covariates ‘time since sunset’ and ‘ambient temperature’. In addition, we included a simple first-order autoregressive correlation structure (corAR1) in the residuals to account for temporal autocorrelation between sequential temperature measurements (Zuur et al., 2009). Our dataset was unbalanced (60 rattlesnake trials, 17 control trials), which can prevent correct model fitting. We fit all models in R v3.2.3 (R Core Team, 2015) using the package *nlme* (Pinheiro et al., 2016), which is able to handle unbalanced datasets well (Pinheiro and Bates, 2000). Nevertheless, we verified that models fitted correctly by checking the confidence intervals of the variance components after fitting the models. We did not detect unusually large intervals, indicating that the models fitted correctly (Pinheiro and Bates, 2000). Means and standard errors for plotting were calculated using the *effects* package (Fox, 2003).

Results

Kangaroo rat body surface temperatures changed significantly after rattlesnake discovery at all measured body locations except for the tail base (Table 2, Figure 3a-d). After rattlesnake presentation, head temperature decreased 0.24°C (± 0.06 SE, $p < 0.001$), snout temp decreased 0.22°C (± 0.11 SE, $p < 0.001$), and hind leg temperature increased 0.19°C (± 0.07 SE, $p = 0.007$); average tail base temperature did not change ($p = 0.158$). We did not test for significance of the individual ID random effect because we lacked a sufficient number of individuals that were tested multiple times. ‘Ambient temperature’ was a significant predictor of surface temperature at all locations. The effect of ‘time since sunset’ was very small in all cases, but significant in the model of hind leg temperature. Control trials with a novel but non-threatening stimulus (plastic jar) elicited no surface temperature changes (Table 2, Figure 3). Kangaroo rats initially behaved the same way to the jar as they did to a rattlesnake, by performing a jump-back

maneuver and then approaching the object. However, kangaroo rats quickly resumed feeding and/or caching seeds after this initial inspection.

Discussion

We found strong evidence of a change in kangaroo rat body surface temperature upon discovery of a rattlesnake. Head and snout temperatures decreased when there was a rattlesnake present, while hind leg temperature increased. Temperature changes were small ($0.19 - 0.24^{\circ}\text{C}$) but highly significant (Table 2); the magnitude of the observed changes is similar to the magnitude of body temperature changes found in chipmunks after handling stress (*Tamias striatus*; Careau et al., 2011). We did not observe body surface temperature changes when kangaroo rats discovered a novel but harmless object, giving us confidence that observed changes were due to predator presence. We conclude that in kangaroo rats, exposure to a rattlesnake predator leads to physiological responses that manifest as changes in body surface temperature.

Contrary to our predictions, we found that head and snout temperatures decreased after exposure to a rattlesnake, whereas other studies have found that stress leads to increases in body temperature (Korhonen et al., 2000; Careau et al., 2011). Stress-induced hyperthermia should increase body temperature by temporarily elevating metabolic rate (Carere and van Oers, 2004; Zhang et al., 2010; Careau et al., 2011). However, other studies have found that body surface temperatures can decrease during stress (Baker et al., 1976; Nakayama et al., 2005). The decrease in head temperature could be explained by a decrease in overall head surface temperature as a side effect of decreased snout temperature. The observed decrease in snout temperature likely resulted from evaporative cooling due to an increase in breathing rate. Elevated ventilation rates in response to acute stress have been observed in birds and fish (e.g. Carere and van Oers, 2004; Sunardi et al., 2007) and presumably also take place in kangaroo rats. Increases in breathing rate have also been directly linked to lower rostrum temperatures in rattlesnakes (*C. durissus*; Cadena et al., 2013). Hind leg temperature increased as a function of rattlesnake presence. This change is most likely due to blood shunting to the primary locomotory

muscles as a part of the stress response (Cannon, 1915; Willmer et al., 2009). Different mechanisms are not mutually exclusive and may even interact. For example, redistribution of blood could have amplified temperature changes at the snout in kangaroo rats (Nakayama et al., 2005; Hawlena and Schmitz, 2010). Overall, changes in head, snout, and hind leg temperatures point to blood redistribution, blood shunting, stress-induced hyperthermia and/or elevated breathing as possible physiological mechanisms.

Unlike head, snout, and hind leg temperatures, tail base temperature was not affected by rattlesnake presentation. This was surprising since we expected temperature changes to be most easily measurable at the mostly furless and uninsulated tail. Ambient temperature had by far the largest effect in the tail base model. We therefore suspect that tail temperature in kangaroo rats is closely linked to ambient temperature, perhaps as part of a thermoregulatory mechanism as found in other rodents (Raman et al., 1983). Ambient temperature was a significant predictor of body surface temperature in all models. This was not surprising since the surface temperature of even an imperfectly insulated animal should vary with ambient temperature to an extent (e.g. Hill et al., 1980; Shine and Sun, 2003; Willmer et al., 2009). The estimate of the effect of ‘time since sunset’ was small in all cases, and it was a significant predictor only of hind leg temperature.

Since temperature changes in kangaroo rats were small, it is unclear if rattlesnakes could use this information to evaluate prey before striking. Rattlesnakes can detect infrared radiation from prey animals (Goris, 2011) and electrophysiological studies have shown that the pit membrane is sensitive to temperature changes as small as 0.003°C (Bullock and Diecke, 1956; de Cock Buning et al., 1981). A behavioral study on western diamondback rattlesnakes (*C. atrox*; Ebert and Westhoff, 2006) suggested pit organs are even more sensitive than the threshold reported by Bullock and Diecke (1956). Rattlesnakes can also detect thermal contrast (difference in temperature between an object and its surroundings; Van Dyke and Grace, 2010); temperature decreases in some body parts should therefore lead to an altered thermal outline of the prey animal against the background. Although it is unlikely that a prey temperature difference of 0.2°C at a typical striking distance of about 30 cm would translate to a 0.003°C temperature change on the pit organ membrane (Bakken and Krochmal, 2007), our estimates of the magnitude of body temperature changes in kangaroo rats are coarse. We used values taken from

periodic scan samples of body temperature for statistical analyses. It is possible that this methodology masks fluctuations in small mammal body temperature that occur for brief periods, so that the full degree of thermal heterogeneity is not captured by our methodology. Additionally, our current understanding of snake infrared detection is primarily based on models rather than measured empirically; more experiments in ecologically relevant settings are needed to evaluate the sensory parameters of this system and its role in predatory behavior.

If snakes can detect the small temperature changes observed in this study, they may use this information to evaluate potential prey. Predators strategically evaluate the condition or degree of vigilance of potential prey during predator-prey interactions (Bradbury and Vehrencamp, 2011). Accurate prey evaluation is critical to hunting success, and predators profit from making accurate hunting decisions informed by prey cues (Lima, 2002). Rattlesnake strike success rate in the wild is typically less than 50%, with most missed strikes being attributable to rapid evasive maneuvers by prey (Clark et al., 2012). Furthermore, kangaroo rats and ground squirrels that have recently interacted with rattlesnakes and are in a state of heightened vigilance exhibit quicker reaction times and evasive startle movements in response to surprise attacks (Putman and Clark, 2015; Freymiller et al., in review). Therefore, accurate evaluation of prey vigilance is especially important for sit-and-wait predators like rattlesnakes. The temperature changes we observed in kangaroo rats are likely indicative of physiological arousal and predator awareness. Importantly, temperature changes would be relevant to rattlesnakes regardless of whether the cause of arousal was a predator or another stimulus in the environment. Stress mediated changes in body surface temperature like the ones reported here may cause an endothermic animal to essentially ‘leak’ information to snake predators. Snakes might take advantage of this leaked information by adjusting predatory behavior accordingly, for example by deciding to only strike from shorter distances at animals that appear to be aroused. To date there is only one study that has included data relevant to this possibility: tail flagging ground squirrels exhibit tail temperatures up to 3°C higher when interacting with rattlesnakes in captivity, who in turn show a behavioral response by retreating from the squirrel (Rundus et al., 2007). Although this temperature change is larger than the changes elicited in this study, it highlights the possibility of a much broader phenomenon involving interactions between pit

vipers and their endothermic prey. Temperature-mediated pursuit deterrence signals, or perhaps even thermal camouflage, remain avenues of research open to exploration. Further studies will address whether snakes respond to the temperature changes we observed in kangaroo rats.

Acknowledgements

Regina Spranger, Frank Accardo, Colin Goodman, Shannon Whelan, Grace Freymiller, and Malachi Whitford marked kangaroo rats and aided in carrying out trials. The Chiricahua Desert Museum provided logistic support and two of the rattlesnakes used. Bill and Jill Cavaliere granted permission to work on their land. We thank Shannon Whelan, Gail Patricelli, members of the Clark and Patricelli labs, and two anonymous reviewers for insightful comments on study design and earlier versions of this manuscript.

References

- Baker, M. A., Cronin, M. J. and Mountjoy, D. G. (1976). Variability of skin temperature in the waking monkey. *Am. J. Physiol.* 230, 449-455.
- Bakken, G. S. and Krochmal, A. R. (2007). The imaging properties and sensitivity of the facial pits of pitvipers as determined by optical and heat-transfer analysis. *J. Exp. Biol.* 210, 2801–2810.
- Bradbury, J. W. and Vehrencamp, S. L. (2011). *Principles of Animal Communication*. Sunderland, USA: Sinauer Associates
- Bolker, B. M., Brooks, M. E., Clark, C. J., Geange, S. W., Poulsen, J. R., Stevens, M. H. H. and White, J.-S. S. (2009). Generalized linear mixed models: a practical guide for ecology and evolution. *Trends Ecol. Evol.* 24, 127–135.
- Briese, E. and Cabanac, M. (1991). Stress hyperthermia: physiological arguments that it is a fever. *Physiol. Behav.* 49, 1153–1157.
- Bullock, T. and Diecke, F. (1956). Properties of an infra-red receptor. *J. Physiol.* 134, 47–87.
- Cadena, V., Andrade, D. V., Bovo, R. P. and Tattersall, G. J. (2013). Evaporative respiratory cooling augments pit organ thermal detection in rattlesnakes. *J. Comp. Physiol. A, Neuroethol. Sensory, Neural, Behav. Physiol.* 199, 1093–1104.
- Cannon, W. (1915). *Bodily changes in pain, hunger, fear, and rage*. New York, USA: Appleton-Century-Crofts.

Careau, V., Réale, D., Garant, D., Speakman, J. R. and Humphries, M. M. (2011). Stress-induced rise in body temperature is repeatable in free-ranging Eastern chipmunks (*Tamias striatus*). *J. Comp. Physiol. B, Biochem. Syst. Environ. Physiol.* 182, 403–414.

Carere, C. and van Oers, K. (2004). Shy and bold great tits (*Parus major*): body temperature and breath rate in response to handling stress. *Physiol. Behav.* 82, 905–912.

Chabot, D., Gagnon, P. and Dixon, E. A. (1996). Effect of predator odors on heart rate and metabolic rate of wapiti (*Cervus elaphus canadensis*). *J. Chem. Ecol.* 22, 839–868.

Clark, R., Tangco, S. and Barbour, M. (2012). Field video recordings reveal factors influencing predatory strike success of free-ranging rattlesnakes (*Crotalus* spp.). *Anim. Behav.* 84, 183–190.

Clark, R. W., Dorr, S. W., Whitford, M. D., Freymiller, G. A. and Putman, B. J. (2016). Activity cycles and foraging behaviors of free-ranging sidewinder rattlesnakes (*Crotalus cerastes*): the ontogeny of hunting in a precocial vertebrate. *Zoology* 119, 196–206.

Clinchy, M., Sheriff, M. J. and Zanette, L. Y. (2013). Predator-induced stress and the ecology of fear. *Funct. Ecol.* 27, 56–65.

de Cock Buning, T., Terashima, S. and Goris, R. C. (1981). Crotaline pit organs analyzed as warm receptors. *Cell. Mol. Neurobiol.* 1, 69–85.

de Cock Buning, T. (1983). Thermal Sensitivity as a Specialization for Prey Capture and Feeding in Snakes. *Am. Zool.* 23, 363–375.

Ebert, J. and Westhoff, G. (2006). Behavioural examination of the infrared sensitivity of rattlesnakes (*Crotalus atrox*). *J. Comp. Physiol. A* 192, 941–7.

Fox, J. (2003). Effect Displays in R for Generalised Linear Models. *J. Stat. Softw.* 8, 1–27.

Freymler, G. A., Whitford, M., Higham, T. E. and Clark, R. W. Recent interactions with snakes enhance escape performance of desert kangaroo rats exposed to a simulated attack. *Funct. Ecol.*, In Review.

Goris, R. C. (2011). Infrared Organs of Snakes: An Integral Part of Vision. *J. Herpetol.* 45, 2–14.

Hawlena, D. and Schmitz, O. J. (2010). Physiological Stress as a Fundamental Mechanism Linking Predation to Ecosystem Functioning. *Am. Nat.* 176, 537–556.

Hill, R. W., Beaver, D. L. and Veghte, J. H. (1980). Body Surface Temperatures and Thermoregulation in the Black-Capped Chickadee (*Parus atricapillus*). *Physiol. Zool.* 53, 305–321.

Kardong, K. V. and Smith, T. L. (2002). Proximate Factors Involved in Rattlesnake Predatory Behavior: A Review. In *Biology of the Vipers* (ed. G. W. Schuett, M. Hoggren, M. E. Douglas and H. W. Greene), pp. 253–266. Eagle Mountain, USA: Eagle Mountain Publishing.

Korhonen, H., Hansen, S., Malmkvist, J. and Houbak, B. (2000). Effect of capture, immobilization and handling on rectal temperatures of confident and fearful male mink. *J. Anim. Breed. Genet.* 117, 337–345.

Lima, S. L. (2002). Putting predators back into behavioral predator – prey interactions. *Trends Ecol. Evol.* 17, 70–75.

Nakayama, K., Goto, S., Kuraoka, K. and Nakamura, K. (2005). Decrease in nasal temperature of rhesus monkeys (*Macaca mulatta*) in negative emotional state. *Physiol. Behav.* 84, 783–790.

Noble, G. and Schmidt, A. (1937). The Structure and Function of the Facial and Labial Pits of Snakes. *Proc. Am. Philos. Soc.* 77, 263–288.

Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D. and R Core Team. (2016). nlme: Linear and Nonlinear Mixed Effects Models. R package version 3.1-124, <http://CRAN.R-project.org/package=nlme>

Preisser, E. L., Bolnick, D. I. and Benard, M. F. (2005). Scared to death? The effects of intimidation and consumption in predator-prey interactions. *Ecology* 86, 501–509.

Putman, B. J. and Clark, R. W. (2015). The fear of unseen predators: ground squirrel tail flagging in the absence of snakes signals vigilance. *Behav. Ecol.* 26, 185–193.

R Core Team. (2015). R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing. <http://www.R-project.org/>

Raman, E. R., Roberts, M. F. and Vanhuyse, V. J. (1983). Body temperature control of rat tail blood flow. *Am. J. Physiol - Reg I* 245, 426-432.

Randall, J., Hatch, S. and Hekkala, E. (1995). Inter-specific variation in anti-predator behavior in sympatric species of kangaroo rat. *Behav. Ecol. Sociobiol.* 36, 243–250.

Rundus, A. S., Owings, D. H., Joshi, S. S., Chinn, E. and Giannini, N. (2007). Ground squirrels use an infrared signal to deter rattlesnake predation. *Proc. Natl. Acad. Sci. U. S. A.* 104, 14372–14376.

Shine, R. and Sun, L. X. (2003). Attack strategies of an ambush predator: which attributes of the prey trigger a pit-viper's strike? *Funct. Ecol.* 17, 340- 348.

Speakman, J. R. and Ward, S. (1998). Infrared Thermography: Principles and Applications. *Zoology* 101, 224–232.

Sunardi, Asaeda, T. and Manatunge, J. (2007). Physiological responses of topmouth gudgeon, *Pseudorasbora parva*, to predator cues and variation of current velocity. *Aquat. Ecol.* 41, 111–118.

Tattersall, G. J., Milsom, W. K., Abe, A. S., Brito, S. P. and Andrade, D. V. (2004). The thermogenesis of digestion in rattlesnakes. *J. Exp. Biol.* 207, 579–585.

Van Dyke, J. U. and Grace, M. S. (2010). The role of thermal contrast in infrared-based defensive targeting by the copperhead, *Agkistrodon contortrix*. *Anim. Behav.* 79, 993–999.

Ward, J. F., MacDonald, D. W., Doncaster, C. P. and Mauget, C. (1996). Physiological response of the European hedgehog to predator and nonpredator odour. *Physiol. Behav.* 60, 1469–1472.

Willmer, P., Stone G. and Johnston I. (2009). *Environmental Physiology of Animals*. Hoboken, USA: Wiley-Blackwell.

Wingfield, J. C. and Ramenofsky, M. (1999). Hormones and the behavioral ecology of stress. In *Stress Physiology of Animals* (ed. P. H. M. Baum), pp. 1-51. Sheffield, UK: Sheffield Academic

Zethof, T. J., van der Heyeden, J. A., Tolboom, J. T. and Olivier, B. (1994). Stress-induced hyperthermia in mice: a methodological study. *Physiol. Behav.* 55, 109–115.

Zhang, W., Sunanaga, J., Takahashi, Y., Mori, T., Sakurai, T., Kanmura, Y. and Kuwaki, T. (2010). Orexin neurons are indispensable for stress-induced thermogenesis in mice. *J. Physiol.* 588, 4117–4129.

Zuur, A. F., Ieno, E. N., Walker, N. J., Savaliev, A. A., Smith, G. M. (2009). *Mixed Effects Models and Extensions in Ecology with R*. New York, USA: Springer Science+Business Media.

Figures

Figure 1: Timeline of kangaroo rat trapping, attracting, and trials. The first 10 min recording period is referred to as ‘baseline’, the second one as ‘rattlesnake’ or ‘control’. In control trials, we presented the kangaroo rat with a plastic jar instead of a rattlesnake during the second 10 min recording period.

Figure 2: Sample infrared images of kangaroo rat with measurement locations: (A) snout, (B), head maximum, (C) hind leg, and (D) tail base. The metal stakes used for tethering rattlesnakes are visible in the image on the right. Note that the temperature scales are slightly different in the two images.

Figure 3: Compared to baseline, kangaroo rat surface temperature changes when discovering a rattlesnake at (A) head, (B) snout, and (C) hind leg, but not (D) tail base. Temperatures did not change between baseline and control conditions. Error bars represent standard errors. Refer to Table 2 for statistical test results.

Tables

Table 1: Number of experimental trials, kangaroo rats, and temperature measurements taken at each body location.

Body location	Rattlesnake trials			Control trials		
	<i># measurements</i>	<i># individuals</i>	<i># trials</i>	<i># measurements</i>	<i># individuals</i>	<i># trials</i>
Head	1860	53	60	381	16	17
Snout	466	51	53	97	16	17
Hind leg	1142	53	60	262	16	17
Tail base	753	51	57	228	16	17
TOTAL		59	60		16	17

Table 2: Results of linear mixed models comparing baseline surface temperatures of kangaroo rats to temperatures after presentation of a rattlesnake and presentation of an inanimate object (control), with covariates for time since sunset and ambient temperature

	Estimate	Std. Error	t-value	p-value
<i>Head</i>				
Intercept	34.286	1.194	28.727	< 0.001
Rattlesnake	-0.240	0.060	-4.016	< 0.001
Control	0.020	0.116	0.175	0.861
Time since sunset	-0.002	0.001	-1.791	0.073
Ambient temperature	0.091	0.041	2.216	0.027
<i>Snout</i>				
Intercept	22.675	1.878	12.077	< 0.001
Rattlesnake	-0.224	0.107	-2.106	0.036
Control	0.059	0.209	0.283	0.777
Time since sunset	-0.003	0.002	-1.480	0.140
Ambient temperature	0.172	0.065	2.663	0.008
<i>Hind leg</i>				
Intercept	24.690	1.987	12.424	< 0.001
Rattlesnake	0.192	0.071	2.711	0.007
Control	0.230	0.129	1.779	0.075
Time since sunset	-0.007	0.002	-3.619	< 0.001
Ambient temperature	0.156	0.068	2.314	0.021
<i>Tail base</i>				
Intercept	19.014	2.953	6.438	< 0.001
Rattlesnake	-0.233	0.165	-1.412	0.158
Control	0.432	0.277	1.563	0.118
Time since sunset	-0.003	0.003	-1.037	0.300
Ambient temperature	0.413	0.101	4.091	< 0.001