1 Cheirolepidiaceae diversity: an anatomically preserved pollen cone
2 from the Lower Jurassic of southern Victoria Land, Antarctica

3
4
5 Timothy J. Hieger^{1*}, Rudolph Serbet¹, Carla J. Harper¹, Thomas N. Taylor¹, Edith L. Taylor¹,
6 Erik L. Gulbranson²

7
8
9
10 ¹Department of Ecology and Evolutionary Biology, and Natural History Museum and
11 Biodiversity Institute, University of Kansas, Lawrence, KS 66045-7534, U.S.A.

12 ²Department of Geosciences, University of Wisconsin-Milwaukee, Milwaukee, WI 53201-0413,
13 U.S.A.
14

15
16
17 * Corresponding author. Tel.: ++1 785-864-4255; E-mail address: tihm@ku.edu
18
19
20
21
22
23
24

1. Introduction

Jurassic floras are well known from the Antarctica Peninsula area. Members of the Swedish South Polar Expedition (1901-1903) collected some of the first plant specimens from Hope Bay on Trinity Peninsula (Halle, 1913; Cantrill and Poole, 2012), which has been variously dated as Lower Jurassic to Lower Cretaceous (Gee, 1987; 1989; Rees and Cleal, 2004). Petrographic and geochronological analyses by Hunter et al. (2005) indicated that the Botany Bay Group is Middle Jurassic. The floras are diverse, e.g., liverworts (*Schizolepidella* spp.), sphenophytes (*Equisetum* spp.), ferns (e.g., *Cladophlebis*, *Coniopteris*, *Dictyophyllum*, *Goeppertella*, *Hausmannia*, *Matonidium*, *Spesia*, *Sphenopteris*, *Todites*), seed ferns (e.g., *Archangelskya*, *Dicroidium*, *Komlopteris*, *Pachypteris*), Caytoniales (*Caytonanthus*), cycadophyte foliage (e.g., *Ctenis*, *Otozamites*, *Pseudoctenis*, *Taeniopteris*, *Weltrichia*, *Williamsonia*, *Zamites*), conifers (e.g., *Araucarites*, *Brachyphyllum*, *Elatocladus*, *Pagiophyllum*) (Gee, 1987, 1989; Rees and Cleal, 2004, and references therein; Ociepa, 2007; Ociepa and Barbacka, 2011).

In contrast to the Antarctic Peninsula, little is known about Jurassic plant life on the Antarctic continent. The Early Jurassic was the beginning of the break-up of Gondwana. Extensive volcanism at this time led to the emplacement of the Ferrar Group, a large igneous province that is related to the continental flood basalts of South Africa (Karoo) and South America (Chon Aike) (Riley and Knight, 2001), along the central Transantarctic Mountains. Fossil plants of Jurassic age on the continent are found only in a few, exceptional sites where sedimentary interbeds occur within the volcanics. Compression and permineralized plant fossils have been described from the Lower Jurassic Shafer Peak Formation in northern Victoria Land (Bomfleur and Kerp, 2010; Bomfleur et al., 2007, 2011a, 2011b; Schöner, et al. 2007). In

addition, several examples of silicified conifers have been described from the Jurassic of Antarctica, including trees engulfed by lava flows from northern Victoria Land (Jefferson et al., 1983), silicified tree trunks from Brebbia and Ramírez Nunataks (Antarctic Peninsula; del Valle et al., 1997), a single tree stump from the Coombs Hills in southern Victoria Land (Garland et al., 2007), and permineralized woody axes from Suture Bench east of the Gair Mesa in northern Victoria Land (Bomfleur et al., 2011b; Harper et al., 2012). Permineralized peat deposits from Storm Peak in the central Transantarctic Mountains have yielded an anatomically preserved fern petiole, *Polyphacelus stormensis* (Yao et al., 1991). Of these rare Jurassic sites, to date, the most extensive and fossiliferous deposit is at Carapace Nunatak in southern Victoria Land (Townrow, 1967a; Ball et al., 1979; Babcock et al., 2006).

In addition to silicified Jurassic wood, there is evidence for reproductive remains of conifers from Antarctica (e.g., Plumstead, 1962, Townrow 1967a; 1967b). Dispersed staminate cones belonging to *Classostrobus* Alvin, Spicer et Watson (Alvin et al., 1978) and containing *in situ* *Classopollis* pollen, and other cheirolepidiaceous cones have been found on other continents (see Van Konijnenburg-van Cittert 1987; Watson 1988; supplemental material); these are often attached to or associated with other macrofossils belonging to the family. Most of what is known about cheirolepidiaceous fossils is from compression-type preservation; discoveries and descriptions of permineralizations are rare. Only a single permineralized cheirolepidiaceous pollen cone has been described previously, *Classostrobus crossii* Rothwell et al. (2007), from Middle Jurassic calcarerous nodules from southern England (Rothwell et al., 2007). Here we describe a new species of anatomically preserved conifer pollen cone based on five specimens, belonging to the Cheirolepidiaceae, that were discovered in Lower Jurassic sedimentary interbeds on Carapace Nunatak, southern Victoria Land, Antarctica. The specimens described in

this report are the first examples of permineralized cheirolepidiaceae pollen cones from the Southern Hemisphere, and offer additional details about the anatomy and histology of the staminate cones of this extinct family.

Geological setting, materials, and methods

2.1 Geological setting

The collection site is located within the Ferrar Large Igneous Province (FLIP), which is a linear 3500 km belt of tholeiitic rocks (Elliot and Fleming, 2004) that contains occasional permineralizations. The cones are preserved in lake sediments within laminated chert. The Lower Jurassic chert blocks containing the fossils used in this study were collected from a moraine-debris field on the E–NE side of Carapace Nunatak, southern Victoria Land, central Transantarctic Mountains, Antarctica (76° 53' S, 159° 25' E, elevation 2149 m, Convoy Range Quadrangle) during the 1993–1994, 1997–1998, and 2011–2012 field seasons (Figure 1, A–C). The rocks of the Kirkpatrick Basalt and Carapace Sandstone Member of the Mawson Formation at Carapace Nunatak are cross-cut by several faults (Bradshaw, 1987), but generally consist of subaerial basalt flows overlain by the volcanoclastic sandstone and breccia of the Carapace Sandstone. This is overlain by subaerial basaltic flows and columns of palagonite and basaltic pillows of the Kirkpatrick Basalt (Ballance and Watters, 1971). The fossil-bearing cherts occur as 2–6 m long pods or lenses set between palagonitic breccia and pillow basalts (Figure 1, D). The cherts display several physical features (Figure 1, E): millimeter-scale laminations, flame structures, diapirs, millimeter-scale normal faults that penetrate some laminae, rupturing of lamina sets, and intense folding of the lamina sets. A rough stratification of the chert and

limestone lenses exists, but lateral continuity of the deposits does not occur due to the prevalence of subaqueous basaltic lava flows in this stratigraphic interval.

The deformation features indicate the material was initially soft and underwent a loss in volume prior to or concurrent with lithification. These lenses are interpreted as either rafts of lacustrine sediment during a single prolonged emplacement of a lava delta, or as rafts of material that was deposited in between two successive lava flows (Bradshaw, 1987). The fossil-bearing cherts are interpreted as freshwater (lacustrine) deposits that have been affected by various aspects of active volcanism (Ballance and Watters, 1971). Although our recognition of different scales of soft-sediment deformation and volume-loss features within the chert layers is consistent with Magadi-type chert (Eugster, 1967; Surdam et al., 1972; Krainer and Spötl, 1998), which forms from the early diagenetic conversion of the Na-rich phyllosilicate mineral magadiite to chert in alkaline lakes or from alkaline groundwaters (Rooney et al., 1969). Magadi-type chert is unique among abiogenic cherts in that it preserves evidence of soft-sediment deformation and desiccation due to its formation from soft beds of magadiite (Krainer and Spötl, 1998). At the type locality, Lake Magadi in the Kenyan Rift Valley, magadiite is thought to form via a Na-rich influx of hydrothermal fluid and a high silicon concentration (Jones et al., 1967). The presence of Magadi-type chert in these lacustrine deposits would suggest that the Early Jurassic lake near Carapace Nunatak was likely alkaline, possibly fed by hydrothermal fluids. The presence of limestone lenses between palagonitic and basaltic pillows attests to lake water pH values near neutral, and likely indicates temporal variation, at an unknown scale, in the geochemistry of these lake waters.

2.2 Materials

The fossiliferous chert blocks from Carapace Nunatak were initially described by Gunn and Warren (1962), and preliminary studies describing and discussing the fossil floras and faunas have been investigated by multiple authors (e.g., Plumstead, 1962, Townrow, 1967a;1967b, Babcock et al., 2006, Ribecai, 2007). Within the chert blocks used in this study, we have observed an abundance of fusainized plant material and animal remains. The fine-grained matrix has preserved several layers that contain abundant plant fossils, including various taxa of fern pinnules, rachides and rhizomes, at least three conifer taxa represented by vegetative and reproductive structures, isolated ovules, and cycadophyte foliage. Intermixed within this flora are various arthropods—an abundance of shallow-water dwelling conchostracans and other ephemeral freshwater crustaceans (Ballance and Watters, 1971; Ball et al., 1979; Shen, 1994).

2.3 Methods

Five permineralized pollen cones were analyzed in this study. The pollen cones are preserved in various planes of section and range from complete to fragmentary. Individual slabs containing two cones on the same surface are indicated as alpha and beta. Chert blocks were cut, polished, etched with 48% hydrofluoric acid, rinsed, and peeled using the standard acetate peel technique (Joy et al., 1956; Galtier and Phillips, 1999). A total of 177 serial peels were mounted on glass slides with Eukitt® mounting medium (O. Kindler, Freiburg, Germany). Pollen diameters were acquired using a Zeiss transmitted-light compound microscope. Scanning electron microscopy (SEM) images were acquired from peels mounted on SEM stubs with two-sided tape and were sputter-coated with gold (15 nm) using a Q150T Turbo-pumped Sputter

Coater (Quorum Technologies). Images were captured by a Versa 3D dual beam Scanning Electron Microscope / Focused Ion Beam (FEI, Hillsboro, OR, USA). All SEM images were obtained at an acceleration voltage of 20 kV, spot size 4.0, and were acquired with an Everhart Thornley (ET) detector using built-in software (XT Microscope) of the FEI company. Digital images of surface peels were taken on a Leica MZ16 manual fluorescence stereomicroscope with an attached Leica DC500 digital camera. Images of slide-mounted peels were captured with a Leica DC500 digital camera on a Leica DM5000B compound microscope. Additional measurements were acquired using ImageJ 1.43u software (Abràmoff et al., 2004) and edited (i.e., brightness, contrast, and color balance) using Adobe Photoshop CS4 version 11.0.2 (1990–2010, © Adobe Systems). Pollen terminology is in concordance with Punt et al. (2007); note that terms sexine 1, 2, 3 are in accordance with Punt et al. (2007) and differ from the same terms in Taylor and Alvin (1984). Specimens include accession numbers: 17570 G bot, 17607 A1 and B top, 17613 F top alpha and beta, 17654 F3 bot, peels, SEM stub: J 2138, and slides: 29,698–29,719, 29,744–29,856, and 29,910–29,951 and are housed in the Division of Paleobotany Collections, Natural History Museum and Biodiversity Institute, University of Kansas (KUPB).

3. Results

3.1 Systematic paleobotany

Division: Coniferophyta

Class: Coniferopsida

Order: Coniferales

158 **Family:** Cheirolepidiaceae Takhtajan

159 **Genus:** *Classostrobus* Alvin, Spicer et Watson

160 **Species:** *Classostrobus elliotii* Hieger, Serbet, Harper, Taylor, Taylor et Gulbranson, sp. nov.

161 **Specific Diagnosis:** Cones ellipsoidal with bluntly pointed apex; microsporophylls stalked,
162 peltate (laminae rhomboidal distally), and helically arranged on the main axis. Sporophyll stalks
163 inflated at the base; comprised of a singular vascular strand and thick- and thin-walled
164 parenchymatous cells with interspersed transfusion tracheids. Distal from main cone axis,
165 sporophyll lamina with transfusion tracheids and some thick-walled parenchymatous cells.
166 Pollen sacs abaxial, vertically deflexed, arranged in an annular cluster with a central vascular
167 strand, attached to sporophyll stalk midway between cone axis and sporophyll lamina. Sporangia
168 bear *in situ* *Classopollis* pollen; cryptopore on distal surface, prominent subequatorial rimula
169 present, trilete on proximal surface, supratectal elements spinulose.

170

171 **Etymology:** The specific epithet is named in honor of Dr. David H. Elliot, for his numerous
172 contributions over the past 40 years that enhance our understanding of the geology and
173 stratigraphy of continental Antarctica, especially of Jurassic volcanic deposits.

174 **Holotype** (*hic designatus*): 17,654 F bot [Accession: slide nos. 29,910–29,951, SEM stub no. J
175 2138] (Plate I, 2, 4–7, Plate II, 1–12) this paper.

176 **Paratypes:** 17,613 F top alpha (Plate I, 3, 8, 9, 10) 17,613 F top beta (Plate I, 1)

177 **Repository:** Natural History Museum and Biodiversity Institute, Division of Paleobotany
178 Collections (KUPB), University of Kansas, Lawrence, KS, USA

179 **Type Locality:** Carapace Nunatak, southern Victoria Land, Transantarctic Mountains, Antarctica
180 (76° 53' S, 159° 25' E, elevation 2149 m, Convoy Range Quadrangle)

181 **Stratigraphic horizon:** Carapace Sandstone Member, Mawson Formation, Ferrar Group

182 **Age:** Early Jurassic (Toarcian) (Elliot, 2013)

183

184 3.2 Description

185 The cones are ellipsoidal in longitudinal section, up to 5.5 mm in length with an average
186 diameter of 2.8 mm (1.0–5.2 mm, n=5), and the cone apex is bluntly pointed with helically
187 arranged imbricate, peltate-shaped microsporophylls. The distal lamina is rhomboidal in face
188 view with attenuate tips. The cone axis (Plate I, 1) is circular, approximately 0.5 mm in diameter
189 and up to 4.4 mm long. Sporophylls (Plate I, 1, 2) diverge from the main axis at approximately
190 90 degrees. Vascular tissue is poorly preserved in both longitudinal and transverse sections of the
191 cone axis. Surrounding the vascular tissue are isodiametric parenchymatous cells 68 µm (60–76
192 µm) in diameter with several cells containing opaque contents. Vascular tissue in the sporophyll
193 stalk (Plate I, 6) consists of a single bundle varying from 8 to 12 tracheids, each approximately 7
194 µm (5–10 µm) in diameter. The basal portion of stalk is about 0.7 mm in diameter narrowing to
195 approximately 0.5 mm and flaring distally (Plate I, 1, 2). At this point, the vascular strand
196 becomes diffuse and is gradually replaced by transfusion tracheids. Ultimately the transfusion
197 tissue (Plate I, 7) comprises about half of the sporophyll lamina and terminates approximately

198 halfway through the distal lamina. Transfusion cells are ovoid, range up to 51 μm in length and
199 up to 24 μm in diameter, and have scalariform wall thickenings (Plate I, 7). Resin ducts were not
200 observed in any of the cones.

201 Microsporophylls consist of an expanded base, a circular stalk, and an expanded distal
202 lamina. Distal laminae (Plate I, 4) are up to 2.7 mm long and 2.5 mm wide, and oriented
203 perpendicular to the long axis of the stalk. The sporophyll epidermis (Plate I, 5) is divided into
204 two parts on the outer distal lamina: an outer epidermis (Plate I, 5) made up of an uniseriate layer
205 of thin-walled cells 17 μm (10–33 μm) in diameter and an inner hypodermal layer (Plate I, 5)
206 constructed of thicker-walled columnar cells 21 μm (12–27 μm) long by 12 μm (8–15 μm) wide.
207 In some sections, an amorphous layer covers the epidermis and is interpreted as the remains of
208 cuticle and/or epicuticular wax. Transfusion tracheids appear clustered near the distal end of the
209 stalk along the vascular trace and extend into the distal lamina in a fan-like configuration among
210 the parenchymatous cells of the sporophyll head. Isodiametric parenchymatous cells of the
211 sporophylls are typically empty but are occasionally filled with opaque contents (Plate I, 4).
212 These cells vary from 40 to 116 μm in diameter.

213 Multiple sequences of serial peels indicate that a cluster of 4 to 7 ellipsoidal
214 microsporangia in an annular arrangement (Plate I, 2, 3) is attached to the abaxial surface of the
215 sporophyll stalk, equidistantly positioned between the distal lamina and the cone central axis
216 (Figure 2). Microsporangia average 0.5 mm long and 0.3 mm in diameter. Vascular tissue is
217 found in the center of the ring of sporangia (Plate I, 3 and 8). Some pollen sacs are devoid of
218 pollen while others contain densely packed well-preserved or degraded grains. Sporangial wall
219 cells are columnar and approximately 34 μm (20–45 μm) long by 9 μm (5–13 μm) wide. These
220 cells are primarily filled with opaque contents (Plate I, 3) or may occasionally be empty (Plate I,

9). Some sporangial walls are intact while others are ruptured possibly indicating pollen dispersal or damage to the sporangial wall. Bordering the microsporangial walls are numerous spheroidal tapetal orbicules (Plate II, 8, 12) that range from 0.8 μm to 1.9 μm in diameter.

Pollen sacs of *Classostrobilus elliotii* are densely packed with pollen grains; identical pollen is (Plate II, 1) often found dispersed in other areas of the pollen cone, e.g., along the cone axis, outer sporangial walls, sporophyll stalks, and the abaxial surfaces of the sporophyll laminae. Some of the grains are in tetrads (Plate II, 2), but most occur as individual grains. In equatorial view, the pollen is globose and shows a size difference from the proximal to the distal (or vice-versa) poles; equatorial diameters of the individual grains are approximately 30 μm (18–60 μm ; $n=123$). Examination of these grains indicates features that are consistent with *Classopollis* pollen, including a distal cryptopore (Plate II, 3, 4, 5), subequatorial furrow or rimula (Plate II, 3, 6, 7), internal equatorial striae (Plate II, 6, 7, 9), and a proximal trilete (Plate II, 8, 9). The cryptopore is 3.5–4.4 μm in diameter. The rimula extends around the pollen grain and is approximately 1.5 μm (1.0–1.8 μm) wide. In proximal view, laesurae of the trilete (Plate II, 8) are 6.6–12.8 μm long. Anastomosing bands of striae (Plate II, 6) are rarely seen in external view; however, internal views from SEM images reveal linear arrangements of infratectal columellae indicative of striations.

Details of the pollen wall are well preserved and consist of an outer sexine and inner nexine layer, and the infrastructure of the grains is reticulate to pseudoreticulate. Layers 1–3 of the sexine (Plate II, 10) are approximately 1.9 μm (1.7–2.1 μm) thick all together and are represented by infratectal columellae (sexine 1), an inner tectum (sexine 2), and supratectal spinules (sexine 3) in a discordant pattern. Supratectal spinules are conical, wide at the base and thinly tapered toward the tip. These wall structures average approximately 320 nm (240–530 nm)

long. The distal and proximal surfaces have numerous columellae (sexine 1) that protrude inwards toward the nexine and are about 890 nm (740–1000 nm) long by 450 nm (260–740 nm) wide. The tectum (sexine 2) between the spinules and columns of the sexine is approximately 330 nm (210–440 nm) thick. In some grains, the nexine (Plate II, 5, 11) is around 280 nm (210–330 nm) thick. Broken grains allow for the internal examination of structures such as the proximal trilete (Plate II, 9). This triangular structure has equidistant arms that are about 12 μ m long. Internal views of the grains reveal linear arrangements (Plate II, 7, 9) of the infratectal columellae, and 3–5 equatorial striations that appear between the columellae below the rimula.

Up to five *Brachyphyllum*-type leaves are in close association with these cones (Plate I, 2, 10). There was no direct anatomical connection between the leaves and the pollen cones used in this study. These leaves appear near the basal region of the cone and attenuate before extending the entire length of the cones; the leaves are incomplete but include some morphological and anatomical features. Leaves are 0.4–0.5 mm thick and 1.3–1.9 mm wide. Ground tissue is of isodiametric cells, 66 μ m (30–123 μ m) in diameter, which are primarily filled with opaque contents. Each leaf contains a single vascular strand (Plate I, 10), consisting of about nine metaxylem tracheids 5–6 μ m in diameter together with a few protoxylem tracheids. There is a space next to the xylem tissue that presumably represented the position of the phloem. Surrounding the vascular tissue of several associated leaves are round cells (Plate I, 10) of unknown affinities that may represent some type of secretory cell or a form of transfusion tissue. Such cells contain contents which may also represent some type of thick-walled fungal hyphae. These cells are between 10–25 μ m in diameter with walls that are 2–5 μ m thick.

4. Discussion

The Cheirolepidiaceae is a large family of important extinct Mesozoic conifers (Alvin, 1982; Watson, 1988; Axsmith and Jacobs, 2005; Escapa et al., 2012; Stockey and Rothwell, 2013), which produce a diverse range of growth habits (Watson, 1988; Taylor et al., 2009). Spanning the Late Triassic to the Cretaceous, cheirolepidiaceous plants were components of coastal environments inundated by tides (Upchurch and Doyle, 1981) and one of the dominant components of forests on several continents (Vakhrameev, 1970; Alvin, 1982; Watson, 1988; Taylor et al., 2009; Stockey and Rothwell, 2013). Cheirolepidiaceous plants were widely distributed geographically with most examples at low paleolatitudes (Alvin et al., 1978; Alvin, 1982). To date, the majority of reports of cheirolepidiaceous macrofossils are from the Northern Hemisphere with fewer examples from the Southern Hemisphere (Alvin, 1982; Francis, 1983; Watson 1988; Taylor et al., 2009; Escapa et al., 2012). *Classopollis* pollen is distributed worldwide, but is rare at higher paleolatitudes (Vakhrameev, 1970; Brenner, 1976; Alvin, 1983). The pollen type of the Cheirolepidiaceae (*Classopollis*) is present on all Gondwana continents during the Early Jurassic (Grant-Mackie et al., 2000). Therefore, the discovery of the permineralized cheirolepidiaceous pollen cones provides evidence of their presence in Antarctica during the Jurassic.

4.1 General comparisons

Previous to this study, there was a single coniferous pollen cone, *Masculostrobus warrenii* Townrow, described from the Carapace Nunatak locality (Townrow, 1967b). This cone is 8 mm long and 5 mm wide with rhomboidal microsporophyll distal laminae, but it contains

saccate pollen. Since one of the unifying characters of the Cheirolepidiaceae is *Classopollis*-type pollen (Alvin et al., 1978), *M. warrenii* clearly cannot be assigned to the Cheirolepidiaceae and was suggested by Townrow (1967b) to share affinities with the Podocarpaceae.

Classostrobus elliotii cones contain many unique characters that provide the basis for a new species, such as helically arranged microsporophylls, and a cluster of four to seven pollen sacs that are attached approximately midway on the abaxial microsporophyll stalk in an annular arrangement. The specimens used in this study are also similar in some characters to other cheirolepidiaceous pollen cones from different localities (see Van Konijnenburg-van Cittert, 1987), i.e., cones are small and simple, have helically arranged peltate microsporophylls with rhomboidal laminar heads situated around a central axis, and contain *Classopollis* pollen (Alvin, 1983; Rothwell et al., 2007). Cheirolepidiaceous cones have been found at other localities attached to or associated with vegetative remains (e.g., Barnard, 1968; Lorch, 1968; Barnard and Miller, 1976; Alvin et al., 1978, 1994; Watson, 1982; Francis, 1983; Van Konijnenburg-van Cittert, 1987; Watson, 1988; Thévenard, 1993; Gomez et al., 2002; Axsmith et al., 2004; Barbacka et al., 2007; Yang, 2008; Kim et al., 2012; see supplemental material). Isolated pollen cones that contain *Classopollis* pollen are placed within the genus *Classostrobus* (Alvin et al., 1978). Comparison is difficult with other cheirolepidiaceous cones due to the lack of stomatal information or compressional views of the cones, since most cones have been described from compression/impressions (Rothwell et al. 2007; Archangelsky and Del Fueyo, 2010).

There has been a single report of an anatomically preserved, permineralized cheirolepidiaceous cone, *Classostrobus crossii*, of southern England (Rothwell et al., 2007), which is from the Middle Jurassic. *Classostrobus crossii* is larger than *C. elliotii*, with a length of 20 mm and width of 18 mm; microsporophyll laminae are up to 11 mm long (Rothwell et al.,

2007), whereas microsporophyll laminae of *C. elliotii* are up to 2.7 mm long and the cones have a maximum diameter of 5.2 mm and length of 5.5 mm. The number of pollen sacs in *C. crossii* is described as several and the microsporangia are situated abaxially at the juncture of the stalk and basal heel of the distal lamina. The pollen sacs of *C. elliotii* occur approximately midway between the microsporophyll head and cone axis and are attached to and the abaxial surface of the microsporophyll stalk.

Characters available for comparison with previously described cones lacking internal anatomy are cone shape and size, shape of the sporophyll head, pollen sac arrangement and position, and pollen features (Rothwell et al., 2007). The largest pollen cone, *Classostrobus cathayanus* Zhou (Zhou, 1983), is 23 mm in diameter and more than 20 mm long. Cones of *Tomaxiella biforme* Archangelsky (Archangelsky and Gamero, 1967) are the smallest, 2-3 mm long and 1 mm wide. *Classostrobus elliotii* is up to 5.5 mm long and 5.2 mm wide with an average diameter of 2.8 mm. The overall shape of the microsporophylls is peltate, and the microsporophyll heads are rhomboidal in face view like most other cheirolepidiaceous cones. For further comparisons of all cheirolepidiaceous cones, see supplemental material.

4.2 Microsporangia

The pollen sacs of the Cheirolepidiaceae are typically situated on the abaxial surface of the microsporophyll attached near the microsporophyll head. For the type species, *Classostrobus rishra* (Barnard) Alvin, Spicer et Watson (Alvin et al., 1978), six to eight pollen sacs are layered on the abaxial surface of the microsporophyll stalk. The microsporangia of *C. arkansensis* Axsmith, Krings et Waselkov (Axsmith et al., 2004) are located on the abaxial side of the

sporophyll stalk near the basal inner region of the microsporophyll head. Abaxial positioning of pollen sacs is also present in cones associated with the foliage taxa *Hirmeriella airelensis* Muir et Van Konijnenburg-van Cittert (Muir and Van Konijnenburg-van Cittert, 1970), and *Frenelopsis* (Schenk) sp. (Barale et al., 1988).

The number of pollen sacs per microsporophyll is variable within *Classostrobus elliotii* cones, i.e., four to seven vertically deflexed pollen sacs, which occur in an annular cluster attached to the abaxial sporophyll stalk. When compared to other cheirolepidiaceous cones, the annular arrangement of the pollen sacs is a unique character of *C. elliotii*. Where observed, most other staminate cones of the Cheirolepidiaceae have two to three pollen sacs per microsporophyll or the number is unknown (Van Konijnenburg-van Cittert, 1987). Up to four pollen sacs are observed in *C. dalatzensis* Yang (Yang, 2008), and cones belonging to *Hirmeriella muensteri* (Schenk) Jung (Jung, 1968) have up to twelve microsporangia per microsporophyll. However, Van Konijnenburg-van Cittert (1987) observed six microsporangia per microsporophyll in *H. muensteri*.

In general, Jurassic cheirolepidiaceous pollen cones contain more pollen sacs per microsporophyll relative to Cretaceous pollen cones (see supplemental material and references therein). To date, evolutionary inferences and trends on the number and positioning of microsporangia in the Cheirolepidiaceae are based on limited reports. Increased datasets on pollen cones from permineralized material such as specimens from Carapace Nunatak may shed additional light on this aspect of pollen cone evolution within this conifer family.

4.3 Pollen

Classopollis pollen is known from the Early Triassic through the Late Cretaceous and is numerous in deposits worldwide (Taylor et al., 2009; Zavialova et al., 2010a; 2010b). Previous studies of palynofloras from Carapace Nunatak have been interpreted as late Middle Jurassic (Shang, 1997; Ribecai, 2007), which conflicts with Early Jurassic radiometric age constraints on the timing of lava emplacement (Elliot, 2013). A more recent revision of the palynoflora of Carapace Nunatak indicates that it is dominated by *Classopollis* pollen, which suggests that members of the Cheirolepidiaceae were present (Bomfleur et al., 2014). *Classopollis* is said to be the one reliable characteristic that allows assignment to the Cheirolepidiaceae, including pollen cones (Watson, 1988), and the presence of *in situ* pollen of this type is a consistent characteristic of *Classostrobus* cones (Alvin et al., 1978). *Classostrobus elliotii* contains numerous *in situ* pollen grains that would be assigned to *Classopollis*, as they exhibit a distal cryptopore, subequatorial furrow (rimula), equatorial striae, and a proximal trilete (Plate II). Striations are rarely seen in light transmitted light microscopy, but are revealed as patterns in infratectal columellae under SEM.

The average pollen diameter (30 μm) of the Antarctic grains is comparable to that of most other cheirolepidiaceous cones that contain *Classopollis* along with those from dispersed pollen assemblages. Pollen grains associated with *Classostrobus dalatzensis* are reported to be as small as 13 μm (Axsmith et al., 2004). The largest pollen grains associated with cheirolepidiaceous cones reported to date is 65 μm from cones associated with *Frenelopsis alata* (Hlustik and Konzalova, 1976). Equatorial diameters of the solitary, mature pollen within *Classostrobus elliotii* are as small as 18 μm and can be as large as 60 μm . Smaller, dispersed pollen grains associated with *C. elliotii* may be interpreted as aborted, but larger grains may be interpreted as possible forms of polyploid pollen (see Kürschner et al., 2013). The infrastructure

of *C. elliotii in situ* pollen grains is reticulate to pseudoreticulate making the striae appear to be absent or vague, which is similar to other cheirolepidaceous pollen grains from the Early Jurassic. The pollen striae of *C. elliotii* are similar to the *sporae dispersae* species *C. chateaunovi* and *C. kieseri* Reyre (Reyre, 1970) that lack or have vague striations. The combination of no striations to vague striations and possession of reticulate infratectal patterns among species with similar supratectal elements is common among Early Jurassic *Classopollis* species (Reyre, 1970). Similar to these Early Jurassic species of *Classopollis*, the striae of *C. elliotii* pollen grains are difficult to discern under light microscopy and are revealed as sinuous linear arrangements of infratectal columellae under SEM. When compared to the Middle Jurassic *Classopollis* pollen of southern England, supratectal elements of *Classostrobus crossii* grains described as spines (Rothwell et al., 2007) are similar in shape to the surface ornaments of *C. elliotii* pollen, but the infratectal columellae of *C. crossii* is differentiated so that anastomosing striae are distinct. *In situ* mature pollen grains of *C. comptonensis* from the Lower Cretaceous Wealden beds provided the opportunity of ultrastructural studies by Taylor and Alvin (1984), and revealed that the grains exhibit distinct striae with supratectal spinules. Pollen of *Classostrobus arkansensis* Axsmith et al. (2004) of the United States shares these characteristics with *C. comptonensis in situ* grains (Axsmith et al., 2004). The pollen of *C. comptonensis* is similar to the pollen of the Early Cretaceous pollen cones of Spain, *C. turolensis* and *C. ugnaensis* Gomez et al. (2002), in possessing distinct striae and microechinate supratectal ornaments, which were described based on *in situ* and dispersed pollen from the same locality (Gomez et al., 2002).

Infrastructure, supratectal ornamentation, and striations have recently been used in Argentina to assign taxonomic value to *Classopollis sporae dispersae* discussed by Reyre (1970)

and to introduce new species (Villar de Seoane, 2014), and the differentiation of infrastructure among species of *Classopollis* has been discussed by Zavialova et al. (2010a, 2010b). One concern about taxonomic placement into dispersed species based on these pollen characters is that they may represent different stages in pollen development, but the pollen characters of *Classostrobus comptonensis* discussed by Taylor and Alvin (1984) were based on samples from interpreted mature and immature cones, which helped to clarify differences between developmental morphologies of the pollen wall. Therefore, the pollen cones of *C. elliotii* (Early Jurassic), *C. crossii* (Middle Jurassic), *C. comptonensis* (Early Cretaceous) may all be interpreted as mature (i.e., complete tetrad development) with developed pollen walls, thus providing insight into ultrastructural differences of *in situ* pollen. Through comparisons with ultrastructure of pollen from other cheirolepidiaceous cones, the pollen characters of *C. elliotii* assist in supporting the hypothesis of Reyre (1970) that in the early evolution of the Cheirolepidiaceae (Late Triassic–Early Jurassic), the pollen cones produced pollen having a less differentiated arrangement of infratectal columellae than in more evolved (i.e., later) forms (Middle Jurassic–Cretaceous) yet similar supracteal elements, in this case, echinate features. If factors other than grain maturity, i.e., taphonomy, preservation, and preparation, have little effect on these aspects of *Classopollis* ultrastructure, this adds further support for the numerous different species of identified *Classopollis* pollen by advancement of morphological traits and strengthens the morphological diversity known for this extinct conifer family.

4.4 Paleobiogeography and paleoenvironment

Classostrobus elliotii extends the biogeographical distribution of macrofossils of the Cheirolepidiaceae to the southernmost Lower Jurassic paleolatitude to date. During the Lower Jurassic, Carapace Nunatak was located at ~60°S paleolatitude. Other occurrences have been noted as far south as Patagonia, Argentina (e.g., Archangelsky, 1968; Archangelsky and Gamarro, 1967; Escapa et al., 2012, 2013) and Australia (e.g., Pole, 2000; Tosolini et al., 2015). This is also the first report of macrofossil evidence of the Cheirolepidiaceae from Antarctica; therefore, the specimens used in this study extend the biogeographical distribution of this family to all present-day continents.

A number of cheirolepidiacean plants are considered to have been halophytes and were clearly thermophilous (Alvin, 1982), although the family was also successful in temperate climates (Watson, 1988). Crustaceans, insects, and plants collected from Carapace Nunatak suggest a non-arid, temperate or possibly warmer climate, based on comparisons to extant representatives (Ball et al., 1979). Deposits represent freshwater, ephemeral pond, and deep-lake environments (Stigall et al., 2008). Plants have been used as proxies for paleoclimate and as paleoenvironmental indicators (Rees et al., 2000; Taylor and Ryberg 2007). The Cheirolepidiaceae can also be used as a proxy for the Carapace Nunatak locality during the Early Jurassic, which suggests that it may have been a xeric or halophytic habitat based on specific characters of this Mesozoic conifer family. The Cheirolepidiaceae typically show xeromorphic characters, such as thick, succulent shoots, and reduced scale-like leaves, and this may explain how this plant was able to grow in a warm environment close to active volcanism. Although more paleobotanical data, e.g., cuticle and stomatal information, are needed to further understand the paleoenvironment of the locality, this discovery adds to the wide range of habitats occupied by this important Mesozoic conifer family.

445

446 6. Conclusions

447 The presence of a cheirolepidiacean pollen cone in Lower Jurassic rocks from Carapace
448 Nunatak, Antarctica bridges the gap between southern and northern Gondwanan continents and
449 provides evolutionary insight into this biogeographically important extinct conifer group. This
450 discovery extends the biogeographical range of macrofossils of the Cheirolepidiaceae during the
451 Early Jurassic to its southernmost point and is the highest paleolatitude, i.e., ~60°S paleolatitude,
452 occurrence. *Classostrobus elliotii* is the first anatomically preserved cheirolepidiacean cone
453 discovered in the Southern Hemisphere. The cones are small and simple, similar to other cones
454 of the Cheirolepidiaceae. *In situ* pollen of Early Jurassic cheirolepidiacean pollen cones like that
455 of *C. elliotii* provides additional data for preexisting hypotheses of *Classopollis* ultrastructural
456 evolution. The number and position of pollen sacs is considered taxonomically important for
457 understanding the evolution of pollen cones in conifers, and this study provides new information
458 on key taxonomic characters for a group of conifers, which is phylogenetically poorly resolved,
459 in order to better understand the evolution on the morphology of pollen cones belonging to the
460 Cheirolepidiaceae.

461 Plumstead (1962) described a small shoot belonging to *Brachyphyllum* cf. *expansum*
462 Sternberg from the Carapace Nunatak locality (Townrow 1967a). The associated leaves of
463 *Classostrobus elliotii* resemble those of *Brachyphyllum* and appear to be thick, fleshy, and
464 triangulate. Whether *C. elliotii* may have been produced by a shoot that bears this type of leaf
465 merits further investigation. Additionally, isolated ovules have been observed in cherts from the
466 Carapace Nunatak locality, which increases the likelihood of finding ovulate cones or the seed-

bearing reproductive organs of these plants. We are hopeful that additional studies in progress will likely provide the opportunity of developing a whole-plant concept for this extinct conifer.

5. Acknowledgments

Financial support was provided by the National Science Foundation (Grant ANT-1142495 to E.L.T. and T.N.T.). T.J.H. extends appreciation for summer funding through the KU Undergraduate Biology Program Research Award. We thank Dr. Prem Thappa and Ms. Heather Shinogle of the KU Bioimaging Facility for assistance in acquiring SEM images, Dr. A.-L. Decombeix (Montpellier, France) for assistance with translations, Dr. Gar W. Rothwell (Athens, OH, USA), Dr. Ruth A. Stockey (Corvallis, OR, USA) and Dr. B. Bomfleur (Stockholm, Sweden) for discussions and constructive comments on this paper, Sara L. Taliaferro for the illustration, and two anonymous reviewers for their helpful comments on an earlier draft of this manuscript.

7. References

- Abràmoff, M.D., Magalhães, P.J., Ram, S.J., 2004. Image processing with ImageJ. *Biophotonics International* 11, 36–42.
- Alvin, K.L., 1982. Cheirolepidiaceae: biology, structure and paleoecology. Review of *Palaeobotany and Palynology* 37, 71–98.
- Alvin K.L., 1983. Reconstruction of a Lower Cretaceous conifer. *Botanical Journal of the Linnean Society* 86, 169–176.

488 Alvin, K.L., Spicer, R.A., Watson, J., 1978. A *Classopollis*-containing male cone associated with
 489 *Pseudofrenelopsis*. *Palaeontology* 21, 847–856.

490 Archangelsky, S., 1968. On the genus *Tomaxellia* (Coniferae) from the Lower Cretaceous of
 491 Patagonia (Argentina) and its male and female cones. *Botanical Journal of the Linnean*
 492 *Society of London* 61, 153–165.

493 Archangelsky, S., Gamero, J.C., 1967. Pollen grains found in coniferous cones from the Lower
 494 Cretaceous of Patagonia (Argentina). *Review of Palaeobotany and Palynology* 5, 179–
 495 182.

496 Archangelsky S., Del Fueyo G.M., 2010. Endemism of Early Cretaceous conifers in western
 497 Gondwana. In: Gee C.T., (ed.), *Plants in Mesozoic Time: Morphological Innovation,*
 498 *Phylogeny, Ecosystems*, Indiana University Press, Bloomington, IN, pp. 247–270.

499 Axsmith, B.J., Krings, M., Waselkov, K., 2004. Conifer pollen cones from the Cretaceous of
 500 Arkansas: implications for diversity and reproduction in the Cheirolepidiaceae. *Journal of*
 501 *Paleontology* 78, 402–409.

502 Axsmith, B.J., Jacobs, B.F., 2005. The conifer *Frenelopsis ramosissima* (Cheirolepidiaceae) in
 503 the Lower Cretaceous of Texas: systematic, biogeographical, and paleoecological
 504 implications. *International Journal of Plant Sciences* 166, 327–337.

505 Babcock, L.E., Leslie, S.A., Elliot, D.H., Stigall, A.L., Ford, L.A., Briggs, D.E.G., 2006. The
 506 “preservation paradox”: microbes as a key to exceptional fossil preservation in the
 507 Kirkpatrick Basalt (Jurassic), Antarctica. *The Sedimentary Record* 4, 4–8.

508 Ball, H.W., Borns Jr, H.W., Hall, B.A., Brooks, H.K., Carpenter, F.M., Delevoryas, T., 1979.
 509 Biota, age, and significance of lake deposits, Carapace Nunatak, Victoria Land,
 510 Antarctica. In: Laskar, B., Rao, C.S.R. (Eds.), *Fourth International Gondwana*

511 Symposium: Papers, vol. 1. Hindustan Publishing Corporation, New Delhi, India, pp.
 512 166–175.

513 Ballance, P.F., Watters, W.A., 1971. The Mawson Diamictite and the Carapace Sandstone,
 514 formations of the Ferrar Group at Allan Hills and Carapace Nunatak, Victoria Land,
 515 Antarctica. New Zealand Journal of Geology and Geophysics 14, 512–527.

516 Barale, G., Fernandez Marron, T., Alvarez Ramis, C., 1988. Étude de cônes mâles de
 517 *Frenelopsis* Schenk emend. Reymanówna et Watson, 1976 (Cheirolepidiaceae) du
 518 Crétacé supérieur de Torrelaguna (Province de Madrid–Espagne). Géobios 21, 187–199.

519 Bomfleur, B., Kerp, H., 2010. The first record of the dipterid fern leaf *Clathropteris* Brongniart
 520 from Antarctica and its relation to *Polyphacelus stormensis* Yao, Taylor et Taylor nov.
 521 emend. Review of Palaeobotany and Palynology 160, 143–153.

522 Bomfleur, B., Schneider, J., Schöner, R., Viereck-Götte, L., Kerp, H., 2007. Exceptionally well-
 523 preserved Triassic and Early Jurassic floras from North Victoria Land, Antarctica. USGS
 524 Open-File Reports 2007–1047 (ea034): 4 pp. [Antarctica: A Keystone in a Changing
 525 World-Online Proceedings of the 10th ISAES].

526 Bomfleur, B., Pott, C., Kerp, H., 2011a. Plant assemblages from the Shafer Peak Formation
 527 (Lower Jurassic), north Victoria Land, Transantarctic Mountains. Antarctic Science 23,
 528 188–208.

529 Bomfleur, B., Schneider, J.W., Schöner, R., Viereck-Götte, L., Kerp, H., 2011b. Fossil sites in
 530 the Continental Victoria and Ferrar groups (Triassic–Jurassic) of North Victoria Land,
 531 Antarctica. Polarforschung 80, 88–99.

532 Bomfleur, B., Schöner, R., Schneider, J.W., Viereck, L., Kerp, H., McKellar, J.L., 2014. From the
 533 Transantarctic Basin to the Ferrar Large Igneous Province–New palynostratigraphic age

534 constraints for Triassic-Jurassic sedimentation and magmatism in East Antarctica.

535 Review of Palaeobotany and Palynology 207, 18–37.

536 Bradshaw, M.A., 1987. Additional field interpretation of the Jurassic sequence at Carapace

537 Nunatak and Coombs Hills, south Victoria Land, Antarctica. New Zealand Journal of

538 Geology and Geophysics 30, 37–49.

539 Brenner, G.J., 1976. Middle Cretaceous floral provinces and early migrations of angiosperms. In:

540 C.B. Beck (ed.), Origin and Early Evolution of Angiosperms, Columbia University Press,

541 NY. pp. 23–47.

542 Cantrill, D.J., 2000. A new macroflora from the South Orkney Islands, Antarctica: evidence of

543 an Early to Middle Jurassic age for the Powell Island Conglomerate. Antarctic Science

544 12, 185–195.

545 Cantrill D.J., Poole I., 2012. The Vegetation of Antarctica Through Geological Time. Cambridge

546 University Press, New York, NY, 487 pp.

547 del Valle, R.A., Lirio, J.M., Lusky, J.C., Morelli, J.R., Nuñez, H.J., 1997. Jurassic trees at Jason

548 Peninsula, Antarctica. Antarctic Science 9, 443–444.

549 Elliot, D.H., 2013. The geological and tectonic evolution of the Transantarctic Mountains: a

550 review. In: Hambrey, M.J., Barker, P.F., Barrett, P.J., Bowman, V., Davies, B., Smellie,

551 J.L., Tranter, M. (eds.), Antarctic Palaeoenvironments and Earth–Surface Processes.

552 Geological Society, London, Special Publications 381, 7–35.

553 Elliot, D.H., Fleming, T.H., 2004. Occurrence and dispersal of magmas in the Jurassic Ferrar

554 Large Igneous Province, Antarctica. Gondwana Research 7, 223–237.

555 Escapa, I.H., Rothwell, G.W., Stockey, R.A., Cúneo, N.R., 2012. Seed cone anatomy of
 556 Cheirolepidiaceae (Coniferales): reinterpreting *Pararaucaria patagonica* Wieland.
 557 American Journal of Botany 99, 1058–1068.

558 Escapa, I.H., Cúneo, N.R., Rothwell, G., Stockey, R.A., 2013. *Pararaucaria delfueyoi* sp. nov.
 559 from the Late Jurassic Cañadón Calcáreo Formation, Chubut, Argentina: insights into the
 560 evolution of the Cheirolepidiaceae. International Journal of Plant Sciences 174, 458–470.

561 Eugster, H.P., 1967. Hydrous sodium silicates from Lake Magadi, Kenya: precursors of bedded
 562 chert. Science 157, 1177–1180.

563 Francis, J.E., 1983. The dominant conifer of the Jurassic Purbeck formation, England.
 564 Palaeontology 26, 277–294.

565 Galtier, J., Phillips, T.L., 1999. The acetate peel technique. In: Jones, T.P., Rowe, N.P. (eds.),
 566 Fossil Plants and Spores: Modern Techniques. The Geological Society, London, pp. 67–
 567 70.

568 Garland, M.J., Bannister, J.M., Lee, D.E., White, J.D.L., 2007. A coniferous tree stump of late
 569 Early Jurassic age from the Ferrar Basalt, Coombs Hills, southern Victoria Land,
 570 Antarctica. New Zealand Journal of Geology and Geophysics 50, 263–269.

571 Gee, C.T., 1987. Revision of the Early Cretaceous flora from Hope Bay, Antarctica. University
 572 of Texas, Austin, Texas, USA. Ph.D. Dissertation. 165 pp.

573 Gee, C.T., 1989. Revision of the Late Jurassic/Early Cretaceous flora from Hope Bay,
 574 Antarctica. Palaeontographica. Abteilung B, Paläophytologie 213, 149–214.

575 Grant-Mackie, J.A., Aita, Y., Balme, B.E., Campbell, H.J., Challinor, A.B., MacFarlan, D.A.B.,
 576 Molnar, R.E., Stevens, G.R., Thulborn, R.A., 2000. Jurassic palaeobiogeography of
 577 Australasia. In: Wright, A.J., Talent, J.A., Young, G.C., Laurie, J.R., (eds.),

578 Palaeobiogeography of Australasian Faunas and Floras, Memoir of the Association of
 579 Australasian Palaeontologists 23, 311–354.

580 Gunn, B.M., Warren, G., 1962. Geology of Victoria Land between the Mawson and Mulock
 581 Glaciers, Antarctica. New Zealand Geological Survey Bulletin 71, 1–157.

582 Halle, T.G., 1913. The Mesozoic flora of Graham Land. Wissenschaftliche Ergebnisse der
 583 Schwedischen Südpolar-Expedition 1901–1903 3, 3–124.

584 Harper, C.J., Bomfleur, B., Decombeix, A.-L., Taylor, E.L., Taylor, T.N., Krings, M., 2012.
 585 Tylosis formation and fungal interactions in an Early Jurassic conifer from northern
 586 Victoria Land, Antarctica. Review of Palaeobotany and Palynology 175, 25–31.

587 Hlušík, A., Konzalová, M., 1976. Polliniferous cones of *Frenelopsis alata* (K. Feistm.)
 588 Knobloch from the Cenomanian of Czechoslovakia. Věstník Ústředního ústavu
 589 Geologického 51, 37–46.

590 Hunter M.A., Cantrill D.J., Flowerdew M.J., Millar I.L., 2005. Mid-Jurassic age for the Botany
 591 Bay Group: Implications for Weddell Sea Basin creation and southern hemisphere
 592 biostratigraphy. Journal of the Geological Society of London 162, 745–748.

593 Jefferson, T.H., Siders, M.A., Haban, M.A., 1983. Jurassic trees engulfed by lavas of the
 594 Kirkpatrick Basalt Group, northern Victoria Land. Antarctic Journal of the United States
 595 18(5), 14–16.

596 Jones, B.F., Rettig, S.L., Eugster, H.P., 1967. Silicate in alkaline brines. Science, 158, 1310–
 597 1314.

598 Joy, K.W., Willis, A.J., Lacey, W.S., 1956. A rapid cellulose peel technique in palaeobotany.
 599 Annals of Botany 20, 635–637.

600 Jung, W., 1968. *Hirmerella münsteri* (Schenk) Jung nov. comb., eine bedeutsame Konifere des
 601 Mesozoïkums. Palaeontographica Abteilung B Paläophytologie 122, 55–93.
 602 Krainer, K., Spötl, C., 1998. Abiogenic silica layers within a fluvio–lacustrine succession,
 603 Bolzano Volcanic Complex, northern Italy: a Permian analogue for Magadi-type cherts?
 604 Sedimentology 45, 489–505.

 605 Kürschner, W.M., Batenburg, S.J., Mander, L., 2013. Aberrant *Classopollis* pollen reveals
 606 evidence for unreduced (2n) pollen in the conifer family Cheirolepidiaceae during the
 607 Triassic–Jurassic transition. Proceedings of the Royal Society B 280, 20131708.
 608 <http://dx.doi.org/10.1098/rspb.2013.1708>

 609 Muir, M., Van Konijnenburg-van Cittert, J., 1970. A Rhaeto–Liassic flora from Ariel, northern
 610 France. Palaeontology 13, 433–442.
 611 Ociepa, A.M., 2007. Jurassic liverworts from Mount Flora, Hope Bay, Antarctic Peninsula.
 612 Polish Polar Research 28, 31–36.
 613 Ociepa, A.M., Barbacka, M., 2011. *Spesia antarctica* gen. et sp. nov.—a new fertile fern spike
 614 from the Jurassic of Antarctica, Polish Polar Research 32, 59–66.
 615 Plumstead, E.P., 1962. Fossil floras from Antarctica. Trans–Antarctic Expedition 1955–1958,
 616 Scientific Reports 9, 1–132.
 617 Pole, M.S., 2000. Mid–Cretaceous conifers from the Eromanga Basin, Australia. Australian
 618 Systematic Botany 13, 153–197.
 619 Punt, W., Hoen, P.P., Blackmore, S., Nilsson, S., Le Thomas, A., 2007. Glossary of pollen and
 620 spore terminology. Review of Palaeobotany and Palynology 143, 1–81.

621 Rees, P.M., Ziegler, A.M., Valdes, P.J., 2000. Jurassic phytogeography and climates: new data
 622 and model comparisons. In: Huber, B.T., MacLeod, K.G., Wing, S.L., (eds.), Warm
 623 Climates in Earth History, Cambridge University Press, Cambridge, U.K., pp. 297–318.

624 Rees, P.M., Cleal, C.J., 2004. Lower Jurassic floras from Hope Bay and Botany Bay, Antarctica.
 625 Special Papers in Palaeontology 72, 1–90.

626 Reyre, Y., 1970. Stereoscan observations on the pollen genus *Classopollis* Pflug 1953.
 627 Palaeontology 13, 303–322.

628 Ribecai, C., 2007. Early Jurassic miospores from Ferrar Group of Carapace Nunatak, South
 629 Victoria Land, Antarctica. Review of Palaeobotany and Palynology 144, 3–12.

630 Riley, T.R., Knight, K.B., 2001. Age of pre-break-up Gondwana magmatism. Antarctic Science
 631 13, 99–110.

632 Rooney, T.P., Jones, B.F., Neal, J.T., 1969. Magadiite from Alkali Lake, Oregon. American
 633 Mineralogist 54, 1034–1043.

634 Rothwell, G.W., Mapes, G., Hilton, J., Hollingworth, N.T., 2007. Pollen cone anatomy of
 635 *Classostrobus crossii* sp. nov. (Cheirolepidiaceae). International Journal of Coal Geology
 636 69, 55–67.

637 Schöner, R., Viereck-Göette, L., Schneider, J., Bomfleur, B., 2007. Triassic-Jurassic sediments
 638 and multiple volcanic events in North Victoria Land, Antarctica: A revised stratigraphic
 639 model. USGS Open-File Reports 2007–1047 (SRP102): 5 pp. [Antarctica: A Keystone in
 640 a Changing World–Online Proceedings of the 10th ISAES].

641 Shang, Y.-K. 1997. Middle Jurassic palynology of Carapace Nunatak, Victoria Land, Antarctica.
 642 Acta Palaeontologica Sinica 36, 170–186.

643 Shen, Y., 1994. Jurassic conchostracans from Carapace Nunatak, southern Victoria Land,
644 Antarctica. *Antarctic Science* 6, 105–113.

645 Surdam, R.C., Eugster, H.P., Mariner, R.H., 1972. Magadi-type chert in Jurassic and Eocene to
646 Pleistocene rocks, Wyoming. *GSA Bulletin* 83, 2261–2266.

647 Stigall, A.L., Babcock, L.E., Briggs, D.E.G., Leslie, S.A. 2008. Taphonomy of lacustrine
648 interbeds in the Kirkpatrick Basalt (Jurassic), Antarctica. *PALAIOS* 23, 344–355.

649 Stockey, R.A., Rothwell, G.W. 2013. *Pararaucaria carrii* sp. nov., anatomically preserved
650 evidence for the conifer family Cheirolepidiaceae in the Northern Hemisphere.
651 *International Journal of Plant Sciences* 174, 445–457.

652 Taylor, T.N., Alvin, K.L., 1984. Ultrastructure and development of Mesozoic pollen:
653 *Classopollis*. *American Journal of Botany* 71, 575–587.

654 Taylor, E.L., Ryberg, P.E., 2007. Tree growth at polar latitudes based on fossil tree ring analysis.
655 *Palaeogeography, Palaeoclimatology, Palaeoecology* 255, 246–264.

656 Taylor, T.N., Taylor, E.L., Krings, M., 2009. *Paleobotany: The Biology and Evolution of Fossil*
657 *Plants*, Second Edition. Academic Press, Amsterdam, The Netherlands, 1230 pp.

658 Tosolini, A.-M.P., McLoughlin, S., Wagstaff, B.E., Cantrill, D.J., Gallagher, S.J., 2015.
659 Cheirolepidiacean foliage and pollen from Cretaceous high-latitudes of southeastern
660 Australia. *Gondwana Research* 27, 960–977.

661 Townrow, J.A., 1967a. Fossil plants from Allan and Carapace Nunataks, and from the upper Mill
662 and Shackleton glaciers, Antarctica. *New Zealand Journal of Geology and Geophysics*
663 10, 456–473.

664 Townrow, J.A., 1967b. On a conifer from the Jurassic of East Antarctica. *Papers and*

665 Proceedings of the Royal Society of Tasmania 101, 137–149.

666 Upchurch, G.R., Doyle, J.A., 1981. Paleoecology of the conifers *Frenelopsis* and
667 *Pseudofrenelopsis* (Cheirolepidiaceae) from the Cretaceous Potomac Group of Maryland
668 and Virginia. In R.C. Romans (ed.), Geobotany. Plenum, New York, pp. 165-172.

669 Vakhrameev, V., 1970. Range and paleoecology of Mesozoic conifers: the Cheirolepidiaceae.
670 Paleontological Journal 4, 12–25.

671 Van Konijnenburg-van Cittert, J.H.A., 1987. New data on *Pagiophyllum maculosum* Kendall and
672 its male cone from the Jurassic of north Yorkshire. Review of Palaeobotany and
673 Palynology 51, 95–105.

674 Venkatachala, B., Góczán, F., 1964. The spore-pollen flora of the Hungarian 'Kössen Facies'.
675 Acta Geologica Hungarica 8, 203–228.

676 Villar de Seoane, L., 2014. Estudio morfológico del género *Classopollis* (Pflug) Pocock &
677 Jansonius (Cretácico, Argentina). Revista Brasileira de Paleontologia 17, 91–104.

678 Watson, J., 1988. The Cheirolepidiaceae. In: Beck, C.B. (ed.), Origin and Evolution of
679 Gymnosperms, Columbia University Press, NY, pp. 382–447.

680 Yang, X.-J., 2008. A male cone of *Pseudofrenelopsis dalatzensis* with *in situ* pollen grains from
681 the Lower Cretaceous of northeast China. Géobios 41, 689–698.

682 Yao, X., Taylor, T.N., Taylor, E.L., 1991. Silicified dipterid ferns from the Jurassic of
683 Antarctica. Review of Palaeobotany and Palynology 67, 353–362.

684 Zavialova, N.E., Tekleva, M.V., Smirnova, S.B., Mroueh, M., 2010a. Exine ultrastructure in
685 pollen grains of *Classopollis* Pflug from the Cretaceous of Lebanon. Paleontological
686 Journal 44, 1353–1367.

687 Zavialova, N., Buratti, N., Roghi G., 2010b. The ultrastructure of some Rhaetian Circumpolles
688 from southern England. *Grana* 49, 281–299.

689 Zhou, Z., 1983. A heterophyllous cheirolepidiaceous conifer from the Cretaceous of East China.
690 *Palaeontology* 26, 789–811.

691

692 **Figure Captions**

693 **Figure 1:** Maps and stratigraphy of the study area.

694 **A.** General map of Antarctica showing the study area in the Ross Sea region; red box
695 highlighting the study site at Carapace Nunatak.

696 **B.** Detailed map of red box area from A. Composite satellite image of the Ross Sea region,
697 included in the Taylor Glacier USGS topographic map, highlighting the study site at Carapace
698 Nunatak.

699 **C.** Geologic map of Carapace Nunatak, modified from Bradshaw (1987); samples were collected
700 from the ice-cored moraine on the E-NE side of Carapace Nunatak.

701 **D.** Generalized stratigraphic column of the Mawson Formation and Kirkpatrick Basalt at
702 Carapace Nunatak; fossiliferous chert and limestone occur as lenses within the basalt and
703 hyaloclastite that overlie the Carapace Sandstone Member of the Mawson Formation.

704 **E.** Rock sample of chert displaying soft sediment deformation and normal faults. Scale = 1 cm.

705

706 **Figure 2:** Suggested reconstruction of *Classostrobus elliotii*. The left side is an external view,
707 and the right side displays internal details such as the position of the microsporangia. Illustration
708 by Sara L. Taliaferro.

709

710 **Plate I:** Components of *Classostrobus elliotii* pollen cone from permineralizations, Early
711 Jurassic of Antarctica.

712 Fig. 1. Apical portion of a longitudinal section of pollen cone. Note helical arrangement of
713 microsporophylls (S = microsporophyll, C = cone axis). Scale = 1.0 mm. Slide no. 29,833.

714 Fig. 2. Transverse section of pollen cone about halfway through specimen. Note masses of pollen
715 grains (amber in color) within microsporangia, and annular arrangement of pollen sacs. Image
716 taken of a peel still attached to the surface of the specimen (S = microsporophyll, L = associated
717 leaf). Scale = 2.0 mm. 17654 F3 bot peel no. 35. Slide no. 29,944.

718 Fig. 3. Annular cluster of 5 microsporangia with presumed microsporophyll stalk in the center
719 (arrow), recognized by the presence of tracheids (see enlargement in Fig. 8). Scale = 0.3 mm.
720 17613 F top alpha slide no. 22.

721 Fig. 4. Transverse section of a microsporophyll with rhomboidal head at the top left. Note
722 epidermis (arrow) and opaque parenchymatous cells; cells below these are transfusion tissue (T).
723 Scale = 0.5 mm. Slide no. 29,912.

724 Fig. 5. Dermal layers of microsporophyll (E = epidermis, H = hypodermis). Scale = 25 μ m. Slide
725 no. 29,912.

726 Fig. 6. Single vascular trace from approximately the middle of microsporophyll stalk, showing
727 metaxylem tracheids with scalariform thickenings and possible protoxylem tracheids with
728 helical-annual thickenings (arrow). Scale = 50 μm . Slide no. 29,915.

729 Fig. 7. Transfusion tracheids of the microsporophyll. Note scalariform pitting (arrow). Scale = 50
730 μm . Slide no. 29,913.

731 Fig. 8. Higher magnification of central area indicated by arrow in Fig. 3, showing part of a
732 tracheid with scalariform thickenings (arrow). Scale = 30 μm . Slide no. 29,822.

733 Fig. 9. Uniseriate wall of pollen sac showing sporangial walls lacking opaque contents. Scale =
734 50 μm . slide no. 29,813.

735 Fig. 10. Detail of vascular bundle of one of the associated *Brachyphyllum*-type leaves.
736 Surrounding the vascular tissue are round cells (indicated by *) of unknown affinities that may
737 represent some type of secretory cell, form of transfusion tissue, or some type of thick-walled
738 fungal hyphae. Scale = 50 μm . Slide no. 29,812.

739

740 **Plate II:** Pollen of *Classostrobilus elliotii*. All SEM images from SEM stub no. J 2138.

741 Fig. 1. Light micrograph (LM) of numerous pollen grains within a microsporangium. Scale = 50
742 μm . Slide no. 29,940.

743 Fig. 2. LM of pollen grains still within a tetrad. Scale = 10 μm . Slide no. 29,937.

744 Fig. 3. Scanning electron micrograph (SEM) of distal view of pollen grain. Note cryptopore (C)
745 and furrow (F). Scale = 5 μm .

746 Fig. 4. LM of side view displaying distal cryptopore (arrow). Scale = 10 μ m. Slide no. 29,937.

747 Fig. 5. SEM of distal view of pollen grain. Sexine (S) is partially separated revealing preserved
748 nexine (N) layer displaying the distal cryptopore (C). Note perforations on nexine from contact
749 with columellae of the sexine. Scale = 10 μ m.

750 Fig. 6. LM of side view of pollen grain showing rimula (arrow) and striae in equatorial region.
751 Scale = 10 μ m. Slide no. 29,937.

752 Fig. 7. SEM of internal view of pollen grain showing rimula (arrow) and equatorial striae. Scale
753 = 10 μ m.

754 Fig. 8. LM of proximal view of trilete. Orbicules are indicated at arrow. Scale = 10 μ m. Slide no.
755 29,939.

756 Fig. 9. SEM of internal view of pollen grain showing trilete mark. Note linear striation patterns
757 (arrow) of sexine columellae. Scale = 10 μ m.

758 Fig. 10. Section of sexine layer displaying internal columellae on the left and external protruding
759 spinules on the right. Scale = 2 μ m.

760 Fig. 11. Internal view of smooth surface of nexine layer. Scale = 2 μ m.

761 Fig. 12. Orbicules with spinules. Scale = 2 μ m.

Figure 2