Temperature sensitivity of organic matter decomposition of permafrost-region soils during laboratory incubations

Rosvel Bracho^{1, 2 *}

¹. Department of Biology, University of Florida, Gainesville, FL 32611, USA.

². School of Forest Resources and Conservation, University of Florida, Gainesville, FL 32611, USA.

Susan Natali³

³Woods Hole Research Center, Falmouth, MA

Elaine Pegoraro^{1, 4}

¹. Department of Biology, University of Florida, Gainesville, FL 32611, USA.

⁴. Center for Ecosystem Science and Society, Northern Arizona University, Flagstaff, AZ

Kathryn G. Crummer^{1, 2}

¹. Department of Biology, University of Florida, Gainesville, FL 32611, USA.

². School of Forest Resources and Conservation, University of Florida, Gainesville, FL 32611, USA.

Christina Schädel^{1, 4}

¹. Department of Biology, University of Florida, Gainesville, FL 32611, USA.

⁴. Center for Ecosystem Science and Society, Northern Arizona University, Flagstaff, AZ

Gerardo Celis¹,

¹. Department of Biology, University of Florida, Gainesville, FL 32611, USA.

Lauren Hale⁵,

⁵. Institute for Environmental Genomics, University of Oklahoma, Norman, OK.

Liyu Wu^{5, 6},

⁵. Institute for Environmental Genomics, University of Oklahoma, Norman, OK.

⁶. Department of Botany and Microbiology, University of Oklahoma, Norman, OK.

Huaqun Yin^{5, 6}

⁵. Institute for Environmental Genomics, University of Oklahoma, Norman, OK.

⁶. Department of Botany and Microbiology, University of Oklahoma, Norman, OK.

James M. Tiedje⁷,

⁷. Center for Microbial Ecology, Michigan State University, East Lansing, MI.

Konstantinos T. Konstantinidis⁸

⁸. School of Civil and Environmental Engineering and School of Biology, Georgia Institute of Technology, Atlanta, GA.

Yiqi Luo⁶,

⁶. Department of Botany and Microbiology, University of Oklahoma, Norman, OK.

Jizhong Zhou^{5, 6, 9, 10}

⁵. Institute for Environmental Genomics, University of Oklahoma, Norman, OK.

⁶. Department of Botany and Microbiology, University of Oklahoma, Norman, OK.

⁹. State Key Joint Laboratory of Environment Simulation and Pollution Control, School of Environment, Tsinghua University, Beijing 100084, China.

¹⁰. Earth Science Division, Lawrence Berkeley National Laboratory, Berkeley, CA, 94270, USA.

Edward. A. G. Schuur^{1, 4}

¹. Department of Biology, University of Florida, Gainesville, FL 32611, USA.

⁴. Center for Ecosystem Science and Society, Northern Arizona University, Flagstaff, AZ

*Corresponding author: rbracho@ufl.edu. Current address School of Forest Resources and Conservation, University of Florida, Gainesville, FL 32611, USA.

Keywords

Temperature sensitivity (Q_{10})
Tundra
Organic matter decomposition
Carbon pools
GeoChip

Abstract

Permafrost soils contain more than 1300 Pg of carbon (C), twice the amount of C in the atmosphere. Temperatures in higher latitudes are increasing, inducing permafrost thaw and subsequent microbial decomposition of previously frozen C, which will most likely feed back to climate warming through release of the greenhouse gases CO₂ and CH₄. Understanding the temperature sensitivity (Q₁₀) and dynamics of soil organic matter (SOM) decomposition under warming is essential to predict the future state of the climate system. Alaskan tundra soils from the discontinuous permafrost zone were exposed to *in situ* experimental warming for two consecutive winters, increasing soil temperature by 2.3°C down to 40 cm in the soil profile. Soils obtained at three depths (0–15, 15–25 and 45–55 cm) from the experimental warming site were incubated under aerobic conditions at 15°C and 25°C over 365 days in the laboratory. Carbon fluxes were measured periodically and dynamics of SOM decomposition, C pool sizes, and decay rates were estimated. Q₁₀ was estimated using both a short-term temperature manipulation (Q_{10-ST}) performed at 14, 100 and 280 days of incubation and via the equal C method (Q_{10-EC}, ratio of time taken for a soil to respire a given amount of C), calculated continuously. At the same time points, functional diversities of the soil microbial communities were monitored for all incubation samples using a microbial functional gene array, GeoChip 5.0. Each array contains over 80,000 probes targeting microbial functional genes involved in biogeochemical cycling of major nutrients, remediation strategies, pathogenicity and other important environmental functions. Of these, over 20,000 probes target genes involved in the degradation of varying C substrates and can be used to quantify the relative gene abundances and functional gene diversities related to soil organic matter turnover. The slow decomposing C pool (C_s), which represented close to 95% of total C in the top 25 cm soils, had a higher Q₁₀ than the

fast decomposing C pool (C_F) and also dominated the total amount of C released by the end of the incubation. Overall, C_S had temperature sensitivities of $Q_{10-ST} = 2.55 \pm 0.03$ and $Q_{10-EC} = 2.19 \pm 0.13$, while the C_F had a temperature sensitivity of $Q_{10-EC} = 1.16 \pm 0.30$. In contrast to the 15°C incubations, the 25°C microbial communities showed reduced diversities of C-degradation functional genes in the early stage of the incubations. However, as the incubations continued the 25°C communities more closely paralleled the 15°C communities with respect to the detection of microbial genes utilized in the degradation of labile to recalcitrant C substrates. Two winter seasons of experimental warming did not affect the dynamics and temperature sensitivity of SOM decomposition or the microbial C-degradation genes during incubation. However, under the projected sustained warming attributable to climate change, we might expect increased contribution of C_S to organic matter decomposition. Because of the higher Q_{10} and the large pool size of C_S , increased soil organic matter release under warmer temperatures will contribute towards accelerating climate change.

1. Introduction

Permafrost zone soils contain approximately 1330-1580 Pg of organic C, which is twice the amount of atmospheric C (Schuur et al., 2008; Tarnocai et al., 2009; Hugelius et al., 2014; Schuur et al., 2015). Though they cover less than 15% of global soil area, permafrost zone soils store about one-third of total global soil C to 3m depth (Schuur et al., 2015). Temperatures in high latitude regions are increasing faster than in the rest of the world (Hassol, 2004; Fyfe et al., 2013) and future climate projections indicate a potential increase between 7 – 8 °C by the end of the 21st century (Trenberth et al., 2007, IPCC 2013). Sustained warming thaws permafrost

(Romanovsky et al., 2010; Smith et al., 2010; Koven et al., 2013), leading to a thicker seasonal active layer that exposes a large pool of previously frozen organic C to microbial decomposition (Harden et al., 2012). The release of CO₂ and CH₄ from this newly thawed C by increased microbial activity could add significant quantities of C to the atmosphere. Recent efforts to model permafrost C in response to warming project a shift from a C sink to a source in the arctic and sub-arctic regions by the end of the 21st century, leading to a positive feedback to a warming climate (Koven et al., 2011; Schaefer et al., 2011; Schaefer et al., 2014)

Soil organic matter is composed of a continuum of C compounds. For simplicity, it is often conceptualized as fast, slow, or passively decomposing C pools (Trumbore, 1997; Amundson, 2001; Schädel et al., 2014). In permafrost zone soils, the fast C pool, with turnover times of a few days to weeks at laboratory temperatures (Schädel et al., 2014), represents less than 10% of total soil C, while the majority belongs to the slow C pool, with turnover times from years to decades (Knoblauch et al., 2013; Schädel et al., 2014). Since slow C is a large proportion of total soil C and has a long residence time, it will dominate the long term response of permafrost soil C decomposition to warming (Schuur et al., 2007; Sistla et al., 2013; Sistla et al., 2014).

Soil organic matter decomposition in arctic and subarctic ecosystems undergoing permafrost thaw is controlled by a complex of biophysical interactions including soil temperature, soil moisture, physical and chemical protection, C quality, changes in microbial biomass and microbial communities, and the dominant plant community composition (Hirsch et al., 2002; Wickland and Neff, 2008; Karhu et al., 2010; Waldrop et al., 2010; O'Donnell et al., 2011; Schmidt et al., 2011; Hugelius et al., 2012; Sistla et al., 2013; Sistla et al., 2014). Changes in the biophysical factors that control SOM decomposition will likely be reflected most rapidly

in the abundance and structure of the microbial communities as they adapt to their new physical and chemical environment (Deslippe et al., 2011; Rinnan et al., 2011; Sistla et al., 2013; Sistla et al., 2014). Functional adaptations of microbial communities to seasonal environmental changes have been documented in alpine and arctic tundra soils; winter microbial biomass is fungus dominated, while growing season microbial biomass is bacteria dominated (Schadt et al., 2003; Wallenstein et al., 2007; Buckeridge et al., 2013). New dominant microbial communities and the C substrates they decompose may have different temperature sensitivities, and small changes could have a significant effect on the C balance of permafrost soils (Davidson and Janssens, 2006; Fan et al., 2008).

Field warming experiments often show an initial burst of respiration after the application of the warming manipulation associated with consumption of the fast C pool, followed by a decline in C release rates as slow C increasingly dominates respiration (Kirschbaum, 1995; Melillo et al., 2002; Kirschbaum, 2004; Eliasson et al., 2005; Knorr et al., 2005; Hartley et al., 2007; Hartley et al., 2009; Streit et al., 2014). In these field warming experiments, microbes can acclimate to warming by adjusting their metabolism to the new temperature regime, thus reducing their respiration rate at a given temperature and improving their carbon use efficiency (CUE) (Luo et al., 2001; Barcenas-Moreno et al., 2009). Microbial communities may also shift in composition as a new C balance is established perhaps reflecting the changes in the environment as well as in C availability. Experimental field warming in arctic and low arctic ecosystems have been shown to shift microbial communities toward dominance of fungi over bacteria leading towards increased use of more recalcitrant slow decomposing C and change in the plant-microbial associations that accompany shifts in plant community composition and

productivity (Deslippe et al., 2011; Deslippe and Simard, 2011; Natali et al., 2011; Deslippe et al., 2012; Natali et al., 2012; Sistla et al., 2013; Sistla et al., 2014).

Accurate measurements of the turnover rates and temperature sensitivity of fast and slow C decomposition are difficult in field conditions. Field studies measure only the apparent temperature sensitivity because of environmental constraints and the confounding effects of different C pools' contribution to total respiration (Davidson and Janssens, 2006; Kirschbaum, 2013). Laboratory soil incubations are a valuable means for estimating the long-term potential for C release from thawing permafrost soils because environmental constraints over SOM decomposition can be carefully controlled (Holland et al., 2000; Reichstein et al., 2000; Dutta et al., 2006). Continuous C loss measurements from long-term incubation experiments provide information on potential C release, kinetics of soil organic matter decomposition, separation of different C pools comprising the SOM, their decay rates and temperature sensitivities, and the composition and abundance of associated microbial populations (Dutta et al., 2006; Karhu et al., 2010; Lavoie et al., 2011).

While many connections between microbial community compositions and ecosystem functions have been made, it has been recently suggested that ecosystem processes may be more dependent on the abundance and diversity of related functional genes rather than the phylogenetic structures of communities (Philippot et al., 2013; Paula et al., 2014). The utilization of the GeoChip functional gene array provides a platform in which a variety of important functional genes, including critical genes for C-turnover, can be detected even when present at low abundances. The sensitivity of the GeoChip arrays was reported previously over half of probes producing positive signal with 10 or fewer pg of DNA template (He et al., 2007). It is also specific as probe design accounts for minimal-to-no cross-hybridization of sequences with

less than 90% similarity and was designated a quantitative tool in which R^2 values averaged 0.93 for signal intensity and DNA concentration correlations (Liebich et al., 2006; Wu et al., 2006; Zhou et al., 2010; Zhou et al., 2012). These arrays allowed for comparison of samples throughout the incubation process with respect to microbial functional diversity, which is the variety of functional genes detected in each sample. We used all functional genes covered on the arrays to broadly compare communities. For analyses centered on traits involved in tundra C degradation we focused on 20,000 probes targeting hundreds of bacterial and fungal genes involved in the degradation of various C substrates.

The main objectives of this incubation study of permafrost zone soils were: (1) to assess the effects of *in situ* experimental soil warming on initial soil C pools, and to estimate C pool sizes and their relative turnover rates, (2) to characterize the relative abundance and structure of microbial functional genes associated with fast and slow C pool decomposition, as well as the response to field warming, (3) to determine the temperature sensitivity of SOM microbial decomposition (Q_{10}) for the fast and the slow C pools, and to characterize how Q_{10} responds to field warming. We expected warming-induced changes in microbial community functional gene abundances and diversities to accompany changes in the kinetics of SOM decomposition. On this matter, we hypothesized that both the abundances and diversities of genes involved in the degradation of fast decomposing C would decrease as these C pools were depleted over the course of the incubation study, whereas genes involved in slow decomposing C would show a relative increase. We also expected that there could be differential temperature sensitivities, with a higher Q_{10} for the slow C pool compared to the fast C pool.

2. Materials and methods

2.1. Site description and experimental design

The Carbon in Permafrost Experimental Heating Research project (CiPEHR) was established in September 2008 (Natali et al., 2012; Natali et al., 2014) on moist acidic subarctic tundra within the Eight Mile Lake research watershed (63°52'59''N, 149°13'32''W) (Schuur et al., 2007; Schuur et al., 2009). Vascular plant cover is dominated by the tussock-forming sedge *Eriophorum vaginatum* and the deciduous shrubs *Betula nana* and *Vaccinium uliginosum*, non-vascular biomass is dominated by mosses and lichens (Natali et al., 2012). Soils are Gelisols (Soil Survey Staff, 1999) with a thick organic horizon (0.45 – 0.65 m) on top of cryoturbated mineral soil and a C content between 50 and 70 kg m⁻² down to 1 m depth (Pries et al., 2012). Long-term mean annual temperature is -1.0°C; mean growing season (May-September) air temperature is 11.2°C and monthly averages range from +15 °C in July to -16°C in December (Schuur et al., 2009). Average annual precipitation is 378 mm with a growing season mean of 245 mm (National Climatic data center, NOAA). The soil warming treatment was applied by increasing snow cover behind snow fences during winter time, coupled with early spring snow removal to keep water input and snow melt timing similar to control plots (Natali et al., 2011). As a result of this treatment, soil warming plots reached an average of -3.9°C in the soil profile (5-40 cm depth) during the winter time, 2.3°C warmer than control plots, this warming effect continued into the growing season when temperature at 40 cm reached 2.9°C in warmed soils and 1.6°C in control soils,

2.2. Soil sampling and preparation

Soil cores were collected frozen from control (n=6) and field warmed (n=6) plots in May 2010 after exposure to two consecutive winter warming seasons (winters of 2008-2009 and 2009-2010) and one full summer growing season (2009), and kept frozen until further processing. Soil cores were separated in the lab into three depth increments for incubation: 0-15 cm, a fibric Oi-Oe horizon, with a lot of litter and coarse roots, 15-25 cm, Oe-Oa horizon with less coarse roots and more decomposed soil, and 45-55 cm, soils with a mineral-organic mix located just above the surface permafrost because the surface soil thaws each summer to about 55-65 cm and refreezes in the winter. Each soil layer increment was split longitudinally into four sections: one for each lab incubation temperature (15°C and 25°C), one for microbial analysis, and one for soil physical characterization. Soil moisture content, bulk density, and mass-based nitrogen (N) and C content (ECS 4010, Costech Analytical, Valencia, CA) were measured for each soil increment (Table 1).

2.3. Incubation design and soil carbon fluxes

Each incubation sample was further divided into 8 subsamples of the same wet weight (~10g). Each subsample was kept largely intact and placed into a perforated foil cup over a bed of 3 mm glass beads to allow drainage and maintain the soils at field capacity soil moisture. Each incubation sample thus consisted of eight 30 cm³ vials inside a 1 L mason jar. Each jar was covered with a perforated lid to allow air exchange while minimizing contamination. Control and field warmed soils were aerobically incubated at each of two laboratory incubation temperatures (15°C and 25°C) for 365 days. Optimal temperature for field microbial growth is well above the mean annual air temperature and can be considered as the average highest temperature during the year (Rousk and Baath, 2011), except for when Q₁₀ was being assessed. Soil temperature at the research site can reach up to 25°C in the surface few centimeters of soils exposed to direct

sunlight during the growing season; laboratory temperatures were chosen to span the soil temperatures found in the soil profile while keeping them warm enough to observe changing dynamics through time.

Jars were placed in a water bath set to the incubation temperature and connected to an automatic soil incubation system (ASIS). Air from each sequential jar was circulated by a pump through an infrared gas analyzer (IRGA, Li-820 Licor, Lincoln, Nebraska) at 0.9 L min^{-1} . A mass flow controller (Mass Flow meter GFM, Aalborg Instruments & Control) maintained a constant flow, while CO_2 concentration and pressure inside each jar were measured by the IRGA and recorded every three seconds by a datalogger (CR1000, Campbell Scientific, Logan UT) over 8 minutes. Jars were flushed with CO_2 -free air at any point when CO_2 concentration in the jar head space reached 10,000 ppm in order to avoid suppression of microbial activity due to oxygen deprivation. Carbon fluxes (F_c) were calculated as the rate of CO_2 increase in the headspace of the jars over time after at least four cycles of eight and an half hours each, and expressed in micrograms of C per gram of initial soil C per day ($\mu\text{gC gC}_{\text{initial}}^{-1} \text{ d}^{-1}$). Fluxes were measured every 48 hours during the first two weeks of incubation in order to capture the initial phase of SOM decomposition, then twice a week up to 45 days of incubation, biweekly up to 180 days, and at least once per month up to one year of total incubation time. Total amount of C respired (C_R) at any measurement point was estimated by linear interpolation between measurement points.

Total C respired (C_R , $\text{mgC gC}_{\text{initial}}^{-1}$) through time (t) up to 365 days was fitted with a two-pool C model (Andren and Paustian, 1987) (Proc NLIN, SAS 9.3. SAS Institute Inc. North Carolina):

$$C_R(t) = C_F(1 - e^{-k_F t}) + C_S(1 - e^{-k_S t}) \quad (1)$$

where C_F and C_S are the sizes of the fast and slow decomposing C pools relative to the initial C content ($\text{mgC gC}_{\text{initial}}^{-1}$), and k_F , and k_S are the first order kinetic decomposition rate constants for the fast and slow decomposing C pools (day^{-1}). This model assumes that C_F and C_S sum to 1000 mg C per gram of initial C and $k_F > 0$ and $k_S > 0$.

2.4. Temperature sensitivity of soil organic matter decomposition (Q_{10}).

In order to determine if and how temperature sensitivity of SOM decomposition was affected by field warming, we calculated the change in decomposition rate in response to an increase in temperature of 10°C (Q_{10}) using two different methods. Short-term Q_{10} ($Q_{10\text{-ST}}$) was calculated by measuring respiration while simultaneously exposing soils to the range of 5 to 30°C via 5°C steps over the total period one week, ~ 1 day at each temperature level (Hamdi et al., 2013). Short-term Q_{10} was calculated by exposing soils to this temperature range at 14, 100 and 280 days following initiation of the incubation (DOI). We expected the contribution of C_F to dominate the total C flux at the beginning of incubation and because of this, the $Q_{10\text{-ST}}$ at 14 DOI would reflect primarily the temperature sensitivity of C_F . Similarly, more than 95% of C_F was expected to be consumed by 280 DOI (Schädel et al., 2014) thus the $Q_{10\text{-ST}}$ at 280 DOI in contrast was expected to reflect primarily the temperature sensitivity of C_S .

Measured C fluxes were fitted to an exponential function to describe the short-term temperature sensitivity of microbial decomposition ($Q_{10\text{-ST}}$) (Proc NLIN, SAS 9.3. SAS Institute Inc. North Carolina):

$$F_C = ae^{bT} \quad (2)$$

Where a and b are estimated parameters; a is the basal C flux at 0°C, T is the incubation temperature and b is the parameter related to the sensitivity of microbial respiration, where:

$$Q_{10-ST} = e^{(10b)} \quad (3)$$

Temperature sensitivity of SOM decomposition was also calculated using a second methodology: the equal C method (Q_{10-EC}) (Rey and Jarvis, 2006; Conant et al., 2008a). Q_{10-EC} was calculated as the ratio of time taken for a soil to respire a given amount of C. For example, comparing rates at steps of 1 mg (or 0.1 % of total C) at each of the two constant incubation temperatures (15°C and 25°C) throughout the incubation experiment. The Q_{10-EC} method allowed us to track the trend in Q_{10} continuously as organic matter was decomposed. This method assumes that a given amount of respired C comes from the same C pool across different temperatures. At the early stages of incubation, when only a small amount of C has been respired (i.e. the first 0.5 – 1% of initial C), whole soil Q_{10} is often used as a proxy for the temperature sensitivity of the fast C pool (Rey and Jarvis, 2006; Conant et al., 2008b; Haddix et al., 2011; Zhu and Cheng, 2011). However, contributions of the slow C pool to total C flux can also be significant at these early stages of the incubation (Schädel et al., 2014; Liang et al., 2015), potentially confounding the estimated Q_{10} for the fast pool alone. To avoid this bias, we used equation 1 to separate the slow and fast C pools, then tracked total C respired from each C pool over time, and estimated Q_{10-EC} for each C pool separately (Conant et al., 2010). Since soils below 15 cm had very low C_F , no Q_{10-EC} was calculated.

2.5. Microbial analysis

One vial containing a soil subsample from each depth was removed from each jar after 14, 90, and 280 days of incubation for microbial analysis. A PowerSoil® DNA isolation kit was

used to extract microbial community DNA from soil subsamples following standard procedures (MoBio Laboratories, Inc, Carlsbad, California). In some samples, DNA of high purity (Nanodrop 260/280 and 260/230 absorbance ratios above 1.70) could not be obtained via the kit alone so a freeze-grind method was used to obtain raw DNA (Zhou et al., 1996), which was subsequently purified with the PowerSoil® kit. We assessed the microbial functional gene structure using GeoChip 5.0, which contains over 80,000 probes targeting microbial environmental functional genes (Wu et al., 2006; Zhou et al., 2012; Xue et al., 2013; Tu et al., 2014; Zhou et al., 2014). For this, 500 ng of soil community DNA was labeled with the fluorescent dye Cy-3, hybridized to GeoChip 5.0 60K microarrays, and scanned with a NimbleGen MS200 Microarray Scanner using techniques described previously (Cong et al., 2015). The images data were processed using the Agilent Feature Extraction program that designates values for probe signal intensities and background (noise) signal intensities based on the scanned images. Extracted data was then loaded onto an in-house GeoChip data analysis pipeline (ieg.ou.edu/microarray/). Data normalization and quality filtering were performed with multiple standard steps (Liang et al., 2010; Deng and He, 2013). Briefly, poor quality spots were removed, the signals of all spots were transferred into relative abundances, and spots with signal-to-noise (SNR) ratios less than 2 were set to 0 or removed, depending on the analysis. Probes with positive signal in only 2 or fewer samples were removed. Probes with high signal intensities reflect a greater amount of hybridization and targeted genes more abundant in the sample. We analyzed the microarray data in three ways. First, we used microarray data from all probes, which included genes involved in C fixation and degradation, nitrogen and sulfur cycling, antibiotic resistance, and contaminant remediation or degradation. These data represent a broad picture of the microbial functional communities in each sample. Next, we analyzed only probes

targeting genes involved in C-degradation. For this, we focused on the abundances of C degradation genes present in each sample by quantifying the signal intensities of only probes with positive signal. We also reflected the diversity of C-degradation genes in a sample by incorporating the probes that did not have an SNR >2 into the data as having a signal intensity of 0. Thus, samples with gene categories containing more 0's represented conditions wherein there were lower functional diversities of the given gene.

2.6. Statistical analysis

We investigated the effects of field warming and sampling depth on soil properties (%C, %N, C:N, bulk density and pH) using linear mixed effects models in R (R Development Core Team, 2012). The same analysis was applied for parameters (C_F , C_S , k_F , k_S), total C respired and Q_{10-ST} , using field warming, sampling depth, incubation temperature, C:N, bulk density and pH as fixed factors. We used graphical inspection and variance inflation factors (VIF) with the full model to avoid variable collinearity. The cutoff VIF value to drop a variable was 3 (Zuur et al., 2010). After selecting the fixed factors we used the full model with all selected fixed factors to optimize the random structure using the 'lmer' function in the lme4 package (Bates et al., 2014), which was field replicate (fence) and soil core nested within field replicate (fence). Once the random structure was optimized, we performed stepwise multiple regression analysis using the 'lme' command with restricted maximum likelihood from the 'nlme' package in R (Pinheiro et al., 2012). For each model the optimal model was selected by dropping the least significant individual explanatory variable one by one and refitting the model every time with the model selection criterion being the smallest Akaike Information Criterion (AIC) as described in Zuur et

al. (2010). For short-term Q_{10} (Q_{10-ST}), we additionally added two-way interactions after graphical exploration of the data in order to test if the effect of a main factor on the dependent variable was affected by another independent factor. Data were log or arcsine transformed as needed and residuals were checked for normality and heterogeneity of variance. For Q_{10-EC} analysis, a linear mixed effects model was used with field warming and C_R as fixed effects and soil core as random effect. Q_{10-EC} was analyzed for each depth. For 0-15 cm soils Q_{10-EC} was compared at different amounts of C_R (1, 25, 45, 65 and 90 mgC gC_{initial}⁻¹) and at 1, 5 and 10 mgC gC_{initial}⁻¹ for 15-25cm and 45-55 cm soils. Q_{10-EC} for all depths was compared after C_S contributed to 75% or more to C_R .

The GeoChip data was analyzed for the entire array (containing probes targeting genes involved in C, nitrogen, phosphorus, and sulfur cycling as well as bioremediation processes) and for C gene targets alone. Samples collected at 2 weeks, 3 months, and 9 months from both incubation temperatures and all 3 depths were used for all analyses. Dissimilarity tests were based on Bray-Curtis dissimilarity index using non-parametric multivariate analysis of variance (ADONIS) (Anderson, 2001) under the package Vegan (v.2.0-3) (Oksanen et al., 2012)). Detrended correspondence analysis (DCA) was used to determine the overall functional changes in the microbial communities with respect to environmental and temporal variables (Hill and Gauch, 1980). All DCA plots were prepared in R package vegan and contained data from all GeoChip probes include those with SNR<2 filled as 0's. The signal intensities for GeoChip probes in subcategories "carbon degradation" and "organic remediation" were isolated for all analyses focused on traits involved in tundra C degradation. We removed all probes with SNR <2 for all analyses focused on gene abundances and included them as 0's for all analyses focused on C-degradation functional gene diversity. Mann-Whitney-Wilcoxon tests, performed in R,

were used to determine statically significant differences between categorized probes in different treatments and diversity and evenness indices of C-degradation functional genes.

3. Results

3.1. Initial soil properties

In general, initial mass based soil C (%C) and nitrogen (%N) concentrations varied systematically by depth (Table 1), but were not affected by field warming over this initial phase of the tundra warming experiment (Table 2). Carbon concentration decreased from about 40% in the 0-15 cm layer, to 35% in 15-25 cm, and to less than 20% in the 45-55 cm layer (Table 1). Nitrogen concentration also significantly changed with depth, with the highest value in the intermediate layer (15-25 cm). Carbon to nitrogen ratio (C:N) slightly decreased with depth although in general, the lowest values were found in the 15-25 cm depth interval. Bulk density and pH both increased with depth with no difference between control and field warmed soils, although there was a trend towards higher bulk density with warming at the deepest soil depth (Table 1, 2).

3.2. Carbon pools, decay rates and total carbon respired

The fast decomposing C pool (C_F) accounted for less than 5% of the total initial C pool and decreased from 30-50 $\text{mgC gC}_{\text{initial}}^{-1}$ in 0-15 cm soils to 5-10 $\text{mgC gC}_{\text{initial}}^{-1}$ in 45-55 cm (Figure 1a, Supplemental Table 1) whereas the slow decomposing C pool size (C_S) accounted for 950-990 $\text{mgC gC}_{\text{initial}}^{-1}$ (Figure 1b). Both C pools significantly changed with soil depth and incubation temperature (Table 3). Because the sum of both pools is defined as 1000 $\text{mgC gC}_{\text{initial}}^{-1}$

¹, they showed opposite patterns versus depth and incubation temperature: C_F decreased with depth but increased with incubation temperature, while C_S increased with depth and decreased with incubation temperature (Figure 1, Table 3). Carbon pool sizes were not affected by field warming in this initial phase of the experiment. Of all the variables that went into the mixed effects regression model, only C:N showed a significant relationship with both C pool sizes ($p < 0.0001$, Table 3), with higher C:N related to larger C_F and the reverse for C_S .

The decomposition rate of the fast decomposing C pool (k_F) ranged from 0.004-0.1 day⁻¹ (i.e. turnover times of 10-250 days, Fig. 2a, Supplemental Table 1) and was negatively influenced by incubation temperature and bulk density (Table 3). On average, 40% and 60% of C_F from control and field warmed soils was consumed by 14 DOI at 15°C incubation temperature. By day 100, 85% and 95% of C_F was consumed from soils incubated at 15°C and 25°C, respectively. By day 280 of incubation virtually all of C_F was consumed from all soils. The decomposition rate of the slow C pool (k_S , Fig 2b, Supplemental Table 1) ranged from 0.00003 to 0.0012 d⁻¹ (i.e. turnover times of 2.3-97.8 years). The slow pool decomposition rate decreased three-fold from 0-15 cm to 15-25 cm soils, and doubled when incubation temperature was increased from 15 to 25°C (Table 3, Supplemental Table 1); k_S also increased with C:N while pH had a marginal negative effect on k_S .

Total cumulative C respired (C_R) in 0-15 cm soils after 365 days of incubation reached an overall average of 137.27 ± 15 and 232.60 ± 24 mgC gC_{initial}⁻¹ for 15°C and 25°C incubations (Fig. 3, Supplemental Table 1). The total amount of C respired significantly decreased with depth and increased with incubation temperature, but no significant field warming effect was detected (Figure 3, Table 3, Supplemental Figure 1). Linear mixed effects regression results

identified C:N as a significant predictor for total C loss over the one year incubation period (Table 3), with higher initial C:N associated with increased total C loss.

3.3. Temperature sensitivity (Q_{10})

Short-term calculated temperature sensitivity of SOM decomposition (Q_{10-ST}) ranged from 1.6 to 4.8, with an average value of 2.55 ± 0.03 across all soils (Supplemental Table 2). Field warming seems to have an effect on Q_{10-ST} at 14 DOI in 0-15 cm soils (Supplemental Table 2, Figure 4), however, a significant interaction was seen between field warming and depth on Q_{10-ST} (Table 3, Figure 4). The strongest effect of field warming was observed for the 45-55 cm depth interval, where a 20% reduction in Q_{10-ST} was observed in field warmed soils. We expected to observe Q_{10-ST} dominated by C_F at 14 DOI however, time of incubation had only a marginal to no effect on Q_{10-ST} . On average, more than 50% of C_F was already consumed in 0-15 cm soils at both temperatures of incubation by 14 DOI, and C_S already dominated total C flux by that time. This effect was stronger in deeper soils, where C_S dominated respiration even at the very beginning of the incubation.

Temperature sensitivity of SOM decomposition estimated by the equal C method (Q_{10-EC}) in 0-15 cm soils showed a significant increase ($p = 0.0097$) from 1.18 ± 0.37 to 2.40 ± 0.37 as the total amount of C respired (C_R) changed from 1 to 90 $\text{mgC gC}_{\text{initial}}^{-1}$ (Figure 5a, d), no significant field warming effect was detected. Contribution of C_S to the first mg of C_R in 0-15 cm soils was in average 24.6% at both temperatures of incubation, and 87% and 97% at 60 and 90 mg of C_R , respectively. In soils below 15 cm (Figure 5b, c; e, f), Q_{10-EC} was marginally affected by C_R ($p < 0.1 > 0.05$), contribution of C_S to the first mg of C respired was in average 50% and 70% for 15-25 and 45-55 cm soils, respectively. However, Q_{10-EC} was higher ($p < 0.01$) for

field warmed than control at both 15-25 cm (3.09 ± 0.58 vs. 2.07 ± 0.54) and 45 – 55 cm (2.93 ± 0.18 vs. 1.98 ± 0.19 vs.) soils, respectively (Figure 5b, e; c, f). Q_{10-EC} after C_S contribution to C_R was greater than 75% was not affected by depth or field warming ($p>0.05$) showing an overall average of 2.66 ± 0.40 .

Temperature sensitivity for each C pool independently was estimated using equation 1 (Supplemental Table 2). Q_{10-EC} for C_F was estimated for the 0-15cm soils as the ratio of time to respire the first 10 mg of C_F . Surface soils' Q_{10-EC} for C_F (1.16 ± 0.30) was not significantly affected by field warming ($p = 0.28$), but was significantly less than Q_{10-EC} for C_S at the same depth (2.05 ± 0.23). In general, estimated Q_{10-EC} for C_S was not significantly affected by field warming ($p=0.21$) or by depth ($p=0.13$), with an overall average of 2.19 ± 0.13 .

3.4. Total microbial functional traits analysis

Dissimilarity analysis for overall microbial functional traits, as identified by GeoChip 5.0, revealed that functional diversity of microbial communities in incubated soils was significantly altered (Table 4, $p = 0.001$) with soil depth, incubation time and temperature of incubation. While there was no overall field warming effect on functional diversity (Table 4, $p > 0.1$), there was a marginal significant interaction between field warming and incubation time ($p=0.08$). This showed that dissimilarities in functional diversity with respect to field warming tended to increase as the incubation progressed and happened in combination with the depletion of C_F from the soil.

Detrended correspondence analysis (DCA) of GeoChip data indicated that the structure of the microbial functional communities in specific soil samples were separated with respect to the variables depth, time and temperature of incubation. For example, throughout the incubation,

distinct microbial communities dominated at 14, 100 and 280 days in 15 – 25 cm soils incubated at 25°C, with a more clear separation in field warmed than control soils (Figure 6). Functional communities in the top (0-15 cm) differed from those in the bottom (45- 55 cm) in field warmed soils (Supplemental Figure 2). Also, temperature of incubation selected different microbial communities in 15 – 25 cm in both control and field warmed soils (Supplemental Figure 3). To some extent, field warming tended to select different microbial communities in the soil profile in samples incubated at 25°C for 280 days although individual outliers samples are present at both depths (Supplemental Figure 4).

3.5. Microbial traits involved in tundra carbon degradation

Focusing on genes involved in C-degradation, we did not see significant differences in the gene abundances or diversities in field warmed or control soils. However, after two weeks of incubation soils incubated at 25°C had lower mean signal intensity values for the overall set of genes involved in C-degradation that were explored here as compared to the 15°C incubations (Figure 7). This was largely caused by higher numbers of absent (zero) signals for genes involved in C-degradation in the 25°C incubations (Figure 8). This trend was consistent for probes targeting genes involved in the degradation of a wide range of C substrates and was strongest in the surface soils and less so at depth. However, by 3 and 9 months the microbial communities from soils incubated at 15°C and 25°C showed similar mean and median signal intensities for C-degradation genes. When only probes with positive signal were considered (0's excluded), the difference in gene abundances between the 15°C and 25°C 2-week samples was no longer apparent (Figure 9). Furthermore, microbial communities assayed from the 2 weeks, 25°C incubations, which corresponded to the highest measured daily C fluxes, also displayed the lowest alpha diversity estimations for functional genes as determined by Shannon and Simpson

indices and Pielou and Simpson evenness (Supplemental Table 3). These direct and quantitative measures show that the assayed genes at the two weeks of incubation point were not equivalently abundant in the samples at the different incubation temperatures. This indicates that the 10°C temperature difference used in the incubation had a short-term, but highly significant effect on the functional diversity of microbial C degradation genes.

4. Discussion

4.1. Dynamics of soil organic matter decomposition

Decomposition of SOM in tundra from permafrost zone soils incubated under controlled conditions was affected by differences in substrate quality (%C, %N) and C availability (Total C) in the soil profile, as well as incubation temperature and time of incubation. These differences were reflected in microbial community diversity, decay rates, and in total C respired at the end of the incubation experiment. Slow decomposing C contributed between 75% and 90% to total C respired by the end of the incubation term, even in 0-15 cm soils where fast decomposing C was relatively higher compared to deeper in the soil profile, stressing the importance of C_s in long-term C release from these soils. The field warming manipulation did not have much effect either on the soil properties or the decomposition dynamics in the laboratory but this was not surprising because the field warming had been recently implemented and so there was not a long period of time for substantial change to the bulk soil pool. Soils were collected before the beginning of the second growing season after the field warming experiment was implemented and changes to the bulk soil might be expected to take multiple years to manifest (Sistla et al., 2013; Sistla et al.,

2014. The one exception to this was a change in bulk density at the deepest layer that was likely a result of decreasing ice/water content as the thaw depth increased in the warming manipulation.

Size and distribution of C_F and C_S in the soil profile of this research site is typical for permafrost zone soils (Knoblauch et al., 2013; Schädel et al., 2014). The largest proportion of C_F was found in 0-15 cm soils is due to fresh litter inputs, microbial biomass, and the presence of roots and root exudates. In tundra soils, more than 80% of root biomass is found in the top 20 cm (Dennis and Johnson, 1970; Dennis, 1977; Jackson et al., 1996; van Wijk et al., 2003). Root exudates comprise low molecular weight C compounds that can be easily assimilated by microbes. As well, microbial biomass turnover due to winter soil freezing likely contributes additional substrate to C_F (Schimel and Mikan 2005). However the contribution of C_S was still higher than contribution of C_F to total C respired, even in 0-15 cm soils at the end of the incubation, as C_F was depleted rapidly by new microbial activity in the incubation.

Although C_F and C_S have complementary patterns with depth relative to initial C content, both decreased on dry soil basis as total soil C decreased with depth. These depth patterns in C were accompanied by changes in functional microbial diversity (Conant et al., 2011; Sistla et al., 2013), affecting mostly decomposition rate for C_S . A higher k_S at the surface than at depth is likely a function of the fact that these soils accrete vertically over time such that C_S in the deeper layers has already been through significant previous decomposition and remaining SOM has slower turnover, reflected in the significant relationship with C:N. At all depths, k_S was higher in 25°C relative to 15 °C incubation reflecting standard temperature effects on microbial metabolism.

The two fold increase in k_S with an increase of 10°C of incubation was driven by increased activation energy and enzymatic activity, eliminating physical protection and breaking

down complex organic C into more simple molecules (Marschner and Bredow, 2002; Davidson and Janssens, 2006; Allison et al., 2010). As result, more C was transferred to C_F and C_R doubled at 25°C incubation. In contrast to k_S , k_F was not sensitive to depth nor C:N and was seemingly less sensitive to incubation temperature. This suggests that C respired from this fast decomposing pool have a similar molecular composition through the soil profile (Mueller et al., 2015), and perhaps that temperature sensitivity of this material was less. It is important to recognize though, that estimates for pool sizes and turnover rates are not completely independent using these modeling procedure (Katterer et al., 1998).

There did not appear to be variation in specific functional genes along with the phases of C degradation measured during these incubations, which was contrary to our original hypothesis. In fact, functional groups representing all gene subcategories present on the functional gene array were abundant throughout the incubation, rather than genes associated with labile C degradation declining over time. This likely reflects concurrent C_S and C_F decomposition and the ability of the microbial communities to perform both functions simultaneously at all points during the incubation. It has been shown that microbial biomass and their products, as well as plant detritus, are simultaneously contributing to total microbial respiration, albeit at different rates, as wet arctic tundra surface soils thaw (Schimel and Mikan, 2005). These substrates contrast greatly in C quality and their decomposition should be reflected in varied microbial functional genes involved in their degradation. Additional support of these findings was indicated in a previous study wherein microbial communities from different depths in moist acidic tundra were able to decompose a wide range of C substrates with increasing recalcitrance and this capacity increased with temperature (Ernakovich and Wallenstein, 2015).

4.2. Microbial function during OM decomposition

As a whole, the DCA clustering plots reflect that microbial functional community structure varied with the divergent environmental parameters, but that the incubation conditions resulted in greater dissimilarities than did field treatment or depth. Soils also differed in the state of C quality as a result of these parameters. While there was some separation in microbial functional communities from field warmed and control plot soils they were often not complete, with individual outlier replicates consistently observed through the dataset. In contrast, separation between communities based on the variables time and incubation temperature were notably clearer, in particular when field warmed soils were considered. This greater separation may indicate some initial functional community shifts that were present in the field warming samples, which then became more apparent as C quality declined, although this was not directly tested here.

In contrast to some findings in literature, we did not find a significant difference in C-degradation genes as a product of field warming as determined by GeoChip analyses. Previous studies showed microbial functional diversity to decrease in response to warmer soils (Yang et al., 2014; Yergeau et al., 2012) and other reports quantified increased abundances of functional genes in response to warming (Wang et al., 2012), particularly those involved in labile C-degradation (Zhou et al., 2012). A common trend was a correlation between plant biomass (Yang et al., 2014; Yergeau et al., 2012) and C₄ plants (Zhou et al., 2012) to microbial functional gene abundances and diversities in warming experiments. However, changes in plant biomass or aboveground net primary productivity had not yet been detected in this study at the time when soils were sampled (Natali et al., 2012; Salmon et al 2016). Consequently, changes in soil

microbial communities would more likely be tied to environmental changes imposed by the experiment rather than in response to changing plant processes.

The monitoring of the SOM decomposition kinetics provided us a unique opportunity to analyze the corresponding microbial communities. One interesting finding was that after two weeks of incubation, communities with higher respiration rates had lower functional diversity than the communities present in cases of both 1) greater abundance of C_F and lower incubation temperatures and 2) lower abundance of C_F and high incubation temperatures. Thus the temperature-stimulated communities may have had some key degraders that initially dominated the C degradation activity as opposed to an equivalent increases in biomass across a diverse flora. As C_F became less available these opportunistic organisms may have declined in abundance. There were no C inputs to the soils during these incubations therefore the temporal increase in functional diversity of the 25°C samples reflected the presence of diverse functional genes and the degradation capacity of these communities that were potentially overshadowed by genetic material from fast-growing organisms earlier in the incubation. Prior studies support this idea, reporting that bacterial diversity was positively correlated with time of litter decomposition and the reduced substrate quality at later stages of decomposition (Dilly et al., 2004). Another study showed soil bacterial diversity to decrease following exposure to freeze-thaw cycles, even though total microbial abundance was unaffected (Sharma et al., 2006). That study also noted that fungal communities were more stable under freeze-thaw conditions and suggested that stronger shifts in bacterial diversity could be associated with the ability of bacteria to respond to C substrates mobilized following thaw.

An alternative explanation for the lower gene signal intensity may be that many groups were hindered at this higher 25°C temperature, rather than some individual groups of organisms

flourishing. The success of microorganisms in tundra can be attributed to freeze-tolerance (Schimel et al., 2005) and the organisms best acclimated to this environment may not thrive at 25°C. This explanation corresponds to the observation that representative functional diversity was low after two weeks at 25°C. Over time, organisms may have acclimated to the 25°C temperatures and thus, their DNA reached detectable quantities, which was observed at levels similar to the 15°C incubations later in the experiment. Previously, Steinweg et al. (2008) found that microbial carbon use efficiency (CUE) declined with warmer temperatures, resulting in less C allocated to cellular growth and more lost as CO₂, but this reduced efficiency was not maintained over time. Thus the high respiration rates of the 0-15 cm, 2 week, 25°C incubation soils could have actually reflected soils with lower biomass, wherein many groups of microbes were inactive under the unusual temperature and those active had low CUE's leading to higher respiration rates.

We expected to see variation in the abundances of fungal and bacterial genes throughout the incubation process. We also anticipated an initial rise in labile C-degradation genes, followed by increased recalcitrant-degradation genes after 280 days of incubation. However, our data actually reflect consistency among fungi and bacteria as well as across the suite of C-degradation genes in response to the effects of time and C pools. This was reflected by lower functional diversities of genes degrading all subcategories of C substrates observed after two weeks 25°C incubation. Additionally, when we focused on C-degradation gene abundances there are not significant differences among the soil depths, incubation times or temperatures.

4.3. Temperature sensitivity of SOM decomposition.

Temperature sensitivity of microbial SOM decomposition reported here is the intrinsic Q_{10} (Davidson and Janssens, 2006; Karhu et al., 2010; Deslippe et al., 2012; Sistla et al., 2014). The temperature sensitivity for C_F estimated using the equal C method was $Q_{10-EC} = 1.16 \pm 0.30$, (Supplemental Table 2) indicating that decomposition of the fast C pool is not temperature sensitive or it may even decrease as shown by the negative effect of increasing temperature on the decay rate (Table 3). It was previously shown that microbes from the organic active layer are able to use the same amount of substrates when temperature of incubation changed from 1°C to 10°C and even 20°C (Ernakovich and Wallenstein, 2015). On the other hand, decomposition of the slow decomposing C pool was affected by changes in temperature with Q_{10} overall average values greater than 2, as calculated by two different approaches, the short term ($Q_{10-ST} = 2.55 \pm 0.03$) and the equal C ($Q_{10-EC} = 2.19 \pm 0.13$). Temperature sensitivity obtained using the short term method primarily corresponded to C_S since this C pool dominated C fluxes the first time Q_{10} was estimated at DOI 14 in all depths and throughout the incubation period. A similar conclusion can be made when a $Q_{10-EC} = 2.66 \pm 0.40$ was estimated across all samples after normalizing contribution of C_S pool to total C respired exceeded 75%. This value was not affected by depth or field treatment and may explain the overall lack of the effect of DOI/time and depth on Q_{10-ST} . The one shift in Q_{10-ST} was the interaction of field warming and depth, where field warmed soils in the deepest depth had a slightly lower temperature response (Figure 4). This corresponds to a mineral soil layer, which also had higher bulk density at this same sample. It is not clear if bulk density was a direct or indirect driving factor, or what the mechanism would be if these were actually linked. Our results indicate that significant amount of C can be released from this slow C pool with an increase of soil temperature even if flux rates are small, considering that substrate availability will not be a limiting factor. The higher

temperature sensitivity for the slow decomposing C pool from tundra soils is consistent with the enzyme kinetics, where high activation energy is required to break down more complex C molecules (Davidson and Janssens, 2006; Craine et al., 2010a; Craine et al., 2010b; Wetterstedt et al., 2010; Conant et al., 2011).

Increases in Q_{10} (Q_{10-EC}), as C_R increased through time, were accompanied by changes in substrate use toward slower decomposable C (Streit et al., 2014), and the consequent shift in microbial communities (Conant et al., 2011; Deslippe et al., 2012; Sistla et al., 2014). This was reflected here by increased functional diversity of communities degrading C_S , primarily, which corresponded to changes in functional community structures with time of incubation and depth. When contributions of C_S to C_R reaches 50%, slow C decomposers probably dominate C decomposition showing a Q_{10} typical for the slow C pool. These microbes are able to use C compounds in the whole range of recalcitrance in the temperature range they were incubated (Ernakovich and Wallenstein, 2015), probably competing and dominating over the fast C decomposers. This is even more significant when contributions of C_S to C_R exceeds 60% and Q_{10-EC} is the same independently of the depth the soil is coming. In general, shifts in microbial communities associated with the slow decomposing C mediate increases in temperature sensitivity of soil organic matter decomposition (Karhu et al., 2014), although we didn't see the shifts in C degradation genes that were expected. This increased temperature sensitivity of SOM decomposition as slow decomposing C becomes dominant (lower organic matter quality) has also been observed in soils from ecosystems with lower soil C content than permafrost soils (Conant et al., 2008a; Conant et al., 2008b; Hartley and Ineson, 2008; Karhu et al., 2010).

Incubation results need to be extrapolated to the field with some caution, because for example, factors such as fresh soil C inputs are completely eliminated in laboratory incubations.

C inputs in the field are not likely to remain the same and it is not completely clear how warming will affect soil C balance. Our results suggest that sustained warming appears to have more effect on the dominant slow C pool as compared to the fast C pool. Increased plant input due to sustained warming (Chapin et al 2005; Tape et al 2006; Schuur et al., 2007; Pearson et al., 2013; Sistla et al., 2013; Sistla et al., 2014) could help to refill or maintain soil pools and counterbalance some soil C loss. Assuming both the quantity and quality of C inputs change or stay the same but temperature increases, microbial communities will shift towards slow C decomposers with a temperature sensitivity greater than 2 but, also with capacity to degrade a wider range of C compounds. This process will be amplified when additional slow C is exposed to decomposition by a deeper thaw as permafrost degrades (Natali et al., 2011). The microbial response to increased temperatures and the exposure of slow C will control regional C feedback to the atmosphere under the projected warming scenarios of 7 – 8 °C by the end of the 21st century (Trenberth et al., 2007, IPCC 2013).

Acknowledgements

This study was financially supported by the US Department of Energy, Terrestrial Ecosystem Sciences grant DE-SC0006982 and Biological Systems Research on the Role of Microbial Communities in Carbon Cycling Program grants DE-SC0004601 and DE-SC0010715. Other support was provided by the National Science Foundation CAREER program, Award #0747195; National Parks Inventory and Monitoring Program; National Science Foundation Bonanza Creek LTER program, Award #1026415; National Science Foundation Office of Polar Programs, Award #1203777”

705

706 REFERENCES

707

708 Allison, S.D., Wallenstein, M.D., Bradford, M.A., 2010. Soil-carbon response to warming dependent on
709 microbial physiology. *Nature Geoscience* 3, 336-340.

710 Amundson, R., 2001. The carbon budget in soils. *Annual Review of Earth and Planetary Sciences* 29, 535-
711 562.

712 Anderson, M.J., 2001. A new method for non-parametric multivariate analysis of variance. *Austral*
713 *Ecology* 26, 32-46.

714 Andren, O., Paustian, K., 1987. BARLEY STRAW DECOMPOSITION IN THE FIELD - A COMPARISON OF
715 MODELS. *Ecology* 68, 1190-1200.

716 Barcenas-Moreno, G., Gomez-Brandon, M., Rousk, J., Baath, E., 2009. Adaptation of soil microbial
717 communities to temperature: comparison of fungi and bacteria in a laboratory experiment. *Global*
718 *Change Biology* 15, 2950-2957.

719 Bates, D., Maechler, M., Bolker, B., Walker, S., 2014. lme4: Linear mixed-effects models using Eigen and
720 S4. R package version 1.0-6.

721 Buckeridge, K.M., Banerjee, S., Siciliano, S.D., Grogan, P., 2013. The seasonal pattern of soil microbial
722 community structure in mesic low arctic tundra. *Soil Biology & Biochemistry* 65, 338-347.

723 Chapin, F.S., Sturm, M., Serreze, M.C., McFadden, J.P., Key, J.R., Lloyd, A.H., McGuire, A.D., Rupp, T.S.,
724 Lynch, A.H., Schimel, J.P., Beringer, J., Chapman, W.L., Epstein, H.E., Euskirchen, E.S., Hinzman, L.D., Jia,
725 G., Ping, C.L., Tape, K.D., Thompson, C.D.C., Walker, D.A., Welker, J.M., 2005. Role of land-surface
726 changes in Arctic summer warming. *Science* 310, 657-660.

727 Conant, R.T., Drijber, R.A., Haddix, M.L., Parton, W.J., Paul, E.A., Plante, A.F., Six, J., Steinweg, J.M.,
728 2008a. Sensitivity of organic matter decomposition to warming varies with its quality. *Global Change*
729 *Biology* 14, 868-877.

730 Conant, R.T., Haddix, M., Paustian, K., 2010. Partitioning soil carbon responses to warming: Model-
731 derived guidance for data interpretation. *Soil Biology & Biochemistry* 42, 2034-2036.

732 Conant, R.T., Ryan, M.G., Agren, G.I., Birge, H.E., Davidson, E.A., Eliasson, P.E., Evans, S.E., Frey, S.D.,
733 Giardina, C.P., Hopkins, F.M., Hyvonen, R., Kirschbaum, M.U.F., Lavalley, J.M., Leifeld, J., Parton, W.J.,
734 Steinweg, J.M., Wallenstein, M.D., Wetterstedt, J.A.M., Bradford, M.A., 2011. Temperature and soil
735 organic matter decomposition rates - synthesis of current knowledge and a way forward. *Global Change*
736 *Biology* 17, 3392-3404.

737 Conant, R.T., Steinweg, J.M., Haddix, M.L., Paul, E.A., Plante, A.F., Six, J., 2008b. Experimental warming
738 shows that decomposition temperature sensitivity increases with soil organic matter recalcitrance.
739 Ecology 89, 2384-2391.

740 Cong, J., Liu, X., Lu, H., Xu, H., Li, Y., Deng, Y., Li, D., Zhang, Y., 2015. Analyses of the influencing factors of
741 soil microbial functional gene diversity in tropical rainforest based on GeoChip 5.0. Genomics Data 5,
742 397-398.

743 Craine, J., Spurr, R., McLauchlan, K., Fierer, N., 2010a. Landscape-level variation in temperature
744 sensitivity of soil organic carbon decomposition. Soil Biology & Biochemistry 42, 373-375.

745 Craine, J.M., Fierer, N., McLauchlan, K.K., 2010b. Widespread coupling between the rate and
746 temperature sensitivity of organic matter decay. Nature Geoscience 3, 854-857.

747 Davidson, E.A., Janssens, I.A., 2006. Temperature sensitivity of soil carbon decomposition and feedbacks
748 to climate change. Nature 440, 165-173.

749 Deng, Y and Z He. 2014. Microarray data analysis. In: Z He (ed) Microarrays: Current Technology,
750 Innovations and Applications, Caister Academic Press, <http://www.horizonpress.com/microarrays2>.

751 Dennis, J.G., 1977. DISTRIBUTION PATTERNS OF BELOW GROUND STANDING CROP IN ARCTIC TUNDRA
752 AT BARROW, ALASKA. Arctic and Alpine Research 9, 113-127.

753 Dennis, J.G., Johnson, P.L., 1970. SHOOT AND RHIZOME ROOT STANDING CROPS OF TUNDRA
754 VEGETATION AT BARROW ALASKA. Arctic and Alpine Research 2, 253-266.

755 Deslippe, J.R., Hartmann, M., Mohn, W.W., Simard, S.W., 2011. Long-term experimental manipulation of
756 climate alters the ectomycorrhizal community of *Betula nana* in Arctic tundra. Global Change Biology 17,
757 1625-1636.

758 Deslippe, J.R., Hartmann, M., Simard, S.W., Mohn, W.W., 2012. Long-term warming alters the
759 composition of Arctic soil microbial communities. Fems Microbiology Ecology 82, 303-315.

760 Deslippe, J.R., Simard, S.W., 2011. Below-ground carbon transfer among *Betula nana* may increase with
761 warming in Arctic tundra. New Phytologist 192, 689-698.

762 Dilly, O., Bloem, J., Vos, A., Munch, J.C., 2004. Bacterial diversity in agricultural soils during litter
763 decomposition. Applied and Environmental Microbiology 70, 468-474.

764 Dutta, K., Schuur, E.A.G., Neff, J.C., Zimov, S.A., 2006. Potential carbon release from permafrost soils of
765 Northeastern Siberia. Global Change Biology 12, 2336-2351.

766 Eliasson, P.E., McMurtrie, R.E., Pepper, D.A., Stromgren, M., Linder, S., Agren, G.I., 2005. The response of
767 heterotrophic CO₂ flux to soil warming. *Global Change Biology* 11, 167-181.

768 Ernakovich, J.G., Wallenstein, M.D., 2015. Permafrost microbial community traits and functional
769 diversity indicate low activity at in situ thaw temperatures. *Soil Biology & Biochemistry* 87, 78-89.

770 Fan, Z., Neff, J.C., Harden, J.W., Wickland, K.P., 2008. Boreal soil carbon dynamics under a changing
771 climate: A model inversion approach. *Journal of Geophysical Research-Biogeosciences* 113.

772 Fyfe, J.C., von Salzen, K., Gillett, N.P., Arora, V.K., Flato, G.M., McConnell, J.R., 2013. One hundred years
773 of Arctic surface temperature variation due to anthropogenic influence. *Scientific Reports* 3.

774 Haddix, M.L., Plante, A.F., Conant, R.T., Six, J., Steinweg, J.M., Magrini-Bair, K., Drijber, R.A., Morris, S.J.,
775 Paul, E.A., 2011. The Role of Soil Characteristics on Temperature Sensitivity of Soil Organic Matter. *Soil*
776 *Science Society of America Journal* 75, 56-68.

777 Hamdi, S., Moyano, F., Sall, S., Bernoux, M., Chevallier, T., 2013. Synthesis analysis of the temperature
778 sensitivity of soil respiration from laboratory studies in relation to incubation methods and soil
779 conditions. *Soil Biology & Biochemistry* 58, 115-126.

780 Harden, J.W., Koven, C.D., Ping, C.-L., Hugelius, G., McGuire, A.D., Camill, P., Jorgenson, T., Kuhry, P.,
781 Michaelson, G.J., O'Donnell, J.A., Schuur, E.A.G., Tarnocai, C., Johnson, K., Grosse, G., 2012. Field
782 information links permafrost carbon to physical vulnerabilities of thawing. *Geophysical Research Letters*
783 39.

784 Hartley, I.P., Heinemeyer, A., Ineson, P., 2007. Effects of three years of soil warming and shading on the
785 rate of soil respiration: substrate availability and not thermal acclimation mediates observed response.
786 *Global Change Biology* 13, 1761-1770.

787 Hartley, I.P., Hopkins, D.W., Garnett, M.H., Sommerkorn, M., Wookey, P.A., 2009. No evidence for
788 compensatory thermal adaptation of soil microbial respiration in the study of Bradford et al. (2008).
789 *Ecology Letters* 12, E12-E14.

790 Hartley, I.P., Ineson, P., 2008. Substrate quality and the temperature sensitivity of soil organic matter
791 decomposition. *Soil Biology & Biochemistry* 40, 1567-1574.

792 Hassol, S.J., 2004. Impacts of a warming Arctic: Arctic Climate Impact Assessment. Impacts of a warming
793 Arctic: Arctic Climate Impact Assessment., i-iii, 1-139.

794 He, Z., 2007. GeoChip: A comprehensive microarray for investigating biogeochemical, ecological and
795 environmental processes. *ISME J.* 1, 67-77.

796

797 Hill, M.O., Gauch, H.G., 1980. DETRENDED CORRESPONDENCE-ANALYSIS - AN IMPROVED ORDINATION
798 TECHNIQUE. *Vegetatio* 42, 47-58.

799 Hirsch, A.I., Trumbore, S.E., Goulden, M.L., 2002. Direct measurement of the deep soil respiration
800 accompanying seasonal thawing of a boreal forest soil. *Journal of Geophysical Research-Atmospheres*
801 108.

802 Holland, E.A., Neff, J.C., Townsend, A.R., McKeown, B., 2000. Uncertainties in the temperature sensitivity
803 of decomposition in tropical and subtropical ecosystems: Implications for models. *Global*
804 *Biogeochemical Cycles* 14, 1137-1151.

805 Hugelius, G., Routh, J., Kuhry, P., Crill, P., 2012. Mapping the degree of decomposition and thaw
806 remobilization potential of soil organic matter in discontinuous permafrost terrain. *Journal of*
807 *Geophysical Research-Biogeosciences* 117.

808 Hugelius, G., Strauss, J., Zubrzycki, S., Harden, J.W., Schuur, E.A.G., Ping, C.L., Schirrmeister, L., Grosse,
809 G., Michaelson, G.J., Koven, C.D., O'Donnell, J.A., Elberling, B., Mishra, U., Camill, P., Yu, Z., Palmtag, J.,
810 Kuhry, P., 2014. Estimated stocks of circumpolar permafrost carbon with quantified uncertainty ranges
811 and identified data gaps. *Biogeosciences* 11, 6573-6593.

812 IPCC, 2013: Annex I: Atlas of Global and Regional Climate Projections [van Oldenborgh, G.J., M. Collins, J.
813 Arblaster, J.H. Christensen, J. Marotzke, S.B. Power, M. Rummukainen and T. Zhou (eds.)]. In: *Climate*
814 *Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment*
815 *Report of the Intergovernmental Panel on Climate Change* [Stocker, T.F., D. Qin, G.-K. Plattner, M.
816 Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University
817 Press, Cambridge, United Kingdom and New York, NY, USA.

818 Jackson, R.B., Canadell, J., Ehleringer, J.R., Mooney, H.A., Sala, O.E., Schulze, E.D., 1996. A global analysis
819 of root distributions for terrestrial biomes. *Oecologia* 108, 389-411.

820 Karhu, K., Auffret, M.D., Dungait, J.A.J., Hopkins, D.W., Prosser, J.I., Singh, B.K., Subke, J.-A., Wookey,
821 P.A., Agren, G.I., Sebastia, M.-T., Gouriveau, F., Bergkvist, G., Meir, P., Nottingham, A.T., Salinas, N.,
822 Hartley, I.P., 2014. Temperature sensitivity of soil respiration rates enhanced by microbial community
823 response. *Nature* 513, 81-+.

824 Karhu, K., Fritze, H., Tuomi, M., Vanhala, P., Spetz, P., Kitunen, V., Liski, J., 2010. Temperature sensitivity
825 of organic matter decomposition in two boreal forest soil profiles. *Soil Biology & Biochemistry* 42, 72-82.

826 Katterer, T., Reichstein, M., Andren, O., Lomander, A., 1998. Temperature dependence of organic matter
827 decomposition: a critical review using literature data analyzed with different models. *Biology and*
828 *Fertility of Soils* 27, 258-262.

829 Kirschbaum, M.U.F., 1995. THE TEMPERATURE-DEPENDENCE OF SOIL ORGANIC-MATTER
830 DECOMPOSITION, AND THE EFFECT OF GLOBAL WARMING ON SOIL ORGANIC-C STORAGE. *Soil Biology &*
831 *Biochemistry* 27, 753-760.

832 Kirschbaum, M.U.F., 2004. Soil respiration under prolonged soil warming: are rate reductions caused by
833 acclimation or substrate loss? *Global Change Biology* 10, 1870-1877.

834 Kirschbaum, M.U.F., 2013. Seasonal variations in the availability of labile substrate confound the
835 temperature dependence of organic matter decomposition. *Soil Biology & Biochemistry* 57, 568-576.

836 Knoblauch, C., Beer, C., Sosnin, A., Wagner, D., Pfeiffer, E.-M., 2013. Predicting long-term carbon
837 mineralization and trace gas production from thawing permafrost of Northeast Siberia. *Global Change*
838 *Biology* 19, 1160-1172.

839 Knorr, W., Prentice, I.C., House, J.I., Holland, E.A., 2005. Long-term sensitivity of soil carbon turnover to
840 warming. *Nature* 433, 298-301.

841 Koven, C.D., Riley, W.J., Stern, A., 2013. Analysis of Permafrost Thermal Dynamics and Response to
842 Climate Change in the CMIP5 Earth System Models. *Journal of Climate* 26, 1877-1900.

843 Koven, C.D., Ringeval, B., Friedlingstein, P., Ciais, P., Cadule, P., Khvorostyanov, D., Krinner, G., Tarnocai,
844 C., 2011. Permafrost carbon-climate feedbacks accelerate global warming. *Proceedings of the National*
845 *Academy of Sciences of the United States of America* 108, 14769-14774.

846 Lavoie, M., Mack, M.C., Schuur, E.A.G., 2011. Effects of elevated nitrogen and temperature on carbon
847 and nitrogen dynamics in Alaskan arctic and boreal soils. *Journal of Geophysical Research-*
848 *Biogeosciences* 116.

849 Liang, Y.T., Z. H. He, L. Y. Wu, Y. Deng, G.H. Li, and J.-Z. Zhou. 2010. Development of a Common
850 Oligonucleotide Reference Standard for Microarray Data Normalization and Comparison across Different
851 Microbial Communities. *Applied and Environmental Microbiology*, 76(4):1088-94.

852 Liang, J., Li, D., Shi, Z., Tiedje, J.M., Zhou, J., Schuur, E.A.G., Konstantinidis, K.T., Luo, Y., 2015. Methods
853 for estimating temperature sensitivity of soil organic matter based on incubation data: A comparative
854 evaluation. *Soil Biology & Biochemistry* 80, 127-135.

855 Liebich, J., Schadt, C.W., Chong, S.C., He, Z.L., Rhee, S.K., Zhou, J.Z., 2006. Improvement of
856 oligonucleotide probe design criteria for functional gene microarrays in environmental applications.
857 *Applied and Environmental Microbiology* 72, 1688-1691.

858 Luo, Y.Q., Wan, S.Q., Hui, D.F., Wallace, L.L., 2001. Acclimatization of soil respiration to warming in a tall
859 grass prairie. *Nature* 413, 622-625.

860 Marschner, B., Bredow, A., 2002. Temperature effects on release and ecologically relevant properties of
861 dissolved organic carbon in sterilised and biologically active soil samples. *Soil Biology & Biochemistry* 34,
862 459-466.

863 Melillo, J.M., Steudler, P.A., Aber, J.D., Newkirk, K., Lux, H., Bowles, F.P., Catricala, C., Magill, A., Ahrens,
864 T., Morrisseau, S., 2002. Soil warming and carbon-cycle feedbacks to the climate system. *Science* 298,
865 2173-2176.

866 Mueller, C.W., Rethemeyer, J., Kao-Kniffin, J., Loeppmann, S., Hinkel, K.M., G. Bockheim, J., 2015. Large
867 amounts of labile organic carbon in permafrost soils of northern Alaska. *Global Change Biology* 21, 2804-
868 2817.

869 Natali, S.M., Schuur, E.A.G., Rubin, R.L., 2012. Increased plant productivity in Alaskan tundra as a result
870 of experimental warming of soil and permafrost. *Journal of Ecology* 100, 488-498.

871 Natali, S.M., Schuur, E.A.G., Trucco, C., Pries, C.E.H., Crummer, K.G., Lopez, A.F.B., 2011. Effects of
872 experimental warming of air, soil and permafrost on carbon balance in Alaskan tundra. *Global Change*
873 *Biology* 17, 1394-1407.

874 Natali, S.M., Schuur, E.A.G., Webb, E.E., Pries, C.E.H., Crummer, K.G., 2014. Permafrost degradation
875 stimulates carbon loss from experimentally warmed tundra. *Ecology* 95, 602-608.

876 O'Donnell, J.A., Harden, J.W., McGuire, A.D., Romanovsky, V.E., 2011. Exploring the sensitivity of soil
877 carbon dynamics to climate change, fire disturbance and permafrost thaw in a black spruce ecosystem.
878 *Biogeosciences* 8, 1367-1382.

879 Oksanen, J., Blanchet, F.G., Kindt, R., Legendre, P., Minchin, P.R., O'Hara, R.B., Simpson, G.L., Solymos,
880 P., Stevens, M.H.H., Wagner, H., 2012. *Vegan: Community Ecology Package*. R package version 2.0-3.

881 Paula, F.S., Rodrigues, J.L.M., Zhou, J., Wu, L., Mueller, R.C., Mirza, B.S., Bohannon, B.J.M., Nuesslein, K.,
882 Deng, Y., Tiedje, J.M., Pellizari, V.H., 2014. Land use change alters functional gene diversity, composition
883 and abundance in Amazon forest soil microbial communities. *Molecular Ecology* 23, 2988-2999.

884 Pearson, R.G., Phillips, S.J., Loranty, M.M., Beck, P.S.A., Damoulas, T., Knight, S.J., Goetz, S.J., 2013. Shifts
885 in Arctic vegetation and associated feedbacks under climate change. *Nature Climate Change* 3, 673-677.

886 Philippot, L., Spor, A., Henault, C., Bru, D., Bizouard, F., Jones, C.M., Sarr, A., Maron, P.-A., 2013. Loss in
887 microbial diversity affects nitrogen cycling in soil. *ISME Journal* 7, 1609-1619.

888 Pinheiro J, Bates D, DebRoy S, Sarkar D, Team, R.D.C., 2012. *nlme: Linear and Nonlinear Mixed Effects*
889 *Models*, R package version 3.1-105 ed.

890 Pries, C.E.H., Schuur, E.A.G., Crummer, K.G., 2012. Holocene Carbon Stocks and Carbon Accumulation
891 Rates Altered in Soils Undergoing Permafrost Thaw. *Ecosystems* 15.

892 Reichstein, M., Bednorz, F., Broll, G., Katterer, T., 2000. Temperature dependence of carbon
893 mineralisation: conclusions from a long-term incubation of subalpine soil samples. *Soil Biology &*
894 *Biochemistry* 32, 947-958.

895 Rey, A., Jarvis, P., 2006. Modelling the effect of temperature on carbon mineralization rates across a
896 network of European forest sites (FORCAST). *Global Change Biology* 12, 1894-1908.

897 Rinnan, R., Michelsen, A., Baath, E., 2011. Long-term warming of a subarctic heath decreases soil
898 bacterial community growth but has no effects on its temperature adaptation. *Applied Soil Ecology* 47,
899 217-220.

900 Romanovsky, V.E., Drozdov, D.S., Oberman, N.G., Malkova, G.V., Kholodov, A.L., Marchenko, S.S.,
901 Moskalenko, N.G., Sergeev, D.O., Ukraintseva, N.G., Abramov, A.A., Gilichinsky, D.A., Vasiliev, A.A., 2010.
902 Thermal State of Permafrost in Russia. *Permafrost and Periglacial Processes* 21, 136-155.

903 Rousk, J., Baath, E., 2011. Growth of saprotrophic fungi and bacteria in soil. *Fems Microbiology Ecology*
904 78, 17-30.

905 Salmon, V.G., Soucy, P., Mauritz, M., Celis, G., Natali, S.M., Mack, M.C., Schuur, E.A.G., 2016. Nitrogen
906 availability increases in a tundra ecosystem during five years of experimental permafrost thaw. *Global*
907 *Change Biology* doi:10.1111/gcb.13204.

908 Schadt, C.W., Martin, A.P., Lipson, D.A., Schmidt, S.K., 2003. Seasonal dynamics of previously unknown
909 fungal lineages in tundra soils. *Science* 301, 1359-1361.

910 Schädel, C., Schuur, E.A.G., Bracho, R., Elberling, B., Knoblauch, C., Lee, H., Luo, Y., Shaver, G.R., Turetsky,
911 M.R., 2014. Circumpolar assessment of permafrost C quality and its vulnerability over time using long-
912 term incubation data. *Global Change Biology* 20, 641-652.

913 Schaefer, K., Lantuit, H., Romanovsky, V.E., Schuur, E.A.G., Witt, R., 2014. The impact of the permafrost
914 carbon feedback on global climate. *Environmental Research Letters* 9.

915 Schaefer, K., Zhang, T., Bruhwiler, L., Barrett, A.P., 2011. Amount and timing of permafrost carbon
916 release in response to climate warming. *Tellus Series B-Chemical and Physical Meteorology* 63, 165-180.

917 Schimel, J.P., Mikan, C., 2005. Changing microbial substrate use in Arctic tundra soils through a freeze-
918 thaw cycle. *Soil Biology & Biochemistry* 37, 1411-1418.

919 Schmidt, M.W.I., Torn, M.S., Abiven, S., Dittmar, T., Guggenberger, G., Janssens, I.A., Kleber, M., Koegel-
920 Knabner, I., Lehmann, J., Manning, D.A.C., Nannipieri, P., Rasse, D.P., Weiner, S., Trumbore, S.E., 2011.
921 Persistence of soil organic matter as an ecosystem property. *Nature* 478, 49-56.

922 Schuur, E.A.G., Bockheim, J., Canadell, J.G., Euskirchen, E., Field, C.B., Goryachkin, S.V., Hagemann, S.,
 923 Kuhry, P., Lafleur, P.M., Lee, H., Mazhitova, G., Nelson, F.E., Rinke, A., Romanovsky, V.E., Shiklomanov,
 924 N., Tarnocai, C., Venevsky, S., Vogel, J.G., Zimov, S.A., 2008. Vulnerability of permafrost carbon to
 925 climate change: Implications for the global carbon cycle. *Bioscience* 58, 701-714.

926 Schuur, E.A.G., Crummer, K.G., Vogel, J.G., Mack, M.C., 2007. Plant species composition and productivity
 927 following permafrost thaw and thermokarst in alaskan tundra. *Ecosystems* 10, 280-292.

928 Schuur, E.A.G., McGuire, A.D., Schädel, C., Grosse, G., Harden, J.W., Hayes, D.J., Hugelius, G., Koven,
 929 C.D., Kuhry, P., Lawrence, D.M., Natali, S.M., Olefeldt, D., Romanovsky, V.E., Schaefer, K., Turetsky, M.R.,
 930 Treat, C.C., Vonk, J.E., 2015. Climate change and the permafrost carbon feedback. *Nature* 520, 171-179.

931 Schuur, E.A.G., Vogel, J.G., Crummer, K.G., Lee, H., Sickman, J.O., Osterkamp, T.E., 2009. The effect of
 932 permafrost thaw on old carbon release and net carbon exchange from tundra. *Nature* 459, 556-559.

933 Sharma, S., Szele, Z., Schilling, R., Munch, J.C., Schlöter, M., 2006. Influence of freeze-thaw stress on the
 934 structure and function of microbial communities and denitrifying populations in soil. *Applied and*
 935 *Environmental Microbiology* 72, 2148-2154.

936 Sistla, S.A., Moore, J.C., Simpson, R.T., Gough, L., Shaver, G.R., Schimel, J.P., 2013. Long-term warming
 937 restructures Arctic tundra without changing net soil carbon storage. *Nature* 497, 615-+.

938 Sistla, S.A., Rastetter, E.B., Schimel, J.P., 2014. Responses of a tundra system to warming using SCAMPS:
 939 a stoichiometrically coupled, acclimating microbe-plant-soil model. *Ecological Monographs* 84, 151-170.

940 Smith, S.L., Romanovsky, V.E., Lewkowicz, A.G., Burn, C.R., Allard, M., Clow, G.D., Yoshikawa, K., Throop,
 941 J., 2010. Thermal State of Permafrost in North America: A Contribution to the International Polar Year.
 942 *Permafrost and Periglacial Processes* 21, 117-135.

943 Soil Survey Staff (1999), *Soil Taxonomy: A Basic System of Soil Classification for Making and Interpreting*
 944 *Soil Surveys*, U.S. Govt. Print. Off., Washington, D. C.
 945

946 Steinweg, J.M., Plante, A.F., Conant, R.T., Paul, E.A., Tanaka, D.L., 2008. Patterns of substrate utilization
 947 during long-term incubations at different temperatures. *Soil Biology & Biochemistry* 40, 2722-2728.

948 Streit, K., Hagedorn, F., Hiltbrunner, D., Portmann, M., Saurer, M., Buchmann, N., Wild, B., Richter, A.,
 949 Wipf, S., Siegwolf, R.T.W., 2014. Soil warming alters microbial substrate use in alpine soils. *Global*
 950 *Change Biology* 20, 1327-1338.

951 Tape, K., Sturm, M., Racine, C., 2006. The evidence for shrub expansion in Northern Alaska and the Pan-
 952 Arctic. *Global Change Biology* 12, 686-702.

953 Tarnocai, C., Canadell, J.G., Schuur, E.A.G., Kuhry, P., Mazhitova, G., Zimov, S., 2009. Soil organic carbon
 954 pools in the northern circumpolar permafrost region. *Global Biogeochemical Cycles* 23.

955 Trenberth, K.E., Jones, P.D., Ambenje, P., Bojariu, R., Easterling, D., Klein Tank, A., Parker, D.,
956 Rahimzadeh, F., Rusticucci, M., Soden, B., Zhai, P. 2007. Observations: Surface and Atmospheric Climate
957 Change. In: Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the
958 Fourth Assessment Report of the Intergovernmental Panel on Climate Change [Solomon, S., D. Qin, M.
959 Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor and H.L. Miller (eds)]. Cambridge University Press,
960 Cambridge, United Kingdom and New York, NY, USA.

961 Trumbore, S.E., 1997. Potential responses of soil organic carbon to global environmental change.
962 Proceedings of the National Academy of Sciences of the United States of America 94, 8284-8291.

963 Tu, Q., Yu, H., He, Z., Deng, Y., Wu, L., Van Nostrand, J.D., Zhou, A., Voordeckers, J., Lee, Y.-J., Qin, Y.,
964 Hemme, C.L., Shi, Z., Xue, K., Yuan, T., Wang, A., Zhou, J., 2014. GeoChip 4: a functional gene-array-
965 based high-throughput environmental technology for microbial community analysis. Molecular Ecology
966 Resources 14, 914-928.

967 van Wijk, M.T., Williams, M., Gough, L., Hobbie, S.E., Shaver, G.R., 2003. Luxury consumption of soil
968 nutrients: a possible competitive strategy in above-ground and below-ground biomass allocation and
969 root morphology for slow-growing arctic vegetation? Journal of Ecology 91, 664-676.

970 Waldrop, M.P., Wickland, K.P., White, R., III, Berhe, A.A., Harden, J.W., Romanovsky, V.E., 2010.
971 Molecular investigations into a globally important carbon pool: permafrost-protected carbon in Alaskan
972 soils. Global Change Biology 16, 2543-2554.

973 Wallenstein, M.D., McMahon, S., Schimel, J., 2007. Bacterial and fungal community structure in Arctic
974 tundra tussock and shrub soils. Fems Microbiology Ecology 59, 428-435.

975 Wang, H., He, Z., Lu, Z., Zhou, J., Van Nostrand, J.D., Xu, X., Zhang, Z., 2012. Genetic Linkage of Soil
976 Carbon Pools and Microbial Functions in Subtropical Freshwater Wetlands in Response to Experimental
977 Warming. Applied and Environmental Microbiology 78, 7652-7661.

978 Wetterstedt, J.A.M., Persson, T., Agren, G.I., 2010. Temperature sensitivity and substrate quality in soil
979 organic matter decomposition: results of an incubation study with three substrates. Global Change
980 Biology 16, 1806-1819.

981 Wickland, K.P., Neff, J.C., 2008. Decomposition of soil organic matter from boreal black spruce forest:
982 environmental and chemical controls. Biogeochemistry 87, 29-47.

983 Wu, L., Liu, X., Schadt, C.W., Zhou, J., 2006. Microarray-based analysis of subnanogram quantities of
984 microbial community DNAs by using whole-community genome amplification. Applied and
985 Environmental Microbiology 72, 4931-4941.

986 Xue, K., Wu, L., Deng, Y., He, Z., Van Nostrand, J., Robertson, P.G., Schmidt, T.M., Zhou, J., 2013.
987 Functional Gene Differences in Soil Microbial Communities from Conventional, Low-Input, and Organic
988 Farmlands. Applied and Environmental Microbiology 79, 1284-1292.

989 Yang, Y., Gao, Y., Wang, S., Xu, D., Yu, H., Wu, L., Lin, Q., Hu, Y., Li, X., He, Z., Deng, Y., Zhou, J., 2014. The
990 microbial gene diversity along an elevation gradient of the Tibetan grassland. *Isme Journal* 8, 430-440.

991 Yergeau, E., Bokhorst, S., Kang, S., Zhou, J., Greer, C.W., Aerts, R., Kowalchuk, G.A., 2012. Shifts in soil
992 microorganisms in response to warming are consistent across a range of Antarctic environments. *ISME*
993 *Journal* 6, 692-702.

994 Zhou, J., Deng, Y., Zhang, P., Xue, K., Liang, Y., Van Nostrand, J.D., Yang, Y., He, Z., Wu, L., Stahl, D.A.,
995 Hazen, T.C., Tiedje, J.M., Arkin, A.P., 2014. Stochasticity, succession, and environmental perturbations in
996 a fluidic ecosystem. *Proceedings of the National Academy of Sciences of the United States of America*
997 111, E836-E845.

998 Zhou, J., Xue, K., Xie, J., Deng, Y., Wu, L., Cheng, X., Fei, S., Deng, S., He, Z., Van Nostrand, J.D., Luo, Y.,
999 2012. Microbial mediation of carbon-cycle feedbacks to climate warming. *Nature Climate Change* 2, 106-
1000 110.

1001 Zhou, J., He, Z., Van Nostrand, J.D., Wu, L.Y., Deng, Y. 2010. Applying GeoChip Analysis to Disparate
1002 Microbial Communities. *Microbes*, 5:60-65.

1003 Zhou, J.Z., Bruns, M.A., Tiedje, J.M., 1996. DNA recovery from soils of diverse composition. *Appl.*
1004 *Environ. Microbiol.* 62, 316-322.

1005 Zhu, B., Cheng, W., 2011. Constant and diurnally-varying temperature regimes lead to different
1006 temperature sensitivities of soil organic carbon decomposition. *Soil Biology & Biochemistry* 43, 866-869.

1007 Zuur, A.F., Ieno, E.N., Walker, N.J., Saveliev, A.A., Smith, G.M., 2010. *Mixed Effects Models and*
1008 *Extensions in Ecology with R.* Springer N.Y.

1009

1010

1011

1012

1013

1014

Table 1. Initial properties of soils from a tundra warming experiment (mean $\pm$ SE; N=6)

Treatment	Depth (cm)	%C	%N	C:N	Bulk density	pH
Control	0-15	41.80 $\pm$ 0.35	1.16 $\pm$ 0.09	37.21 $\pm$ 3.03	0.09 $\pm$ 0.01	4.67 $\pm$ 0.03
	15-25	34.83 $\pm$ 4.14	1.69 $\pm$ 0.21	20.89 $\pm$ 0.55	0.30 $\pm$ 0.11	4.73 $\pm$ 0.09
	45-55	17.70 $\pm$ 2.97	0.74 $\pm$ 0.17	25.50 $\pm$ 1.56	0.42 $\pm$ 0.05	5.15 $\pm$ 0.14
Field Warming	0-15	39.88 $\pm$ 0.69	1.27 $\pm$ 0.04	31.66 $\pm$ 1.34	0.12 $\pm$ 0.03	4.71 $\pm$ 0.06
	15-25	33.67 $\pm$ 2.96	1.66 $\pm$ 0.17	20.54 $\pm$ 0.58	0.30 $\pm$ 0.07	4.97 $\pm$ 0.07
	45-55	12.53 $\pm$ 2.51	0.48 $\pm$ 0.09	26.42 $\pm$ 1.44	0.86 $\pm$ 0.18	5.05 $\pm$ 0.10

Table 2. Statistics for initial properties of soils from a tundra warming experiment.

Model	Coefficient	Estimate	SE	Df	t-value	P-value
Soil properties						
C (%)	Intercept	0.809	0.028	29	29.318	<0.0001
	Depth	-0.008	0.001	29	-9.886	<0.0001
N (%)	Intercept	0.139	0.007	29	18.472	<0.0001
	Depth	-0.001	0.000	29	-5.127	<0.0001
Bulk density	Intercept	-1.231	0.094	28	-13.130	<0.0001
	Depth	0.017	0.002	28	7.472	<0.0001
	Field warming	0.131	0.078	28	1.676	0.1050
pH	Intercept	0.659	0.009	28	73.171	<0.0001
	Depth	0.001	0.000	28	5.555	<0.0001

1023 Table 3. Statistics for parameters derived from laboratory incubations of soils from an Alaskan tundra warming experiment incubated at
1024 15°C and 25°C for 365 days.

Model	Coefficient	Estimate	SE	Df	t-value	P-value
Parameters						
C_F	Intercept	-0.039	0.142	60	-0.277	0.7830
	Depth	-0.008	0.001	60	-5.953	<0.0001
	Temp _{incubation}	0.015	0.004	60	3.474	0.0010
	C:N	0.015	0.003	60	4.739	<0.0001
C_S	Intercept	1.566	0.037	60	42.107	<0.0001
	Depth	0.002	0.000	60	6.323	<0.0001
	Temp _{incubation}	-0.004	0.001	60	-3.622	<0.0001
	C:N	-0.004	0.001	60	-4.713	<0.0001
k_F	Intercept	0.282	0.031	61	9.115	<0.0001
	Temp _{incubation}	-0.004	0.001	61	-2.959	0.0040
	Bulk density	-0.069	0.019	61	-3.580	0.0010
k_S	Intercept	-3.517	0.592	59	-5.945	<0.0001
	Depth	-0.012	0.002	59	-6.676	<0.0001
	Temp _{incubation}	0.032	0.004	59	7.408	<0.0001
	C:N	0.020	0.003	59	6.257	<0.0001
	pH	-0.224	0.123	59	-1.824	0.0730
C_{respired}	Intercept	1.913	0.531	59	3.606	0.0010
	Depth	-0.012	0.002	59	-7.846	<0.0001
	Temp _{incubation}	0.031	0.004	59	8.042	<0.0001
	C:N	0.019	0.003	59	6.470	<0.0001
	pH	-0.170	0.110	59	-1.539	0.1290
$Q_{10\text{-ST}}$	Intercept	0.392	0.018	194	22.277	<0.0001
	Field warming	0.066	0.023	194	2.861	0.0050
	Depth	0.001	0.000	194	1.637	0.1030
	Day of incubation	0.000	0.000	194	-1.597	0.1120
	Field warming:Depth	-0.002	0.001	194	-3.827	<0.0001

1025

Table 4. Dissimilarity analysis of the microbial communities based on Geochip data by Adonis in Alaskan tundra soils incubated at 15°C and 25°C for 365 days.

Geochip			
	F.Model	R2	P-value
Plot	0.803	0.01225	0.611
Depth	13.289	0.04053	0.001
Field warming	1.459	0.00445	0.198
DOI	34.552	0.10538	0.001
Temp _{incubation}	31.161	0.09504	0.001
Field warming : depth	1.662	0.00507	0.169
Depth : DOI	1.036	0.00316	0.297
Field warming : DOI	2.327	0.0071	0.084
Depth : Temp _{incubation}	1.048	0.0032	0.325
Field warming : Temp _{incubation}	1.808	0.00551	0.142
DOI : Temp _{incubation}	33.635	0.10258	0.001

DOI = Day of incubation

Temp_{incubation} = Temperature of incubation

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Figure 6.

Figure 7.

Figure 8.

Figure 9.

Figure 1. Size of the fast (C_F) (a) and slow (C_S) (b) decomposing carbon pools ($\text{mgC/gC}_{\text{initial}}$) for field control (Ctrl) and field warmed (Warm) Alaskan tundra soils incubated at 15°C and 25°C (hatched bars). (Mean $\pm$ SE, $n = 6$).

Figure 2. Decay rates for the fast (k_F) (a) and slow (k_S) (b) decomposing carbon pools for field control (Ctrl) and field warmed (Warm) Alaskan tundra soils incubated at 15°C and 25°C (hatched bars). (Mean $\pm$ SE, $n = 6$).

Figure 3. Total carbon respired, C_R ($\text{mgC/Cg}_{\text{initial}}$) for field control (Ctrl) and field warmed (Warm) Alaskan tundra soils from different depths incubated at 15°C and 25°C (hatched bars) after one year of incubation. (Mean $\pm$ SE, $n = 6$).

Figure 4. Overall average $Q_{10\text{-ST}}$ (15°C & 25°C) for field control (Ctrl) and field warmed (Warm) Alaskan tundra soils from different depths after one year of incubation. (Mean $\pm$ SE, $n = 12$).

Figure 5. Relationship between $Q_{10\text{-EC}}$ (continuous line) and total amount of carbon respired (C_R , $\text{mgC/g}_{\text{initial_C}}$), and contribution of the slow decomposing C pool ($\%C_S$) (right Y axis) to C_R at 15°C incubation (dotted line) and 25°C incubation (dashed line). Control soils (Panels a- c, left), field warmed soils (Panels d – f, right). Top, intermediate, and bottom panels correspond to 0-15, 15 -25 and 45-55 cm depths. Data are the average of six replicates, error bars omitted for simplicity of visualization.

Figure 6. DCA plots reflecting effects of time of incubation on microbial functional communities for 15 – 25 cm Alaskan tundra soils incubated at 25°C. The placement of the samples along the axes are output of the DCA ordination algorithm, which functions to find maximum correlations between all GeoChip probe scores and sample scores.

Figure 7. Mean signal intensities of GeoChip 5.0 probes targeting genes involved in C-degradation. Solid fill (white and black): 0-15cm, stripped fill: 15-25cm, dotted fill: 45 – 55cm.

Figure 8. Mean signal intensities from GeoChip 5.0. Values represent means and standard deviations from 6 replicate soils and all probes categorized as targeting genes involved in degradation of denoted carbon sources. Carbon substrates range from fast to slow decomposing C (left to right).

Figure 9. Mean signal intensities from GeoChip 5.0 for only probes showing positive signal intensity. Values represent means and standard deviations from 6 replicate soils and all probes categorized as targeting genes involved in degradation of denoted carbon sources. Carbon substrates range from fast to slow decomposing C (left to right).