1   **The nature and dynamics of soil organic matter: Plant inputs,**
2   **microbial transformations, and organic matter stabilization**

3
4   **Eldor A. Paul**

5
6   Natural Resource Ecology Laboratory and Department of Soil and Crop Sciences

7   Colorado State University

8   Fort Collins, CO 80523-1499, USA

9   Eldor.paul@colostate.edu

10

11 Abstract

12 This review covers historical perspectives, the role of plant inputs, and the nature and
13 dynamics of soil organic matter (SOM), often known as humus. Information on turnover
14 of organic matter components, the role of microbial products, and modeling of SOM, and
15 tracer research should help us to anticipate what future research may answer today's
16 challenges. Our globe's most important natural resource is best studied relative to its
17 chemistry, dynamics, matrix interactions, and microbial transformations. Humus has
18 similar, worldwide characteristics, but varies with abiotic controls, soil type, vegetation
19 inputs and composition, and the soil biota. It contains carbohydrates, proteins, lipids,
20 phenol-aromatics, protein-derived and cyclic nitrogenous compounds, and some still
21 unknown compounds. Protection of transformed plant residues and microbial products
22 occurs through spatial inaccessibility-resource availability, aggregation of mineral and
23 organic constituents, and interactions with sesquioxides, cations, silts, and clays. Tracers
24 that became available in the mid-20th century made the study of SOM dynamics possible.
25 Carbon dating identified resistant, often mineral-associated, materials to be thousands of
26 years old, especially at depth in the profile. The ¹³C associated with C₃-C₄ plant switches
27 characterized slow turnover pools with ages ranging from dozens to hundreds of years.
28 Added tracers, in conjunction with compound-specific product analysis and incubation,
29 identified active pools with fast turnover rates. Physical fractionations of the intra- and
30 inter-aggregate materials, and those associated with silt and clay, showed that all pools
31 contain both old and young materials. Charcoal is old but not inert. The C:N ratio
32 changes from 25 to 70:1 for plant residues to 6 to 9:1 for soil biota and microbial

products associated with soil minerals. Active, slow and passive (resistant) pool concepts have been well used in biogeochemical models. The concepts discussed herein have implications for today's challenges in nutrient cycling, biogeochemistry, soil ecosystem functioning, pollution control, feeding the expanding global population and global change.

1. Introduction

Soils play a major role in ecosystem dynamics, biotic diversity, water infiltration, erosion control, global food security, biofuels, the control of and response to global change, and the Earth's fresh water quality (Fang et al., 2005; Cheeke et al., 2013; Coleman and Wall, 2014). Decomposition must be slightly lower than plant inputs, on a long-term basis, to produce the soil organic matter (SOM) found in the world's biomes. Our present, industrial society is based on the fossil fuels produced during the Carboniferous and Permian eras. The precursors of the fossil fuels were deposited over millions of years under conditions of high plant productivity with limited decomposition due to high moisture contents and low, fungal lignase activity (Horwath, 2015). These hydrocarbons are now being returned to the atmosphere in excess amounts. Carbon dioxide, the most important plant nutrient, is the major atmospheric component of the global C cycle and a greenhouse gas. The simultaneous advent of agriculture, which occurred 10,000 yr ago on three continents, could be attributable to an increase in atmospheric CO₂ to above 200 ppmv (Sage, 1995). A level below 200 could be too low to allow for domestication of crops. Increased CO₂ levels, rising to today's 400+ ppmv, together with other

components of the green revolution such as improved plant genetics and fertilization, have played a part in the three-fold increase in global grain production and associated plant residue inputs since 1950. Future levels of atmospheric CO₂, as well as predicted increased temperatures will also impact SOM dynamics (Batjes, 2014).

A great wealth of information on soil organic carbon (SOC) constituents and dynamics can be found in the literature of the mid-20th century when tracers first became available. Hopefully, this information is not lost, repeated, or disregarded in today's experiments. This review highlights some of the original work, relates it to modern concepts, and will hopefully lead to improved future research. The tables, figures and associated information in this paper are primarily based on the authors' work, but reflect the literature in general. It is not possible to cover all the literature over this extended period. The selected references of both the older and modern literature hopefully can lead interested scientists to the relevant papers.

The composition and global distribution of SOM is a valuable storehouse of information on vegetation, parent materials, climate, and disturbance (Table 1). De Deyn et al. (2008) published a review of plant, functional traits and soil C sequestration in contrasting biomes. Tropical forests constitute 12.2% of the Earth's land area with high net primary productivity (NPP) and annual C inputs of 2.03 kg C m⁻². Their soils, long thought to be SOM deficient, are now known to have fairly high SOC contents of 12 kg C m⁻² or 692 PgC (10¹⁵g) on a global basis when analyzed to the depth 3 m.

74 A useful estimate of gross SOC turnover, at an ecosystem level, can be attained by
75 relating the SOC level to litter inputs (SOC content/annual litter input) on the assumption
76 of steady state conditions. Tropical, forest soils, characterized by high rainfall and
77 temperatures, trees with endotrophic mycorrhiza, lower C:N ratios and abundant soil
78 fauna, have gross turnover times of 5.3 yr in the surface and 22 yr over the 3 m profile
79 depth. Thus, they are major contributors to the annual CO₂ flux. The lower inputs for
80 temperate forests with diverse trees and high litter decomposability, from broad leaved
81 trees, result in lower SOC stocks that however can have reasonable soil ages at the
82 surface (Mathieu et al., 2015). The higher SOC stocks in boreal forests reflect inputs
83 from mostly coniferous trees with ectomycorrhizal fungi, lower temperatures and often
84 wet conditions that result in high C:N ratios (Aitkenhead and McDowell, 2000).
85 Podzolization can affect the SOC and soil organic nitrogen (SON) contents under such
86 conditions. Croplands span the full range of mean annual, global temperatures.
87 Disturbance, such as that caused by cultivation, usually results in decreased stocks of
88 SOC with higher turnover times (Follett et al., 1997; Lal et al., 2015), whereas
89 afforestation of agricultural land can lead to increased stocks (Morris et al., 2011; Mellor
90 et al., 2013).

91 The deserts are affected by abiotic as well as biotic decomposition resulting in low SOC
92 per unit area and low C:N ratios (Adair et al., 2008). Grasslands with high, beneath- to
93 above- ground plant inputs usually have high SOC levels. In addition to the climatic-
94 grass root-faunal interactions, there are effects of Ca concentrations (O'Brien et al., 2015)

and the fact that smectitic clays, protective of SOC, are formed in grasslands (Yuan and Theng, 2012).

The tundra soils at 19.6 kg C m^{-1} (Table 1) contain a total of 818 Pg SOC with a turnover time of 165 yr. They have slow turnover rates in both the surface and subsurface even though mineral associations are less than those in other biomes (Tarnocai et al., 2009; Schuur et al., 2015). The tundra SOC stocks are continually being revised as more information becomes available. Hugelius et al. (2014) estimate the northern, circumpolar permafrost SOC stock to 3 m at $1035 \pm 130 \text{ Pg}$ rather than the 818 shown in Table 1. The large stock of SOC in the tundra soil, associated with ice and in estuarine sediments, could be very sensitive to warming (Ping et al., 2015). Many of the upland soils are expected to become drier. The wet environment in many lowland tundra sites prevents decomposition. Management scenarios that prevent drainage and maintain wetlands under global warming of the Arctic should be considered. Wetlands in all temperature regimes across the globe still retain high SOC levels with high C:N ratios attributable to the effect of anaerobiosis on decomposition (Aitkenhead and McDowell, 2000).

The gross, turnover time does not represent the true mean residence time (MRT) of the SOC due to the vast range in MRTs of the different pools-components (Bruun et al., 2005). Carbon dates (Campbell et al., 1967; Paul et al., 2001a; Trumbore, 2009) show that measured SOC MRTs are much higher than the gross turnover times in Table 1. The majority of plant residues are decomposed in the first year, especially in the tropics (Trumbore, 1993). ^{13}C studies have shown that plant residues are transformed during

decomposition into microbial products that interact with the soil matrix to protect 5 to 10% of the added grassland, plant residues as SOM over a 100 yr period (Follett et al., 1997). Approximately 10% of added ^{14}C is found in microbial tissue one year after addition, dropping to 4% after four years (Jenkinson and Ladd, 1981).

2. Plant inputs

Plant inputs drive decomposition and SOM formation and could be viewed as the controlling pump in the C cycle. There are indications that the native biota, adapted to a particular site, decomposes the litter from that site more rapidly than if the litter was placed on another site, i.e., has a home field advantage (Wallenstein et al., 2013). An adequate understanding of root inputs continues to be a problem (Follett et al., 1997; Adair et al., 2008; Hinsinger et al., 2012). Root-derived materials occur as exudates, fine and coarse root detritus, plant polysaccharides at the growing root tip, and mycorrhizal hypha (Calderón et al., 2012; Balestrini et al., 2015). On average, cereals, due to genetic selection for more above-ground growth, transfer 20 to 30% of their assimilated C to roots compared to the 30 to 50% for pasture grasses. However, because of the increased growth potential of cultivated cereals, the amount of the root-derived C is considered similar (Kuzyakov, 2010). This review showed root respiration accounted for 6 to 17% of the photosynthate for different plants, with 3 to 6% of the assimilated C transferred to soil. Another review (Jones et al., 2009) concluded that 20 to 40% of photosynthetic C is translocated underground. Of this, 5 to 55% remains in the roots (average of $19 \pm 8\%$), 2 to 18% forms SOM, and $12 \pm 5\%$ is respired. Symbiotic root microbes, such as N fixers and

mycorrhiza require extensive substrate (Paul and Kucey, 1981), but are so closely intertwined with root physiology that they are probably more closely associated with plant (autotrophic) than the soil-associated (heterotrophic) respiration.

The basic controls on plant photosynthesis, C translocation (Horwath et al., 1994; Rasse et al., 2005), and below-ground processes are similar throughout the large number of plant-microbial interactions in different ecosystems. The great range in the transfer of C to the roots, rhizosphere organisms, and soils is associated with differences in the plant-food web-abiotic controls in specific ecosystems (Moore and de Ruiter, 2012). Therefore, it is crucial to understand the factors involved in different ecosystems (Drigo et al., 2012).

An increased knowledge of the genetic information concerning symbiotic and rhizosphere organisms and their genes is giving us a better understanding of the C cycle (Balestrini et al., 2015). There is now a rich data source on gene expression and sequencing information, which provides insight into the controls on plant and microbial tissue decomposition, nutrient transfers, signal molecules, and enzyme production at the soil-root interface (Hinsinger et al., 2012; Pietramella et al., 2012).

Dahlman and Kucera (1968) used ^{14}C to show that the roots of a C_4 dominated tall-grass prairie had three times the biomass of the above-ground standing crop, but had a turnover time of three to four years. The above-ground biomass of a northern C_3 grassland was much smaller, but accounted for ~50% of the incorporated label; roots and rhizomes accounted for 25% and below-ground respiration and transfer to the soil for 25%

(Warembourg and Paul, 1977). Sampling through the growing season made it possible to measure a root turnover of 155 days. Wheat shoots under similar conditions incorporated 69% of the label above-ground, 14% below-ground, and respired 17%.

Trees store much of their assimilated C in perennial wood. Data for a 5-yr-old poplar plantation (Table 2) showed that leaves and litter, which accounted for 37% of the biomass, contained 24% of the ^{14}C immediately after labeling. The label distribution in other plant parts was: 48% in stems and branches, 1.7% in fine roots, and 8.1% in coarse roots (Horwath et al., 1994). Root-soil-associated respiration accounted for 7.7% of the ^{14}C label; only 0.45% of the label went directly into the microbial biomass. Measurement after 328 days showed the expected loss of ^{14}C from the leaves after leaf fall, with 26% retention in the stem and branches and 9.7 % in the roots. The root-soil respiration accounted for 15.3% of the label after one year, and the microbial biomass represented 0.3%. The microbial biomass lost 25% of its label after one year, verifying the slow microbial biomass turnover rates found in other studies (Paul and Voroney, 1980).

The microorganisms involved in symbiotic systems, such as N_2 fixation and mycorrhizal-P uptake, require extensive amounts of energy. A ^{14}C study showed that nodules in faba beans (*Vicia faba*) utilized 12% of the photosynthate, whereas the mycorrhiza utilized only 4% (Paul and Kucey, 1981). In both systems, the amount of ^{14}C attributed to symbiotic respiration-nutrient uptake was much greater than that fixed in the nodule or mycorrhizal tissues. Plants have some ability to compensate for the needs of their microbial partners by increased photosynthesis (Kucey and Paul, 1982). Calderón et al.

179 (2012) applied a $^{14}\text{CO}_2$ pulse exposure to a *Sorghum bicolor*-*Glomus* association and
180 found that mycorrhizal plants fixed 21% more CO_2 than non-mycorrhizal plants, but still
181 had somewhat lower biomass production indicating there was an incomplete
182 compensation of the needs of the mycorrhizal partners. This was also noted by Snellgrove
183 et al. (1982). The mycorrhizal fatty acid, 16:1 $\omega 5$, comprising 30% of the fatty acids of
184 mycorrhizal roots, had a turnover time of 210 hr^{-1} . This was similar to the turnover time
185 of fungal arbuscules (Cox et al., 1975) showing the importance of this fatty acid in
186 endomycorrhizal-plant C transfers. The sorghum-mycorrhizal association transferred
187 6.3% of the tracer to the soil relative to only 2.3% in the non-mycorrhizal plants
188 indicating the potential of mycorrhizal fungi to transfer assimilated C to soil.

189 There also are numerous indications that plant inputs can decrease SOM levels, such as in
190 N mining and priming of the decomposition of native SOM (Kuzyakof and Domanski,
191 2000; Fontaine et al., 2003). Measurements with tracers and mass balance calculations
192 that determine a net loss of SOM in the presence of added substrate show that priming is
193 dependent on external factors (Bingeman et al., 1953). An example of N mining was
194 shown when a corn crop, grown on soils with an extensive, active SOM pool, built up by
195 previous legume crops or compost addition, contained more N than an adjoining wheat
196 crop or even the N mineralized in a fallow plot (Sanchez et al., 2002). Roots can affect
197 rhizosphere effects by drying the soil resulting in decreased decomposition, and dissolved
198 organic matter in the soil can move to the root surfaces by mass flow.

The significance of plant-functional traits (De Deyn et al., 2008) and plant-derived C on soil biota and SOM formation (Bardgett, 2005) is not yet fully understood. Plant-microbial-SOM interactions are complex (Balestrini et al., 2015) and transfer rates vary widely with different symbiotic partners (Hinsinger et al., 2012) and in different ecosystems (Jones et al., 2009). Fortunately some of the techniques in use today will be of use in our research. Spence et al. (2011), using HR-MAS NMR, concluded that 50% of SOM is derived from microbial products. Workers such as Pisana et al. (2013) have used biological marker analysis to identify plant residue products, such as suberin and cutin and long chain aliphatics, in SOM. Simpson et al. (2011) reviewed such studies relative to molecular interactions that affect global processes. Gillespie et al. (2014) combined XANES spectroscopy (Lehman et al. 2005) at the C and N edges with microbial biomarkers to measure the biochemical composition of the SOM of the soil fine fraction and the relative contribution of fungi and bacteria to this SOM. Chenu et al. (2015) provides information on other techniques, such as NanoSIMS, IR spectroscopy and electron and atomic microscopy that can help determine the fate of plant residues in soil.

3. The Nature and Formation of Soil Organic Matter

3.1 Historical perspectives

Ancient Chinese dynasties (2357 to 2261 BCE) recognized differences in soil color in soil classification for taxation purposes (Coleman et al., 2004). The ancient Vidic civilization of India classified soils by color, land forms, erosion, vegetation, land use, and human health implications (Paul, 2007). Waksman (1938), in his treatise entitled

220 “Humus,” stated that the literature from Theophrastus (373-328 BCE) to Wallerius
 221 (1709-1778 CE) equated the words “humus” with the color of the surface, soil layer.
 222 Dark colored soils were known to be more fertile, more easily cultivated, and to hold
 223 more water. Linneaus (1707-1778), the great Swedish botanist, classified garden soils as
 224 *Humus daedalea* and field soils as *Humus rualis* (Feller, 1997). He possibly foretold the
 225 difficulties in microbial classification, before the advent of the molecular age, when he
 226 placed all organisms, seen under the microscope by Leeuwenhoek, under the genus
 227 *Chaos* (Paul, 2007).

228 Archard in 1786 and Berzelius in 1806 extracted humic substances from soil with alkali
 229 and showed their interaction with metals (Paul, 2007; Warkentin, 2006). Müller’s studies
 230 in 1881 showed the role of earthworms in the production of nutrient-rich soil (Mull)
 231 relative to the nutrient poor litter layer (Mor), with little biological activity (Hillel, 1991;
 232 Feller, 1997, Feller et al., 2003). The work of Dokochaev and Kustyechev in 1883
 233 (Kononova, 1961) led to modern concepts of soil formation and ecosystem studies,
 234 including the interactions of climate, vegetation (biota), parent materials, topography (the
 235 landscape), and time. Hilgaard recognized the role of cations, especially calcium, in SOM
 236 stabilization. Heber, Deherain, and Omylyanski established the role of biological-abiotic
 237 controls on humus formation, plant residue structure, and decomposition (Paul, 2007).
 238 Schreiner and Shorey (see Waksman, 1932) used available biochemical fractionation
 239 techniques, to isolate 40 identifiable materials including hydrocarbons, sterols, organic
 240 acids, aldehydes, carbohydrates, and organic P compounds (Stevenson, 1994). However,
 241 the amounts extracted were too small to assign a structure to SOM. Löhnis and Fred

(1923) suggested that old humus is richer in N and much of the CO₂ is derived from fresh litter with higher C:N ratios. Waksman and Tenney (1927) contributed greatly to the knowledge of residue composition and decomposition. Waksman (1932) gives detailed data on the chemical composition of various plant parts at different stages of decomposition and said, “This mass of undecomposed, partially decomposed, and transformed materials makes up soil organic matter which is being modified constantly.” He also noted the production of stable soil nitrogenous compounds from amino acids of the proteins in plants and stressed the importance of fungi in SOM formation.

3.2 Humus structure

Today, we again are looking at what constitutes humus-SOM (Schmidt et al., 2011; Chenu et al., 2015; Horwath, 2015). Piccolo (2001) suggested that humic substances are composed of supramolecular structures of self-assembling, relatively small, heterogeneous molecules. This involves weak dispersive rather than covalent bonding. He suggested that the large molecular weights, previously demonstrated by ultracentrifugation and gel chromatograph, were artifacts. Schnitzer and Monreal (2011) agreed, suggesting molecular weights of 6400 to 7800 arising from 755 to 950 atoms. These molecular weights place a restriction on the involvement of larger-molecular weight, aromatic structures, or proteins. Pisana et al. (2013) listed the average proportions of the major groups in SOM identified by nuclear magnetic resonance (NMR) as alkyl = 31%, O-alkyl = 39%, aromatic = 19%, and carbonyl = 10%. The long chain alkyl compounds increased linearly with an increase in MAT from 2 to 26°C.

263 Kononova (1961) and Schnitzer and Monreal (2011) argue for the occurrence of humic
264 materials in soils. They recognized that there is some uptake of O₂ during NaOH
265 fractionation, but stated that this can be eliminated by extraction under N₂. NMR data that
266 identifies primarily functional groups (Simpson et al., 2011) has been used to argue for a
267 non-humic nature of SOM (Feng and Simpson, 2011). These studies are usually
268 conducted on soils previously treated with HF that results in a loss of 10 to 30% of its
269 SOC (Chenu et al., 2015). The material lost is associated with the labile pool composed
270 largely of carbohydrates and N compounds, which are primarily responsible for soil
271 fertility (McLauchlan and Hobbie, 2004; Zegouagh et al., 2004).

272 The big difference between the views expressed by Kononova (1961) and Schnitzer and
273 Monreal (2011), relative to those of Schmidt et al. (2011), Simpson et al. (2011), and
274 Lehmann and Kleber (2015), involves the extent to which covalent bonds hold the
275 superstructure, humus molecule, or micelle together. Clapp et al. (2005) quote Burdon
276 who in 2001 wrote that there is no biochemical or chemical knowledge to suggest that
277 humic structures would be formed in the soil environment. On the other hand, Haider et
278 al. (1975) reference many articles where enzymatic and autoxidation of phenols to
279 quinones leads to covalent bond formation with amino acids. Many fungi and bacteria
280 were found to increase the reactions as were clays. Clemente and Simpson (2013) suggest
281 that organic protection by lignin or lignin-oxidation products against chemical oxidation
282 does occur. The hydrophobicity of the lipids is also a factor (Kögel-Knabner et al., 2008),
283 as are the stabilizing effects of cations (Stevenson, 1994). Some of the stable,

284 hydrophobic materials are often similar to plant waxes, cutins, and suberins (Simpson et
285 al., 2011; Baldock and Broos, 2012).

286 The extraction of humic compounds by alkali or resins has provided information on
287 molecular weights, optical and colloidal properties, and chemical composition
288 (Stevenson, 1994; Clapp et al., 2005). The words “humic compounds” tend to be
289 associated with specific types of extraction. “Humus” is a more useful general term. The
290 words “macromolecules,” “organic aggregates,” and “micelles” have been suggested as
291 other alternatives. In 1930, Schmook (see Waksman, 1938) described the nature and
292 structure of humus as mixtures of related compounds having similar structures, with N
293 coming from proteins as an integral part of the structure. Lignin was thought to be a
294 precursor of SOM after it was modified and interacted with microbial products (Paul and
295 Clark, 1986; Stevenson, 1994). Now lignin is considered to be decomposed in soil and
296 microbial products arising from the transformation of non-lignin plant residues are
297 considered more important (Cotrufo et al., 2013). However, lignin and the lignin-to-N
298 ratios are still considered to be major control agents in litter decomposition and
299 stabilization (Parton et al., 2015).

300 The nature of SOM is best understood through a knowledge of the controls on its
301 formation and dynamics (Figure 1). Chemical composition (recalcitrance), physical
302 protection, and matrix interactions (clay, silts, sesquioxides, and cations) are often cited
303 as the major controls on SOM formation and dynamics (Stevenson, 1994; von Lützow et
304 al., 2007). In many soils, silts comprised primarily of quartz and feldspar, contain one-

quarter to one half the amount of SOC per unit weight as the clay. Brady (1984) states that the silts also contain approximately one-third the sesquioxides and equivalent Ca and Mg as the clays, possibly in fine clay coatings of chemically-oxidized quartz and feldspars. Spatial effects and resource availability are important especially in the subsurface (Chabbi et al., 2009) and in aggregates of surface soils (O'Brien and Jastrow 2013; Segoli et al., 2013). Clays produce higher microbial biomass, stabilized microbial products (Sorensen, 1972; Amatto and Ladd, 1992), enzymatic interactions (Ladd and Butler, 1975; Sinsabaugh and Follstad Shah, 2012), and increased quinone formation (Haider et al., 1975).

A generalized structural diagram, adapted from Horwath (2015), represents present knowledge about the general composition of SOM including its interactions, bonding mechanisms, cations, and clay effects (Figure 2). The diagram illustrates the aromatic and aliphatic nature, the relationships among functional groups, and organo-mineral interactions. Near edge structure X-ray absorption, fine structure (NEXAFS), synchrotron near edge structure (XANES), and Fourier transform infrared (FTIR) showed interior pore regions of micro-aggregates where SOM is complexed with minerals (Gillespie et al., 2014; Horwath, 2015). Outer layers contain more oxidized C dominated by carboxyl and O-Alkyl functional groups. The preferential adsorption of amino compounds to clay surfaces (Nannipieri and Paul, 2009) results in enrichment of ^{15}N relative to ^{13}C or ^{14}C compounds (Sorensen, 1975). These compounds are thought to form a stable inner core to which other organics are sorbed (Sollins et al., 2006).

326 The role of the complex interactions of SOM in ecosystem functioning (Kögel-Knabner
327 et al., 2008; Cheeke et al., 2013) can best be established by a combination of
328 fractionation, incubation, and chemical characterization methodologies. Combining tracer
329 studies with observations at the μm scale with NanoSIMS made it possible to examine
330 the locations of these interactions (Chenu et al., 2015). This verified earlier electron
331 micrographs that cell debris, such as cell walls and mucilage, are often directly associated
332 with mineral particles (Foster and Martin, 1981; Campbell and Porter, 1982; Foster et al.,
333 1983). Pyrolysis-molecular beam, mass spectroscopy (py-MBMS) (Hoover et al., 2002)
334 of soil showed that the ~50% of the SOM that is pyrolyzed (Paul et al., 2011) contains: 5
335 to 10% aromatics and lignin monomers, 10 to 15% carbohydrates, 25 to 35% N
336 compounds, 7 to 10% lipids, and 5 to 10% phenols (Haddix et al., 2011).

337 Acid hydrolysis, together with ^{14}C carbon dating, can be used to determine the resistant
338 fraction that on average is 1400 to 1800 yr older than the total SOC (Leavitt et al., 1997;
339 Paul et al., 2006). The N of even the non-hydrolyzable fraction is largely amide in nature
340 (Plante et al., 2009). Mid-infrared spectroscopy (MiDIR) (Calderón et al., 2013) shows
341 the effects of soil minerals and adsorption and is useful for measuring the characteristics
342 of the LF and POM fractions, which are so important in C cycling. There are at least 10^6
343 natural products, including secondary metabolites of microbial origin (Alexander, 1961;
344 Waide, 1975; Bérdy, 2005). Schnitzer and Monreal (2011) emphasized the role of these
345 secondary microbial metabolites in SOM formation stating that abundant arylaromatic
346 products (polyketides) could form a significant core of the thousand-year old humic acids

that have been isolated from many soils (Campbell et al., 1967; Scharpenseel and Schiffman, 1977; Scharpenseel et al., 1989).

4. Turnover of specific SOM components

4.1 Historical perspectives

Measures of SOM dynamics could not be conducted until the advent of tracers when Norman and Werkman (1943) incorporated ^{15}N -labeled soybean residues into soil. Their mass spectrometer measurements of the stable isotope showed that 26% of the labeled N was recovered by a subsequent crop, with most of the tracer remaining in the soil. The growth of plant materials in a ^{14}C atmosphere and incorporation of the residues into soil in both laboratory and field studies made it possible to follow both the labeled and non-labeled SOC (Coleman and Fry, 1991). Thus, both the C and N components of SOM could be followed (Norman 1946; Broadbent and Norman, 1946; Broadbent, 1947).

The basic equations for tracer use (Broadbent and Bartholomew, 1948; Bingeman et al., 1953; Kirkham and Bartholomew, 1954, 1955) still apply today. These workers also redeveloped the original Löhnis concept of priming as the additional decomposition of native SOM in the presence of the added substrate. The word priming can be used in two contexts: 1) the use of ^{13}C , ^{14}C , or ^{15}N data to measure the increased decomposition (turnover) of the native SOM in the presence of the added substrate, and 2) the use of the tracer to measure not only the increased turnover of SOM, but also to calculate whether there was a true net change of soil C in the presence of an added substrate. This requires

367 consideration of the amount of added C or N remaining and involves the use of a mass
 368 balance equation to calculate the total SOM gains or losses.

369 The proportion of added residue remaining in various soils and climates is similar if the
 370 incubations have been carried out long enough to establish a steady state for that
 371 environment (Jenkinson, 1965, 1971, 1990). The addition of plant residues or microbial
 372 materials, and the measurement of their decomposition rates in soils (Mayaudon and
 373 Simonart, 1958, 1963), showed that the ^{14}C from both substrates rapidly entered all
 374 fractions, but to different extents, depending on that soil fraction's turnover rate.

375 There are a number of reviews that provide access to the original ^{14}C papers. Some of
 376 these include: Chahal (1968) "Biosynthesis and characterization of soil polysaccharides;"
 377 Wagner (1968) "Significance of microbial tissue to soil organic matter;" Finch et al.
 378 (1971) "The biochemistry of soil polysaccharides;" Marshall (1971) "Sorptive reactions
 379 between microorganisms and clay particles;" Haider et al. (1975) "Humus biochemistry;"
 380 Wagner (1975) "Microbial growth and carbon turnover;" Sorensen (1977) "Factors
 381 affecting the biostability of metabolic materials in soil;" Paul and van Veen (1978) "The
 382 use of tracers to determine the dynamic nature of organic matter;" Foster and Martin
 383 (1981) "*In situ* analysis of soil components of biological origin;" Jenkinson and Ladd
 384 (1981) "Microbial biomass in soil: measurement and turnover;" Stout et al. (1981)
 385 "Chemistry and turnover of naturally occurring resistant organic compounds in soil;"
 386 Ladd and Martin (1984) "Soil organic matter studies;" Dinel et al. (1990) "Soil lipids:
 387 origin, nature content, decomposition and effect on soil properties;" Smith and Paul

(1990) “The significance of soil microbial biomass estimates;” and Kögel-Knabner and Kleber (2012) “Mineralogical, physiochemical and microbiological controls on soil organic matter stabilization and turnover.”

4.2 Soil polysaccharides

Plant carbohydrates, comprised primarily of glucose from cellulose, and xylose and arabinose from hemi-cellulose, are the primary source of energy in soil (Chahal, 1968). The traditional method of measuring SOC polysaccharides, involving hydrolysis to constituent monomers, showed some xylose and arabinose, which indicated plant remains (Finch et al., 1971). Mannose, fucose, and rhamnose are widespread in fungi. The primary sugars in bacteria are glucose, fructose, mannose, galactose, rhamnose, and fucose. The production of these compounds can be used to estimate microbial growth, but may also accumulate in the microbial products, especially in extracellular slimes-biofilms (Redmile-Gordon et al., 2014). They also and react with humic substances (Cheshire et al., 1974). The role of polysaccharides in aggregation is well-established (Acton et al., 1963; Six et al., 2002a, Fry, 2015). The N-acetylglucosamine of chitin, in fungi cell walls, and N-acetylmuramic acid, of bacterial cell walls, can be used in specific compound, stable isotope probing to determine fungal to bacterial ratios (Gillespie et al., 2014; Kandeler, 2015).

4.3 Soil lipids

407 Lipids, such as free fatty acids and phospholipids, are of interest because specific lipids
408 (Wagner and Muzerewa, 1977) are associated with various microbial groups, and their
409 tracer contents can be readily measured (Vestal and White, 1989). The separation of ^{13}C -
410 labeled, fatty acid-phospholipids (PFLAs) thought to occur only in living cells (Kandeler,
411 2015) is fast, reproducible, and easily measured by GC-C-IRMS. Interpretation is limited
412 by the somewhat restricted number of peaks that identify major groups of organisms,
413 their variable occurrence in cells (Dinel et al., 1990), the somewhat erratic ^{13}C natural
414 abundance in different components (Pelz et al., 2005), and the difficulty in distinguishing
415 between plant and microbial fatty acids. The incorporation of the tracer indicative of
416 actively growing cells can be a useful adjunct to soil metagenomic-pyrotagged
417 sequencing (Myrold et al., 2013) to determine community composition (Bottomley et al.,
418 2012).

419 McMahon et al. (2005) following microbial community dynamics via PLFA analysis
420 after the addition of ^{13}C -wheat straw and incubation confirmed earlier observations that
421 Gr^- bacteria were most active when growing on soluble extract. The fungi, represented by
422 18:w6.9, were dominant in decomposition of insoluble residues. Labeled ergosterol can
423 be measured as a fungal marker (Kandeler, 2015). The determination of waxes and resins
424 that comprise 1 to 5% of SOC in mineral soils and up to 20% of organic deposits
425 (Wagner and Muzorewa, 1977) require pretreatments, such as NaOH hydrolysis before
426 GC analysis. pyMBMS (Hoover et al., 2002) can identify them directly. Waxes and
427 resins play special roles in the formation of SOM micelles and hydrophobicity (Kögel-

Knabner et al., 2008). Higher molecular weight lipids from cutins and suberins can differentiate root from leaf residue (Feng and Simpson, 2011).

4.4 Soil phenol-aromatics

The transformations of aromatics in SOM have received attention due to their similarity with what were thought to be humic constituents (Clapp et al., 2005). Labeled (^{14}C) cellulose decomposed four times as fast as labeled lignin even though they are physically related in plants (Crawford et al., 1977). Lignins are usually degraded by cometabolism and phenol-oxidase enzymes with little or no production of microbial biomass or incorporation into resistant SOM (Haider et al., 1975). During degradation, the aromatic component of lignin molecules is often broken up while still attached to a larger unit (von Lützow et al., 2007). Thus, lignin itself is not a major constituent of SOM, but is still considered of importance in modeling litter dynamics (Parton et al., 2015).

The brown-black color of humus has been related to light absorption at specific wavelengths (Kononova, 1961; Shields et al., 1968). White rot fungi, which completely degrade lignin and its byproducts, do not form colored compounds in soils (Haider et al., 1975). Brown rot fungi, which only partially degrade aromatics and attack side chains, produce many brown-colored intermediates (Haider, 1992; Stevenson, 1994) that may undergo further oxidation-reduction in soils (Horwath, 2015). Phenolic compounds and lipids are abundant in silts, whereas N components and carbohydrates tend to concentrate in the clays. This conclusion is supported by the observation that silts contain more non-hydrolyzable C than clays (Paul et al., 2006; Plante et al., 2009).

449 4.5 N compounds

450 Nitrogenous compounds comprise approximately 20 to 35% of the SOC (Bremner and
451 Hauck, 1982; Haddix et al., 2011). Amino sugars are important C and N constituents of
452 SOM, and if associated with resistant, melanic, fungal constituents from dark-colored
453 fungi, can be quite resistant to decomposition (Wagner, 1968, 1975). The C:N ratio of
454 plant residues varies from 250:1 to 20:1 depending on the plant and plant part. Fungi are
455 nearer to 10 to 15:1 and bacteria closer to 5:1. Soil organic matter, normally said to have
456 a C:N ratio of 10 to 13:1, approaches 8 to 10:1 when the plant residues in the light
457 fraction (LF) and particulate SOM (POM) are removed. Clays in cultivated soils
458 approach 6:1. Such a low ratio does not allow for the presence of much N-free
459 carbohydrates, lipids, or phenols-aromatics in the clay fraction (Paul et al., 2011).
460 However, the presence of fixed ammonia in the clay fraction can affect the C:N ratio of
461 this fraction (Jansson, 1958, 1963).

462 Earlier humic-formation theories (Kononova, 1961) postulated interactions of amino
463 acids with sugars through the Maillard reaction and with phenols through condensation
464 reactions (Haider et al., 1975; Horwath, 2015). Today's theories of SOM stabilization
465 postulate direct adsorption of amino acids to clays (Sollins et al., 2006; Yuan and Theng,
466 2012) and the general belief that most soil N is composed of proteinaceous-derived and
467 cell wall constituents (Stevenson, 1994; Di Costy et al., 2003; Nannipieri and Paul,
468 2009). Pyrolysis analysis (Schulten and Leinweber, 1993; Schulten, 1996) shows
469 significant concentrations of cyclic N monomers in soil. The possibility that these are

formed during analysis has been considered (Schulten and Leinweber, 1999); however, it is now believed that considerable quantities of these non-plant N compounds do occur in soil (Gillespie et al., 2014).

Derivatization of amino sugars allows for the separation and measurement of the compounds and associated tracers by gas chromatography-mass spectrometry (Kandeler, 2015; Bai et al., 2013). Liquid chromatography eliminates contamination of SOC with non-labeled C compounds associated with derivatization. Microbial products (Sorensen, 1975, 1977) rapidly enter what is known as the slow fraction of SOM with only a small amount being identified as microbial biomass. All of this helps explain why the modern literature still agrees with Löhnis and Fred (1923), that on average, 1% of the soil N becomes mineralized over a year's time (Robertson and Groffman, 2015).

5. Turnover of soil organic matter-humus constituents

5.1 Tracer applications

A further understanding of humus turnover in ecosystem dynamics and sustainability, global change, and soil fertility requires us to go beyond the simple gross turnover estimates of Table 1. Meaningful pool sizes and reaction rates are required (Manzoni and Porporato, 2009; Manzoni et al., 2012). The measurement of SOM fractions has proven successful when used together with: 1) modeling (Jenkinson and Rayner, 1977; Parton et al., 2015), 2) comparison of management techniques (Paul et al., 1997), 3) determination of pedogenic-soil forming factors (Scharpenseel and Schiffman, 1977; Trumbore, 1993),

490 4) depth (Stout et al., 1981; Goh, 1991), and 5) mineral interactions (Anderson and Paul,
491 1984; Torn et al., 1997).

492 The dynamics of field soils are best determined by two techniques: 1) carbon dating that
493 utilizes the ^{14}C produced in the atmosphere by solar radiation, or 2) C_3 - C_4 plant switches
494 that provide a ^{13}C signal. Radiocarbon (^{14}C) has a long half-life, and such measurements
495 are significant over periods of 100 to 50,000 yr (Stout et al., 1981). The production of ^{14}C
496 during hydrogen bomb testing produced a spike of ^{14}C beginning in 1954, peaking in
497 1963-1964, and gradually declining to where it is now immeasurable (O'Brien and Stout,
498 1978; Chenu et al., 2015). This has proven useful for the determination of turnover times
499 of SOC components with a rapid turnover, such as roots and litter (Trumbore, 2009).

500 Balesdent et al. (1988) demonstrated that the different discrimination against the naturally
501 occurring ^{13}C isotope during photosynthesis by C_3 and C_4 plants produced a usable signal
502 in the SOM if there was a C_3 - C_4 plant switch, such as the growth of C_4 maize on a
503 previous C_3 forested soil or the growth of C_3 wheat on a previous C_4 grassland (Boutton,
504 1996). Automatic instrumentation for measuring $^{13}\text{CO}_2$ evolution, and the measurement
505 of ^{13}C of specific compounds, such as fatty acids and chitin, are making rapid analysis of
506 SOM dynamics and the identification of specific SOM constituents in stable isotope
507 probing possible (Chen et al., 2012; Kandeler, 2015).

508 The use of mass spectrometers coupled to pyrolysis units has shown that the ^{13}C content
509 of the pyrolysis products mirror their nonvolatile precursors allowing this technique to be
510 used to determine the dynamics of soil constituents (Gleixner, 2013). Another use of ^{13}C

occurs when the DNA of soil organisms is exposed to a ^{13}C substrate. The heavy DNA produced by the incorporation of the isotope is separated by ultracentrifugation and analyzed for microbial diversity using nucleic acid, molecular techniques (Thies, 2015). Another ^{13}C opportunity occurred when CO_2 derived from fossil fuels, thus having a different ^{13}C signature, was used in free atmosphere CO_2 enhancement (FACE) experiments under field conditions (Leavitt et al., 1997; Norby and Zak, 2011). Controls on decomposition and SOM stabilization (Figure 1), such as soil texture, plant type, moisture, and N availability, have been shown to have interactive effects on the SOC dynamics of different sites (Hungate et al., 1996; Leavitt et al., 2001).

The turnover dynamics measured with tracers are dependent on the length of the tracer exposure. MRTs measured in laboratory incubations that are limited to a few years are shorter than those measured in the field where the $\text{C}_3\text{-C}_4$ plant switch is usually measured in decades (Collins et al., 2000). Carbon dates measure SOM dynamic of constituents, which can be millennia in length. However, they also reflect shorter term disturbance, such as management (Campbell et al., 1967, Mathieu et al., 2015). Paul et al. (2006) related field ^{13}C measurements that averaged 20 yr of tracer exposure to ^{14}C carbon dates from surface and depth measurements made on five soil types in the US Corn Belt to obtain the following equation:

$$^{14}\text{C MRT} = 176(^{13}\text{C MRT})^{0.54} \quad (1)$$

This equation has the potential to relate the extensive number of ^{13}C field measurements to their longer, much more difficult to obtain carbon-dating equivalents. The length of the

¹³C field or laboratory exposure should be considered if different from the 20yr used to establish this equation.

5.2. Soil fraction dynamics

An understanding of SOC-fraction dynamics is best achieved by examining fraction-tracer data from different soils. A Canadian, Chernozemic (Mollisolic) soil with a mean annual temperature (MAT) of -2°C and 5.6% SOC had an overall MRT of 870 yr (Table 3). In the absence of a C₃-C₄ vegetation switch in this cool climate, the data for the LF and POM fractions were derived from SOC-fraction changes during long-term incubation (Paul et al. 2004) and comparison to sites with ¹³C data (Gregorich et al., 1995; Haile-Mariam et al., 2008). The light fraction (LF), comprising intra-aggregate plant residues (Hurisso et al., 2013), was the youngest fraction with an MRT of 3 yr. The inter-aggregate particulate organic matter (POM) comprising the other portion of fairly decomposable SOC (Jastrow, 1996; Jastrow and Miller, 1998) had an MRT of 12 yr. The MRT of the humic acid fraction at 1235 yr. was much older than fulvic acids at 470 yr showing that this fractionation does differentiate on an MRT basis (Martel and Paul, 1974 a, b). Hydrolysis of the old humic acids increased the MRT by 165 yr, while producing a soluble fraction of 25 yr (Campbell et al., 1967), showing that although the humic acid was an old, measurable fraction, it contained a small proportion of young SOC (Paul et al., 2006). The coarse clay comprised 31% of the SOC and dated 1255 ± 80 yr. The Ca humates isolated with by humic fractionation after removal of Ca were similar in MRT to the nonhydrolyzable humic acids (Table 3). The fine clay, comprising 8% of the soil

553 mass, adsorbed microbial metabolites and dated only 170 ± 50 yr (Anderson and Paul,
554 1984).

555 Analysis of SOC dynamics for a warmer grassland (MAT of 9°C) with 1.1% SOC
556 included vegetation comparisons, long-term incubation, and ^{14}C and ^{13}C together with
557 fractionation (Table 4). The SOC had a field MRT of 1015 yr; the 60% of the SOC that
558 was non-hydrolyzable (Poirier et al., 2005) had an MRT of 2658 yr (Follett et al., 2007).
559 In spite of the great MRT of both the whole soil and the non-hydrolyzable fraction, this
560 soil evolved 33% of its SOC during an 842-day incubation. The LF and some of the
561 POM, in addition to the labile microbial biomass, are responsible for much of the
562 mineralizable C of this soil. This soil also has a low level of aggregation, which has been
563 shown in other soils to protect the labile pools (Mikha et al., 2013). Measurement of
564 microbial biomass kinetics (Follett et al., 2007; Paul et al., 2011) showed that 1.2% of its
565 SOC occurred in a labile microbial fraction with an MRT of 0.6 yr; another 2.3% of the
566 SOC was stable microbial biomass with a field equivalent MRT of 14 yr. The drop in the
567 microbial biomass C accounted for 24% of the mineralized C during the first 172 days of
568 incubation. From day 332 to 842, the drop in biomass contributed just 6% of the
569 mineralized C. The LF and POM of this soil, together representing 19% of the SOC,
570 contained some prairie-derived SOC attributable in part to their 7% pyrogenic C
571 (charcoal) (Lavalley et al., 2015).

572 The data from Table 4 can be used to test Equation 1. Conversion of the ^{13}C data for the
573 total soil to a ^{14}C field-equivalent via this equation yielded a MRT of 2435 yr (data not

shown) relative to the measured carbon date of 1020 yr. The average of silt and coarse clays with a calculated, ^{13}C equivalent age of 2565 and 3478 yr, respectively, was similar to the carbon date of the non-hydrolysable C at 2658 yr. This badly degraded soil had a very old, fine clay fraction as measured by ^{13}C . Equation 1 was developed on soils with an average of a 20 yr $\text{C}_3\text{-C}_4$ switch rather than the 100 yr switch on this site. This confirms earlier comments that equation 1 and all other tracer measurements should be interpreted relative to the time of the experiment.

A third study involved acid hydrolysis, ^{14}C dating, and ^{13}C measurement of the C_3 and C_4 dynamics using data from long-term incubation. The native, Ohio Mollisol was developed under a poorly drained, maple forest. The cultivated, continuous corn on this site had 1.96 % SOC with an MRT of 920 yr (Table 5). The 45% of the SOC that was non-hydrolyzable at 1770 yr MRT was assumed to be the resistant SOC pool for curve fitting purposes. The active pool accounted for 3% of the SOC with an MRT of 91 days. Fractionation and ^{13}C analysis (Table 5) showed that the light fraction, with a C:N ratio of 20 (Haile-Mariam et al., 2008), accounted for 3.5% of the SOC at the initiation of the incubation, with 84% of its C derived from corn. The slow pool (Gregorich et al., 1995, 1996), with 52% of the SOC and an MRT of 45 yr, is the seat of soil fertility. The POM plus sand, had 56% of its C originating from corn, and had a corn-derived $\text{C}_4\text{-MRT}$ of 7.8 yr. The non-corn C_3 of this fraction was five times as old. Black carbon (Krull et al., 2006; Cheng et al., 2008) constituted 3.9% of the C of this fraction, with a ^{13}C signature of -24‰, showing that it was derived primarily from the original forest vegetation (Lavallee et al., 2015). The silt had more of its C derived from corn than did the clay

596 fraction. The C:N ratios decreased from LF at 20:1 to clay at 6.5:1 showing the effects of
597 decomposition and the importance of proteinaceous material stabilized by clay. Bradford
598 reactive soil protein or glomalin (Rillig et al., 2003), said to be formed by arbuscular
599 mycorrhizal fungi or extracted as humic substances (Schindler et al., 2007), constituted
600 5% of the SOC (Haile–Mariam et al., 2008) and contained only 10% C₄-corn C.

601 The MRT of the SOC of this soil increased by a factor of 5 or more at depth
602 (Scharpenseel et al., 1989; Paul et al., 2001a) (Table 6). The MRT of the residue of acid
603 hydrolysis doubled at depths for all soils showing that the procedure removed significant
604 amounts of young C. However, the amount of non-hydrolyzable SOC decreased with
605 depth, confirming previous observations (Martel and Paul, 1974a, b), that factors other
606 than recalcitrance are involved in the great age. The observations (Chabbi et al. 2009;
607 Rumpel et al., 2011; Dungait et al., 2012) that much of this great age is not due to
608 recalcitrance, but rather inaccessibility, are supported by the similar percentage of SOC
609 (Table 6) that is released by mixing-incubation of the subsurface and surface soils
610 (Collins et al., 2000). Mathieu et al. (2015) through a meta-analysis of carbon dates
611 concluded that while the age of topsoil SOC is controlled primarily by climate and
612 cultivation, the age of deeper horizons is dependent on pedological traits, such as clay
613 content and mineralogy that influence accessibility.

614 Lajtha et al. (2014), in a similar study, used ¹³C and ¹⁴C together with incubation and
615 density fractionation of forest and restored grassland plots. Root and litter inputs on their

prairie site appeared to have similar effects on SOC levels. The estimates of labile C were similar between control and litter plots.

The above data shows that although a number of fractionation tracer techniques are required to characterize all fractions there is a consistency between soils and fractions. The subsurface soils are much older and have different controls than the surface. There also is agreement between these data and other literature. This all gives support to the hope that actual pools and fluxes can be used for modeling purposes to better understand ecosystem dynamics and responses to global change.

6. The role of microbial products in SOC formation and dynamics

Microbial products have long been considered to be the major components of SOM (Löhnis and Fred, 1923; Waksman, 1932; Wagner, 1975; Miltner et al., 2011), yet we don't fully know how they are produced or where and how they are stabilized (Cotrufo et al., 2013). Measurement of biomass size changes, during long-term laboratory incubation (Table 4), showed that 35% of the original biomass had a field-adjusted MRT of 0.6 yr. The other 65% was old with a field equivalent MRT of 14 yr. The microbial biomass C at 3.8% of this SOC is in the upper range of 2 to 5% found for most soils (Schnurer et al., 1985; Smith and Paul, 1990). The microbial biomass, as a percent of the SOC, increases with soil quality as affected by manure, compost additions, or rotations, including legumes (Anderson and Domsch, 1989). Also, the proportion of the biomass that is labile with incubation is more sensitive to soil quality than the total biomass (Alvarez et al., 1995; Follett et al., 2007). Other factors affecting the biomass include the clay content and pH

values. There is both a greater biomass and a greater carbon utilization efficiency (CUE) in clay than sandy soils (Amato and Ladd, 1992). Also, acidic soils have different decomposition kinetics (Jenkinson and Ladd, 1981) and a different community composition (Fierer et al., 2009) than those with higher pH values. Fungi (especially mycorrhizal fungi) are more common in litter layers, at the soil surface and on the surface of aggregates (Fry, 2015). Bacteria and archaea are more often found at depth and soil biota are restricted to pores $>0.8 \mu\text{m}$, so only 20 to 50% of the soil pore volumes are available for decomposition.

The experiments of Oades and Wagner (1970), Jenkinson (1971), Martin et al. (1974), and Sorensen (1972, 1981), and reviews (Webley and Jones, 1971; Paul and van Veen, 1978), delineate the principles affecting microbial growth and its effect on SOM formation. These established that: 1) the level of substrate addition does not appreciably affect the amount remaining, 2) priming of *in situ* SOM at times does occur, 3) acid hydrolysis is superior to humic fractionation for separating differentially labeled constituents, but not satisfactory for determination of pool sizes, 4) the label rapidly enters all soil fractions, and 5) different temperatures of incubation, while following Arrhenius (Q_{10}) kinetics, do not appreciably alter the amount of added substrate remaining after steady state is achieved. These observations have been confirmed in the more recent experiments on the effects of global warming (Alvarez et al., 1995; Steinweg et al., 2008; Conant et al., 2011).

657 An extensive series of experiments, with ^{14}C - labeled plant residues, glucose, and
658 cellulose that were incubated for up to 6yr (Sorensen, 1972, 1975, 1977, and 1981)
659 showed that ^{14}C cellulose produced less amino acid ^{14}C than that added as ^{14}C glucose.
660 Due to the recycling of ^{15}N in soil during microbial growth, and the preferential retention
661 of amino compounds in clays, ^{15}N compounds have greater MRTs than ^{14}C labeled
662 compounds. Sorensen and Paul (1971) found that after a 200-day incubation with ^{14}C and
663 ^{15}N , the remaining ^{14}C had an MRT of 3.2 yr compared to a 6.4 yr MRT for ^{15}N . Some of
664 the ^{14}C and ^{15}N was associated with fine clay, nano-sized ($<0.04\mu\text{m}$) particles (McGill et
665 al., 1975). Enzyme production (Ladd and Paul, 1973), estimated at 2 to 5% of the
666 microbial biomass (Sinsabaugh et al., 2013), could be one of the components of the
667 microbial products. Plants and soil fauna are also sources of soil enzymes (Nannipieri and
668 Small, 2006; Gianfreda et al., 2012).

669 Further indication of the role of microbial products comes from a 7 yr field study
670 involving glucose and wheat straw additions (Voroney et al., 1989). Although the
671 decomposition rate of the glucose was faster initially, a greater amount of stable SOC
672 was formed from the ^{14}C glucose than from the ^{14}C wheat straw (Figure 3) with a field
673 decomposition rate constant $T_{1/2}$ of 7.9 yr for the glucose-derived ^{14}C , corresponding to
674 an MRT of 11.5 yr.

675 Visual proof of associations between bacterial cell components with clays and humified
676 materials was demonstrated by Foster and Martin's (1981) electron, microscope photos.
677 Miltner et al. (2011) provided further visual examples in arguing for a significant

contribution of microbial biomass to SOM formation. Clay-SOM interactions are usually associated with adsorbed humus, sesquioxides, and cations (Yuan and Theng, 2012) that affect the soil biota, its substrate, and extracellular enzymes (Marshall, 1971; Gianfreda et al., 2012; Kögel-Knabner and Kleber, 2012).

Much of the biomass is at a resting state at any one time. Paul and Voroney (1980) added 1600 $\mu\text{g } ^{14}\text{C}$ as glucose to soil in a laboratory incubation. At the end of day one, there was no measurable glucose, but 950 $\mu\text{g } ^{14}\text{C}$ remained in microbial cells, showing 60% CUE (Figure 4). The original ^{12}C biomass lost 12% of its C in the presence of the labeled glucose with a decay rate of $1.27 \times 10^{-1} \text{ day}^{-1}$. The decay rate constant of the non-active pool of the unlabeled biomass in the presence of glucose at $5.5 \times 10^{-3} \text{ t}$ was equivalent to an MRT ($1/k$) of 182 days. The labeled biomass could be separated into three pools. The active component accounted for 40% of the glucose-C immobilized and decayed at a rate constant of 1.2 day^{-1} . The smaller, second component had a decay rate of $9.08 \times 10^{-2} \text{ day}^{-1}$. The third component at 534 $\mu\text{g } ^{14}\text{C}$ had a decay rate of $5.5 \times 10^{-3} \text{ t}$. This was equivalent to the biomass decay rate of the unlabeled soil. Others confirm that only 2 to 14% of the total microbial biomass is active at any one time (Stursova et al., 2012).

Further information on organism growth and organic matter interactions was shown in a study where long-term incubation reduced the mineralizable SOC by 18 to 20% with a 50% drop in the chloroform-extractable biomass (Birge et al., 2015). The respiration of CO_2 from added corn or wheat residue, however, was the same (~40%) in the fresh and the biomass-depleted soil. The insensitivity of decomposition to microbial biomass size

699 also has been demonstrated where chloroform treatment for microbial biomass
700 measurements, killed 99.9% of the culturable biomass (Shields et al., 1973; Kemmitt et
701 al., 2008). Reinoculation was not required to achieve decomposition with mineralization of
702 ~45% of the killed biomass-C during 10 days at 25°C (Jenkinson and Ladd, 1981; Smith and
703 Paul, 1990; Hurisso et al., 2013).

704 The above data is relevant to questions about how microbial biomass estimates should be
705 incorporated into modeling SOM dynamics relative to microbial growth (Sinsabaugh and
706 Follstad Shah, 2012), CUE (Sinsabaugh et al., 2016), and enzyme kinetics (Schimel, 2001) or
707 first order dynamics (Parton et al., 2015). It also gives support to the Kemmitt et al. (2008)
708 hypothesis that physical characteristics, not the microbial biomass, controls decomposition in
709 soils. Also to be considered is the possibility that external enzymes independent of the
710 biomass are contributing to decomposition (Nannipieri and Small, 2006; Gianfreda et al.,
711 2012.)

712 The interactions of organism growth, organic matter, and tracers have recently become
713 more tightly fused through the use of stable-isotope probing (SIP) (Chen et al., 2012).
714 The incorporation of substrate-¹³C through added labeled compounds into the
715 rhizosphere, mycorrhizal fungi, or plant litter can now be used to: 1) measure the
716 relationship between CUE and SOM formation (Sinsabaugh et al., 2016), 2) identify
717 community composition through DNA and RNA-SIP, gene expression (Pietramella et al.,
718 2012), as well as PLFA formation (Myrold et al., 2013), and 3) movement of plant
719 residues and microbial products into and through soil humus (Kandeler, 2015; Thies,
720 2015). Organisms in the β- and γ- subgroups of Protobacteria (Killham and Prosser,

2015) utilize soluble constituents and are representative of Winogradsky's zymogenous bacteria (Waksman, 1932; Alexander, 1961). Phyla, such as Actinobacteria, Planctomycetes, and Firmicutes, are involved in complex molecule decomposition and are therefore representative of Winogradsky's autochthonous organisms. The Ascomycota fungi represent the dominant fungal forms and are also active decomposers. *Fusarium* which is often associated with plant diseases (Taylor and Sinsabaugh, 2015) has been shown to be a major straw decomposition organism (Chattopadhyay et al., 2016).

7. Modeling in soil organic matter and tracer research

Models organize and analyze data, test and develop concepts, predict future processes, and expand understanding across scales (Campbell and Paustian, 2015). They are central to the understanding of soil organism-organic matter interactions, which is why there are >250 published models (Manzoni and Porporato, 2009). The publications by Smith et al. (1997), Schimel (2001), Manzoni et al. (2012), Moore and de Ruiter (2012), Sinsabaugh and Follstad Shah (2012), and Parton et al. (2015) give background information and comparisons between models. These models explore a variety of topics including: substrate-enzyme-microbe interactions (Sinsabaugh et al., 2016), food web-faunal effects (Osler and Sommerkorn, 2007), and global climate change (EPA, 2015).

Manzoni and Porporato (2009) note that in 1936, Nikiforoff suggested that humus formation involved pools with different turnover times, although the term "active organic matter" was used much earlier (Waksman, 1932). Jenny (1941) restated Dokuchaev's

five soil forming factors of climate, parent material, topography, organisms, and time in an equation format. Jenkinson and Rayner (1977) used long-term Rothamsted field, ^{14}C addition, and carbon dating data to publish a simulation model that described organic matter turnover. Their model, now adapted as Roth C (Jenkinson, 1990), used an inert fraction that was included to help the model describe the high ^{14}C MRTs found in SOM, especially at depth. Previous discussions in this and other publications point out that there is no inert pool, not even black carbon is inert (Kögel-Knabner et al., 2008; Lavallee et al., 2015).

The McGill et al. (1975) model included a sub-model that described the transformation of microbial biomass. The van Veen and Paul (1981) model that used first order kinetics, with a microbial biomass pool, recognized the effects of physical protection by aggregates and clay-associated materials. The values for the effects of protection were determined by comparing the decomposition of amino acids in the presence and absence of clays. The effects of cultivation were modeled by using long-term climate data and alteration of the physical protection factors due to disturbance and erosion (Voroney et al., 1981). Aggregates are an important part of physical protection (Six et al., 2002a). Segoli et al. (2013) utilized data from incubation experiments to calculate and validate biological and environmental influences on the rates of formation and breakdown of micro and macroaggregates. Their model estimated the turnover rate for macroaggregates at 31 days and 181 days for microaggregates. These were laboratory incubations, the turnover rates measured in the field with tracers would be much longer (Collins et al., 2000).

Biological decomposition is enzyme mediated; therefore, one must ask how can a SOM decomposition model not include them? However, 50% of the models listed by Manzoni and Porporato (2009) are based on first order kinetics. Some models, such as Century (Parton et al., 1987, 2015), use first order kinetics with an implicit recognition of microbial enzymes in the decay functions for the various pools. Moisture and temperature are among the major control variables of all the models tested. Their overriding importance in decomposition (Adair et al., 2008), together with the extensive mathematical fine tuning of decomposition rates, have helped many of the models to be reasonably predictive of natural processes.

Extensive field and laboratory information is required to obtain actual pools and fluxes. Paul et al. (1999) inserted analytically-derived, kinetic pools and fluxes into the daily version of the Century model (Parton et al., 1998, 2015) to simulate daily field CO₂ evolution rates over a season in a number of cropping systems. Long-term incubations (Collins et al., 1999), ¹³C estimates of field turnover rates (Collins et al., 2000), together with acid hydrolysis (Leavitt et al., 1997) and carbon dating (Paul et al., 2001a) provided an estimate of the active slow and resistant (passive) pools. The model accurately predicted field-measured CO₂ evolution rates on a daily basis over the growing season. It overestimated CO₂ fluxes after harvest in the fall. This was attributed to the fact that the model did not consider a lag period during which the above-ground residues lose their hydrophobicity, and are comminuted and colonized by the soil biota before the residue is in intimate contact with the soil biota and its enzymes.

The extensive studies and models on organism-enzyme interactions (Weintraub and Schimel, 2003; Sinsabaugh and Follstad Shah, 2012) usually involve Michealis-Menten kinetics (Schimel, 2001). The amount of substrate turned into microbial products is central to SOM dynamics (Cotrufo et al., 2013, Sinsabaugh et al., 2016) and can be handled directly (Sinsabaugh et al., 2013) or indirectly as in the decomposition rate constants in first order models (Jenkinson and Rayner, 1977; Parton et al., 1987). Biotic information is most often included in food webs (Zwart et al., 1994; Coleman et al., 2004) and can become complicated if not process-function based (Moore and de Ruiter, 2012).

In addition to modeling, we need to ensure our data helps us interpret information on ecosystem dynamics. We can test our knowledge about organism biomass activity and SOM dynamics by relating it to measured field respiration rates (Fisher, 2004) and ecosystem productivity. The CO₂ flux of 5.6 Mg C ha⁻¹ yr⁻¹ released from poplar soils (Figure 5), relative to 4.2 Mg C ha⁻¹ yr⁻¹ for a corn-soybean rotation (Hamilton et al., 2015) reflects the higher NPP in the poplar plantation. The higher levels of SOM on the poplar site altered the kinetically-determined sizes and dynamics of the active and slow pools. (Collins et al., 1999; Paul et al., 1999, 2001b). The figure in tying the various sections of this review together points out the relationship between plant growth, microbial respiration, SOM pool sizes, and even the effects of symbiotic organisms. The data for the contributions of the fauna, bacteria, fungi, and roots (Figure 5) are based on the knowledge of biomass size obtained via microscopy and chloroform-fumigation (Horwath and Paul, 1994) and knowledge of microbial growth determined by ¹⁴C

thymidine incorporation (Harris and Paul, 1994). The numbers reflect the increase in fungal biomass in the poplar plantation as the trees switched from a vesicular arbuscular to an ectotrophic mycorrhizal symbiosis at ~5 yr in age, which affected the underground C transfers (Horwath et al., 1994).

8. Looking forward

Plants, as ecosystem C pumps, produce the substrate for soil biotic activity and SOM formation. Over the long term, net primary production must slightly exceed decomposition for SOM to accumulate. Even though much is known about plant-soil - biota controls (Balestrini et al., 2015), their interactive effects make it difficult to predict individual ecosystem effects (Hinsinger et al., 2012). These effects include: 1) the range in amounts and chemistry of above and below-ground plant C allocation with plant type, soil depth and season, 2) the ability of plants to compensate, in part, for the needs of symbiotic partners, 3) the soil drying effects on the soil biota and the mass movement of DOC as the plants withdraw water, and 4) the interactions with root-associated organisms including food webs.

The role of non-culturable organisms and genes can now be better understood with the finding of huge reservoirs of previously unknown genetic material (Pietramella et al., 2012; Taylor and Sinsabaugh, 2015). Physiological, gene expression, enzymatic, and food web data are also needed to gain a better understanding of microbial products in humus formation. The role of resting organisms (Jenkinson and Ladd, 1981), maintenance energy (Babiuk and Paul, 1970), enzyme stabilization (Sinsabaugh et al.,

2013), priming (Kirkham and Bartholomew, 1954), CUE (Sinsabaugh et al., 2016), and how microbial products are stabilized (Miltner et al., 2011), must be better understood. Compound specific isotope studies (Chen et al., 2012; Gleixner, 2013) and tracer dynamics (Chenu et al., 2015), conducted together with enzyme (Sinsabaugh and Follstad Shah, 2012; Kandeler, 2015) and microbial community studies (Thies, 2015), will prove most useful in addressing these questions.

The concepts discussed in this review have implications for soil ecosystem functioning, nutrient cycling, pollution control, biogeochemistry, feeding an expanding global population and global change. Additional areas where more information is needed include: the timing and distribution of above- and beneath-ground plant inputs, residue and SOM composition (Chenu et al., 2015), temporal and spatial scaling (Hinckley et al., 2014), and microbial community composition (Kong et al., 2011). We are still not able to reliably control the effects of root-associated diseases, the ability of plants to compensate for C requirements of symbionts, or the effects of competition in attempts to inoculate desirable organisms into soil.

Microbial parameters still not well understood include CUE relative to SOM formation, maintenance energy of soil organisms, carbon overflow, and polysaccharide excretion under nutrient deficiency and cometabolism. We are finally able to characterize the vast numbers of organisms in soil. But, how important is community diversity? How can we express soil biota activities and interactions relative to food webs, predators, and parasites, including relative unknowns such as viruses? There is a great deal of microbial-

enzyme data that is still difficult to interpret, such as: the role of resting soil populations, gene control by induction and inhibition, external enzyme diffusion and stoichiometry, degradation of and inhibition by specific organics, as well as the Arrhenius-based activation energy effects which are now being investigated in regard to increased temperatures across the globe.

Hopefully, this review helped identify the extensive historical literature, especially that on tracers, which can further develop future basic concepts about decomposition, microbial growth, and microbial product formation relative to SOM dynamics. The case studies were included to show the usefulness of multiple data sets. Some of the available information discussed in this review includes:

- 1) The combination of methods, including tracer additions, long-term incubation, ^{14}C carbon dating, and ^{13}C natural abundance measurements, which have helped to develop kinetically-defined SOC pool sizes and turnover rates.
- 2) Physical fractionation, especially when combined with tracers, to provide pool size and turnover information on actual SOM fractions, such as the LF, POM, silt, and clay.
- 3) All SOC pools contain young and old constituents due to their dynamic nature.
- 4) Physically isolated fractions reflect decomposition stages with decreasing C:N ratios and increasing MRT as the fractions move from the LF to the clays.
- 5) In some soils, fine clays are associated with recent microbial products and have low MRTs; in other soils this is not the case.

- 870 6) Much of the pyrogenic or black carbon is associated with the silt and clay-sized
871 pools and has high MRTs, but it is not inert.
- 872 7) Tracer measurements are dependent on the time length of exposure. Data should
873 be expressed relative to the length of exposure. The empirically-derived equation
874 that relates ^{14}C dates to field-determined ^{13}C rates requires more validation at
875 different time scales to relate data from short-term experiments to the pedological
876 controlled values that occur in the field.
- 877 8) Silt-sized materials have less SOC per unit weight than clays. Their MRTs are
878 often slightly lower than those of clays. We still don't have definitive knowledge
879 concerning the mechanisms of SOC protection by silt-sized particles.
- 880 9) Since N is a strong controller of SOC dynamics, further integration of the data
881 from both C and N tracers would be very useful.
- 882 10) The soils at depth are thousands of years older than those found at the surface, yet
883 when brought to the surface, they often are fairly rapidly decomposed indicating
884 that decomposition at depth is heavily controlled by spatial effects-resource
885 availability.
- 886 11) All soils have similar controls allowing for the development of overall concepts,
887 modeling, and extrapolations to larger landscapes. The expression of the controls
888 at different sites results in the individual site characteristics.

889

890 Acknowledgements. This review was conducted with funding from the NSF-LTER
891 OPUS Grant #DEB-1256754 entitled “Soil organisms, carbon and nitrogen interact in the
892 control of soil organic matter dynamics in ecosystem functioning.” I greatly appreciate
893 the interactions with the many past and present excellent students and coworkers for the
894 ideas and scientific findings that are so necessary for us to advance our understanding of
895 SOM formation and dynamics. Especially appreciated is the excellent editing by Laurie
896 Richards and the very helpful reviews.

897   **References**

- 898   Acton, C.J., Rennie, D.A., Paul, E.A., 1963. The relationship of polysaccharides to soil
899   aggregation. Canadian Journal of Soil Science 43, 201-209.
- 900   Adair, E.C., Parton, W.J., del Grosso, S.J., Silver, W.L., Harmon, M.E., Hall, S.A.,
901   Burke, I.C., Hart, S.C., 2008. Simple three pool kinetic model describes patterns of long-
902   term litter decomposition in diverse climates. Global Change Biology 14, 2636-2660.
- 903   Aitkenhead, J.A, McDowell, W.H., 2000. Soil C:N ratio as a predictor of annual riverine
904   DOC flux at local and global scales. Global Biogeochemical Cycles 14,127-138
- 905   Alexander, M., 1961. Introduction to Soil Microbiology. Wiley, New York.
- 906   Alvarez, R., Santanatoglia, J.J., Garcia R., 1995. Effect of temperature on soil microbial
907   biomass and its metabolic quotient *in situ* under different tillage systems. Biology and
908   Fertility of Soils 19, 227-230.
- 909   Amato, M., Ladd, J.N., 1992. Decomposition of ¹³C labeled glucose and legume
910   materials in soils: Properties influencing the accumulation of organic residue and
911   microbial biomass. Soil Biology and Biochemistry 23, 455-464.
- 912   Anderson, D., Paul, E.A., 1984. Organo-mineral complexes and their study by
913   radiocarbon dating. Soil Science Society of America Journal 48, 298-301.
- 914   Anderson, J.F., Domsch, K.H., 1980. Quantities of plant nutrients in the microbial
915   biomass of selected soils. Soil Science 130, 211-216.

916 Anderson, J.F., Domsch, K.H., 1989. Ratios of microbial biomass C to total organic C in
 917 arable soils. *Soil Biology and Biochemistry* 21, 471-480.

918 Babiuk, L.A., Paul, E.A., 1970. The use of fluorescesin isothiocyanate in the
 919 determination of the microbial biomass of grassland soils. *Canadian Journal of*
 920 *Microbiology* 16, 57-62.

921 Bai, Z., Bode, S., Huygens, D., Zhang, X.D., Boeckx, P., 2013. Kinetics of amino sugar
 922 formation from organic residues of different quality. *Soil Biology and Biochemistry* 57,
 923 814-821.

924 Baldock, J.A, Broos, K., 2012. Soil organic matter. In: Huang, P.M., Li, Y., Sumner, M.
 925 (Eds.), *Handbook of Soil Sciences: Properties and Processes*, second ed. CRC Press,
 926 Boca Raton, pp. 11-1 to 11-52.

927 Balesdent, J., Wagner, G.H., Mariotti, A., 1988. Soil organic matter turnover in long-
 928 term field experiments as revealed by carbon-13 natural abundance. *Soil Science Society*
 929 *of America Journal* 52, 118-124.

930 Balestrini, R., Lumini, E., Borriello, R., Bianciotto, V., 2015. Plant-soil Biota
 931 Interactions. In: Paul, E.A. (Ed.), *Soil Microbiology, Ecology and Biochemistry*, fourth
 932 ed. Academic Press, San Diego, pp. 311-338.

933 Bardgett, R.D., 2005. *The Biology of Soil. A community and ecosystem approach.*
 934 Oxford University Press, London, UK

- 935 Batjes, N.H., 2014. Total carbon and nitrogen in the soils of the world. *European Journal*
936 *of Soil Science* 65, 10-25.
- 937 Bérdy, J., 2005. Bioactive materials. *The Journal of Antibiotics* 58, 1-26.
- 938 Bingeman, W., Varner, J.E., Martin, W.P., 1953. The effect of the addition of organic
939 materials on the decomposition of an organic soil. *Soil Science of America Proceedings*
940 17, 34-38.
- 941 Birge, H.E., Conant, R.T., Follett, R F., Haddix, M.L., Morris, S J., Snap, S.S.,
942 Wallenstein, M.D., Paul, E.A., 2015. Soil respiration is not limited by reductions in
943 microbial biomass during long-term incubation. *Soil Biology and Biochemistry* 81, 304-
944 310.
- 945 Bottomley, P.J., Taylor, A.E., Myrold, D.D., 2012. A consideration of the relative
946 contributions of different microbial subpopulations to the soil N cycle. *Frontiers in*
947 *Microbiology* Vol 3, 1-7.
- 948 Boutton, T.W., 1996. Stable carbon isotope ratios of soil organic matter and their use as
949 indicators of vegetation and climate change. In: Boutton, T.W., Yamasaki, S. (Eds.),
950 *Mass Spectrometry of Soils*. Marcel Dekker, New York, pp. 47-82.
- 951 Brady, N.C., 1984. *The Nature and Properties of Soils*, ninth ed. Macmillan, New York.

952 Bremner, J.M., Hauck, R.D., 1982. Advances in methodology for research on nitrogen
 953 transformations in soil. In: Stevenson, F.J. (Ed.), Nitrogen in Agricultural Soils,
 954 American Society of Agronomy, Madison, pp. 467-501.

955 Broadbent, F.E., 1947. Nitrogen release and carbon loss from soil organic matter during
 956 decomposition of added plant materials. Soil Science Society of America Proceedings 12,
 957 246-249.

958 Broadbent, F.E., Norman, A.G., 1946. Some factors affecting the availability of organic
 959 nitrogen in soil. A preliminary report. Soil Science Society of America Proceedings 11,
 960 359-363.

961 Broadbent, F.E., Bartholomew, W.V., 1948. The effect of quantity of plant material
 962 added to soil on its rate of decomposition. Soil Science Society of America Proceedings
 963 13, 271-274.

964 Bruun, S., Six, J., Jensen, L.S., Paustian, K., 2005. Estimating turnover of soil organic
 965 carbon fractions based on radiocarbon measurements. Radiocarbon 47, 93-113.

966 Calderón, F.J., Schultz, D., Paul, E.A., 2012. Carbon allocation, below ground transfers
 967 and lipid turnover in a plant-microbial association. Soil Science Society of America
 968 Proceedings 76, 1614-1623.

969 Calderón F.J., Haddix, M.L., Conant, R.T., Magrini-Blair, K., Paul, E.A., 2013. Mid-
 970 infrared spectroscopy as a method of characterizing changes in soil organic matter. Soil
 971 Science Society of America Proceedings 77, 1591-1600.

972 Campbell, C.A., Paul, E.A., Rennie, D.A., McCallum, R.J., 1967. Applicability of the
 973 carbon dating method of analysis to soil humus studies. *Soil Science* 104, 217-224.

974 Campbell, E., Paustian, K., 2015. Current developments in soil organic matter modeling
 975 and expansions of models: a review. *Environmental Research Letters* 10, 123004.

976 Campbell, R., Porter, R., 1982. Low temperature scanning electron microscopy of
 977 microorganisms in soils. *Soil Biology and Biochemistry* 14, 241-245.

978 Chabbi, A., Kögel Knabner, C.I., Rumpel, C., 2009. Stabilized carbon in subsoil horizons
 979 is located in distinct parts of the soil profile. *Soil Biology and Biochemistry* 41, 256-261.

980 Chahal, K.S., 1968. Biosynthesis and characterization of soil polysaccharides. In:
 981 Isotopes and Radiation in Soil Organic Matter Studies. Proceedings Symposium
 982 FAO/IAEA, Vienna, pp. 207-217.

983 Chattopadhyay, S., Morris, S.J., Paul, E.A., Blackwood, C.B., 2016. Do microbial guilds
 984 track organic carbon content in soil? Unpublished results.

985 Cheeke, T.C., Coleman, D.C., Wall, D.H., 2013. *Microbial Ecology in Sustainable*
 986 *Agroecosystems*. CRC Press, Boca Raton.

987 Chen, Y., Ascher, J., Cecherinni, M., 2012. Stable isotope probing and its application to
 988 soil microbiology. In: Huang, P.M., Li, Y., Sumner, M. (Eds.), *Handbook of Soil*
 989 *Sciences: Properties and Processes*, second ed. CRC Press, Boca Raton, pp. 28-12 to 28-
 990 31.

- 991 Cheng, C-H., Lehman, J., Englehard, M.H., 2008. Natural oxidation of black carbon in
992 soil: changes in molecular form and surface charge along a climo sequence. *Geochimica*
993 *et Cosmochimica Acta* 72, 1598-1610.
- 994 Chenu, C., Rumpel, C., Lehmann, J., 2015. Methods for studying soil organic matter:
995 nature, dynamics, spatial accessibility and interactions with minerals. In: Paul, E.A.
996 (Ed.), *Soil Microbiology, Ecology and Biochemistry*, fourth ed. Academic Press, San
997 Diego, pp. 383-419.
- 998 Cheshire, M.V., Mundie, C.M., Shepard, W., 1974. Transformation of sugars when rye
999 cellulose labeled with ^{14}C decomposes in soil. *Journal of Soil Science* 25, 90-98.
- 1000 Clapp, C.C., Hayes, M.H.B., Simpson, A.K., Kingery, W.L., 2005. Chemistry of soil
1001 organic matter. In: Tabatabai, M.A., Sparks, D.L. (Eds.), *Chemical properties of soils*.
1002 Soil Science Society of America, Book Series, 8, 1-150.
- 1003 Clemente, J.S., Simpson, M.J., 2013. Physical protection of lignin by organic matter and
1004 clay minerals from chemical oxidation. *Organic Geochemistry* 58, 1-12.
- 1005 Coleman, D.C., Fry, B., 1991. *Carbon Isotope Techniques*. Academic Press, San Diego
- 1006 Coleman, D.C., Wall, D., 2014. Soil Fauna: occurrence, biodiversity and roles in
1007 ecosystem function. In: Paul, E.A. (Ed.), *Soil Microbiology, Ecology and Biochemistry*,
1008 fourth ed. Academic Press, San Diego, pp. 111-150.

1009 Coleman, D.C., Crossley, D.A., Hendrix P.F., 2004. Fundamentals of Soil Ecology.
 1010 Elsevier, New York.

1011 Collins, H.P., Blevins, R.J., Bundy, L.G., Christenson, D.R., Dick, W.A., Higgins, D.R.,
 1012 Paul, E.A., 1999. Soil carbon dynamics in corn-based ecosystems: Results from carbon-
 1013 13 abundance. Soil Science Society of America Journal 63, 584-589.

1014 Collins, H.P., Elliott, E.T., Paustian, K., Bundy, L.G., Dick, W.A., Huggins, D.R.,
 1015 Smucker, A.J.M., Paul, E.A., 2000. Soil carbon pools and fluxes in long-term corn-belt
 1016 agroecosystems. Soil Biology and Biochemistry 32, 157-168.

1017 Conant, R.T., Ryan, M.G., Ågren, G.I., Birge, H.E., Davidson, E.A., Eliasson, P.E.,
 1018 Evans, S.E., Frey, S.D., Giardina, C.P., Hopkins, F., Hyvönen, R., Kirschbaum, M.U.F.,
 1019 Lavalley, J.M., Leifeld, J., Parton, W.J., Steinweg, M., Wallenstein, M.D., Wetterstedt,
 1020 J.A.M., Bradford, M.A., 2011. Temperature and soil organic matter decomposition –
 1021 synthesis of current knowledge and a way forward. Global Change Biology 17, 3392-
 1022 3404.

1023 Cotrufo, F., Wallenstein, M., Boot, C., Deneff, K., Paul, E.A., 2013. The molecular
 1024 efficiency-matrix stabilization (MEMS) framework integrates plant litter decomposition
 1025 with soil organic matter stabilization. Do labile plant inputs form stable soil organic
 1026 matter? Global Change Biology 19, 988–995, doi10.1111 GCB 12113.

1027 Cox, G.C., Sanders, F.E., Tinker, F.E., Wild, J., 1975. Ultrastructural evidence relating to
 1028 host-endophyte transfer in a vesicular-arbuscular mycorrhiza. In: Sanders, F.E., Mosse,
 1029 B., Tinker, P.B. (Eds.), *Endomycorrhizae*. Academic Press, London, pp. 297-312.

1030 Crawford, D.I., Crawford, R.L., Pomatto, A.L., 1977. Preparation of specifically labeled
 1031 ^{14}C (lignin) and ^{14}C (cellulose, lignocellulose), and their decomposition by the microflora
 1032 of soils. *Applied and Environmental Microbiology* 33, 1247-1251.

1033 Dahlman, R., Kucera, C.L., 1968. Tagging native grassland vegetation with carbon-14.
 1034 *Ecology* 49, 1199-1203.

1035 De Deyn, G.B., Cornelissen, J.H.C., Bardgett, R.D., 2008. Plant functional traits and soil
 1036 carbon sequestration in contrasting biomes. *Ecology Letters* 11, 516-541.

1037 Di Costy, R.J., Welicky, D.P., Anderson, S.J., Paul, E.A., 2003. ^{15}N CPAMS nuclear
 1038 magnetic resonance spectroscopy and biological stability of soil organic matter in whole
 1039 soil and particle size fractions. *Organic Geochemistry* 34, 1635-1650.

1040 Dinel, H.H., Schnitzer, M., Mehuys, G.R., 1990. Soil lipids: origin, nature, content,
 1041 decomposition and effect on soil physical properties. In: Bollag, J., Stotsky, G. (Eds.),
 1042 *Soil Biochemistry*, volume.6. Marcel Dekker, New York, pp. 397-430.

1043 Drigo, B.J., Anderson, C., Kannangara, G.S.K., Cairney, J.W.G., Johnson, D., 2012.
 1044 Rapid incorporation of carbon from ectomycorrhizal mycelia necromass into soil fungal
 1045 communities. *Soil Biology and Biochemistry* 40, 4-10.

1046 Dungait, J., Hopkins, D., Gregory, A., Whitmore, A., 2012. Soil organic matter turnover is
 1047 governed by accessibility not recalcitrance. *Global Change Biology* 18, 1781-1796.

1048 EPA. 2015. Future climate change <http://www3.epa.gov/climatechange/science/future.html>.

1049 Fang, C., Smith, P., Moncrief, J., Smith, J.U., 2005. Similar response of labile and
 1050 resistant organic pools to changes in temperature. *Nature* 433, 57-59.

1051 Feller, C.L., 1997. The concept of humus in the past three centuries. In: Yaalon, D.H.,
 1052 Berkowicz, S. (Eds.), *History of Soil Science, Advances in Geoecology*, Catena Verlag
 1053 GMBH, Reiskirchen, Germany, ISBN: 9783923381401.

1054 Feller, C.L., Thuriès, L.J.-M., Manlay, R.J., Robin, P., Frossard, E., 2003. The principles
 1055 of rational agriculture” by Albrecht Daniel Thaer (1752–1828), An approach to the
 1056 sustainability of cropping systems at the beginning of the 19th century. *Journal of Plant*
 1057 *Nutrition and Soil Science* 166, 687–698.

1058 Feng, X., Simpson, M.J., 2011. Molecular-level methods for monitoring soil organic
 1059 matter responses to global climate change (invited review). *Journal of Environmental*
 1060 *Monitoring* 13, 1246-1254.

1061 Fierer, N., Grandy, S., Six, J., Paul E.A., 2009. Searching for unifying principles in soil
 1062 ecology. *Soil Biology and Biochemistry* 41, 2249-2256.

1063 Finch, P., Hayes, M.B.H., Stacey, M., 1971. The biochemistry of soil polysaccharides. In:
 1064 McLaren, A.D., Skujins, J. (Eds.), Soil Biochemistry, Volume 2. Marcel Dekker, New
 1065 York, pp. 258-319.

1066 Follett, R.F., Paul, E.A., Leavitt, S.W., Halvorson, A., Lyon, D., Peterson, G.A., 1997.
 1067 Carbon isotope ratios of Great Plains soils and in wheat-fallow systems. Soil Science
 1068 Society of America Journal 61, 1068-1077.

1069 Follett, R.F., Paul, E.A., Pruessner, E.G., 2007. Soil carbon dynamics during a long-term
 1070 incubation study involving ^{13}C and ^{14}C measurements. Soil Science 172, 189-208.

1071 Fontaine, S., Mariotti, A., Abbadie, L., 2003. The priming effect of organic matter: A
 1072 question of microbial competition? Soil Biology and Biochemistry 35, 837-843.

1073 Foster, R.C., Martin, J.K., 1981. *In situ* analysis of soil components of biological origin.
 1074 In: Paul, E.A., Ladd, J.N. (Eds.), Soil Biochemistry, volume 5. Marcel Dekker, New
 1075 York, pp. 75-112.

1076 Foster, R.C., Rovira, A.D., Cook, T.W., 1983. Ultrastructure of the root soil interface.
 1077 The American Phytopathological Society, St. Paul, Minnesota.

1078 Fry, S.D., 2015. The spatial distribution of soil biota. In: Paul, E.A. (Ed.), Soil
 1079 Microbiology, Ecology and Biochemistry, fourth ed. Academic Press, San Diego, pp.
 1080 223-244.

1081 Gianfreda, I., Rao, M.A., Mora, M., 2012. Enzymatic activity as influenced by soil
 1082 mineral and humic colloids and its impact on biogeochemical processes. In: Huang, P.M.,
 1083 Li, Y., Sumner, M. (Eds.), Handbook of Soil Sciences: Resource Management and
 1084 Environmental Impacts, second ed. CRC Press, Taylor and Francis, pp. 5-1 to 5-24.

1085 Gillespie, W., Diochon, A.B., Ma, B.A., Morrison, M.J., Kellman, L., Walley, F.L.,
 1086 Regier, T.Z., Chevrier, D., Dynes, J.J., Gregorich, E.G., 2014. Nitrogen input quality
 1087 changes the biochemical composition of soil organic matter stabilized in the fine fraction:
 1088 a long-term study. Biogeochemistry 117, 337-352.

1089 Gleixner, G., 2013. Soil organic matter dynamics: A biological perspective derived from
 1090 the use of compound specific isotope studies. Ecological Research 28, 683-695.

1091 Goh, K.M., 1991. Carbon dating. In: Coleman, D.C, Fry, B. (Eds.), Carbon Isotope
 1092 Techniques. Academic Press, San Diego, pp. 125-146.

1093 Gregorich, E.G., Ellert, B.H., Monreal, C.M., 1995. Turnover of soil organic matter and
 1094 storage of corn residue carbon estimated from natural ^{13}C abundance. Canadian Journal
 1095 of Soil Science 75, 161-167.

1096 Gregorich, E.G., Monreal, C.M., Schnitzer, M., Schulten, H.R., 1996. Transformation of
 1097 plant residues into soil organic matter: Chemical characterization of plant tissue, isolated
 1098 soil fractions, and whole soils. Soil Science 161, 680-693.

1099 Haddix, M.L., Plante, A.F., Conant, R.T., Paul, E.A., Six, J., Steinweg, J.M., Drijber,
 1100 R.A., Morris S.J., 2011. The role of soil characteristics on temperature sensitivity of soil
 1101 organic matter. *Soil Science Society of America Journal* 75, 56-68.

1102 Haider, K., 1992. Problems related to the humification processes in soils of temperate
 1103 climates. In: Stotzky, G., Bollag, J.M. (Eds.), *Soil Biochemistry, Volume 7*. Marcel
 1104 Dekker, New York. pp. 55-94.

1105 Haider, K., Martin, J.P., Filip, Z., 1975. Humus biochemistry. In: Paul, E.A., McLaren,
 1106 A.D. (Eds.), *Soil Biochemistry, Volume 4*. Marcel Dekker, New York, pp. 195-244.

1107 Haile-Mariam, S., Collins, H.P., Paul, E.A., 2008. Dynamics of soil organic matter
 1108 constituents determined by ^{13}C changes during long-term incubation. *Soil Science*
 1109 *Society of America Journal* 72, 370-378.

1110 Hamilton, S.K., Doll, J.E., Robertson, G.P., 2015. *The Ecology of Agricultural*
 1111 *Landscapes: Long-Term Research on the Path to Sustainability*. Oxford University Press,
 1112 New York.

1113 Harris, D., Paul, E.A., 1994. Measurement of bacterial growth rates in soil. *Applied Soil*
 1114 *Ecology* 1, 277-290.

1115 Hillel, D., 1991. *Out of the Earth, Civilization and the Life of the Soil*. University of
 1116 California Press, Berkeley.

1117 Hinckley, E.S., Wieder, W., Fierer, N., Paul, E.A., 2014. Digging into the soil beneath
 1118 our feet: Bridging knowledge across scales in the age of global change. EOS
 1119 Transactions, American Geophysical Union 95, 11, 95-97.

1120 Hinsinger, P., Jones, D.L., Marschner, P., 2012. Biogeochemical, biophysical and
 1121 biological processes in the rhizosphere. In: Huang, P.M., Li, Y., Sumner, M. (Eds.),
 1122 Handbook of Soil Sciences: Resource Management and Environmental Impacts, second
 1123 ed. CRC Press, Boca Raton, pp. 6-1 to 6-30.

1124 Hoover, C., Magrini, K., Evans, R., 2002. Use of pyrolysis molecular beam mass
 1125 spectrometry (py-MBMS) to characterize forest soil carbon: Preliminary results from an
 1126 old-growth forest in Northwestern Pennsylvania. Environmental Pollution 116, S269-
 1127 S275.

1128 Horwath, W.R., 2015. Carbon cycling: the dynamics and formation of organic matter. In:
 1129 Paul, E.A. (Ed.), Soil Microbiology, Ecology and Biochemistry, fourth ed. Academic
 1130 Press, San Diego, pp. 339-359.

1131 Horwath, W.R., Paul, E.A., 1994. Microbial biomass. In: Weaver, R.W., Angle, S.,
 1132 Bottomley, P. (Eds.), Biochemical and Microbiological Properties of Soil. Methods of
 1133 Soil Analyses, Part 2. Soil Science Society of America Journal, Madison, pp. 753-773.

1134 Horwath, W.R., Pregitzer, K.S., Paul, E.A., 1994. ^{14}C allocation in tree-soil systems. Tree
 1135 Physiology 14, 1163-1176.

1136 Hugelius, G., Strauss, J., Zubrzycki, S., Harden, J.W., Schuur, E.A., Ping, C.-L.,
 1137 Schirrmeister, L., Grosse, G., Michaelson, G.J., Koven, C.D., O'Donnell, J.A., Elberling,
 1138 B., Mishra, U., Camill, P., Yu, Z., Palmtag, J., Kuhry, P., 2014. Estimated stocks of
 1139 circumpolar permafrost carbon with quantified uncertainty ranges and identified data
 1140 gaps. *Biogeosciences* 11, 6573-6593.

1141 Hungate, B.A., Jackson, R.B., Field, C.B., Chapin, F.S. III., 1996. Detecting changes in
 1142 soil carbon in CO₂ enrichment experiments. *Plant and Soil* 187:135-145.

1143 Hurisso, T.T., Davis, J.G., Brummer, J.E., Stromberger, M.E., Mikha, M.M., Haddix,
 1144 M.L., Booher, M.R., Paul, E.A., 2013. Rapid changes in microbial biomass and aggregate
 1145 size distribution in response to changes in organic matter management in grass pasture.
 1146 *Geoderma* 193-194, 68-75.

1147 Jansson, S.L., 1958. Tracer studies on nitrogen transformations in soil with special
 1148 attention to mineralization-immobilization relationships. *Kungj. Landbrukshögskolans*
 1149 *Annaler* 24, 101-164.

1150 Jansson, S.L., 1963. Balance sheet and residual effects of fertilizer in a six year study
 1151 with ¹⁵N. *Soil Science* 95, 31-37.

1152 Jastrow, J.D., 1996. Soil aggregate formation and the accrual of particulate and mineral
 1153 associated organic matter. *Soil Biology and Biochemistry* 28, 656-676.

1154 Jastrow, J.W., Miller, R.M., 1998. Soil aggregate stabilization and carbon sequestration;
 1155 feedback through organo-mineral associations. In. Lal, R., Kimble, J.M., Follett, R.P.,

1156 Stewart, B.A., (Eds.), Soil Processes and the Carbon Cycle. CRC Press, Boca Raton, pp.
 1157 207-223.

1158 Jenkinson, D.S., 1965. Studies on the decomposition of plant material in soil. I. Losses of
 1159 carbon from ^{14}C -labelled ryegrass incubated in the field. Journal of Soil Science 16, 104-
 1160 115.

1161 Jenkinson, D.S., 1971. Studies of the decomposition of C^{14} labelled organic matter in
 1162 soils. Soil Science 111, 64-69.

1163 Jenkinson, D.S., 1990. The turnover of organic carbon and nitrogen in soil. Philosophical
 1164 Transactions of the Royal Society B 320, 361-368.

1165 Jenkinson, D.S., Ladd, J.N., 1981. Microbial biomass in soil: Measurement and turnover.
 1166 In: Paul, E.A., Ladd, J.N. (Eds.), Soil Biochemistry, Volume 5. Marcel Dekker, New
 1167 York, pp. 415-472.

1168 Jenkinson, D.S., Rayner, J.H., 1977. The turnover of soil organic matter in some of the
 1169 Rothamsted classical experiments. Soil Science 123, 298-305.

1170 Jenny, H., 1941. Factors of Soil Formation. McGraw Hill, New York.

1171 Jones, D.L., Nguyen, C., Finlay, R.D., 2009. Carbon flow in the rhizosphere: Carbon
 1172 trading at the soil-root interface. Plant and Soil 321, 5-33.

1173 Kandeler, E., 2015. Physiological and biochemical methods for studying soil biota and
 1174 their functions. In: Paul, E.A. (Ed.), *Soil Microbiology, Ecology and Biochemistry*,
 1175 fourth ed. Academic Press, San Diego, pp. 188-222.

1176 Kemmitt, S.J., Lanyon, C.V., Waite, I.S., Wen, Q., Addiscott, T.M., Bird, N.R.A.,
 1177 O'Donnell, A.G., Brookes, P.C., 2008. Mineralization of native soil organic matter is not
 1178 regulated by the size, activity or composition of the soil microbial biomass – a new
 1179 perspective. *Soil Biology and Biochemistry* 40, 61-73.

1180 Killham, K., Prosser, J.I., 2015. The bacteria and archaea. In: Paul, E.A. (Ed.), *Soil*
 1181 *Microbiology, Ecology and Biochemistry*, fourth ed. Academic Press, San Diego, pp. 41-
 1182 76.

1183 Kirkham, D., Bartholomew, W.V., 1954. Equations following nutrient transformations in
 1184 soil, utilizing tracer data. 1. *Soil Science Society of America Proceedings* 18, 189-192.

1185 Kirkham, D., Bartholomew, W.V., 1955. Equations following nutrient transformations in
 1186 soil, utilizing tracer data. 11. *Soil Science Society of America Proceedings* 19, 33-34.

1187 Kögel-Knabner, I., Kleber, M., 2012. Mineralogical, physiochemical and microbiological
 1188 controls on soil organic matter stabilization and turnover. In: Huang, P.M., Li, Y.,
 1189 Sumner, M. (Eds.), *Handbook of Soil Sciences: Resource Management and*
 1190 *Environmental Impacts*, second ed. CRC Press, pp. 7-1 to 7-21.

1191 Kögel-Knabner, I., Guggenberger, G.M., Kleber, M., Kandellar, E., Kalbitz, K., Scheu,
 1192 S., Eusterhaus, K., Leinweiber, R., 2008. Organo mineral associations in temperate soils:

1193 Integrating biology, mineralogy and organic chemistry. Journal of Plant
 1194 Nutrition and Soil Science 171, 61-82.

1195 Kong, A.Y., Scow, K.M., Holmes, W.E., Six, J., 2011. Microbial community composition
 1196 and carbon cycling within soil microenvironments of conventional, low-input, and
 1197 organic cropping systems. Soil Biology and Biochemistry 43, 20-30.

1198 Kononova, M.M., 1961. Soil Organic Matter. It's Nature. It's Role in Soil Formation and
 1199 Soil Fertility. Pergammon Press, Oxford UK.

1200 Krull, E.S., Swanson, C.W., Skjemstad, J.O., 2006. Importance of charcoal in
 1201 determining the age and chemistry of organic carbon in surface soils. Journal of
 1202 Geophysical Research 111, G04001.

1203 Kucey, R.M.N., Paul, E.A., 1982. Carbon flow, photosynthesis and N₂ fixation in
 1204 mycorrhizal and nodulated faba beans, *Vicia faba*. Soil Biology and Biochemistry 14,
 1205 407-412.

1206 Kuzyakof, Y., Domanski, G., 2000. Carbon inputs by plants into soil. Review, Journal of
 1207 Plant Nutrition and Soil Science 163, 421-431.

1208 Kuzyakov, Y., 2010. Priming effects: Interactions between living and dead organic
 1209 matter. Soil Biology and Biochemistry 42, 1363-1371.

1210 Ladd, J.N., Butler, J.H.A., 1975. Humus enzyme systems and synthetic, organic polymer-
 1211 enzyme molecules. In: Paul, E.A., Ladd, J.N. (Eds.), Soil Biochemistry, Vol. 4. Marcel
 1212 Dekker, New York, pp. 143-180.

1213 Ladd, J.N., Martin, J.K., 1984. Soil organic matter studies. In: L'Annunziata, M.F., Legg,
 1214 J.O. (Eds.), Isotopes and Radiation in Agricultural Sciences, Vol. 1. Academic Press, San
 1215 Diego, pp. 1-34.

1216 Ladd, J.N., Paul, E.A., 1973. Changes in enzyme activity and distribution of acid-soluble,
 1217 amino acid-nitrogen in soil during nitrogen immobilization and mineralization. Soil
 1218 Biology and Biochemistry, 5, 825-840.

1219 Lajtha, K., Townsend, K. L., Kramer, M. G., Swanston, C., Bowden, R.D., Nadelhoffer,
 1220 K., 2014. Changes in particulate versus mineral-associated soil carbon after 50 years of
 1221 litter accumulation in forest and prairie experimental ecosystems. Biogeochemistry
 1222 119:341-360.

1223 Lal, R. Nagassa, W. Lorenz, K., 2015. Carbon sequestration in soils. Current Opinions in
 1224 Environmental Sustainability 15, 79-88.

1225 Lavallee, J.M., Haddix, M.L., Conant, R.T., Bird, M.I., Paul, E.A., 2015. Contribution of
 1226 pyrogenic carbon to soil organic matter fractions in agricultural soils and its influence on
 1227 calculated turnover times. Unpublished results.

1228 Leavitt, S.W., Paul, E.A., Pendall, E., Pinter, Jr., P.J., Kimball, B.A., 1997. Field
 1229 variability of carbon isotopes in soil organic carbon. *Nuclear Instruments and Methods in*
 1230 *Physics Research B* 123, 451-454.

1231 Leavitt, S.W., Pendall, E., Paul, E.A., Brooks, T., Kimball, B.A., Pinter, Jr., P.J.,
 1232 Johnson, H.B., Matthias, A., Wall, G.W., LaMorte, R.L., 2001. Stable-carbon isotopes
 1233 and soil organic carbon in wheat under CO₂ enrichment. *New Phytologist* 150, 305-314.

1234 Lehmann, J., Kleber, M., 2015. The contentious nature of soil organic matter. *Nature* 528,
 1235 60-68.

1236 Lehmann, J., Liang, B., Solomon, D., Lerotic, M., Luizão, F., Kinyangi, F., Schäfer, T.,
 1237 Wirick, S., Jacobsen, C., 2005. Near-edge X-ray absorption resource fine structure
 1238 (NEXAFS) spectroscopy for mapping nano-scale distribution of organic carbon forms in
 1239 soil: application to black carbon particles. *Global Biogeochemical Cycles* 19, GB 1013.

1240 Löhnis, F., Fred, E.B., 1923. *Textbook of Agricultural Bacteriology*. McGraw Hill, New
 1241 York.

1242 Manzoni, S., Porporato, A., 2009. Soil carbon and nitrogen mineralization: Theory and
 1243 models across scales. *Soil Biology and Biochemistry* 41, 1355-1379.

1244 Manzoni, S., Piñeiro, G., Jackson, R.B., Jobbá, E.J., Kim, E.G., Porporato, A., 2012.
 1245 Analytical models of soil and litter decomposition: Solutions for mass loss and time
 1246 dependent decay rates. *Soil Biology and Biochemistry* 5, 66-76.

1247 Marshall, K.C., 1971. Sorptive interactions between soil particles and microorganism. In:
 1248 McLaren, A.D., Skujins, J. (Ed.), Soil Biochemistry, Vol. 2. Marcel Dekker, New York,
 1249 pp. 409-445.

1250 Martel, Y.A., Paul, E.A., 1974a. The use of radiocarbon dating of organic matter in the
 1251 study of soil genesis. Soil Science Society of America Proceedings 38, 501-506.

1252 Martel, Y.A., Paul, E.A., 1974b. Effects of cultivation on the organic matter of grassland
 1253 soils as determined by the fractionation and radiocarbon dating. Canadian Journal of Soil
 1254 Science 54, 419-426.

1255 Martin, J.P., Haider, K., Farmer, W.J., Mathon, E.F., 1974. Decomposition and
 1256 distribution of residual activity of some ^{14}C -microbial polysaccharides and cells and
 1257 glucose cellulose and wheat straw in soil. Soil Biology and Biochemistry 6, 221-230.

1258 Mathieu, J.A., Hatté, C., Balesdent, J., Parent É., 2015. Deep soil carbon dynamics are
 1259 driven more by soil type than by climate: a worldwide meta-analysis of radiocarbon
 1260 profiles. Global Change Biology 21, 4278-4292.

1261 Mayaudon, J., Simonart P., 1958. Étude de la decomposition de la matière organique dans le
 1262 sol de moyen du carbone radioactif II. Plant and Soil 9, 381-384.

1263 Mayaudon, J., Simonart, P., 1963. Humification des microorganismes marqués par ^{14}C dans
 1264 le sol. Annales de l'Institut Pasteur 105, 257-266.

1265 McGill, W.B., Shields, J.A., Paul, E.A., 1975. Relation between carbon and nitrogen
1266 turnover in soil organic fractions of microbial origin. *Soil Biology and Biochemistry* 7,
1267 57-63.

1268 McLauchlan, K.K., Hobbie, S.A., 2004. Comparison of labile soil organic matter
1269 fractionation techniques. *Soil Science Society of America Journal* 68, 1612-1625.

1270 McMahon, S.K., Williams, M.A., Bottomley, P.J., Myrold, D.A., 2005. Dynamics of
1271 microbial communities during decomposition of carbon-13 labeled ryegrass fractions in
1272 soil. *Soil Science Society of America Journal* 69, 1238-1247.

1273 Mellor, N.J., Hellerich, J., Drijber, R., Stromberg, M., Morris, S.J., Paul, E.A., 2013.
1274 Changes in ecosystem carbon following afforestation of native sand prairie. *Soil Society*
1275 *of America Journal* 77, 1613- 1624.

1276 Mikha, M.M., Vigil, M.F., Benjamin, J.G., 2013. Long-term tillage impacts on soil
1277 aggregation and carbon dynamics under wheat-fallow in the central Great Plains. *Soil*
1278 *Society of America Journal* 77, 594-605.

1279 Miltner, A., Bombach, P., Schmidt-Brücken, B., Kastner, M., 2011. SOM genesis:
1280 microbial biomass as a significant source. *Biogeochemistry* 111, 41-56.

1281 Moore, J.C., de Ruiter, P.C., 2012. *Energy Food Webs: An Analysis of Real and Model*
1282 *Ecosystems*. Oxford University Press, Oxford, UK.

1283 Morris, S.J., Conant, R., Mellor, N., Brewer, E., Paul, E.A., 2011. Controls on soil carbon
 1284 sequestration and dynamics: Lessons from land use change. *Nematology* 72, 78-83.

1285 Myrold, D.D., Zeglin, L.H., Janssen, J.K., 2013. The potential of metagenomic processes
 1286 for understanding soil microbial approaches. *Soil Science of America Journal* 78, 3-10.

1287 Nannipieri, P., Small, K., 2006. *Nucleic Acids and Proteins in Soil*. *Soil Biology*,
 1288 Springer Verlag, Berlin, 8.

1289 Nannipieri, P., Paul, E.A., 2009. The chemical and functional characteristics of soil
 1290 organic N and its biotic components. *Soil Biology and Biochemistry* 41, 2357-2369.

1291 Norby, R.J., Zak, D.R., 2011. Ecological lessons from free-air CO₂ enrichment (FACE)
 1292 experiments. *Annual Review of Ecology, Evolution, and Systematics* 42, 181.

1293 Norman, A.G., 1946. Recent advances in soil microbiology. *Soil Science Society of*
 1294 *America Proceedings* 11, 9-15.

1295 Norman, A.G., Werkman, C.H., 1943. The use of the nitrogen isotope N¹⁵ in determining
 1296 nitrogen recovery from plant materials decomposing in soil. *Journal of American Society*
 1297 *of Agronomy* 35, 1023-1025.

1298 Oades, J.M., Wagner, G.H., 1970. Incorporation of ¹⁴C into sugars in a soil incubated
 1299 with ¹⁴C glucose. *Geoderma* 4, 417-423.

1300 O'Brien, S.L., Jastrow, J.D., 2013. Physical and chemical protection in hierarchical soil
 1301 aggregates regulates soil carbon and nitrogen recovery in restored perennial grasslands.
 1302 *Soil Biology and Biochemistry* 61, 1-13.

1303 O'Brien, B.J., Stout, J.D., 1978. Movement and turnover of soil organic matter as
 1304 indicated by carbon isotopes measurement. *Soil Biology and Biochemistry* 10, 309-317.

1305 O'Brien, S., Jastrow, J.D. Grimely, D.A., Gonzalez-Meir, M.A., 2015. Edaphic
 1306 controls on soil organic carbon stocks in restored grasslands. *Geoderma*
 1307 251-252, 117-123.

1308 Osler, G. H.R., Sommerkorn, M., 2007. Toward a completed soil C and N cycle
 1309 incorporating the soil fauna. *Ecology* 88:1611-1621.

1310 Parton, W.J., Schimel, D.S., Cole, C.V., Ojima, D.S., 1987. Analysis of factors
 1311 controlling soil organic matter levels in Great Plains grasslands. *Soil Science Society of*
 1312 *America Journal* 51, 1173-1179.

1313 Parton, W.J., Hartman, M., Ojima, D., Schimel, S., 1998. DAYCENT and its land surface
 1314 sub model: description and testing. *Global and Planetary Change* 19, 32-48.

1315 Parton, W.J., Del Grosso, S.J., Plante, A.F., Carol, E., 2015. The dynamics of soil organic
 1316 matter and nutrient cycling. In: Paul, E.A. (Ed.), *Soil Microbiology, Ecology and*
 1317 *Biochemistry*, fourth ed. Academic Press, San Diego, pp. 505-538.

1318 Paul, E.A., 2007. Soil microbiology, ecology and biochemistry in perspective. In: Paul,
 1319 E.A. (Ed.), Soil Microbiology Ecology and Biochemistry, third ed. Academic Press, San
 1320 Diego, pp. 3-24.

1321 Paul, E.A., 2015. Soil microbiology, ecology and biochemistry: An exciting present and
 1322 great future built on basic knowledge and unifying concepts. In: Paul, E.A. (Ed.), Soil
 1323 Microbiology, Ecology and Biochemistry, fourth ed. Academic Press, San Diego, pp. 1-
 1324 15.

1325 Paul, E.A., Clark, F.E., 1986. Soil Microbiology and Biochemistry. Academic Press, San
 1326 Diego.

1327 Paul, E.A., Kucey, R.M.N., 1981. Carbon flow in plant microbial associations. Science
 1328 213, 473-474.

1329 Paul, E.A., van Veen, J.A., 1978. The use of tracers to determine the dynamic nature of
 1330 organic matter. Proceedings of 11th International Society of Soil Science, Trans.
 1331 (Edmonton), June, 1978, pp. 61-102.

1332 Paul, E.A., Voroney, R.P., 1980. Nutrient and energy flows through soil microbial
 1333 biomass. In: Ellwood, R.D., Hedger, N., Lynch, M.J., Slater, J.H. (Eds.), Contemporary
 1334 Microbial Ecology. Academic Press, New York, pp. 215-237.

1335 Paul, E.A., Follett, R.F., Leavitt, S.W., Halvorson, A., Peterson, G.A., Lyon, D.J., 1997.
 1336 Radiocarbon dating for determination of soil organic matter pool sizes and dynamics.
 1337 Soil Science Society of America Journal 61, 1058-1067.

1338 Paul, E.A., Harris, D., Collins, H.P., Schulthess, U., Robertson, G.P., 1999. Evolution of
 1339 CO₂ and soil carbon dynamics in biologically managed, row-crop agroecosystems.
 1340 Applied Soil Ecology 11, 53-65.

1341 Paul, E.A., Collins, H.P., Leavitt, S.W., 2001a. Dynamics of resistant soil carbon of
 1342 Midwestern agricultural soils measured by naturally-occurring ¹⁴C abundance. Geoderma
 1343 104, 239-256.

1344 Paul, E.A., Morris, S.J., Böhm, S., 2001b. The determination of soil C pool sizes and
 1345 turnover rates: biophysical fractionation and tracers. In: Lal, R., Kimble, J.M., Follett,
 1346 R.F., Stewart, B.A. (Eds.), Assessment Methods for Soil Carbon. Lewis Publishers, Boca
 1347 Raton, pp. 193-205.

1348 Paul, E.A., Collins, H.P., Elliott, E.T., Frey, S., Juma, N., Janzen, H., Campbell, C.A.,
 1349 Zentner, R.P., La Fond, G.P., Moulin, A.P., 2004. Management effects on the dynamics
 1350 and storage rates of organic matter in long-term rotations. Canadian Journal of Soil
 1351 Science 84, 49-61.

1352 Paul, E.A., Morris, S.J., Conant, R.T., Plante, A.F., 2006. Does the acid hydrolysis-
 1353 incubation method measure meaningful soil organic matter pools? A review of the
 1354 literature. Soil Science Society of America Journal 70, 1023-1035.

1355 Paul, E.A., Follett, R.F., Haddix, M., Preussner, E., 2011. Soil N dynamics related to soil
 1356 C and microbial changes during long-term incubation. Soil Science 176, 527-536.

1357 Pelz, O., Abraham, W., Saurer, M., Siegwolf, R., Zeyer, J., 2005. Microbial assimilation
 1358 of plant-derived carbon in soil traced by isotope analysis. *Biology and Fertility of Soils*
 1359 41, 153-162.

1360 Piccolo, A. 2001. The supramolecular structure of humic substances. *Soil Science* 166,
 1361 810-832.

1362 Ping, C.L., Jastrow, J.D., Jorgenson, M.T., Michaelson, G.J., and Schuur, Y.L., 2015.
 1363 Permafrost soils and carbon cycling, *SOIL* 1, 147-171.

1364 Pisana, O., Hills, K.M., Courtier-Murias, D., Simpson, A.J., Mellor, N.J., Paul, E.A.,
 1365 Morris, S.J., Simpson, M.J., 2013. Molecular level analysis of long-term vegetation shifts
 1366 and relation to soil organic matter composition. *Organic Biogeochemistry* 62, 7-16.

1367 Pisana, O., Hills, K.M., Courtier-Murias, D., Haddix, M.L., Paul, E.A., Conant, R.T.,
 1368 Simpson, A.J., Arhonditsis, B., Simpson, M.J., 2014. Accumulation of aliphatic
 1369 compounds in soil with increasing mean annual temperature. *Organic Geochemistry* 76,
 1370 118-127.

1371 Pisani, O., Haddix, M.L., Conant, R.T., Paul, E.A., Simpson, M.J., 2015. Molecular level
 1372 shifts in soil organic matter composition with cultivation along a mean annual
 1373 temperature gradient. *Plant and Soil*, Unpublished.

1374 Pietramella, G., Ascher, J.C., Ceccherini, M., 2012. Expression of genes in soil. In:
 1375 Huang, P.M., Li, Y., Sumner, M. (Eds.), *Handbook of Soil Sciences: Properties and*
 1376 *Processes*, second ed. CRC Press, Boca Raton, pp. 28-12 to 28-31.

1377 Plante, A.F., Magrini-Baer, K., Vigil, M., Paul, E.A., 2009. Pyrolysis-molecular beam
 1378 mass spectrometry to characterize soil organic matter composition in chemically isolated
 1379 fractions from different land uses. *Biogeochemistry* 92, 145-161.

1380 Poirier, N., Derenne, S., Balesdent, J., Chenu, C., Bardoux, G., Mariotti, A., Largeau, C.,
 1381 2005. Dynamics and origin of the non-hydrolyzable organic fraction in a forest and a
 1382 cultivated temperate soil as determined by isotopic and microscopic studies. *European*
 1383 *Journal of Soil Science* DOI: 10.1111/j.1365-2389.2005.00764.x

1384 Rasse, D.P., Rumpel, C., Dignac, M-F., 2005. Is soil carbon mostly root carbon?
 1385 Mechanisms for a specific stabilization. *Plant and Soil* 269, 341-356.

1386 Redmile-Gordon, M.A., Brookes, P.C., Evershed, R.P., Goulding, K.W.T., Hirsch, P.R.,
 1387 2014. Measuring the soil-microbial interface: Extraction of extracellular polymeric
 1388 substances (EPS) from soil biofilms. *Soil Biology Biochemistry* 72:163-171.

1389 Rillig, M.C., Ramsey, P.W., Morris, S.J., Paul, E.A., 2003. Glomalin, an arbuscular-
 1390 mycorrhizal soil protein. *Plant and Soil* 253, 293-299.

1391 Robertson, G.P., Groffman, P.M., 2015. Nitrogen transformations in soil. In: Paul, E.A.
 1392 (Ed.), *Soil Microbiology, Ecology and Biochemistry*, fourth ed. Academic Press, San
 1393 Diego, pp. 421-446.

1394 Rumpel C, Kögel-Knabner, I. 2011. Deep soil organic matter-a key, but poorly
 1395 understood component of terrestrial C cycle. *Plant and Soil* 338, 143-158.

1396 Sage, R.F., 1995. Was low atmospheric CO₂ during the Pleistocene a limiting factor for
 1397 the origin of agriculture? *Global Change Biology* 1, 93-106.

1398 Sanchez, J.E., Paul, E.A., Willson, T.C., Smeenk, J., Harwood, R.R., 2002. Corn root
 1399 effects on the nitrogen supplying power of a conditioned soil. *Agronomy Journal* 94, 391-
 1400 396.

1401 Scharpenseel, H.W., Schiffman, H., 1977. Radiocarbon dating of soils: A review.
 1402 *Zeitschrift für Pflanzenernährung und Bodenkunde* 140, 159-174.

1403 Scharpenseel, H.W., Becker-Heidmann, P., Neue, H.U., Tsutsuki, K., 1989. Bomb-
 1404 carbon, ¹⁴C dating and ¹³C measurements as tracers of organic matter dynamics as well as
 1405 of morphogenetic and turbation processes. *Science of the Total Environment* 81, 99-110.

1406 Schimel, J., 2001. Biogeochemical models implicit versus explicit microbiology. In:
 1407 Schultz, E.D., Harrison, S.P., Heiman, M. et al. (Eds.), *Global biogeochemical models in*
 1408 *the climatic system*. Academic Press, San Diego, pp. 177-183.

1409 Schindler, P.V., Mercer, E.J., Rice, J.A., 2007. Chemical characterization of glomalin –
 1410 related soil protein (GRSP) extracted from soils of various organic matter contents. *Soil*
 1411 *Biology Biochemistry* 39, 320-329.

1412 Schmidt, M., Torn, W.I., Abiven, M.S., Dittmar, S., Guggenberger, T., Janssens, G.,
 1413 Kleber, I.A., Kögel-Knabner, M., Lehmann, I., Manning, J., Nannipieri, P., Rasse, D.P.,
 1414 Weiner, S., Trumbore, S.E., 2011. Persistence of soil organic matter as an ecosystem
 1415 property. *Nature* 478 (7367), 49-56.

1416 Schnitzer, M., Monreal, J.A., 2011. Quo vadis soil organic matter research? A biological
 1417 link to the chemistry of humification. *Advances in Agronomy* 113, 139-213.

1418 Schnurer, J.M., Clarholm, M., Rosswall, T., 1985. Microbial biomass and activity in
 1419 agricultural soil with different organic matter contents. *Soil Biology and Biochemistry*
 1420 17, 611-618.

1421 Schulten, H.R., 1996. Direct pyrolysis-mass spectrometry of soils: A novel tool in
 1422 agriculture, ecology, forestry, and soil science. In: Boutton, T.W., Yamasaki, S.H. (Eds.),
 1423 *Mass Spectrometry of Soils*. Marcel Dekker, Inc., New York, pp. 373-436.

1424 Schulten, H.R., Leinweber, P., 1993. Influence of the mineral matrix on the formation
 1425 and molecular composition of soil organic matter in a long-term, agricultural experiment.
 1426 *Biogeochemistry* 22, 1-22.

1427 Schulten, H.R., Leinweber, P., 1999. Advances in analytical pyrolysis of soil organic
 1428 matter. *Journal of Analytical and Applied Pyrolysis* 49 (1-2), 359-383.

1429 Schuur, E.A.G., McGuire, A.D., Schädel, C., Grosse, G., Harden, J.W., Hayes, D.J.,
 1430 Hugelius, G., Kovenm, C.D., Kuhry, P., Lawrence, D.M., Natali, S.M., Olefeldt, C.,
 1431 Romanovsky, V.E., Schaefer, K., Turetsky, M.R., Treat, C.C., Vonk, J.E., 2015. Climate
 1432 change and the permafrost carbon feedback. *Nature* 520, 171-179.

1433 Segoli, M., De Gryze, S., Dou, F., Lee, J., Post, W.M., Denef, K., Six, J., 2013.
 1434 *AggModel*: A soil organic matter model with measurable pools for use in incubation
 1435 studies. *Ecological Modeling* 263, 1-9.

1436 Shields, J.A., Paul, E.A., St. Arnaud, R.J., Head, W.K., 1968. Spectrophotometric
 1437 measurement of soil color and its relationship to moisture and organic matter. *Canadian*
 1438 *Journal of Soil Science* 48, 271-280.

1439 Shields, J.A., Paul, E.A., Lowe, W.E., Parkinson, D., 1973. Turnover of microbial tissue
 1440 in soil under field conditions. *Soil Biology and Biochemistry* 5, 753-764.

1441 Simpson, A.J., McNally, D.J., Simpson, M.J., 2011. NMR spectroscopy in environmental
 1442 research: From molecular interactions to global processes. *Progress in Molecular*
 1443 *Magnetic Spectroscopy* 58, 97-175.

1444 Sinsabaugh, R., Follstad Shah, J.J., 2012. Ecoenzymatic stoichiometry and ecological
 1445 theory. *Annual Review of Ecology and Systematics* 43, 313-343.

1446 Sinsabaugh, R., Manzoni, D., Moorehead, D., Richter, A., 2013. Carbon use efficiency of
 1447 microbial communities: stoichiometry, methodology and modeling. *Ecology Letters*
 1448 doi:10.2111/ele.123113.

1449 Sinsabaugh, R.L., Turner, B.T., Talbot, J.M., Waring, B., Powers, J.S., Kuske, C.R.,
 1450 Moorhead, D., Follstad Shah, J.F., 2016. Stoichiometry of microbial carbon use
 1451 efficiency in soils. *Ecological Monographs*, Unpublished results.

1452 Six, J., Feller, C., Denef, K., Ogle, S.M., de Morales, J.O., Albrecht, A., 2002a. Soil
 1453 organic matter, biota and aggregation in temperate and tropical soils-Effects of no tillage.
 1454 *Agronomie* 22, 755-775.

1455 Six, J., Conant, R.T., Paul, E.A., Paustian, K., 2002b. Stabilization mechanisms of soil
 1456 organic matter: Implications for C-saturation of soils. *Plant and Soil* 241, 155-176.

1457 Smith, J.L., Paul, E.A., 1990. The significance of soil microbial biomass estimations. In:
 1458 Skujins, J., Bollag, J. (Eds.), *Soil Biochemistry*, vol. 6. Marcel Dekker, Inc., New York,
 1459 pp. 357-396.

1460 Smith, P., Smith, J.U., Powlson, D.S., McGill, W.B., Arah, J.R., Chertov, O.G.,
 1461 Coleman, K., Franko, U., Fröklings, S., Jenkinson, D.S., Jensen, K.L., Kelly, R.H., Klein-
 1462 Gunneweik, H., Komerov, A.S., Li, C., Molina, J., Parton, W.J., Thornley, J., Whitmore,
 1463 A., 1997. A comparison of the performance of nine soil organic matter models using data
 1464 sets from seven long-term experiments. *Geoderma* 81, 153-225.

1465 Snellgrove, R., Splittstoesser, W., Stribley, D., Tinker, P.B., 1982. The distribution of
 1466 carbon and demand of the fungal symbiont in leek plants infected with arbuscular
 1467 mycorrhizae. *New Phytologist* 92, 75-87.

1468 Sollins, P., Swanston, C., Kleber, M., Filley, T., Kramer, M., Crow, S., Caldwell, B.A.,
 1469 Lajtha, K., Bowden, R., 2006. Organic C and N stabilization in a forest soil: Evidence
 1470 from sequential density fractionation. *Soil Biology and Biochemistry* 38, 3313-3324.

1471 Sorensen, L.H., 1972. Role of amino acids in the formation of soil organic matter. *Soil*
 1472 *Biology and Biochemistry* 4, 245-253.

1473 Sorensen, L.H., 1975. The influence of clay on the rate of decay of amino acids
 1474 metabolites synthesized in soil during decomposition of cellulose. *Soil Biology and*
 1475 *Biochemistry* 7, 171-177.

1476 Sorensen, L.H., 1977. Factors affecting the biostability of metabolic materials in soil. In:
 1477 *Soil Organic Matter Studies*, Vol. 2. International Atomic Energy Agency, Vienna, pp. 3-
 1478 14.

1479 Sorensen, L.H., 1981. Carbon- nitrogen relationships during the humification of cellulose
 1480 in soils containing different amount of clay. *Soil Biology and Biochemistry* 13, 313-321.

1481 Sorensen, L.H., Paul, E.A., 1971. Transformation of acetate carbon into carbohydrate and
 1482 amino acid metabolites during decomposition in soil. *Soil Biology and Biochemistry* 3,
 1483 173-180.

1484 Spence, A., Simpson, A., Menally D.J., Moran B.W., McFaul, M.V., Hart, K., Paull, B.
 1485 Kelleher, B.P., 2011. The degradation characteristics of microbial biomass in soil.
 1486 *Geochimica Cosmochimica Acta* 75, 2571-2581.

1487 Steinweg, J.M., Plante, A.F., Conant, R.T., Paul, E.A., Tanaka, D., 2008. Patterns of
 1488 substrate utilization during long-term incubations at different temperatures. *Soil Biology*
 1489 *and Biochemistry* 40, 2722-2728.

1490 Stevenson, F.J., 1994. *Humus Chemistry. Concepts, Composition, Reactions*. John Wiley
 1491 and Sons, New York.

1492 Stout, J D., Goh, K.M., Rafter, T.A., 1981. Chemistry and turnover of naturally occurring
 1493 resistant organic compounds in soil. In: Paul, E.A., Ladd, J.N. (Eds.), Soil Biochemistry,
 1494 vol. 5. Marcel Dekker, New York, pp. 1-74.

1495 Stursova, M., Zifcafova, I., Leigh, M.B., Burgess, R., Baldrian, P., 2012. Cellulose
 1496 utilization in forest litters and soils: identification of bacterial and fungal decomposers.
 1497 FEMS Microbiology Ecology 80, 735-746.

1498 Tarnocai, C., Canadell, J.G., Schuur, E.A.G., Kuhry, P., Mazhitova, G., Zimov, S., 2009.
 1499 Soil organic carbon pools in the northern circumpolar permafrost region. Global
 1500 Biogeochemical Cycles 23, GB2023.

1501 Taylor, D.L., Sinsabaugh, R., 2015. The soil fungi, occurrence, phylogeny and ecology.
 1502 In: Paul, E.A. (Ed.), Soil Microbiology, Ecology and Biochemistry, fourth ed. Academic
 1503 Press, San Diego, pp. 77-110.

1504 Thies, J., 2015. Molecular approaches to studying the soil biota. In: Paul, E.A. (Ed.), Soil
 1505 Microbiology, Ecology and Biochemistry, fourth ed. Academic Press, San Diego, pp.
 1506 151-168.

1507 Torn, M.S., Trumbore, S.E., Chadwick, O.A., Vitousek, P., Hendricks, D.M., 1997.
 1508 Mineral control of soil organic matter storage and turnover. Nature 389, 170-173.

1509 Trumbore, S.E., 1993. Comparison of carbon dynamics in tropical and temperate soils
 1510 using radiocarbon measurements. Global Biogeochemistry 7, 275-290.

1511 Trumbore, S., 2009. Radiocarbon and soil carbon dynamics. *Annual Review of Earth and*
1512 *Planetary Sciences* 37, 47-66.

1513 van Veen, J.A., Paul, E.A., 1981. Organic carbon dynamics in grassland soil. Background
1514 simulation and model verification. *Canadian Journal of Soil Science* 61, 185-201.

1515 Vestal, J.R., White, D.C., 1989. Lipid analysis in microbial ecology. *Bioscience* 39, 535-
1516 541.

1517 von Lützow, C., Kögel-Knabner, I., Elsk Schmidt, K., Flessa, H., Guggenberger, G.,
1518 Matzner, E., Marschner, B., 2007. SOM fractionation methods: Relevance to functional
1519 pools and to stabilization mechanisms. *Soil Biology and Biochemistry* 39, 2183-2207.

1520 Voroney, R.P., van Veen, J.A., Paul, E.A., 1981. Organic carbon dynamics in grassland
1521 soils. 2. Model verification of cultivation effects and prediction of long term erosion
1522 influences. *Canadian Journal of Soil Science* 62, 211-224.

1523 Voroney, R.P., Anderson, D.W., Paul, E.A., 1989. Decomposition of wheat straw and
1524 stabilization of microbial products. *Canadian Journal of Soil Science* 69, 63-77.

1525 Wagner, G.H., 1968. Significance of microbial tissue to soil organic matter. In: *Isotopes*
1526 *and Radiation in Soil Organic Matter Studies*. International Atomic Energy Agency,
1527 Vienna, STI/PUB/190, pp. 197-204.

1528 Wagner, G.H., 1975. Microbial growth and carbon turnover. In: Paul, E.A., McLaren, D.
1529 (Eds.), *Soil Biochemistry*, vol. 3. Marcel Dekker, New York, pp. 269-306.

- 1530 Wagner, G.H., Muzorewa, E.I., 1977. Lipids of microbial origin soil organic matter in
1531 soil organic matter studies. Proceedings of a Symposium. Braunshweig. International
1532 Atomic Energy Agency, Vienna, pp. 99-104.
- 1533 Waide, J.S., 1975. Hydroxyamides in soil systems. In: Paul, E.A., McLaren, A.D. (Eds.),
1534 Soil Biochemistry, vol. 4, Marcel Dekker, New York, pp. 65-102.
- 1535 Waksman, S.A., 1932. Principles of Soil Microbiology. Williams and Wilkins, Baltimore.
- 1536 Waksman, S.A., 1938. Humus - Origin, Chemical, Composition, and Importance in
1537 Nature, second ed. Williams and Wilkins, Baltimore.
- 1538 Waksman, S.A., Tenney, F.G., 1927. The composition of natural products materials and
1539 their decomposition in soil. Soil Science 24, 317-333.
- 1540 Wallenstein, M.D., Haddix, M.L., Ayres, E., Steltzer, H., Magrini-Baire, K.A., Paul,
1541 E.A., 2013. Litter chemistry change more rapidly when decomposed at home but
1542 converges during decomposition-transformation. Soil Biology and Biochemistry 57, 311-
1543 319.
- 1544 Warembourg, F.R., Paul, E.A., 1977. Seasonal transfers of assimilated ^{14}C in grassland:
1545 Plant production and turnover, soil and plant respiration. Soil Biology and Biochemistry
1546 9, 295-301.
- 1547 Warkentin, B.P. (Ed.), 2006. Footprints in the Soil. Elsevier, Amsterdam.

1548 Webley, D.M., Jones, D., 1971. Biological transformations of microbial residues in soil.
 1549 In: McLaren, A.J., Skujins, M.J., (Eds.) Soil Biochemistry, Vol. 2. Marcel Dekker, New
 1550 York, pp. 446-485.

1551 Weintraub, M.N., Schimel, J., 2003. Interactions between carbon and nitrogen
 1552 mineralization and soil organic matter chemistry in Arctic tundra soils. *Ecosystems* 6,
 1553 120-143.

1554 Yuan, G., Theng, B.K.G., 2012. Clay-organic reactions in the soil environment. In:
 1555 Huang, P.M., Li, Y., Sumner, M. (Eds.), *Handbook of Soil Science: Resource*
 1556 *Management and Environmental Impacts*, second ed. CRC Press, Boca Raton, pp 2-1 to
 1557 2-30.

1558 Zegouagh, Y., Derenne, S., Dignac, M.F., Baruiso, E., Marriott, A., Largeau, C., 2004.
 1559 Demineralization of a crop soil by mild hydrofluoric acid treatment: Influence on organic
 1560 matter composition and pyrolysis. *Journal of Analytical and Applied Pyrolysis* 71, 119-
 1561 135.

1562 Zwart, K.B., Burgers, S., Bloem, J., Bouwman, L., Brussard, L., Lebbink, G., Didden,
 1563 W., Marinisen, J., Vreeken-Buijs, M., de Ruiter, P., 1994. Population dynamics in the
 1564 belowground food webs of two different agricultural systems. *Agriculture, Ecosystems*
 1565 *and Environment* 51, 187-198.

1566

1567   Figure Captions

1568   Figure 1. The biotic and abiotic controls on soil organic matter formation and dynamics
1569   are interactive and highly integrated.

1570   Figure 2. Suggested structures for soil organic matter. The flocculation process of
1571   dissolved SOM with metals, such as Fe, is one mechanism leading to stable soil C. The
1572   flocs interact with SOM on minerals and aggregates. The SOM interacts with minerals by
1573   forming layers of different chemical properties (adapted from Horwath, 2015).

1574   Figure 3. The relative decomposition of ^{14}C glucose and ^{14}C wheat straw in a northern
1575   Prairie soil (from Voroney et al., 1989).

1576   Figure 4. Decay rates and pool sizes of microbial biomass and microbial products present
1577   in soil after growth on ^{14}C glucose in a laboratory incubation (from Paul and Voroney,
1578   1980).

1579   Figure 5. Soil organic matter pools (0-25cm) and atmospheric C fluxes (% of total)
1580   through bacteria, fungi fauna, symbionts, and roots for a poplar plantation and corn-
1581   soybean rotation at KBS-LTER MI.

Table 1. The global area, abiotic controls, net primary productivity, C inputs, terrestrial C, and turnover rates of soil C (adapted from Batjes, 2014; De Deyn et al., 2008; Horwath, 2015).

Biome	Area	MAP	MAT	Mean soil C	NPP	C inputs	Terrestrial C Stock (PgC)	Gross SOC turnover time yr	
	% of Total	(mm) yr ⁻¹	°C	(kg m ⁻²)	(PgCyr ⁻¹)	(kg m ⁻² yr ⁻¹)	Soil (0-3m)	0-20 cm	0-300 cm
Tropical forests	12.2	1400-4500	23-28.5	12.0	20.1	2.03	692	5.3	22
Temperate forests	9.5	750-2500	9-14.5	8.7	7.4	0.85	262	9.5	26
Boreal forests	8.8	600-1800	-5-5	16.4	2.4	0.50	150	10	27
Tropical savannas	19.0	500-1350	18-28	5.4	13.7	0.48	345	6.2	30
Temperate grasslands	7.1	450-1400	-2-15	13.3	5.1	0.30	172	15.1	63
Mediterranean shrubland	3.1	400-600	15-20	7.6	1.3	0.46	124	16.7	70
Deserts	14.4	125-500	-4-25	3.4	3.2	0.08	208	25.4	144
Tundra	7.0	250-1500	-15--5	19.6	0.5	0.10	818	52.3	165
Croplands	16.8	150-4500	-3-28.5	7.9	3.8	0.48	248	6.3	24
Wetlands g	2.2	250-4500	-2-28.5	72.3	4.3	0.17	450	150	945
Total	100			16.7	61.8	0.05	3051	NA	NA

MAP = mean annual precipitation; MAT = mean annual temperature; NPP = net annual primary productivity for the biome; NA = not applicable

Table 2. Recovery of ^{14}C labeled C in different components of a poplar plantation (data from Horwath and Paul, 1994).

	Total C		^{14}C distribution	
			At labeling	After 328 days
	gC m ⁻²	% of total	%	%
Leaves/litter	448	37.0	24.0	0.2
Stems/branches	520	44.0	48.0	26.0
Roots < 0.5 mm	52	4.4	1.7	1.3
Roots > 0.5 mm	166	14.0	8.1	8.4
Root – soil respiration	--	--	7.7	15.4
Microbial biomass	103	1.5	0.4	0.3
Soluble C	75	--	--	--
Soil C	6788	--	1.2	1.4

Table 3

Dynamics of soil fractions of a northern cultivated grassland (Mollisol) with 5.6% SOC (adapted from Campbell et al., 1967; Anderson and Paul, 1984; Paul et al., 2004).

Fraction	% SOC	MRT yr	Fraction	% SOC	MRT yr
LF	5	3	Whole soil	100	870
POM	10	12	Fine clay	8	170
Acid extract	14	325	Course clay	31	1255
Fulvic acids	15	495	Fine silt	29	968
Humic acids	40	1235	Coarse silt	25	800
Humin	31	1140	Non-hyd. Humin	23	1230
Non-hyd. HA	29	1400	Non-hyd. SOC	57	1350
Hyd. HA	11	25	Ca humates	21	1410

LF = light fraction; POM = particulate organic matter; HA = humic acids; Non-hyd. = Non hydrolyzable

Table 4. Use of long-term incubation, acid hydrolysis, ^{13}C plant switches, and ^{14}C carbon dating to measure the kinetics of SOC pools in a cultivated, Colorado grassland soil with 1.1% C, a microbial biomass of $377\mu\text{g Cg}^{-1}$ soil and 110yr of cultivation-plant switch (data from Follett et al., 2007; Paul et al., 1997, 2011).

	% of total soil	% native vegetation	MRT
	C	remaining after cultivation	(yr)
Cult. Soil (^{14}C)	100	-	1015
Microbial, Labile	2	-	0.6
Microbial, Non-labile	2	-	14
Active Soil C (Inc)	8	-	11
Slow Soil C (Inc)	32	-	72
Resistant Soil C (^{14}C)	60	-	2658
Cult. Whole Soil (^{13}C)	100	70	186
Light Fraction (^{13}C)	11	16	108
POM (^{13}C)	8	17	110
Silt (^{13}C)	30	61	199
Coarse Clay (^{13}C)	22	81	304
Fine Clay (^{13}C)	15	85	540

(^{13}C) (^{14}C) =method of measurement

Table 5. The use of ^{13}C changes of Hoytville soil fractions during long-term incubation of a corn-field soil (1.96% SOC after 30yr continuous corn) derived from a forest to determine the C_3 and C_4 dynamics of the light fraction (LF), POM + sand, silt, and clay (data from Haile-Mariam et al., 2008; Lavallee et al., 2015).

	Whole Soil	LF	POM plus sand	Silt	Clay
SOC day 0 (%)	100.0	3.5	11.0	23.0	53.0
SOC day 800 (%)	100.0	1.0	14.0	26.0	51.0
C:N day 0	9.6	20.0	15.0	10.0	6.5
C_4 C from corn (%)	23.0	84.0	56.0	20.0	11.0
C_4 MRT yr	18.2	3.7	7.8	12.8	26.8
^{13}C Day 0 ‰	-22.1	-14.1	-17.8	-22.4	-23.5
C_3 MRT yr	173.0	17.0	38.0	139.0	261.0
Black C%	5.0	3.9	2.0	6.3	10.6

The SOC dropped from 1.9% C at day 0 to 1.6% C at day 800. The sum of fractions does not add to 100 because of SOC losses during the fractionation, especially during the density fractionation.

Table 6. The role of depth and non-hydrolyzable C on the dynamics of SOC (data from Paul et al., 2001a).

Soil Type %	Depth	SOC %	SOC MRT	NHC %	NHC	CO ₂
	cm		yr		MRT yr	MIN%
Northern US	0-20	1.75	792	50	2180	6.8
Prairie						
(Mollisols)	25-50	0.86	2860	42	5881	5.6
	50-100	0.36	5260	29	10480	4.4
E. Dec. Forest	0-20	1.31	794	45	1928	6.3
(Alfisols)	25-50	0.47	2113	35	4856	7.2
	50-100	0.26	4128	34	8520	6.9

MIN=Mineralized. All MRT are based on carbon dating

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5