1 **Iron addition to soil specifically stabilized lignin**

2 Steven J. Hall*¹, Whendee L. Silver², Vitaliy I. Timokhin³, Kenneth E. Hammel^{4,5}

3
4 ¹Department of Ecology, Evolution, and Organismal Biology, Iowa State University, Ames, IA,
5 50011

6 ²Department of Environmental Science, Policy, and Management, University of California,
7 Berkeley, CA 94720

8 ³Department of Biochemistry, Wisconsin Energy Institute, University of Wisconsin, Madison,
9 WI 53706

10 ⁴US Forest Products Laboratory, Madison, Wisconsin 53726

11 ⁵Department of Bacteriology, University of Wisconsin, Madison, Wisconsin 53706

12
13 Corresponding author*: stevenjh@iastate.edu, 515-294-7650

14
15
16 Short communication for Soil Biology and Biochemistry

17
18
19
20 **Abstract**

The importance of lignin as a recalcitrant constituent of soil organic matter (SOM) remains contested. Associations with iron (Fe) oxides have been proposed to specifically protect lignin from decomposition, but impacts of Fe-lignin interactions on mineralization rates remain unclear. Oxygen (O₂) fluctuations characteristic of humid tropical soils drive reductive Fe dissolution and precipitation, facilitating multiple types of Fe-lignin interactions that could variably decompose or protect lignin. We tested impacts of Fe addition on ¹³C methoxyl-labeled lignin mineralization in soils that were exposed to static or fluctuating O₂. Iron addition suppressed lignin mineralization to 21% of controls, regardless of O₂ availability. However, Fe addition had no effect on soil CO₂ production, implying that Fe oxides specifically protected lignin methoxyls but not bulk SOM. Iron oxide-lignin interactions represent a specific mechanism for lignin stabilization, linking SOM biochemical composition to turnover via geochemistry.

Key words: carbon stabilization, iron, lignin, recalcitrance, redox, soil organic matter

Article text

The new paradigm of soil organic matter (SOM) dynamics posits that biochemical differences among compounds that influence short-term decomposition do not impact long-term persistence in SOM (Grandy and Neff, 2008; Marschner et al., 2008; Schmidt et al., 2011). These findings challenged the doctrine of humus formation, namely that recalcitrant biomolecules derived from lignin contribute disproportionately to SOM (Bollag et al., 1997). In support of the new paradigm, several studies reported that lignin might decompose as fast or faster than bulk SOM (Dignac et al., 2005; Gleixner et al., 1999; Guggenberger et al., 1994;

Heim and Schmidt, 2007; Kiem and Kögel-Knabner, 2003). Yet, these findings are difficult to reconcile with other studies demonstrating preferential association of aromatic lignin constituents with minerals, especially Fe oxides (Chorover and Amistadi, 2001; Gu et al., 1995; Hernes et al., 2013; Huang et al., 1977; Kaiser, 2003; Riedel et al., 2013). These associations would presumably decrease lignin mineralization, as has been observed for coprecipitates of lignin and ferrihydrite (Eusterhues et al., 2014). Thus, a fundamental question remains: can associations with minerals, particularly Fe oxides, decrease the decomposition of lignin relative to bulk SOM?

Iron-lignin interactions in soils are also complicated by potential *stimulation* of decomposition by solid-phase and soluble Fe. Fenton reactions driven by coupled biotic/abiotic Fe redox cycling can decompose lignin (Yelle et al., 2011), and may be important in Fe-rich soils that experience fluctuating O₂ (Hall et al., 2015; Hall and Silver, 2013). Also, Fe minerals such as goethite can abiotically oxidize SOM (Chorover and Amistadi, 2001). Finally, Fe reduction dominates anaerobic metabolism in many Fe-rich soils (Dubinsky et al., 2010), solubilizing Fe and exposing adsorbed or co-precipitated organic matter to enzymatic attack.

Our previous work (Hall et al., 2015) demonstrated that redox fluctuations increased the contribution of methoxyl groups in lignin to respiration from a tropical forest soil relative to static aerobic conditions. Here, we report an additional experiment where soils were amended with aqueous Fe(II) to simulate a highly reduced microsite, and then exposed to aerobic or fluctuating aerobic/hypoxic conditions. Added Fe(II) reacted with O₂ to form Fe(III) oxides, a portion of which was again reduced under subsequent hypoxic conditions. We evaluated three alternative hypotheses: H1) the precipitation of Fe oxides following Fe(II) oxidation disproportionately protects lignin relative to bulk SOM; H2) precipitation of Fe oxides protects

lignin and bulk SOM equally; H3) oxidation of added Fe(II) stimulates organic matter decomposition.

Soil was sampled from a tropical montane forest in the Luquillo Experimental Forest, Puerto Rico (Bisley Watershed), with 3500 mm of annual precipitation and mean annual temperature of 22 °C (Hall and Silver, 2015). This Aquic Hapludox was derived from volcaniclastic sediments and contained substantial short-range-order Fe oxides (24 mg Fe g⁻¹) and Fe(II) (0.2 – 0.8 mg g⁻¹). Iron(II) concentrations were much higher in reducing microsites, occasionally exceeding 5 mg g⁻¹ (Hall and Silver, 2013).

A macromolecular synthetic guaiacyl lignin labeled with ¹³C at the methoxyl position was produced (Kirk et al., 1975) to provide a sensitive index of mineralization. Lignin was precipitated onto ground leaf litter and gently homogenized with soils (20 mg lignin and 1000 mg soil) following Hall et al. (2015). We used a synthetic ¹³C-labeled lignin because these polymers yield similar long-term mineralization estimates as plant-derived lignins (Haider and Martin, 1981), while eliminating the labeled polysaccharides that inevitably contaminate labeled lignins produced *in planta* (Crawford, 1981). We recognize that the physical state of synthetic lignins does not completely represent that of natural lignins, which are intimately associated with other organic compounds (e.g. hemicellulose) in plant biomass. However, the fact that lignin biodegradation is a highly nonspecific oxidative process (Ruiz- Dueñas and Martínez, 2009) indicates that the precise spatial orientation of the lignin is probably not a critical factor, especially considering that extensive fragmentation of litter occurs during its breakdown in soils.

Samples were incubated under one of four headspace and Fe addition treatment combinations (n = 5 per treatment): static aerobic + aqueous Fe(II), static aerobic + water,

fluctuating aerobic/hypoxic + aqueous Fe(II), fluctuating aerobic/hypoxic + water. The fluctuating aerobic/hypoxic treatment experienced three days of aerobic conditions followed by four days of hypoxia (N₂ atmosphere). Before the experiment, samples received either N₂-flushed water (controls), or 5 mg Fe(II) g soil⁻¹ as aqueous FeCl₂ (Fe addition treatment) in an anaerobic chamber. Total CO₂ and ¹³CO₂ production were measured by gas chromatography and isotope ratio mass spectrometry over 3-4 day intervals. Additional subsamples were destructively sampled for Fe(II) analysis in 0.5M HCl using a modified ferrozine method (Hall et al., 2015).

In the Fe addition treatment, most (> 70 %) of added Fe(II) oxidized to Fe(III) or precipitated in non-HCl extractable phases within the first three days (Fig. 1). Iron addition increased subsequent Fe(II) production during the hypoxic phase of the fluctuating treatment 2-5 fold relative to the control (Fig. 1), likely due to the facile reduction of newly-formed short-range-ordered Fe(III) oxides (i.e., phases with nm-scale structure). This indicated that the availability of short-range-ordered Fe phases likely limited Fe reduction even in this Fe-rich soil. Iron(II) decreased over time in all treatments, and was ultimately similar between Fe addition and control treatments.

Iron addition consistently and significantly ($p < 0.0001$) decreased the contribution of lignin methoxyl C to soil respiration (p_{lignin}) to 15-55 % of the control (Fig. 2a). Hypoxia also decreased p_{lignin} in both treatments on days 7 and 14 ($p < 0.05$; Fig. 2a). However, O₂ effects were dwarfed by Fe addition, especially for cumulative mineralization (Fig. 2b). Iron-amended soils cumulatively mineralized only 21 % of lignin methoxyls relative to the controls ($p < 0.0001$). Cumulative mineralization did not differ between O₂ treatments (Fig. 2b).

We modeled cumulative lignin methoxyl mineralization in the aerobic treatments (Fig. 2b) using first-order decay after discarding the first observation ($R^2 = 0.93$ and 0.92 , for Fe addition and controls, respectively). A two-pool model did not improve the Akaike Information Criterion. Decomposition constants for Fe addition and control soils measured 0.0191 ± 0.0003 and $0.0762 \pm 0.0009 \text{ y}^{-1}$, respectively, implying mean turnover times of 52 and 13 y. These were similar to previous estimates of total lignin turnover in other ecosystems (10-40 y; Bahri et al., 2008; Heim and Schmidt, 2007; Rasse et al., 2006), and of slow-pool mineral-associated C in this ecosystem (11 – 26 y; Hall et al., 2015b). Because we measured the most labile constituent of lignin, its methoxyl group (Kirk et al., 1975), our results suggest that the more recalcitrant aromatic C of lignin could persist in soils for decades because of interactions with Fe oxides.

Trends in soil respiration differed from those of lignin mineralization (Fig. 2c,d). Total CO_2 production from Fe-amended soils was less than controls on day 3 ($p < 0.0001$), potentially due to direct toxicity of soluble Fe or reactive oxygen species produced during Fe(II) oxidation. Both mechanisms may have been important given the Fe(II) concentrations added here (Dunning et al., 1998). However, by day 10 Fe addition did not affect CO_2 production (Fig. 1c). Cumulative CO_2 production was similar between Fe addition and controls ($p = 0.15$; Fig. 1d). The similar CO_2 production from Fe addition and control treatments suggests no general suppression of organic matter mineralization upon interaction with Fe oxides as observed elsewhere (Eusterhues et al., 2014). Rather, availability of Fe(III) oxides as a terminal electron acceptor (Sutton-Grier et al., 2011), or shifts in pH, dissolved organic C, or reactive oxygen species known to be associated with Fe reduction and oxidation (Hall and Silver, 2013; Thompson et al., 2006) might have sustained CO_2 production despite protective effects of Fe-C interactions.

Our data thus support H1: Fe addition disproportionately inhibited the mineralization of lignin methoxyls relative to bulk SOM, irrespective of O₂ availability. This is consistent with preferential interactions between Fe oxide minerals and aromatic lignin constituents (Gu et al., 1995; Kaiser, 2003; Riedel et al., 2013). Given that short-range-order Fe oxides are present in most soils (Reyes and Torrent, 1997), Fe-lignin interactions deserve attention as contributors to “hidden lignin” pools that have potentially been obscured due to methodological artifacts associated with common lignin measurements (Hernes et al., 2013). Lignins synthesized with labeled isotopes provide one method to surmount these analytical challenges. We propose that consideration of specific interactions between recalcitrant biomolecules and soil geochemical constituents may prove useful in bridging old and new paradigms of SOM.

Acknowledgments

We thank H. Dang, C. Torrens, and the US Forest Service International Institute of Tropical Forestry. NSF grant DEB-1457805 to WLS and SJH, DOE Biological and Environmental Research grant DE-SC0006929 to KEH, and the NSF Luquillo Critical Zone Observatory (EAR-0722476) and LTER (DEB-0620910) supported this work.

References

- Bahri, H., Rasse, D.P., Rumpel, C., Dignac, M.-F., Bardoux, G., Mariotti, A., 2008. Lignin degradation during a laboratory incubation followed by ¹³C isotope analysis. *Soil Biology and Biochemistry* 40, 1916–1922. doi:10.1016/j.soilbio.2008.04.002
- Bollag, J.-M., Dec, J., Huang, P.M., 1997. Formation mechanisms of complex organic structures in soil habitats, in: Donald L. Sparks (Ed.), *Advances in Agronomy*. Academic Press, pp. 237–266.

154 Chorover, J., Amistadi, M.K., 2001. Reaction of forest floor organic matter at goethite, birnessite
 155 and smectite surfaces. *Geochimica et Cosmochimica Acta* 65, 95–109.
 156 doi:10.1016/S0016-7037(00)00511-1

157 Crawford, R.L., 1981. Lignin biodegradation and transformation. Wiley, New York.

158 Dignac, M.-F., Bahri, H., Rumpel, C., Rasse, D.P., Bardoux, G., Balesdent, J., Girardin, C.,
 159 Chenu, C., Mariotti, A., 2005. Carbon-13 natural abundance as a tool to study the
 160 dynamics of lignin monomers in soil: an appraisal at the Closeaux experimental field
 161 (France). *Geoderma, Mechanisms and regulation of organic matter stabilisation in soils*
 162 128, 3–17. doi:10.1016/j.geoderma.2004.12.022

163 Dubinsky, E.A., Silver, W.L., Firestone, M.K., 2010. Tropical forest soil microbial communities
 164 couple iron and carbon biogeochemistry. *Ecology* 91, 2604–2612. doi:10.1890/09-1365.1

165 Dunning, J.C., Ma, Y., Marquis, R.E., 1998. Anaerobic killing of oral Streptococci by reduced,
 166 transition metal cations. *Applied and Environmental Microbiology* 64, 27–33.

167 Eusterhues, K., Neidhardt, J., Hädrich, A., Küsel, K., Totsche, K.U., 2014. Biodegradation of
 168 ferrihydrite-associated organic matter. *Biogeochemistry* 119, 45–50. doi:10.1007/s10533-
 169 013-9943-0

170 Gleixner, G., Bol, R., Balesdent, J., 1999. Molecular insight into soil carbon turnover. *Rapid*
 171 *Communications in Mass Spectrometry* 13, 1278–1283. doi:10.1002/(SICI)1097-
 172 0231(19990715)13:13<1278::AID-RCM649>3.0.CO;2-N

173 Grandy, A.S., Neff, J.C., 2008. Molecular C dynamics downstream: The biochemical
 174 decomposition sequence and its impact on soil organic matter structure and function.
 175 *Science of The Total Environment* 404, 297–307. doi:10.1016/j.scitotenv.2007.11.013

176 Gu, B., Schmitt, J., Chen, Z., Liang, L., McCarthy, J.F., 1995. Adsorption and desorption of
 177 different organic matter fractions on iron oxide. *Geochimica et Cosmochimica Acta* 59,
 178 219–229. doi:10.1016/0016-7037(94)00282-Q
 179 Guggenberger, G., Christensen, B.T., Zech, W., 1994. Land-use effects on the composition of
 180 organic matter in particle-size separates of soil: I. Lignin and carbohydrate signature.
 181 *European Journal of Soil Science* 45, 449–458. doi:10.1111/j.1365-2389.1994.tb00530.x
 182 Haider, K., Martin, J.P., 1981. Decomposition in soil of specifically ¹⁴C-labeled model and
 183 cornstalk lignins and coniferyl alcohol over two years as influenced by drying, rewetting,
 184 and additions of an available C substrate. *Soil Biology and Biochemistry* 13, 447–450.
 185 doi:10.1016/0038-0717(81)90032-8
 186 Hall, S.J., Silver, W.L., 2015. Reducing conditions, reactive metals, and their interactions can
 187 explain spatial patterns of surface soil carbon in a humid tropical forest. *Biogeochemistry*
 188 125, 149–165. doi:10.1007/s10533-015-0120-5
 189 Hall, S.J., Silver, W.L., 2013. Iron oxidation stimulates organic matter decomposition in humid
 190 tropical forest soils. *Global Change Biology* 19, 2804–2813. doi:10.1111/gcb.12229
 191 Hall, S.J., Silver, W.L., Timokhin, V.I., Hammel, K.E., 2015. Lignin decomposition is sustained
 192 under fluctuating redox conditions in humid tropical forest soils. *Global Change Biology*
 193 21, 2818–2828. doi:10.1111/gcb.12908
 194 Heim, A., Schmidt, M.W.I., 2007. Lignin turnover in arable soil and grassland analysed with two
 195 different labelling approaches. *European Journal of Soil Science* 58, 599–608.
 196 doi:10.1111/j.1365-2389.2006.00848.x

197 Hernes, P.J., Kaiser, K., Dyda, R.Y., Cerli, C., 2013. Molecular trickery in soil organic matter:
 198 hidden lignin. *Environmental Science & Technology* 47, 9077–9085.
 199 doi:10.1021/es401019n

200 Huang, P.M., Wang, T.S.C., Wang, M.K., Wu, M.H., Hsu, N.W., 1977. Retention of phenolic
 201 acids by noncrystalline hydroxy-aluminum and -iron compounds and clay minerals of
 202 soils. *Soil Science* 123, 213–219.

203 Kaiser, K., 2003. Sorption of natural organic matter fractions to goethite (α -FeOOH): effect of
 204 chemical composition as revealed by liquid-state ^{13}C NMR and wet-chemical analysis.
 205 *Organic Geochemistry* 34, 1569–1579. doi:10.1016/S0146-6380(03)00120-7

206 Kiem, R., Kögel-Knabner, I., 2003. Contribution of lignin and polysaccharides to the refractory
 207 carbon pool in C-depleted arable soils. *Soil Biology and Biochemistry* 35, 101–118.
 208 doi:10.1016/S0038-0717(02)00242-0

209 Kirk, T.K., Connors, W.J., Blear, R.D., Hackett, W.F., Zeikus, J.G., 1975. Preparation and
 210 microbial decomposition of synthetic [^{14}C]lignins. *Proceedings of the National Academy*
 211 *of Sciences (USA)* 72, 2515–2519.

212 Marschner, B., Brodowski, S., Dreves, A., Gleixner, G., Gude, A., Grootes, P.M., Hamer, U.,
 213 Heim, A., Jandl, G., Ji, R., Kaiser, K., Kalbitz, K., Kramer, C., Leinweber, P.,
 214 Rethemeyer, J., Schäffer, A., Schmidt, M.W.I., Schwark, L., Wiesenberger, G.L.B., 2008.
 215 How relevant is recalcitrance for the stabilization of organic matter in soils? *Journal of*
 216 *Plant Nutrition and Soil Science* 171, 91–110. doi:10.1002/jpln.200700049

217 Rasse, D.P., Dignac, M.-F., Bahri, H., Rumpel, C., Mariotti, A., Chenu, C., 2006. Lignin
 218 turnover in an agricultural field: from plant residues to soil-protected fractions. *European*
 219 *Journal of Soil Science* 57, 530–538. doi:10.1111/j.1365-2389.2006.00806.x

220 Reyes, I., Torrent, J., 1997. Citrate-ascorbate as a highly selective extractant for poorly
 221 crystalline iron oxides. *Soil Science Society of America Journal* 61, 1647–1654.
 222 Riedel, T., Zak, D., Biester, H., Dittmar, T., 2013. Iron traps terrestrially derived dissolved
 223 organic matter at redox interfaces. *Proceedings of the National Academy of Sciences*
 224 10101–10105. doi:10.1073/pnas.1221487110
 225 Ruiz- Dueñas, F.J., Martínez, Á.T., 2009. Microbial degradation of lignin: how a bulky
 226 recalcitrant polymer is efficiently recycled in nature and how we can take advantage of
 227 this. *Microbial Biotechnology* 2, 164–177. doi:10.1111/j.1751-7915.2008.00078.x
 228 Schmidt, M.W.I., Torn, M.S., Abiven, S., Dittmar, T., Guggenberger, G., Janssens, I.A., Kleber,
 229 M., Kogel-Knabner, I., Lehmann, J., Manning, D.A.C., Nannipieri, P., Rasse, D.P.,
 230 Weiner, S., Trumbore, S.E., 2011. Persistence of soil organic matter as an ecosystem
 231 property. *Nature* 478, 49–56. doi:10.1038/nature10386
 232 Sutton-Grier, A.E., Keller, J.K., Koch, R., Gilmour, C., Megonigal, J.P., 2011. Electron donors
 233 and acceptors influence anaerobic soil organic matter mineralization in tidal marshes.
 234 *Soil Biology and Biochemistry* 43, 1576–1583. doi:10.1016/j.soilbio.2011.04.008
 235 Thompson, A., Chadwick, O.A., Boman, S., Chorover, J., 2006. Colloid mobilization during soil
 236 iron redox oscillations. *Environmental Science & Technology* 40, 5743–5749.
 237 doi:10.1021/es061203b
 238 Yelle, D.J., Wei, D., Ralph, J., Hammel, K.E., 2011. Multidimensional NMR analysis reveals
 239 truncated lignin structures in wood decayed by the brown rot basidiomycete *Postia*
 240 *placenta*. *Environmental Microbiology* 13, 1091–1100. doi:10.1111/j.1462-
 241 2920.2010.02417.x
 242

243 **Figure captions:**

244 Figure 1: Soil Fe(II) concentrations in 0.5M HCl extractions by treatment and day of experiment
245 (n = 1 per treatment/day). The grey bars represent the hypoxic phase of the fluctuating headspace
246 treatment. In the Fe addition treatments, 5000 $\mu\text{g Fe(II) g}^{-1}$ soil were added at the beginning of
247 the experiment.

248 Figure 2: a) Production of CO₂ from ¹³C methoxyl lignin as a percent of total soil respiration. b)
249 Total mineralization of methoxyl lignin C. c) Soil CO₂ production by sampling date. d)
250 Cumulative CO₂ production. Bars represent standard errors with n = 5 per treatment. See the text
251 for a description of the treatments.