Title: Methodological uncertainty in estimating carbon turnover times of soil fractions

Running head: Carbon turnover times of soil fractions

Authors: Wenting Feng^{1*}, Zheng Shi¹, Jiang Jiang^{1, 2}, Jianyang Xia³, Junyi Liang¹, Jizhong Zhou¹, Yiqi Luo¹

¹ Department of Microbiology & Plant Biology, University of Oklahoma, Norman, Oklahoma, USA

² Key Laboratory of Soil and Water Conservation and Ecological Restoration in Jiangsu Province, Forestry College of Nanjing Forestry University, Nanjing, China

³ School of Ecological and Environmental Sciences, East China Normal University, Shanghai, China

Corresponding author: Wenting Feng, wenting.fwt@gmail.com, tel: +1 405 325 6519, fax: +1 405 325 7519, 101 David L Boren BLVD, Norman, OK 73019; Yiqi Luo, yluo@ou.edu, tel: +1 405 325 1651, fax: +1 405 325 7519, 101 David L Boren BLVD, Norman, OK 73019

Abstract

Improving predictions of soil organic carbon (SOC) dynamics by multi-compartment models requires validation of turnover times of different SOC pools. Techniques such as laboratory incubation and isotope analysis have been adopted to estimate C turnover times, yet no studies have systematically compared these techniques and assessed the uncertainties associated with them. Here, we tested whether C turnover times of soil fractions were biased by methodology, and how this changed across soil particle sizes and ecosystems. We identified 52 studies that quantified C turnover times in different soil particles either fractionated according to aggregate size (*e.g.*, macro- versus micro-aggregates) or according to soil texture (*e.g.*, sand versus silt versus clay). C turnover times of these soil fractions were estimated by one of three methods: laboratory incubation (16 studies), $\delta^{13}\text{C}$ shift due to C_3 - C_4 vegetation change (25 studies), and ^{14}C dating (19 studies). All methods showed that C turnover times of soil fractions generally increase with decreasing soil particle size. However, estimates of C turnover times within soil fractions differed significantly among methods, with incubation estimating the shortest turnover times and ^{14}C the longest. The short C turnover times estimated by incubation are likely due to optimal environmental conditions for microbial decomposition existing in these studies, which is often a poor representation of field conditions. The ^{13}C method can only be used when documenting a successive C_3 versus C_4 vegetation shift. C turnover times estimated by ^{14}C were systematically higher than those estimated by ^{13}C , especially for fine soil fractions (*i.e.*, silt and clay). Overall, our findings highlight methodological uncertainties in estimating C turnover times of soil fractions, and correction factors should be explored to account for methodological bias when C turnover times estimated from different methods are used to parameterize soil C models.

Keywords: soil organic carbon; turnover; ^{13}C ; ^{14}C ; incubation; fraction

1. Introduction

Uncertainty in predicting carbon–climate feedbacks largely stems from poor representation of soil organic carbon (SOC) pools. This is an important consideration as SOC is the largest C pool in terrestrial ecosystems and perturbation of it strongly modulates climate change (Todd-Brown et al., 2013; Koven et al., 2015; Luo et al., 2016). SOC is heterogeneous in terms of composition, structure, location, and stabilization mechanism (Stevenson, 1994; Sollins et al., 1996; Schmidt et al., 2011; Lehmann & Kleber 2015). Conventional soil C models classify SOC into multiple conceptual pools with different turnover times based on their resistance to microbial decomposition (Jenkinson & Rayner, 1977; Parton et al., 1987). A growing body of research calls for mechanistic representations of SOC processes in Earth System Models, such as protection by physical isolation and mineral sorption (Sulman et al., 2014; Wieder et al., 2014; Tang & Riley, 2015). Therefore, attention should be paid to physically fractionated SOC fractions which are measurable and could represent soil organic matter (SOM) protection mechanisms (Christensen, 1996; von Lützow et al., 2007; Schmidt et al., 2011). Quantifying C turnover times of these soil fractions is important for models which integrate explicit mineral protection processes. Until now there has been no consensus on the turnover times of various measurable SOC fractions, due to various methodologies being used to estimate C turnover times.

There are three commonly used methods for assessing SOC turnover times: the laboratory incubation (Christensen, 1987), shifts in natural ^{13}C abundance after C_3 - C_4 vegetation change (Balesdent et al., 1987), and ^{14}C dating (O'Brien & Stout, 1978; Trumbore, 2000). The laboratory incubation directly quantifies biological decomposition of isolated soil fractions under controlled optimal conditions. This method is easy to conduct and has been widely used. In contrast, the ^{13}C and ^{14}C methods trace C isotopes during decomposition and stabilization

processes to estimate C turnover times (O'Brien & Stout 1978; Balesdent et al., 1990). The ^{13}C method can only be used in studies where there are $\delta^{13}\text{C}$ shifts after years of successive $\text{C}_3\text{-C}_4$ vegetation change and requires careful C inventory measurements of disturbed and undisturbed soils (Balesdent et al., 1987; Zhang et al., 2015). The ^{14}C dating method assumes that SOC fractions are at equilibrium between input and decay, and that all the C inputs to soils enter the system at the same time or are constant (Trumbore, 1993; Bruun et al., 2005). These assumptions are often not met in reality and soil ^{14}C is expensive to measure. Due to these differences in methodology, the three methods likely generate different estimates of SOC turnover times. For instance, the turnover times of mineral associated organic matter (MOM) at 0-10 cm depth has been reported to be 8-43 years using the laboratory incubation method (Rabbi et al., 2014), 53-63 years using the ^{13}C abundance after $\text{C}_3\text{-C}_4$ vegetation change (Dalal et al., 2013; Liang et al., 2014), and 52-381 years when using ^{14}C dating (Budge et al., 2011).

Bulk soil can be separated into soil fractions using the physical, chemical, density, and combined fractionation methods, among which the physical fractionation is able to generate soil fractions with distinct C turnover times (Christensen, 2001; Mikutta et al., 2006; von Lützow et al., 2007). Variation in C turnover times results from different SOC protection mechanisms associated with soil particles as well as inconsistent methods used to estimate C turnover times (Bird et al., 2002; Tan et al., 2013; Yonekura et al., 2013; Beniston et al., 2014). Physically fractionated soil particles are often obtained according to soil aggregate size or soil texture. According to soil aggregates size, C in macro-aggregates (*i.e.*, coarse organic matter, COM) turns over fast, while C in the micro-aggregates (*i.e.*, fine organic matter, FOM) and MOM is supposed to represent C that is primarily protected by physical isolation and mineral matrix, respectively (Six et al., 1998; Baldock & Skjemstad, 2000; von Lützow et al., 2007). According

to soil texture, C in the sand fraction has a short turnover time and C associated with the silt and clay fractions is considered as mineral associated OM in models (Parton et al., 1987; Beniston et al., 2014; Tang & Riley, 2014; Wieder et al., 2014). However, we still do not know whether different classifications to separate soil fractions can differentiate their C turnover times.

By synthesizing published studies, we compared C turnover times of physically fractionated soil particles (*i.e.*, COM – FOM – MOM or sand – silt – clay) across ecosystems. We aimed to test whether C turnover times of soil fractions estimated using the laboratory incubation, ^{13}C , and ^{14}C were different, and how this changed with soil particle size and ecosystems. We predicted that turnover times estimated using the laboratory incubation would be shorter than those using the C isotope methods, and that C turnover times based on soil fractions would increase with decreasing particle size.

2. Material & methods

2.1 Data sources

We searched the literature to find information that included: (1) used at least one of the following physically fractionated soil particles as study materials: macro-aggregates (coarse organic matter, COM, 250-2000 μm), micro-aggregates (fine organic matter, FOM, 20/53/63-250 μm), MOM (<20/53/63 μm), sand (20/53/63-2000 μm), silt (2-20/53/63 μm), and clay (<2 μm), and (2) measured CO_2 flux multiple times over the time course of laboratory incubations, or assessed C turnover rates or times based on the $\delta^{13}\text{C}$ difference after years of successive $\text{C}_3\text{-C}_4$ vegetation change, or estimated mean residence times based on $\Delta^{14}\text{C}$ activity. Detailed information of the selected studies can be found in Table 1 and the supplementary materials (Supplementary material Table S1). We extracted information on 537 soil fractions from 52

studies around the world (Fig. 1). For all the studies identified, we also gathered the information regarding soil fraction classification used, the coordinates, climate, soil depth, soil type, and vegetation at soil sampling sites, and the mass proportion and organic C concentration or content of each soil fraction (Supplementary material Table S1).

2.2 Carbon turnover estimate

For the studies using laboratory incubations to estimate C turnover time, we generated a sub-dataset that included the following data for each soil fraction: the date of measurement, initial organic C concentration or content, and CO₂ respiration rate or cumulative CO₂ respiration at each time point. We used the two-pool rather than one-pool exponential decomposition model to estimate C turnover times of soil fractions, because C in soil fractions is not homogeneous and so the two-pool model could more accurately describe decomposition than the one-pool model (Derrien & Amelung, 2011). For comparison, we converted values of cumulative CO₂ respiration from the original unit (mg CO₂-C g⁻¹ sample) to mg CO₂-C per gram of initial organic C concentration of a sample.

$$C_t = f_l \times (1 - e^{-k_l \times t}) + (1 - f_l) \times (1 - e^{-k_s \times t}) \quad \text{Equation (1)}$$

was used to estimate C turnover times of soil fractions, where C_t is the cumulative CO₂ respired, f_l is the proportion of labile SOC pool, and k_l and k_s are the decomposition constants of labile and stable SOC pools. The turnover times of labile (τ_l) and stable (τ_s) SOC are the reciprocal of k_l and k_s , respectively. Given that stable SOC accounts for a large proportion of total SOC and τ_l is similar for the studied soil fractions from a variety of ecosystems, using τ_s instead of τ_l is much more representative to characterize C turnover of the entire SOC. Therefore, τ_s values of soil fractions were used to compare whether the three methods provide different C turnover times

values. Parameters in the two-pool model were estimated using probabilistic inversion approach (Xu et al., 2006; Weng & Luo, 2011), which was performed using the Metropolis-Hastings (M-H) algorithm – a Markov Chain Monte Carlo (MCMC) technique (Metropolis et al., 1953; Hastings, 1970). Rationale and details about this technique can be seen in Schädel et al. (2013).

For the studies using $\delta^{13}\text{C}$ after $\text{C}_3\text{-C}_4$ vegetation change to estimate C turnover time, we collected the data of turnover time (τ , year) or decomposition constant (k , year^{-1}) for all of the six soil fractions (*i.e.*, COM-FOM-MOM and sand-silt-clay). In the studies where neither k nor τ were reported, we calculated k using equations (2) or (3) according to the data available in selected studies.

$$k = \frac{-\ln(\text{proportion of old C})}{\text{period of C}_3\text{-C}_4 \text{ vegetation change}} \quad \text{Equation (2)}$$

$$A_t = A_0 \times e^{-k \times t} \quad \text{Equation (3)}$$

where A_0 is the initial SOC stock of soil fraction, and A_t is old C stock of soil fraction at time t in years since $\text{C}_3\text{-C}_4$ vegetation change (Balesdent et al., 1987).

Logarithmic transformation of equation (3) is essentially the same as equation (2). But when studies only measure $\delta^{13}\text{C}$ twice before and after the $\text{C}_3\text{-C}_4$ vegetation change, we necessarily used equation (2) to calculate k . When studies measure $\delta^{13}\text{C}$ multiple times after the $\text{C}_3\text{-C}_4$ vegetation change, k was assessed using equation (3), due to higher confidence in estimates obtained with this equation. This is due to the fact that calculations of k using equation (2) overestimates when using $\delta^{13}\text{C}$ measured at an early stage after the $\text{C}_3\text{-C}_4$ vegetation change, and underestimates when measuring $\delta^{13}\text{C}$ at a late stage after the vegetation change (Skjemstad et al., 1990; Liao et al., 2006; Derrien & Amelung, 2011). Calculating k according to equation (2) is the two-point ^{13}C method, and the calculation according to equation (3) is the multi-point ^{13}C

method. According to these two calculation methods, we separated studies that report k or τ values to two groups, to test whether these two calculations generate different k estimates.

For studies that use ^{14}C dating techniques, there are also two distinct approaches to estimate C turnover times of soil fractions - the conventional ^{14}C model and the bomb ^{14}C model. The conventional ^{14}C method assumes that all C atoms in a sample entered soils at the same time and the measured SOC fraction is in steady state between input and decay (Talma & Vogel, 1993; Bruun et al., 2005), and calculates C turnover time (τ) by

$$\tau = \frac{1}{\lambda} \ln\left(\frac{A_{abs}}{A_t}\right) \quad \text{Equation (4)}$$

λ is the decay rate constant of ^{14}C , and A_{abs} is defined as 95% of the activity in 1950 of an oxalic acid standard, A_t is the ^{14}C activity of soil sample. But the assumptions in the conventional ^{14}C dating are mostly untrue for modern soils except for buried paleosols. Meanwhile, the bomb ^{14}C model uses the natural decay of atmospheric ^{14}C activity generated in the 1950s and 1960s bomb tests to estimate C turnover times (O'Brien & Stout, 1978; Trumbore, 1993; Rabbi et al., 2013). This model assumes that SOC decomposition follows the first order law and is at steady state, where C turnover time is described by

$$^{14}C_t = ^{14}C_{atm_{t-lag}} \times k + ^{14}C_{t-l} \times (1 - k - \lambda) \quad \text{Equation (5)}$$

where $^{14}C_t$ and $^{14}C_{t-l}$ are the $\Delta^{14}\text{C}$ activities at years t and $t-l$, $^{14}C_{atm_{t-lag}}$ is the $\Delta^{14}\text{C}$ of the atmosphere, k is the decomposition constant, and λ is the ^{14}C decay constant. Here, we grouped studies into those that used the conventional ^{14}C model or those that used the bomb ^{14}C model, aiming to find whether these two methods provide different C turnover times of soil fractions.

2.3 Statistical analysis

Multiple comparison was used to examine whether the laboratory incubation, ^{13}C , and ^{14}C methods generated different C turnover times for each of the six soil fractions, and to test whether C turnover times estimated by the same method are significantly different among COM, FOM, and MOM, and among the sand, silt, and clay fractions. In the multiple comparison to examine whether C turnover times estimated by the three methods were different, C turnover estimated by the two-point and multi-point ^{13}C methods were compiled, but only the estimates by the bomb ^{14}C model were used, since the two-point and multi-point ^{13}C methods did not generate significantly different estimates, but the conventional ^{14}C dating showed remarkably longer C turnover times compared to the bomb ^{14}C model. Since C turnover times were not normally distributed, a Mann-Whitney rank test was used for the multiple comparison by using the nparcomp R package (Konietschke et al., 2015). All differences were tested at the significance level of 0.05.

3. Results

Carbon turnover times differed with soil fractions and method used (Fig. 2). When bulk soils were separated into the COM, FOM, and MOM fractions, C turnover times estimated by the ^{13}C and ^{14}C methods were significantly longer than those using the laboratory incubation, but the estimates by the former two methods showed no significant difference (Fig. 2). The results of the laboratory incubation show that turnover times of stable SOC pool (mean $\pm$ SE) were significantly longer in the FOM (31.5 ± 12.9 yr) and MOM (30.9 ± 17.6 yr) fractions than in the COM fraction (8.6 ± 2.3 yr). C turnover times were significantly longer in the MOM (31.5 ± 12.9 yr) fraction than in the FOM and COM fractions when using the ^{13}C and ^{14}C methods (Fig. 2). When bulk soils were separated to the sand, silt, and clay fractions, C turnover times

estimated by the three methods were significantly different from each other, following the order: incubation $< {}^{13}\text{C} < {}^{14}\text{C}$ (Fig. 2). Regardless of estimate methods, C turnover times of the silt and clay fractions were similar to each other, both of which were significantly longer than those of the sand fraction (Fig. 2).

Using the ${}^{13}\text{C}$ after $\text{C}_3\text{-C}_4$ vegetation change, t-test results show that the multi-point and two-point calculation methods generated similar C turnover times of soil fractions, although estimates by the two-point method tended to be slightly lower (Fig. 3). Using the ${}^{14}\text{C}$ dating method, the two calculation methods (*i.e.*, ${}^{14}\text{C}$ conventional and ${}^{14}\text{C}$ bomb) provided significantly different values of C turnover times of soil fractions (Fig. 4). C turnover times of small soil particles (*i.e.*, FOM-MOM and silt-clay) estimated by the $\Delta{}^{14}\text{C}$ conventional method were 665-2047 years, compared to 149-431 years estimated by using the $\Delta{}^{14}\text{C}$ bomb model (Fig. 4).

4. Discussion

Estimations of C turnover times of measurable soil fractions are important for incorporation into newly emerging soil C models that explicitly include interactions between organic matter and soil minerals. Our study shows that C turnover times of physically fractionated soil particles generally increase with decreasing particle size, following the order: $\text{COM} \approx \text{FOM} < \text{MOM}$ and $\text{sand} < \text{silt} \approx \text{clay}$ (Fig. 2), suggesting that fine soil fractions (*i.e.*, FOM-MOM and silt-clay) allow a higher organic C preservation. These results agree with the reported range of C turnover times of these soil fractions: 3-203 years for the COM fraction, 1.2 to 374 years for the FOM fraction, 63-125 years for the MOM fraction, 8-1660 years for the silt fraction, and 33-4409 years for the clay fraction (Feller & Beare, 1997; Six et al., 2002; von

Lützow et al., 2007). Although other studies have addressed C turnover times across soil fractions (Christensen, 1987; Feller & Beare, 1997; Bird et al., 2002; Six et al., 2002; von Lützow et al., 2007; Rabbi et al., 2014), this study has the advantage of including a large sample size for each of the six soil fractions and for each C turnover estimate method (Fig. 1; Table 1). Moreover, these soil fractions are from a wide variety of ecosystem types and span a substantial latitudinal gradient (68.10° N to 40.38° S (Fig. 1)).

Although the patterns of how C turnover times change with soil particle size is similar regardless of the estimate methods, the laboratory incubation, the ^{13}C method, and the ^{14}C dating method provide different mean values of C turnover times of soil fractions (Fig. 2), in the order: incubation < ^{13}C < ^{14}C . This difference highlights methodological uncertainties in estimating C turnover times of soil fractions. Special attention should be paid when parameterizing soil C turnover times to simulate SOC dynamics. Short C turnover times estimated via laboratory incubation might be due to microbial decomposition rates at optimal temperature and moisture, unrealistic under climatic limitations present in natural systems. Sieving and rewetting soils, that routinely occurs before incubation, has been found to increase C mineralization (Fierer and Schimel, 2002; Miller et al., 2005), and thus could lead to the underestimate of C turnover times. In contrast, the ^{13}C and ^{14}C methods estimates SOC in the field where the climate likely constrains microbial decomposition. Another reason for short C turnover times of soil fractions estimated by the laboratory incubation could be that soils used in incubations are often from top soil layers, and C in shallow soils has shorter turnover times than deeper soils (Rumpel et al., 2002; Mathieu et al., 2015). Additionally, the fractionation procedure may redistribute C in different soil particles and accelerate C decomposition (Christensen, 1987; Parfitt & Salt, 2001; Benbi et al., 2014). So, C turnover times of soil particles estimated by the ^{13}C and ^{14}C methods

are likely more representative of actual values in field. However, laboratory incubations are still useful to elucidate how factors other than climate might affect C turnover.

When using the ^{13}C method to estimate C turnover times of soil fractions, the two-point and the multi-point calculations generate similar values, although the former method estimates were slightly shorter C turnover times than the latter (Fig. 3). This finding demonstrates that the repeatability is high when using the ^{13}C method to estimate C turnover times of soil fractions (Fig. 2). The multi-point ^{13}C method is recommended to calculate soil C turnover, because it generates results with higher confidence. Derrien & Amelung (2011) also found that multiple-time measurements of $\delta^{13}\text{C}$ are better for estimating C turnover times, because this method can assess C turnover times at both steady and non-steady states while the two time-point measurements cannot. This study suggests that when multiple-time point measurements of $\delta^{13}\text{C}$ of soil fractions are not available, two time-point measurements can be used as a substitute to give reasonable estimates of C turnover times.

Using $\Delta^{14}\text{C}$ to estimate C turnover times of soil fractions, caution should be exercised concerning the calculation approach used. We found that C turnover times of all studied soil fractions estimated by the conventional ^{14}C model were 4-5 folds longer than those by the bomb ^{14}C model (Fig. 4), and were also longer than the values used in current multi-compartment soil C models (Jenkinson & Rayon, 1977; Parton et al., 1987). Thus, C turnover times of soil fractions estimated by the bomb ^{14}C model are recommended when simulating SOC dynamics by using multi-compartment models. This estimate divergence probably results from different assumptions of these two models. The conventional ^{14}C model assumes that C in different soil fractions are formed directly from external C sources with the age of zero, but C in some soil particles (*e.g.*, the silt and clay sized particles) may be formed from the transfer of C in coarse

soil particles with the age larger than zero (Trumbore, 1993; Bruun et al., 2005). The bomb ^{14}C model that considers continuous C inputs to soils is more realistic, because it uses abundant ^{14}C derived from the 1950s bomb test as a tracer and the numerical solution to estimate C turnover times are more accurate when compared to the conventional ^{14}C method (Trumbore, 1993; Bruun et al., 2005). However, the steady state assumption may underestimate the turnover times of SOC fractions which need a long time to reach equilibrium (Bruun et al., 2005).

Even using the same estimate method, C turnover times of the same soil fraction still vary greatly (Fig. 2). This is likely because soils come from a variety of environments, where climate, vegetation, microbial community, and soil mineralogy and depth likely influence C turnover times in soils. Among these factors, soil depth is important in impacting C turnover time of soil fractions. We observed that at the same site, C turnover times of a given soil fraction generally increase with depth, regardless of the estimate method used (Skjemstad et al., 1990; Schöning & Kögel-Knabner, 2006; Yonekura et al., 2013; Dalal et al., 2013; Beniston et al., 2014; Liang et al., 2014). This finding is consistent with other studies (Rumpel et al., 2002; Rumpel & Kögel-Knabner, 2011; Mathieu et al., 2015). We did not observe longer C turnover times of soil fractions at high latitude than at low latitude. It is likely that local environments at studied sites, such as SOM chemistry and soil properties, cause large variations of C turnover times of soil fractions, which masks the influences of latitude and associated climate on C turnover times. The reason might also be that there is not sufficient data on C turnover times along a latitude gradient to generalize patterns of how it changes with climate.

Although our study has used soil fractions from locations worldwide to estimate C turnover times by the laboratory incubation, the $\delta^{13}\text{C}$ after $\text{C}_3\text{-C}_4$ vegetation change, and the ^{14}C dating (Fig. 1), we are aware that these soil fractions used for estimation by these three methods

are not the same or well paired. So, we cannot attribute the variations of C turnover times of soil fractions solely to different estimation methods. Other factors, such as temperature, precipitation, soil depth, and soil texture, that has been found to influence C turnover times of bulk soil might impact C turnover times of soil fractions as well (Carvalhais et al., 2014; Mathieu et al., 2015; Xu et al., 2016). To parameterize soil C turnover times in the multi-compartment models, we highly recommend studies that assess C turnover times of the same physically fractionated soil particles by using different methods.

This synthesis study compared C turnover times of physically fractionated soil fractions estimated using three methods: (1) laboratory incubation, (2) $\delta^{13}\text{C}$ after C_3 - C_4 vegetation change, and (3) ^{14}C dating. We found that estimated C turnover times of soil fractions differed significantly among methods. We suggested that the relatively fast soil C turnover time found by the incubation studies under optimal environmental conditions are likely an overestimate of C turnover rates under field conditions, as soil moisture and temperature are not always at optimum levels in nature. Estimates derived from $\delta^{13}\text{C}$ and $\Delta^{14}\text{C}$ are likely closer to actual C turnover rates found in the field. However, the ^{13}C method can only be used when there are detectable changes in $\delta^{13}\text{C}$ after years of successive C_3 versus C_4 vegetation change, and ^{14}C dating could more accurately estimate C turnover of soil fractions when soils are under steady-state conditions or ^{14}C inputs derived from atmosphere and vegetation are well documented. It is noticeable that when using the ^{14}C dating method the presence of black C in soils could bias C turnover times of coarse organic matter, which is considered to be labile and has short turnover times (Baisden et al., 2002; Leifeld, 2008; Leifeld et al., 2015). Overall, these findings suggest that consideration should be given to methodological differences when using C turnover data to inform and parameterize soil C models.

309

310 **Acknowledgements**

311 We would like to thank Dr. Kevin Wilcox and two anonymous reviewers for their insightful
312 comments and suggestions on this article. This research is financially supported by the United
313 States Department of Energy, Terrestrial Ecosystem Sciences grant DESC0008270, and
314 Biological Systems Research on the Role of Microbial Communities in Carbon Cycling Program
315 grants DE-SC0004601 and DE-SC0010715.

316

317 **References**

- 318 Baisden, W.T., Amundson, R., Cook, A.C., Brenner, D.L., 2002. Turnover and storage of C and
319 N in five density fractions from California annual grassland surface soils. *Global*
320 *Biogeochemical Cycles* 16, 64-61-64-16.
- 321 Baldock, J.A., Skjemstad, J.O., 2000. Role of the soil matrix and minerals in protecting natural
322 organic materials against biological attack. *Organic Geochemistry* 31, 697-710.
- 323 Balesdent, J., Mariotti, A., Boisgontier, D., 1990. Effect of tillage on soil organic carbon
324 mineralization estimated from ^{13}C abundance in maize fields. *Journal of Soil Science* 41,
325 587-596.
- 326 Balesdent, J., Mariotti, A., Guillet, B., 1987. Natural C-13 abundance as a tracer for studies of
327 soil organic matter dynamics. *Soil Biology & Biochemistry* 19, 25-30.
- 328 Benbi, D.K., Boparai, A.K., Brar, K., 2014. Decomposition of particulate organic matter is more
329 sensitive to temperature than the mineral associated organic matter. *Soil Biology &*
330 *Biochemistry* 70, 183-192.

331 Beniston, J.W., Dupont, S.T., Glover, J.D., Lal, R., Dungait, J.J., 2014. Soil organic carbon
 332 dynamics 75 years after land-use change in perennial grassland and annual wheat
 333 agricultural systems. *Biogeochemistry* 120, 37-49.

334 Bird, M., Santruckova, H., Lloyd, J., Lawson, E., 2002. The isotopic composition of soil organic
 335 carbon on a north-south transect in western Canada. *European Journal of Soil Science* 53,
 336 393-403.

337 Bruun, S., Six, J., Jensen, L.S., Paustian, K., 2005. Estimating turnover of soil organic carbon
 338 fractions based on radiocarbon measurements. *Radiocarbon* 47, 99-113.

339 Budge, K., Leifeld, J., Hiltbrunner, E., Fuhrer, J., 2011. Alpine grassland soils contain large
 340 proportion of labile carbon but indicate long turnover times. *Biogeosciences* 8, 1911-
 341 1923.

342 Carvalhais, N., Forkel, M., Khomik, M., Bellarby, J., Jung, M., Migliavacca, M., u, M., Saatchi,
 343 S., Santoro, M., Thurner, M., Weber, U., Ahrens, B., Beer, C., Cescatti, A., Randerson,
 344 J.T., Reichstein, M., 2014. Global covariation of carbon turnover times with climate in
 345 terrestrial ecosystems. *Nature* 514, 213-217.

346 Christensen, B.T., 1996. Matching measurable soil organic matter fractions with conceptual
 347 pools in simulation models of carbon turnover: revision of model structure. In: evaluation
 348 of soil organic matter models. (eds Powlson D, Smith P, Smith J). Springer Berlin
 349 Heidelberg.

350 Christensen, B.T., 1987. Decomposability of organic matter in particle size fractions from field
 351 soils with straw incorporation. *Soil Biology & Biochemistry* 19, 429-435.

352 Christensen, B.T., 2001. Physical fractionation of soil and structural and functional complexity
 353 in organic matter turnover. *European Journal of Soil Science* 52, 345-353.

354 Dalal, R.C., Thornton, C.M., Cowie, B.A., 2013. Turnover of organic carbon and nitrogen in soil
 355 assessed from $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ changes under pasture and cropping practices and estimates
 356 of greenhouse gas emissions. *Science of the Total Environment* 465, 26-35.

357 Derrien, D., Amelung, W., 2011. Computing the mean residence time of soil carbon fractions
 358 using stable isotopes: impacts of the model framework. *European Journal of Soil Science*
 359 62, 237-252.

360 Feller, C., Beare, M.H., 1997. Physical control of soil organic matter dynamics in the tropics.
 361 *Geoderma* 79, 69-116.

362 Fierer, N., Schimel, J.P., 2002. Effects of drying-rewetting frequency on soil carbon and nitrogen
 363 transformations. *Soil Biology & Biochemistry* 34, 777-787.

364 Hastings, W.K., 1970. Monte-Carlo sampling methods using Markov chains and their
 365 applications. *Biometrika* 57, 97-107.

366 Jenkinson, D.S., Rayner, J.H., 1977. Turnover of soil organic matter in some of Rothamsted
 367 classical experiments. *Soil Science* 123, 298-305.

368 Konietzschke, F., Placzek, M., Schaarschmidt, F., Hothorn, L.A., 2015. Nparcomp: An R software
 369 package for nonparametric multiple comparisons and simultaneous confidence intervals.
 370 *Journal of Statistical Software* 64, 1-17

371 Koven, C.D., Chambers, J.Q., Georgiou, K., Knox, R., Negron-Juarez, R., Riley, W.J., Arora,
 372 V.K., Brovkin, V., Friedlingstein, P., Jones, C.D., 2015. Controls on terrestrial carbon
 373 feedbacks by productivity versus turnover in the CMIP5 Earth System Models.
 374 *Biogeosciences* 12, 5211-5228.

375 Lehmann, J., Kleber, M., 2015. The contentious nature of soil organic matter. *Nature* 528, 60-68.

376 Leifeld, J., 2008. Biased ^{14}C -derived organic carbon turnover estimates following black carbon
 377 input to soil: an exploration with RothC. *Biogeochemistry* 88, 205-211.

378 Leifeld, J., Heiling, M., Hajdas, I., 2015. Age and thermal stability of particulate organic matter
 379 fractions indicate the presence of black carbon in soil. *Radiocarbon* 57, 99-107.

380 Liang, A., Chen, S., Zhang, X., Chen, X., 2014. Short-term effects of tillage practices on soil
 381 organic carbon turnover assessed by $\delta^{13}\text{C}$ abundance in particle-size fractions of black
 382 soils from northeast China. *The Scientific World Journal*, DOI: 10.1155/2014/514183.

383 Liao, J.D., Boutton, T.W., Jastrow, J.D., 2006. Organic matter turnover in soil physical fractions
 384 following woody plant invasion of grassland: Evidence from natural ^{13}C and ^{15}N . *Soil*
 385 *Biology & Biochemistry* 38, 3197-3210.

386 Luo, Y., Ahlström, A., Allison, S.D., Batjes, N.H., Brovkin, V., Carvalhais, N., Chappell, A.,
 387 Ciais, P., Davidson, E.A., Finzi, A., Georgiou, K., Guenet, B., Hararuk, O., Harden, J.W.,
 388 He, Y., Hopkins, F., Jiang, L., Koven, C., Jackson, R.B., Jones, C.D., Lara, M.J., Liang,
 389 J., McGuire, A.D., Parton, W., Peng, C., Randerson, J.T., Salazar, A., Sierra, C.A., Smith,
 390 M.J., Tian, H., Todd-Brown, K.E.O., Torn, M., van Groenigen, K.J., Wang, Y.P., West,
 391 T.O., Wei, Y., Wieder, W.R., Xia, J., Xu, X., Xu, X., Zhou, T., 2016. Toward more
 392 realistic projections of soil carbon dynamics by Earth system models. *Global*
 393 *Biogeochemical Cycles* 30, 40–56.

394 Mathieu, J.A., Hatté, C., Balesdent, J., Parent, É., 2015. Deep soil carbon dynamics are driven
 395 more by soil type than by climate: a worldwide meta-analysis of radiocarbon profiles.
 396 *Global Change Biology*, 4278–4292.

397 Metropolis, N., Rosenbluth, A.W., Rosenbluth, M.N., Teller, A.H., Teller, E., 1953. Equation of
398 state calculations by fast computing machines. *Journal of Chemical Physics* 21, 1087-
399 1092.

400 Mikutta, R., Kleber, M., Torn, M.S., Jahn, R., 2006. Stabilization of soil organic matter:
401 Association with minerals or chemical recalcitrance? *Biogeochemistry* 77, 25-56.

402 Miller, A.E., Schimel, J.P., Meixner, T., Sickman, J.O., Melack, J.M., 2005. Episodic rewetting
403 enhances carbon and nitrogen release from chaparral soils. *Soil Biology and*
404 *Biochemistry* 37, 2195-2204.

405 O'Brien, B.J., Stout, J.D., 1978. Movement and turnover of soil organic matter as indicated by
406 carbon isotope measurements. *Soil Biology & Biochemistry* 10, 309-317.

407 Parfitt, R.L., Salt, G.J., 2001. Carbon and nitrogen mineralisation in sand, silt, and clay fractions
408 of soils under maize and pasture. *Soil Research* 39, 361-371.

409 Parton, W.J., Schimel, D.S., Cole, C.V., Ojima, D.S., 1987. Analysis of factors controlling soil
410 organic matter levels in Great Plains grasslands. *Soil Science Society of America Journal*
411 51, 1173-1179.

412 Rabbi, S.M.F., Hua, Q., Daniel, H., Lockwood, P.V., Wilson, B.R., Young, I.M., 2013. Mean
413 residence time of soil organic carbon in aggregates under contrasting land uses based on
414 radiocarbon measurements. *Radiocarbon* 55, 127-139.

415 Rabbi, S.M.F., Wilson, B.R., Lockwood, P.V., Daniel, H., Young, I.M., 2014. Soil organic
416 carbon mineralization rates in aggregates under contrasting land uses. *Geoderma* 216, 10-
417 18.

418 Rumpel, C., Kögel-Knabner, I., 2011. Deep soil organic matter-a key but poorly understood
419 component of terrestrial C cycle. *Plant and Soil* 338, 143-158.

420 Rumpel, C., Kögel-Knabner, I., Bruhn, F., 2002. Vertical distribution, age, and chemical
 421 composition of organic carbon in two forest soils of different pedogenesis. *Organic*
 422 *Geochemistry* 33, 1131-1142.

423 Schädel, C., Luo, Y., Evans, R.D., Fei, S., Schaeffer, S.M., 2013. Separating soil CO₂ efflux into
 424 C-pool-specific decay rates via inverse analysis of soil incubation data. *Oecologia* 171,
 425 721-732.

426 Schmidt, M.W.I., Torn, M.S., Abiven, S., Dittmar, T., Guggenberger, G., Janssens, I.A., Kleber,
 427 M., Kögel-Knabner, I., Lehmann, J., Manning, D.A.C., Nannipieri, P., Rasse, D.P.,
 428 Weiner, S., Trumbore, S.E., 2011. Persistence of soil organic matter as an ecosystem
 429 property. *Nature* 478, 49-56.

430 Schöning, I., Kögel-Knabner, I., 2006. Chemical composition of young and old carbon pools
 431 throughout Cambisol and Luvisol profiles under forests. *Soil Biology & Biochemistry* 38,
 432 2411-2424.

433 Six, J., Elliott, E.T., Paustian, K., Doran, J.W., 1998. Aggregation and soil organic matter
 434 accumulation in cultivated and native grassland soils. *Soil Science Society of America*
 435 *Journal* 62, 1367-1377.

436 Six, J., Feller, C., Denef, K., Ogle, S.M., Sa, J.C.D., Albrecht, A., 2002. Soil organic matter,
 437 biota and aggregation in temperate and tropical soils - Effects of no-tillage. *Agronomie*
 438 22, 755-775.

439 Skjemstad, J.O., Lefeuvre, R.P., Prebble, R.E., 1990. Turnover of soil organic matter under
 440 pasture as determined by C-13 natural abundance. *Australian Journal of Soil Research* 28,
 441 267-276.

442 Sollins, P., Homann, P., Caldwell, B.A., 1996. Stabilization and destabilization of soil organic
 443 matter: Mechanisms and controls. *Geoderma* 74, 65-105.

444 Stevenson, F.J., 1994. Humus chemistry: genesis, composition, reactions, New York, John Wiley
 445 & Sons.

446 Sulman, B.N., Phillips, R.P., Oishi, A.C., Shevliakova, E., Pacala, S.W., 2014. Microbe-driven
 447 turnover offsets mineral-mediated storage of soil carbon under elevated CO₂. *Nature*
 448 *Climate Change* 4, 1099-1102.

449 Talma, A.S., Vogel, J.C., 1993. A simplified approach to calibrating ¹⁴C dates. *Radiocarbon* 35,
 450 317-322.

451 Tan, W., Zhou, L., Liu, K., 2013. Soil aggregate fraction-based C-14 analysis and its application
 452 in the study of soil organic carbon turnover under forests of different ages. *Chinese*
 453 *Science Bulletin* 58, 1936-1947.

454 Tang, J., Riley, W.J., 2015. Weaker soil carbon-climate feedbacks resulting from microbial and
 455 abiotic interactions. *Nature Climate Change* 5, 56-60.

456 Todd-Brown, K.E.O., Randerson, J.T., Post, W.M., Hoffman, F.M., Tarnocai, C., Schuur, E.A.G.,
 457 Allison, S.D., 2013. Causes of variation in soil carbon simulations from CMIP5 Earth
 458 system models and comparison with observations. *Biogeosciences* 10, 1717-1736.

459 Trumbore, S.E., 2000. Age of soil organic matter and soil respiration: radiocarbon constraints on
 460 belowground C dynamics. *Ecological Applications* 10, 399-411.

461 Trumbore, S.E., 1993. Comparison of carbon dynamics in tropical and temperate soils using
 462 radiocarbon measurements. *Global Biogeochemical Cycles* 7, 275-290.

463 von Lützow, M., Kögel-Knabner, I., Ekschmitt, K., Flessa, H., Guggenberger, G., Matzner, E.,
 464 Marschner, B., 2007. SOM fractionation methods: Relevance to functional pools and to
 465 stabilization mechanisms. *Soil Biology & Biochemistry* 39, 2183-2207.

466 Weng, E., Luo, Y., 2011. Relative information contributions of model vs. data to short- and long-
 467 term forecasts of forest carbon dynamics. *Ecological Applications* 21, 1490-1505.

468 Wieder, W.R., Grandy, A.S., Kallenbach, C.M., Bonan, G.B., 2014. Integrating microbial
 469 physiology and physio-chemical principles in soils with the MICrobial-MIneral Carbon
 470 Stabilization (MIMICS) model. *Biogeosciences* 11, 3899-3917.

471 Xu, T., White, L., Hui, D., Luo, Y., 2006. Probabilistic inversion of a terrestrial ecosystem model:
 472 Analysis of uncertainty in parameter estimation and model prediction. *Global*
 473 *Biogeochemical Cycles* 20, GB2007.

474 Xu, X., Shi, Z., Li, D., Rey, A., Ruan, H., Craine, J.M., Liang, J., Zhou, J., Luo, Y., 2016. Soil
 475 properties control decomposition of soil organic carbon: Results from data-assimilation
 476 analysis. *Geoderma* 262, 235-242.

477 Yonekura, Y., Ohta, S., Kiyono, Y., Aksa, D., Morisada, K., Tanaka, N., Tayasu, I., 2013. Soil
 478 organic matter dynamics in density and particle-size fractions following destruction of
 479 tropical rainforest and the subsequent establishment of Imperata grassland in Indonesian
 480 Borneo using stable carbon isotopes. *Plant and Soil* 372, 683-699.

481 Zhang, K., Dang, H., Zhang, Q., Cheng, X., 2015. Soil carbon dynamics following land-use
 482 change varied with temperature and precipitation gradients: evidence from stable isotopes.
 483 *Global Change Biology* 21, 2762-2772

Table 1

Study sites and data distribution of C turnover times of soil fractions estimated by the laboratory incubation, $\delta^{13}\text{C}$ after successive $\text{C}_3\text{-C}_4$ vegetation change, and the ^{14}C dating methods. COM: coarse organic matter, 250-2000 μm ; FOM: fine organic matter, 20/53/63-250 μm ; MOM, mineral associated organic matter, <20/53/63 μm ; sand: 20/53/63-2000 μm ; silt: 2-20/53/63 μm ; clay: <2 μm .

Methods	Fraction	Location (latitude, longitude)	sample size	C turnover times			
				mean	Min	Max	Coefficient variation
Incubation	COM	-30.47-64.87°, -147.72-151.65°	28	8.6	0.5	51	140%
	FOM	-30.47-54.20°, -114.13-151.65°	31	31.5	3.6	342	227%
	MOM	-30.47-64.87°, -147.72-151.65°	37	30.9	3.8	662	347%
	Sand	-40.38-56.08°, -82.73-175.6°	16	6.8	1.6	38	130%
	Silt	-40.38-56.08°, -82.73-175.6°	16	17.9	4.7	123	159%
	Clay	-40.38-56.08°, -82.73-175.6°	16	24.0	6.2	122	118%
^{13}C	COM	-28.67-51.87°, -98.20-153.33°	50	54.1	3.0	278	108%
	FOM	-24.81-51.87°, -98.20-149.8°	38	83.9	6.0	429	133%
	MOM	-24.81-48.36°, 4.40-33.10°	36	210.0	29.0	3124	260%
	Sand	-22.72-48.35°, -88.31-13.18°	15	35.4	9.0	80	62%
	Silt	-10.50-51.87°, -75.35-117.93°	28	116.8	8.0	677	117%
	Clay	-22.72-51.87°, -75.35-117.93°	28	125.4	24.0	357	67%
^{14}C	COM	-30.44-68.10°, -115.30-152.69°	30	286.8	23.0	1265	114%
	FOM	-34.32-68.10°, -115.30-149.98°	28	646.3	42.0	2585	116%
	MOM	-34.32-64.47°, -75.35-149.98°	40	1179.7	26.0	6905	133%
	Sand						
	Silt	34.18-68.10°, -117.77-117.65°	37	415.0	18.0	1660	92%
	Clay	-29.37-68.10°, -117.77-117.65°	55	708.7	8.0	4745	131%

Figure captions

Fig. 1. Geographic locations of soil sampling sites to determine C turnover time of soil fractions.

Triangles represent sampling sites for the laboratory incubations, circles for the ^{13}C method, and diamonds for the ^{14}C method.

Fig. 2. Carbon turnover time of soil fractions estimated by the laboratory incubation and the ^{13}C and ^{14}C methods. COM: coarse organic matter, 250-2000 μm ; FOM: fine organic matter, 20/53/63-250 μm ; MOM, mineral associated organic matter, <20/53/63 μm ; sand: 20/53/63-2000 μm ; silt: 2-20/53/63 μm ; clay: <2 μm . Data are mean $\pm$ SE. Different uppercase letters indicate that C turnover times estimated by the same method significantly differ among soil fractions, and different lowercase letters mean significantly different C turnover times among methods.

Fig. 3. Carbon turnover time of soil fractions estimated by using the two-point and multi-point calculations based on $\delta^{13}\text{C}$ shift after $\text{C}_3\text{-C}_4$ vegetation change. COM: coarse organic matter, 250-2000 μm ; FOM: fine organic matter, 20/53/63-250 μm ; MOM, mineral associated organic matter, <20/53/63 μm ; sand: 20/53/63-2000 μm ; silt: 2-20/53/63 μm ; clay: <2 μm . Data are mean $\pm$ SE.

Fig. 4. Carbon turnover time of soil fractions estimated by using the conventional ^{14}C model and the bomb ^{14}C model. COM: coarse organic matter, 250-2000 μm ; FOM: fine organic matter, 20/53/63-250 μm ; MOM, mineral associated organic matter, <20/53/63 μm ; sand: 20/53/63-2000 μm ; silt: 2-20/53/63 μm ; clay: <2 μm . Data are mean $\pm$ SE. * means values are statistically different between estimate methods.

Figure1

Figure2

Figure3

Figure4