Continental deformation accommodated by non-rigid passive bookshelf faulting: An example from the Cenozoic tectonic development of northern Tibet

Andrew V. Zuza^{1*} (azuza@ucla.edu; avz5818@gmail.com) and An Yin^{1,2}

¹*Department of Earth, Planetary, and Space Sciences, University of California, Los Angeles, CA 90095-1567, USA*

²*Structural Geology Group, China University of Geosciences (Beijing), Beijing 10085, China*

** Corresponding author*

Submitted to: **Tectonophysics**

Submission Date: **June 8th, 2015**

Submission of revised manuscript: **December 7th, 2015**

Submission of second revision: **March 21st, 2016**

ABSTRACT

Collision-induced continental deformation commonly involves complex interactions between strike-slip faulting and off-fault deformation, yet this relationship has rarely been quantified. In northern Tibet, Cenozoic deformation is expressed by the development of the >1000-km-long east-striking left-slip Kunlun, Qinling, and Haiyuan faults. Each have a maximum slip in the central fault segment exceeding 10s to ~100 km but a much smaller slip magnitude (~< 10% of the maximum slip) at their terminations. The along-strike variation of fault offsets and pervasive off-fault deformation create a strain pattern that departs from the expectations of the classic plate-like rigid-body motion and flow-like distributed deformation approximations for continental deformation. Here we propose a non-rigid bookshelf-fault model for the Cenozoic development of northern Tibet. Our model, relating discrete left-slip faulting to distributed off-fault deformation during regional clockwise rotation, explains several puzzling features, including the: (1) clockwise rotation of east-striking left-slip faults against the northeast-striking left-slip Altyn Tagh fault along the northwestern margin of the Tibetan Plateau, (2) alternating fault-parallel extension and shortening in the off-fault regions, and (3) eastward-tapering Qimen Tagh, Qaidam, and Qilian Shan thrust belts that link with the three left-slip faults. We refer to this non-rigid bookshelf fault system as passive bookshelf-faulting because the rotating bookshelf panels are detached from the rigid bounding domains. As a consequence, the wallrock of the strike-slip faults deforms to accommodate both the clockwise rotation of the left-slip faults and the space issues that arise at the bookshelf-panel ends. The model suggests that thrust and strike-slip faulting are coeval and kinematically linked. Another implication of our model is that the style and magnitude of Cenozoic deformation in northern Tibet vary considerably in the east-west direction. Thus, any single north-south cross section and

its kinematic reconstruction through the region does not properly quantify the complex deformational processes of plateau construction.

Key Words: Bookshelf faulting; Tibetan Plateau; Strike-slip faulting; Kunlun fault; Qinling fault; Haiyuan fault

1. Introduction

The fundamental mode of continental tectonics has been characterized by two end-member processes: plate-like rigid-body motion (e.g., Tapponnier et al., 1982; Weldon and Humphreys, 1986; Avouac and Tapponnier et al., 1993; Meade, 2007) and distributed deformation via viscous flow (e.g., England and Houseman, 1986; Yin and Taylor, 2011). In the rigid-plate model, continental deformation is quantified by rigid block rotation on a sphere about their respective Euler poles; the horizontal dimension of the blocks is much greater than the width of faults/shear zones that bound the blocks (e.g., Avouac and Tapponnier, 1993). In contrast, the viscous-flow model quantifies continental deformation by solving a boundary-value problem that requires the knowledge of lithospheric rheology (e.g., England and Houseman, 1986). This model envisions distributed continental deformation, with major faults approximated as zones of high strain within a continuum. These two end-member models have been extensively tested in Tibet against structures created during the Cenozoic India-Eurasia collision (e.g., Yin and Harrison, 2000; Zhu et al., 2005; Yin, 2010a; van Hinsbergen et al., 2011; Yin and Taylor, 2011). Debates have been centered on whether the >1000-km-long east-striking left-slip Haiyuan, Qinling, and Kunlun faults in northern Tibet (**Fig. 1**) have acted as rigid-block boundaries (Tapponnier et al., 1982; Avouac and Tapponnier, 1993; Tapponnier et al., 2001) or

transfer-fault structures linking dip-slip fault systems (e.g., Burchfiel et al., 1991; Yin, 2000; Duvall and Clark, 2010).

One form of rigid-block models for deformation in northern Tibet is bookshelf faulting, which requires that the observed left-slip faulting is driven by regional right-lateral shear (Cobbold and Davy, 1988; England and Molnar, 1990). Applying the classic rigid bookshelf-fault model (Freund 1970) to explain the tectonic development of northern Tibet raises several important questions that have not yet been addressed: (1) Why do the parallel east-striking left-slip faults terminate at the northeast-striking left-slip Altyn Tagh fault rather than a right-slip northeast-striking shear zone as required by the bookshelf-fault model (**Fig. 1**)? (2) How are the required lithospheric-scale “gaps” and “overlaps” at the ends of the rotating blocks accommodated by off-strike-slip-fault deformation (e.g., Luyendyk et al., 1980; Onderdonk, 2005; Platt and Becker, 2013) (**Fig. 2**)? (3) What is the kinematic relationship between the east-striking left-slip faults and the triangular eastward-tapering thrust belts at the western ends of the strike-slip faults (i.e., the Qimen Tagh, Qaidam, and Qilian Shan-Nan Shan thrust belts) (**Fig. 1**)?

In this study we propose a non-rigid bookshelf-fault model (e.g., Yin and Pappalardo, 2015) to resolve the above issues. Specifically, we show that an eastward decrease in Cenozoic strain results in clockwise rotation and left-slip bookshelf faulting across northern Tibet. The detached rotation of these bookshelf faults against rigid bounding domains, which we refer to as passive bookshelf faulting, may explain why a left-slip bookshelf fault system is bounded by the left-slip Altyn Tagh fault (**Fig. 1**). Non-rigid wallrock deformation within the strike-slip-fault-bounded regions accommodates both the clockwise rotation of the strike-slip faults and the space issues that arise at the ends of the bookshelf panels. The model implies that thrust belt development and strike-slip faulting in the region are coeval and kinematically linked, which

contrasts to an earlier suggestion that they represent two distinct stages of plateau development (e.g., Yuan et al., 2013).

2. Cenozoic Left-slip Faults in Northern Tibet

The ~N110°E-striking left-slip Kunlun, Qinling, and Haiyuan faults, extending for ~1500, ~1000, and ~1000 km respectively, are by far the longest and most continuous structures in northern Tibet (**Fig. 1**) (Tapponnier et al., 2001; Taylor and Yin, 2009). The faults are lithospheric structures (Wang et al., 2011; Gao et al., 2013) that closely follow the surface trace of the Qilian, Qinling, and Kunlun suture zones (Yin and Harrison, 2000; Wu et al., 2016) (**Fig. 1**).

The kinematics of these major east-striking faults has been related to lateral extrusion (Tapponnier et al., 1982; 2001; Cheng et al., 2015), strain transfer between thrust belts (Burchfiel et al., 1991; Zhang et al., 1991; Duvall and Clark, 2010), and bookshelf faulting associated with clockwise fault rotation induced by broad north-trending right-lateral shear (England and Molnar, 1990; Zuza and Yin, 2013) (**Fig. 3a**). The extrusion fault model requires high slip rates (> 10-20 mm/yr), large fault offsets (100s km) (Tapponnier et al., 1982; Avouac and Tapponnier, 1993), the presence of zipper thrusts at the western end of the left-slip faults (Peltzer and Tapponnier, 1988; Cheng et al., 2015), and a conjugate right-slip fault with a similar slip magnitude in the west to assist eastward lateral extrusion (scenario 1 in **Fig. 3a**). In contrast, the transfer-fault model predicts transpressional deformation at the termination thrusts oriented obliquely to the strike-slip faults (scenario 2 in **Fig. 3a**). Finally, the bookshelf fault model predicts clockwise rotation of the left-slip faults and the fault-bounded wallrock. If the bookshelf panels are rigid, the model predicts the formation of “gaps” and/or “overlaps” at the end of the

rotating blocks (scenario 3 in **Fig. 3a**); these gaps and overlaps can be reconciled by fault-parallel extension and/or shortening with non-rigid off-fault deformation (scenario 4 in **Fig. 3a**). Proposed solutions to the space issues of rigid bookshelf faulting (scenario 3 in **Fig. 3a**) include (1) extension leading to basin formation (Luyendyk et al., 1980) (**Fig. 2b**), (2) shortening within the terminal regions of the rotated blocks (e.g., Onderdonk, 2005) (**Fig. 2c**), or (3) fault-parallel stretching across the rotated blocks (e.g., Platt and Becker, 2010, 2013) (**Fig. 2d**).

2.1 Initiation age and along-strike variation of fault offsets and slip rates

A summary of inferred fault initiation ages across northern Tibet is shown in Figure 3b (Yuan et al., 2013). The Haiyuan fault, which initiated at ~17-12 Ma in the west (Duvall et al., 2013) and ~8 Ma in the east (Zheng et al., 2006; Yuan et al., 2013) (**Fig. 3b**), is estimated to have 95 ± 15 km left-lateral offset at the center (Gaudemer et al., 1995) and 10-15 km slip at the eastern end (Burchfiel et al., 1991; Zhang et al., 1991) (**Fig. 1**). Slip along the range-bounding thrusts at the western termination of the Haiyuan fault is ≤ 10 km (Reith, 2013; Zuza et al., 2014, 2016) (**Fig. 1**), requiring a significant decrease in fault slip from the center to the western end. The bidirectional decrease in the total fault offset along the Haiyuan fault correlates with the along-fault variation of Quaternary and global positional system (GPS) velocity field based slip rates: 11-19 mm/yr at the center (Gaudemer et al., 1995; Lasserre et al., 1999, 2002), < 5 mm/yr in the east (Zhang et al., 1991; Cavalié et al., 2008; Li et al., 2009), and 2-4 mm/yr in the west (Duvall and Clark, 2010).

The western segment of the Qinling fault is a transpressional structure that initiated at ~50 Ma (Duvall et al., 2013; Yuan et al., 2013) (**Fig. 3b**), whereas the eastern segment of the Qinling fault is a transtensional structure initiated in the Eocene with accelerated motion in the

late Miocene (Mercier et al., 2013) (**Fig. 1**). Slip along the central and eastern sections of the Qinling fault is 20-30 km (Ratschbacher et al., 2003) with a Quaternary slip rate of 7.2 ± 2.2 mm/yr (Zhang et al., 1995). The total fault offset in the west is constrained by east-west shortening magnitude estimates of 10-15 km across both the Jishi Shan and Liupan Shan thrust belts (**Fig. 1**) (Zhang et al. 1991; Burchfiel et al. 1991; Lease et al., 2012). The Jishi Shan and Liupan Shan thrust belts initiated at ~13 Ma and ~8 Ma respectively (Zheng et al., 2006; Godard et al., 2009; Lease et al., 2012), corresponding to an average slip rate on the western Qinling fault of about 1-1.5 mm/yr. This is significantly less than that of the central and eastern fault segments.

The Kunlun fault initiated diachronously at 35-30 Ma in the west (Mock et al., 1999; Jolivet et al., 2003; Clark et al., 2010), 20-15 Ma along its central segment (Yuan et al., 2006; Duvall et al., 2013), and ~8 Ma in the east (Duvall et al., 2013) (**Fig. 3b**). Offset along the fault decreases bidirectionally from ~120 km in the center (Kidd and Molnar, 1988) to <10s km at the eastern and western ends when it links with minor dip-slip fault systems (Jolivet et al., 2003; Fu and Awata, 2007) (**Fig. 1**). Quaternary slip rates along the central Kunlun fault of 10-12 mm/yr (Van der Woerd et al., 1998, 2000, 2002; Fu et al., 2005; Li et al., 2005) decrease to < 2 mm/yr along the eastern segment (Kirby et al., 2007; Harkins and Kirby, 2008; Harkins et al., 2010). Given the fault initiation age of 35-30 Ma in the west and 20-30 km of slip along the fault (Mock et al., 1999; Jolivet et al., 2003), the average slip rate of the Kunlun fault in the west is also < 2 mm/yr. Thus, slip rates along the Kunlun fault must decrease bidirectionally from the center.

These east-striking left-slip faults terminate in the west at the northwest-trending Qilian Shan-Nan Shan, Qaidam Basin, and Qimen Tagh thrust belts and in the east at the north-trending Liupan Shan and Longmen Shan thrust belts (**Fig. 1**). Thrusting first started at ~50 Ma in the

southern Qilian Shan and northern Qaidam Basin thrust systems, which was followed by activity in the Qimen Tagh and the northern Qilian Shan thrust belts at 25-20 Ma (e.g., Yin et al., 2008a; Duvall et al., 2013; Cheng et al., 2014) (**Fig. 3b**). The Longmen Shan thrust belt, which involves dip-slip thrusting and minor right-slip faulting (Densmore et al., 2007; Yin, 2010b), began to develop at 30-25 Ma and accelerated deformation occurred at 15-10 Ma (Godard et al., 2009; Wang et al., 2012). The Liupan Shan thrust belt initiated at ~8 Ma (Zheng et al. 2006). Existing studies indicate a north-south shortening magnitude of 200-450 km across the three northwest-trending thrust belts (i.e., the Qimen Tagh, Qaidam Basin, and Qilian Shan-Nan Shan) (Gaudemer et al., 1995; Meyer et al., 1998; Yin et al., 2007; Yin et al., 2008a, 2008b; Gao et al., 2013; Cheng et al., 2014; Zuza et al., 2016), which contrasts sharply to the 10-40 km of shortening across the north-trending Liupan Shan and Longmen Shan thrust belts (**Fig. 1**) (Zhang et al., 1991; Burchfiel et al., 1995; Hubbard and Shaw, 2009).

The major left-slip faults in northern Tibet also link with extensional zones and right-slip faults (**Fig. 1**). The Haiyuan fault merges with the north-trending Yinchuan rift system in the east and southeast-striking right-slip Elashan and Riyueshan faults in the west (e.g., Zhang et al., 1998), the Qinling fault links with the Shanxi rift system in the east and right-slip Riyueshan-Gonghe fault in the west (e.g., Zhang et al., 1998), and the Kunlun fault links with central Tibetan rifts in the west (Yin, 2000) and the right-slip Elashan and Gonghe faults along its eastern segment (**Fig. 1**). Note that both the obliquely oriented western thrust belts and right-slip faults and the orthogonally oriented rifts and eastern thrust belts accommodate stretching and shortening parallel to the east-striking left-slip faults (**Fig. 1**). This off-fault strain pattern is inconsistent with the extrusion and transfer-fault models for the tectonic development of left-slip faulting in northern Tibet (scenarios 1 and 2 in **Fig. 3a**).

2.2 Protracted regional clockwise rotation of northern Tibet

The restoration of a series of balanced cross sections indicates an eastward decrease in north-south Cenozoic shortening across the Qaidam Basin thrust belt (Yin et al., 2008a,b). A similar eastward decrease in Cenozoic shortening strain may be inferred across the Qilian Shan and Qimen Tagh thrust belts on the basis of an eastward decrease in the (1) number of thrusts, (2) thrust belt width, and (3) average elevation assuming all thrusts initiated at the same elevation and have experienced the same amount of erosion (**Fig. 1**). The eastward decrease in shortening strain implies clockwise rotation across northern Tibet throughout much of the Cenozoic (i.e., since thrust initiation at 50–45 Ma). This observation is consistent with paleomagnetic data that suggests 15°–20° total clockwise rotation across northern Tibet with respect to the Eurasian reference pole since the Cretaceous (**Fig. 4b**) at a rate of 0.3 to 0.5°/Ma (Frost et al., 1995; Halim et al., 1998, 2003; Cogné et al., 1999; Dupont-Nivet et al., 2004; Chen et al., 2002a, 2002b; Sun et al., 2006). Variability in rotation magnitudes, ranging from 0° to >20° across the region, results from heterogeneous Cenozoic deformation (Yin et al., 2008a).

Geodetic measurements (Gan et al., 2007) show clockwise rotation of northern Tibet about an Euler pole located ~500 km to the southeast of the Eastern Himalayan Syntaxis (**Fig. 4**). Other workers have placed the rotation poles for northern Tibet in the western South China Sea (Peltzer and Saucier, 1996) and southwest China (Thatcher, 2007) to fit the global positional system (GPS) data with clockwise rotation rates of 0.5 to 1°/Ma.

We examine the velocity field across the region using the published GPS data of Gan et al. (2007) to constrain the rotation rates of the east-striking left-slip faults in northern Tibet. With this data, we derive the relative rotation and shear strain rates for four fault-bounded domains:

the Qilian Shan (QS), North Qaidam (NQ), Qaidam Basin (QB), and East Central Tibet (ECT). The rotation rates of the above fault-bounded domains may approximate the fault rotation rates if present-day fault-perpendicular shortening and extension are negligible. Using the northern foreland of the Tibetan Plateau in the North China craton as a fixed reference frame (i.e., the star in **Fig. 4b**), we project fault-perpendicular velocities (i.e., N20°E, referred to as the y -direction) of each fault-bounded domain onto a line trending N110°E (referred to as the x -direction) that approximates the strike-slip fault trend (**Fig. 4b**). Note that we only picked stations that are to the north or south of the major left-slip faults (e.g., stations in the North Qaidam thrust belt were excluded) and are located more than ~50 km from the active strike-slip faults to avoid complications due to elastic behavior along locked faults. Each domain displays a relatively linear east-west velocity gradient, shear strain rate ($\dot{\gamma} = 2\dot{\epsilon}_{xy}$), and rotation rate ($\dot{\omega}$) (**Fig. 4c**). Clockwise rotation rates are similar for the East Central Tibet, Qaidam Basin, and North Qaidam domains (i.e., 0.59°/Ma, 0.66°/Ma, and 0.64°/Ma, respectively) and decrease to 0.41°/Ma for the Qilian Shan domain in the north (**Fig. 4c**). The decrease in rotation rate across the Qilian Shan implies that central and northern Tibet are rotating in a clockwise sense relative to North China. The magnitudes of north-south right-lateral shear strain rates ($\dot{\epsilon}_{xy}$) vary between $1.0 \times 10^{-16} \text{ s}^{-1}$ and $2.5 \times 10^{-16} \text{ s}^{-1}$ (**Fig. 4c**). These present-day rotation rates are similar with those that are geologically and paleomagnetically derived (see above), although less than then 1-2°/Ma estimated by England and Molnar (1990).

2.3 Extension and shortening within fault-bounded domains

The wallrock of the major left-slip faults, defined as the crust bounded by the strike-slip faults, displays a distinctive four-quadrant strain distribution (**Fig. 5**). Each strike-slip fault can

be divided into four quadrants that alternate between domains of fault-parallel stretching and domains fault-parallel shortening (**Fig. 5a**). The northern wallrock of the Haiyuan fault experiences fault-parallel shortening through thrusting across the obliquely oriented Qilian Shan-Nan Shan thrust belt in the west and fault-parallel extension across the orthogonally oriented Yinchuan rift in the east (**Figs. 1 and 5b**). The southern wallrock of the Haiyuan fault experiences fault-parallel stretching via right-slip faulting in the west and fault-parallel shortening across the Liupan Shan thrust belt in the east (**Figs. 1 and 5b**). The northern and southern wallrock panels of the Qinling fault are dominated by fault-parallel contraction across the Jishi Shan and Liupan Shan thrust belts in the northwest and a contractional zone bounding the southeastern edges of the Qinling and Daba Shan in the east (**Figs. 1 and 5**). In contrast, the Qinling wallrock is characterized by fault-parallel extension along the Shanxi rift system to the northeast and a right-slip fault system to the southwest (**Fig. 5b**). The Kunlun fault also displays a four-quadrant deformation pattern expressed by shortening across the Qimen Tagh and Min Shan-Longmen Shan thrust belts and extension across north-striking Tibetan rifts and right-slip faulting (Kirby et al., 2000; Jolivet et al., 2003; Wang and Burchfiel, 2004; Yin et al., 2007, 2008b; Duvall and Clark, 2010; Cheng et al., 2014) (**Fig. 5b**).

The southeast-striking right-slip Elashan, Riyueshan and Gonghe faults may represent a secondary bookshelf faults within this larger bookshelf fault system (e.g., Duvall and Clark, 2010) (**Figs. 1 and 5b**). The fault geometries and right-slip kinematics, consistent with left-slip shear between the Kunlun and Haiyuan faults, also suggest north-south shortening and east-west stretching (i.e., strike-slip fault-parallel extension and fault-perpendicular shortening).

3. Non-rigid Bookshelf Fault Model

As summarized above, existing work across northern Tibet indicates: (1) an eastward decrease in Cenozoic north-south shortening accompanied by clockwise rotation of the east-striking left-slip faults and their wallrock since the Eocene, (2) a bilateral decrease in fault offset and slip rate along the left-slip faults that eventually almost disappears at the fault ends, and (3) a distinctive four-quadrant strain pattern in the wallrock of these faults. These observations can be tested against the existing models for the role of strike-slip faulting in accommodating India-Eurasia convergence across northern Tibet (**Fig. 3a**). The observed fault rotation is not explicitly required by the extrusion-fault and transfer-fault models (scenarios 1 and 2 in **Fig. 3a**), and neither of these models predict a four-quadrant strain pattern along the left-slip faults (**Fig. 5**). The observed bidirectional decrease in fault slip is also inconsistent with these models. The extrusion-fault model (scenario 1 in **Fig. 3a**) requires conjugate faults with comparable slip magnitudes, which is not observed in northern or central Tibet (**Fig. 1**) (cf. Cheng et al., 2015). Lastly, dip-slip north-south shortening in the western Qilian Shan-Nan Shan thrust belt near the western end of the Haiyuan fault, as indicated by fault plane solutions (**Fig. 1**) (e.g., Taylor and Yin, 2009) and field observations (Yin et al., 2008a; Reith, 2013; Zuza et al., 2014), is inconsistent with the transfer fault model that predicts left-slip transpression across this termination thrust system (scenario 2 in **Fig. 3a**).

At face value, the observations from northern Tibet may best be explained by a rotated-fault model. The rigid bookshelf-fault model (England and Molnar, 1990) (scenario 3 in **Fig. 3a**), based on the classic work of bookshelf faulting by Freund (1970), does not account for off-fault deformation and fails to quantify the relationship between strike-slip and dip-slip faulting. Below

we propose a new non-rigid bookshelf-fault model for the development of linked left-slip faults and eastward-tapering thrust belts in northern Tibet.

3.1 Active and passive bookshelf faulting

Traditionally, the sense of shear within a bookshelf fault system is opposite that of the bounding shear zone that drives bookshelf panel rotation (Freund, 1970). For example, a left-slip bookshelf fault system requires a right-slip bounding shear zone to drive clockwise panel rotation. In this case, right-lateral shear “drags” the fault-bounded domains in a clockwise fashion. We herein refer to this process as *active* bookshelf faulting, because at least one of the bounding “walls” is moving to drive panel rotation (**Fig. 6a**). Alternatively, we introduce a new self-rotating *passive* bookshelf fault system, where the rotating panels are detached from the fixed boundary “walls” (e.g., Yin and Pappalardo, 2015) (**Fig. 6b**). In this model, the kinematics of the boundary shear zones and the individual bookshelf faults can be the same when the rotation axis is fixed in the center of the bookshelf fault system and the bounding walls are fixed (middle panel of **Fig. 6b**). Qualitatively, if the active bookshelf system is driven by drag along the edges, then the passive system is driven by a push from behind (or upward as in **Fig. 6b**) or rotation from below. Our model predicts that the top bounding surface in **Figure 6b** absorbs shortening strain, and the magnitude of this strain varies laterally due to block rotation. Note that there are many possible variations of our passive bookshelf-fault model, depending on the relative magnitude of rigid-block rotation and rigid-block translation. For example, if the bookshelf fault system experiences both self-rotation at the panel centers and significant translation (upward in **Fig. 6b**), the sense of shear along one of the boundary shear zones may

switch (bottom panel in **Fig. 6b**). Depending on the magnitude of translation, there may be little observable shear on the right boundary zone in **Figure 6b**.

The geometric relationships between the initial width of the system in the x -direction (L_o), width of the panels in the y -direction (W), panel rotation magnitude (α), translation velocity magnitude (v), and boundary shear slip (e.g., S_o , S_l , S_2 , etc.) are quantified, assuming rigid-body motion, in **Figure 6**. For the active bookshelf fault system shown in **Figure 6a**, translation of the left wall by S_o relative to the right wall leads to right lateral shear strain (total shear strain of S_o/L) and clockwise panel rotation (α), so that the variable width of the system (L) is

$$L = W \sin \alpha + L_o \cos \alpha . \quad (1)$$

Half of the corners of the bookshelf panels remain in contact with the bounding-edge of the walls but they may slip in the y -direction (i.e., S_l and S_6 in **Fig. 6a**), so that the system can experience more shear than the panel rotation expresses. Given that all of left-side bookshelf panel corners move relative to the left boundary wall in the y -direction by S_l , and $S_l \geq 0$ (**Fig. 6a**), the slip of the other left-side corners relative to the left boundary wall is

$$S_2 = W / (\cos \alpha) - W + S_1 \quad (2a)$$

$$S_3 = 2W / (\cos \alpha) - 2W + S_1. \quad (2b)$$

Slip of the right-side corners relative to the right boundary wall is similar, except that the magnitudes are related to S_6 instead of S_l (**Fig. 6a**).

The width of the passive bookshelf fault system remains constant ($L = L_o$) (**Fig. 6b**). Rotation of the panels in the passive bookshelf fault system of **Figure 6b** results in boundary zone slip magnitudes (i.e., S_l and S_2 on the left and right sides, respectively) of

$$S_1 = (2 \tan \alpha) / L_o \quad (3a)$$

$$S_2 = -(2 \tan \alpha) / L_o. \quad (3b)$$

Alternatively, if there is a component of translation in the y -direction (v) (bottom panel of **Fig. 6b**), the resulting boundary zone slip magnitudes are described by:

$$S_1 = (2 \tan \alpha)/L_o + v \quad (4a)$$

$$S_2 = -(2 \tan \alpha)/L_o + v. \quad (4b)$$

First-order deformation in either the passive or active systems can be quantified as continuous velocity fields. The idealized velocity field (i.e., no discrete faults) across an active bookshelf system is

$$u = 0 \quad (5a)$$

$$v = -\frac{v_o}{h}x \quad (5b)$$

where u and v are the velocity components in the x and y directions perpendicular and parallel to the shear zone, respectively, h is the width of the shear zone, and v_o is the velocity on one side of the shear zone at $x = -h$ relative to the other side of the shear zone (**Fig. 7a**). Note that in all of our derivations, the origin is placed at the right-side boundary, so that the left boundary is located at $x = -h$ (**Fig. 7**). The corresponding strain-rate-tensor components are

$$\dot{\epsilon}_{xx} = \frac{\partial u}{\partial x} = 0 \quad (6a)$$

$$\dot{\epsilon}_{yy} = \frac{\partial v}{\partial y} = 0 \quad (6b)$$

$$\dot{\epsilon}_{xy} = \frac{1}{2} \left(\frac{\partial u}{\partial y} + \frac{\partial v}{\partial x} \right) = -\frac{v_o}{2h}, \quad (6c)$$

where $\dot{\epsilon}_{xx}$ and $\dot{\epsilon}_{yy}$ are extension strain rates in the x and y directions, perpendicular and parallel to the shear zone, and $\dot{\epsilon}_{xy}$ is the shear strain rate (e.g., Ramsay and Huber, 1983). The rotation rate may be derived as follows:

$$\dot{\epsilon}_{xy} = \frac{\dot{\gamma}}{2} = \frac{\tan \omega}{2} \quad (7a)$$

$$\dot{\omega} = \tan^{-1} \frac{-v_o}{h} \quad (7b)$$

where $\dot{\omega}$ is rotation rate and $\dot{\gamma}$ is engineering or total shear strain rate (**Fig. 7a**).

Deformation for the passive bookshelf system (**Fig. 7b**) can be quantified in a similar manner, except that the velocity in the y -direction is the sum of velocities related to translation and rotation (**Figs. 6b** and **7b**), such that

$$v = v_r + v_t \quad (8a)$$

$$v_r = -v_{ro} \left(\frac{2x}{h} + 1 \right) \quad (8b)$$

$$v_t \geq 0 \quad (8c)$$

where v is the velocity component in the y -direction parallel to the shear zone, h is the width of the shear zone, v_r is velocity component related to rotation, v_{ro} is the rotation-related velocity at one side of the shear zone at $y = -h$, and v_t is the velocity component related to translation. In this approximation, we are only considering the y -component of the rotational velocity (v_r). The complete corresponding velocity field is

$$u = 0 \quad (9a)$$

$$v = -v_{ro} \left(\frac{2x}{h} + 1 \right) + v_t \quad (9b)$$

where u is the velocity component in the x -direction perpendicular to the shear zone, and the other variables are the same as above (**Fig. 7b**). The strain-rate-tensor components are as follows:

$$\dot{\epsilon}_{xx} = \frac{\partial u}{\partial x} = 0 \quad (10a)$$

$$\dot{\epsilon}_{yy} = \frac{\partial v}{\partial y} = 0 \quad (10b)$$

$$\dot{\epsilon}_{xy} = \frac{1}{2} \left(\frac{\partial u}{\partial y} + \frac{\partial v}{\partial x} \right) = \frac{-v_{ro}}{h} \quad (10c)$$

from which the rotation rate is given by

$$\dot{\omega} = \tan^{-1} \frac{-2v_{ro}}{h}. \quad (11)$$

3.2 Non-rigid bookshelf faulting in northern Tibet: passive and stretchy

We propose that the clockwise rotating east-striking left-slip faults in northern Tibet are part of a passive bookshelf system (**Fig. 6b**) because they are bounded to the northwest by the left-slip Altyn Tagh fault (**Fig. 1**). An active left-slip bookshelf fault system (**Fig. 6a**) would require a right-slip bounding shear zone to drive clockwise panel rotation, whereas the passive system allows the panels to rotate in a clockwise fashion against a rigid boundary to the west, creating a left-slip shear zone. This simple model involving clockwise rotation across the region may explain why Tibet is bounded by left-slip fault system to the northwest (i.e., the Altyn Tagh fault) and minor-to-insignificant right-slip faulting to the southeast (i.e., within the Longmen Shan) (**Fig. 1**), rather than the expected broad right-slip shear zones as predicted by the classic rigid-bookshelf model (England and Molnar, 1990).

The predicted velocity field from the passive bookshelf system (Equation 9) can be compared to the GPS-derived velocity field across northern Tibet (**Fig. 4**). The eastern boundary of northern Tibet does not have a significant north-striking strike-slip bounding fault like the western boundary (i.e., the Altyn Tagh fault); only minor right-slip faulting is observed in the Longmen Shan (Densmore et al., 2007; Yin, 2010b) (**Fig. 1**). Because of this, we can infer that the N-S velocity due to clockwise rotation is matched by the northward translation velocity, so that v_{ro} is equal but opposite to v_t at $x = 0$ (the eastern boundary) (**Fig. 7b**) in Equation 9. The N20°E velocity gradient plotted in **Figure 4c** shows a maximum velocity near the western boundary ($x = -h$) of ~16 mm/yr, which requires v_r and v_t of Equation 9 to equal ~8 mm/yr. A northward translational velocity (v_t) of 8 mm/yr is reasonable given that the northward GPS

velocities just south of the Kunlun fault are 7-15 mm/yr (Zhang et al., 2004; Yin and Taylor, 2011). If we assume that the fixed width of the system h is 1400 km (**Fig. 1**), the N-S velocity field can be described by $v = (-2.46 \times 10^{-8})x \text{ yr}^{-1}$ (**Fig. 7b**). The resulting shear strain rate and clockwise rotation rate are $\dot{\epsilon}_{xy} = -1.9 \times 10^{-16} \text{ s}^{-1}$ and $\dot{\omega} = 0.68^\circ/\text{Ma}$ respectively. These rates are comparable to the paleomagnetic, geologic, and geodetic data discussed above.

Since the boundary domains in the active or passive bookshelf system are relatively rigid, the bookshelf panels must be non-rigid to resolve the space issues that arise at the panel ends (**Figs. 2 and 6**) (e.g., Luyenduk et al., 1980; Onderdonk, 2005; Platt and Becker, 2013). The necessary fault-parallel stretching or shortening of the fault-bounded domains accommodates negligible slip at the ends of each rotating fault. The strike-slip faults in a non-rigid bookshelf system, bounded by deformable rather than rigid wallrocks, are referred to as stretching faults (Means et al., 1989, 1990).

Unlike the models of either Onderdonk (2005) or Platt and Becker (2013), the bookshelf panels in our non-rigid passive bookshelf model do not have to either *all stretch* or *all shorten* to accommodate rotation in a deforming material. Instead, each quadrant of wallrock deforms according to the kinematics of the bounding strike-slip faults, so that off-strike-slip-fault wallrock deformation accommodates negligible slip at the strike-slip faults tips (**Fig. 5**). An example of a possible interaction along the left-most boundary is shown in **Figure 8**. Given the small angle of clockwise rotation of the left-slip faults ($\theta \ll 1^\circ$), the resulting left-lateral slip (d) must be accommodated by fault-parallel stretching *and* shortening strain (ϵ_{xx}) in the panels adjacent to the strike-slip fault (i.e., panels A and B in **Fig. 8**). Assuming that the resulting strain is homogenous throughout one-half of the bookshelf panel and affects both panels A and B by the same magnitude, this strain can be written as:

$$\varepsilon^A_{xx} = \frac{1}{2} \frac{d}{S} \quad (12a)$$

$$\varepsilon^B_{xx} = -\frac{1}{2} \frac{d}{S} \quad (12b)$$

381 where ε^A_{xx} and ε^B_{xx} are the extension strain in panel A and panel B, respectively, d is the left-
 382 lateral slip, and S is the half the length of the bookshelf panel ($S = L/2$), all in the x -direction
 383 (**Fig. 8**). Because this left-slip (d) is caused by clockwise rotation, we can relate this strain to
 384 rotation rate, and derive the resulting fault-parallel strain rate:

$$\dot{\varepsilon}^A_{xx} = \frac{1}{2} \frac{W}{S} \frac{\tan(\Delta\theta)}{(\Delta t)} \quad (13a)$$

$$\dot{\varepsilon}^B_{xx} = -\frac{1}{2} \frac{W}{S} \frac{\tan(\Delta\theta)}{(\Delta t)} \quad (13b)$$

385 where W is the width of the bookshelf panel in the y -direction, $\Delta\theta$ is the small angle of clockwise
 386 rotation of the left-slip faults over a short timescale (Δt), and the other variables are the same as
 387 above. The corresponding velocity field may be written as

$$u^A_x = \frac{x}{2} \frac{W}{S} \frac{\tan(\Delta\theta)}{\Delta t} + c \quad (14a)$$

$$u^B_x = -\frac{x}{2} \frac{W}{S} \frac{\tan(\Delta\theta)}{\Delta t} + c \quad (14b)$$

388 where u^A_x and u^B_x are the velocity components in the x -direction in panels A and B respectively,
 389 and c is a constant of integration may be dropped by setting the boundary condition of $u_x = 0$ at x
 390 $= 0$ (**Fig. 8**). Combining Equations 14 and 11 yields an expression for fault-parallel stretching
 391 rate (u_x) as a function of v_{ro} :

$$u^A_x = \frac{x}{2} \frac{W}{S} \tan^{-1} \frac{-2v_{ro}}{h} \quad (15a)$$

$$u^B_x = -\frac{x}{2} \frac{W}{S} \tan^{-1} \frac{-2v_{ro}}{h}. \quad (15b)$$

392 If we take v_{ro} to be 8 mm/yr (as discussed above), the bookshelf panel between the
 393 Kunlun and Qinling faults to have a width of 150 km and length of 1300 km ($S = 650$ km) (**Fig.**

1), and the width of the bookshelf system h as 1300 km, we arrive at fault-parallel stretching rates of ~ 1 mm/yr. For the Kunlun fault, as an example, this should be accommodated by shortening in the Qimen Tagh and extension to the south in central Tibet at a rate of ~ 1 mm/yr (**Fig. 1**).

4. Discussion

4.1 Driving mechanisms for clockwise rotation

Applying the passive bookshelf fault model to northern Tibet requires clockwise rotation of the left-slip faults and fault-bounded wallrock relative to the relatively fixed Tarim and North China (i.e., the top, left, and right sides of the passive bookshelf fault system in **Fig. 6b**). This system can be driven by either rotation of the bottom boundary (i.e., rotation of central and southern Tibet) or basal drag beneath the system. In the simplest case, clockwise rotation of southern Tibet and/or the Himalaya relative to North China (**Fig. 9a**) can drive clockwise rotation of northern Tibet. Due to the lack of paleomagnetic and GPS data across central and southern Tibet, this model is difficult to test with confidence (**Figs. 1** and **4a**). However, existing geologic data from central and southern Tibet suggest that the regions are actually experiencing counterclockwise rotation (e.g., Yin, 2006; Burgess et al., 2012), which is inconsistent with the model requirements.

Alternatively, rotation of northern Tibet could have been driven by an eastward decrease in gravitational potential energy (GPE) and GPE gradient in central Tibet (Molnar and England, 1997) (**Fig. 9b**). This GPE distribution may lead to faster spreading of the thickened lithosphere in the west than that in the east, causing clockwise rotation of northern Tibet. One potential problem is that this model would require north-south extension across central Tibet that is not

observed. However, the spreading-induced extensional strain may be overwhelmed by India-indentation-induced shortening strain in the region. It is also possible that fault rotation in northern Tibet is driven by a toroidal asthenospheric flow (**Fig. 9c**) that develops as the Burma trench, which is located just south of the Eastern Himalayan Syntaxis, retreats westward. The roll-back of the Burma slab causes poloidal flow of the asthenosphere around the slab, which draws in a toroidal flow around its northern edge. This inference is consistent with the current knowledge of mantle seismic anisotropy in the region (e.g., León Soto et al., 2012).

4.2 Implications for the Cenozoic development of northern Tibet

Recent work on the timing of structures in northern Tibet suggests a two-stage development of the region (**Fig. 3b**), with a change from thrust-dominated to strike-slip-dominated deformation at 20 to 15 Ma (e.g., Craddock et al., 2011; Duvall et al., 2013) due to a change in boundary conditions and stress regime (Yuan et al., 2013). As discussed by Clark (2012), the average bulk strain rate (i.e., $\dot{\epsilon} = v/L$) across the Himalayan-Tibetan orogen has remained constant since the onset of collision ($\dot{\epsilon} = \sim 7 \times 10^{-16} \text{ s}^{-1}$); deformation and narrowing of the orogen (L) has been balanced by slowed plate convergence (v) as a result of viscous mantle resistance. A constant strain for a viscous medium (i.e., the Tibetan-Eurasian lithosphere) equates to constant contractional stress acting on it, regardless of a shift in the geodynamics (e.g., crustal thickness, rheology, boundary conditions). Our calculated present-day shear strain rates and clockwise rotation rates across northern Tibet are consistent with geologic and paleomagnetic rotation rates. This similarity suggests that strain rates have remained constant throughout the Cenozoic, which is inconsistent with the discrete two-stage deformation model for the development of northern Tibet (e.g., Yuan et al., 2013) that requires a change in boundary

conditions. Alternatively, the non-rigid passive bookshelf model proposed in this study implies that coeval and kinematically linked thrust and strike-slip faulting could have formed under a constant stress regime.

The fundamental mode of Cenozoic deformation in northern Tibet appears to be protracted north-trending right-lateral simple shear (England and Molnar, 1990), nearly parallel to the northeast-striking Altyn Tagh fault, since the onset (or soon after) of the India-Eurasian collision at 60-55 Ma (e.g., Zhu et al., 2005; Green et al., 2008; Dupont-Nivet et al., 2010) (**Fig. 10**). Right-lateral shear strain was accommodated in the Eocene-Oligocene to early Miocene by distributed shortening and crustal-thickening in the Qimen Tagh, Qaidam Basin, and southern Qilian Shan thrust belts (**Fig. 10a**). An eastward decrease in north-south shortening strain was associated with the passive clockwise rotation of northern Tibet against the left-slip Altyn Tagh fault.

Crustal thickening led to a reorientation of the intermediate compressive stress orientation from horizontal to vertical by 20 to 15 Ma that shifted deformation to a mixed-mode of thrusting and left-slip faulting along preexisting weaknesses (i.e., Phanerozoic sutures) (**Fig. 10b**). The clockwise rotation and northward translation of the east-striking left-slip Haiyuan, Qinling, and Kunlun faults against the left-slip Altyn Tagh bounding fault zone in the northwest and the Longmen Shan in the southeast operated as passive bookshelf fault system. Combined northward translation and clockwise rotation of northern Tibet lead to minor observable right-slip shear along the eastern northeast-striking bounding faults (i.e., right-slip faulting within the thrust belt) (**Figs. 6b and 10b**). The rotating fault-bounded domains, surrounded by a relatively rigid medium (i.e., a strong Tarim and North China) (e.g., Molnar and Tapponnier, 1981), require fault-parallel deformation to resolve the space issues at the panel ends (**Fig. 10c**). Note

that the Altyn Tagh fault is oblique to the regional shear direction, and this orientation results in more shortening in the west than the east as the left-slip faults rotate against rigid Tarim and the Altyn Tagh boundary shear zone. Wallrock deformation and subsequent fault stretching (Means, 1989) results in negligible fault slip at the ends of each east-striking left-slip fault.

Several testable predictions emerge from our proposed model. First, the east-striking left-slip faults should be relatively weak. The western and central sections of the faults strike at a high angle to the maximum compressive stress direction (**Fig. 4b**) which indicates that minimal shear stress is acting on the fault planes. The faults also initiated along sutures (**Fig. 1**) that are presumably zones of preexisting weaknesses. The orthogonal orientation of the strike-slip faults to the maximum compressive stress direction in the west allows for the development of thrust belts that obliquely merge with the left-slip faults, as opposed to the eastern thrust belts that develop orthogonally to the strike-slip faults (**Fig. 1**). Second, the distinct two stages of deformation in northern Tibet (e.g., Yuan et al., 2013) may be apparent as a result of limited sampling and loose timing constraints. For example, we note that there are no direct age constraints for the initiation of the Haiyuan fault (Yuan et al., 2013) (**Fig. 3b**). The complex deformational patterns in northern Tibet make differentiating between thrust and strike-slip fault initiation difficult with current thermochronology techniques. Lastly, detailed studies of crustal strain can reveal if the fault-parallel shortening and extension in the off-fault regions equal offset on the strike-slip faults. There is currently little detailed structural data across the Qimen Tagh and Qilian Shan thrust belts to convincingly test this hypothesis, and GPS stations are sparse in these regions (**Fig. 4a**).

An important implication of the non-rigid bookshelf-fault model is that the style and magnitude of Cenozoic deformation in northern Tibet vary considerably in the east-west

direction. In order to understand processes of continental deformation and plateau construction, future geologic studies must establish along-strike variations of the structural framework along the left-slip faults to examine the relationship between discrete left-slip faulting and distributed off-fault deformation. It is clear that a single north-south cross section and its kinematic reconstruction across the northeastern margin of the Tibetan Plateau would not properly quantify the complex deformational processes of plateau formation.

5. Conclusions

The Cenozoic tectonic development of northern Tibet is characterized by the clockwise rotation of three major east-striking left-slip faults. Similar rotation and shear strain rates from paleomagnetic, geologic, and geodetic datasets (e.g., $\dot{\epsilon}_{xy} = \sim 10^{-16} \text{ s}^{-1}$) suggest continuous clockwise rotation of the region since the Eocene, which is consistent with a constant bulk strain rate across the Himalayan-Tibetan orogen since the onset of India-Eurasia collision (Clark, 2012). Our proposed non-rigid passive bookshelf-fault model, which relates discrete left-slip faulting to distributed off-fault deformation during regional clockwise rotation, accounts for the rotation of major left-slip faults, bidirectional decrease in slip from the central segments of these faults, eastward-tapering northwest-trending thrust belts, and four-quadrant strain pattern in the wallrock of the strike-slip faults. This model quantifies the relationship between strike-slip faulting and off-fault deformation during processes of continental tectonics and highlights the complexity of Cenozoic continental deformation in northern Tibet.

Acknowledgements. An Yin's research in Tibet has been supported by grants from the Tectonics Program of the US National Science Foundation. We thank Paul Mann and an

509 anonymous reviewer, as well as editor Jean-Philippe Avouac, for their constructive comments on
510 the manuscript.

511

References

- Avouac, J. P., and Tapponnier, P., 1993, Kinematic model of active deformation in central Asia: *Geophysical Research Letters*, v. 20, no. 10, p. 895-898.
- Burchfiel, B.C., Zhang, P. Wang, Y., Zhang, W., Song, F., Deng, Q., Molnar, P., and Royden, L., 1991, Geology of the Haiyuan fault zone, Ningxia Hui Autonomous Region, China, and its relation to the evolution of the northeastern margin of the Tibetan Plateau: *Tectonics*, v. 10, p. 1091-1110.
- Burchfiel, B.C., Chen, Z., Liu, Y., and Royden, L.H., 1995, Tectonics of the Longmen Shan and adjacent regions: *International Geology Review*, v. 37p. 661-735.
- Burgess, W.P., Yin, A., Dubey, C.S., Shen, Z.K., and Kelty, T. K., 2012, Holocene shortening across the Main Frontal Thrust zone in the eastern Himalaya: *Earth and Planetary Science Letters*, v. 357, p. 152-167.
- Cavalié, O., Lasserre, C., Doin, M. P., Peltzer, G., Sun, J., Xu, X., and Shen, Z. K., 2008, Measurement of interseismic strain across the Haiyuan fault (Gansu, China), by InSAR: *Earth and Planetary Science Letters*, v. 275, no. 3, p. 246-257.
- Chen, Y., Gilder, S., Halim, N., Cogné, J. P., Courtillot, V., 2002a, New paleomagnetic constraints on central Asian kinematics: Displacement along the Altyn Tagh fault and rotation of the Qaidam Basin: *Tectonics*, v. 21, no. 5, p. 6-1.
- Chen, Y., Wu, H., Courtillot, V., and Gilder, S., 2002b, Large N–S convergence at the northern edge of the Tibetan plateau? New Early Cretaceous paleomagnetic data from Hexi Corridor, NW China: *Earth and Planetary Science Letters*, v. 201, no. 2, p. 293-307.
- Cheng, F., Jolivet, M., Fu, S., Zhang, Q., Guan, S., Yu, X., and Guo, Z., 2014, Northward growth of the Qimen Tagh Range: A new model accounting for the Late Neogene strike-slip deformation of the SW Qaidam Basin: *Tectonophysics*, v. 632, p. 32-47.
- Cheng, F., Jolivet, M., Dupont-Nivet, G., Wang, L., Yu, X., and Guo, Z., 2015, Lateral extrusion along the Altyn Tagh Fault, Qilian Shan (NE Tibet): insight from a 3D crustal budget: *Terra Nova*, v. 27, no. 6, p. 416-425.

- 539 Clark, M.K., 2012, Continental collision slowing due to viscous mantle lithosphere rather than
540 topography: *Nature*, v. 483, no. 7387, p. 74-77.
- 541 Clark, M.K., Farley, K.A., Zheng, D., Wang, Z., and Duvall, A.R., 2010, Early Cenozoic faulting
542 of the northern Tibetan Plateau margin from apatite (U–Th)/He ages: *Earth and Planetary*
543 *Science Letters*, v. 296, no. 1, p. 78-88.
- 544 Cobbold, P.R., and Davy, P.H., 1988, Indentation tectonics in nature and experiment. 2. Central
545 Asia: *Bull. Geol. Inst. Univ. Uppsala*, v. 14, p. 143-162.
- 546 Cogné, J.P., Halim, N., Chen, Y., and Courtillot, V., 1999, Resolving the problem of shallow
547 magnetizations of Tertiary age in Asia: insights from paleomagnetic data from the
548 Qiangtang, Kunlun, and Qaidam blocks (Tibet, China), and a new hypothesis: *Journal of*
549 *Geophysical Research: Solid Earth*, v. 104, no. B8, p.17715-17734.
- 550 Craddock, W., Kirby, E., and Zhang, H., 2011, Late Miocene–Pliocene range growth in the
551 interior of the northeastern Tibetan Plateau: *Lithosphere*, v. 3, no. 6, p. 420-438.
- 552 Densmore, A.L., Ellis, M.A., Li, Y., Zhou, R., Hancock, G.S., and Richardson, N., 2007, Active
553 tectonics of the Beichuan and Pengguan faults at the eastern margin of the Tibetan
554 Plateau: *Tectonics*, v. 26, no. 4.
- 555 Dupont-Nivet, G., Guo, Z., Butler, R.F., Jia, C., 2002, Discordant paleomagnetic direction in
556 Miocene rocks from the central Tarim Basin: evidence for local deformation and
557 inclination shallowing: *Earth and Planetary Science Letters*, v. 199, no. 3, p. 473-482.
- 558 Dupont-Nivet, G., Horton, B.K., Butler, R.F., Wang, J., Zhou, J., and Waanders, G.L., 2004,
559 Paleogene clockwise tectonic rotation of the Xining-Lanzhou region, northeastern
560 Tibetan Plateau. *Journal of Geophysical Research*, v. 109, DOI: 10.1029/2003JB002620.
- 561 Dupont-Nivet, G., Lippert, P.C., Van Hinsbergen, D.J., Meijers, M.J., and Kapp, P., 2010,
562 Palaeolatitude and age of the Indo–Asia collision: palaeomagnetic constraints:
563 *Geophysical Journal International*, v. 182, no. 3, p. 1189-1198.
- 564 Duvall, A.R., and Clark, M.K., 2010, Dissipation of fast strike-slip faulting within and beyond
565 northeastern Tibet: *Geology*, v. 38, p. 223-226.

566 Duvall, A.R., Clark, M.K., Kirby, E., Farley, K.A., Craddock, W.H., Li, C., and Yuan, D.Y.,
567 2013, Low-temperature thermochronometry along the Kunlun and Haiyuan faults, NE
568 Tibetan Plateau: Evidence for kinematic change during late-stage orogenesis: *Tectonics*,
569 32(5), p. 1190-1211.

570 England, P., and Houseman, G., 1986, Finite strain calculations of continental deformation: 2.
571 Comparison with the India-Asia collision zone: *Journal of Geophysical Research: Solid*
572 *Earth*, v. 91, no. B3, p. 3664-3676.

573 England, P., and Molnar, P., 1990, Right-lateral shear and rotation as the explanation for strike-
574 slip faulting in eastern Tibet: *Nature*, v. 344, p. 140-142.

575 England, P., and Molnar, P., 1997, Active deformation of Asia: From kinematics to dynamics:
576 *Science*, v. 278, p. 647-650.

577 Freund, R., 1970, Rotation of strike slip faults in Sistan, southeast Iran: *Journal of Geology*, v.
578 78, p. 188-200.

579 Frost, G.M., et al., 1995, Preliminary early Cretaceous paleomagnetic results from the Gansu
580 Corridor, China: *Earth and Planetary Science Letters*, v. 129, no. 1, p. 217-232.

581 Fu, B. and Awata, Y., 2007, Displacement and timing of left-lateral faulting in the Kunlun Fault
582 Zone, northern Tibet, inferred from geologic and geomorphic features: *Journal of Asian*
583 *Earth Sciences*, v. 29, p. 253-265.

584 Fu, B., Awata, Y., Du, J., and He, W., 2005, Late Quaternary systematic stream offsets caused
585 by repeated large seismic events along the Kunlun fault, northern Tibet: *Geomorphology*,
586 v. 71, no. 3, p. 278-292.

587 Gan, W., Zhang, P., Shen, Z.-K., Niu, Z., Wang, M., Wan, Y., Zhou, D., and Cheng, J., 2007,
588 Present-day crustal motion within the Tibetan Plateau inferred from GPS measurements:
589 *Journal of Geophysical Research*, v. 112, no. B8.

590 Gao, R., Wang, H., Yin, A., Dong, S., Kuang, Z., Zuza, A. V., Li, W., and Xiong, X., 2013,
591 Tectonic development of the northeastern Tibetan Plateau as constrained by high-
592 resolution deep seismic-reflection data: *Lithosphere*, v. 5, no. 6, p. 555-574.

593 Gaudemer, Y., Tapponnier, P., Meyer, B., Peltzer, G., Guo, S., Chen, Z., Dai, H, and Cifuentes,
594 I., 1995, Partitioning of crustal slip between linked, active faults in the eastern Qilian
595 Shan, and evidence for a major seismic gap, the ‘Tianzhu gap’, on the western Haiyuan
596 Fault, Gansu (China): *Geophysical Journal International*, v. 120, p. 599-645.

597 Godard, V., Pik, R., Lavé, J., Cattin, R., Tibari, B., De Sigoyer, J., Pubellier, M., and Zhu, J,
598 2009, Late Cenozoic evolution of the central Longmen Shan, eastern Tibet: insight from
599 (U-Th)/He thermochronometry: *Tectonics*, v. 28, no. 5.

600 Green, O.R., Searle, M.P., Corfield, R.I., and Corfield, R.M., 2008, Cretaceous-Tertiary
601 carbonate platform evolution and the age of the India-Asia collision along the Ladakh
602 Himalaya (Northwest India): *The Journal of Geology*, v. 116, no. 4, p. 331-353.

603 Halim, N., et al., 1998, New Cretaceous and Early Tertiary paleomagnetic results from Xining-
604 Lanzhou basin, Kunlun and Qiangtang blocks, China: Implications on the geodynamic
605 evolution of Asia. *Journal of Geophysical Research: Solid Earth*, v. 103, no. B9, p.
606 21025-21045.

607 Halim, N., Chen, Y., and Cogné, J.P., 2003, A first palaeomagnetic study of Jurassic formations
608 from the Qaidam basin, Northeastern Tibet, China—tectonic implications: *Geophysical*
609 *Journal International*, v. 153, no. 1, p. 20-26.

610 Harkins, N., and Kirby, E., 2008, Fluvial terrace riser degradation and determination of slip rates
611 on strike - slip faults: An example from the Kunlun fault, China: *Geophysical Research*
612 *Letters*, v. 35, no. 5.

613 Harkins, N., Kirby, E., Shi, X., Wang, E., Burbank, D., and Chun, F., 2010, Millennial slip rates
614 along the eastern Kunlun fault: Implications for the dynamics of intracontinental
615 deformation in Asia: *Lithosphere*, v. 2, p. 247-266.

616 Heidbach, O., Tingay, M., Barth, A., Reinecker, J., and Kurfeß, D., Müller, B., 2010, Global
617 crustal stress pattern based on the World Stress Map database release 2008:
618 *Tectonophysics*, v. 482, no. 1, p. 3-15.

619 Hubbard, J. and Shaw, J. H., 2009, Uplift of the Longmen Shan and Tibetan plateau, and the
620 2008 Wenchuan (M= 7.9) earthquake: *Nature*, v. 458, no. 7235, p. 194-197.

621 Jolivet, M., Brunel, M., Seward, D., Xu, Z., Yang, J., Malavieille, J., Roger, F., Leyreloup, A.,
622 Arnaud, N., and Wu, C., 2003, Neogene extension and volcanism in the Kunlun fault
623 zone, northern Tibet: New constraints on the age of the Kunlun fault: *Tectonics*, v. 22,
624 DOI: 10.1029/2002TC001428.

625 Kidd, W.S., and Molnar, P., 1988, Quaternary and active faulting observed on the 1985
626 Academia Sinica--Royal society geotraverse of Tibet: *Philosophical Transactions of the*
627 *Royal Society of London. Series A*, v. 327, 337-363.

628 Kirby, E., Whipple, K. X., Burchfiel, B. C., Tang, W., Berger, G., Sun, Z., and Chen, Z., 2000,
629 Neotectonics of the Min Shan, China: Implications for mechanisms driving Quaternary
630 deformation along the eastern margin of the Tibetan Plateau: *Geological Society of*
631 *America Bulletin*, v. 112, no. 3, p. 375-393.

632 Kirby, E., Harkins, N., Wang, E., Shi, X., Fan, C., and Burbank, D., 2007, Slip rate gradients
633 along the eastern Kunlun fault: *Tectonics*, v. 26, no. 2.

634 Lasserre, C. et al., 1999, Postglacial left slip rate and past occurrence of M> 8 earthquakes on the
635 western Haiyuan fault, Gansu, China: *Journal of Geophysical Research: Solid Earth*, v.
636 104, no. B8, p. 17633-17651.

637 Lasserre, C., Gaudemer, Y., Tapponnier, P., Mériaux, A. S., Van der Woerd, J., Daoyang, Y.,
638 Ryerson, F. J., Finkel, R. C., and Caffee, M. W., 2002, Fast late Pleistocene slip rate on
639 the Leng Long Ling segment of the Haiyuan fault, Qinghai, China: *Journal of*
640 *Geophysical Research: Solid Earth*, v. 107, no. B11.

641 Lease, R.O., Burbank, D.W., Zhang, H., Liu, J., and Yuan, D., 2012, Cenozoic shortening budget
642 for the northeastern edge of the Tibetan Plateau: Is lower crustal flow necessary?:
643 *Tectonics*, v. 31, no. 3.

644 León Soto, G., Sandvol, E., Ni, J.F., Flesch, L., Hearn, T.M., Tilmann, F., Chen, J., and Brown,
645 L.D., 2012, Significant and vertically coherent seismic anisotropy beneath eastern Tibet:
646 Journal of Geophysical Research: Solid Earth, v. 117, no. B5.

647 Li, C., Zhang, P., Yin, J., and Min, W., 2009, Late Quaternary left-lateral slip rate of the Haiyuan
648 fault, northeastern margin of the Tibetan Plateau: Tectonics, v. 28.

649 Li, H., et al., 2005, Slip rate on the Kunlun fault at Hongshui Gou, and recurrence time of great
650 events comparable to the 14/11/2001, Mw ~7.9 Kokoxili earthquake, Earth and Planetary
651 Science Letters, v. 237, p. 285-299.

652 Luyendyk, B.P., Kamerling, M.J., and Terres, R., 1980, Geometric model for Neogene crustal
653 rotations in southern California: Geological Society of America Bulletin, v. 91, no. 4, p.
654 211-217.

655 Meade, B.J., 2007, Present-day kinematics at the India-Asia collision zone: Geology, v. 35, no.
656 1, p. 81-84.

657 Means, W.D., 1989, Stretching faults: Geology, v. 17, v. 10, p. 893-896.

658 Means, W.D., 1990, One-dimensional kinematics of stretching faults: Journal of Structural
659 Geology, v. 12, no. 2, p. 267-272.

660 Mercier, J.L., Vergely, P., Zhang, Y.Q., Hou, M.J., Bellier, O., and Wang, Y.M., 2013,
661 Structural records of the Late Cretaceous–Cenozoic extension in Eastern China and the
662 kinematics of the Southern Tan-Lu and Qinling Fault Zone (Anhui and Shaanxi
663 provinces, PR China): Tectonophysics, v. 582, p. 50-75.

664 Meyer, B., Tapponnier, P., Bourjot, L., Métivier, F., Gaudemer, Y., Peltzer, G., Shunmin, G.,
665 and Zhitai, C., 1998, Crustal thickening in Gansu-Qinghai, lithospheric mantle
666 subduction, and oblique, strike-slip controlled growth of the Tibet Plateau: Geophysical
667 Journal International, v. 135, no. 1, p. 1–47, doi:10.1046/j.1365-246X.1998.00567.x.

668 Mock, C., Arnaud, N. O., and Cantagrel, J. M., 1999, An early unroofing in northeastern Tibet?
669 Constraints from $^{40}\text{Ar}/^{39}\text{Ar}$ thermochronology on granitoids from the eastern Kunlun

670 range (Qianghai, NW China): *Earth and Planetary Science Letters*, v. 171, no. 1, p. 107-
671 122.

672 Molnar, P., and Tapponnier, P., 1981, A possible dependence of tectonic strength on the age of
673 the crust in Asia: *Earth and Planetary Science Letters*, v. 52, p. 107–114.

674 Onderdonk, N.W, 2005, Structures that accommodated differential vertical axis rotation of the
675 western Transverse Ranges, California: *Tectonics*, v. 24, no. 4.

676 Peltzer, G. and Saucier, F., 1996, Present-day kinematics of Asia derived from geologic fault
677 rates: *Journal of Geophysical Research: Solid Earth*, v. 101, no. B12, p. 27943-27956.

678 Peltzer, G. and Tapponnier, P., 1988, Formation and evolution of strike - slip faults, rifts, and
679 basins during the India - Asia collision: An experimental approach: *Journal of*
680 *Geophysical Research: Solid Earth*, v. 93, no. B12, p. 15085-15117.

681 Platt, J.P. and Becker, T. W., 2010, Where is the real transform boundary in California?:
682 *Geochemistry, Geophysics, Geosystems*, v. 11, no. 6.

683 Platt, J.P. and Becker, T. W., 2013, Kinematics of rotating panels of E–W faults in the San
684 Andreas system: what can we tell from geodesy?: *Geophysical Journal International*, v.
685 194, no. 3, p. 1295-1301.

686 Ramsay, J.G., and Huber, M.I., 1983, *The techniques of modern structural geology. Vol. 1.*
687 *Strain Analysis*: Academic Press, London.

688 Ratschbacher, L., Hacker, B.R., Calvert, A., Webb, L.E., Grimmer, J.C., McWilliams, M.O,
689 Ireland, T., Dong, S., and Hu, J., Tectonics of the Qinling (Central China):
690 tectonostratigraphy, geochronology, and deformation history: *Tectonophysics*, v. 366, p.
691 1-53.

692 Reith, R.C., 2013, *Structural Geology of a Central Segment of the Qilian Shan-Nan Shan Thrust*
693 *Belt: Implications for the Magnitude of Cenozoic Shortening in the Northeastern Tibetan*
694 *Plateau [M.S. thesis]: Los Angeles, University of California, 73 p.*

695 Ryan, W.B.F., Carbotte, S.M., Coplan, J.O., O'Hara, S., Melkonian, A., Arko, R., Weissel, R.A.,
696 Ferrini, V., Goodwillie, A. Nitsche, F., Bonczkowski, J., and Zemsky, R., 2009, Global
697 Multi-Resolution Topography synthesis: *Geochemistry Geophysics Geosystems*, v. 10,
698 no. 8.

699 Sun, Z.M., Yang, Z.Y., Pei, J.L., Yang, T.S., and Wang, X.S., 2006, New Early Cretaceous
700 paleomagnetic data from volcanic and red beds of the eastern Qaidam Block and its
701 implications for tectonics of Central Asia: *Earth and Planetary Science Letters*, v. 243, p.
702 268-281.

703 Tapponnier, P., Peltzer, G., Le Dain, A.Y., Armijo, R., and Cobbold, P., 1982, Propagating
704 extrusion tectonics in Asia: new insights from simple experiments with plasticine:
705 *Geology*, v. 10, no. 12, p. 611-616.

706 Tapponnier, P., Xu, Z., Roger, F., Meyer, B., Arnaud, N., Wittlinger, G., and Yang, J., 2001,
707 Oblique stepwise rise and growth of the Tibet Plateau: *Science*, v. 294, p. 1671-1677.

708 Taylor, M., Yin, A., 2009, Active structures of the Himalayan-Tibetan orogen and their
709 relationships to earthquake distribution, contemporary strain field, and Cenozoic
710 volcanism: *Geosphere*, v. 5, p. 199-214.

711 Thatcher, W., 2007, Microplate model for the present - day deformation of Tibet. *Journal of*
712 *Geophysical Research: Solid Earth*, v. 112, no. B1.

713 Van Der Woerd, J., Ryerson, F.J., Tapponnier, P., Gaudemer, Y., Finkel, R., Mériaux, A.S.,
714 Caffee, M., Guoguang, Z., and Qunlu, H., 1998, Holocene left-slip rate determined by
715 cosmogenic surface dating on the Xidatan segment of the Kunlun fault (Qinghai, China):
716 *Geology*, v. 26, no. 8, p. 695-698.

717 Van Der Woerd, J., Ryerson, F.J., Tapponnier, P., Meriaux, A.S., Gaudemer, Y., Meyer, B.,
718 Finkel, R., Caffee, M., Guoguang, Z., and Zhiqin, X., 2000, Uniform slip-rate along the
719 Kunlun Fault: Implications for seismic behaviour and large-scale tectonics: *Geophysical*
720 *Research Letters*, v. 27, no. 16, p. 2353-2356.

721 Van Der Woerd, J. et al., 2002, Uniform postglacial slip-rate along the central 600 km of the
722 Kunlun Fault (Tibet), from ²⁶Al, ¹⁰Be, and ¹⁴C dating of riser offsets, and climatic
723 origin of the regional morphology: *Geophysical Journal International*, v. 148, no. 3, p.
724 356-388.

725 Van Hinsbergen, D. J., Kapp, P., Dupont - Nivet, G., Lippert, P. C., DeCelles, P. G., and
726 Torsvik, T. H., 2011, Restoration of Cenozoic deformation in Asia and the size of Greater
727 India: *Tectonics*, v. 30, no. 5.

728 Wang, E. and Burchfiel, B.C, 2004, Late Cenozoic right-lateral movement along the Wenquan
729 Fault and associated deformation: implications for the kinematic history of the Qaidam
730 Basin, Northeastern Tibetan Plateau: *International Geology Review*, v. 46, no. 10, p. 861-
731 879.

732 Wang, C., Gao, R., Yin, A., Wang, H., Zhang, Y., Guo, T., Li, Q., and Li, Y., 2011, A mid-
733 crustal strain-transfer model for continental deformation: A new perspective from high-
734 resolution deep seismic-reflection profiling across NE Tibet: *Earth and Planetary Science*
735 *Letters*, v. 306, no. 3, p. 279-288.

736 Wang, E. et al., 2012, Two-phase growth of high topography in eastern Tibet during the
737 Cenozoic: *Nature Geoscience*, v. 5, no. 9, p. 640-645.

738 Weldon, R. and Humphreys, E., 1986, A kinematic model of southern California: *Tectonics*, v. 5,
739 no. 1, p. 33-48.

740 Wu, C., Yin, A., Zuza, A. V., Zhang, J., Liu, W., and Ding, L., 2016, Pre-Cenozoic Geologic
741 History of the Central and Northern Tibetan Plateau and the Role of Wilson Cycles in
742 Constructing the Tethyan Orogenic System: *Lithosphere*, in press.

743 Yin, A., 2000, Mode of Cenozoic east-west extension in Tibet suggesting a common origin of
744 rifts in Asia during the Indo-Asian collision. *Journal of Geophysical Research: Solid*
745 *Earth*, v. 105, no. B9, p. 21745-21759.

746 Yin, A., 2006, Cenozoic tectonic evolution of the Himalayan orogen as constrained by along
747 strike variation of structural geometry, exhumation history, and foreland sedimentation:
748 Earth-Science Reviews, v. 76, no. 1, p. 1-131.

749 Yin, A., 2010a, Cenozoic tectonic evolution of Asia: A preliminary synthesis: Tectonophysics, v.
750 488, no. 1, p. 293-325.

751 Yin, A., 2010b, A special issue on the great 12 May 2008 Wenchuan earthquake (M_w 7.9):
752 Observations and unanswered questions: Tectonophysics, v. 491, p. 1-9.

753 Yin, A., Dang, Y.-Q., Zhang, M., McRivette, M.W., Burgess, W.P., and Chen, X.-H., 2007,
754 Cenozoic tectonic evolution of Qaidam Basin and its surrounding regions (part 2): Wedge
755 tectonics in southern Qaidam Basin and the Eastern Kunlun Range, in Sears, J.W.,
756 Harms, T.A., and Evenchick, C.A., eds., Whence the Mountains? Inquiries into the
757 Evolution of Orogenic Systems: A Volume in Honor of Raymond A. Price: Geological
758 Society of America Special Paper 433, p. 369-390.

759 Yin, A., Dang, Y.-Q., Wang, L.-C., Jiang, W.-M., Zhou, S.-P, Chen, X. -H., Gehrels, G.E., and
760 McRivette, M.W., 2008a, Cenozoic tectonic evolution of Qaidam Basin and its
761 surrounding regions (part 1): The southern Qilian Shan–Nan Shan thrust belt and
762 northern Qaidam Basin: Geological Society of America Bulletin, v. 120, no. 7/8, p. 813–
763 846, doi: 10.1130.B26180.1.

764 Yin, A., Dang, Y.Q., Zhang, M., Chen, X.H., and McRivette, M.W., 2008b. Cenozoic tectonic
765 evolution of the Qaidam basin and its surrounding regions (Part 3): Structural geology,
766 sedimentation, and regional tectonic reconstruction. Geological Society of America
767 Bulletin, v. 120, no. 7-8, p. 847-876.

768 Yin, A. and Taylor, M.H., 2011, Mechanics of V-shaped conjugate strike-slip faults and the
769 corresponding continuum mode of continental deformation: Geological Society of
770 America Bulletin, v. 123, no. 9-10, p. 1798-1821.

771 Yin, A., and Pappalardo, R.T., 2015, Gravitational spreading, bookshelf faulting, and tectonic
772 evolution of the South Polar Terrain of Saturn’s moon Enceladus: Icarus, v. 260, p. 409-
773 439.

774 Yuan, W., Dong, J., Shicheng, W., and Carter, A., 2006, Apatite fission track evidence for
775 Neogene uplift in the eastern Kunlun Mountains, northern Qinghai–Tibet Plateau, China:
776 *Journal of Asian Earth Sciences*, v. 27, no. 6, p. 847-856.

777 Yuan et al., 2013, The growth of northeastern Tibet and its relevance to large-scale continental
778 geodynamics: A review of recent studies: *Tectonics*, v. 32, p. 1358-1370.

779 Zhang, P. et al., 1991, Amount and style of late Cenozoic deformation in the Liupan Shan area,
780 Ningxia Autonomous Region, China: *Tectonics*, v. 10, no. 6, p. 1111-1129.

781 Zhang, Y.Q., Mercier, J. L., and Vergély, P., 1998, Extension in the graben systems around the
782 Ordos (China), and its contribution to the extrusion tectonics of south China with respect
783 to Gobi-Mongolia: *Tectonophysics*, v. 285, no. 1, p. 41-75.

784 Zhang, P.Z. et al., 2004, Continuous deformation of the Tibetan Plateau from global positioning
785 system data: *Geology*, v. 32, no. 9, p. 809-812.

786 Zhang, Y.Q., Vergely, P., and Mercier, J., 1995, Active faulting in and along the Qinling Range
787 (China) inferred from SPOT imagery analysis and extrusion tectonics of south China:
788 *Tectonophysics*, v. 243, no. 1, p. 69-95.

789 Zheng, D. et al., 2006, Rapid exhumation at~ 8 Ma on the Liupan Shan thrust fault from apatite
790 fission-track thermochronology: Implications for growth of the northeastern Tibetan
791 Plateau margin: *Earth and Planetary Science Letters*, v. 248, no. 1, p. 198-208.

792 Zheng, D., Clark, M.K., Zhang, P., Zheng, W., and Farley, K.A., 2010, Erosion, fault initiation
793 and topographic growth of the North Qilian Shan (northern Tibetan Plateau): *Geosphere*,
794 v. 6, p. 937-941.

795 Zhu, B., Kidd, W.S.F., Rowley, D.B., Currie, B.S., and Shafique, N., 2005, Age of initiation of
796 the India–Asia collision in the east-central Himalaya: *J. Geol.*, v. 113, p. 265–285.

797 Zuza, A.V., Cheng, X., Yin, A., 2016, Testing models of Tibetan Plateau formation with
798 Cenozoic shortening estimates across the Qilian Shan-Nan Shan thrust belt: *Geosphere*,
799 v. 12, no. 2, p. 1-32.

800 Zuza, A.V., Yin, A., 2013, Ductile bookshelf faulting: A new kinematic model for Cenozoic
801 deformation in northern Tibet: AGU Fall Meeting Abstracts, v. 1, p. 2.

802 Zuza, A.V., Yin, A., Li, Jianhua, 2014, Deciphering the coupled Paleozoic and Cenozoic tectonic
803 history of the Qilian Shan, northeastern Tibetan Plateau: AGU Fall Meeting
804 Abstracts, v.1 p. 4563.

805

806

Figure Captions

Figure 1. Color-shaded relief map showing Cenozoic faults related to the India-Eurasia collision zone after Yin et al. (2008a), Taylor and Yin (2009), and Zuza et al. (2016). Also shown are Harvard centroid moment tensor (CMT) earthquake focal mechanisms from 1 January 1977 to 1 January 2009 of events $>M5.5$. Inset shows location of northern Tibet in the context of the Himalayan-Tibetan Orogen. The digital topographic basemap is from GeoMapApp software (Ryan et al., 2005) available at <http://www.geomapapp.org/>.

Figure 2. Conceptual models for the space issue that arises in a bookshelf fault system given an (a) initial configuration with three bookshelf panels bounded on the bottom and top with rigid but movable blocks. Three mechanisms have been proposed to conserve area at the ends of the panels, including (b) lithospheric-scale basin development (e.g., Luyendyk et al., 1980), (c) crustal shortening at the ends of the panels (e.g., Onderdonk, 2005), and (d) fault-parallel stretching throughout the panels (e.g., Platt and Becker, 2013). Note that in each case, the width of the system (W) is not constant.

Figure 3. (a) Simplified sketches showing end-member models for the role of the east-striking left-slip faults (i.e., the Haiyuan, Qinling, and Kunlun faults) in the Cenozoic tectonic development of northern Tibet. Only one left-slip fault is shown in each scenario but the same model predictions apply to each of left-slip faults in northern Tibet. Annotations show, as an example, how predicted structures match the geology for Haiyun fault and linked termination structures. The (1) extrusion fault model (e.g., Tapponnier et al., 1982) predicts the presence of a conjugate right-slip fault with a similar slip magnitude near the western end the fault, the (2)

transfer fault model (e.g., Burchfiel et al., 1991; Duvall and Clark, 2010) requires a left-slip component across the termination thrusts, the (3) rigid bookshelf fault model (e.g., England and Molnar, 1990) predicts gaps and/or overlaps at the ends of the rigid blocks, and the (4) non-rigid bookshelf fault model predicts fault-parallel stretching and/or shortening to preserve mass at the panel ends (this study). (b) Ages of deformation in northern Tibet from numerous sources. Ages represent initiation of fault activity. Faults initiated in the Paleogene are in blue whereas faults initiated in the Neogene are red. Modified from Yuan et al. (2013).

Figure 4. (a) Global positioning system (GPS) velocity field across Tibet relative to stable Eurasia modified from Gan et al. (2007). Also shown is the location of Figure 4b. (b) GPS velocity field relative to stable Eurasia with data from Gan et al. (2007). Colored vectors represent stations from four distinct domains that bounded by the left-slip Kunlun, Qinling, and Haiyuan faults, from north to south to north: East Central Tibet (ECT), Qaidam Basin (QB), North Qaidam (NQ), and Qilian Shan (QS). Paleomagnetic rotation magnitudes are shown with white arrows along with the lower age bound for the onset of rotation. The averaged maximum compressive-stress directions from the World Stress Map are plotted as green lines (Heidbach et al., 2008). Gold star shows the reference location for analyses shown in Figure 4c. Also shown is the x - y coordinate system, with the x -axis parallel to N110°E. Paleomagnetic rotation data are from (1) Halim et al. (2003), (2) Chen et al. (2002a), (3) Chen et al. (2002b), (4) Frost et al. (1995), (5) Halim et al. (1998), (6) Cogné et al. (1999), (7) Dupont-Nivet et al., (2004), and (8) Dupont-Nivet et al. (2002). (c) GPS velocity component in the N20°E direction perpendicular to the general trend of the left-slip Kunlun, Qinling, and Haiyuan faults in northern Tibet. The

N20°E velocity components relative to a fixed point in the Tibetan Plateau foreland (i.e., gold star in Figure 4b) in each fault-bounded domain (color codes corresponding to those in Figure 4b) are projected onto a N110°E-trending line. Best-fit (blue, yellow and red dashed lines) velocity gradients are bounded by shear strain rates ($\dot{\epsilon}_{xy}$) of 10^{-16} s^{-1} and $5 \times 10^{-16} \text{ s}^{-1}$ shown as thick black lines.

Figure 5. (a) Simplified diagram highlighting a possible four-quadrant deformation pattern with a strike-slip fault and its coeval termination structures. Note that fault-parallel extension can be accommodated by either normal or strike-slip faulting. (b) Sketch map highlighting the relationship between east-striking left-slip faults and off-fault wallrock deformation. The Haiyuan, Qinling, and Kunlun faults all display a four-quadrant strain distribution, with fault-parallel shortening in the northwest and southeast quadrants and fault-parallel extension in the southwest and northeast quadrants. Inset shows how the non-rigid bookshelf faulting can generate the observed four-quadrant strain distribution. Fault-perpendicular deformation accommodates clockwise rotation of the faults whereas fault-parallel deformation is required to resolve the “gap” and “overlap” issues that arise at the panel ends. Note that the specific strain asymmetry corresponds to a clockwise fault rotation direction.

Figure 6. (a) Conceptual model for the active bookshelf fault system (Freund, 1970) where shear is driven by drag along the boundary walls and the sense of shear of the bounding shear zones must be opposite of the slip between the bookshelf panels. (b) Conceptual model for a passive bookshelf fault system, where panels rotating about the center of the system are driven by a push from the bottom. The panels can both rotate and translate. Note that at some point between stages

(2) and (3), the easternmost bounding fault transitions from left-slip to right slip, and at this point there is negligible strike-slip offset at this right-side boundary.

Figure 7. Velocity fields for the (a) active and (b) passive bookshelf fault systems. Note that the velocity field in the passive system involves the summation of translational (v_t) and rotational (v_r) velocities. See text for discussion.

Figure 8. Kinematic model illustrating the required fault-parallel strain at the western end of one passive bookshelf fault. Given a rigid left-side boundary, left-lateral offset on the fault (d) caused by clockwise fault rotation requires fault-parallel shortening and stretching strain (ϵ_{xx}), for panels A and B respectively, related to approximately one-half the fault slip divided by half the length of each panel, L : $\epsilon^A_{xx} \approx 0.5d/L$ and $\epsilon^B_{xx} \approx -0.5d/L$. In this example, the angle of rotation, θ , is very small, such that $\tan\theta$ can be approximated as θ . See text for discussion.

Figure 9. Passive bookshelf faulting in northern Tibet may be driven by (a) boundary-zone rotation to the south, (b) gravitational spreading of thick central Tibet lithosphere with the distribution of gravitational potential energy after England and Molnar (1997), or (c) a toroidal flow in the asthenosphere (e.g., León Soto et al., 2012). See text for further discussion.

Figure 10. Cenozoic tectonic evolution of northern Tibet, showing the timing of major fault initiation where grey structures are already active at given time interval. Note orientation of north arrows. (a) Right-lateral shear, driven from the south, begins in the Eocene and is expressed by north-south shortening in the Qimen Tagh, North Qaidam, and southern Qilian

899 Shan thrust belts. An eastward decrease in north-south shortening magnitude and thrust spacing
900 indicates clockwise rotation of northern Tibet. (b) Progressive crustal thickening causes the
901 initiation of the east-striking left-slip faults along preexisting weaknesses. Continued right-lateral
902 shear drives clockwise rotation of these faults and their wallrock passively against the left-slip
903 Atlyn Tagh bounding fault zone. (c) The left-slip faults in this passive bookshelf fault system
904 develop a four-quadrant deformation pattern with domains of fault-parallel stretching and
905 shortening. Age data from Craddock et al. (2011), Yuan et al. (2013), Duvall et al. (2013), and
906 references therein.

Zuza and Yin - Figure 1

Figure 1. Color-shaded relief map showing Cenozoic faults related to the Indo-Asian collision zone after Yin et al. (2008a), Taylor and Yin (2009), and Zuza et al. (2016). Also shown are Harvard centroid moment tensor (CMT) earthquake focal mechanisms from 1 January 1977 to 1 January 2009 of events $>M 5.5$. Inset shows location of northern Tibet in the context of the Himalayan-Tibetan Orogen. The digital topographic basemap is from GeoMapApp software (Ryan et al., 2005) available at <http://www.geomapapp.org/>.

(a) Initial configuration

(b) Basin development to conserve area (e.g., Luyendyk et al., 1980)

(c) Shortening to conserve area (e.g., Onderdonk, 2005)

(d) Fault-parallel stretching (e.g., Platt and Becker, 2013)

Zuza and Yin - Figure 2

Figure 2. Conceptual models for the space issue that arises in a bookshelf fault system given an (a) initial configuration with three bookshelf panels bounded on the bottom and top with rigid but movable blocks. Three mechanisms have been proposed to conserve area at the ends of the panels, including (b) lithospheric-scale basin development (e.g., Luyendyk et al., 1980), (c) crustal shortening at the ends of the panels (e.g., Onderdonk, 2005), and (d) fault-parallel stretching throughout the panels (e.g., Platt and Becker, 2013). Note that in each case, the width of the system (W) is not constant.

Zuza and Yin - Figure 3

Figure 3. (a) Simplified sketches showing end-member models for the role of the east-striking left-slip faults (i.e., the Haiyuan, Qinling, and Kunlun faults) in the Cenozoic tectonic development of northern Tibet. Only one left-slip fault is shown in each scenario but the same model predictions apply to each of left-slip faults in northern Tibet. Annotations show, as an example, how predicted structures match the geology for Haiyuan fault and linked termination structures. The (1) extrusion fault model (e.g., Tapponnier et al., 1982) predicts the presence of a conjugate right-slip fault with a similar slip magnitude near the western end the fault, the (2) transfer fault model (e.g., Burchfiel et al., 1991; Duvall and Clark, 2010) requires a left-slip component across the termination thrusts, the (3) rigid bookshelf fault model (e.g., England and Molnar, 1990) predicts gaps and/or overlaps at the ends of the rigid blocks, and the (4) non-rigid bookshelf fault model predicts fault-parallel stretching and/or shortening to preserve mass at the panel ends (this study). (b) Ages of deformation in northern Tibet from numerous sources. Ages represent initiation of fault activity. Faults initiated in the Paleogene are in blue whereas faults initiated in the Neogene are red. Modified from Yuan et al. (2013).

Zuza and Yin - Figure 4

Figure 4. (a) Global positioning system (GPS) velocity field across Tibet relative to stable Eurasia modified from Gan et al. (2007). Also shown is the location of Figure 4b. (b) GPS velocity field relative to stable Eurasia with data from Gan et al. (2007). Colored vectors represent stations from four distinct domains that bounded by the left-slip Kunlun, Qinling, and Haiyuan faults, from north to south to north: East Central Tibet (ECT), Qaidam Basin (QB), North Qaidam (NQ), and Qilian Shan (QS). Paleomagnetic rotation magnitudes are shown with white arrows along with the lower age bound for the onset of rotation. The averaged maximum compressive-stress directions from the World Stress Map are plotted as green lines (Heidbach et al., 2008). Gold star shows the reference location for analyses shown in Figure 4c. Also shown is the x - y coordinate system, with the x -axis parallel to N110°E. Paleomagnetic rotation data are from (1) Halim et al. (2003), (2) Chen et al. (2002a), (3) Chen et al. (2002b), (4) Frost et al. (1995), (5) Halim et al. (1998), (6) Cogné et al. (1999), (7) Dupont-Nivet et al., (2004), and (8) Dupont-Nivet et al. (2002). (c) GPS velocity component in the N20°E direction perpendicular to the general trend of the left-slip Kunlun, Qinling, and Haiyuan faults in northern Tibet. The N20°E velocity components relative to a fixed point in the Tibetan Plateau foreland (i.e., gold star in Figure 4b) in each fault-bounded domain (color codes corresponding to those in Figure 4b) are projected onto a N110°E-trending line. Best-fit (blue, yellow and red dashed lines) velocity gradients are bounded by shear strain rates (ϵ_{xy}) of 10^{-16} s^{-1} and $5 \times 10^{-16} \text{ s}^{-1}$ shown as thick black lines.

Zuza and Yin - Figure 5

Figure 5. (a) Simplified diagram highlighting a possible four-quadrant deformation pattern with a strike-slip fault and its coeval termination structures. Note that fault-parallel extension can be accommodated by either normal or strike-slip faulting. (b) Sketch map highlighting the relationship between east-striking left-slip faults and off-fault wallrock deformation. The Haiyuan, Qinling, and Kunlun faults all display a four-quadrant strain distribution, with fault-parallel shortening in the northwest and southeast quadrants and fault-parallel extension in the southwest and northeast quadrants. Inset shows how the non-rigid bookshelf faulting can generate the observed four-quadrant strain distribution. Fault-perpendicular deformation accommodates clockwise rotation of the faults whereas fault-parallel deformation is required to resolve the “gap” and “overlap” issues that arise at the panel ends. Note that the specific strain asymmetry corresponds to a clockwise fault rotation direction.

Zuza and Yin - Figure 6

Figure 6. (a) Conceptual model for the active bookshelf fault system (Freund, 1970) where shear is driven by drag along the boundary walls and the sense of shear of the bounding shear zones must be opposite of the slip between the bookshelf panels. (b) Conceptual model for a passive bookshelf fault system, where panels rotating about the center of the system are driven by a push from the bottom. The panels can both rotate and translate. Note that at some point between stages (2) and (3), the easternmost bounding fault transitions from left-slip to right slip, and at this point there is negligible strike-slip offset at this right-side boundary.

(a) Velocity field across an active bookshelf fault system

Linear velocity and constant shear strain

(b) Velocity field across a passive bookshelf fault system

Rotational velocity

v_r : Velocity due to rotation

Translational velocity

v_t : Velocity due to translation

Combined linear velocity and constant shear strain

v_r : Velocity due to translation

v_r : Velocity due to rotation

Zuza and Yin - Figure 7

Figure 7. Velocity fields for the (a) active and (b) passive bookshelf fault systems. Note that the velocity field in the passive system involves the summation of translational (v_t) and rotational (v_r) velocities. See text for discussion.

Zuza and Yin - Figure 8

Figure 8. Kinematic model illustrating the required fault-parallel strain at the western end of one passive bookshelf fault. Given a rigid left-side boundary, left-lateral offset on the fault (d) caused by clockwise fault rotation requires fault-parallel shortening and stretching strain (ϵ_{xx}), for panels A and B respectively, related to approximately one-half the fault slip divided by half the length of each panel, L : $\epsilon_{xx}^A \approx 0.5 d/L$ and $\epsilon_{xx}^B \approx -0.5 d/L$. In this example, the angle of rotation, θ , is very small, such that $\tan\theta$ can be approximated as θ . See text for discussion.

Zuza and Yin - Figure 9

Figure 9. Passive bookshelf faulting in northern Tibet may be driven by (a) boundary-zone rotation to the south, (b) gravitational spreading of thick central Tibet lithosphere with the distribution of gravitational potential energy after England and Molnar (1997), or (c) a toroidal flow in the asthenosphere (e.g., León Soto et al., 2012). See text for further discussion.

Zuza and Yin - Figure 10

Figure 10. Cenozoic tectonic evolution of northern Tibet, showing the timing of major fault initiation where grey structures are already active at given time interval. Note orientation of north arrows. (a) Right-lateral shear, driven from the south, begins in the Eocene and is expressed by north-south shortening in the Qimen Tagh, North Qaidam, and southern Qilian Shan thrust belts. An eastward decrease in north-south shortening magnitude and thrust spacing indicates clockwise rotation of northern Tibet. (b) Progressive crustal thickening causes the initiation of the east-striking left-slip faults along preexisting weaknesses. Continued right-lateral shear drives clockwise rotation of these faults and their wallrock passively against the left-slip Altyn Tagh bounding fault zone. (c) The left-slip faults in this passive bookshelf fault system develop a four-quadrant deformation pattern with domains of fault-parallel stretching and shortening. Age data from Craddock et al. (2011), Yuan et al. (2013), Duvall et al. (2013), and references therein.