Graphical Abstract

To create your abstract, type over the instructions in the template box below. Fonts or abstract dimensions should not be changed or altered.

1, 2, 3-Triazole Amine as Directing Group in Promoting Catalytic Oxidative C-H Olefination under Aerobic Conditions

Leave this area blank for abstract info.

Xiaohan Ye^a, Yanwei Zhang^b, Ying He^a, and Xiaodong Shi^{a, *}

^a Department of Chemistry, University of South Florida, 4202 E. Fowler Avenue, Tampa, Florida 33620, United States

^b C. Eugene Bennett Department of Chemistry, West Virginia University, Morgantown, West Virginia 26506, United States

Tetrahedron

journal homepage: www.elsevier.com

1, 2, 3-Triazole Amine as Directing Group in Promoting Catalytic Oxidative C-H Olefination under Aerobic Conditions

Xiaohan Ye^a, Yanwei Zhang^b, Ying He^a, and Xiaodong Shi^{a,*}^a Department of Chemistry, University of South Florida, 4202 E. Fowler Avenue, Tampa, Florida 33620, United States^b C. Eugene Bennett Department of Chemistry, West Virginia University, Morgantown, West Virginia 26506, United States

ARTICLE INFO

Article history:

Received

Received in revised form

Accepted

Available online

ABSTRACT

A practical *ortho* C-H olefination of phenyl acetic acid derivatives was achieved with a 1,2,3-triazole auxiliary directing group. Good to excellent yields were achieved with O₂ as terminal oxidant. Other bidentate directing groups, such as 8-aminoquinoline, gave poor reactivity under this aerobic condition, highlighting the unique reactivity of triazole in promoting C-H activation.

2009 Elsevier Ltd. All rights reserved.

Keywords:

C-H activation

1, 2, 3-triazole

Oxidative heck

Aerobic olefination

Directing group

* Corresponding author. Tel.: +1-813-974-7249; fax: +1-813-974-3203; e-mail: xmshi@usf.edu

1. Introduction

During the past decade, transition metal catalyzed C-H activation has been widely applied as an efficient approach in complex molecule synthesis.¹ Pd-mediated C-H oxidative olefination, known as the Fujiwara-Moritani oxidative Heck reaction, is one of the most important transformations for arene functionalization.^{2,3} However, this transformation generally suffers from poor selectivity (relying on the different C-H bond reactivity).⁴ The directing group strategy provides an alternative approach to overcome this issue by offering highly selective functionalization at the *ortho* C-H position. So far, several directing groups (DGs) have been reported in promoting the C-H olefination, including OR, COOH and pyridine.⁵ Although excellent selectivity was achieved, most of the reported directing groups suffered one major drawback: the use of stoichiometric amounts of expensive oxidant (such as AgOAc, Selectfluor, etc.) were required. Thus, our interests were triggered to develop a new directing group system, which allows this transformation under aerobic conditions.

Limitations:
 excess amount of metal oxidant is needed

Figure 1. Selective C-H olefination using directing groups

Besides the requirement of strong oxidant, another limitation of directing group strategy was that the DG has to be part of the substrate. Thus, to enhance this powerful directing C-H activation strategy, removable bidentate-auxiliary directing groups have been developed. According to literature, pyridine and quinoline modified protecting groups are the most popular heterocycles in facilitating this type of transformation.⁶ Recently, our group reported the 1,2,3-triazole-4-carboxylic acid as another versatile removable directing group (formation of triazole amide) in promoting selective C-H olefination.⁷ This new triazole directing group not only gives the desired products in excellent yields (up to 95%), but also allows molecular oxygen (1 atm O₂) to be used as terminal oxidant, emphasizing the high efficiency of triazole DG in promoting inactive C-H functionalization (**Figure 2A**).

(A) TA-Promoted Oxidative C-H olefination of amine derivatives

(B) Design for revised TA amine directing group

Figure 2. Triazole group directed C-H activation

2. Results and discussion

In 2014, the Ackermann group introduced a novel TA-amine directing group using easily accessible 1,2,3-triazole for Iron catalyzed C-H activation.⁸ Encouraged by the result that TA amide could successfully promote the C-H olefination under aerobic conditions, we postulate that revised TA-amine may be a

suitable directing group to promote the acid directed C-H olefination under aerobic condition without using any expensive external oxidant (**Figure 2B**).⁹ To verify this idea, several triazole-amine modified substrates were prepared and reacted with alkene **2a** under standard oxidation conditions. The results are summarized in **Figure 3**.

Figure 3. Triazole-amine directed aerobic C-H olefination

First, using previously optimized conditions with triazole acid directing group, 10% Pd(OAc)₂, 10% Cu(OTf)₂, 2.5 eq HOAc in 1,4-dioxane with 1 atm O₂, the desired C-H olefination product **3a** was formed with the triazole-amine directing group, though with much lower yield (45%, 50% conversion of **1a**). Interestingly, increasing the amount of Cu(OTf)₂, which proved to be an effective additive for previous cases, shut down the overall reaction with only trace product **3a** obtained.¹⁰ We postulated that it might be due to a stronger coordination of copper metal towards triazole amine (unlike triazole acid), which inhibit the Pd coordination/C-H activation. In fact, without copper salt as additive, the reaction achieved 95% conversion and 85% yield. Alternatively, tuning the electron properties by switching benzyl group on N1 position to phenyl lead to the complete conversion of **1a** with 95% isolated yield of olefination product **3a**, even at lower temperature (80 °C).¹¹ Notably, triazole C-H olefination was not observed, which highlighted the mild conditions in this aerobic oxidation condition.¹²

To compare and evaluate the reactivity of the triazole amine directing group, substrates containing other common directing groups were prepared and evaluated under the same conditions. As shown in **Figure 3**, N1 phenyl substitution was identified to be the optimal DG with balanced electronic effect in promoting this transformation.¹³ Notably, 8-aminoquinoline and 2-aminophenylthio ether, which have been used in the olefination with strong oxidant, gave poor results under the aerobic oxidation conditions.¹⁴ These results highlighted the unique reactivity of 1,2,3-triazole directing groups in promoting C-H activation.

As shown in **Figure 4**, the TAM-directing group tolerated a large number of substrates, furnishing the corresponding olefination in good to excellent yields. Both EDG (**3a**) and EWG (**3c, d**) modified arenes were suitable for this reaction. Although

substitution occurs primarily at the less hindered *ortho*-carbons position (**3e-3f**) for meta-substituted benzene ring, the di-olefination was also observed. Typically, the selectivity of mono and di-olefination would be poor when both *ortho*-C-Hs were present (**3g-3i**). This problem was overcome through increasing the steric bulkiness at the benzylic position. As shown in **3j**, incorporation of a methyl group effectively improved the mono:di selectivity to >20:1. Meanwhile, using excess amount of alkene (4 equiv.) gave the di-olefination in good yields (**3m-3o**). However, benzoic acid derivatives are proved to be ineffective under this condition as one limitation for this transformation.

General reaction conditions: **1** (0.2 mmol), alkene (0.4 mmol), HOAc (0.5 mmol) and catalyst in 1,4-Dioxane (1.0 mL), and Condition A: 7.5 mol % $Pd(OAc)_2$, O_2 balloon, 80 °C, 12 h; Condition B: 10 mol % $Pd(OAc)_2$, 1 atm O_2 charged in seal tube, 100 °C, 12 h; Condition C: 10 mol % $Pd(OAc)_2$, 0.8 mmol *n*-butyl acrylate, 1 atm O_2 charged 80 °C, 12 h. Isolated yield. ^b the ratio of mono and di substitution was determined by ¹H NMR. ^c Nonactivated alkene, such as 1-hexene gave no reaction under current conditions.

Figure 4. Reaction substrate scope

The scope of alkene was also evaluated. Besides acrylate, most of other activated alkenes, such as, vinyl phosphate (**5d**), vinyl sulfone (**5e**) are suitable for this transformation, giving the desired olefins in excellent yields. Impressively, non-active olefins, such as styrene derivatives, could also undergo this transformation with good to excellent yields (**5f-5g**). Moreover, the coupling with estrone derivatives was also successfully achieved, showing the potential application of this method. However, similar to other reported oxidative Heck-type reactions, α,β substituted alkenes failed to give the desired products due to steric hindrance on either alkene insertion or β -hydride elimination steps. Nevertheless, the broad substrate scope, simple reaction conditions and use of atmosphere oxygen as the oxidant highlighted the advantages of this new triazole-directing group in promoting selective C-H olefination.

3. Conclusion

In conclusion, we have unveiled 1,2,3-triazole as an effective directing group in promoting aerobic C-H oxidative olefination. This study not only revealed a new efficient and economic approach (using 1 atm O_2 as the terminal oxidant) in achieving selective C-H olefination, but also confirmed the versatile reactivity of 1,2,3-triazole in promoting metal catalyzed C-H activation. Other synthetic applications using this new TA-directed C-H activation strategy are currently under investigation.

4. Experimental section

4.1 General experimental methods

All of the reactions dealing with air and/or moisture-sensitive reactions were carried out under an atmosphere of nitrogen using oven/flame-dried glassware and standard syringe/septa techniques. Unless otherwise noted, all commercial reagents and solvents were obtained from the commercial provider and used without further purification. ¹H NMR and ¹³C NMR spectra were recorded on Agilent 400 MHz spectrometers/ Varian 500 MHz spectrometers. Chemical shifts were reported relative to internal tetramethylsilane (δ 0.00 ppm) or $CDCl_3$ (δ 7.26 ppm) for ¹H and $CDCl_3$ (δ 77.00 ppm) for ¹³C. Flash column chromatography was performed on 230-430 mesh silica gel. Analytical thin layer chromatography was performed with precoated glass baked plates (250 μ) and visualized by fluorescence and by charring after treatment with potassium permanganate stain. HRMS were recorded on Agilent 6540 LC/QTOF spectrometer.

4.2. General procedure for Pd catalyzed C-H olefination

Condition A: A 20 mL schlenk tube was charged with the triazole amide (0.2 mmol, 1.0 equiv), alkene (0.4 mmol, 2.0 equiv), $Pd(OAc)_2$ (0.015 mmol, 7.5 mol %) and HOAc (0.5 mmol, 2.5 equiv) in 1 mL anhydrous 1,4-Dioxane. The tube was placed under vacuum and purged with O_2 balloon. The reaction was run at 80 °C for 12 h. After the reaction was completed, the solvent was removed under reduced pressure and the residue was purified by flash chromatography on silica gel to give desired olefination product.

Condition B: A 20 mL seal tube was charged with the triazole amide (0.2 mmol, 1.0 equiv), alkene (0.4 mmol, 2.0 equiv), $Pd(OAc)_2$ (0.02 mmol, 10 mol %), and HOAc (0.5 mmol, 2.5 equiv) in 1 mL anhydrous 1,4-Dioxane. The tube was placed under vacuum and purged with O_2 twice. The reaction was run at 100 °C for 12 h. After the reaction was completed, the solvent was removed under reduced pressure and the residue was purified by flash chromatography on silica gel to give desired olefination product.

Condition C: A 20 mL schlenk tube was charged with the triazole amide (0.2 mmol, 1.0 equiv), alkene (0.8 mmol, 4 equiv), Pd(OAc)₂ (0.015 mmol, 7.5 mol %), and HOAc (0.5 mmol, 2.5 equiv) in 1 mL anhydrous 1,4-Dioxane. The tube was placed under vacuum and purged with O₂ balloon. The reaction was run at 80 °C for 20h. After the reaction was completed (12 h), the solvent was removed under reduced pressure and the residue was purified by flash chromatography on silica gel to give desired olefination product.

4.2.1 (E)-butyl 3-(3-methyl-2-(2-oxo-2-((2-(1-phenyl-1H-1,2,3-triazol-4-yl)propan-2-yl)amino)ethyl)phenyl)acrylate (3a) was prepared following the general procedure A and purified by flash chromatography (Hexane-EtOAc, v/v 2/1). ¹H-NMR (500 MHz; CDCl₃): δ 7.97 (d, *J* = 15.7 Hz, 1H), 7.91 (s, 1H), 7.73-7.71 (m, 2H), 7.51-7.41 (m, 4H), 7.23 (dt, *J* = 14.7, 8.1 Hz, 2H), 6.34 (d, *J* = 15.7 Hz, 1H), 6.07 (s, 1H), 4.17 (t, *J* = 6.7 Hz, 2H), 3.71 (s, 2H), 2.33 (d, *J* = 23.9 Hz, 3H), 1.74 (s, 6H), 1.68-1.62 (m, 2H), 1.40 (dd, *J* = 15.0, 7.5 Hz, 2H), 0.94 (t, *J* = 7.4 Hz, 3H); ¹³C NMR (126 MHz; CDCl₃): δ 169.0, 166.7, 153.4, 142.2, 138.2, 137.0, 134.6, 132.8, 132.3, 129.5, 128.5, 127.7, 125.1, 121.2, 120.48, 120.46, 118.8, 64.4, 51.5, 37.9, 30.7, 27.8, 20.3, 19.1, 13.7; HRMS Calculated for C₂₇H₃₂N₄O₃ [M+H]⁺: 461.2547, Found: 461.2568.

4.2.2 (E)-butyl 3-(3-methoxy-2-(2-oxo-2-((2-(1-phenyl-1H-1,2,3-triazol-4-yl)propan-2-yl)amino)ethyl)phenyl)acrylate (3b) was prepared following the general procedure A and purified by flash chromatography (Hexane-EtOAc, v/v 2/1) (inseparable mixture of the starting material and product with ratio= 3.2:1); ¹H-NMR (500 MHz; CDCl₃): δ 8.03 (d, *J* = 15.8 Hz, 1H), 7.86 (s, 1H), 7.80 (s, 3.2H), 7.70-7.66 (m, 8.4H), 7.50-7.47 (m, 8.4H), 7.41 (m, 4.2H), 7.29-7.21 (m, 9.4H), 6.96-6.90 (m, 7.4H), 6.46 (s, 3.2H), 6.37-6.35 (m, 2H), 4.17 (t, *J* = 6.8 Hz, 2H), 3.86 (s, 12.6H), 3.72 (s, 2H), 3.52 (s, 6.4H), 1.77 (s, 25.2H), 1.69-1.62 (m, 2H), 1.41 (q, *J* = 7.5 Hz, 2H), 0.94 (t, *J* = 7.4 Hz, 3H); ¹³C NMR (126 MHz; CDCl₃): δ 170.8, 169.8, 166.7, 157.5, 157.0, 153.8, 141.8, 137.1, 135.6, 131.2, 129.55, 129.51, 128.67, 128.48, 128.31, 124.0, 123.6, 121.4, 121.0, 120.51, 120.47, 119.4, 118.84, 118.82, 111.8, 110.6, 64.5, 55.8, 55.3, 51.37, 51.17, 40.0, 34.9, 30.7, 27.99, 27.96, 19.1, 13.7; HRMS Calculated for C₂₇H₃₂N₄O₄ [M+H]⁺: 477.2496, Found: 477.2520.

4.2.3 ((E)-butyl 3-(3-fluoro-2-(2-oxo-2-((2-(1-phenyl-1H-1,2,3-triazol-4-yl)propan-2-yl)amino)ethyl)phenyl)acrylate (3c) was prepared following the general procedure A and purified by flash chromatography (Hexane-EtOAc, v/v 2/1); ¹H-NMR (500 MHz; CDCl₃): δ 7.92-7.88 (m, 2H), 7.72-7.70 (m, 2H), 7.51-7.48 (m, 2H), 7.43-7.39 (m, 2H), 7.30-7.26 (m, 1H), 7.13-7.09 (m, 1H), 6.39 (d, *J* = 15.8 Hz, 1H), 6.28 (s, 1H), 4.17 (t, *J* = 6.7 Hz, 2H), 3.73 (d, *J* = 1.5 Hz, 2H), 1.79 (s, 6H), 1.68-1.62 (m, 2H), 1.44-1.36 (m, 2H), 0.94 (t, *J* = 7.4 Hz, 3H); ¹³C NMR (126 MHz; CDCl₃): δ 168.3, 166.4, 160.3, 153.5, 140.51, 140.48, 137.1, 136.51, 136.48, 129.6, 128.84, 128.77, 128.5, 122.69, 122.66, 122.02, 121.93, 121.89, 120.6, 118.8, 116.67, 116.48, 77.3, 77.0, 76.7, 64.6, 51.8, 33.62, 33.58, 30.7, 27.9, 19.1, 13.7; HRMS Calculated for C₂₆H₂₉FN₄O₃ [M+H]⁺: 465.2296, Found: 465.2310.

4.2.4 (E)-butyl 3-(3-nitro-2-(2-oxo-2-((2-(1-phenyl-1H-1,2,3-triazol-4-yl)propan-2-yl)amino)ethyl)phenyl)acrylate (3d) was prepared following the general procedure A and purified by flash chromatography (Hexane-EtOAc, v/v 2/1); ¹H-NMR (500 MHz; CDCl₃): δ 7.99 (d, *J* = 15.8 Hz, 1H), 7.95-7.92 (m, 2H), 7.78-7.73 (m, 3H), 7.52-7.40 (m, 4H), 6.37 (d, *J* = 15.8 Hz, 2H), 4.19 (t, *J* = 6.7 Hz, 2H), 3.91 (s, 2H), 1.83 (s, 6H), 1.69-1.64 (m, 3H), 1.45-1.37 (m, 2H), 0.94 (t, *J* = 7.4 Hz, 3H); ¹³C NMR (126 MHz; CDCl₃): δ 167.3, 166.0, 150.9, 140.6, 138.1, 137.1, 131.7,

129.6, 128.59, 128.51, 128.2, 125.6, 124.1, 120.6, 118.8, 64.8, 52.1, 37.0, 35.8, 30.6, 28.1, 19.1, 13.7; HRMS Calculated for C₂₆H₂₉N₅O₅ [M+H]⁺: 492.2241, Found: 492.2260.

4.2.5 (E)-butyl 3-(4-methyl-2-(2-oxo-2-((2-(1-phenyl-1H-1,2,3-triazol-4-yl)propan-2-yl)amino)ethyl)phenyl)acrylate (3e) was prepared following the general procedure A and purified by flash chromatography (Hexane-EtOAc, v/v 2/1); ¹H-NMR (500 MHz; CDCl₃): δ 7.92 (s, 1H), 7.89 (d, *J* = 15.8 Hz, 1H), 7.71 (d, *J* = 8.0 Hz, 2H), 7.53-7.47 (m, 3H), 7.41 (d, *J* = 7.4 Hz, 1H), 7.12 (d, *J* = 8.0 Hz, 1H), 7.08 (s, 1H), 6.35 (d, *J* = 15.7 Hz, 1H), 6.14 (s, 1H), 4.15 (t, *J* = 6.7 Hz, 2H), 3.64 (s, 2H), 2.35 (s, 3H), 1.76 (s, 6H), 1.64 (quintet, *J* = 7.3 Hz, 2H), 1.40 (sextet, *J* = 7.5 Hz, 2H), 0.94 (t, *J* = 7.4 Hz, 3H); ¹³C NMR (126 MHz; CDCl₃): δ 169.4, 166.9, 153.4, 141.2, 140.8, 137.1, 134.4, 132.1, 130.9, 129.5, 128.8, 128.5, 126.9, 120.5, 119.4, 119.0, 51.6, 42.0, 30.7, 27.9, 21.3, 19.1, 13.7; HRMS Calculated for C₂₇H₃₂N₄O₃ [M+H]⁺: 461.2547, Found: 461.2560.

4.2.6 (E)-butyl 3-(3-methoxy-2-(2-oxo-2-((2-(1-phenyl-1H-1,2,3-triazol-4-yl)propan-2-yl)amino)ethyl)phenyl)acrylate (3e-1) was prepared following the general procedure A and purified by flash chromatography (Hexane-EtOAc, v/v 2/1); ¹H-NMR (500 MHz; CDCl₃): δ 7.95-7.90 (m, 2H), 7.80-7.73 (m, 3H), 7.51-7.48 (m, 3H), 7.41 (t, *J* = 7.4 Hz, 1H), 7.19 (d, *J* = 8.0 Hz, 1H), 6.34 (d, *J* = 15.7 Hz, 1H), 6.15 (s, 1H), 6.04 (d, *J* = 16.3 Hz, 1H), 4.18-4.14 (m, 4H), 3.72 (s, 2H), 2.32 (s, 3H), 1.79 (s, 6H), 1.67-1.62 (m, 4H), 1.41-1.37 (m, 4H), 0.95-0.91 (m, 6H); ¹³C NMR (126 MHz; CDCl₃): δ 168.9, 166.7, 166.1, 153.5, 142.8, 141.8, 138.7, 137.1, 136.6, 132.9, 132.1, 129.8, 129.5, 128.5, 126.8, 126.0, 120.65, 120.59, 118.8, 64.68, 64.48, 51.8, 38.8, 30.69, 30.61, 27.9, 21.1, 19.14, 19.13, 13.7 δ 160.2, 159.7, 140.4, 139.2, 138.9, 134.7, 129.93, 129.87, 129.6, 128.8, 128.45, 128.33, 127.4, 125.9, 121.0, 114.2, 112.1, 55.1, 52.0, 40.2, 36.0; HRMS Calculated for C₃₄H₄₂N₄O₅ [M+H]⁺: 587.3228, Found: 587.3237

4.2.7 (E)-butyl 3-(4-methoxy-2-(2-oxo-2-((2-(1-phenyl-1H-1,2,3-triazol-4-yl)propan-2-yl)amino)ethyl)phenyl)acrylate (3f) was prepared following the general procedure A and purified by flash chromatography (Hexane-EtOAc, v/v 2/1); ¹H-NMR (500 MHz; CDCl₃): δ 7.92 (s, 1H), 7.87-7.84 (m, 1H), 7.73-7.71 (m, 2H), 7.60 (d, *J* = 8.7 Hz, 1H), 7.51-7.48 (m, 2H), 7.41 (t, *J* = 7.4 Hz, 1H), 6.86 (dd, *J* = 8.7, 2.6 Hz, 1H), 6.80 (d, *J* = 2.6 Hz, 1H), 6.30 (d, *J* = 15.7 Hz, 1H), 6.08 (s, 1H), 4.15 (t, *J* = 6.7 Hz, 2H), 3.83 (s, 3H), 3.67 (s, 2H), 1.77 (s, 6H), 1.67-1.61 (m, 3H), 1.40 (sextet, *J* = 7.5 Hz, 2H), 0.94 (t, *J* = 7.4 Hz, 3H); ¹³C NMR (126 MHz; CDCl₃): δ 169.1, 167.1, 161.2, 153.4, 140.8, 137.1, 136.4, 129.5, 128.50, 128.48, 126.2, 120.50, 120.42, 118.9, 117.9, 116.3, 113.9, 64.3, 55.3, 51.6, 42.2, 30.7, 27.9, 19.1, 13.7; HRMS Calculated for C₂₇H₃₂N₄O₄ [M+H]⁺: 477.2496, Found: 477.2509.

4.2.8 (2E,2'E)-dibutyl 3,3'-(4-methoxy-2-(2-oxo-2-((2-(1-phenyl-1H-1,2,3-triazol-4-yl)propan-2-yl)amino)ethyl)-1,3-phenylene)diacrylate (3f-1) was prepared following the general procedure A and purified by flash chromatography (Hexane-EtOAc, v/v 2/1); ¹H-NMR (500 MHz; CDCl₃): δ 7.97 (s, 1H), 7.91 (d, *J* = 15.7 Hz, 1H), 7.80 (d, *J* = 16.2 Hz, 1H), 7.75-7.73 (m, 2H), 7.60 (d, *J* = 8.8 Hz, 1H), 7.51-7.47 (m, 2H), 7.42-7.39 (m, 1H), 6.90 (d, *J* = 8.8 Hz, 1H), 6.42 (d, *J* = 16.2 Hz, 1H), 6.28 (d, *J* = 15.6 Hz, 1H), 6.17 (s, 1H), 4.15 (td, *J* = 6.8, 2.4 Hz, 4H), 3.89 (s, 3H), 3.80 (s, 2H), 1.79 (s, 6H), 1.67-1.60 (m, 4H), 1.41-1.36 (m, 4H), 0.95-0.91 (m, 6H); ¹³C NMR (126 MHz; CDCl₃): δ 168.6, 167.01, 166.83, 159.6, 153.5, 141.6, 138.4, 137.2, 134.7, 129.5, 129.1, 128.5, 127.8, 124.95, 124.86, 120.6, 119.5, 118.9, 110.2, 64.49, 64.42, 55.7, 52.0, 38.4, 30.71, 30.67, 27.9, 19.14, 19.14, 13.7; HRMS Calculated for C₃₄H₄₂N₄O₆ [M+H]⁺: 603.3177, Found: 603.3187.

4.2.9 (E)-butyl 3-(5-methyl-2-(2-oxo-2-((2-(1-phenyl-1H-1,2,3-triazol-4-yl)propan-2-yl)amino)ethyl)phenyl)acrylate (3g) was prepared following the general procedure A and purified by flash chromatography (Hexane-EtOAc, v/v 2/1); ¹H-NMR (500 MHz; CDCl₃): δ 7.90-7.87 (m, 2H), 7.73-7.70 (m, 2H), 7.51-7.48 (m, 2H), 7.43-7.41 (m, 2H), 7.18-7.16 (m, 3H), 6.38 (d, *J* = 15.7 Hz, 1H), 6.02 (s, 1H), 4.15 (t, *J* = 6.7 Hz, 2H), 3.64 (s, 2H), 2.36 (s, 3H), 1.75 (s, 6H), 1.68 (m, *J* = 19.4 Hz, 2H), 1.40 (ddd, *J* = 7.9, 7.1, 6.9 Hz, 2H), 0.94 (t, *J* = 7.4 Hz, 3H); ¹³C NMR (126 MHz; CDCl₃): δ 169.5, 166.8, 153.4, 141.4, 137.7, 137.1, 133.7, 131.47, 131.34, 131.32, 129.63, 129.60, 129.56, 129.2, 128.5, 127.6, 120.57, 120.53, 118.9, 64.5, 51.6, 44.2, 41.7, 30.7, 27.9, 21.1, 19.1, 13.7; HRMS Calculated for C₂₇H₃₂N₄O₃ [M+H]⁺: 461.2547, Found: 461.2561.

4.2.10 (E)-butyl 3-(5-methoxy-2-(2-oxo-2-((2-(1-phenyl-1H-1,2,3-triazol-4-yl)propan-2-yl)amino)ethyl)phenyl)acrylate (3h) was prepared following the general procedure A and purified by flash chromatography (Hexane-EtOAc, v/v 2/1); ¹H-NMR (500 MHz; CDCl₃): δ 7.91-7.85 (m, 2H), 7.73-7.70 (m, 2H), 7.52-7.47 (m, 2H), 7.42 (dt, *J* = 7.5, 1.6 Hz, 1H), 7.19 (d, *J* = 8.4 Hz, 1H), 7.12 (d, *J* = 2.7 Hz, 1H), 6.92 (dd, *J* = 8.4, 2.7 Hz, 1H), 6.36 (d, *J* = 15.7 Hz, 1H), 6.07 (d, *J* = 0.8 Hz, 1H), 4.16 (t, *J* = 6.7 Hz, 3H), 3.83 (s, 4H), 3.61 (s, 2H), 1.75 (s, 6H), 1.70-1.61 (m, 2H), 1.40 (td, *J* = 8.0, 7.0 Hz, 2H), 0.94 (t, *J* = 7.4 Hz, 3H); ¹³C NMR (126 MHz; CDCl₃): δ 169.7, 166.6, 159.0, 153.4, 141.3, 137.1, 134.9, 132.5, 129.5, 128.5, 126.7, 120.72, 120.54, 120.45, 118.9, 116.4, 111.9, 64.5, 55.4, 51.5, 41.2, 30.7, 27.9, 19.1, 13.7; HRMS Calculated for C₂₇H₃₂N₄O₄ [M+H]⁺: 477.2496, Found: 477.2509.

4.2.11 (E)-butyl 3-(5-fluoro-2-(2-oxo-2-((2-(1-phenyl-1H-1,2,3-triazol-4-yl)propan-2-yl)amino)ethyl)phenyl)acrylate (3i) was prepared following the general procedure A and purified by flash chromatography (Hexane-EtOAc, v/v 2/1); ¹H-NMR (500 MHz; CDCl₃): δ 7.91 (s, 1H), 7.84 (dd, *J* = 15.8, 1.4 Hz, 1H), 7.73-7.71 (m, 2H), 7.52-7.49 (m, 2H), 7.44-7.40 (m, 1H), 7.30 (d, *J* = 2.7 Hz, 1H), 7.23 (s, 1H), 7.06 (td, *J* = 8.2, 2.7 Hz, 1H), 6.36 (d, *J* = 15.7 Hz, 1H), 6.18 (s, 1H), 4.17 (t, *J* = 6.7 Hz, 2H), 3.64 (s, 2H), 1.78 (s, 6H), 1.68-1.62 (m, 2H), 1.44-1.37 (m, 2H), 0.94 (t, *J* = 7.4 Hz, 3H); ¹³C NMR (126 MHz; CDCl₃): δ 169.1, 166.4, 153.4, 140.3, 137.1, 135.8, 133.0, 130.3, 129.6, 128.6, 121.7, 120.6, 118.9, 117.32, 117.15, 113.60, 113.42, 64.7, 51.7, 41.2, 30.7, 27.9, 19.1, 13.7; HRMS Calculated for C₂₆H₂₉FN₄O₃ [M+H]⁺: 465.2296, Found: 465.2307.

4.2.12 (E)-butyl 3-(2-(1-oxo-1-((2-(1-phenyl-1H-1,2,3-triazol-4-yl)propan-2-yl)amino)propan-2-yl)phenyl)acrylate (3j) was prepared following the general procedure A and purified by flash chromatography (Hexane-EtOAc, v/v 2/1); ¹H-NMR (500 MHz; CDCl₃): δ 8.02 (d, *J* = 15.7 Hz, 1H), 7.85 (s, 1H), 7.72-7.68 (m, 2H), 7.57 (d, *J* = 7.7 Hz, 1H), 7.52-7.48 (m, 2H), 7.43-7.37 (m, 3H), 7.30 (ddd, *J* = 10.8, 6.8, 3.5 Hz, 1H), 6.37 (d, *J* = 15.7 Hz, 1H), 5.99 (s, 1H), 4.19 (t, *J* = 6.7 Hz, 2H), 3.91 (q, *J* = 7.1 Hz, 1H), 1.75 (d, *J* = 3.0 Hz, 6H), 1.71-1.65 (m, 2H), 1.47 (d, *J* = 7.1 Hz, 4H), 1.45-1.40 (m, 2H), 0.96 (t, *J* = 7.4 Hz, 3H); ¹³C NMR (126 MHz; CDCl₃): δ 172.7, 166.7, 153.5, 141.4, 140.6, 137.1, 133.2, 130.5, 129.56, 129.54, 128.9, 128.5, 127.64, 127.50, 127.47, 127.2, 121.2, 120.49, 120.46, 118.7, 64.6, 51.4, 43.5, 30.7, 27.90, 27.78, 19.1, 18.4, 13.7; HRMS Calculated for C₂₇H₃₂N₄O₃ [M+H]⁺: 461.2547, Found: 461.2560.

4.2.13 (E)-butyl 3-(1-(2-oxo-2-((2-(1-phenyl-1H-1,2,3-triazol-4-yl)propan-2-yl)amino)ethyl)naphthalen-2-yl)acrylate (3k) was prepared following the general procedure A and purified by flash chromatography (Hexane-EtOAc, v/v 2/1); ¹H-NMR (500 MHz; CDCl₃): δ 8.22 (d, *J* = 15.7 Hz, 1H), 8.08 (d, *J* = 8.1 Hz, 1H), 7.86-7.81 (m, 2H), 7.77 (s, 1H), 7.70 (d, *J* = 8.7 Hz, 1H), 7.66-

7.64 (m, 2H), 7.55 (quintet, *J* = 7.4, 1.5 Hz, 2H), 7.50-7.47 (m, 2H), 7.42-7.39 (m, 1H), 6.52 (d, *J* = 15.7 Hz, 1H), 5.99 (s, 1H), 4.21 (dd, *J* = 12.1, 5.3 Hz, 4H), 1.72-1.66 (m, 8H), 1.43 (dq, *J* = 15.0, 7.5 Hz, 2H), 0.96 (d, *J* = 14.8 Hz, 3H); ¹³C NMR (126 MHz; CDCl₃): δ 169.2, 166.8, 153.4, 141.6, 137.0, 134.3, 132.6, 131.6, 131.2, 129.5, 128.73, 128.64, 128.49, 127.4, 127.0, 124.6, 123.8, 121.8, 120.5, 118.7, 64.6, 51.7, 37.3, 30.7, 27.9, 19.2, 13.7; HRMS Calculated for C₃₀H₃₂N₄O₃ [M+H]⁺: 497.2547, Found: 497.2561.

4.2.14 (2E,2'E)-dibutyl 3,3'-(5-methyl-2-(2-oxo-2-((2-(1-phenyl-1H-1,2,3-triazol-4-yl)propan-2-yl)amino)ethyl)-1,3-phenylene)diacrylate (4a) was prepared following the general procedure C and purified by flash chromatography (Hexane-EtOAc, v/v 4/1); ¹H-NMR (500 MHz; CDCl₃): δ 7.94 (d, *J* = 15.6 Hz, 3H), 7.72 (d, *J* = 7.7 Hz, 2H), 7.50-7.47 (m, 2H), 7.41-7.39 (m, 3H), 6.34 (d, *J* = 15.7 Hz, 2H), 6.17 (s, 1H), 4.16 (t, *J* = 6.7 Hz, 4H), 3.78 (s, 2H), 2.37 (s, 3H), 1.77 (s, 6H), 1.68-1.62 (m, 4H), 1.40 (dd, *J* = 15.0, 7.5 Hz, 4H), 0.93 (t, *J* = 7.4 Hz, 6H); ¹³C NMR (126 MHz; CDCl₃): δ 168.5, 166.5, 153.4, 141.8, 137.6, 137.1, 135.5, 130.4, 129.50, 129.45, 128.5, 121.8, 120.6, 118.8, 64.5, 51.8, 37.2, 35.3, 30.7, 27.9, 21.0, 19.1, 13.7; HRMS Calculated for C₃₄H₄₂N₄O₅ [M+H]⁺: 587.3228, Found: 587.3241.

4.2.15 (2E,2'E)-dibutyl 3,3'-(5-methoxy-2-(2-oxo-2-((2-(1-phenyl-1H-1,2,3-triazol-4-yl)propan-2-yl)amino)ethyl)-1,3-phenylene)diacrylate (4b) was prepared following the general procedure C and purified by flash chromatography (Hexane-EtOAc, v/v 4/1); ¹H-NMR (500 MHz; CDCl₃): δ 7.95-7.92 (m, 3H), 7.73-7.71 (m, 2H), 7.50-7.47 (m, 2H), 7.42-7.39 (m, 1H), 7.12 (s, 2H), 6.33 (dd, *J* = 15.7, 1.0 Hz, 2H), 6.24 (s, 1H), 4.17 (t, *J* = 6.7 Hz, 4H), 3.84 (d, *J* = 1.0 Hz, 3H), 3.74 (s, 2H), 1.77 (s, 6H), 1.65 (qd, *J* = 7.7, 5.9 Hz, 4H), 1.40 (q, *J* = 7.5 Hz, 4H), 0.93 (td, *J* = 7.4, 1.0 Hz, 6H); ¹³C NMR (126 MHz; CDCl₃): δ 168.7, 166.4, 158.6, 153.4, 141.7, 137.1, 136.8, 129.5, 128.4, 125.8, 122.1, 120.5, 118.8, 114.0, 64.6, 55.4, 51.8, 36.9, 30.6, 27.8, 19.1, 13.6; HRMS Calculated for C₃₄H₄₂N₄O₆ [M+H]⁺: 603.3177, Found: 603.3189.

4.2.16 (2E,2'E)-dibutyl 3,3'-(5-fluoro-2-(2-oxo-2-((2-(1-phenyl-1H-1,2,3-triazol-4-yl)propan-2-yl)amino)ethyl)-1,3-phenylene)diacrylate (4c) was prepared following the general procedure C and purified by flash chromatography (Hexane-EtOAc, v/v 4/1); ¹H-NMR (500 MHz; CDCl₃): δ 7.93-7.89 (m, 3H), 7.73-7.71 (m, 2H), 7.50 (dd, *J* = 8.4, 7.2 Hz, 2H), 7.43-7.41 (m, 1H), 7.29-7.27 (m, 2H), 6.33 (d, *J* = 15.7 Hz, 3H), 4.18 (t, *J* = 6.7 Hz, 4H), 3.76 (s, 2H), 1.79 (s, 6H), 1.68-1.62 (m, 4H), 1.42-1.38 (m, 4H), 0.94 (t, *J* = 7.4 Hz, 6H); ¹³C NMR (126 MHz; CDCl₃): δ 168.1, 166.2, 160.7, 153.4, 140.71, 140.69, 137.81, 137.75, 137.1, 129.6, 129.36, 129.33, 128.6, 123.1, 120.6, 118.8, 115.26, 115.08, 115.06, 64.7, 51.9, 36.9, 30.6, 27.8, 19.1, 13.7; HRMS Calculated for C₃₃H₃₉FN₄O₅ [M+H]⁺: 591.2977, Found: 591.2992.

4.2.17 (E)-methyl 3-(2-(2-oxo-2-((2-(1-phenyl-1H-1,2,3-triazol-4-yl)propan-2-yl)amino)ethyl)phenyl) (5a) was prepared following the general procedure B and purified by flash chromatography (Hexane-EtOAc, v/v 2/1); ¹H-NMR (500 MHz; CDCl₃): δ 7.98 (d, *J* = 15.8 Hz, 1H), 7.91 (s, 1H), 7.73-7.70 (m, 2H), 7.51-7.48 (m, 2H), 7.45-7.39 (m, 2H), 7.27-7.21 (m, 2H), 6.34 (d, *J* = 15.7 Hz, 1H), 6.07 (s, 1H), 3.75 (s, 3H), 3.71 (s, 2H), 2.34 (s, 3H), 1.75 (s, 7H); ¹³C NMR (126 MHz; CDCl₃): δ 169.0, 167.0, 153.4, 142.5, 138.2, 137.0, 134.6, 132.8, 132.3, 129.5, 128.5, 127.7, 125.1, 120.68, 120.49, 118.8, 51.7, 51.5, 37.9, 27.8, 20.3; HRMS Calculated for C₂₄H₂₆N₄O₃ [M+H]⁺: 419.2078, Found: 419.2098.

4.2.18 (*E*)-benzyl 3-(3-methyl-2-(2-oxo-2-((2-(1-phenyl-1*H*-1,2,3-triazol-4-yl)propan-2-yl)amino)ethyl)phenyl)acrylate (**5b**) was prepared following the general procedure A and purified by flash chromatography (Hexane-EtOAc, v/v 2/1); ¹H-NMR (500 MHz; CDCl₃): δ 8.03 (d, *J* = 15.7 Hz, 1H), 7.89 (s, 1H), 7.72-7.70 (m, 2H), 7.50-7.47 (m, 2H), 7.44-7.31 (m, 7H), 7.22 (dd, *J* = 15.7, 8.1 Hz, 2H), 6.39 (d, *J* = 15.7 Hz, 1H), 6.01 (s, 1H), 5.21 (s, 2H), 3.71 (s, 2H), 2.34 (s, 3H), 1.72 (s, 6H); ¹³C NMR (126 MHz; CDCl₃): δ 168.9, 166.4, 142.9, 138.3, 137.1, 135.9, 134.5, 132.9, 132.4, 129.6, 128.54, 128.52, 128.21, 128.20, 127.7, 125.1, 120.7, 120.5, 118.8, 66.3, 51.5, 38.0, 27.9, 20.4; HRMS Calculated for C₃₀H₃₀N₄O₃ [M+H]⁺: 495.2391, Found: 495.2408.

4.2.19 (*E*)-tert-butyl 3-(3-methyl-2-(2-oxo-2-((2-(1-phenyl-1*H*-1,2,3-triazol-4-yl)propan-2-yl)amino)ethyl)phenyl)acrylate (**5c**) was prepared following the general procedure A and purified by flash chromatography (Hexane-EtOAc, v/v 2/1); ¹H-NMR (500 MHz; CDCl₃): δ 7.89 (d, *J* = 15.7 Hz, 2H), 7.74-7.72 (m, 2H), 7.51-7.48 (m, 2H), 7.44-7.41 (m, 2H), 7.26-7.21 (m, 3H), 6.27 (d, *J* = 15.7 Hz, 1H), 5.98 (s, 1H), 3.70 (s, 2H), 2.33 (s, 3H), 1.74 (s, 7H), 1.50 (s, 9H); ¹³C NMR (126 MHz; CDCl₃): δ 169.0, 165.9, 153.4, 141.3, 138.2, 137.1, 134.9, 132.6, 132.1, 129.6, 128.5, 127.7, 125.2, 123.1, 120.5, 118.8, 80.6, 51.5, 38.1, 28.1, 27.9, 20.4; HRMS Calculated for C₂₇H₃₂N₄O₃ [M+H]⁺: 461.2547, Found: 461.2576.

4.2.20 (*E*)-diethyl 3-methyl-2-(2-oxo-2-((2-(1-phenyl-1*H*-1,2,3-triazol-4-yl)propan-2-yl)amino)ethyl)styrylphosphonate (**5d**) was prepared following the general procedure B and purified by flash chromatography (Hexane-EtOAc, v/v 1/2); ¹H-NMR (500 MHz; CDCl₃): δ 7.94 (s, 1H), 7.85-7.79 (m, 1H), 7.75-7.71 (m, 2H), 7.51-7.47 (m, 2H), 7.42-7.38 (m, 2H), 7.25-7.20 (m, 2H), 6.26-6.15 (m, 2H), 4.15-4.07 (m, 5H), 3.70 (s, 2H), 2.33 (s, 3H), 1.76 (s, 6H), 1.34-1.30 (m, 6H); ¹³C NMR (126 MHz; CDCl₃): δ 168.9, 153.5, 146.59, 146.58, 146.54, 146.53, 138.2, 137.0, 135.28, 135.11, 132.59, 132.57, 132.2, 129.5, 128.4, 127.5, 124.8, 120.4, 118.8, 118.0, 116.5, 61.88, 61.84, 51.4, 37.6, 27.8, 20.3, 16.33, 16.28; HRMS Calculated for C₂₆H₃₃N₄O₄P [M+H]⁺: 497.2312, Found: 497.2328.

4.2.21 (*E*)-2-(2-methyl-6-(2-(phenylsulfonyl)vinyl)phenyl)-*N*-(2-(1-phenyl-1*H*-1,2,3-triazol-4-yl)propan-2-yl)acetamide (**5e**) was prepared following the general procedure B and purified by flash chromatography (Hexane-EtOAc, v/v 2/1); ¹H-NMR (500 MHz; CDCl₃): δ 8.00 (d, *J* = 15.2 Hz, 1H), 7.93 (dt, *J* = 6.7, 1.7 Hz, 3H), 7.73-7.71 (m, 1H), 7.60-7.56 (m, 1H), 7.52-7.47 (m, 4H), 7.42-7.39 (m, 1H), 7.32 (d, *J* = 7.7 Hz, 1H), 7.26 (t, *J* = 3.7 Hz, 1H), 7.18 (t, *J* = 7.7 Hz, 1H), 6.79 (d, *J* = 15.1 Hz, 1H), 6.20 (s, 1H), 3.72 (s, 2H), 2.34 (s, 3H), 1.77 (s, 7H). ¹³C NMR (126 MHz; CDCl₃): δ 168.5, 153.5, 140.56, 140.41, 138.6, 137.0, 133.48, 133.34, 133.1, 132.5, 129.67, 129.59, 129.3, 128.6, 127.70, 127.66, 125.4, 120.5, 118.7, 51.7, 37.8, 27.9, 20.4. HRMS Calculated for C₂₈H₂₈N₄O₃S [M+H]⁺: 501.1955, Found: 501.1966.

4.2.22 (*E*)-2-(2-(4-chlorostyryl)-6-methylphenyl)-*N*-(2-(1-phenyl-1*H*-1,2,3-triazol-4-yl)propan-2-yl)acetamide (**5f**) was prepared following the general procedure B and purified by flash chromatography (Hexane-EtOAc, v/v 2/1); ¹H-NMR (500 MHz; CDCl₃): δ 7.84 (s, 1H), 7.64-7.62 (m, 2H), 7.49-7.45 (m, 3H), 7.43-7.38 (m, 3H), 7.31-7.18 (m, 6H), 6.89 (d, *J* = 16.0 Hz, 1H), 5.94 (s, 1H), 3.71 (s, 2H), 2.37 (s, 3H), 1.69 (s, 6H); ¹³C NMR (126 MHz; CDCl₃): δ 169.7, 153.4, 137.59, 137.53, 137.0, 135.6, 133.4, 131.3, 130.7, 130.3, 129.6, 128.8, 128.5, 127.8, 126.9, 124.7, 120.4, 118.7, 51.3, 38.6, 27.9, 20.4; HRMS Calculated for C₂₈H₂₇ClN₄O [M+H]⁺: 471.1946, Found: 471.1958.

4.2.23 (*E*)-2-(2-(4-(tert-butyl)styryl)-6-methylphenyl)-*N*-(2-(1-phenyl-1*H*-1,2,3-triazol-4-yl)propan-2-yl)acetamide (**5g**) was prepared following the general procedure B and purified by flash chromatography (Hexane-EtOAc, v/v 2/1); ¹H-NMR (500 MHz; CDCl₃): δ 7.86 (s, 1H), 7.66 (d, *J* = 7.5 Hz, 2H), 7.50-7.39 (m, 6H), 7.34-7.23 (m, 4H), 7.17 (d, *J* = 7.4 Hz, 1H), 6.96 (d, *J* = 16.0 Hz, 1H), 5.95 (s, 1H), 3.72 (s, 2H), 2.36 (s, 3H), 1.70 (s, 6H), 1.31 (s, 9H); ¹³C NMR (126 MHz; CDCl₃): δ 169.9, 153.4, 151.0, 137.9, 137.6, 137.1, 134.4, 131.9, 131.1, 130.0, 129.6, 128.5, 127.7, 126.4, 125.62, 125.47, 124.7, 120.4, 118.8, 51.3, 38.6, 34.6, 31.2, 27.9, 20.4; HRMS Calculated for C₃₂H₃₆N₄O [M+H]⁺: 493.2962, Found: 493.2977.

4.2.24 (*E*)-(8*R*,9*S*,13*S*)-13-methyl-17-oxo-7,8,9,11,12,13,14,15,16,17-decahydro-6*H*-cyclopenta[*a*]phenanthren-3-yl 3-(3-methyl-2-(2-oxo-2-((2-(1-phenyl-1*H*-1,2,3-triazol-4-yl)propan-2-yl)amino)ethyl)phenyl)acrylate (**5h**) was prepared following the general procedure A and purified by flash chromatography (Hexane-EtOAc, v/v 2/1); ¹H-NMR (500 MHz; CDCl₃): δ 8.14 (d, *J* = 15.7 Hz, 1H), 7.90 (s, 1H), 7.69-7.68 (m, 2H), 7.53-7.51 (m, 1H), 7.47-7.43 (m, 2H), 7.40-7.37 (m, 1H), 7.30-7.25 (m, 3H), 6.91 (dd, *J* = 8.4, 2.5 Hz, 1H), 6.87 (d, *J* = 2.4 Hz, 1H), 6.53 (d, *J* = 15.7 Hz, 1H), 6.11 (s, 1H), 3.74 (s, 2H), 2.90 (dd, *J* = 9.9, 5.0 Hz, 2H), 2.52 (t, *J* = 9.6 Hz, 1H), 2.43-2.39 (m, 1H), 2.36 (s, 3H), 2.29 (td, *J* = 10.7, 4.2 Hz, 1H), 2.16 (dd, *J* = 18.6, 9.4 Hz, 1H), 2.09-1.96 (m, 4H), 1.76 (s, 6H), 1.68-1.42 (m, 7H), 0.92 (s, 3H); ¹³C NMR (126 MHz; CDCl₃): δ 220.8, 168.9, 165.3, 153.4, 148.6, 144.1, 138.4, 137.9, 137.3, 137.0, 134.4, 133.1, 132.7, 129.5, 128.5, 127.8, 126.3, 125.3, 121.6, 120.5, 120.2, 118.77, 118.76, 60.4, 51.5, 50.4, 47.9, 44.1, 37.98, 37.95, 35.8, 31.5, 29.4, 27.9, 26.3, 25.7, 21.6, 21.0, 20.4, 14.2, 13.8; HRMS Calculated for C₄₁H₄₄N₄O₄ [M+H]⁺: 657.3435, Found: 657.3453.

Acknowledgments

We thank the NSF (CHE-1362057), NSFC (21228204), and University of South Florida for financial support.

References and notes

Supplementary Material

Supplementary material that may be helpful in the review process should be prepared and provided as a separate electronic file. That file can then be transformed into PDF format and submitted along with the manuscript and graphic files to the appropriate editorial office.

- (a) Labinger, J. A.; Bercaw, J. E. *Nature* **2002**, *417*, 507-514.; (b) Godula, K.; Sames, D. *Science* **2006**, *312*, 67-72.; (c) Alberico, D.; Scott, M. E.; Lautens, M. *Chem. Rev.* **2007**, *107*, 174-238.; (d) Seregin, I. V.; Gevorgyan, V. *Chem. Soc. Rev.* **2007**, *36*, 1173-1193.; (e) Giri, R.; Shi, B.-F.; Engle, K. M.; Mangel, N.; Yu, J.-Q. *Chem. Soc. Rev.* **2009**, *38*, 3242-3272.; (f) Colby, D. A.; Bergman, R. G.; Ellman, J. A. *Chem. Rev.* **2010**, *110*, 624-655.; (g) Cho, S. H.; Kim, J. Y.; Kwak, J.; Chang, S. *Chem. Soc. Rev.* **2011**, *40*, 5068-5083.; (h) Arockiam, P. B.; Bruneau, C.; Dixneuf, P. H. *Chem. Rev.* **2012**, *112*, 5879-5918.; (i) Kuhl, N.; Hopkinson, M. N.; Wencel-Delord, J.; Glorius, F. *Angew. Chem. Int. Ed.* **2012**, *51*, 10236-10254.; (j) Yeung, C. S.; Dong, V. M. *Chem. Rev.* **2011**, *111*, 1215.; (k) Song, G.; Wang, F.; Li, X. *Chem. Soc. Rev.* **2012**, *41*, 3651.; (l) Li, B.-J.; Shi, Z.-J. *Chem. Soc. Rev.* **2012**, *41*, 5588-5598.

- ² Moritani, I.; Fujiwara, Y. *Tetrahedron Lett.* **1967**, 1119-1122.
- ³ Le Bras, J.; Muzart, J. *Chem. Rev.* **2011**, *111*, 1170-1214
- ⁴ For recent examples of transition metal catalyzed non-directed C-H olefination of arene, see: (a) Zhang, Y.-H.; Shi, B.-F.; Yu, J.-Q. *J. Am. Chem. Soc.* **2009**, *131*, 5072-5074.; (b) Kubota, A.; Emmert, M. H.; Sanford, M. S. *Org. Lett.* **2012**, *14*, 1760-1763.; (c) Ying, C.-H.; Yan, S.-B.; Duan, W.-L. *Org. Lett.* **2014**, *16*, 500-503.; (d) Vora, H. U.; Silvestri, A. P.; Engelin, C. J.; Yu, J.-Q. *Angew. Chem. Int. Ed.* **2014**, *53*, 2683-2686.
- ⁵ For recent examples of Pd catalyzed directed oxidative C-H olefination of arene, see: (a) Engle, K. M.; Wang, D.-H.; Yu, J.-Q. *J. Am. Chem. Soc.* **2010**, *132*, 14137-14151.; (b) Lu, Y.; Wang, D.-H.; Engle, K. M.; Yu, J.-Q. *J. Am. Chem. Soc.* **2010**, *132*, 5916-5921.; (c) Miyasaka, M.; Hirano, K.; Satoh, T.; Miura, M. *J. Org. Chem.* **2010**, *75*, 5421-5424.; (d) Garcia-Rubia, A.; Urones, B.; Gomez Arrayas, R.; Carretero, J. C. *Angew. Chem. Int. Ed.* **2011**, *50*, 10927-10931.; (e) Huang, C.; Chattopadhyay, B.; Gevorgyan, V. *J. Am. Chem. Soc.* **2011**, *133*, 12406-12409.; (f) Wang, C.; Ge, H. *Chem. Eur. J.* **2011**, *17*, 14371-14374.; (g) Yu, M.; Liang, Z.; Wang, Y.; Zhang, Y. *J. Org. Chem.* **2011**, *76*, 4987-4994.; (h) Yu, M.; Xie, Y.; Xie, C.; Zhang, Y. *Org. Lett.* **2012**, *14*, 2164-2167.; (i) Chan, L. Y.; Kim, S.; Ryu, T.; Lee, P. H. *Chem. Commun.* **2013**, *49*, 4682-4684.; (j) Li, G.; Leow, D.; Wan, L.; Yu, J.-Q. *Angew. Chem. Int. Ed.* **2013**, *52*, 1245-1247.; (k) Hu, J.; Guan, M.; Han, J.; Huang, Z.-B.; Shi, D.-Q.; Zhao, Y. *J. Org. Chem.* **2015**, *80*, 7896-7904.
- ⁶ For reviews, see: (a) Rouquet, G.; Chatani, N. *Angew. Chem. Int. Ed.* **2013**, *52*, 11726-11743.; (b) Zhang, B.; Guan, H.; Liu, B.; Shi, B. *Chinese J. Org. Chem.* **2014**, *34*, 1487-1498.; (c) Castro, L. C. M.; Chatani, N. *Chem. Lett.* **2015**, *44*, 410-421.; (d) Daugulis, O.; Roane, J.; Tran, L. D. *Acc. Chem. Res.* **2015**, *48*, 1053-1064. (e) Yadav, M. R.; Rit, R. K.; Shankar, M.; Sahoo, A. K. *Asian J. Org. Chem.* **2015**, *4*, 846-864.
- ⁷ (a) Ye, X.; He, Z. R.; Ahmed, T.; Weise, K.; Akhmedov, N. G.; Petersen, J. L.; Shi, X. D. *Chem. Sci.* **2013**, *4*, 3712.; (b) Ye, X.; Shi, X. *Org. Lett.* **2014**, *16*, 4448-4451.; (c) Ye, X.; Petersen, J. L.; Shi, X. *Chem. Commun.* **2015**, *51*, 7863 - 7866.
- ⁸ For examples of triazole amine directed C-H activation, see: (a) Gu, Q.; Al Mamari, H. H.; Graczyk, K.; Diers, E.; Ackermann, L. *Angew. Chem. Int. Ed.* **2014**, *53*, 3868-3871.; (b) Al Mamari, H. H.; Diers, E.; Ackermann, L. *Chem. Eur. J.* **2014**, *20*, 9739-9743.; (c) Graczyk, K.; Haven, T.; Ackermann, L. *Chem. Eur. J.* **2015**, *21*, 8812-8815.; (d) Zhang, G.; Xie, X.; Zhu, J.; Li, S.; Ding, C.; Ding, P. *Org. Biomol. Chem.* **2015**, *13*, 5444-5449.
- ⁹ For examples of Pd-catalyzed C-H olefination using O₂ as terminal oxidant, see: (a) Beck, E. M.; Grimster, N. P.; Hatley, R.; Gaunt, M. J. *J. Am. Chem. Soc.* **2006**, *128*, 2528-2529.; (b) Cong, X.; You, J.; Gao, G.; Lan, J. *Chem. Commun.* **2013**, *49*, 662-664.; (c) Liu, B.; Jiang, H.-Z.; Shi, B.-F. *J. Org. Chem.* **2014**, *79*, 1521-1526.
- ¹⁰ (a) Werner, E. W.; Sigman, M. S. *J. Am. Chem. Soc.* **2010**, *132*, 13981.; (b) Mei, T.-S.; Werner, E. W.; Burckle, A. J.; Sigman, M. S. *J. Am. Chem. Soc.* **2013**, *135*, 6830.
- ¹¹ The reaction under air atmosphere resulted a lower conversion (70%) and quicker decomposition of Pd catalyst by the formation of Pd black.
- ¹² For examples of C-H functionalization on C-4 of 1,2,3-triazole, see: (a) Chuprakov, S.; Chernyak, N.; Dudnik, A. S.; Gevorgyan, V. *Org. Lett.* **2007**, *9*, 2333.; (b) Iwasaki, M.; Yorimitsu, H.; Oshima, K. *Chem. Asian. J.* **2007**, *2*, 1430.; (c) Ackermann, L.; Vicente, R.; Born, R. *Adv. Synth. Catal.* **2008**, *350*, 741. ; (d) Jiang, H.; Feng, Z.; Wang, A.; Liu, X.; Chen, Z. *Eur. J. Org. Chem.* **2010**, *2010*, 1227.; (e) Liu, W.; Li, Y.; Xu, B.; Kuang, C. *Org. Lett.* **2013**, *15*, 2342.
- ¹³ Wang, K.; Chen, M.; Wang, Q.; Shi, X.; Lee, J. K. *J. Org. Chem.* **2013**, *78*, 7249-7258.
- ¹⁴ Deb, A.; Bag, S.; Kancherla, R.; Maiti, D. *J. Am. Chem. Soc.* **2014**, *136*, 13602-13605.