Graphical Abstract

Stereospecific synthesis of *cis*-stilbenes from benzaldehydes and phenylacetic acids via sequential Perkin condensation and decarboxylation

Leave this area blank for abstract info.

Clayton M. Thompson, Matthew D. Orellana, Shannon E. Lloyd and Weiming Wu*

Department of Chemistry and Biochemistry, San Francisco State University, San Francisco, CA 94132 USA

Stereospecific synthesis of *cis*-stilbenes from benzaldehydes and phenylacetic acids via sequential Perkin condensation and decarboxylation

Clayton M. Thompson, Matthew D. Orellana, Shannon E. Lloyd and Weiming Wu*

Department of Chemistry and Biochemistry, San Francisco State University, San Francisco, CA 94132 USA

ARTICLE INFO

Article history:

Received

Received in revised form

Accepted

Available online

Keywords:

cis-stilbene

α -aryl cinnamic acid

stereospecific

decarboxylation

Perkin condensation

ABSTRACT

cis-Stilbenes were synthesized stereospecifically in high yield from the cuprous oxide-catalyzed decarboxylation of α -aryl-*trans*-cinnamic acids that were prepared from the Perkin condensation of benzaldehydes and phenylacetic acids.

2009 Elsevier Ltd. All rights reserved.

1. Introduction

Stilbene derivatives have many applications in industry as dyes, optical brighteners, phosphors, and scintillators.¹ Stilbene moieties are also common in natural products such as resveratrol and pterostilbene.^{2,3} The *trans*- and *cis*-stilbenes are also used for the study of the transmission of substituent effects from one benzene ring to another through a double bond.⁴⁻⁶

The synthesis of stilbene has been extensively studied and discussed.⁷ The common methods include the Wittig reaction and related reactions⁸⁻¹⁰, the reduction of benzoin^{11,12}, the reaction of benzaldehydes with benzylmagnesium halides followed by dehydration¹³, the Meerwein reaction of cinnamic acids with aromatic diazonium salt¹⁴⁻¹⁷, and the decarboxylation of α -aryl cinnamic acids from the Perkin condensation of benzaldehydes with phenylacetic acids.¹⁸⁻²⁰ These methods do not usually give exclusively one isomer.

The decarboxylation of α -aryl cinnamic acids have recently been revisited as a method to prepare resveratrol derivatives.²¹⁻²³ In this *Letter*, we describe an improved method for the stereospecific synthesis of *cis*-stilbene derivatives.

2. Results and Discussion

The Perkin condensation between benzaldehydes and phenylacetic acids to form α -aryl cinnamic acids (Scheme 1) have been found to be moderate in yield and stereospecific in some cases^{18,20} but not others.²¹⁻²³ In this work, the yields were

improved by adding triethylamine slowly and by designing a simpler purification procedure.²⁴ The reaction was complete in about five hours, but overnight reaction did not seem to hurt the yield. A dilute HCl solution was added to the reaction mixture to precipitate the product α -aryl cinnamic acids **1**. Upon filtration and drying in the same Buchner funnel, the solid was washed with a 9:1 mixture of hexanes and ethyl acetate to remove the trace amount of unreacted aldehydes. The solids thus isolated were found to be pure as judged by NMR spectroscopy. Only the α -aryl substituted *trans*-cinnamic acids **1** were produced. The stereochemistry of **1** was deduced from that of the product stilbene **2**.

Scheme 1

* Corresponding author. Tel.: +0-415-338-1436; fax: +0-415-338-2384; e-mail: wuw@sfsu.edu.

Table 1. Synthesis of *cis*-stilbenes **2** from benzaldehydes and phenylacetic acids through *E*- α -aryl-*trans*-cinnamic acids **1**

Entries	X=	Y=	Yield (1 , isolated, %)	Yield (2 , isolated, %)
a	4-chloro	H	80	59
b	4-chloro	4-methoxy	92	82
c	3-cyano	H	100	85
d	3-cyano	4-fluoro	99	85
e	3-cyano	4-chloro	86	66
f	3-cyano	4-bromo	99	76
g	3-cyano	4-iodo	86	87
h	3-cyano	4-methoxy	93	95
i	2-cyano	4-bromo	99	65
j	3-methoxy	4-methoxy	80	69
k	3-methoxy	4-bromo	99	55
l	3,5-dimethoxy	4-methoxy	94	76

^aThe reaction^bReaction produced

The yields were generally excellent for various substituents as seen in Table 1. Benzaldehydes and phenylacetic acids with strong electronwithdrawing *p*-substituents such as cyano and nitro groups were not examined because they were reported to form stilbene product under Perkin condensation conditions.^{7,17} The 3-cyano derivatives were used in many cases due to our interest in the property of the stilbene-3-carboxylic acids in connection with another project.

The reported decarboxylation of α -aryl-*trans*-cinnamic acids **1** to stilbenes **2** was usually carried out in quinoline using copper or copper chromite catalysts at high temperature (~260 °C).^{7,19,22} At these temperatures, the decarboxylation product could be transformed into the more stable *trans*-isomer. Thus, the formation of a mixture of *cis*- and *trans*-stilbenes had been observed.^{7,22}

A somewhat milder condition for the decarboxylation would preserve the *cis*-configuration. It has been reported that cuprous oxide is an efficient catalyst for the decarboxylation of benzoic acids.²⁵ Different amines were thus tested as base with cuprous oxide as catalyst for the efficient decarboxylation of α -aryl-*trans*-cinnamic acids. Tributylamine turned out to be the best since it has a boiling point of slightly higher than the reaction temperature and is easily removed by washing with dilute acid. N-Methyl-2-pyrrolidone (NMP) was found to be an excellent solvent for the reaction.

The reaction condition was thus applied to various α -aryl-*trans*-cinnamic acids. Good to excellent yields were obtained regardless of the properties of the substituents as seen in Table 1. The stereochemistry of the stilbenes was established by the chemical shifts and the characteristic vicinal ¹H-¹H-coupling constants of the stilbene double bond protons. The two protons

of the *cis*-isomers have lower chemical shifts and smaller coupling constants (about 12 Hz) than the *trans*-isomers. All isolated stilbenes were found to be *cis*-isomers and were judged to be pure by thin-layer chromatography (TLC) and NMR. The *cis*-stilbene products can be conveniently converted to the *trans*-isomer if desired.^{7,22} The described method can be useful in the synthesis of derivatives of resveratrol, as evident from the synthesis of resveratrol precursor **2l**. It should also be pointed out that this synthetic method is flexible in terms of substitution patterns of benzaldehydes and phenylacetic acids starting material. In *cis*-stilbene products, substituents X and Y may stem from either benzaldehydes or phenylacetic acids, depending on the availability of these starting materials.

3. Conclusion

In summary, *cis*-stilbenes were synthesized stereospecifically from the decarboxylation of α -aryl-*trans*-cinnamic acids, which was formed from the Perkin condensation of benzaldehydes and phenylacetic acids. The synthetic method allows the mix-and-match of different substituents on the two phenyl groups. Overall, the two-step sequence represents a convenient, stereospecific, flexible and high-yielding method for the synthesis of symmetrically and especially unsymmetrically substituted *cis*-stilbenes.

Acknowledgments

This investigation was supported by the SFSU Office of Research and Sponsored Programs (W.W.) and the Department of Chemistry and Biochemistry Summer Research Fellowship (C.M.T.). The NMR facility was funded by the National Science Foundation (DUE-9451624 and DBI 0521342). We also thank

Professor Ihsan Erden for helpful discussions and ACS Project SEED student Junyi Liu for technical assistance.

References and Notes

1. Likhtenshtein, G. I. *Stilbenes Synthesis and Applications* In *Kirk-Othmer Encyclopedia of Chemical Technology*, Wiley: 2012; DOI: 10.1002/0471238961.stillkh.a01.
2. Shen, T.; Wang, X. N.; Lou, H. X. *Nat. Prod. Rep.* **2009**, *26*, 916-935.
3. Bavaresco, L.; Fregoni, C.; Cantù, E.; Trevisan, M. *Drugs Exp. Clin. Res.* **1999**, *25*, 57-63.
4. Beale, R. N.; Roe, E. M. F. *J. Am. Chem. Soc.* **1952**, *74*, 2302-2307.
5. Riezebos, G.; Havinga, E. *Rec. Trav. Chim.* **1961**, *80*, 446-467.
6. Jaffe, H. H.; Orchin, M. *J. Chem. Soc.* **1960**, 1078-1087.
7. Wheeler, O. H.; Battle de Pabon, H. N. *J. Org. Chem.* **1965**, *30*, 1473-1477.
8. Wittig, G.; Haag, W. *Chem. Ber.* **1955**, *88*, 1654-1666.
9. Horner, L.; Hoffmann, H.; Wippel, H. G.; Klahre, G. *Chem. Ber.* **1959**, *92*, 2499-2505.
10. Wadsworth, W. S.; Emmons, W. D. *J. Am. Chem. Soc.* **1961**, *83*, 1733-1738.
11. Ballard, D. A.; Dehn, W. M. *J. Am. Chem. Soc.* **1932**, *64*, 3969-3971.
12. Shriner, R. L.; Burger, A. *Org. Syn. Coll. Vol. III*, 786-788.
13. Dodds, E. C.; Goldberg, L.; Lawson, W.; Robinson, R. *Proc. Royal Soc* **1939**, *B127*, 140-167.
14. Meerwein, H.; Buchner, E.; Van Emster, K. *J. Prakt. Chem.* **1939**, *152*, 237-266.
15. Bergmann, F.; Schapiro, D. *J. Org. Chem.* **1947**, *12*, 57-66.
16. L'Ecuier, P.; Turcotte, F.; Giguere, J.; Olivier, C. A.; Roberge, P. *Can. J. Res.* **1948**, *26B*, 70-80.
17. Bell, F.; Waring, D. H. *J. Chem. Soc.* **1948**, 1024-1026.
18. Buckles, R. E.; Bremer, K. *Org. Syn. Coll. Vol. IV*, 777-779.
19. Buckles, R. E.; Wheeler, N. G. *Org. Syn. Coll. Vol. IV*, 857-859.
20. Buckles, R. E.; Cooper, J. A. *J. Org. Chem.* **1965**, *30*, 1588-1592.
21. Kumar, V.; Sharma, A.; Sharma, A.; Sinha, A. K. *Tetrahedron* **2007**, *63*, 7640-7646.
22. Solladie, G.; Pasturel-Jacope, Y.; Maignan, J. *Tetrahedron* **2003**, *59*, 3315-3321.
23. Sinha, A. K.; Kumar, V.; Sharma, A.; Sharma, A.; Kumar, R. *Tetrahedron* **2007**, *63*, 11070-11077.
24. *Experimental details*: All reagents were obtained from commercial sources and used without further purification. Typical experimental procedures are described below using the syntheses of α -phenyl-3-cyano-*trans*-cinnamic acid (**1c**) and 3-cyano-*cis*-stilbene (**2c**) as examples. In some cases, the crude cinnamic acids appeared as oil and liquid-liquid extraction was necessary to separate the product. Upon removal of solvent, the crude product was washed with 9:1 hexane/ethyl acetate solution to yield the products as solids.
 α -Phenyl-3-cyano-*trans*-cinnamic acid (**1c**): Phenylacetic acid (274 mg, 2 mmol) and 3-cyanobenzaldehyde (264 mg, 2 mmol) were mixed with acetic anhydride (300 mg, 3 mmol) in a round-bottom flask. The mixture was heated at 65 °C for 15 minutes before triethylamine (400 mg, 4 mmol) was added dropwise. The reaction mixture was heated overnight at 65 °C. Upon cooling to room temperature, the reaction mixture was poured into a dilute aqueous HCl solution. The solid precipitated was suction-filtered and washed with water, suction-dried on the Buchner funnel under vacuum for 1 hour, and washed with a hexane/ethyl acetate solution (9:1) to remove trace starting materials. An off-white solid (501 mg, 100%) was obtained and was found to be pure by NMR.
3-Cyano-*cis*-stilbene (**2c**): The acid **1c** (495 mg, 2 mmol) was mixed with cuprous oxide (15 mg, 5 mol%), tributylamine (185 mg, 1 mmol) in 3 mL of N-methyl-2-pyrrolidone (NMP). The reaction mixture was heated at 200 °C for 90 minutes. Upon cooling to room temperature, the reaction was partitioned between a dilute HCl solution in brine and ethyl acetate. The organic layer was washed with water three times, dried over anhydrous sodium sulfate, and concentrated to give a crude product. The product was further purified by column chromatography to give the product as a clear oil (346 mg, 85%).
25. Cahiez, G.; Moyeux, A.; Gager, O.; Poizat, M. *Adv. Synth. Catal.* **2013**, *355*, 790-796.