Stereoselective β -Mannosylation via Anomeric *O*-Alkylation: Formal Synthesis of Potent Calcium Signal Modulator Acremomannolipin A

Xiaohua Li,* Nader Berry, Kevin Saybolt, Uddin Ahmed, Yue Yuan

Department of Natural Sciences, University of Michigan-Dearborn, 4901 Evergreen Road, Dearborn, Michigan 48128, United States.

ARTICLE INFO

Article history:

Received
 Received in revised form
 Accepted
 Available online

Keywords:

β -Mannosylation
 Anomeric *O*-Alkylation
 Synthesis
 Acremomannolipin A

ABSTRACT

Stereoselective β -mannosylation has been investigated via cesium carbonate-mediated anomeric *O*-alkylation of D-mannose-derived lactol with various electrophiles. It was found that electrophiles bearing trifluoromethanesulfonate (triflate) as the leaving group are most reactive. In addition, a highly efficient formal synthesis of potent calcium signal modulator acremomannolipin A has been achieved using this β -mannosylation method.

2013 Elsevier Ltd. All rights reserved.

The glycolipid acremomannolipin A (**1**) was isolated from a filamentous fungus *Acromonium strictum*.¹ Structurally, acremomannolipin A contains a D-mannopyranoside β -linked to a D-mannitol and all the hydroxyls in the mannose are acylated with saturated aliphatic acids (**Figure 1**). Therefore, the D-mannose moiety is made hydrophobic, whereas the D-mannitol portion is hydrophilic. The structure of acremomannolipin A was elucidated on the basis of intensive spectroscopic analyses as well as its degradation studies. Biologically, acremomannolipin A showed the interesting activity at 200 nM enabling calcineurin deletion mutant cells to grow in the presence of Cl^- , which would be caused by calcium signal modulating.¹ As a potential calcium signal modulator, acremomannolipin A is considered an attractive target for biologists as well as synthetic chemists as acremomannolipin A and its synthetic analogs may be of significance for therapeutic and biotechnological purposes.

The structural features of acremomannolipin A have posed significant difficulties for the total synthesis, mainly due to the presence of a β -mannoside which is known to be one of the most synthetically challenging glycosidic linkages.² Previously, acremomannolipin A (**1**) was first synthesized by Muraoka and co-workers in 2013 (**Scheme 1**).³ Based on Crich β -mannosylation protocol,⁴ key intermediate β -mannoside **4** was obtained from 4,6-*O*-benzylidene-protected D-mannose donor **2** and acceptor **3** in 71% yield ($\beta/\alpha = 30/1$). Later, Muraoka and co-workers also prepared 1'-*epi*-

acremomannolipin A, the α -anomer of acremomannolipin A, which showed reduced activity.⁵ In 2015, the same group also prepared five homologs of acremomannolipin A bearing alditols of different length and found that the length of the alditol side chain was a crucial determinant for the potent calcium signal modulating activity.⁶ Early 2015, Li and co-workers reported another total synthesis of acremomannolipin A in which key intermediate β -mannoside **7** was obtained via gold(I)-catalyzed glycosylation⁷ between 4,6-*O*-benzylidene-protected D-mannose-derived *ortho*-alkynylbenzoate donor **5** and acceptor **6** in 85% yield ($\beta/\alpha = 13/1$).⁸ In 2016, the Toshima group described the third total synthesis of acremomannolipin A in which key intermediate β -mannoside **10** was obtained via borinic acid-catalyzed glycosylation between 1,2-anhydromannose donor **8** and acceptor **9** in 99% yield (β only).⁹

Figure 1. The structure of acremomannolipin A (**1**).

* Corresponding author. Tel.: +1-313-593-5154; fax: +1-313-593-4937; e-mail: shannli@umich.edu.

Scheme 1. Previous synthesis of acremomannolipin A and our strategy.

Early in 2016 we disclosed a new method for stereoselective construction of β -mannosides via cesium carbonate-mediated anomeric *O*-alkylation of D-mannose-derived lactols.¹⁰ In this Communication, we would like to report our efforts in the synthesis of acremomannolipin A in which the key intermediate β -mannoside **10** was prepared from known D-mannose-derived lactol **11**¹¹ and D-mannitol-derived primary triflate **12** via cesium carbonate-mediated anomeric *O*-alkylation.

In our previous report,¹⁰ only sugar-derived primary and secondary alkyl triflates, e.g. **14**, were studied as electrophiles for cesium carbonate-mediated anomeric *O*-alkylation with D-mannose-derived lactols. For example, when C6-primary triflate **14** was employed, β -D-mannoside **16** was obtained in 93% yield (β only, entry 1).¹⁰ We wondered if other primary electrophiles bearing different leaving groups other than triflates would also react with lactol **11** in this type of β -mannosylation. As shown in **Table 1**, it was found that only the use of excess 1-iodopentane and Cs_2CO_3 afforded desired β -D-mannoside **17** in 21% yield (β only, entry 2), while there was no detectable product when 1-bromopentane, *n*-pentyl mesylate or tosylate was employed (entries 3–5). Elevation of the reaction temperature to 50 °C or use of other solvents did not improve the reaction outcome. Use of methyl iodide gave methyl β -D-mannoside **18** in 71% yield (β only, entry 6). Over-methylation at O2 could be seen if the reaction was allowed to proceed longer. When activated alkyl halides, such as allyl bromide and benzyl bromide, were used, corresponding desired β -D-mannoside **19** and **20** were produced in 94% and 64% yields, respectively (β only, entries 7 and 8). These studies indicated that non-activated primary alkyl triflates are required to react with mannose-derived lactols in the presence of cesium carbonate for efficient synthesis of β -mannosides.

Table 1. Studies of stereoselective β -mannosylation via anomeric *O*-alkylation involving various electrophiles.^a

			15a X = I 15b X = Br 15c X = OMs 15d X = OTs
entry	electrophile/condition	product, yield, ^b α/β ratio	
1 ¹⁰	14 / condition A	 16 93%, β only	
2	15a / condition B	 17 21%, β only	
3	15b / condition B	17 0%	
4	15c / condition B	17 0%	
5	15d / condition B	17 0%	
6	CH₃I / condition B	 18 71%, β only	
7	allylBr / condition B	 19 94%, β only	
8	BnBr / condition B	 20 64%, β only	

^a Condition A: **11** (1.0 eq.), **14** (1.5 eq.), Cs₂CO₃ (2.0 eq.), ClCH₂CH₂Cl, 40 °C, 24 h; Condition B: **11** (1.0 eq.), electrophile (2.5 eq.), Cs₂CO₃ (3.0 eq.), ClCH₂CH₂Cl, 40 °C, 24 h. ^b Isolated yield.

Next, we aimed at the synthesis of acremomannolipin A. Starting from D-mannitol, the triflate acceptor **12** can be prepared in six steps by adopting the known procedures.^{9,12} As shown in **Scheme 2**, conversion of commercially available D-mannitol **21** into its corresponding tri-acetonide (88%) followed by regioselective deprotection of one of the terminal acetonides afforded diol **22** (91%).¹² Regioselective benzylation¹¹ of the primary alcohol of **22** (73%) followed by silylation⁹ of the secondary alcohol gave rise to **23** (85%). Next, removal of the benzyl ether of **23** via palladium-catalyzed hydrogenolysis furnished the known primary alcohol **24** in 83% yield.⁹ This primary alcohol **24** was then subjected to standard triflation (triflic anhydride, pyridine, dichloromethane, 0 °C) to afford desired triflate **12** in 96% yield.¹³ Under our recently developed optimal β -mannosylation condition,¹⁰ known D-mannose-derived lactol **11**¹¹ reacted with triflate acceptor **12** in dichloroethane in the presence of cesium carbonate at 40 °C for 24 hours afforded the desired key β -mannoside **10** in 87% yield (β only).¹⁴ The R_f, ¹H and ¹³C NMR, optical rotation, and HRMS data of our synthesized β -mannoside **10** were found to be identical to those reported in the literature.⁹ For the synthesis of acremomannolipin A (**1**), β -mannoside **10** would just need to undergo standard acylations and deprotections.⁹ Thus, our efforts constitute a highly efficient formal synthesis of acremomannolipin A (**1**).

Scheme 2. Synthesis of acremomannolipin A.

In conclusion, stereoselective β -mannosylation has been studied via cesium carbonate-mediated anomeric *O*-alkylation of D-mannose-derived lactol with various electrophiles. It was found that electrophiles bearing triflate as the leaving group

are most reactive and efficient for this type of β -mannosylation. In addition, a highly efficient formal synthesis of potent calcium signal modulator acremomannolipin A has been achieved using this β -mannosylation method.

Acknowledgments

We are grateful to National Science Foundation (CHE-1464787) and University of Michigan-Dearborn for supporting this research. We would also like to thank Professor Jianglong Zhu (Department of Chemistry and Biochemistry, University of Toledo) for helpful discussions.

References and notes

- 1 Sugiura, R.; Kita, A.; Tsutsui, N.; Muraoka, O.; Hagihara, K.; Umeda, N.; Kunoh, T.; Takada, H.; Hirose, D. *Bioorg. Med. Chem. Lett.* **2012**, *22*, 6735–6739.
- 2 (a) Nigudkar, Swati S.; Demchenko, Alexei V. *Chem. Sci.* **2015**, *6*, 2687–2704. (b) Ishiwata, A.; Lee, Y. J.; Ito, Y. *Org. Biomol. Chem.* **2010**, *8*, 3596–3608. (c) Barresi, F.; Hindsgaul, O. *Modern Methods in Carbohydrate Synthesis*; Khan, S. H.; O'Neill, R. A. Eds.; Harwood Academic Publishers: Amsterdam, **1996**; 251–276. (d) Toshima, K.; Tatsuta, K. *Chem. Rev.* **1993**, *93*, 1503–1531. (e) Paulsen, H. *Angew. Chem., Int. Ed.* **1982**, *21*, 155–173.
- 3 Tsutsui, N.; Tanabe, G.; Kita, A.; Sugiura, R.; Muraoka, O. *Tetrahedron Lett.* **2013**, *54*, 451–453.
- 4 (a) Crich, D.; Sun, S. *J. Org. Chem.* **1996**, *61*, 4506–4507. (b) Crich, D.; Sun, S. *Tetrahedron* **1998**, *54*, 8321–8348. (c) Crich, D.; Sun, S. *J. Am. Chem. Soc.* **1997**, *119*, 11217–11223.
- 5 Tsutsui, N.; Tanabe, G.; Gotoh, A.; Kita, A.; Sugiura, R.; Muraoka, O. *Tetrahedron* **2013**, *69*, 9917–9930.
- 6 Tsutsui, N.; Tanabe, G.; Morita, N.; Okayama, Y.; Kita, A.; Sugiura, R.; Muraoka, O. *Bioorg. Med. Chem.* **2015**, *23*, 3761–3773.
- 7 (a) Li, Y.; Yang, Y.; Yu, B. *Tetrahedron Lett.* **2008**, *49*, 3604–3608. (b) Yu, B.; Sun, J.; Yang, X. *Acc. Chem. Res.* **2012**, *45*, 1227–1236.
- 8 Sun, P.; Wang, P.; Zhang, Y.; Zhang, X.; Wang, C.; Liu, S.; Lu, J.; Li, M. *J. Org. Chem.* **2015**, *80*, 4164–4175.
- 9 Tanaka, M.; Nashida, J.; Takahashi, D.; Toshima, K. *Org. Lett.* **2016**, *18*, 2288–2291.
- 10 Nguyen, H.; Zhu, D.; Li, X.; Zhu, J. *Angew. Chem., Int. Ed.* **2016**, *55*, 4767–4771.
- 11 Jonke, S.; Liu, K.-G.; Schmidt, R. R. *Chem. Eur. J.* **2006**, *12*, 1274–1290.
- 12 Vidyasagar, A.; Sureshan, K. M. *Eur. J. Org. Chem.* **2014**, 2349–2356.

- 13 Preparation of triflate **12**: To known alcohol **24**⁹ (50.0 mg, 0.13 mmol) in 1.0 mL anhydrous CH₂Cl₂ cooled at 0 °C was added anhydrous pyridine (0.39 mmol, 0.03 mL) followed by the addition of triflic anhydride (0.2 mmol, 0.03 mL). The mixture was stirred at 0 °C for 30 minutes before being quenched with water. The crude reaction mixture was diluted with CH₂Cl₂ and washed sequentially with saturated CuSO₄ aqueous solution, water, and brine. The organic solution was dried over sodium sulfate, filtered, and concentrated in vacuo to afford 64 mg of the desired triflate **12** (96% yield) which was sufficiently pure for the next step. ¹H NMR (400 MHz, CDCl₃) δ 4.65-4.62 (dd, *J*=3.7, 10.3 Hz, 1H), 4.56-4.51 (dd, *J*=7.0, 10.3 Hz, 1H), 4.26-4.23 (m, 1H), 4.19- 4.15 (dd, *J*=6.2, 8.4 Hz, 1H), 4.09- 4.04 (m, 1H), 4.02-3.99 (dd, *J*=3.2, 7.0 Hz, 1H), 3.97- 3.93 (dd, *J*=5.9, 8.4 Hz, 1H), 3.92- 3.89 (*J*=7.2, 15.0 Hz, 1H), 1.43 (s, 3H), 1.42 (s, 3H), 1.37 (s, 3H), 1.36 (s, 3H), 0.94 (s, 9H), 0.16 (s, 3H), 0.15 (s, 3H). ¹³C NMR (100 MHz, CDCl₃) δ 112.3, 109.9, 109.0, 82.7, 81.0, 80.3, 73.2, 73.1, 70.2, 67.7, 66.8, 26.8, 26.4, 25.9, 25.7, 25.2, 24.7, 24.5.
- 14 Synthesis of key β-mannoside **10**: To a mixture of known D-mannose-derived lactol **11**¹¹ (45 mg, 0.1 mmol), sugar-derived triflate **12** (76 mg, 0.15 mmol), and cesium carbonate (65 mg, 0.2 mmol) was added 1,2-dichloroethane (1.0 mL). The reaction mixture was stirred at 40°C for 24 hours under Argon. The crude reaction mixture was diluted with 1 mL dichloromethane and directly purified by preparative thin layer chromatography (hexanes: EtOAc = 1:1) to furnish 70 mg of β-mannoside **7** (87% yield). The R_f, ¹H and ¹³C NMR, optical rotation, and HRMS data of our synthetic β-mannoside **10** were found to be identical to those reported in the literature.⁹