1 **Temporal variations of disinfection byproduct precursors in wildfire detritus**

2

3 Jun-Jian Wang¹, Randy A. Dahlgren², Mahmut S. Erşan³, Tanju Karanfil³, Alex T.
4 Chow^{1,3,*}

5

6 ¹Baruch Institute of Coastal Ecology & Forest Science, Clemson University, Georgetown,
7 South Carolina, 29442, USA;

8 ²Department of Land, Air and Water Resources, University of California – Davis,
9 California, 95616, USA;

10 ³Department of Environmental Engineering and Earth Sciences, Clemson University,
11 Anderson, SC 29625, USA;

12 *Correspondence to: A. T. Chow (achow@clemson.edu), P.O. BOX 596, Georgetown,
13 SC, 29442, USA. Tel.: +1 843 546 1013x232; fax: +1 843 546 6296.

Abstract

The Rim Fire ignited on August 17, 2013 and became the third largest wildfire in California history. The fire consumed 104,131 ha of forested watersheds that were the drinking water source for 2.6 million residents in the San Francisco Bay area. To understand temporal variations in dissolved organic matter (DOM) after the wildfire and its potential impacts on disinfection byproduct (DBP) formation in source water supply, we collected the 0-5 cm ash/soil layer with surface deposits of white ash (high burn severity) and black ash (moderate burn severity) within the Rim Fire perimeter in Oct 2013 (pre-rainfall) for five sequential extractions, and in Dec 2013 (~87 mm cumulative precipitation) and Aug 2014 (~617 mm cumulative precipitation) for a single water extraction. Water-extractable DOM was characterized by absorption and fluorescence spectroscopy and DBP formation tests. Both increasing cumulative precipitation in the field or number of extractions in the lab resulted in a significant decrease in specific conductivity, dissolved organic carbon, and DBP formation potential, but an increase in DOM aromaticity (reflected by specific UV absorbance). However, the lab sequential leaching failed to capture the increase of the $\text{NO}_x^- \text{-N} / \text{NH}_4^+ \text{-N}$ ratio and the decrease in pH and dissolved organic carbon / nitrogen ratio of ash/soil extracts from Oct 2013 to Aug 2014. Increasing cumulative precipitation, inferring an increase in leaching after fire, led to an increase in DOM reactivity to form trihalomethanes, haloacetic acids, and chloral hydrate, but not for haloketones, haloacetonitrile, or *N*-nitrosodimethylamine, which were more related to the original burn severity. This study highlights that fire-affected DBP precursors for different DBP species have distinct temporal variation possibly due to their various sensitivity to biogeochemical alterations.

- 37 **Keywords:** dissolved organic matter; forest fires, water quality; source water treatment;
- 38 trihalomethanes

1. Introduction

Wildfire is one of the strongest disturbances to forests in western North America and is likely to have more frequent and severe impacts on forest health and water supply as a result of global climate change and increasing human activities (Jolly et al. 2015, van Mantgem et al. 2013, Westerling et al. 2006). As more than 120 million people depend on water supplied from national forests in the USA (United States Forest Service 2007), understanding how wildfires affect drinking water quality is critical for effective management of drinking water supply (Bladon et al. 2014). Studies across a spectrum of geographical locations have collectively recognized that wildfires have the potential to cause water quality deterioration in terms of increased loads of sediment, nutrients, metals, and persistent organic pollutants (Bladon et al. 2014, Bladon et al. 2008, Smith et al. 2011). As demonstrated by previous post-fire water quality studies of the 2002 Hayman Fire (55,893 ha), Colorado (Rhoades et al. 2011) and 2003 Lost Creek Wildfire (21,000 ha) in Alberta, Canada (Emelko et al. 2011), wildfires are likely to have persistent effects on water quality (e.g., sediments and nutrients) ranging from a few years to more than a decade.

Compared to these relatively known water quality effects, less information is available on how wildfires can potentially affect dissolved organic matter (DOM) quantity and chemistry and disinfection byproduct (DBP) precursors in burned forest watersheds (Bladon et al. 2014). DBPs are a group of potentially harmful chemicals formed during drinking water treatment where the oxidants (e.g., chlorine, chloramine, and ozone) react with DBP precursors like natural organic matter and inorganic constituents in source waters (Karanfil et al. 2008, Xie 2004). Trihalomethanes (THMs)

and haloacetic acids (HAAs) are the carbonaceous DBPs regulated by the US-EPA with maximum contamination levels of 80 µg/L and 60 µg/L, respectively (Richardson et al. 2007). The highly toxic nitrogenous DBPs such as haloacetonitriles (HANs) and *N*-nitrosodimethylamine (NDMA) (Plewa and Wagner 2009), although unregulated, are often detected in chlorinated and chloraminated drinking waters, respectively (Lee et al. 2007, Shah and Mitch 2012). Previous studies have clearly shown that different DOM chemical compositions have a wide range of reactivity with oxidants to form DBPs (Dotson et al. 2009, Lee and Hur 2014, Wang et al. 2012). Wildfires cause major transformations in DOM quantity and chemical composition depending on fire temperature and oxygen availability (Bladon et al. 2014, Santin et al. 2015, Wang et al. 2015a,b). Thus, understanding the load and reactivity of DOM following wildfires is critical for assessing the potential effects of wildfire on DBP formation in water supplies.

Following wildfires, a significant fraction of terrestrial DOM can enter surface waters via surface runoff and soil leaching contributing to increased DBP precursor concentrations in water supplies (Chow et al. 2009, Revchuk and Suffet 2014). In a study of the 2013 Rim Fire in California (104,131 ha), Wang et al. (2015a) found distinct changes in the chemistry of water extractable DOM with burn severity being an important variable affecting the chemical characteristics of the DOM and its reactivity to form DBPs. With higher burn severity, the reactivity of DOM from surface ash/soil extracts to form THMs and HAAs was decreased; however, the formation of the more toxic HANs and NDMA was increased. After the High Park Fire in Colorado (35,323 ha), Writer et al. (2014) studied the temporal variation of stream water quality and the associated DBP formation potential in burned and unburned sites using monthly

sampling. They found that the first storm event caused high concentrations of dissolved organic carbon (DOC; up to 13.7 mg/L) and DBP precursors (>250 µg/L THMs; >400 µg/L HAAs; 32.1 µg/L HANs) in burned watersheds. The reactivity of fire-affected DOM showed higher reactivity to form HANs, but comparable reactivity to form THMs and HAAs relative to the unburned site. Revchuk and Suffet (2014) found that the average DOC in source water was elevated by the 2007 Zaca Fire in California (97,208 ha) from 2.8 mg/L pre-fire to ~4.8 mg/L one year post-fire (~70% increase), and the THM formation potential was increased to ~200-290 µg/L (~49% increase).

Despite the information gained from these previous studies, there remain significant knowledge gaps hindering our understanding of the chemical, hydrological, and biological mechanisms regulating the “black box” with wildfire as input and drinking water DBPs as output. A mechanistic understanding calls for systematic studies on variations in temporal terrestrial and aquatic DBP precursors and the effects of site-specific meteorology, topography, ecosystem types, and hydrological conditions. Revchuk and Suffet (2014) proposed that precipitation leaching and biological consumption were the primary factors affecting DOC dynamics from wildfire ash. However, it remains unquantified how the leaching process affects the fire-affected terrestrial precursors for different types of DBPs. There are several important questions yet to be answered, such as how do terrestrial DOM chemistry and chlorine reactivity change with increasing time and increasing leaching after fire alteration of the forest floor detritus layer, i.e., the surface ash/soil (AS) layer? Do temporal variations in DOM leaching depend on sites and burn severity? Do carbonaceous and nitrogenous DBP precursors change synchronously with increasing time after fire?

To explore these questions, we investigated the temporal variation in characteristics and chlorine reactivity of water extractable DOM in surface deposits with white ash (AS-W; indicator of high burn severity) and black ash (AS-B; indicator of moderate burn severity) from conifer and chaparral vegetation within the 2013 Rim Fire perimeter in California. This study used two approaches to evaluate the variation of DBP precursors with increased leaching in AS extracts: 1) temporal sampling of AS samples that were exposed to natural precipitation, biological activity, and other environmental factors for a one-year period; and 2) a laboratory sequential leaching of the AS (AS-W vs AS-B; conifer vs chaparral) collected before any precipitation events to examine whether the lab dissolution/leaching process was feasible for simulating the temporal variations in the field.

2. Materials and methods

2.1. Site information and ash/soil sampling

The Rim Fire was the largest wildfire on record in the Sierra Nevada Mountains and the third largest wildfire in California history. The effects of the Rim Fire on water quality raised considerable public concern as it occurred in watersheds supporting the drinking water supply for 2.6 million residents in the San Francisco Bay area. In the Rim Fire perimeter, one mixed conifer forest site (37°54'41.1"N, 119°57'56.8"W; 1020 m elevation; ~30% west-facing slope) and one chaparral site (37°50'50.5"N, 120° 0'1.9"W; 1080 m elevation; ~10% southwest-facing slope) in the Tuolumne River Watershed in the Stanislaus National Forest were selected (Fig. S1). The conifer forest site was dominated by ponderosa pine (*Pinus ponderosa*), incense cedar (*Calocedrus decurrens*), and white fir (*Abies concolor*), and the chaparral site by *Ceanothus* spp., manzanita (*Arctostaphylos*

spp.), and chamise (*Adenostoma* spp.) as identified from the remaining portions of the burned trees and shrubs. The mean annual precipitation in the region was ~920 mm, and mean annual temperature ~15 °C (7 °C for January and 24 °C for July).

As black ash and white ash are commonly formed from burning with moderate and high burn severity, respectively (Cilimburg and Short 2005), visual ash characteristics have been adopted as an indicator of burn severity (or fuel combustion completeness; Bodi et al. 2014). Ash/soil composite samples (0-5 cm) with surface deposits of white ash (AS-W) or black ash (AS-B) were collected from the conifer and chaparral sites, on October 2, 2013 (no rainfall), December 19, 2013 (~87 mm cumulative precipitation), and August 5, 2014 (~617 mm cumulative precipitation; http://www.cnrfc.noaa.gov/arc_search.php). Three AS-W and three AS-B composite samples were collected from each site, and each composite sample was a mixture of five randomly collected ash/soil cores (7.6 cm diameter × 5.0 cm depth). It should be noted that some of the AS-W samples may have contained some recognizable black ash at the interface with the mineral soil; however, the majority of the ash in the sampled materials was distinctly white with little mixing of black ash. The black ash samples contained no visual white ash materials. The thickness of ash layer in each soil core ranged from about 2.5 to 4 cm with an estimated mean thickness of 3 cm for the composite samples. The AS carbon and nitrogen concentrations were 3.6-7.1% and 0.1-0.8%, respectively (Table S1). Both sites were burned with relatively high burn severity as all surface detritus was consumed by the fire (Fig. S1), and there was complete vegetation mortality. The black and white ash coverage accounted for ~90% and ~10% respectively of the area at each site.

2.2. DOM preparation and characterization

Water-extractable DOM was obtained by mixing 40.0 g of the air-dried and sieved AS (<2 mm) with 200 mL Milli-Q water in a 250 mL Erlenmeyer flask at room temperature for 2 h at a rotary speed of 250 rpm. After filtration using a pre-rinsed 0.45 μ m polyethersulfone filter, the filtrate was analyzed for pH and specific conductivity (SC) using an Accumet XL60 dual channel pH/Ion/Conductivity meter and the remaining filtrate was stored at 4 °C in an amber glass bottle for further analyses.

To simulate the effects of multiple rainfall events and dissolution/leaching processes on the DOM chemistry and quantity from the AS, we conducted five sequential leaching events ([Chow et al. 2003](#)) on the air-dried samples collected in October 2013. Each sample was extracted with water and filtered using the aforementioned method to obtain the filtrate, and before the extraction the total weight of flask (~140 g), sample (40.0 g), and water (200.0 mL) was recorded. The residual on the filter was carefully transferred back to the Erlenmeyer flask and less than 0.1 g solid sample was lost in each extraction. For each extraction, 175 $\pm$ 5 mL filtrate was obtained depending on the water holding capacity of the materials. The extracted AS residue was retained and stored at 4 °C for ~18 h. Then, the sample was warmed for 2 h to reach room temperature, Milli-Q water added to maintain a constant total weight (flask + sample + water), and shaken prior to extraction and filtration. Five sequential extracts were obtained for each AS sample. In addition, to investigate the temporal variation of field samples, the first extracts of AS samples collected in Oct 2013, Dec 2013, and Aug 2014 were compared. The temporal samples (Oct 2013 $\rightarrow$ Dec 2013 $\rightarrow$ Aug 2014) were noted as “TS” and the

sequentially leached samples (from the first to fifth extractions of the Oct 2013 AS samples) were noted as “SLS”.

For all water extracts, we measured DOC and TDN using a Shimadzu TOC/TN analyzer, and ammonia/ammonium-N (NH_4^+ -N) and nitrate/nitrite-N (NO_x^- -N) using a Syntea® EasychemTM discrete analyzer (Chow et al. 2013). The UV-VIS absorption and fluorescence excitation-emission matrix (EEM) of the DOM was determined by a Shimadzu UV-1800 spectrophotometer (scan range: 200-700 nm) and Shimadzu spectrofluorometer RF5301 (emission, or Em: 200-550 nm; excitation, or Ex: 220-450 nm; slit width: 5 nm; Zhou et al. 2013). Based on the results, optical indices reflecting the DOM composition were calculated. Specific UV absorbance at 254 nm (SUVA_{254} in $\text{L mg-C}^{-1} \text{ m}^{-1}$), an indicator for DOM aromaticity, was calculated by dividing the UV absorbance at 254 nm by DOC concentration (Karanfil et al. 2002). The E2/E3 ratio, an indicator of molecular size of DOM, was calculated as absorbance at 254 nm divided by absorbance at 365 nm (Peuravuori and Pihlaja 1997). The raw EEM was corrected (Murphy et al. 2010) and the fluorescence regional integration based on Simpson’s rule (Zhou et al. 2013) was used to divide the EEM into five operationally-defined regions: I, tyrosine-like; II, tryptophan-like; III, fulvic acid-like; IV, soluble microbial byproduct-like; and V, humic acid-like. The percent fluorescent response in each region i ($P_{i,n}$) was determined as the proportion of area-normalized volume in region i to the entire region.

Several fluorescence-based indices, including fluorescence index, freshness index, and humification index, were calculated following Fellman et al. (2010). Specifically, the fluorescence index (FI), a common index to differentiate microbial versus terrestrial origins of DOM (~ 1.8 for microbial origin and ~ 1.2 for terrestrial origin), was determined

as the ratio of fluorescence signals at Em 470 and Em 520 nm, at Ex 370 nm (Cory and McKnight 2005). The humification index (HIX), an index for the relative abundance of humic substances, was calculated as the area under the emission spectra 435–480 nm divided by the peak area 300–345 nm, at Ex 254 nm (Ohno 2002). The freshness index (β/α), an index for the relative abundance of recently produced autochthonous DOM, was calculated as the ratio of the fluorescence signal at Em 380 nm divided by the maximum signal between Em 420 and 435 nm, at Ex 310 nm (Wilson and Xenopoulos 2009).

2.3. Disinfection byproduct formation

The chlorine reactivity of the DOM was evaluated using the disinfection formation potential (FP) test (Krasner and Scilimenti 1993). Specifically, all samples were diluted to a DOC concentration of ≤ 3 mg/L, buffered to pH 8.0 ($\text{H}_3\text{BO}_3/\text{NaOH}$), and chlorinated with freshly prepared $\text{NaOCl}/\text{H}_3\text{BO}_3$ solution (pH 8.0) at 25 °C in the dark for 24 h without headspace. The chlorine dose was based on the organic and inorganic chlorine demand calculated as $[\text{Cl}_2] = (3 \times [\text{DOC}]) + (7.6 \times [\text{NH}_3])$ based on mass concentration (Krasner and Scilimenti 1993). After quenching the residual chlorine with 10% Na_2SO_3 solution, DBPs were liquid-liquid extracted with MTBE and quantified by GC-ECD (Agilent 7890) strictly following US-EPA method 551.1. All samples were quantified for four THMs (trichloro-, dichlorobromo-, dibromochloro-, and tribromo-methanes), four HANs (trichloro-, dichloro-, bromochloro-, and dibromo- acetonitriles), three haloketones (HKs; 1,1-dichloro-2-, 1,1,1-trichloro-2-, 1,2,3-trichloro- propanones), and chloral hydrate (CHD). Five HAAs (chloro-, dichloro-, trichloro-, bromo-, and dibromo- acetic acids) were measured for one water extract of each AS type from the

different sampling times based on US-EPA method 552.2 ($n = 12$). The minimum reporting levels (MRLs) for all DBP species were approximately 0.1-0.3 $\mu\text{g/L}$.

As a commonly found and toxic DBP species in drinking water disinfected with chloramine (Shah and Mitch 2012), *N*-nitrosodimethylamine was measured for the first extracts for all AS samples after chloramination following US-EPA method 521 (Selbes et al. 2013). Specifically, AS extracts were diluted to ~ 3 mg/L DOC and then reacted with excessive chloramine (100 mg/L as Cl_2) at pH 7.8 (phosphate buffered) in the dark at 22 $^{\circ}\text{C}$ for 5 days. After quenching the residual chloramine, the samples were extracted using solid-phase extraction employing cartridges pre-packed with 2 g of coconut charcoal and eluted with dichloromethane. The extracts were analyzed using a Varian GC 3800-MS/MS 4000 equipped with a RTX-5MS column (Restek 30 m length, 0.25 mm id, 0.25 μm thickness) with a MRL of 3 ng/L.

The DOM reactivity to form DBPs was expressed as specific DBP formation potential (SDBP-FP; $\mu\text{g-DBP/mg-DOC}$), which was calculated as the ratio of the DBP concentration and initial DOC concentration. The DBP-FP per mass of AS (in $\mu\text{g-DBP/g-AS}$), was calculated as the SDBP-FP in $\mu\text{g/mg-DOC}$ multiplied by the water extracted DOC yield of the source material (mg-DOC/g-AS).

2.4. Statistical analyses.

The effects of sampling time in TS (or leaching order for SLS) and AS type (AS-W vs AS-B; conifer vs chaparral) were evaluated for all water quality parameters using two-way ANOVA with Tukey's HSD test. In this study, we primarily focused on temporal variation of DOM as DBP precursors in AS. Correlations between the SDBP-FP and the

optical indices of the AS DOM were evaluated using the two-tailed Pearson's method. All optical indices and SDBP-FP for AS DOM for TS were standardized and analyzed using principal component analysis (PCA). The SHAA-FP was not included in the PCA as we did not analyze HAA formation for all replicates, but it is expected that the SHAA-FP has a similar PCA loading to STHM-FP as it was highly correlated with STHM-FP ($r=0.71$; $P<0.001$).

3. Results and discussions

3.1. General extract quality

It has been widely reported that with higher soil burn severity, there will be lower organic matter remaining and higher soil pH due to the higher degree of mineralization and release of acid volatiles such as CO₂ (Certini 2005, Wang et al. 2015a). For water extracts from October 2013 AS samples in this study, the chaparral AS-W extract had the lowest DOC (93.3 ± 19.6 mg/L) and DON (5.0 ± 2.6 mg/L) levels (Fig. 1) but the highest pH (11.8 ± 0.7), which collectively suggest the highest degree of detritus mineralization. There was little difference in extract quality between the conifer and chaparral AS-B samples, but large differences between the conifer and chaparral AS-W samples for most parameters (Table S2). The differences between AS-W samples are likely due to a higher soil surface temperature in the chaparral fire than in the coniferous forest fire (up to 700 °C for chaparral and 200-400 °C for coniferous forest; Rundel 1983). Comparing all sample types, the chaparral AS-W had the lowest potential to contribute DOM to waterbodies.

From Oct 2013 to Aug 2014, all TS (temporal sample) extracts displayed decreasing pH, SC, DOC, and DOC/DON molar ratios with increasing post-fire leaching

(Fig. S2 & Table S2). In this study, the decreases (mean $\pm$ standard derivation) were ΔpH = 1.2 ± 1.2 , ΔSC = 0.69 ± 0.51 mS/cm (equivalent to $62 \pm 26\%$ decrease), ΔDOC = 161 ± 116 mg/L ($74 \pm 17\%$), and $\Delta(\text{DOC}/\text{DON})$ = 16 ± 6 . The AS sample type commonly interacted with the sampling time to affect the general extract water quality (most $P < 0.01$; Table S4). The AS-W showed larger percentage loss in SC and DOC with increasing post-fire leaching compared to the AS-B (Fig. 1). For SLS (sequentially leached samples), pH and DOC/DON did not change significantly while the SC, DOC, TDN, DON, $\text{NH}_4^+\text{-N}$, and $\text{NO}_x^-\text{-N}$ consistently decreased with increasing leaching from the 1st to 5th extracts (Figs. 1 & S3; Table S3).

Both TS and SLS showed decreasing SC and DOC levels with increasing leaching (Figs. S2 & S3), suggesting that the dissolution/leaching process has contributed to the loss of soluble inorganic and organic materials from the AS to water bodies. The decrease in DOC agrees with Revchuk and Suffet (2014), which showed higher DOC in forest detritus 2 months after fire than detritus 2 years after fire, although the two samples were not from a same site. These observations can explain the highest stream inorganic and DOC concentrations in the first two storm events after fire relative to the later storm events in fire-affected watersheds (Audry et al. 2014; Writer et al. 2014). Most field studies observe elevated DOC concentrations in stream water following wild and prescribed fires (Gerla and Galloway 1998, Mast and Clow 2008, McEachern et al. 2000, Revchuk and Suffet 2014, Writer et al. 2014), probably due to a higher degree of allochthonous contribution following fires (Shakesby 2011).

Regarding other parameters, the SLS failed to capture the temporal trends of ash/soil acidification and organic matter decomposition/transformation associated with

294 nitrogen leaching, DOC/DON ratio, and nitrogen speciation. First, the pH decreased in
 295 TS, but it was relatively stable for SLS (Figs. S2A & S3A). The acidic precipitation
 296 (Lajtha and Jones 2013) over the winter season could be responsible for the temporal
 297 decrease in pH of the AS extracts; however, soil pH usually takes years to recover to pre-
 298 fire levels (Granged et al. 2011). Compared to non-burned sites in the same region (Wang
 299 et al. 2015a), our study suggests that within 1 year after fire the soil acidification had not
 300 yet recovered to pre-fire levels. The relatively stable pH for SLS in this study on ash or
 301 soil could not capture the temporal variation of field samples possibly because the short-
 302 term lab processes have excluded long-term biogeochemical processes. In addition, TS
 303 showed no distinct trends in dissolved nitrogen concentrations (TDN, DON, $\text{NH}_4^+\text{-N}$, and
 304 $\text{NO}_x^-\text{-N}$) but an increase of the $\text{NO}_x^-\text{-N}/\text{NH}_4^+\text{-N}$ ratio and a decrease of the DOC/DON
 305 ratio, whereas SLS showed decreases in all dissolved measures of nitrogen but no distinct
 306 trend for $\text{NO}_x^-\text{-N}/\text{NH}_4^+\text{-N}$ or DOC/DON ratios (Table S2 & S3). Wan et al. (2001)
 307 conducted a meta-analysis on 87 fire studies and noted that soil $\text{NH}_4^+\text{-N}$ levels decreased
 308 but $\text{NO}_x^-\text{-N}$ levels increased within one year after fire. In this study, whereas a stable
 309 DOC/DON ratio in SLS indicates low microbial alteration, the lower DOC/DON ratio but
 310 higher $\text{NO}_x^-\text{-N} / \text{NH}_4^+\text{-N}$ ratio observed in Aug 2014 compared to Oct 2013 clearly reflect
 311 organic matter decomposition (organic carbon and nitrogen mineralization) and
 312 nitrification by microbes (Choromanska and DeLuca 2002; Murphy et al. 2006; Wan et
 313 al. 2001). Overall, these contrasting results between TS and SLS indicate that field
 314 exposure of AS to natural processes (e.g., leaching, new input from plant recovery,
 315 microbial alternation and sunlight irradiation) has an impact on AS DOM properties that
 316 is not manifested in the laboratory sequential leaching technique.

3.2. Optical characteristics of ash/soil extracts

Most optical parameters of TS extracts showed continuous changes with increasing leaching (Fig. S2 H-L & Table S5). For example, the $SUVA_{254}$ and HIX both increased with increasing post-fire leaching, from 1.81-3.92 to 2.93-4.55 L/mg-C/m and from 3.39-7.51 to 9.24-15.4, respectively. The E2/E3 decreased only for the AS-W extracts (from 9.47-11.2 to 6.01-6.29). Fluorescence regional integration (Fig. S4) showed decreased $P_{II,n}$ (from 25.5-33.7% to 22.0-24.7%) and $P_{IV,n}$ (from 11.5-12.4% to 10.1-12.1%) and increased $P_{III,n}$ (from 37.0-41.6% to 42.7-45.9%) and $P_{V,n}$ (from 11.8-14.1% to 17.6-18.7%) for all extracts. In contrast, $P_{I,n}$ decreased only for AS-B extracts (from 8.18-10.8% to 2.86-5.05%). Although the FI decreased for the chaparral AS-W extract, there was no consistent overall trend for FI or β/α across all extracts. Highly consistent with the TS, the SLS showed consistent trends for several optical parameters (Fig. S3 H-L & Table S6). Both $SUVA_{254}$ and HIX increased; E2/E3 decreased only for the AS-W extracts; $P_{II,n}$ and $P_{IV,n}$ decreased, $P_{III,n}$ and $P_{V,n}$ increased, and $P_{I,n}$ decreased for AS-B samples; FI decreased only for the chaparral AS-W extracts; and there was no significant change for FI or β/α across all samples. The AS sample type always interacted with the sampling time to affect the optical characteristics of extracts (all $P < 0.05$; Table S7).

As $SUVA_{254}$ is an index for aromaticity and HIX is an index for the degree of humification, the increases in these parameters for both TS and SLS suggest that the DOM from the early leachates after fire are less aromatic and humified than in the DOM from later leachates. Similarly, Revchuk and Suffet (2014) found an increase in $SUVA_{254}$

of detritus leachates two years after fire versus two months after fire in the Zaca and Jesusita fires (California) although they were not from exactly the same sites. As the E2/E3 ratio is usually inversely correlated with the molecular size of DOM (Peuravuori and Pihlaja 1997), the decreasing E2/E3 of AS-W extracts in both TS and SLS suggests that the molecular size of the DOM in AS-W extracts is lower in the early leachates compared to the later leachates.

Revchuk and Suffet (2014) found that fluorescence region I (tyrosine-like) was significantly lower in detritus leachates two years after fire versus two months after fire. This phenomenon was the case for the AS-B extracts from the Rim Fire which showed a sharp drop in $P_{I,n}$ from Oct 2013 to Dec 2013 in TS, and from the 1st to the 2nd sequential extracts in SLS. However, it was not the case in the AS-W extracts. Similar to the red shift in emission of fluorescence EEM in the present study (decreased $P_{II,n}$ and $P_{IV,n}$; increased $P_{III,n}$ and $P_{V,n}$), Revchuk and Suffet (2014; see their Fig. 8) observed a shift in the EEM of detritus leachates two years after fire versus two months after fire. As found in previous studies, the humic-like component of DOM from fire-affected ash/soil had a blue shift in emission, from “terrestrial humic-like” to “marine humic-like”, compared to DOM from non-burned detritus (Vergnoux et al. 2011, Wang et al. 2015a,b). The red shift in emission for the later leachates compared to the early leachates indicates a dissolution or degradation in DOM composition that transitions toward the composition of non-burned detritus over time (Revchuk and Suffet 2014).

In sum, all the optical indices suggest that the dissolution/leaching process associated with long-term field exposure of AS preferentially removes the less aromatic, less humified, low-molecular-weight, and blue-shifted fluoro-active DOM components,

and the remaining DOM in the AS is more aromatic and contains a higher humic-like fraction.

3.3. DBP precursors in water extracts from ash/soil

For the TS, DBP formation potential per unit mass of AS continuously decreased with increasing leaching from Oct 2013 to Aug 2014 (Fig. 2A-F) for all DBP species: THMs from 5.69-56.0 to 4.15-18.1 µg/g-AS (14-89% decrease), HAAs from 14.1-81.9 to 4.61-17.1 µg/g-AS (45-94%), HANs from 1.52-5.95 to 0.19-0.69 µg/g-AS (58-94%), CHD from 1.63-4.70 to 0.40-1.39 µg/g-AS (42-88%), and HKs from 0.11-0.78 to 0.03-0.13 µg/g-AS (67-94%). There was a similar decreasing trend for NDMA from 7.04-21.4 to 1.62-7.75 ng/g-AS (48-90%) based on the chloramination test. For SLS, as the DOC level decreased with increased leaching (Fig. 1), the average DBP-FP decreased markedly for all DBP species (Fig. 2G-J).

For the TS, the SDBP-FP did not show a consistent temporal trend across all DBP species (Fig. 3 A-F). The STHM-FP followed an increasing trend, from 11.9-33.0 to 32.1-55.4 µg/mg-DOC ($P<0.001$; Table S8). The SHAA-FP and SCHD-FP were higher in Dec 2013 than Oct 2013, but there was no difference between Dec 2013 and Aug 2014 samples. The SHK-FP, SHAN-FP, and SNDMA-FP showed no statistical differences from Oct 2013 to Aug 2014. In contrast, for the SLS, the average SDBP-FP continuously increased for all DBP species (Fig. 3 G-J). There was no statistical difference in SDBP-FP for either THMs or CHD between the 5th sequential extract and the Aug 2014 extract representing ~1 year of field exposure. In contrast, the SHAN-FP and SHK-FP of the 5th sequential extract were 3.3 and 2.5 times higher than the Aug 2014 AS extract,

respectively. The large difference in SDBP-FP between the TS and SLS extracts suggested that the temporal changes in DOM reactivity to form DBP could not be explained by the simple laboratory dissolution/leaching processes.

Similar to the temporal patterns in TDN species (DON , $\text{NH}_4^+\text{-N}$, and $\text{NO}_x^-\text{-N}$) in AS extracts, the temporal variation in DOM reactivity to form HANs was not explained by dissolution/leaching processes alone. Relatively high SHAN-FP levels in DOM from AS extracts (particularly AS-W extracts) were previously observed in lab and field fire studies (Majidzadeh et al. 2015, Wang et al. 2015a,b). HAN concentrations were commonly $>2.5 \mu\text{g-HAN/mg-DOC}$ for field AS-W extracts and up to $17 \mu\text{g-HAN/mg-DOC}$ for lab burned litter. It suggests that there may be a potential concern for nitrogenous DBPs in managing drinking water supply following wildfires, especially during the first leaching events. In this study, the TS extracts did not show as high SHAN-FP as later extracts of SLS, suggesting that the reactive fire-produced HAN precursors in the AS were possibly photo-/bio-degraded/transformed, or rapidly diluted with less reactive precursors such as the recovered plant/soil-derived carbon-rich DOM (Wang et al. 2012) under field conditions. Thus, although HANs are an important concern as they are more potent carcinogens than common carbonaceous DBPs such as THMs or HAAs (Richardson et al. 2007) and high SHAN-FP has been found in lab-burned ashes ($>10 \mu\text{g-HAN/mg-DOC}$; Majidzadeh et al. 2015, Wang et al. 2015b), the extremely high reactivity of DOM to form HANs observed for the sequential AS extracts may not persist in field AS samples ($<5 \mu\text{g-HAN/mg-DOC}$ in Wang et al. 2015a) or source waters ($\leq 2.7 \mu\text{g-HAN/mg-DOC}$ in Writer et al. (2014)).

3.4. Correlations among SDBP-FP and optical properties

For SLS, most of the SDBP-FP and optical properties changed synchronously and thus there were remarkably high correlations among these parameters (Table S9). In contrast, TS displayed obvious disparities among the major carbonaceous DBPs (THMs, HAAs, and CHD) and nitrogenous DBPs (HANs and NDMA) resulting in low correlations of SDBP-FP among DBP species and correlations of SDBP-FP to different respective optical properties compared to those found for SLS extracts (Table S10). These results indicate that the DBP precursors for different DBP species changed differentially over time after fire and cannot be explained by laboratory dissolution/leaching processes alone.

The PCA of spectroscopic characteristics and chlorine/chloramine reactivity of DOM for the TS showed clear relations among different DOM properties (Fig. 4). While the STHM-FP, HIX, $P_{III,n}$, and $P_{V,n}$ showed high loading on the first principal component (PC1; 32.5% of total variance), the SHK-FP, SNDMA-FP, $SUVA_{254}$, E2/E3, FI, β/α , and $P_{I,n}$ showed high loading on the second principal component (PC2; 29.2% of total variance; Table S11). Whereas PC1 can be considered as the major effect of increased leaching, PC2 reflects the effect of soil burn severity. The PCA indicates that SDBP-FP for the major nitrogenous DBPs (HANs and NDMA) and a group of minor carbonaceous DBPs (HKs) are independent from the SDBP-FP for major carbonaceous DBPs (THMs, HAAs, and CHD) as they have considerably different loadings on the principal components (Fig. 4 & Table S11). This observation agrees with a recent study (Wang et al. 2015b) that showed THM and CHD precursors were more dependent on heating temperature, while HAN and HK precursors were highly dependent on oxygen

availability. Based on the PCA results, the trends determined for the commonly monitored/regulated THM and HAA concentrations may not reflect the temporal trends for nitrogenous DBPs, such as HAN and NDMA concentrations.

Based on the component scores for different time categories, the DOM from Oct 2013 to Aug 2014 increased along PC1, indicating an increased humic-like DOM fraction contributing to DOM reactivity to preferentially form THMs with increasing time after fire (Fig. 4A). It is not surprising as the humic-like DOM component has long been identified as a reactive DBP precursor (Singer 1999). Regarding the difference in DOM between AS-W and AS-B, the DOM from AS-W showed higher PC2 scores than DOM from AS-B (Fig. 4B). These findings support the contention that the DOM from AS-W is more aromatic, smaller in molecular size, and less reactive in the formation of NDMA and HKs (Wang et al. 2015a,b). The difference in the PC2 score between AS-W and AS-B became smaller from Oct 2013 to Aug 2014, suggesting that the differences in chemical properties between AS-W and AS-B decreased with increasing time after fire. These decreasing differences between AS-W and AS-B could be due to physical processes in the field, such as wind, precipitation, and erosion, which may mix white and black ash materials.

3.5. Limitation of lab sequential leaching to capture temporal variations

Dissolution/leaching appears to be a major process affecting the quantity and chemical composition of organic and inorganic DBP precursors in the AS materials, as supported by the high consistency between TS and SLS for DOC, SC, optical properties of DOM (Figs. 1, S2 & S3), and DBP-FP (Fig. 2). These variations in leachate chemistry

suggest leaching of an initial pulse of fire ashes may cause significant DOM loss to surface water bodies (Revchuk and Suffet 2014) and can partially explain the higher stream DOM levels in the first several storm events after fire (Audry et al. 2014; Writer et al. 2014). Here, both TS and SLS suggest that AS DOM reactivity to form carbonaceous DBPs (THMs, HAAs, and CHD) can be elevated with increasing time or leaching after fire, as the DOM in early leachates is more hydrophilic, less aromatic, and less humified than that in later leachates. Thus, the lab dissolution/leaching process may partially capture the temporal dynamics of DBP precursor in burnt forest detritus. However, the lab dissolution/leaching processes alone failed to capture several temporal variations of ash/soil pH, DON concentration, DOC/DON ratio, nitrogen speciation (Tables S2 & S3), and DOM reactivity to form HANs (Fig. 3) in field incubated ash materials. One marked disadvantage of the lab dissolution/leaching process is that it cannot simulate several field biogeochemical processes, such as long-term microbial activity, plant recovery, photo-reaction, and other chemical reactions. Microbial processes are possibly important with increasing time after fire as reflected by the decreasing DOC/DON ratio, increasing NO_x^- / NH_4^+ ratio, and increasing humification degree (HIX) of DOM with increasing time and cumulative precipitation after fire.

4. Conclusions

This study investigated the temporal variations of spectroscopic characteristics and chlorine/chloramine reactivity for DOM from AS leachates after wildfire. Increasing cumulative precipitation after fire decreased SC and DOC concentrations in AS leachate, but increased DOM aromaticity, humic-like fraction, and reactivity to form THMs,

HAAs, and CHD. The large losses of inorganic and organic constituents from AS samples for both TS and SLS clearly indicate that there was a large pulse of DOM released from the AS in the first year after the fire. The temporal variation was dependent on sample type with the AS-W (higher burn severity) having a greater percentage loss of SC and DOM when comparing to the AS-B (moderate burn severity). Temporal variations in TDN and the DOM reactivity in forming nitrogenous DBPs did not change significantly with increasing cumulative precipitation after fire, which could not be explained solely by dissolution/leaching loss. Also, DOM reactivity in forming nitrogenous DBPs was found to be more dependent on sample type (AS-W vs. AS-B). In this study, we clearly demonstrated that fire-affected DBP precursors for different DBP species had distinct sensitivity to biogeochemical alterations with increasing cumulative precipitation after fire. Leaching effects only partially explain the temporal variations of DBP precursors in wildfire detritus.

Acknowledgements

This work was funded by NSF RAPID grant (1361678) and partially supported by NIFA/USDA (SC1700489 and 2014-67019-21615) and the Joint Fire Science Program (14-1-06-19), as presented in technical contribution number 6789 of the Clemson University Experiment Station. J.W. appreciated the financial support from the China Scholarship Council (CSC[2011]3010).

References

Audry, S., Akerman, A., Riotte, J., Oliva, P., Marechal, J.C., Fraysse, F., Pokrovsky, O.S.

501 and Braun, J.J. (2014) Contribution of forest fire ash and plant litter decay on
 502 stream dissolved composition in a sub-humid tropical watershed (Mule Hole,
 503 Southern India). *Chemical Geology* 372, 144-161.

504 Bladon, K.D., Emelko, M.B., Silins, U. and Stone, M. (2014) Wildfire and the future of
 505 water supply. *Environmental Science & Technology* 48(16), 8936-8943.

506 Bladon, K.D., Silins, U., Wagner, M.J., Stone, M., Emelko, M.B., Mendoza, C.A.,
 507 Devito, K.J. and Boon, S. (2008) Wildfire impacts on nitrogen concentration and
 508 production from headwater streams in southern Alberta's Rocky Mountains.
 509 *Canadian Journal of Forest Research-Revue Canadienne De Recherche Forestiere*
 510 38(9), 2359-2371.

511 Bodi, M.B., Martin, D.A., Balfour, V.N., Santin, C., Doerr, S.H., Pereira, P., Cerda, A.
 512 and Mataix-Solera, J. (2014) Wild land fire ash: Production, composition and eco-
 513 hydro-geomorphic effects. *Earth-Science Reviews* 130, 103-127.

514 Certini, G. (2005) Effects of fire on properties of forest soils: A review. *Oecologia* 143(1),
 515 1-10.

516 Chen, W., Westerhoff, P., Leenheer, J.A. and Booksh, K. (2003) Fluorescence excitation -
 517 emission matrix regional integration to quantify spectra for dissolved organic
 518 matter. *Environmental Science & Technology* 37(24), 5701-5710.

519 Choromanska, U. and DeLuca, T.H. (2002) Microbial activity and nitrogen
 520 mineralization in forest mineral soils following heating: Evaluation of post-fire
 521 effects. *Soil Biology & Biochemistry* 34(2), 263-271.

522 Chow, A.T., Dai, J.N., Conner, W.H., Hitchcock, D.R. and Wang, J.J. (2013) Dissolved
 523 organic matter and nutrient dynamics of a coastal freshwater forested wetland in

Winyah Bay, South Carolina. Biogeochemistry 112(1-3), 571-587.

Chow, A.T., Lee, S.T., O'Geen, A.T., Orozco, T., Beaudette, D., Wong, P.K., Hernes, P.J., Tate, K.W. and Dahlgren, R.A. (2009) Litter contributions to dissolved organic matter and disinfection byproduct precursors in California oak woodland watersheds. Journal of Environmental Quality 38(6), 2334-2343.

Chow, A.T., Tanji, K.K. and Gao, S.D. (2003) Production of dissolved organic carbon (DOC) and trihalomethane (THM) precursor from peat soils. Water Research 37(18), 4475-4485.

Cilimburg, A.C. and Short, K.C. (2005) Rating the severity of fire in forest soils; in Forest fire in the U. S. Northern Rockies: A primer. <http://www.northernrockiesfire.org/effects/soilrate.htm>. Accessed on Sep 1, 2015.

Cory, R.M. and McKnight, D.M. (2005) Fluorescence spectroscopy reveals ubiquitous presence of oxidized and reduced quinones in dissolved organic matter. Environmental Science & Technology 39(21), 8142-8149.

Dotson, A., Westerhoff, P. and Krasner, S.W. (2009) Nitrogen enriched dissolved organic matter (DOM) isolates and their affinity to form emerging disinfection by-products. Water Science and Technology 60(1), 135-143.

Emelko, M.B., Silins, U., Bladon, K.D. and Stone, M. (2011) Implications of land disturbance on drinking water treatability in a changing climate: Demonstrating the need for "source water supply and protection" strategies. Water Research 45(2), 461-472.

Fellman, J.B., Hood, E. and Spencer, R.G.M. (2010) Fluorescence spectroscopy opens new windows into dissolved organic matter dynamics in freshwater ecosystems: A

547 review. *Limnology and Oceanography* 55(6), 2452-2462.

548 Gerla, P.J. and Galloway, J.M. (1998) Water quality of two streams near Yellowstone
 549 Park, Wyoming, following the 1988 Clover-Mist wildfire. *Environmental*
 550 *Geology* 36(1-2), 127-136.

551 Granged, A.J.P., Zavala, L.M., Jordan, A. and Barcenas-Moreno, G. (2011) Post-fire
 552 evolution of soil properties and vegetation cover in a Mediterranean heathland
 553 after experimental burning: A 3-year study. *Geoderma* 164(1-2), 85-94.

554 Jolly, W.M., Cochrane, M.A., Freeborn, P.H., Holden, Z.A., Brown, T.J., Williamson, G.J.
 555 and Bowman, D.M. (2015) Climate-induced variations in global wildfire danger
 556 from 1979 to 2013. *Nature Communications* 6, 7537, 1-11.

557 Karanfil, T., Krasner, S.W., Westerhoff, P. and Xie, Y.F. (2008) *Disinfection By-products*
 558 *in Drinking Water: Occurrence, Formation, Health Effects, and Control*, Oxford
 559 University Press, Washington, DC, USA.

560 Karanfil, T., Schlautman, M.A. and Erdogan, I. (2002) Survey of DOC and UV
 561 measurement practices with implications for SUVA determination. *Journal*
 562 *American Water Works Association* 94(12), 68-80.

563 Krasner, S. and Scilimenti, M. (1993) Characterization of natural organic matter:
 564 Disinfection by-product analysis, In *Workshop on Natural Organic Matter in*
 565 *Drinking Water: Origin, Characterization, and Removal*, Preceeding, Denver, CO,
 566 USA, p. 9.

567 Lajtha, K. and Jones, J. (2013) Trends in cation, nitrogen, sulfate and hydrogen ion
 568 concentrations in precipitation in the United States and Europe from 1978 to
 569 2010: A new look at an old problem. *Biogeochemistry* 116(1-3), 303-334.

570 Lee, M.H. and Hur, J. (2014) Photodegradation-induced changes in the characteristics of
 571 dissolved organic matter with different sources and their effects on disinfection
 572 by-product formation potential. *Clean-Soil Air Water* 42(5), 552-560.

573 Lee, W., Westerhoff, P. and Croue, J.P. (2007) Dissolved organic nitrogen as a precursor
 574 for chloroform, dichloroacetonitrile, *N*-nitrosodimethylamine, and
 575 trichloronitromethane. *Environmental Science & Technology* 41(15), 5485-5490.

576 Majidzadeh, H., Wang, J.J. and Chow, A.T. (2015) Disinfection By-Products in Drinking
 577 Water Occurrence, Formation, Health Effects, and Control. In Karanfil, T. (ed),
 578 Recent Advances in Disinfection By-Products. American Chemical Society,
 579 Washington, DC, p271-292.

580 Mast, M.A. and Clow, D.W. (2008) Effects of 2003 wildfires on stream chemistry in
 581 Glacier National Park, Montana. *Hydrological Processes* 22(26), 5013-5023.

582 McEachern, P., Prepas, E.E., Gibson, J.J. and Dinsmore, W.P. (2000) Forest fire induced
 583 impacts on phosphorus, nitrogen, and chlorophyll-a concentrations in boreal
 584 subarctic lakes of northern Alberta. *Canadian Journal of Fisheries and Aquatic*
 585 *Sciences* 57(S2), 73-81.

586 Murphy, J.D., Johnson, D.W., Miller, W.W., Walker, R.F., Carroll, E.F. and Blank, R.R.
 587 (2006) Wildfire effects on soil nutrients and leaching in a Tahoe Basin watershed.
 588 *Journal of Environmental Quality* 35(2), 479-489.

589 Murphy, K.R., Butler, K.D., Spencer, R.G.M., Stedmon, C.A., Boehme, J.R. and Aiken,
 590 G.R. (2010) Measurement of dissolved organic matter fluorescence in aquatic
 591 environments: An interlaboratory comparison. *Environmental Science &*
 592 *Technology* 44(24), 9405-9412.

593 Ohno, T. (2002) Fluorescence inner-filtering correction for determining the humification
 594 index of dissolved organic matter. *Environmental Science & Technology* 36(4),
 595 742-746.

596 Peuravuori, J. and Pihlaja, K. (1997) Molecular size distribution and spectroscopic
 597 properties of aquatic humic substances. *Analytica Chimica Acta* 337(2), 133-149.

598 Plewa, M.J. and Wagner, E.D. (2009) *Mammalian Cell Cytotoxicity and Genotoxicity of*
 599 *Disinfection By-products*, Water Research Foundation, Denver, CO, USA.

600 Revchuk, A.D. and Suffet, I.H. (2014) Effect of wildfires on physicochemical changes of
 601 watershed dissolved organic matter. *Water Environment Research* 86(4), 372-381.

602 Rhoades, C.C., Entwistle, D. and Butler, D. (2011) The influence of wildfire extent and
 603 severity on streamwater chemistry, sediment and temperature following the
 604 Hayman Fire, Colorado. *International Journal of Wildland Fire* 20(3), 430-442.

605 Richardson, S.D., Plewa, M.J., Wagner, E.D., Schoeny, R. and DeMarini, D.M. (2007)
 606 Occurrence, genotoxicity, and carcinogenicity of regulated and emerging
 607 disinfection by-products in drinking water: A review and roadmap for research.
 608 *Mutation Research-Reviews in Mutation Research* 636(1-3), 178-242.

609 Rundel, P. (1983) Impact of fire on nutrient cycles in Mediterranean-type ecosystems
 610 with reference to chaparral, in *Mediterranean-type ecosystems*, Springer-Verlag
 611 Berlin, Heidelberg, Germany.

612 Santin, C., Doerr, S.H., Preston, C.M. and Gonzalez-Rodriguez, G. (2015) Pyrogenic
 613 organic matter production from wildfires: A missing sink in the global carbon
 614 cycle. *Global Change Biology* 21(4), 1621-1633.

615 Selbes, M., Kim, D., Ates, N. and Karanfil, T. (2013) The roles of tertiary amine

616 structure, background organic matter and chloramine species on NDMA
 617 formation. Water Research 47(2), 945-953.

618 Shah, A.D. and Mitch, W.A. (2012) Halonitroalkanes, halonitriles, haloamides, and *N*-
 619 nitrosamines: A critical review of nitrogenous disinfection byproduct formation
 620 pathways. Environmental Science & Technology 46(1), 119-131.

621 Shakesby, R.A. (2011) Post-wildfire soil erosion in the Mediterranean: Review and future
 622 research directions. Earth-Science Reviews 105(3-4), 71-100.

623 Singer, P.C. (1999) Humic substances as precursors for potentially harmful disinfection
 624 by-products. Water Science and Technology 40(9), 25-30.

625 Smith, H.G., Sheridan, G.J., Lane, P.N.J., Nyman, P. and Haydon, S. (2011) Wildfire
 626 effects on water quality in forest catchments: A review with implications for water
 627 supply. Journal of Hydrology 396(1-2), 170-192.

628 United States Forest Service (USFS) (2007) The U.S. Forest Service - An Overview.
 629 http://www.fs.fed.us/documents/USFS_An_Overview_0106MJS.pdf. Accessed
 630 Sep 1, 2015.

631 van Mantgem, P.J., Nesmith, J.C.B., Keifer, M., Knapp, E.E., Flint, A. and Flint, L.
 632 (2013) Climatic stress increases forest fire severity across the western United
 633 States. Ecology Letters 16(9), 1151-1156.

634 Vergnoux, A., Di Rocco, R., Domeizel, M., Guiliano, M., Doumenq, P. and Theraulaz, F.
 635 (2011) Effects of forest fires on water extractable organic matter and humic
 636 substances from Mediterranean soils: UV-vis and fluorescence spectroscopy
 637 approaches. Geoderma 160(3-4), 434-443.

638 Wan, S.Q., Hui, D.F. and Luo, Y.Q. (2001) Fire effects on nitrogen pools and dynamics in

639 terrestrial ecosystems: A meta-analysis. *Ecological Applications* 11(5), 1349-
640 1365.

641 Wang, J.J., Chow, A.T., Sweeney, J.M. and Mazet, J.A. (2014) Trihalomethanes in marine
642 mammal aquaria: Occurrences, sources, and health risks. *Water Research* 59, 219-
643 228.

644 Wang, J.J., Dahlgren, R.A., Ersan, M.S., Karanfil, T. and Chow, A.T. (2015a) Wildfire
645 altering terrestrial precursors of disinfection byproducts in forest detritus.
646 *Environmental Science & Technology* 49(10), 5921-5929.

647 Wang, J.J., Dahlgren, R.A. and Chow, A.T. (2015b) Controlled burning of forest detritus
648 altering spectroscopic characteristics and chlorine reactivity of dissolved organic
649 matter: Effects of temperature and oxygen availability. *Environmental Science &*
650 *Technology* 49(24):14019–14027.

651 Wang, J.J., Ng, T.W., Zhang, Q., Yang, X.B., Dahlgren, R.A., Chow, A.T. and Wong, P.K.
652 (2012) Reactivity of C1 and C2 organohalogen formation – from plant litter to
653 bacteria. *Biogeosciences* 9(10), 3721–3727.

654 Westerling, A.L., Hidalgo, H.G., Cayan, D.R. and Swetnam, T.W. (2006) Warming and
655 earlier spring increase western US forest wildfire activity. *Science* 313(5789),
656 940-943.

657 Wilson, H.F. and Xenopoulos, M.A. (2009) Effects of agricultural land use on the
658 composition of fluvial dissolved organic matter. *Nature Geoscience* 2(1), 37-41.

659 Writer, J.H., Hohner, A., Oropeza, J., Schmidt, A., Cawley, K.M. and Rosario-Ortiz, F.L.
660 (2014) Water treatment implications after the High Park Wildfire, Colorado.
661 *Journal American Water Works Association* 106(4), E189-E199.

662 Xie, Y.F. (2004) Disinfection Byproducts in Drinking Water: Formation, Analysis, and
663 Control, CRC Press, Boca Raton, Florida, USA.

664 Zhou, J., Wang, J.J., Baudon, A. and Chow, A.T. (2013) Improved fluorescence
665 excitation-emission matrix regional integration to quantify spectra for fluorescent
666 dissolved organic matter. Journal of Environmental Quality 42(3), 925-930.

Figure legends

Fig. 1. Dissolved organic carbon (DOC) and nitrogen (DON) in ash extracts (40 g detritus: 200 mL water). Error bars refer to the standard deviation (n=3). AS-W: ash/soil with white ash; AS-B: ash/soil with black ash.

Fig. 2. Disinfection byproduct (DBP) formation potential (FP) per mass of ash. A-F: temporal sample (TS) extracts; G-J: sequentially leached sample (SLS) extracts for October 2013 ash/soil samples. Error bars refer to the standard deviation (n=3). Haloacetic acids and *N*-nitrosodimethylamine were not measured for the sequentially leached sample extracts. AS-W: ash/soil with white ash; AS-B: ash/soil with black ash.

Fig. 3. Reactivity of dissolved organic matter from ash extracts to form disinfection byproducts (DBPs) as indicated by specific DBP formation potential (SDBP-FP). A-F: temporal sample (TS) extracts; G-J: sequentially leached sample (SLS) extracts for October 2013 ash/soil samples. Error bars refer to the standard deviation (n=3). Haloacetic acids and *N*-nitrosodimethylamine were not measured for the sequentially leached sample extracts. AS-W: ash/soil covered by white ash; AS-B: ash/soil covered by black ash.

Fig. 4. Principal component analysis (PCA) for spectroscopic characteristics and chlorine/chloramine reactivity of DOM from ash/soil leachates. Panel A shows the component scores of different leaching stages after fire and panel B shows component scores of different sample types. Detailed PC loading for different DOM properties is

690 available in Table S11. AS-W: ash/soil with white ash; AS-B: ash/soil with black ash.

Graphic Abstract

Fig. 1. Dissolved organic carbon (DOC) and nitrogen (DON) in ash extracts (40 g detritus: 200 mL water). Error bars refer to the standard deviation (n=3). AS-W: ash/soil with white ash; AS-B: ash/soil with black ash.

Fig. 2. Disinfection byproduct (DBP) formation potential (FP) per mass of ash. A-F: temporal sample (TS) extracts; G-J: sequentially leached sample (SLS) extracts for October 2013 ash/soil samples. Error bars refer to the standard deviation (n=3). Haloacetic acids and N-nitrosodimethylamine were not measured for the sequentially leached sample extracts. AS-W: ash/soil with white ash; AS-B: ash/soil with black ash.

Fig. 3. Reactivity of dissolved organic matter from ash extracts to form disinfection byproducts (DBPs) as indicated by specific DBP formation potential (SDBP-FP). A-F: temporal sample (TS) extracts; G-J: sequentially leached sample (SLS) extracts for October 2013 ash/soil samples. Error bars refer to the standard deviation (n=3). Haloacetic acids and *N*-nitrosodimethylamine were not measured for the sequentially leached sample extracts. AS-W: ash/soil with white ash; AS-B: ash/soil with black ash.

Fig. 4. Principal component analysis (PCA) for spectroscopic characteristics and chlorine/chloramine reactivity of DOM from ash/soil leachates. Panel A shows the component scores of different leaching stages after fire and panel B shows component scores of different sample types. Detailed PC loading for different DOM properties is available in Table S11. AS-W: ash/soil with white ash; AS-B: ash/soil with black ash.