Effects of feeding frequency of live prey on larval growth, survival, resistance to hyposalinity stress, Na^+/K^+ ATPase activity, and fatty acid profiles in black sea bass *Centropristis striata*

Daniel J. Russo¹, Wade O. Watanabe¹, Stephen T. Kinsey² and Pamela J. Seaton³

¹ *University of North Carolina Wilmington, Center for Marine Science, Aquaculture Program, 601, South College Road, Wilmington, NC, 28403-5927, USA.*

² *University of North Carolina Wilmington, Department of Biology and Marine Biology, 601 South College Road, Wilmington, NC, 28403-5915, USA.*

³ *University of North Carolina Wilmington, Department of Chemistry and Biochemistry, 601, South College Road, Wilmington, NC, 28403-5932, USA.*

Running title: Feeding frequency of live prey affects larval growth and quality in black sea bass

---

Correspondence: Dr. W.O. Watanabe, University of North Carolina Wilmington, Center for Marine Science, Aquaculture Program, 601 South College Road, Wilmington, NC, 28403-5927, USA, Tel: 910-962-2941, Email: watanabew@uncw.edu

Abstract

Feeding live prey organisms in a commercial marine fish hatchery requires significant labor and resources. The effects of feeding frequency of live rotifers and *Artemia* on growth, survival, hyposalinity stress resistance, and Na^+/K^+ ATPase activity in black sea bass larvae *Centropristis striata* were evaluated. Newly hatched larvae (day 0 post-hatching = d0ph) were stocked (53 larvae/L) into twenty-four 30-L tanks supported by a recirculating seawater (35 g/L) system in a controlled-environment laboratory. Temperature (22°C), aeration (225 mL/min), light intensity (~1,000 lx), and photoperiod (18 L: 6 D) were constant. Beginning on d3ph, larvae were fed a prescribed daily ration of live prey (enriched s-type rotifers and/or *Artemia* nauplii), but distributed at different frequencies: (1) 1x/d at 0800 h, (2) 2x/d at 0800 and 1600 h, (3) 3x/d at 0800, 1200 and 1600 h, and (4) 4x/d at 0800, 1200, 1600 and 2000 h, with six replicate tanks per treatment. The daily ration of live prey (maximum of 750,000 rotifers and 90,000 *Artemia*) was divided equally among feedings; therefore, prey density at each feeding (i.e., meal size) decreased at higher frequencies. By d31ph, larvae fed 4x/d showed significantly ($P < 0.05$) greater notochord length and SGR (mean $\pm$ s.e. = 7.05 ± 0.20 mm and $20.1 \pm 0.467\%/d$) compared to larvae fed 1x/d (6.02 ± 0.11 mm, and $18.1 \pm 0.204\%/d$) or 2x/d (6.08 ± 0.12 mm and $18.4 \pm 0.399\%/d$). Feeding frequency had no effect on larval survival (23.8-38.2%) or whole body fatty acid composition. Ability to withstand an acute hyposalinity (10 g/L) challenge on d32ph was significantly poorer in larvae fed 1x/d than in those fed 2x/d, 3x/d or 4x/d. Following transfer to a sub-lethal salinity (15 g/L) on d32ph, Na^+/K^+ ATPase activity ($\mu\text{moles ADP}/\text{mg protein}/\text{h}$) was higher in larvae fed 3x/d (5.61) than in those fed 4x/d (5.22) or 1x/d (5.20). The largest larvae (4x/d treatment) showed reduced Na^+/K^+ ATPase activity (vs 3x/d), likely due to accelerated development and a shifting of the Na^+/K^+ ATPase from a global distribution to

specialized osmoregulatory sites. The results demonstrated that, for a prescribed total daily ration, feeding lower prey densities (smaller meals) at higher frequencies (3x/d or 4x/d) improved larval growth and resistance to hyposalinity stress and may increase osmoregulatory ability, without affecting survival. These findings delineate more judicious use of limited live prey resources to maximize larval growth and vitality during hatchery production of pre-metamorphic stage black sea bass.

Keywords: black sea bass, ration size, feeding regime, feeding frequency, live prey, rotifers, *Artemia*, Na^+/K^+ ATPase, tissue fatty acids, larval growth, survival, stress resistance

1. Introduction

1.1 Black sea bass

Black sea bass *Centropristis striata* (family Serranidae) is a commercially and recreationally important marine finfish that inhabits coastal waters of the eastern US from the Gulf of Maine to central Florida. They are generally sedentary and are associated with hard substrates, rough bottom and natural reefs (Sedberry, 1988). Black sea bass are protogynous hermaphrodites; most fish develop first as females and then mature into males at one to five years of age (Wenner et al., 1986). Stringent regulations on harvesting of wild populations and the potential for limited future supplies and higher market prices of ocean-caught black sea bass have stimulated studies on black sea bass aquaculture to help meet market demand. Techniques for spawning adult black sea bass in captivity (Watanabe et al., 2003; Berlinsky et al., 2005), raising larvae through juvenile stages in a hatchery (Berlinsky et al. 2000, Copeland and Watanabe, 2006; Rezek et al., 2010; Carrier et al., 2011; Watanabe et al., 2015), and for growing

juvenile fish to market sizes (Copeland et al. 2003, 2005; Watanabe, 2011) have been developed in the eastern US for black sea bass, and there is now developed technology to support commercial-scale hatchery operations. Based on research funded in North Carolina by federal and state agencies, premium market size black sea bass are being produced in marine recirculating aquaculture systems at UNCW's Aquaculture Facility (Wrightsville Beach) as well as startup farms in North Carolina, Virginia and Maine. Ultra-fresh cultured black sea bass have been distributed to niche ("upscale") restaurants serving seafood throughout the state of NC as well as to US metropolitan markets (NY City, Philadelphia, Miami and San Francisco) (Wilde et al., 2008; Dumas and Wilde, 2009) to evaluate market potential. These studies have demonstrated a significant demand for farm-raised black sea bass in upscale niche markets at prices that may be profitable to startup growers.

1.2 Feeding strategies for marine finfish larvae

Feeding of the early larval stages is a bottleneck for large scale production of many commercially and recreationally important marine finfish species (Wittenrich et al., 2009; Dhert et al., 2001). This is related to significant larval mortality that is often related to a lack of prey of suitable quantity and quality. In nature, marine finfish larvae consume zooplankton and phytoplankton, including a wide range of organisms such as amphipods, decapods, and copepods (Sedberry, 1988). In aquaculture, it is more practical to feed rotifers *Brachionus rotundiformis* and brine shrimp nauplii *Artemia franciscana* to fish larvae because the techniques to culture these organisms have been standardized (Stottrup and Attramadal, 1992; Dhert, 2001; Bentley et al., 2008; Watanabe, 2011). However, these prey organisms are nutritionally deficient and must be enriched, particularly with essential fatty acids (EFAs), before feeding them to the fish larvae

(Watanabe et al., 1983; Izquierdo et al. 1989; Koven et al., 2001; Sorgeloos et al., 2001, Rezek et al. 2010; Carrier et al., 2011; Watanabe et al., 2014, 2016). Live feed organisms are expensive to produce because they require intensive manpower to grow and to improve nutritional value and are challenging to produce on a consistent basis (Bentley et al., 2008; Slembrouck et al., 2009). Providing an optimal supply of nutritionally-balanced feed is therefore important, not only for the growth and survival of fish larvae, but to minimize the use of expensive zooplanktonic prey and potentially the costs of producing fingerlings in a hatchery (Watanabe et al. 2016). Studies to determine the optimum feeding frequency of live prey for growth of larval fish are of considerable importance both biologically and economically (Rabe and Brown, 2000; Cho et al., 2003).

Larval finfish are often fed at high prey densities to maximize feed consumption, growth rate and survival (Houde and Schekter, 1978; Rabe and Brown, 2000; Slembrouck et al., 2009). Although a number of studies have investigated the effects of feeding frequency on post-larval and juvenile finfish (Folkvord and Ottera, 1993; Dwyer et al., 2002; Cho et al., 2003, Biswas et al. 2010; Luo et al. 2015), definitive experimental data demonstrating optimal feeding frequency of live prey on the performance of the early larval and pre-metamorphic stages of marine finfish is limited. In yellowtail flounder *Pleuronectes ferrugineus* feeding a moderate ration 2x/d, 4x/d or continuously resulted in better growth and survival (vs 1x/d) (Rabe and Brown, 2000). These authors concluded that feeding 2x/d may be more cost-effective than feeding 4x/d or continuously and that prolonged feeding at high prey densities may reduce gut retention time and digestive efficiency, citing earlier work in Pacific herring *Clupea harengus pallasii* (Boehlert and Yoklavich, 1984) and larval walleye *Stizostedion vitreum* (Canino and Bailey, 1995; Johnston and Mathias, 1996). In addition to minimizing wastage of prey, optimizing feeding frequency

may also improve water quality and decrease size variation and cannibalism (Dwyer et al., 2002; Biswas et al., 2010).

Optimum feeding frequency and rate varies with species, ration size, fish size and rearing system (Rabe and Brown, 2000; Biswas et al., 2010). Live prey is often a limiting resource in commercial fish hatcheries, this constraint becoming more severe as scale of operation increases. Experimental data on the effects of feeding frequency of live prey for black sea bass during the early larval and pre-metamorphic stages are needed to optimize larval performance, reduce juvenile production costs, and advance industrial scale hatchery production of this high value marine finfish species. The objectives of this study were to determine, under controlled laboratory conditions, the effects of feeding frequency of live prey, including s-type rotifers *Brachionus rotundiformis* and *Artemia franciscana* on growth, survival, fatty acid composition, resistance to hyposalinity stress, and whole body Na^+/K^+ ATPase activity in black sea bass larvae.

2. Materials and methods

2.1 Experimental animals

Eggs were collected following induced spawning of black sea bass broodstock held at the University of North Carolina Wilmington's Center for Marine Science Aquaculture Facility (Wrightsville Beach, NC, USA) (Watanabe, 2011). Broodstock were maintained outdoors in six 2.46-m diameter cylindrical tanks supplied with recirculating seawater (32-35 g/L) under an artificial photothermal regime simulating ambient offshore conditions in coastal southeastern North Carolina. Seawater was pumped from the Atlantic Intracoastal Waterway adjacent to Wrightsville Beach. Adult females at the post-vitellogenic stage of ovarian development were induced to spawn using luteinizing hormone releasing hormone-analogue (LHRHa) following

established protocols (Watanabe, 2011). Following strip spawning, eggs were transferred to a separatory funnel in full strength seawater where floating (viable) eggs were separated from sinking (non-viable) eggs. Fertilized eggs were transferred to a 125-L incubator at 16°C and 33 g/L salinity for 48-55 h.

2.2 Experimental system

The experimental units consisted of black, 30-L cylindrical (diam. = 38 cm, depth = 40 cm) larval rearing tanks (LRTs) with center draining standpipes and in-flow manifolds. The LRTs were supported by a common recirculating system, including a 113-L bubble bead filter (Aquaculture Systems Technology, New Orleans, LA); a 40-W UV sterilizer (Emperor Aquatics, Pottstown, PA); a foam fractionator (Top Fathom TF300A, 1/10 HP Foam Fractionator Pump) (Orangeburg, NY); ¾ HP pump (Jacuzzi Stingray) (Chino Hills, CA); and a 25-µm cartridge filter. Full strength seawater was recirculated through each tank at an exchange rate of 200% daily from d0ph through d14ph and then increasing to 300% from d15 through d32ph. Water temperature was held constant at 22 °C using a 1-HP heat pump (Aqualogic, San Diego, CA) (Watanabe et al., 2010; Watanabe, 2011). Air was supplied to each tank by two air stones on either side of the center standpipe. To maximize feeding efficiency, illumination was provided at 1,000 lux at the water surface by two 40-W full spectrum fluorescent light bulbs suspended over the tanks (Copeland and Watanabe, 2006). To enable larvae to feed for a prolonged time each day, photoperiod was 18L: 6D starting at 0800 h and ending at 0200 h (Moustakas et al., 2004; Copeland and Watanabe, 2006).

2.3 Experimental design

Newly hatched yolk sac larvae were transferred to twenty-four, 30-L larval rearing tanks at a density of 53 larvae/L. This marked the beginning of the experimental trials and is hereafter referred to as d0ph (day 0 post-hatch). Background microalgae *Nannochloropsis oculata* was added twice daily to maintain a density of 1×10^6 cells/mL. From d1ph to d2ph, all larval rearing tanks were fed rotifers at a density of 5 prey/mL to initiate larval feeding. During this time, rotifer density was maintained by quantifying rotifers in each tank twice daily and adding the appropriate number of enriched rotifers to make up the difference. From d2ph to d4ph, rotifer density was increased 2 prey/mL per day.

To determine the effects of feeding frequency on larval performance, beginning on d5ph larvae were fed a prescribed amount of live prey, enriched rotifers and/or *Artemia* nauplii, under different feeding regimes. Rotifers *Brachionus rotundiformis* were enriched for 16 h with “N-Rich” Algae paste (72 billion cells/mL, EPA: 18.1 mg/g dwt, DHA: 41.9 mg/g dwt) (Reed Mariculture, Campbell, CA, USA). *Artemia* nauplii were enriched for 24 h with A1-DHA Selco (Total HUFA’s: minimum 200 mg/g dwt, DHA/EPA minimum ratio: 2.5) (INVE Aquaculture, Salt Lake City, Utah, USA). To maintain nutritional quality of enriched rotifers and *Artemia*, prey were harvested at 0800 h and a portion fed to the larvae, with the remainder stored at reduced temperatures (i.e. “cold-banked”) of 4°C and 12°C, respectively, until the last feeding at 2000 h (Sorgeloos et al. 2001).

Beginning on d5ph, larvae in all experimental tanks received the same prescribed amount of rotifers (10/mL = 300,000 total prey per tank per day), with the daily ration increasing each day until a maximum density of 25/ mL (750,000 total prey per tank per day) was reached on d13ph. Rotifers were fed using four different feeding regimes: (1) one time/d at 0800 h, (2) two

times/d at 0800 and 1600 h, (3) three times/d at 0800, 1200 and 1600 h, and (4) four times/d at 0800, 1200, 1600 and 2000 h. For each treatment, the total daily ration of rotifers (300,000 - 750,000) was equally divided among the feedings. Six replicate tanks were assigned to each feeding regime.

On d15ph, newly hatched *Artemia* Instar 1 nauplii were added at a density of 0.5 prey/mL, increasing to 1 prey/mL by d17ph. Beginning on d18ph and continuing to d34ph, all experimental tanks received the same prescribed density of *Artemia* nauplii starting at 1 prey/mL (30,000 prey per day) and then increasing to 3 prey/mL (90,000 per day) by d27ph, using the same four feeding regimes described above for rotifers. By d20ph, rotifer feeding was stopped, and larvae were transitioned from *Artemia* Instar I nauplii to *Artemia* Instar II metanauplii enriched with A1-DHA Selco until d34ph.

2.4 Experimental conditions

Temperature at stocking was 16°C and was increased 1°C per day until an optimum rearing temperature of 22°C was reached on d4ph (Watanabe, 2011). Temperature, salinity (35 g/L), dissolved oxygen, and pH were measured daily in each rearing unit using a multi-parameter probe (YSI 556 MPS, YSI Inc., Yellow Springs, OH, USA). Light intensity (1,000 lx) was measured at the water surface of each tank using a light meter (Extech Instruments, Waltham, MA, USA). Aeration was supplied at approximately 226 mL/min (Watanabe, 2011). Exchange rates were kept at 200% (41 mL/min) until d12ph when it was increased to 300% (Watanabe, 2011).

2.5 Sampling and data collection

2.5.1 Growth and survival

To measure growth and survival, 10 larvae were randomly sampled from each replicate tank using volumetric methods on d0, d4, d10, d17, d25 and d31ph at approximately 0830 h (Moustakas et al., 2004). Before sampling, airflow was increased to uniformly distribute the larvae in the tanks. Larvae were sampled by plunging a 2-cm PVC pipe, 46 cm in length, into the water column and then transferring the water and larvae collected with each sample to a graduated cylinder until a total volume of approximately 600 mL and 10 larvae were collected. The number of fish were counted and divided by the water volume removed during sampling to determine sample density. Survival for each tank was calculated by dividing the sample density by the initial density. Larvae were anesthetized (0.5 mg/L of 2-phenoxyethanol) and placed in a 10-well depression slide and photographs were taken using a Nikon camera attached to a dissection microscope. A stage micrometer was also photographed as a calibration reference. Notochord lengths (NL) of larvae were measured by the computer program Image J (Research Services Branch, National Institute of Mental Health, Bethesda, Maryland, USA). Wet weight was determined by placing 10 larvae on a pre-weighed microscope slide after they were poured onto a 23- μ m mesh screen, which was blotted with a sponge from underneath to remove excess water. The slides were then placed into a laboratory oven at 60 °C for 72 h and then reweighed to determine dry weights of larvae. Specific growth rates were determined for d31ph from mean wet weight using this equation:

$$SGR = K' = \frac{\ln(m_{t_2}/m_{t_1})}{t_2 - t_1}$$

Where m_{t_2} is the mean wet weight on d31ph and m_{t_1} is the initial mean wet weight on d4ph, and $t_2 - t_1$ represents the number of days (27).

2.6 Diurnal prey (rotifer and *Artemia*) concentration-time profiles in larval rearing tanks

To elucidate the prey concentration-time profiles in the larval rearing tanks under the different feeding frequency treatments, prey concentrations were monitored *in situ* during the photophase on d12ph (Fig. 5) and on d14ph for rotifers and on d27ph for *Artemia* (Fig. 6). Rotifers and *Artemia* were sampled every 4 h for a total of 12 h using volumetric methods as described above.

2.7 Fatty acid analysis

To determine changes in fatty acid composition of live prey (rotifers and *Artemia*) during storage at reduced temperature, prey were sampled at 0, 4, 8 and 12 h post-enrichment ($N = 3$ per sampling time) to correspond with the times (0800, 1200, 1600 and 2000 h) that prey were fed to the fish larvae. To determine fatty acid composition of whole body tissue of black sea bass larvae, larvae fed enriched rotifers and *Artemia* at four different feeding frequencies were sampled from replicate tanks ($N = 4-5$ per treatment) on d32ph.

Samples of live prey and larvae for fatty acid analysis were prepared from a homogenate created from approximately 10 larvae or ~ 300 mg of rotifers or ~135 mg of *Artemia* sampled on d31ph (Watanabe et al., 1989). Samples were poured onto a 23- μ m mesh screen, which was blotted with a sponge to remove excess water. Samples (approximately 100-150 mg) were then transferred to a 1.2- mL cryogenic vial (Fisher Scientific, Pittsburg, PA, USA), filled with nitrogen gas, and then frozen (-25 °C) before lipid extraction and fatty acid analysis. Total lipid and fatty acid compositions of the prey and larvae were determined by first extracting total lipids into chloroform: methanol (Folch et al. 1957). One mL of a 0.001 g mL⁻¹ solution of C19:0 fatty acid was added to each sample as an internal standard. Lipid fatty acids were converted to their methyl esters (FAMES) for GC analysis as previously described in Watanabe et al. 2016.

2.8 Resistance to acute hyposalinity challenge

At the end of the experiment (d32ph), a stress resistance test was performed on a sub-sample of larvae to measure the ability of the larvae to withstand an acute decrease in salinity from 32 to 10 g/L under zero feeding, conditions lethal to the larvae. Ten larvae were randomly sampled from each replicate tank using volumetric methods and pooled. Twenty larvae from each treatment were stocked in a 15-L tank, with three replicate tanks per treatment. To evaluate larval quality, survival activity index (SAI) was calculated using the following equation:

$$SAI = \frac{1}{N} \sum_{i=1}^k (N - h_i) \times i$$

Where N is the total number of initial larvae, h_i is the cumulative mortality by the i th hour, and k is the number of hours that elapsed until all larvae have died (Matsuo et al., 2006). Using volumetric methods, survival was monitored every 12 h for the duration of the experiment.

2.9 Measurement of whole body Na^+ / K^+ ATPase activity following sub-lethal salinity challenge

At the end of the experiment (d32ph), the remaining larvae in each replicate tank were abruptly transferred to a lower, but sub-lethal salinity of 15 g/L. Using volumetric methods, 50 larvae were sampled volumetrically from each replicate tank at 0 h (immediately before the hyposalinity challenge) and subsequently at 12, 24, and 48 h post-transfer. Sampled larvae were placed in 10 microcentrifuge cryo-vials (5 larvae/vial) and immediately frozen. Samples were stored at -80°C.

Measurement of Na^+/K^+ ATPase activity in larval whole body samples was similar to that described by Penefsky and Bruist (1984) and McCormick and Bern (1989, 1993), modified for 96-well microplates (McCormick, 1993). In this kinetic assay, the ouabain-sensitive hydrolysis of adenosine triphosphate (ATP) is enzymatically coupled to the oxidation of nicotinamide adenine dinucleotide (reduced form, NADH), which was directly measured in a microplate reader. An assay mixture (solution A) containing 4 U lactate dehydrogenase (LDH)/ mL, 5 U pyruvate kinase (PK)/mL, 2.8 mM phosphoenolpyruvate (PEP), 0.7 mM ATP, 0.22 mM NADH, and 50 mM imidazole (pH 7.5) was made just prior to the assay (all biochemical reagents from Sigma, St. Louis, MO, USA, or VWR, Radnor, PA, USA). Assay solution B was as above, but also contained 0.5 mM ouabain. A salt solution containing 189 mM NaCl, 10.5 mM MgCl_2 , 42 mM KCl, and 50 mM imidazole (pH 7.5) was prepared in advance and was stable for several weeks at 4°C. Assay solutions A and B and salt solutions were mixed separately in a 3:1 ratio, and kept on ice.

Whole body larval black sea bass were homogenized for 10-15 s in the presence of 25 μL of a solution (SEI) (pH 7.4) containing 300 mM sucrose, 20 mM disodium ethylenediamine tetraacetate (Na_2EDTA) and 100 mM imidazole (pH 7.1) (Zaugg, 1982) using a motorized pestle (Kontes, Vineland, NJ, USA). Samples were centrifuged at 5,000 g for 30 s and then a 1:10 dilution of the supernatant from centrifuge to SEI was created. For every sample, ten microliters of diluted homogenate was added to each of four wells. Two-hundred microliters of the A-plus-salt mixture was added to two wells per sample, and 200 μL of the B-plus-salt mixture was added to the other two wells per sample. The plate was then placed in a temperature-controlled plate reader (Flex Station 3, Molecular Devices Corp., Menlo Park, CA, USA) and the linear rate of NADH disappearance was measured at 340 nm for 10 min. The linear rate from 2 to 10 min in

each pair of duplicate wells was determined, and Na^+ , K^+ -ATPase activity was calculated as the difference in ATP hydrolysis in the absence and presence of ouabain, expressed as micromoles of ADP per milligram of protein per hour.

2.10 Analytical methods

The effects of feeding frequency on notochord length, wet weight, dry weight, SGR, fatty acid and lipid concentrations, SAI and Na^+/K^+ ATPase activity were expressed as treatment means $\pm$ standard error of the mean (SEM) and compared by a one-way analysis of variance (ANOVA). A two-way ANOVA was used to investigate the interactive effects of sampling time and feeding frequency on Na^+ , K^+ -ATPase activity of whole-body black sea bass larvae. To test the assumptions of ANOVA, all data were analyzed using Levene's homogeneity of variance test prior to the ANOVA. A Tukey-Kramer analysis ($P < 0.05$) was used for multiple comparisons among means. If data did not meet the assumptions for an ANOVA, a non-parametric Wilcoxon/Kruskal-Wallis test was used. Percentage data was arcsine transformed before analysis. Analyses were performed using the JMP IN version 7 statistical software (SAS Institute Inc., Cary, NC, USA).

3. Results

3.1 Growth

3.1.1 Notochord length

Larvae were stocked on d3ph and initial larval length and weight were determined beginning on d4ph. Mean initial notochord length on d4ph was $2.51 \text{ mm} \pm 0.08$ (Fig. 1a). Notochord length increased steadily in all treatments to d10ph (mean = 3.65 mm) and d17ph

(mean = 4.99 mm), with no significant effects of feeding frequency observed. The increase in notochord length among treatments slowed from d17ph, and there was a departure in growth rates among treatments on d25ph (Fig. 1a) when mean notochord length generally increased with higher feeding frequency and was significantly ($P < 0.05$) higher in fish fed 4 x/d (5.39 mm) than in fish fed 1x/d (5.05 mm) (Fig. 1a). This trend toward higher notochord length with increasing feeding frequency was maintained on d31ph, when notochord length of fish fed 4x/d (mean = 7.05 mm) was significantly higher than at 1x/d (mean = 6.02 mm) or 2x/d (mean = 6.08 mm) (Fig. 2a).

3.1.2 Wet and dry weight

On d4ph, wet weights (mean = 0.022 mg) did not differ among treatments (Fig. 1b). Wet weight increased steadily in all treatments on d10ph (mean = 0.29 mg), d17ph (mean = 0.91 mg), and on d25ph (mean = 1.40 mg), with no significant differences. However, there was a clear increase in wet weight and a departure in growth rates among treatments on d31ph (Figs. 1b, 2b) when wet weight increased with increasing feeding frequency and was markedly higher ($P < 0.05$) at 4x/d (5.10 mg) compared to 1x/d (2.86 mg). There were no significant differences between the 2x/d (3.16 mg) and the 3x/d (4.16 mg) treatments (Fig. 2b).

Due to small larval size, accurate measurements of dry weight on d4ph could not be obtained (Table 1). Dry weights increased steadily from d10ph (mean = 0.06 mg) to d17ph (mean = 0.17 mg) and to d25ph (mean = 0.26 mg) in all treatments, with no significant differences. On d31ph, there was an apparent trend toward higher dry weights with increasing feeding frequency from lowest values in larvae fed 1 x/d (mean = 0.55 mg) to maximum values

in larvae fed 4 x/d (mean = 0.96 mg) (Table 1). Due to high inter-replicate variability, no statistical differences were observed.

3.1.3 Specific growth rate (SGR)

A trend toward higher SGR of body weight (wet) with increasing feeding frequency was observed, from a minimum (18.1 %/d) in larvae fed 1x/d to a maximum (20.1 %/d) in larvae fed 4x/d (Table 2). SGR was significantly higher in larvae fed 4x/d than in those fed 1x or 2x/d (Table 2).

3.2 Survival

No significant effects of feeding frequency on larval survival were observed for the duration of the study. From d4ph, survival decreased slowly to d10ph (mean = 87.9%) and to d17ph (mean = 85.8%) (Table 3). Survival decreased sharply in all treatments by d25ph (mean = 50.2%), remained relatively stable through d31ph (mean = 52.6%), and then fell to 31.4% by d35ph (Table 3).

3.3 Resistance to acute hyposalinity challenge

Following acute hyposalinity challenge (from 32 to 10 g/L) on d32ph, larvae fed 2x/d, 3x/d, and 4x/d had significantly ($P < 0.05$) higher survival activity indices (SAI) (526.5, 498.0, 491.9, respectively) than larvae fed 1x/d (244.5) (Table 2).

3.4 Diurnal prey concentration-time profiles in larval rearing tanks

Mean rotifer density on d12ph under the different feeding frequency treatments are shown in Fig. 5. Under the 1x/d feeding treatment, maximum concentration (20 rotifers/mL) was observed at 0800 h, abruptly decreased to 7.6 rotifers/mL within 1 h, and then gradually declined to zero by 9 h post-feeding (Fig. 5). Under the 2x/d feeding treatment, maximum concentration (10 rotifers/mL) was observed at 0800 h, abruptly decreased to 2.3 rotifers/mL within 1 h post-feeding, and then gradually declined to zero by 7 h post-feeding. Rotifer concentration increased to 10 rotifers/mL at the second feeding at 1600 h, decreased abruptly to 2.1 rotifers per mL by 1700 h, and then gradually declined to zero by 2000 h. The 3x/d feeding treatment showed similar trends with a maximum concentration of 6.6 rotifers/mL at 0800 h, 1200 h, and 1600 h, decreasing abruptly within 1 h of each feeding to an average of 0.66 rotifers/mL, then declining to zero over the next 3 h. Under the 4x/d feeding treatment, maximum concentrations (5 rotifers/mL) were observed at 0800 h, 1200 h, 1600 h, and 2000 h with each feeding. After each feeding, rotifer density decreased abruptly within 1 h to an average of 0.26 rotifers/mL and then declined to zero within 1-3 h after each feeding.

On d14ph, total prey concentration was increased from 20 rotifers/mL to 25 rotifers/mL. Similar trends to those found on d12ph were observed in rotifer concentrations across treatments and times, except that peak rotifer concentrations increased to 25, 12.5, 8.33 and 6.25 rotifers/mL at feeding frequencies of 1, 2, 3 and 4x/d, respectively.

Artemia concentration profiles on d27ph (Fig. 6) were similar to those observed for rotifers on d12ph (Fig. 5). Under the 1x/d treatment, maximum concentration (3 *Artemia*/mL) was observed at first feeding (0800 h), abruptly decreased to 1.28 *Artemia*/mL within 1 h, and then gradually decreased to zero by 9 h post-feeding (Fig. 6). Under the 2x/d treatment, maximum concentration (1.5 *Artemia*/mL) was observed at first feeding (0800 h), abruptly

decreased to 0.02 *Artemia*/mL within 1 h, and then decreased to zero after 3 h. *Artemia* concentration increased to 1.5 *Artemia*/mL again at the 1600 h feeding, and then abruptly decreased to zero within 1 h (Fig. 6). Under the 3x/d treatment, maximum concentration (1 *Artemia*/mL) was observed at 0800 h and then decreased to zero within 1 h. This pattern was repeated at 1200 and 1600 h. Under the 4x/d treatment, maximum concentration (0.75 *Artemia*/mL) was observed at 0800, 1200, 1600 and 2000 h, decreasing to zero within 1 h of feeding.

3.5 Fatty acid composition

3.5.1 Fatty acid composition of enriched rotifers

Following enrichment of rotifers, no significant effects of storage time (from 0800 h to 2000 h) at 4 °C were observed on total lipid (mean = 312 mg g⁻¹ dwt) (Table 4). There were no significant differences among storage times in concentrations of any of the fatty acids, including the total saturates, monounsaturates, n-3 or n-6 PUFAs, or ratios of n-3 to n-6 PUFAs (Table 6).

3.5.2 Fatty acid composition of enriched *Artemia*

Following enrichment of *Artemia*, no significant effects of storage time (from 0800 h to 2000 h) at 4°C were observed on total lipid (mean = 431 mg g⁻¹ dwt) (Table 5). There were no significant differences among storage times in concentrations of any of the fatty acids, including the total saturates, monounsaturates, n-3 or n-6 PUFAs, or ratios of n-3 to n-6 PUFAs (Table 5).

3.5.3 Fatty acid composition of larval black sea bass

The fatty acid composition whole body tissue of black sea bass larvae fed at the different feeding frequencies is shown in Table 6. No significant ($P > 0.05$) treatment effects were observed in total lipid (mean = 221.5 mg g⁻¹ dwt) or any of the fatty acids, including ARA (range

= 2.29 – 2.49 %TFA), EPA (7.50 – 7.79 %TFA), DHA (9.27 – 9.57 %TFA), total saturates, monounsaturates, n-3 or n-6 PUFAs, or ratios of n-3 to n-6 PUFAs, DHA to EPA, EPA to ARA, or DHA to ARA (Table 8).

3.6 Na⁺/K⁺ ATPase activity following sub-lethal salinity challenge

Black sea bass larvae abruptly transferred on d32ph from 32 g/L to a sub-lethal salinity of 15 g/L showed significant effects of feeding frequency on whole body Na⁺/K⁺ ATPase activity, but there were no significant effects related to time post-transfer and there were no interactive effects. Hence, the effects of feeding frequency on Na⁺/K⁺ ATPase activity were compared by combining data for all sampling times (Fig. 7). Na⁺/K⁺ ATPase activity (μmol ADP/mg protein/h) was significantly ($P < 0.05$, Student's t-test) higher in the larvae fed 3x/d (5.61 ± 0.13) than in larvae fed 1x/d (5.20 ± 0.10) or 4x/d (5.22 ± 0.11).

4. Discussion

4.1 Growth

The results of this study demonstrated that, given a fixed daily ration of live prey, the frequency of feeding of prey, which required lower prey concentrations (i.e., smaller meal sizes) with increasing feeding frequency, significantly influenced growth of black sea bass larvae from first-feeding through the pre-metamorphic stage. By d31ph, notochord length, wet weight and SGR increased with increasing feeding frequency from a minimum in larvae fed 1x/d to a maximum in larvae fed 4x/d (Figs. 1, 2, 3, and 4). Thus, black sea bass larvae showed slowest growth when fed a high concentration of live prey once per day, but growth improved when larvae were fed the same daily ration distributed over an increasing number of smaller meals.

These results are not directly comparable to earlier studies of feeding frequency in larval finfish in which live prey was fed to satiation at each feeding (Mollah et al. 1988), ration size was the same at each feeding (Rabe et al. 2000), or fish were fed artificial microdiets at each feeding (Xie et al. 2011). For example, yellowtail flounder *Pleuronectes ferrugineus* larvae fed a moderate prey level (8,000/L) 1x/d grew slower than fish fed 2x/d, 4x/d, or continuously (Rabe et al., 2000). However, since meal size was constant at each feeding, total ration in the 2x/d and 4x/d treatments were two and four times higher, respectively, than in the 1x/d treatment (Rabe et al. 2000). Growth of catfish *Clarias microcephalus* (Mollah et al., 1988) and yellow croaker *Pseudosciaena crocea* (Xie et al., 2011) larvae also increased at higher feeding frequencies when larvae were fed to satiety at each feeding. The present study is unique in that it reports higher growth rates in fish fed equivalent daily rations of prey, but at different frequencies and prey concentrations.

Improved growth at higher feeding frequencies in larval black sea bass may have been related to a number of factors. At the lowest feeding frequency (1x/d), prey concentrations were depleted after 8 h, with 10 h of daylight remaining and with 16 h until the next feeding (8 am) (Fig. 5). Likewise, for larvae fed 2x/d, prey concentrations were depleted 6 h after the first feeding, remained at zero for 1 h until the second feeding, and then were depleted again for 15 h, so that larvae were deprived of prey for a total of 16 h in a diurnal cycle (Fig. 5). For the larvae fed 3x/d, larvae were deprived of prey for 1 h after each of the first two feedings and then for 13 h after the final feeding for a total of 15 h of total prey deprivation. Larvae fed 4x/d were deprived of prey for 1 h after the first feeding and second feedings, for 2 h after the third feeding, and then for 12 h after the fourth feeding for a total 16 hours of prey deprivation (Fig. 5). Hence, in all

treatments, the total duration of prey deprivation was comparable, ranging from 9-10 h during the photophase and 15-16 h over a 24-h diurnal cycle.

While total duration of feed deprivation among treatments was comparable, the length of continuous feed deprivation was disparate and likely affected growth. Whereas fish fed 1x/d experienced 10 h of continuous feed deprivation, those fed 2x/d, 3x/d and 4x/d experienced 8 h, 7 h and 6 h, respectively, of continuous feed deprivation. Hence, at lower feeding frequencies (Fig. 5) the longer durations of continuous feed deprivation may have increased competition for limited prey. In contrast, at the highest feeding frequency (4x/d), a shorter continuous duration of feed deprivation alleviated competition and improved growth. In larval yellow croaker *Pseudosciaena crocea*, increasing size variation and slower growth were observed with a decrease in feeding frequency (Xie et al., 2011). It is possible that size heterogeneity of black sea bass in the present study was increased at the lower feeding frequencies which inhibited growth.

The energy that is required to breakdown, absorb, transport and assimilate nutrients from a meal (specific dynamic action of the meal, or SDA) require up to 25% of daily energy expenditures in fish (Secor, 2011). At constant meal type, fish size, and water temperature, an increase in meal size results in a larger SDA (Secor, 2011). While the black sea bass larvae under all treatments in this experiment were fed the same number of prey (i.e. ration size) each day, the allocation per meal was reduced as feeding frequency was increased. Thus, fish fed smaller meals more frequently would have a lower SDA than fish fed a single large meal, consistent with the trend toward faster growth with increasing feeding frequency observed in the present study.

Higher growth at higher feeding frequencies may have also been related in part to water flow and exchange patterns in the larval rearing tank inherent to a recirculating aquaculture system. Daily water exchange ranged from 200 to 300% in each tank throughout the experiment. Although effluent standpipes in each tank were screened to prevent larvae from escaping, smaller prey were lost with water exchange, and recycled water was 1- μ m-filtered to exclude prey. At higher feeding frequencies (3x/d and 4x/d), a rapid and total depletion of prey was observed after each feeding (Figs. 5 and 6), suggesting that larval appetite and prey consumption were maximized and prey loss via water exchange was minimized. In contrast, at the lower feeding frequencies (1x/d and 2x/d), high prey concentrations at each feeding may have rapidly satiated the larvae as evidenced by a rapid decline in prey concentration within 1 h post-feeding (Figs. 5 and 6) followed by a more gradual decline. Thus, a higher percentage of prey was removed from the tank via system water exchange.

The lack of differences in growth related to feeding frequency during through d17ph (i.e., rotifer phase) and the significant departure in growth from d25ph (i.e., *Artemia* phase) (Fig. 1) seem contradictory. As observed in larval red porgy, it is likely that the first 13-d period of feeding on rotifers was insufficient for treatment effects to be clearly demonstrated and that differences in growth rates were accentuated during the *Artemia* phase when larval feeding and specific growth rates (Fig.3) were much higher (Andrade et al., 2012; Watanabe et al., 2016).

4.2 Survival

Final survival on d32ph in all treatments was high (23.4 – 32.5%) and was typical for hatchery rearing of black sea bass (Copeland et al., 2006; Rezek et al., 2010; Carrier et al., 2011; Watanabe, 2011) but was not influenced by feeding frequency (Table 5). Hence, while a

constant daily ration fed under different feeding frequencies caused differential growth of the larvae, the level of nutrition in all treatments met the minimum maintenance requirements for survival. In gilthead sea bream *Sparus aurata* (Goldan et al., 1997) and in yellowtail flounder *Pleuronectes ferrugineus* (Rabe and Brown, 2000), no differences in larval survival were observed as daily ration increased proportionately with increasing feeding frequency.

Behavioral changes may help explain the relatively uniform survival of black sea bass larvae at the different feeding frequencies in the present study. Cod *Gadus morhua* larvae fed at low prey densities maintained good growth and survival by increasing swimming activity and attack rate and decreasing the size selectivity of prey, allowing them to capture more prey (Munk and Kiorboe, 1985; Munk, 1995). In yellowtail kingfish *Seriola lalandi* (Valenciennes, 1833), rotifer consumption and larval survival over the first 12 days post-hatching were unaffected when prey were fed at high or low densities (10-30 or 5-8 individuals/mL, respectively), or at high density until 5 dph and then at a low density thereafter (Woolley and Partridge, 2015), presumably related to behavioral adaptations to prey concentrations. Black sea bass larvae may likewise increase hunting activity and prey consumption at lower prey densities.

During larval culture of black sea bass, artificial microdiets are typically co-fed with both rotifers and *Artemia* beginning on d15ph, when live prey is gradually reduced and artificial feeds increased through d35ph (Watanabe 2011). Microdiets were not included in the present study in order that the effects of manipulating the feeding frequency of live prey could be delineated. Nonetheless, survival (23.4 – 32.5%) was typical for hatchery rearing of black sea bass. The co-feeding of microdiets with live prey at correspondingly high frequencies may further improve growth performance and the efficiency of utilizing live prey resources in a hatchery.

4.3 Fatty acid concentrations of live prey and black sea bass larvae

4.3.1 Fatty acid composition of enriched rotifers

Starvation can cause a loss of essential fatty acids in enriched prey, which can be minimized by storing prey at low temperature (Estevez et al. 1998). Concentrations of DHA, EPA, ARA, and total lipids in enriched rotifers and *Artemia* did not change ($P < 0.05$) over time when held at $\sim 12^{\circ}\text{C}$ for up to 16 h until feeding. This indicates that the effects of feeding frequency on larval performance in this study were not influenced by a reduction in prey nutritional quality post-enrichment. Rainuzzo et al. (1989) found that starvation of rotifers for 49 h at 16°C after enrichment did not change the essential fatty acid (n-3 HUFA) profile of rotifers, although total amounts were decreased by 25%. On the other hand, Birkou et al. (2012) found that *Brachionus* spp. enriched with n-3 PUFAs were able to maintain their improved nutritional value for only several hours after being transferred to a low PUFA-containing media at 25°C .

ARA levels in rotifers ranged from 2.40 %TFA (2.69 mg/g dwt) to 2.95 %TFA (5.59 mg/g dwt), whereas a previous study (Rezek et al., 2010) showed that growth and survival of black sea bass improved as ARA was increased from 0 to 6% TFA (4.37 – 18.0 mg/g dwt). Koven et al. (2001) found that increasing ARA levels in live feeds from 1.0 to 2.75 mg/g dwt did not affect larval growth, but increased tolerance to handling stress in larval gilthead sea bream *Sparus aurata*. In the present study, the ratio of DHA/ARA (1.80 to 4.25) was within the range of 2:1 recommended for many species of larval marine fish (Sargent et al., 1999). The ratio of DHA/EPA (0.324 to 0.538) was below the suggested dietary ratio (2:1) for European sea bass *Dicentrarchus labrax*, suggesting that the rotifer enrichment medium was deficient in DHA, or that the enrichment protocol was not optimized (Sargent et al., 1999). The ratio of EPA/ARA in

rotifers (ranged from 1.79 to 4.25) was higher than the optimal dietary ratio of 1:1 for European sea bass (Sargent et al., 1999).

4.3.2 Fatty acid composition of enriched Artemia

Fatty acid composition of cold-banked *Artemia* did not change significantly over a 12 h period post-enrichment (0800 h - 2000 h). Similar findings were reported by several workers (Coutteau and Mourente, 1997; Han et al., 2001; Koven et al., 2001). The ratio of DHA/EPA found in cold-banked *Artemia* (1.08-1.20) was lower than that in the enrichment medium (minimum DHA/EPA of 2.5). Mean DHA and EPA levels in enriched *Artemia* (38.1 and 33.2 mg/g dwt, respectively) were higher than those previously reported in DHA Selco-enriched *Artemia* nauplii (17.7 and 10.8 mg/g dwt) (Lavens and Sorgeloos, 1996). During starvation, *Artemia* selectively catabolize certain fatty acids such as 18:4n-3, 22:5n-3, and 22:6n-3, while retaining fatty acids such as 18:0, 20:4n-6, and 20:5n-3, but significant losses of fatty acids were found only after 72 h (Coutteau and Mourente, 1997). This is consistent with the stable fatty acid concentrations found in enriched *Artemia* cold-banked for 12 h post-enrichment in the current experiment. Furthermore, retro-conversion of DHA to EPA by *Artemia* via 22:5n-3 (docosapentaenoic acid, DPA) (Sargent et al., 1999; Rezek et al., 2010; Carrier et al., 2011) could explain the low DHA relative to the amount supplied by the enrichment. The EPA/ARA ratio (6.19 - 7.80) was above the 1:1 recommended dietary ratio for European sea bass larvae (Sargent et al., 1999). The DHA/ARA ratio (7.19-9.15) was also above the recommended 2:1 dietary ratio for European sea bass larvae (Sargent et al., 1999).

4.3.3 Fatty acid composition of black sea bass larvae

There were no significant differences in total lipid or fatty acid composition of whole body larval black sea bass fed enriched rotifers and *Artemia* at the four feeding frequencies (1x/d, 2x/d, 3x/d, or 4x/d). This may be explained by increased whole body protein content associated with greater muscle (rather than lipid) deposition in faster-growing fish in the 3x/d and 4x/d treatments. Different fish species use different strategies for meeting energetic demands during food deprivation, including selectively catabolizing carbohydrates, lipids and proteins (Czesny et al., 2003; Barcellos et al., 2010). Some species (eel *Anguilla anguilla*, goldfish *Carassius auratus*, and plaice *Pleuronectes platessa*) use muscle protein as their primary fuel source, whereas others (northern pike *Esox lucius*, roach *Rutilus rutilus*, and golden perch *Macquaria ambigua*) use lipids (see Czesny et al., 2003 for review). Proximate composition data in the present study suggest that lipid was utilized and deposited uniformly among feeding frequency treatments, while more protein was catabolized for energy at lower feeding frequencies, causing reduced muscle growth. This is consistent with what has been reported in larval Ayu (Yao et al., 1994; Cho et al., 2003; Xie et al., 2011), Trairao *Hoplias lacerdae* (Light et al., 2005), and yellow croaker *Pseudosciaena crocea* (Xie et al., 2011) where growth increased with increasing feeding frequency, but without correlation to total lipid or fatty acid profiles. In other species including Ayu (Cho et al., 2003), meager *Argyrosomus regius* (Roo et al., 2010), and red-spotted grouper *Epinephelus akaara* (Kayano et al., 1993; Yao et al., 1994), total lipid has been reported to increase with increasing feeding frequency.

4.4 Resistance to acute hyposalinity challenge (Survival Activity Index, SAI)

The SAI test was designed to reveal differences in the ability of larvae from different feeding frequency treatments to withstand an acute salinity decrease under zero feeding,

conditions prescribed to induce short-term, albeit gradual mortality. Despite uniformly high survival among feeding frequency treatments through the pre-metamorphic stage, stress resistance (i.e., SAI) of d32ph black sea bass larvae abruptly transferred from seawater (32 g/L) to a lethal hyposalinity of 10 g/L under conditions of feed deprivation was poorer in larvae fed only 1x/d than in larvae fed at higher frequencies (Table 2). In tilapia *Oreochromis mossambicus*, feed deprivation prior to salinity challenge impaired the salinity acclimation process, indicating that nutritional state affects osmoregulatory ability (Vijayan et al., 1996). In the present study, it is possible that black sea bass larvae under the lowest feeding frequency (1x/d) were exposed to longer periods of continuous feed deprivation during a diurnal cycle (Fig. 5), reducing their ability to meet the increased metabolic demands of the hyposalinity challenge. Hence, these slowest growing fish succumbed more rapidly (i.e., lower SAI) to the combined stresses of feed deprivation and hyposalinity challenge. Other workers have reported differences in larval stress resistance even when no obvious differences in survival were evident (Ako et al., 1994, Koven et al., 2001, Watanabe et al., 2016).

4.5 Na⁺/K⁺ ATPase activity following sub-lethal salinity challenge

In this study, exposure of 32dph black sea bass larvae to a sub-lethal hyposalinity challenge (i.e., from 32 to 15 g/L) elicited a greater osmoregulatory response (i.e., whole body Na⁺/K⁺ ATPase activity) in larvae fed 3x/d than in those fed 1x/d or 4x/d. Na⁺/K⁺ ATPase is a key enzyme involved in osmoregulation in larval fish and is a component of the plasma membranes of chloride cells which are the cellular sites for the excretion of monovalent ions (primarily Na⁺, Cl⁻ in seawater) (Li et al., 1995; Varsamos et al., 2004, 2005). In most teleost fish larvae, Na⁺/K⁺ ATPase is concentrated in the chloride cells of the skin, but as larvae develop these cells become more localized and are found primarily in the integument, gills,

urinary system, and digestive tract (Hiroi et al., 1998; Varsamos et al., 2002, 2005; Li et al., 2011).

In early larval marine finfish species, such as obscure puffer *Takifugu obscurus* (Li et al., 2011) and European sea bass *Dicentrarchus labrax* (Varsamos, 2002; Varsamos et al., 2004), a hyper- or hypoosmotic challenge elicited an initial increase in whole body Na^+/K^+ ATPase activity within 12-48 h. In the present study, no increase in Na^+/K^+ ATPase in whole body tissues of black sea bass was observed from 12 to 48 h following hyposalinity challenge, possibly due to the magnitude of the salinity challenge conditions used. After transfer of day-old obscure puffer larvae hatched in freshwater to salinities of up to 16 g/L, Na^+/K^+ ATPase activity increased threefold after 48 h at higher salinities of 12-16 g/L, but not at the lower salinities of 0-10 g/L (Li et al., 2011). In European sea bass, early larvae cope more easily with hypoosmotic than with hyperosmotic challenges (Varsamos et al., 2004). This suggests that the modest hyposalinity challenge (i.e., from 32 to 15 g/L) used in the present experiment elicited a weak Na^+/K^+ ATPase response in black sea bass larvae. The selection of more acute salinity challenge conditions may accentuate the Na^+/K^+ ATPase response, but must be weighed against the possibility of overwhelming the larva's osmoregulatory capacity and causing rapid mortality (Varsamos et al., 2004; Carrier et al., 2011).

Developmental stage (i.e., ontogeny) can also influence the magnitude of the Na^+/K^+ ATPase response in larval fish. After European sea bass larvae reached 5 mm (~10dph), Na^+/K^+ ATPase activity increased due to chloride cell accumulation in the gills, and overall osmoregulatory ability increased as the larvae developed the ability to regulate drinking rate (Varsamos et al., 2004; 2005). It is likely that near metamorphic stage (d32ph) black sea bass in

the present study had well-developed osmoregulatory organs and could regulate drinking rate to adapt to the hyposalinity challenge without a need to boost Na^+/K^+ ATPase enzyme activity.

Across all sampling times post salinity challenge, Na^+/K^+ ATPase activity was higher in larvae fed 3x/d than in larvae fed 1x or 2x/d, possibly related to ontogenetic factors. In the present study, the development of gills and chloride cells in branchial filaments may likewise have been greater in larger black sea bass larvae fed 3x/d (vs 2x or 1x/d). On the other hand, the lower Na^+/K^+ ATPase specific activity response in largest fish fed 4x/d compared to fish fed 3x/d seems contradictory. As observed in tilapia *Oreochromis sp.* (Watanabe et al., 1985), increased osmoregulatory ability in black sea bass may be related to fish size rather than age. Although black sea bass in all feeding frequency treatments in the present study were the same age (32dph) at the time of hyposalinity challenge, larger larvae in the 4x/d treatment likely had more well-developed osmoregulatory organs (e.g. gills) and ability to regulate drinking rate, decreasing the need to boost specific Na^+/K^+ ATPase activity.

4.6 Conclusions

The results demonstrate that, given a prescribed total daily ration of live rotifers and *Artemia*, feeding lower prey concentrations (i.e., smaller meal sizes) at higher frequencies, within a range of 1x, 2x, 3x, and 4x/d, increases growth performance and stress resistance of black sea bass larvae through pre-metamorphic stages, with no diminution of survival. The results suggest that, when live prey resources are limiting in a marine fish hatchery, feeding lower concentrations of prey at higher frequencies maximizes growth, survival, and larval vitality and therefore optimizes the efficiency of utilization of expensive planktonic prey while maximizing fingerling yield and quality. These findings have important implications for improving the economics of marine finfish hatchery operations by minimizing the duration of the

larval rearing period to produce healthy fingerlings and to maximize productivity of hatchery and growout operations.

5. Acknowledgements

We thank the United States Department of Agriculture, National Institute of Food and Agriculture (USDA-NIFA Grant No. 2006-38854-03596) and Marine Biotechnology in North Carolina (MARBIONC) for funding support, Ana Jimenez and Joan Willey for helpful advice and Patrick Carroll, Troy Rezek, Amanda Meyers, Jen Gabel, Matt Dawson, Tyler Gibson, Casey Kelley, and Robert Farnell for technical assistance.

6. Literature Cited

- Ako, H. Tamaru, C.S., Bass, P., and Lee, C-S. 1994. Enhancing the resistance to physical stress in larvae of *Mugil cephalus* by the feeding of enriched *Artemia* nauplii. *Aquaculture* 122, 81-90.
- Barcellos, L.J.G., Marqueze, A., Trapp M., Quevedo, R.M., Ferreira, D. 2010. The effects of fasting on cortisol, blood glucose and liver and muscle glycogen in adult jundia *Rhamdia quelen*. *Aquaculture* 300, 231-236.
- Bentley, C.D., Carroll, P.M., Riedel, A.M., Watanabe, W.O. 2008. Intensive rotifer production in a pilot-scale continuous culture recirculating system using condensed microalgae. *Journal of the World Aquaculture Society* 39, 625-635.
- Berlinsky, D.L., King V, W., and Smith, T.I.J. 2005. The use of luteinizing hormone releasing hormone analogue for ovulation induction in black sea bass (*Centropristis striata*). *Aquaculture* 250, 813-822.
- Berlinsky, D., Watson, M., Nardi, G., Bradley, T.M., 2000. Investigations of selected parameters for growth of larval and juvenile black sea bass *Centropristis striata* L. *J. World Aquac. Soc.* 31, 426-435.
- Birkou, M. Bokas, D., Aggelis, G. 2012. Improving fatty acid composition of lipids synthesized by *Brachionus plicatilis* in large scale experiments. *Journal of American Oil Chemists' Society* 89, 2047-2055.
- Biswas, G., Thirunavukkarasu, A.R., Sundaray, J.K., Kailasam, M., 2010. Optimization of feeding frequency of Asian seabass (*Lates calcarifer*) fry reared in net cages under brackishwater environment. *Aquaculture* 305, 26-31.

- Boehlert, G.W., Yoklavich, M.M., 1984. Carbon assimilation as a function of ingestion rate in larval Pacific Herring, *Clupea harengus pallasi* Valenciennes. J. Exp. Marine Biol. and Ecol. 79, 251-262.
- Canino, M.F., Bailey, K.M., 1995. Gut evacuation of walleye pollock larvae in response to feeding conditions. J. Fish Biol. 46, 389-403.
- Carrier, J.K., Watanabe, W.O., Harel, M., Rezek, T.C., Seaton, P.J., Shafer, T.H., 2011. Effects of dietary arachidonic acid on larval performance, fatty acid profiles, stress resistance, and expression of Na⁺/K⁺ ATPase mRNA in black sea bass *Centropristis striata*. Aquaculture 319, 111-121.
- Cho, S.H., Lim, Y.S., Lee, J.H., Lee, J.K., Park, S., Lee, S.M., 2003. Effects of feeding rate and feeding frequency on survival, growth, and body composition of ayu post-larvae *Plecoglossus altivelis*. J. World Aquac. Soc. 34, 85-91.
- Copeland, K.A., Watanabe, W.O., 2006. Light intensity effects on early life stages of black sea bass, *Centropristis striata* (Linnaeus 1758). Aquaculture Research 37, 1458-1463.
- Copeland, K.A., Watanabe, W.O., Dumas, C.F., 2005. Economic evaluation of a small-scale recirculating system for ongrowing of captive wild black sea bass *Centropristis striata* in North Carolina. J. World Aquac. Soc. 36, 489-497.
- Copeland, K.A., Watanabe, W.O., Carroll, P.M., Wheatley, K.S., Losordo, T.M., 2003. Growth and feed utilization of captive wild black sea bass *Centropristis striata* at four different densities in a recirculating tank system. Journal of the World Aquaculture Society 34, 300-307.

- Coutteau, P., Mourente, G., 1997. Lipid classes and their content of n-3 highly unsaturated fatty acids (HUFA) in *Artemia franciscana* after hatching, HUFA-enrichment and subsequent starvation. *Marine Biol.* 130, 81-91.
- Czesny, S., Rinchard, J., Abiado, M.A.G., Dabrowski, K. 2003. The effect of fasting, prolonged swimming, and predator presence on energy utilization and stress in juvenile walleye (*Stizostedion vitreum*). *Physiol. and Behav.*, 79, 597-603.
- Dhert, P., Rombaut, G., Suantika, G., Sorgeloos, P., 2001. Advancement of rotifer culture and manipulation techniques in Europe. *Aquaculture* 200, 129-146.
- Dumas, C.F., Wilde, J.D. 2009. Assessing metro niche market demand for North Carolina mariculture products. North Carolina Biotechnology Center Regional Development Grant Program. Grant No. 2008-RDG-4003 and North Carolina Sea Grant.
- Dwyer, K.S., Brown, J.A., Parrish, C., Lall, S.P., 2002. Feeding frequency affects food consumption, feeding pattern and growth of juvenile yellowtail flounder (*Limanda ferruginea*). *Aquaculture* 213, 279-292.
- Estevez, A., McEvoy, L.A., Bell, J.G., Sargent, J.R., 1998. Effects of temperature and starvation time on the pattern and rate of loss of essential fatty acids in *Artemia nauplii* previously enriched using arachidonic acid and eicosapentaenoic acid-rich emulsions. *Aquaculture*. 165, 295-311.
- Folch, J., Lees, M., Sloane-Stanley, G.H. 1957. A simple method for the isolation and purification of total lipids from animal tissues, *Journal of Biological Chemistry* 226, 497–509.

- Folkvord, A., Ottera, H., 1993. Effects of initial size distribution, day length and feeding frequency on growth, survival, and cannibalism in juvenile Atlantic cod (*Gadus morhua* L.). *Aquaculture* 114, 243-260.
- Goldan, O., Popper, D., Karplus, I., 1997. Management of size variation in juvenile gilthead sea bream (*Sparus aurata*). I: Particle size and frequency of feeding dry and live food. *Aquaculture* 152, 181-190.
- Han, K.M., Geurden, I., Sorgeloos, P., 2001. Fatty acid changes in enriched and subsequently starved *Artemia franciscana* nauplii enriched with different essential fatty acids. *Aquaculture* 199, 93-105.
- Hiroi, J., Kaneko, T., Seikai, T., Tanaka, M. 1998. Developmental sequence of chloride cells in the body skin and gills of Japanese flounder (*Paralichthys olivaceus*) larvae. *Zoological Science* 15, 455-460.
- Houde, E.D., Schekter, R.C., 1978. Simulated food patch and survival of larval bay anchovy, *Anchoa mitchilli*, and sea bream, *Archosargus rhomboidalis*. *Fish. Bull.* 76, 483-487.
- Izquierdo, M.S., Watanabe, T., Takeuchi, T., Arakawa, T. and Kitajima, C. 1989. Requirement of larval red seabream, *Pagrus major*, for essential fatty acids. *Nippon Suisan Gakkaishi* 55, 859-86.
- Johnston, T.A., Mathias, J.A., 1996. Gut evacuation and absorption efficiency of walleye larvae. *J. Fish Biol.* 49, 375-389.
- Kayano, Y., Yao, S., Yamamoto, S., Nakagawa, H. 1993. Effects of feeding frequency on the growth and body constituents of young red-spotted grouper *Epinephelus akaara*. *Aquaculture* 110, 271-278.

- Koven, W., Barr, Y., Lutzky, S., Ben-Atia, I., Weiss, R., Harel, M., Behrens, P., Tandler, A., 2001. The effect of dietary arachidonic acid (20 : 4n-6) on growth, survival and resistance to handling stress in gilthead seabream (*Sparus aurata*) larvae. *Aquaculture* 193, 107-122.
- Lavens, P. , Sorgeloos, P. 1996. Manual on the production and use of live food for aquaculture. FAO Fisheries Technical Paper No. 361. Rome, FAO. 295 p.
- Li, J.J., Liu, Y., Wang, W., Yang, Z., Chen, Y.F. 2011. Survival and Na⁺/K⁺ ATPase specific activity in response to acute salinity change in newly hatched obscure puffer, *Takifugu obscurus*, larvae. *J. World Aquac. Soc.* 42, 275-280.
- Li, J., Eygensteyn, J., Lock, R.A.C., Verbost, P.M., Vanderheijden, A.J.H., Bonga, S.E.W., Flik, G. 1995. Branchial chloride cell in larvae and juveniles of fresh-water tilapia *Oreochromis mossambicus*. *J. Experimental Biol.* 198, 2177-2184.
- Light, R.K., Portella, M.C., 2005. Effects of feeding frequency on larval rearing of trairao (*Hoplias lacerdae*). *Revista Brasileira De Zootecnia-Brazilian J. Animal Science.* 34, 1442-1448.
- Luo, L., Li, T., Xing, W., Xue, M., Ma, Z. Jiang, N. and Li, W. 2015. Effects of feeding rates and feeding frequency on growth performances of juvenile hybrid sturgeon, *Acipenser schrenckii* Brandt (female) x *A. baeri* Brandt (male). *Aquaculture* 448, 229-233.
- Matsuo, Y., Kasahara, Y., Hagiwara, A., Sakakura, Y., Arakawa, T., 2006. Evaluation of larval quality of viviparous scorpionfish *Sebastiscus marmoratus*. *Fish. Science.* 72, 948-954.
- McCormick, S.D. 1993. Methods for nonlethal gill biopsy and measurements of Na⁺, K⁺ ATPase activity. *Can J. Fish. Aquatic Sciences* 50, 656-658.

- McCormick, S.D., Bern, H.A. 1989. *In vitro* stimulation of Na⁺, K⁺ - ATPase activity and ouabain binding by cortisol in coho salmon gill. *Am. J. Physiol.* 256, R707-R715.
- Mollah, M. F. A, Tan, E. S. P. 1988. Effects of feeding frequency on the growth and survival of catfish (*Clarias macrocephalus*) larvae. *Bangladesh J. Fish.* 11.2, 9-14
- Moustakas, C. T., Watanabe, W.O., Copeland, K. A. 2004. Combined effects of photoperiod and salinity on growth, survival, and osmoregulatory ability of southern flounder *Paralichthys lethostigma*. *Aquaculture* 229, 159–179.
- Munk, P., 1995. Foraging behavior of larval cod *Gadus morhua* influenced by prey density and hunger. *Marine Biol.*, 122, 205-212.
- Munk, P., Kiorboe, T., 1985. Feeding behavior and swimming activity of larval herring *Clupea harengus* in relation to density of copepod nauplii. *Marine Ecol. Progress Series.* 24, 15-21.
- Penefsky, H.S., M.F. Bruist. 1984. Adenosinetriphosphatases, p. 324-328. *In* H.U. Bergmeyer [ed.] *Methods of enzymatic analysis.* vol. IV. Cerlag Chemie, Beinhem.
- Rabe, J., Brown, J.A., 2000. A pulse feeding strategy for rearing larval fish: an experiment with yellowtail flounder. *Aquaculture* 191, 289-302.
- Rainuzzo, J.R., Olsen, Y., Rosenlund, G., 1989. The effect of enrichment diets on the fatty-acid composition of the rotifer *Brachionus plicatilis*. *Aquaculture* 79, 157-161.
- Rezek, T.C., Watanabe, W.O., Harel, M., Seaton, P.J., 2010. Effects of dietary docosahexaenoic acid (22:6n-3) and arachidonic acid (20:4n-6) on the growth, survival, stress resistance and fatty acid composition in black sea bass *Centropristis striata* (Linnaeus 1758) larvae. *Aquaculture Research* 41, 1302-1314.

- Roo, J., Hernandez-Cruz, C.M., Borrero, C., Schuchardt, D., Fernandez-Palacios, H., 2010. Effect of larval density and feeding sequence on meagre (*Argyrosomus regius*; Asso, 1801) larval rearing. *Aquaculture*. 302, 82-88.
- Sargent, J., Bell, G., McEvoy, L., Tocher, D., Estevez, A., 1999. Recent developments in the essential fatty acid nutrition of fish. *Aquaculture* 177, 191-199.
- Secor, S.M. 2011. Food acquisition and digestion: Cost of digestion and assimilation. In: Encyclopedia of Fish Physiol. (Editor, Farrell, A.P.). Academic Press, San Diego, pp. 1608-1616.
- Sedberry, G.R., 1988. Food and feeding of black sea bass *Centropristis striata*, in live bottom habitats in the South Atlantic Bight. The Journal of Elisha Mitchell Scientific Society, 35-50.
- Slembrouck, J., Baras, E., Subagja, J., Hung, L.T., Legendre, M., 2009. Survival, growth and food conversion of cultured larvae of *Pangasianodon hypophthalmus*, depending on feeding level, prey density and fish density. *Aquaculture* 294, 52-59.
- Sorgeloos, P., Dhert, P., Candreva, P., 2001. Use of brine shrimp, *Artemia* spp., in marine fish larviculture. *Aquaculture* 200, 147-159.
- Støttrup, J. G., Attramadal, Y. 1992. The influence of different rotifer and *Artemia* enrichment diets on growth, survival and pigmentation in Turbot (*Scophthalmus maximus* L.) larvae. *J. World Aquac. Soc.* 23, 307–316.
- Varsamos, S., Nebel, C., Charmantier, G. 2005. Ontogeny of osmoregulation in postembryonic fish: A review. *Comp. Biochem. Physiol. a- Molec. and Integ. Physiol.* 141, 401-429.
- Varsamos, S., Bonga, S.E.W., Charmantier, G., Flik, C. 2004. Drinking and Na⁺/K⁺ ATPase activity during early development of European sea bass, *Dicentrarchus labrax* -

- Ontogeny and short-term regulation following acute salinity changes. J. Exper. Marine Biol. Ecol. 311, 189-200.
- Varsamos, S., Diaz, J.P., Charmantier, G., Blasco, C., Connes, R., Flik, G. 2002. Location and morphology of chloride cells during the post-embryonic development of the european sea bass, *Dicentrarchus labrax*. Anat. Embry. 205, 203-213.
- Vijayan, M.M., Morgan, J.D., Sakamoto, T., Grau, E.G., Iwama, G.K. 1996. Food-deprivation affects seawater acclimation in tilapia: Hormonal and metabolic changes. J. Exper. Biol., 199, 2467-2475.
- Watanabe, W.O., 2011. Species profile: black sea bass. SRAC publication no. 7207. Southern Regional Aquaculture Center, United States Department of Aquaculture, Stoneville, MS, pp. 1–16.
- Watanabe, T., Kitajima, C., Fujita, S., 1983. Nutritional values of live organisms used in Japan for mass propagation of fish: A review. Aquaculture 34, 115-143.
- Watanabe, W.O., Kuo, C-M., Huang, M-C. 1985. The ontogeny of salinity tolerance in the tilapias *Oreochromis niloticus*, *O. aureus*, and *O. mossambicus* x *O. niloticus* hybrid, spawned and hatched in freshwater. Aquaculture 47: 353-367.
- Watanabe, W.O., Dumas, C.F., Carroll, P.M. and Resimius, C.R. 2015. Production economic analysis of black sea bass juveniles to support finfish mariculture growout industry development in the Southeastern U.S. Aquaculture Economics and Management 19:2, 226-250. <http://www.tandfonline.com/doi/full/10.1080/13657305.2015.1024347>
- Watanabe, W.O., Rezek, T.C., Harel, M., Seaton, P.J. 2014. The effects of dietary arachidonic acid (20:4n-6) on growth, survival and hypersaline stress resistance in larval southern flounder *Paralichthys lethostigma* (Jordan and Gilbert 1884). Chapter 1 (p. 1-27): In

- “Arachidonic Acid: Sources, Biosynthesis and Health Effects”, J.M. O’Keefe, Editor.
Nova Science Publishers Inc. ISBN: 978-1-63117-619-7.
- Watanabe, T., Izquierdo, M.S., Takeuchi, T., Satoh, S.Y. and Kitajima, C. 1989. Comparison between eicosapentaenoic and docosahexaenoic acids in terms of essential fatty acid efficacy in larval red seabream. *Bulletin Japanese Society Scientific Fisheries* 55, 1635-1640.
- Watanabe, W.O., Alam, M.S., Ostrowski, A.D., Montgomery, F.A., Gabel J.E., Morris, J.A. Jr, and Seaton, P.J. 2016. Live prey enrichment and artificial microdiets for larviculture of Atlantic red porgy *Pagrus pagrus*. *Aquaculture Reports* 3, 93-107.
- Watanabe, W.O., Smith, T.I.J., Berlinsky, D.L, Woolridge, C.A., Stuart, K.R., Copeland, K.A., Denson, M.R. 2003. Volitional spawning of black sea bass *Centropristis striata* induced with pelleted luteinizing hormone releasing hormone-analogue. *Journal of the World Aquaculture Society* 34, 319-331.
- Wenner, C.A., Roumillat, W.A., Waltz, C.W., 1986. Contributions to the life-history of black sea bass, *Centropristis striata*, off the Southeastern United States. *Fish. Bull.*, 84, 723-741.
- Wilde, J.D., Dumas, C.F., Watanabe, W.O. 2008. Analyses of the niche restaurant market for farm-raised black sea bass *Centropristis striata* in North Carolina. *World Aquaculture* 2008, Orlando, Florida. MS Thesis. University of North Carolina Wilmington, Center for Marine Science.
- Wittenrich, M.L., Rhody, N.R., Turingan, R.G., Main, K.L., 2009. Coupling osteological development of the feeding apparatus with feeding performance in common snook,

- Centropomus undecimalis, larvae: Identifying morphological constraints to feeding. Aquaculture 294, 221-227.
- Woolley, L.D., Partridge, G.J. 2015. The effect of different rotifer feeding regimes on the growth and survival of yellowtail kingfish *Seriola lalandi* (Valenciennes, 1833) larvae. Aquaculture Research, In press .
- Xie, F., Ai, Q., Mai, K., Xu, W., Ma, H., 2011. The optimal feeding frequency of large yellow croaker (*Pseudosciaena crocea*, Richardson) larvae. Aquaculture. 311, 162-167.
- Yao, S.J., Umino, T., Nakagawa, H., 1994. Effect of feeding frequency on lipid-accumulation in ayu. Fish. Science 60, 667-671.
- Zaugg, W.W. 1982. A simplified preparation for adenosine triphosphatase determination in gill tissue. Can. J. Fish. Aquat. sci. 39: 215-217.

Figure 1. Growth in notochord length (mm) (mean $\pm$ SEM, $N = 6$, except on d4ph when $N = 4$) of larval black sea bass fed enriched rotifers and *Artemia* at different feeding frequencies (1x/d, 2x/d, 3x/d, 4x/d) from d4ph to d31ph. Asterisks indicate significant (* $P < 0.05$, ** $P < 0.01$) treatment differences for that sampling date.

Figure 2. Notochord length (mm) (mean $\pm$ SEM, $N = 6$) of black sea bass larvae fed enriched rotifers and *Artemia* at different feeding frequencies (1x/d, 2x/d, 3x/d, 4x/d) on d31ph. Means not sharing the same letter are significantly ($P < 0.05$) different.

Figure 3. Growth in wet weight (mg) (mean $\pm$ SEM, $N = 6$, except on d4ph when $N = 4$) of black sea bass larvae fed enriched rotifers and *Artemia* at different feeding frequencies (1x/d, 2x/d, 3x/d, 4x/d) from d4ph to d31ph. Asterisks indicate significant (* $P < 0.05$) treatment differences for that sampling date.

Figure 4. Wet weight (mg) (mean $\pm$ SEM, $N = 6$) of black sea bass larvae fed enriched rotifers and *Artemia* at different feeding frequencies (1x/d, 2x/d, 3x/d, 4x/d) on d31ph. Treatments not sharing the same letter are significantly ($P < 0.05$) different.

Figures 5. Mean rotifer density (individuals/mL) ($N = 3$) sampled *in situ* on d12ph over a 12 h sampling period. Rotifers were fed at 08:00 (1x/d), 08:00 and 16:00 h (2x/d), 08:00, 12:00 and 16:00 h (3x/d), and 08:00, 12:00, 16:00 and 20:00 h (4x/d). The gradual increases in rotifer densities preceding feeding at 12:00, 16:00 and 20:00 h are artifacts of the time interval between samples.

Figure 6. Mean *Artemia* density (individuals/mL) ($N = 3$) sampled *in situ* on d27ph over a 12 h sampling period. *Artemia* were fed at 08:00 (1x/d), 08:00 and 16:00 h (2x/d), 08:00, 12:00 and 16:00 h (3x/d), and 08:00, 12:00, 16:00 and 20:00 h (4x/d).

The gradual increases in rotifer densities preceding feeding at 12:00, 16:00 and 20:00 h are artifacts of the time interval between samples.

Figure 7. Na^+/K^+ ATPase activity ($\mu\text{mol ADP/mg protein/hr}$, mean $\pm$ SEM, $N = 6$) of whole body homogenates of d32ph black sea bass larvae fed live prey (rotifers and *Artemia*) at different feeding frequencies (1x/d, 2x/d, 3x/d, 4x/d) following abrupt transfer from 32 g/L to a sub-lethal salinity of 15 g/L. Data for all sampling times (0, 12, 24, and 48 h post-transfer) were combined.

Table 1. Dry weights (mg) (mean $\pm$ SEM, $N = 6$) of black sea bass larvae fed enriched rotifers and *Artemia* at four different feeding frequencies (1x/d, 2x/d, 3x/d, 4x/d) from d4ph to d31ph.

Age (dph)	Feeding Frequency (x/d)				<i>P</i> *
	1	2	3	4	
4	ND	ND	ND	ND	
10	0.05 $\pm$ 0.01	0.06 $\pm$ 0.05	0.06 $\pm$ 0.01	0.07 $\pm$ 0.01	NS
17	0.13 $\pm$ 0.02	0.26 $\pm$ 0.13	0.13 $\pm$ 0.08	0.15 $\pm$ 0.01	NS
25	0.22 $\pm$ 0.03	0.26 $\pm$ 0.01	0.28 $\pm$ 0.02	0.29 $\pm$ 0.02	NS
31	0.55 $\pm$ 0.03	0.69 $\pm$ 0.16	0.80 $\pm$ 0.12	0.96 $\pm$ 0.12	NS

ND = no data

*Probability level of overall ANOVA. NS= not significant.

Table 2. Specific growth rate of wet weight (SGR, %/d) and survival activity index (SAI) (mean $\pm$ SEM, $N = 6$) for black sea bass larvae fed enriched rotifers and *Artemia* at four different feeding frequencies (1x/d, 2x/d, 3x/d, 4x/d) from d4ph to d31ph.

	Feeding Frequency (x/d)				
	1	2	3	4	P^*
SGR ^x (%/d)	18.1 $\pm$ 0.204 ^b	18.4 $\pm$ 0.399 ^b	19.3 $\pm$ 0.507 ^{ab}	20.1 $\pm$ 0.467 ^a	0.01
SAI ^y	244.5 $\pm$ 55.3 ^b	526.5 $\pm$ 48.32 ^a	498.0 $\pm$ 40.23 ^a	491.9 $\pm$ 102.27 ^a	0.05

^x SGR calculated from d4ph to d31ph.

^y SAI calculated following an acute hyposalinity (10 g/L) challenge on d32ph.

*Probability level of overall ANOVA. NS = not significant. Means in the same row not sharing the same letter are significantly different (Tukey-Kramer HSD Test).

Table 3. Survival (%) (mean $\pm$ SEM, $N = 6$) for black sea bass larvae fed enriched rotifers and *Artemia* at four different feeding frequencies (1x/d, 2x/d, 3x/d, 4x/d) from d0ph to d31ph.

Age (dph)	Feeding Frequency (x/d)				<i>P</i> *
	1	2	3	4	
10	80.7 $\pm$ 7.87	87.5 $\pm$ 5.82	92.8 $\pm$ 4.67	90.5 $\pm$ 6.12	NS
17	81.1 $\pm$ 6.10	83.7 $\pm$ 6.37	88.7 $\pm$ 5.47	89.6 $\pm$ 5.20	NS
25	40.1 $\pm$ 5.09	51.4 $\pm$ 4.60	59.4 $\pm$ 8.77	52.9 $\pm$ 6.47	NS
31	55.3 $\pm$ 8.14	52.1 $\pm$ 5.77	53.5 $\pm$ 7.82	49.8 $\pm$ 5.70	NS
35	32.5 $\pm$ 3.31	36.8 $\pm$ 6.23	32.7 $\pm$ 5.70	23.4 $\pm$ 4.69	NS

*Probability level of overall ANOVA. NS = not significant. Means in the same row not sharing the same letter are significantly different (Tukey-Kramer HSD Test).

Table 4. Fatty acid composition (% TFA, mean $\pm$ SEM, $N = 3$) of enriched rotifers *Brachionus rotundiformis* harvested at 08:00 h and then stored at 4 °C (“cold-banked”) until 20:00 h. No significant differences among storage times were observed.

Fatty acids	Time of sampling after storage			
	08:00 h	12:00 h	16:00 h	20:00 h
14:0	3.48 $\pm$ 0.01	3.50 $\pm$ 0.12	3.30 $\pm$ 0.07	3.56 $\pm$ 0.07
16:0	22.0 $\pm$ 0.30	22.1 $\pm$ 0.29	21.8 $\pm$ 0.21	21.8 $\pm$ 0.20
16:1n-7	11.8 $\pm$ 0.38	12.1 $\pm$ 0.44	11.6 $\pm$ 0.65	12.1 $\pm$ 0.67
18:0	5.49 $\pm$ 0.20	5.68 $\pm$ 0.17	5.62 $\pm$ 0.32	5.69 $\pm$ 0.12
18:1n-9	4.70 $\pm$ 0.24	4.84 $\pm$ 0.28	4.58 $\pm$ 0.33	5.58 $\pm$ 0.39
18:2n-6	5.57 $\pm$ 0.66	5.43 $\pm$ 0.80	5.29 $\pm$ 0.73	4.11 $\pm$ 0.54
18:3n-3	12.4 $\pm$ 0.75	11.4 $\pm$ 0.42	12.0 $\pm$ 0.42	11.2 $\pm$ 0.28
18:4n-3	1.95 $\pm$ 0.55	2.00 $\pm$ 0.68	1.54 $\pm$ 0.37	2.44 $\pm$ 0.73
20:1	3.34 $\pm$ 0.44	2.83 $\pm$ 0.10	2.79 $\pm$ 0.06	2.68 $\pm$ 0.08
20:4n-6	2.81 $\pm$ 0.12	2.65 $\pm$ 0.04	2.40 $\pm$ 0.52	2.95 $\pm$ 0.21
20:5n-3	15.4 $\pm$ 0.61	15.5 $\pm$ 0.41	15.4 $\pm$ 0.56	14.7 $\pm$ 0.69
22:5n-3	5.02 $\pm$ 0.18	5.83 $\pm$ 0.71	5.94 $\pm$ 0.81	6.81 $\pm$ 0.22
22:6n-3	5.00 $\pm$ 0.35	6.08 $\pm$ 0.98	8.18 $\pm$ 1.72	6.29 $\pm$ 1.15
Σ Saturates	31.0 $\pm$ 0.51	31.3 $\pm$ 0.35	30.73 $\pm$ 0.36	31.0 $\pm$ 0.20
Σ Monounsaturates	19.9 $\pm$ 0.40	19.8 $\pm$ 0.25	18.9 $\pm$ 0.42	20.4 $\pm$ 0.61
Σ n-3 PUFA	40.8 $\pm$ 0.56	40.9 $\pm$ 1.39	42.6 $\pm$ 1.40	41.5 $\pm$ 1.11
Σ n-6 PUFA	8.39 $\pm$ 0.66	8.08 $\pm$ 0.84	7.69 $\pm$ 0.94	7.06 $\pm$ 0.74
n-3 / n-6 PUFA	4.93 $\pm$ 0.40	5.28 $\pm$ 0.76	5.73 $\pm$ 0.77	6.05 $\pm$ 0.83
DHA/EPA	0.32 $\pm$ 0.01	0.40 $\pm$ 0.07	0.54 $\pm$ 0.12	0.44 $\pm$ 0.09
EPA/ARA	5.50 $\pm$ 0.33	5.85 $\pm$ 0.13	7.19 $\pm$ 1.85	5.02 $\pm$ 0.28
DHA/ARA	1.80 $\pm$ 0.17	2.30 $\pm$ 0.41	4.25 $\pm$ 1.96	2.20 $\pm$ 0.51
Total lipid (mg/g dwt)	309.8 $\pm$ 17.6	375.0 $\pm$ 48.8	271.7 $\pm$ 26.7	291.2 $\pm$ 18.1

Table 5. Fatty acid composition (% TFA, mean $\pm$ SEM, $N = 3$) of enriched *Artemia* cold-banked at $\sim 12^{\circ}\text{C}$ from 08:00 h until 20:00 h. No significant differences among storage times were observed.

Fatty acids	Time of sampling after storage			
	08:00h	12:00h	16:00h	20:00
14:0	1.13 $\pm$ 0.01	1.15 $\pm$ 0.06	1.04 $\pm$ 0.13	1.12 $\pm$ 0.04
16:0	10.7 $\pm$ 0.31	10.4 $\pm$ 0.28	10.2 $\pm$ 0.79	11.312 $\pm$ 0.54
16:1n-7	4.30 $\pm$ 0.03	4.33 $\pm$ 0.08	4.2 $\pm$ 0.20	4.12 $\pm$ 0.19
17:0	0.56 $\pm$ 0.03	0.54 $\pm$ 0.04	0.83 $\pm$ 0.26	0.67 $\pm$ 0.08
18:0	4.98 $\pm$ 0.18	4.54 $\pm$ 0.27	4.8 $\pm$ 0.25	4.95 $\pm$ 0.08
18:1n-9	22.4 $\pm$ 2.89	27.1 $\pm$ 3.88	22.4 $\pm$ 3.19	26.4 $\pm$ 4.04
18:1n-11	5.01 $\pm$ 0.05	4.48 $\pm$ 0.28	6.02 $\pm$ 1.84	5.02 $\pm$ 0.63
18:2n-6	7.28 $\pm$ 0.15	6.9 $\pm$ 0.18	6.72 $\pm$ 0.42	6.22 $\pm$ 0.35
18:3n-3	16.7 $\pm$ 3.87	14.5 $\pm$ 2.28	19.0 $\pm$ 1.88	16.2 $\pm$ 2.56
18:4n-3	3.32 $\pm$ 0.14	3.06 $\pm$ 0.22	3.00 $\pm$ 0.18	3.29 $\pm$ 0.17
20:1	2.44 $\pm$ 0.13	2.55 $\pm$ 0.13	2.89 $\pm$ 0.13	2.65 $\pm$ 0.13
20:4n-6	1.34 $\pm$ 0.09	1.06 $\pm$ 0.18	1.03 $\pm$ 0.05	0.96 $\pm$ 0.13
20:5n-3	8.26 $\pm$ 0.47	7.64 $\pm$ 0.44	7.73 $\pm$ 0.24	7.1 $\pm$ 0.33
22:1	0.84 $\pm$ 0.07	0.95 $\pm$ 0.08	1.00 $\pm$ 0.05	0.91 $\pm$ 0.04
22:5n-3	1.18 $\pm$ 0.42	0.95 $\pm$ 0.07	1.10 $\pm$ 0.18	0.86 $\pm$ 0.14
22:6n-3	9.56 $\pm$ 0.34	8.14 $\pm$ 0.66	9.27 $\pm$ 0.11	8.29 $\pm$ 0.56
DHA/EPA	1.16 $\pm$ 0.03	1.08 $\pm$ 0.12	1.20 $\pm$ 0.03	1.17 $\pm$ 0.03
EPA/ARA	6.19 $\pm$ 0.11	7.51 $\pm$ 0.81	7.73 $\pm$ 0.57	7.80 $\pm$ 1.59
DHA/ARA	7.19 $\pm$ 0.31	8.29 $\pm$ 1.77	9.12 $\pm$ 0.48	9.15 $\pm$ 2.00
Σ Saturates	17.4 $\pm$ 0.46	16.6 $\pm$ 0.51	16.9 $\pm$ 0.74	18.1 $\pm$ 0.58
Σ Monounsaturates	35.0 $\pm$ 2.86	39.5 $\pm$ 3.94	36.6 $\pm$ 2.05	39.1 $\pm$ 3.26
Σ n-3 PUFA	39.0 $\pm$ 3.07	34.2 $\pm$ 1.75	39.1 $\pm$ 1.65	35.7 $\pm$ 3.25
Σ n-6 PUFA	25.0 $\pm$ 3.40	22.1 $\pm$ 2.53	25.7 $\pm$ 1.64	23.3 $\pm$ 2.77
n-3 / n-6 PUFA	1.59 $\pm$ 0.09	1.58 $\pm$ 0.11	1.53 $\pm$ 0.04	1.55 $\pm$ 0.05
Total lipid (mg/g dwt)	570.6 $\pm$ 132.1	500.5 $\pm$ 77.5	303.6 $\pm$ 77.3	352.7 $\pm$ 20.1

Table 6. Fatty acid profile (% total fatty acid) (mean $\pm$ SEM, $N = 5$, except treatment 3, $N = 4$) of whole body tissue of black sea bass *Centropomus striata* larvae (d32ph) fed enriched rotifers *Brachionus rotundiformis* and *Artemia* at four different feeding frequency frequencies (1x/d, 2x/d, 3x/d, 4x/d).

Fatty acids	Feeding Frequency (x/d)			
	1	2	3	4
16:0	16.7 $\pm$ 0.48	15.3 $\pm$ 0.99	15.9 $\pm$ 0.33	14.7 $\pm$ 0.94
16:1n-7	2.27 $\pm$ 0.08	2.63 $\pm$ 0.21	2.54 $\pm$ 0.13	2.51 $\pm$ 0.06
17:0	0.91 $\pm$ 0.05	0.89 $\pm$ 0.04	0.85 $\pm$ 0.02	0.83 $\pm$ 0.03
18:0	10.7 $\pm$ 0.54	10.1 $\pm$ 0.22	9.49 $\pm$ 0.23	9.43 $\pm$ 0.26
18:1n-9	19.1 $\pm$ 0.55	19.4 $\pm$ 1.03	19.2 $\pm$ 1.37	19.6 $\pm$ 0.99
18:1n-11	8.05 $\pm$ 0.56	7.71 $\pm$ 0.68	6.90 $\pm$ 0.51	7.05 $\pm$ 0.29
18:2n-6	5.29 $\pm$ 0.25	5.61 $\pm$ 0.38	6.32 $\pm$ 0.47	6.22 $\pm$ 0.68
18:3n-3	12.5 $\pm$ 0.89	13.1 $\pm$ 0.66	13.6 $\pm$ 1.10	14.2 $\pm$ 0.42
18:4n-3	1.54 $\pm$ 0.90	1.73 $\pm$ 0.10	1.68 $\pm$ 0.12	1.96 $\pm$ 0.16
20:1	1.99 $\pm$ 0.06	2.11 $\pm$ 0.05	2.15 $\pm$ 0.03	2.25 $\pm$ 0.14
20:4n-6	2.40 $\pm$ 0.09	2.49 $\pm$ 0.08	2.29 $\pm$ 0.07	2.24 $\pm$ 0.15
20:5n-3	7.50 $\pm$ 0.10	7.79 $\pm$ 0.07	7.75 $\pm$ 0.103	7.78 $\pm$ 0.11
22:5n-3	1.86 $\pm$ 0.21	1.94 $\pm$ 0.06	1.81 $\pm$ 0.05	1.73 $\pm$ 0.08
22:6n-3	9.27 $\pm$ 0.57	9.30 $\pm$ 0.20	9.57 $\pm$ 0.29	9.45 $\pm$ 0.41
Σ Saturates	28.2 $\pm$ 0.95	26.2 $\pm$ 0.98	26.2 $\pm$ 0.54	24.9 $\pm$ 0.89
Σ Monounsaturates	33.3 $\pm$ 0.98	33.8 $\pm$ 1.21	32.6 $\pm$ 1.65	33.1 $\pm$ 0.96
Σ n-3 PUFA	36.8 $\pm$ 0.47	36.03 $\pm$ 0.90	36.7 $\pm$ 0.23	35.7 $\pm$ 1.07
Σ n-6 PUFA	7.69 $\pm$ 0.30	8.1 $\pm$ 0.39	8.61 $\pm$ 0.53	8.46 $\pm$ 0.63
n-3 / n-6 PUFA	4.81 $\pm$ 0.17	4.48 $\pm$ 0.16	4.301 $\pm$ 0.22	4.29 $\pm$ 0.31
DHA/EPA	1.23 $\pm$ 0.06	1.19 $\pm$ 0.03	1.24 $\pm$ 0.02	1.21 $\pm$ 0.04
EPA/ARA	3.14 $\pm$ 0.12	3.15 $\pm$ 0.08	3.39 $\pm$ 0.07	3.53 $\pm$ 0.21
DHA/ARA	3.87 $\pm$ 0.24	3.76 $\pm$ 0.19	4.18 $\pm$ 0.04	4.30 $\pm$ 0.35
Total lipid (mg/g dwt)	211.0 $\pm$ 21.1	224.7 $\pm$ 19.1	250.6 $\pm$ 22.5	199.6 $\pm$ 36.9