**Microbial toxicity and characterization of DNAN (bio)transformation product
mixtures**

Christopher I. Olivares¹, Reyes Sierra-Alvarez¹, Cristina Alvarez-Nieto¹, Leif
Abrell^{2,3}, Jon Chorover² and Jim A. Field^{1*}

¹ Department of Chemical and Environmental Engineering, University of Arizona,
Tucson, AZ, USA 85721

² Department of Soil, Water & Environmental Science, University of Arizona,
Tucson, AZ, USA 85721

³ Department of Chemistry & Biochemistry, University of Arizona, Tucson, AZ, USA 85721

Corresponding author:

*Jim A. Field (jimfield@email.arizona.edu)

Word count: 4170 words (intro-acknowledgments) + 5 figures/ 2 tables (300 words
each) = 6270 word equivalents

20 **Abstract (245 words)**

21 2,4-Dinitroanisole (DNAN) is an emerging insensitive munitions compound. It undergoes
22 rapid (bio)transformation in soils and anaerobic sludge. The primary transformation pathway
23 catalyzed by a combination of biotic and abiotic factors is nitrogroup reduction followed by
24 coupling of reactive intermediates to form azo-dimers. Additional pathways include N-
25 acetylation and O-demethoxylation. Toxicity due to (bio)transformation products of DNAN
26 has received little attention. In this study, the toxicity of DNAN (bio)transformation
27 monomer products and azo-dimer and trimer surrogates to acetoclastic methanogens and the
28 marine bioluminescent bacterium, *Allivibrio fischeri*, were evaluated. Methanogens were
29 severely inhibited by 3-nitro-4-methoxyaniline (MENA), with a 50%-inhibiting concentration
30 (IC_{50}) of 25 μ M, which is more toxic than DNAN with the same assay, but posed a lower
31 toxicity to *A. fischeri* (IC_{50} = 219 μ M). On the other hand, N-(5-amino-2-methoxyphenyl)
32 acetamide (Ac-DAAN) was the least inhibitory test-compound for both microbial targets.
33 Azo-dimer and trimer surrogates were very highly toxic to both microbial systems, with a
34 toxicity similar or stronger than that of DNAN. A semi-quantitative LC-QTOF-MS method
35 was employed to determine product mixture profiles at different stages of biotransformation,
36 and compared with the microbial toxicity of the product-mixtures formed. Methanogenic
37 toxicity increased due to putative reactive nitroso-intermediates as DNAN was reduced.
38 However, the inhibition later attenuated as dimers became the predominant products in the
39 mixtures. In contrast, *A. fischeri* tolerated the initial biotransformation products but were
40 highly inhibited by the predominant azo-dimer products formed at longer incubation times,
41 suggesting these ultimate products are more toxic than DNAN.

42

43 **Keywords (6 max):** munitions, 2,4-dinitroanisol, biotransformation, microbial toxicity,
44 dimers, liquid chromatography quadrupole time-of-flight mass spectrometry

45

46 **Graphical abstract**

47

48

49 **1. Introduction**

50 2,4-dinitroanisole (DNAN) is an insensitive munitions compound that is less prone to
51 accidental explosions and has emerged as an important replacement of conventional
52 explosives, such as 2,4,6-trinitrotoluene (TNT) (Davies et al., 2006). Once released to the
53 environment, DNAN is readily (bio)transformed via nitro-reduction in natural and engineered
54 systems by a combination of abiotic (Ahn et al., 2011; Niedzwiecka et al., 2013; Hawari et
55 al., 2015; Linker et al., 2015; Olivares et al., 2016) and biotic processes (Platten et al., 2010;
56 Olivares et al., 2013; Olivares et al., 2016). The initial nitro-reduction occurs with a dominant
57 regioselectivity towards the *ortho* nitro group (Platten et al., 2010; Perreault et al., 2012;
58 Hawari et al., 2015; Olivares et al., 2016), yielding 2-methoxy-5-nitroaniline (MENA), but its
59 regioisomer 3-methoxy-4-nitroaniline (iMENA) has been found in few studies (Schroer et al.,
60 2015; Olivares et al., 2016). Continued nitro-reduction leads to the aromatic amine 2,4-
61 diaminoanisole (DAAN) and reactive intermediates that form azo-dimers (Platten et al.,
62 2010; Olivares et al., 2013; Olivares et al., 2016). Parallel anaerobic (bio)transformation
63 pathways have been reported, such as N-acetylation of products of nitrogroup reduction
64 (Perreault et al., 2012; Olivares et al., 2013), as well as N-methylation and step-wise O-
65 demethoxylation of azo-dimers (Olivares et al., 2013; Olivares et al., 2016), creating a large
66 array of possible products.

67 As DNAN is transformed, its products may pose an environmental hazard yet to be
68 characterized. Most toxicity studies have been focused on the parent compound, DNAN
69 (Dodard et al., 2013; Kennedy et al., 2015), but little attention has been given to its
70 (bio)transformation products. While there has been an initial characterization of the microbial
71 toxicity of the primary DNAN nitrogroup-reduction-products (Liang et al., 2013), the toxicity
72 posed by the majority of transformation products is unknown.

Therefore, the occurrence of (bio)transformation products should be assessed and put in context with their toxicity potential. The objectives of this work were: (1) to assay microbial toxicity of individual transformation products and/or best available surrogate compounds to acetoclastic methanogens and the marine bacterium *Allivibrio fischeri*; and (2) to determine biotransformation product profiles for DNAN incubated anaerobically in soils and anaerobic sludge and link temporal changes in product mixture composition to toxicity changes during transformation using acetoclastic methanogens and *A. fischeri*, as microbial toxicity targets.

2. Materials and Methods

2.1 Chemicals and biological materials

Fig. 1 shows the chemical structures of monomer DNAN (bio)transformation products and azo-dimer/trimer surrogates assayed. 3-nitro-4-methoxyaniline (CAS# 577-72-0, denoted “iMENA”, purity 97%) was procured from Accela ChemBio Inc. (San Diego, CA, USA). N-(5-amino-2-methoxyphenyl) acetamide (CAS# 64353-88-4, denoted “Ac-DAAN”, purity 95%) was purchased from ChemBridge Corporation (San Diego, CA, USA). 2,2'-dimethoxy-4,4'-azodianiline (CAS# 6364-31-4, denoted “dimer L”, purity >90%) was acquired from MolMall Sarl (Lonay, Switzerland). Bismarck Brown Y (*m*-Bis(2,4-diaminophenylazo)-benzene, CAS# 8005-77-4, denoted “BBY”, dye purity 46%) was obtained from Chem-Impex International (Wood Dale, IL, USA). Camp Navajo soil (water content = 9.3%) and Camp Butner soil (water content = 20.7%), described in (Krzmarzick et al., 2015; Olivares et al., 2016), were sieved (2 mm) and stored at 4 °C before use. Granular sludge (dry weight (dwt) solids = 11.2%, volatile suspended solids (VSS) = 7.9% per wet weight) was obtained from a full-scale upflow anaerobic sludge blanket reactor located at an industrial brewery

wastewater bioreactor (Mahou, Guadalajara, Spain) and was used for (bio)transformation assays and methanogenic inhibition toxicity. The marine bioluminescent bacterium *A. fischeri* (lyophilized culture NRRL-B-11177, Modern Water Inc., New Castle, DE, USA) was also used in microbial toxicity assays.

2.2 Staggered DNAN soil/sludge (bio)transformation assays

DNAN was incubated anaerobically in soil or sludge microcosms to obtain product mixtures at different (bio)transformation stages. DNAN (500 μ M final concentration) was added to 10 mL of mineral medium (Olivares et al., 2013) (pH =7.2, 18 mM phosphate buffer) amended with 10 mM pyruvate in anaerobic tubes (Bellco Glass Inc., Vineland, NJ, USA). The solutions were inoculated with soil (100 mg wet) or sludge (75 mg wet). The headspace was flushed with He/CO₂ (80/20 %). The tubes were sealed with t-butyl caps and aluminum seals, and incubated in the dark at 30° C in an orbital shaker (115 rpm). Tubes were prepared at different times and incubated immediately so that when samples were collected on the same day, the overall (bio)transformation time elapsed would be 0, 1, 5, 10, 20, 30, 40, or 50 days of anaerobic incubation. Sampling was performed in an anaerobic hood to minimize oxidation of reactive products with oxygen. Samples were centrifuged (10 min, 9,600×g), sealed inside the anaerobic chamber, and kept at 4 °C before methanogenic and *A. fischeri* toxicity assays. All experiments were performed in duplicate. A diagram depicting the staggered (bio)transformation assays and overall workflow is shown in the Supplementary Material, Fig. S-1.

2.3 Microbial toxicity assays

2.3.1 Acetoclastic methanogens

The toxicity of (bio)transformation products/surrogates to acetoclastic methanogenesis was assayed in a methanogenic consortia immobilized in anaerobic granular sludge. The assays were performed in 160 mL serum bottles supplemented with 25 mL basal mineral medium at pH 7.2 (Liang et al., 2013), inoculated with 1.5 g VSS L⁻¹ of anaerobic granular sludge (7.9% VSS of wet weight), and amended with sodium acetate as substrate (2.6 mM). Serum bottles were flushed with He/CO₂ (80/20% v/v) and crimped with aluminum seals and t-butyl rubber septa. The bottles were incubated overnight, and flushed with He/CO₂ (80/20% v/v) before addition of the individual test-compounds: iMENA (10-200 µM), Ac-DAAN (80-8,000 µM), dimer L (4-83 µM), and BBY (0.12-1.92 µM). In addition, mixtures of biotransformation products from a starting DNAN concentration of 500 µM diluted 54-fold in aqueous solutions were utilized in the methanogenic toxicity bioassays. The dilution was chosen based on preliminary experiments to avoid complete methanogenesis inhibition that would prevent toxicity changes to be detected. Separate samples of these aqueous solutions containing mixtures of biotransformation products were harvested at different stages of DNAN (bio)transformation to detect temporal changes in toxicity reflecting the shifting mixture of intermediates and products during (bio)transformation. All experiments were run in duplicate.

CH₄ was monitored for each treatment and the toxicant-free control (Fig. S-2) using a gas chromatograph coupled to a flame ionization detector (GC-FID) (Ochoa-Herrera et al., 2009). The rate of CH₄ production was used to determine acetoclastic methanogen inhibition by comparing the rate of methane production in treatments normalized to a test-compound/mixture-free control.

2.3.2 *A. fischeri* (Microtox)

The Microtox (Bulich and Isenberg, 1981) assay was also used to characterize the toxicity of the (bio)transformation products/surrogates. Bioluminescence after 30 min chemical exposure was monitored. The individual compounds tested included iMENA (2.5-650 μ M), Ac-DAAN (9-2,285 μ M), dimer L (29-226 μ M), and BBY (0.40-103 μ M). As described for the acetoclastic methanogenesis assay, aqueous mixtures of biotransformation compounds from a starting DNAN concentration of 500 μ M that were recovered from different stages of DNAN (bio)transformation at a single 36-fold dilution were also evaluated in the Microtox toxicity bioassays. In preliminary tests, this dilution allowed for toxicity changes to be detected in Microtox without being completely inhibitory. All experiments were run in duplicate.

2.4 Analytical methods to assess (bio)transformation products

2.4.1 UHPLC-DAD

DNAN, MENA, and DAAN were quantified on an ultra-high performance liquid chromatograph coupled to a diode array detector (UHPLC-DAD) with an Acclaim RSLC Explosives E2 column (2.1 x 100 mm, 2.2 μ m) (Thermo Fisher Scientific, Waltham, WA, USA) and a methanol/H₂O eluent (isocratic 40/60, v/v, 0.25 mL min⁻¹) at room temperature. Detection wavelengths and retention times were (nm:min) 300:9, 254:5, and 210:2.3 for DNAN, MENA, and DAAN, respectively.

2.4.2 UHPLC-Q-ToF-MS

MENA, DAAN, and oligomer products in the liquid phase were semi-quantified without internal standards on an UltiMate 3000 UHPLC (Dionex, Sunnyvale, CA) using the same chromatography parameters and column described above for UHPLC-DAD, that was coupled to a TripleTOF 5600 quadrupole time-of-flight mass spectrometer (Q-ToF-MS) (AB Sciex,

Framingham, MA, USA). The Q-ToF-MS was run with an electrospray ionization source in positive mode at 450 °C with a capillary setting of 5.5 kV, and a declustering potential of 80 V. N₂ was used as curtain gas, desolvation gas, and nebulizer gas at 30, 35, and 35 psi, respectively. Transformation products were detected and quantified by integrating accurate, selected parent ion (M+H⁺) mass chromatogram peaks from a survey scan (m/z 30-1000) based on newly detected, and previously characterized (Olivares et al., 2013), transformation products. Both of these are shown in Table 1: previously characterized products (above) and m/z values for the uncharacterized products (below; [M+H]⁺ (retention time, min)). A potential cation [M]⁺ with m/z 247.0425* (retention time 1.52 min) was also detected. Semi-quantitative comparisons of transformation product abundances are based on integrated peak areas of parent mass ions extracted from mass chromatograms using a 10 mDa window centered on the calculated masses. AnalystTF 1.6 with PeakView 1.2.0.3 and MultiQuant version 2.1 were used to develop the ion list and to quantify individual ion masses from integrated mass chromatogram peaks.

Other analyses, including pH and volatile suspended solids content in sludge, were performed according to standard methods (APHA, 2005).

3. Results and Discussion

3.1 Biotransformation product profiles

An overview of (bio)transformation products and transformation pathways observed in this work are shown in Fig. 1 Panel A. DNAN underwent nitro-reduction primarily to MENA (but also to iMENA), and subsequent nitro-reduction led to DAAN, as reported for anaerobic soil and sludge systems (Platten et al., 2010; Olivares et al., 2013; Olivares et al., 2016). Nitro-group reduction is known to occur biologically via the broad family of nitroreductases

(Roldan et al., 2008) or by abiotic factors, such as ferrous iron adsorbed to mineral surfaces (Klausen et al., 1995; Elsner et al., 2004) and sulfide via natural organic matter electron transfer (van Beelen and Burris, 1995). During the course of nitro-group reduction, reactive hydroxylamino and nitroso intermediates are formed (Bryant and DeLuca, 1991; Spain, 1995), which can lead to further reactions. A putative nitroso intermediate coupled with DAAN to form an azo-dimer, 3,3'-diamino-4,4'-dimethoxy-azobenzene (m/z 273). In anaerobic environments, azo-dimerization could be explained by direct coupling of nitroso- and amino-derivatives, which has been used in azo-dye syntheses under reducing conditions (Moglie et al., 2008; Zhao et al., 2011). In some cases, m/z 273 underwent reductive azo-cleavage to DAAN via a hydrazo intermediate, 3,3'-diamino-4,4'-dimethoxy-hydrazobenzene (m/z 275). Reductive azo-cleavage via hydrazo intermediates has been shown to occur in biological reducing environments (Bin et al., 2004). In addition to the main (bio)transformation pathway, there were secondary routes. For instance, the detection of Ac-DAAN indicated DAAN underwent N-acetylation, a reaction reported to occur biologically in anaerobic conditions with aromatic amines (Razo-Flores et al., 1999). Another parallel pathway involved reactions of azo-dimer m/z 273, such as N-methylation to yield 5-((3-Amino-4-methoxyphenyl)diazenyl)-2-methoxy-N-methylethaniline (m/z 285) and O-demethoxylation, forming 4'-methoxy-3-methylamino-3'-methylethylamino-azobenzene (m/z 269).

DNAN was (bio)transformed by microbial and abiotic processes in soil (Hawari et al., 2015; Olivares et al., 2016) and sludge (Platten et al., 2010; Olivares et al., 2013). DNAN was readily reduced within 10 days in the anaerobic sludge and in the Camp Navajo soil, while for the Camp Butner soil, DNAN was completely removed by 30 days (Fig. 2A). The specific rate of reduction per unit dwt solids was highest for the sludge and Camp Navajo soil

216 (45.6 and 56.7 $\mu\text{mol DNAN d}^{-1} \text{dwt g}^{-1}$), while Camp Butner had the slowest (14.3 μmol
217 $\text{DNAN d}^{-1} \text{g dwt soil}^{-1}$).

218 Very limited amounts of MENA and DAAN were detected during (bio)transformation
219 based on HPLC-DAD (Fig. S-3), suggesting that additional transformation products had been
220 formed. Completing a mass balance accounting for DNAN (bio)transformation has been
221 problematic when relying exclusively on HPLC-DAD (Olivares et al., 2013). While
222 additional products can be confidently identified by mass spectrometry techniques, not all of
223 the compounds identified can be quantified (Platten et al., 2010; Perreault et al., 2012;
224 Olivares et al., 2013).

225 The individual profile of (bio)transformation product mixtures formed at different stages
226 of DNAN (bio)transformation for the two soils, Camp Butner (Fig. 2A) and Navajo (Fig.
227 3A), as well as for the anaerobic sludge (Fig. 4A), were determined using LC-QToF-MS to
228 detect products that were below detection in the HPLC-DAD. Overall, the rate of DNAN
229 conversion affected the rate of production of monomer and oligomer products. MENA and
230 DAAN were detected primarily between 10-30d of incubation in Camp Navajo soil, but were
231 detected only sporadically in the Camp Butner soil. Unlike the soils, in the anaerobic sludge a
232 consistent amount of DAAN was detected throughout 0-50d of incubation, accounting for
233 slightly less than a third of the total product peak area (3×10^7 area units for DAAN). A higher
234 amount of DAAN in the system might result from better reducing conditions. For instance,
235 2,4,6-triaminotoluene from TNT reduction has only been reported below -200 mV (Esteve-
236 Núñez et al., 2001). A hydrazo compound (m/z 275 in Fig. 1) had a higher occurrence in the
237 sludge (Fig. 4) and is believed to be an intermediate formed during the reductive cleavage of
238 azo-dimers and a source of DAAN.

Other monomer products, such as iMENA and Ac-DAAN were detected in smaller quantities than MENA or DAAN. For iMENA, the highest amount detected was 2×10^6 area units in Camp Butner soil at 20 d of (bio)transformation, but overall MENA was at least twice as abundant than its isomer. For Ac-DAAN, the largest amount detected was in sludge (8×10^5 area units), but its signal was on average two orders of magnitude smaller than DAAN.

Azo-dimer products were detected concomitantly with the formation of MENA, iMENA, DAAN, and Ac-DAAN detectable by QToF-MS in all three systems studied, suggesting that dimerization occurs rapidly during nitro-reduction. However, at longer incubation times (9-30 days), the major abundant species were products with ion $m/z > 200$, attributed to azo-dimers. Based on common dominant (bio)transformation products detected across all systems, dimers underwent other reactions, such as N-substitution and O-demethoxylation.

Besides common dominant products, the relative abundance of each product varied depending on the system. For example in Camp Butner soil incubations, the most abundant ions included m/z 269 and 285. In Camp Navajo, the primary products were m/z 247 and 274 (putatively with molecular formulas $C_7H_9N_3O_7$ and $C_9H_{11}N_3O_7$, respectively), followed by m/z 269 and 285, like in Camp Butner. The ions m/z 247 and 274 have not been characterized, but could be a product formed during incorporation of DNAN reduced (bio)transformation products with humic substances. Covalent binding between quinone-like and reduced TNT products, a chemical analog to DNAN, have been found in ^{15}N studies with soil humic substances in aerobic conditions (Thorn and Kennedy, 2002). In anaerobic sludge, the primary dimers detected included m/z 285 and 269, with significant contributions of m/z 273 and 275.

The largest cumulative amount of products in the aqueous phase, as quantified by peak area, occurred in Camp Butner soil and in the anaerobic sludge (1.2×10^6 area units). On the other hand, Camp Navajo soil, had two orders of magnitude less products at longer incubation times (40-50d). A higher abundance in Camp Navajo was also observed with compounds with the highest molecular weight in the list, such as m/z 325 and 327, which were detected in all systems more often after DNAN was depleted (Figs. S-5 through S-7). Their concentration was in the range of $\sim 10^5$ area units for Camp Butner and sludge, but $\sim 10^4$ for Camp Navajo. The metabolite m/z 431, a potential trimer, based on the molecular mass, was detected only in Camp Navajo soil at a 2.5-fold higher amount than m/z 325 and 327 (Fig. S-6). Camp Navajo has more organic carbon than Camp Butner (5.24 versus 2.07 % dwt soil) (Krzmarzick et al., 2015), and the difference in amount of products recovered may be due to irreversible adsorption and covalent incorporation into soil humus. In studies with ^{14}C radiolabeled TNT incubated with soil, more than half of the label was incorporated into different soil humic fractions (Drzyzga et al., 1999).

Data obtained from the semi-quantitative mass spectrometry analysis allowed a limited interpretation of the cumulative mass chromatogram peak areas. Because the method lacked calibration standards, it mainly provided temporal comparative information for each compound, since different ionization efficiencies likely exist between compound species; although magnitude differences in abundance between separate species may also be interpreted. And although matrix effects may exist, they were restrained by use of the same matrix (a mineral salts basal medium) in all biotransformation mass spectrometry analyses.

3.2 Monomeric (bio)transformation products as pure compounds

Based on the (bio)transformation products detected in this work, a chemical library with monomer products and best available surrogates for azo-dimer and trimer compounds was

created for microbial toxicity assessment. Most monomeric metabolites from DNAN reduction had decreased microbial toxicity with the exception of the MENA isomers (Table 2, Fig. 5). DAAN and especially Ac-DAAN were many fold less toxic than DNAN, whereas iMENA and MENA were more toxic than DNAN in the case of methanogens and Microtox, respectively. This is consistent with nitroaniline isomers from nitroreduction of 2,4-dinitrotoluene being more toxic than the parent compound in *A. fischeri* (Dodard et al., 1999). In addition, nitroanilines were also the most toxic compounds in an extensive evaluation of the methanogenic toxicity of 24 different amino- and nitro-substituted benzenes (Donlon et al., 1995). During the course of (bio)transformation, regiospecificity in DNAN (bio)conversion could impact the overall toxicity. For instance, nitro-reduction of DNAN has predominantly been reported in the *ortho* group in biological systems (Perreault et al., 2012; Olivares et al., 2013; Hawari et al., 2015), whereas evidence of *para*-nitro-reduction has been found primarily in abiotic systems (Ahn et al., 2011; Hawari et al., 2015), with one account of biological production (Schroer et al., 2015). Dominance of nitroreduction in the *ortho* group has been attributed to overall molecular stability due to H-bonding with the neighboring methoxy group (Hawari et al., 2015). Given these considerations, the toxicity of the nitroaniline isomers formed from DNAN nitro-reduction is expected to be primarily due to MENA.

On the other hand, there are other reactions during (bio)transformation that could yield less toxic products. Complete nitro-reduction to aromatic amines has been considered as a detoxification mechanism (Gorontzy et al., 1993), and aromatic amines were generally less cytotoxic than nitroaromatic counterparts to methanogens (Donlon et al., 1995; Liang et al., 2013) and *A. fischeri* (Neuwoehner et al., 2007; Liang et al., 2013). In a previous study we reported DAAN as less toxic than DNAN or MENA in both microbial assays (Fig. 5) (Liang et al., 2013). In addition, other transformation pathways could further decrease the inhibition

potential of aromatic amines. For instance, DAAN N-acetylation resulted in a considerable decrease in toxicity (Fig. S-3 panels A1,B1 and Fig. 5). Ac-DAAN was the least toxic of all monomers, and did not cause inhibition up to 8 mM to acetoclastic methanogens and had a 50% inhibition concentration (IC_{50}) of 911 μ M in Microtox. It had a much lower inhibition than its non-acetylated chemical analog, DAAN. N-acetylation has been reported as a detoxification mechanism for amine moieties in aromatic compounds (Tweedy et al., 1970; Bruns-Nagel et al., 2000).

3.3 Surrogates of dimers and trimers

The azo-oligomers tested, dimer L and BBY, were among the compounds causing the strongest toxicity. Dimer L was a potent toxicant to *A. fischeri* (IC_{50} = 29.8 μ M) and methanogens (IC_{50} = 65 μ M). While there is limited information on toxicity of oligomers formed during the biotransformation of explosives, a study on the cytotoxicity of TNT and its degradation products to H4IIE cells and Chinese hamster ovary-K1 (CHO) cells found that the azoxy-dimers were as toxic as the parent compound (Honeycutt et al., 1996). BBY was found to be the most toxic of all compounds tested with an IC_{50} in each of the microbial systems of 0.7 μ M. Even though both systems had the same IC_{50} , *A. fischeri* were more sensitive than methanogens, since complete inhibition occurred at 1.4 μ M (Fig. S-3 Panel A2).

The IC_{50} values reported in this work are based on the molarity of the chemical species. However, a comparison of toxicity for micromolar monomer equivalents (two for dimer L, three for BBY), reveals dimer L to be less toxic than DNAN (open symbols, Fig. 5), and BBY to be the most toxic tested in this work, and still considerably more toxic than DNAN (Fig. 5).

335

336 3.4 Exposures to DNAN (bio)transformation mixtures

337 3.4.1 Toxicity profile during (bio)transformation

338 Acetoclastic methanogenic and *A. fischeri* bioluminescent inhibition assays were used to
339 assess the overall toxicity of the aqueous extracts of DNAN (bio)transformation product
340 mixtures sampled from the biotransformation assays at different times of anaerobic
341 incubation. Both microbial toxicity targets were very susceptible to toxicity of the mixture of
342 (bio)transformation products. The sample mixtures were diluted significantly (54-fold for
343 methanogens and 36-fold for *A. fischeri*) before exposure, because at lower dilutions, there
344 was complete inhibition in both assays which did not allow observable differences in
345 inhibition across the (bio)transformation timeline.

346 At early stages of DNAN bioconversion, methanogenic activity dropped sharply across
347 all systems assayed (Figs. 2B-4B). This period of increased inhibition lasted as long as
348 DNAN was detectable in aqueous solutions. The dip in activity was the shortest in anaerobic
349 sludge (1d) (Fig. 4B), while for Camp Butner (with the slowest rate of DNAN
350 (bio)conversion), it lasted for 30 days (Fig. 3B).

351 The increase in methanogen inhibition could be attributed to putative reactive
352 intermediates formed during nitroreduction of DNAN, such as nitroso- and hydroxylamino-
353 derivatives. Nitroso- and hydroxylamino-products have been identified as potentially
354 responsible for methanogen cell lysis (and methanogenic inhibition) during nitroreduction
355 (Gorontzy et al., 1993). Cytotoxicity studies on TNT and its reduced (bio)transformation
356 products, showed that 4-hydroxylamino,2,6-dinitrotoluene could be as toxic or more toxic
357 than TNT (Honeycutt et al., 1996). DNAN nitroso-derivatives have been found in abiotic
358 (Hawari et al., 2015) and biotic systems (Olivares et al., 2016), as well as for other

359 nitroaromatic compounds such as 2,4-dinitrotoluene (Liu et al., 1984). Hydroxylamino
360 products have also been reported in DNAN (Perreault et al., 2012; Olivares et al., 2016) and
361 TNT (Knicker et al., 2001) reductive (bio)transformation studies. Furthermore, azo-dimers
362 detected in this work could be indirect evidence of the occurrence of these intermediates
363 since classic azo formation involves coupling of nitroso-bearing products with either
364 hydroxylamino (Smith and March, 2006) or amino derivatives (Moglie et al., 2008; Zhao et
365 al., 2011).

366 At longer incubation times (30-50d), the diluted mixtures of metabolites present at this
367 stage of (bio)transformation caused very low inhibition to methanogens. Compared to earlier
368 incubation times, when the toxicity was maximum, there was clearly a decrease in toxicity
369 associated with the formation of azo-dimers. This recovery of activity might be attributed to
370 the disappearance of a steady supply of reactive intermediates from nitroreduction once
371 DNAN is depleted from the system. In all cases, after the initial inhibition, there was a
372 consistent recovery and detoxification compared to the initial stage of (bio)transformation,
373 eventually achieving restoration of 100% activity compared to the controls. Thus, the
374 complete toxicity profile of DNAN (bio)transformation towards methanogens could be
375 described by a Big Dipper constellation pattern (particularly in Fig. 2B), with initial
376 inhibition and latter activity recovery.

377 For *A. fischeri*, there was little inhibition during initial (bio)transformation of DNAN in
378 all systems, but inhibition started to increase when more than half of the DNAN had been
379 converted (Figs. 2B-4B). Unlike methanogens, *A. fischeri* may not be affected by the reactive
380 nitroso and hydroxylamino intermediates if coping mechanisms against radicals and N-
381 reactive species are present. Recently, flamohaemoglobin Hmp in *A. fischeri* has been
382 attributed to protect against NO and reactive oxygen produced by the host squid as an
383 antimicrobial agent during colonization in *A. fischeri*-squid symbiosis (Wang et al., 2010).

However, at the latter part of DNAN reduction and after DNAN was completely (bio)transformed, there was increased inhibition of bioluminescence. During these stages of (bio)transformation, there was also significant detection of dimer products ($m/z > 200$), which could be directly responsible for affecting bioluminescence due to inhibition in growth. With Camp Navajo soil, the highest inhibition (as evidenced by the lowest normalized activity) occurred at 50 days (corresponding to 88% inhibition) compared to the toxicant-free control. However, in Camp Butner soil and in the anaerobic sludge, the maximum inhibition was 50%. In the anaerobic sludge there was a temporary decrease in inhibition from 20-40d which occurred at the same time as the amount of detected dimers decreased, and by 50d the inhibition increased again concomitantly with the dimer products. Based on the oscillation of toxicity from 20-50d in the anaerobic sludge system, dimer abundance might be directly related to *A. fischeri* inhibition.

4. Conclusions

A workflow integrating transformation product semi-quantitation with microbial toxicity provided information about scenarios expected in soils and wastewater sludge systems polluted with DNAN. Two key events during (bio)transformation were identified as potential toxicity drivers. Increased methanogenic inhibition during nitroreduction of DNAN. This increase is attributed to putative reactive nitroso-intermediates, while mixtures rich with azo-dimers were severely toxic to *A. fischeri*. On the other hand, N-acetylation greatly reduced toxicity of DNAN (bio)transformation products. N-acetylation and azo-dimerization of amines are transformation pathways that have been reported to be in direct competition with each other (Tweedy et al., 1970). Both of these processes have been identified in microbial systems in the presence of DNAN (Perreault et al., 2012; Olivares et al., 2013), although azo-

dimerization seems to be the dominant route based on semi-quantitative analysis of this work. However, devising strategies to increase N-acetylation reactions could be exploited in bioremediation efforts to yield non-toxicant products in systems polluted with DNAN.

5. Acknowledgments

This study was supported by the Strategic Environmental Research and Development Program (SERDP) project ER-2221 and NSF CBET 1510698. Analyses performed at the Arizona Laboratory for Emerging Contaminants (ALEC) were supported by NSF CBET 0722579, AB Sciex, and additional funding from University of Arizona colleges. CIO was supported by the Mexican National Council for Science and Technology (CONACyT) and the NIEHS Superfund Research Program (P42 ES04940).

6. References

- Ahn, S.C., Cha, D.K., Kim, B.J., Oh, S.-Y., 2011. Detoxification of PAX-21 ammunitions wastewater by zero-valent iron for microbial reduction of perchlorate. *J. Hazard. Mater.* 192, 909-914.
- APHA, 2005. Standard methods for the examination of water & wastewater. American Public Health Association, American Water Works Association, Water Environment Federation., Washington, D.C.
- Bin, Y., Jiti, Z., Jing, W., Cuihong, D., Hongman, H., Zhiyong, S., Yongming, B., 2004. Expression and characteristics of the gene encoding azoreductase from *Rhodobacter sphaeroides* AS1.1737. *FEMS Microbiol. Lett.* 236, 129-136.
- Bruns-Nagel, D., Knicker, H., Drzyzga, O., Bütehorn, U., Steinbach, K., Gemsa, D., von Löw, E., 2000. Characterization of ¹⁵N-TNT residues after an anaerobic/aerobic treatment of soil/molasses mixtures by solid-state ¹⁵N NMR spectroscopy. 2. Systematic investigation of whole soil and different humic fractions. *Environ. Sci. Technol.* 34, 1549-1556.
- Bryant, C., DeLuca, M., 1991. Purification and characterization of an oxygen-insensitive NAD(P)H nitroreductase from *Enterobacter cloacae*. *J. Biol. Chem.* 266, 4119-4125.
- Bulich, A.A., Isenberg, D.L., 1981. Use of the luminescent bacterial system for the rapid assessment of aquatic toxicity. *ISA T.* 20, 29-33.
- Davies, P.J., Provatas, A., Defence, S., Technology Organisation . Weapons Systems, D., 2006. Characterisation of 2,4-dinitroaniline an ingredient for use in low sensitivity melt cast formulations. Science Defence Technology Organization, Edinburgh, Australia.

- Dodard, S.G., Renoux, A.Y., Hawari, J., Ampleman, G., Thiboutot, S., Sunahara, G.I., 1999. Ecotoxicity characterization of dinitrotoluenes and some of their reduced metabolites. *Chemosphere* 38, 2071-2079.
- Dodard, S.G., Sarrazin, M., Hawari, J., Paquet, L., Ampleman, G., Thiboutot, S., Sunahara, G.I., 2013. Ecotoxicological assessment of a high energetic and insensitive munitions compound: 2,4-dinitroanisole (DNAN). *J. Hazard. Mater.* 262, 143-150.
- Donlon, B.A., Razo-Flores, E., Field, J.A., Lettinga, G., 1995. Toxicity of N-substituted aromatics to acetoclastic methanogenic activity in granular sludge. *Appl. Environ. Microb.* 61, 3889-3893.
- Drzyzga, O., Bruns-Nagel, D., Gorontzy, T., Blotevogel, K.H., von Low, E., 1999. Anaerobic incorporation of the radiolabeled explosive TNT and metabolites into the organic soil matrix of contaminated soil after different treatment procedures. *Chemosphere* 38, 2081-2095.
- Elsner, M., Schwarzenbach, R.P., Haderlein, S.B., 2004. Reactivity of Fe(II)-bearing minerals toward reductive transformation of organic contaminants. *Environ. Sci. Technol.* 38, 799-807.
- Esteve-Núñez, A., Caballero, A., Ramos, J.L., 2001. Biological degradation of 2,4,6-trinitrotoluene. *Microbiol. Mol. Biol. R.* 65, 335-352.
- Gorontzy, T., Küver, J., Blotevogel, K.H., 1993. Microbial transformation of nitroaromatic compounds under anaerobic conditions. *J. Gen. Microbiol.* 139, 1331-1336.
- Hawari, J., Monteil-Rivera, F., Perreault, N.N., Halasz, A., Paquet, L., Radovic-Hrapovic, Z., Deschamps, S., Thiboutot, S., Ampleman, G., 2015. Environmental fate of 2,4-dinitroanisole (DNAN) and its reduced products. *Chemosphere* 119, 16-23.
- Honeycutt, M.E., Jarvis, A.S., McFarland, V.A., 1996. Cytotoxicity and mutagenicity of 2,4,6-trinitrotoluene and its metabolites. *Ecotox. Environ. Safe* 35, 282-287.
- Kennedy, A.J., Laird, J.G., Lounds, C., Gong, P., Barker, N.D., Brasfield, S.M., Russell, A.L., Johnson, M.S., 2015. Inter- and intraspecies chemical sensitivity: A case study using 2,4-dinitroanisole. *Environ. Toxicol. Chem.* 34, 402-411.
- Klausen, J., Trober, S.P., Haderlein, S.B., Schwarzenbach, R.P., 1995. Reduction of substituted nitrobenzenes by Fe(II) in aqueous mineral suspensions. *Environ. Sci. Technol.* 29, 2396-2404.
- Knicker, H., Achtnich, C., Lenke, H., 2001. Solid-state nitrogen-15 nuclear magnetic resonance analysis of biologically reduced 2,4,6-trinitrotoluene in a soil slurry remediation. *J. Environ. Qual.* 30, 403-410.
- Krzmarzick, M.J., Khatiwada, R., Olivares, C.I., Abrell, L., Sierra-Alvarez, R., Chorover, J., Field, J.A., 2015. Biotransformation and degradation of the insensitive munitions compound, 3-nitro-1,2,4-triazol-5-one, by soil bacterial communities. *Environ. Sci. Technol.* 49, 5681-5688.
- Liang, J., Olivares, C., Field, J.A., Sierra-Alvarez, R., 2013. Microbial toxicity of the insensitive munitions compound, 2,4-dinitroanisole (DNAN), and its aromatic amine metabolites. *J. Hazard. Mater.* 262, 281-287.
- Linker, B.R., Khatiwada, R., Perdrial, N., Abrell, L., Sierra-Alvarez, R., Field, J.A., Chorover, J., 2015. Adsorption of novel insensitive munitions compounds at clay mineral and metal oxide surfaces. *Environ. Chem.* 12, 74-84.
- Liu, D., Thomson, K., Anderson, A.C., 1984. Identification of nitroso compounds from biotransformation of 2,4-dinitrotoluene. *Appl. Environ. Microb.* 47, 1295-1298.
- Moglie, Y., Vitale, C., Radivoy, G., 2008. Synthesis of azo compounds by nanosized iron-promoted reductive coupling of aromatic nitro compounds. *Tetrahedron Lett.* 49, 1828-1831.

- Neuwoehner, J., Schofer, A., Erlenkaemper, B., Steinbach, K., Hund-Rinke, K., Eisentraeger, A., 2007. Toxicological characterization of 2,4,6-trinitrotoluene, its transformation products, and two nitramine explosives. *Environ. Toxicol. Chem.* 26, 1090-1099.
- Niedzwiecka, J.B., Finneran, K.T., Arnett, C.M., 2013. Biological and chemical mechanisms for 2,4-dinitroanisole (DNAN) degradation. Graduate Research and Discovery Symposium (GRADS). Paper 64. http://tigerprints.clemson.edu/grads_symposium/64.
- Ochoa-Herrera, V., Banihani, Q., León, G., Khatri, C., Field, J.A., Sierra-Alvarez, R., 2009. Toxicity of fluoride to microorganisms in biological wastewater treatment systems. *Water Res.* 43, 3177-3186.
- Olivares, C., Liang, J., Abrell, L., Sierra-Alvarez, R., Field, J.A., 2013. Pathways of reductive 2,4-dinitroanisole (DNAN) biotransformation in sludge. *Biotechnol. Bioeng.* 110, 1595-1604.
- Olivares, C.I., Abrell, L., Khatiwada, R., Chorover, J., Sierra-Alvarez, R., Field, J.A., 2016. (Bio)transformation of 2,4-dinitroanisole (DNAN) in soils. *J. Hazard. Mater.* 304, 214-221.
- Perreault, N.N., Manno, D., Halasz, A., Thiboutot, S., Ampleman, G., Hawari, J., 2012. Aerobic biotransformation of 2, 4-dinitroanisole in soil and soil *Bacillus* sp. *Biodegradation* 23, 287-295.
- Platten, W.E., Bailey, D., Suidan, M.T., Maloney, S.W., 2010. Biological transformation pathways of 2,4-dinitroanisole and N-methylparanitroaniline in anaerobic fluidized-bed bioreactors. *Chemosphere* 81, 1131-1136.
- Razo-Flores, E., Lettinga, G., Field, J.A., 1999. Biotransformation and biodegradation of selected nitroaromatics under anaerobic conditions. *Biotechnol. Progr.* 15, 358-365.
- Roldan, M., Perez-Reinado, E., Castillo, F., Moreno-Vivian, C., 2008. Reduction of polynitroaromatic compounds: the bacterial nitroreductases. *Fems Microbiol. Rev.* 32, 474-500.
- Schroer, H.W., Langenfeld, K.L., Li, X., Lehmler, H.-J., Just, C.L., 2015. Stable isotope-enabled pathway elucidation of 2,4-dinitroanisole metabolized by *Rhizobium litchii*. *Environ. Sci. Technol. Lett.*
- Smith, M.B., March, J., 2006. Aliphatic, Alkenyl, and Alkynyl Substitution, Electrophilic and Organometallic. *March's Advanced Organic Chemistry*. John Wiley & Sons, Inc., pp. 752-852.
- Spain, J.C., 1995. Biodegradation of nitroaromatic compounds. *Annu. Rev. Microbiol.* 49, 523-555.
- Thorn, K.A., Kennedy, K.R., 2002. N-15 NMR investigation of the covalent binding of reduced TNT amines to soil humic acid, model compounds, and lignocellulose. *Environ. Sci. Technol.* 36, 3787-3796.
- Tweedy, B.G., Loeppky, C., Ross, J.A., 1970. Metobromuron: Acetylation of the aniline moiety as a detoxification mechanism. *Science* 168, 482-483.
- van Beelen, P., Burris, D.R., 1995. Reduction of the explosive 2,4,6-trinitrotoluene by enzymes from aquatic sediments. *Environ. Toxicol. Chem.* 14, 2115-2123.
- Wang, Y., Dunn, A.K., Wilneff, J., McFall-Ngai, M.J., Spiro, S., Ruby, E.G., 2010. *Vibrio fischeri* flavohaemoglobin protects against nitric oxide during initiation of the squid-*Vibrio* symbiosis. *Mol. Microbiol.* 78, 903-915.
- Zhao, R., Tan, C., Xie, Y., Gao, C., Liu, H., Jiang, Y., 2011. One step synthesis of azo compounds from nitroaromatics and anilines. *Tetrahedron Lett.* 52, 3805-3809.

7. Figure legends

Fig. 1. Panel A: (Bio)transformation pathways of DNAN in anaerobic incubations of soils and sludge. Microbial toxicity was evaluated for the monomer compounds shown. Panel B: Azo-oligomer surrogate compounds used for microbial toxicity. Notation: OMe = methoxy; Ac = acetyl. MENA and iMENA are isobaric regioisomers.

Fig. 2. DNAN anaerobic (bio)transformation in Camp Butner soil and collective microbial toxicity of the product mixtures during incubation. Panel A: DNAN ($\square$) formed transformation products (stacked area, from light to dark in ascending molecular mass): 139.0866, 165.0659, 169.0608, 181.0972, 185.0652, 193.0607, 228.0768, 243.0877, 243.1241, 245.1300, 247.0425, 259.1190, 267.0975, 269.1397, 273.1347, 274.0715, 275.1503, 285.1347, 299.1179, 301.1289, 313.1289, 325.1659, 327.1452, 431.1569. The predominant $[M+H]^+$ values are underlined and are also labeled in the plot. Panel B: Impact of mixtures of transformation products formed during (bio)transformation on *A. fischeri* bioluminescence ($\blacksquare$) and acetoclastic methanogens ($\bullet$).

Fig. 3. DNAN anaerobic (bio)transformation in Camp Navajo soil and collective microbial toxicity of the product mixtures during incubation. Panel A: DNAN ($\square$) formed transformation products (stacked area, from light to dark in ascending molecular mass): 139.0866, 165.0659, 169.0608, 181.0972, 185.0652, 193.0607, 228.0768, 243.0877, 243.1241, 245.1300, 247.0425, 259.1190, 267.0975, 269.1397, 273.1347, 274.0715, 275.1503, 285.1347, 299.1179, 301.1289, 313.1289, 325.1659, 327.1452, 431.1569. The predominant $[M+H]^+$ values are underlined and are also labeled in the plot. Panel B: Impact

of mixtures of transformation products formed during (bio)transformation on *A. fischeri* bioluminescence (■) and acetoclastic methanogens (●).

Fig. 4. DNAN anaerobic (bio)transformation in anaerobic sludge and collective microbial toxicity of the product mixtures during incubation. Panel A: DNAN (□) formed transformation products (stacked area, from light to dark in ascending molecular mass): 139.0866, 165.0659, 169.0608, 181.0972, 185.0652, 193.0607, 228.0768, 243.0877, 243.1241, 245.1300, 247.0425, 259.1190, 267.0975, 269.1397, 273.1347, 274.0715, 275.1503, 285.1347, 299.1179, 301.1289, 313.1289, 325.1659, 327.1452, 431.1569. The predominant [M+H]⁺ values are underlined and are also labeled in the plot. Panel B: Impact of mixtures of transformation products formed during (bio)transformation on *A. fischeri* bioluminescence (■) and acetoclastic methanogens (●).

Fig. 5. Fifty-percent inhibition concentrations (IC₅₀) for acetoclastic methanogens (■) and *A. fischeri* (●) for DNAN and its (bio)transformation products and azo-oligomer surrogates. Shaded area indicates DNAN IC₅₀ range. Open symbols in dimer L show adjusted concentrations to monomer equivalents. Ac-DAAN did not cause inhibition to methanogens and the highest concentration tested (8000 μM) is shown instead with an asterisk above the symbol. The BBY IC₅₀ for both microbial targets was 0.7 μM.

Table 1. Selected parent ion list for semi-quantitative LC-MS determination of soluble products formed during anaerobic DNAN

(bio)transformation in soil or sludge. A mass range of 0.01 Da was used to select ions $[M+H]^+$ to measure for peak integration.

Chemical compound (identifier); Molecular formula;	CAS #	Calculated $[M+H]^+$	Retention time (min)
2,4-diaminoanisole (DAAN); $C_7H_{10}N_2O$	615-05-4	139.0866	2.51
2-methoxy-5-nitroaniline (MENA); $C_7H_8N_2O_3$	99-59-2	169.0608	6.23
3-nitro-4-methoxyaniline (iMENA); $C_7H_8N_2O_3$	577-72-0	169.0608	2.99
N-(5-amino-2-methoxyphenyl) acetamide (Ac-DAAN); $C_9H_{12}N_2O_2$	64353-88-4	181.0972	2.14
3-amino-3'-nitro-azobenzene (m/z 243); $C_{12}H_{10}N_4O_2$	61390-99-6	243.0877	1.87
4'-methoxy-3-methylamino-3'-methyleamino-azobenzene (m/z 269); $C_{15}H_{16}N_4O$	N/A	269.1397	5.09
3,3'-diamino-4,4'-dimethoxy-azobenzene (m/z 273); $C_{14}H_{16}N_4O_2$	N/A	273.1347	13.75
3,3'-diamino-4,4'-dimethoxy-hydrazobenzene (m/z 275); $C_{14}H_{18}N_4O_2$	N/A	275.1503	3.48
5-((3-Amino-4-methoxyphenyl)diazenyl)-2-methoxy-N-methyleaniline (m/z 285); $C_{15}H_{16}N_4O_2$	N/A	285.1347	10.61
Unassigned parent compounds; m/z (ret. time in min): 165.0659 (2.84), 185.0652 (4.66), 193.0607 (2.52), 228.0768 (2.86), 243.1241 (13.66), 245.1300 (4.13), 247.0425 (1.52), 259.1190 (9.02), 267.0975 (1.63), 274.0715 (11.24), 299.1179 (7.45), 301.1289 (12.3), 313.1289 (3.21), 325.1659 (12.07), 327.1452 (3.81), 431.1569 (1.32).			

584

585

586

587

Table 2. Summary of inhibitory concentrations for DNAN, (bio)transformation products and best available surrogates to acetoclastic methanogens and *A. fischeri* (Microtox)

Compound	Methanogens			Microtox			Source
	IC ₂₀	IC ₅₀	IC ₈₀	IC ₂₀	IC ₅₀	IC ₈₀	
	μM						
DNAN	16	41	70	27	57	110	Liang et al., 2013
MENA	121	175	275	13	48	143	Liang et al., 2013
DAAN	72	176	327	39	155	>302 ^a	Liang et al., 2013
iMENA	8	25	43.4	114	219	>446 ^b	This study
Ac-DAAN	>8000 ^c	>8000 ^c	>8000 ^c	660	911	1266	This study
dimer L	22	65	76	18.6	29.8	49.7	This study
BBY	0.4	0.71	1.2	0.3	0.7	3.14	This study

588

^a302 μM DAAN caused 75% inhibition to *A. fischeri*.

589

^b446 μM iMENA caused 62% inhibition to *A. fischeri*.

590

^c8000 μm Ac-DAAN caused 2% inhibition to acetoclastic methanogens.

Figure 1

Figure 2

Figure 3

606

607

608 **Figure 4**

Figure 5