When mothers need others: the impact of hominin life history evolution on cooperative breeding

Karen L. Kramer ^{a,b,*} and Erik. Otárola-Castillo ^{b,c}

^a *Department of Anthropology, University of Utah, 270 S 1400 E, Salt Lake City, UT 84112*

^b *Department of Human Evolutionary Biology, Harvard University, 11 Divinity Ave., Cambridge, MA 03128*

^c *Department of Ecology, Evolution and Organismal Biology, Iowa State University, Ames, IA 50011*

Corresponding author.

E-mail address: karen.kramer@anthro.utah.edu (K. L. Kramer).

Keywords: Human evolution; Juvenility; Intergenerational transfers; Parental care

Abstract

The evolution of cooperative breeding is particularly complex in humans because many other traits that directly affect parental care (shorter birth intervals, increased offspring survivorship, juvenile dependence, and older ages at dispersal) also emerge during the Pleistocene. If human cooperative breeding is ancient, it likely evolved in a hominin lacking a fully modern life history. However, the impact that changing life history traits have on parental care and cooperative breeding has not been analytically investigated. We develop an exploratory model to simulate an economic problem that would have arisen over the course of hominin life history evolution to identify those transitions that produced the strongest pressures for cooperative childrearing. The model generates two central predictions. First, help within maternal-offspring groups can support early changes in juvenile dependence, dispersal age, birth intervals, and fertility. If so, maternal-juvenile cooperation may be an important but understudied step in the evolution of human cooperative breeding. Second, pressure to recruit adult cooperation is most pronounced under more derived conditions of late dispersal and later ages of juvenile dependence, with a strong interaction at short birth intervals. Our findings indicate that changes in life history traits that affect parental care are critical in considering background selective forces that shaped the evolution of cooperative breeding.

Introduction

Compared to other closely related species, humans wean infants at a young age, have short birth intervals, and raise multiple dependents at the same time. Juveniles depend on others for their well-being and adults cooperate to raise young. Because mothers rely on others to help provision offspring, humans have been characterized as cooperation breeders (Foster and Ratnieks, 2005; Hrdy, 2005a, 2009; Kramer, 2005a, 2010; Sear and Mace, 2008; Burkart et al., 2009; Hill and Hurtado, 2009; Russell and Lummaa, 2009; Strassman and Kurapati, 2010; van Schaik and Burkart, 2010; Crepsi, 2014). Cooperative breeding is a rare mammalian reproductive strategy in which group members other than parents help to support mothers or raise their young (Russell, 2004).¹ However, this suite of traits did not always characterize humans. We likely evolved from an ape-like animal with long birth intervals, independence at weaning, mothers who raised young on their own, and juveniles who were self-sufficient foragers. While cooperative breeding is often presumed to be ancient (i.e., dating to *Homo erectus*-grade hominins), the impact that other evolving life history traits have on its emergence has not been analytically explored. Our goal is to model an economic problem that would have occurred during hominin life history evolution and to estimate the relative influences that these changes have on pressure to recruit allocares.

Evolutionary changes in dependent juveniles, birth intervals, and cooperative breeding

The pace at which mothers reproduce and how long offspring are dependent have considerable effects on parental care and have likely undergone significant transformation during

hominin evolution (Dean, 2006; Hawkes and Paine, 2006; Smith et al., 2010; Kramer, 2011). Great ape mothers nurse infants for 4–6 years (Emery Thompson, 2013). In contrast, modern children are weaned on average between the ages of 2–3 years (Kennedy, 2005; Emery Thompson, 2013). While we cannot know the length of birth intervals in the past, paleontological evidence (Humphrey, 2010; Kelley and Schwartz, 2012) and adult onset of lactase decline (Sellen, 2006) suggest that ancestral weaning ages were late, similar to those in great apes, and gradually declined over time (van Noordwijk et al., 2013). This has important implications for birth intervals and reproductive rates, which are 2–5 times faster in modern natural-fertility populations compared to other primates adjusting for body size and are the highest among the great apes (Alvarez, 2000).

The lives of juveniles have also changed substantially during hominin evolution. Because all other ape juveniles are independent foragers (Knott, 2001), juveniles in the past were likely self-sufficient at younger ages, and the direction of selection was toward a longer period of dependence. While age at independence varies among modern juveniles, even among hunter-gatherer societies considered evolutionarily relevant (Supplementary Online Material [SOM] Fig. 6) weaned children both provide some of their own calories and rely on others for some of their caloric intake. Thus, over the last two million years the duration of lactation and infancy has shortened, while support has been extended to juveniles. This combination of derived traits commits human mothers to raise multiple dependents of different ages.

Raising multi-aged dependents has well-recognized implications for human sociality and is hypothesized to be associated with the emergence of cooperative breeding (Lee and Kramer, 2002; Hrdy, 2005b, 2007, 2009; Kramer, 2005a,b; Gurven and Walker, 2006). While it may seem that a trajectory toward a longer period of dependence and shorter birth intervals would

exceed a mother's ability to raise overlapping dependent offspring on her own, the interaction effects of these life history changes on maternal time and energy budgets have not been investigated under changing evolutionary conditions. Further, because none of these traits leaves unambiguous fossil evidence, an archaeological record, or a genetic signature, the sequence of life history changes and their causal relationships are unknown. Nor is there a cooperative breeding great ape model to guide us in terms of what intermediary forms might look like. We address this empirical gap by developing an exploratory model to assess the impacts that derived life history traits have on the rate of change in the *force of dependence*. We measure the force of dependence as the net cost of overlapping offspring as a function of juvenile dependence, dispersal age, and birth intervals. The model is used to predict those life history transitions under which mothers would be unable to support children on their own and pressure would have been strongest to recruit adult help.

Modeling the force of dependence

The model incorporates several novel elements so that we can formally test the hypothesis that evolutionary changes in the duration of juvenile dependence, dispersal age, and birth intervals have directional effects on the force of dependence. First, we use life history parameters that span the ancestral hominin pattern of long birth intervals and juveniles who are self-provisioning at a young age to a modern pattern of short birth intervals and juveniles who are dependent until older ages. While most cooperative breeding models and cost of children analyses assume a modern life history, our approach allows us to expand beyond relying on contemporary ethnographic examples as a direct analogy to the conditions under which

cooperative breeding evolved. Second, the model accounts for the hominin pattern of raising multiple dependents at the same time—some younger and more dependent on provisioning from others, and some older and self-provisioning. This differs from traditional cost of children analyses that consider the net or cumulative cost of only a single child (SOM Fig. 11). The effects of overlapping young are incorporated into the model by integrating the net cost of all children alive in each year of a mother’s reproductive career. This more closely reflects the demographic pressures that the hominin reproductive pattern generates on mothers and others. Third, we use time allocation as the unit of measure for net cost. Because the human feeding niche is based on high-quality foods that require technology to access, food processing, and many other activities that do not have a caloric value, the time allocated to these activities—not only the calories that children consume—is a more comprehensive measure of the force of dependence and demands for parental or nonparental investment.

Methods

Calculating juvenile dependence

In the first step we develop an underlying model to calculate the duration of juvenile dependence. Since weaning does not correlate with independence in human children, we use the age at which children produce more resources than they consume, which marks the crossover from being a net consumer to a net producer (SOM Part 1, 3; SOM Figs 1-5).

An individual consumes a certain amount and produces another amount according to his or her age. Although specific values may vary, the general shape of these functions for human

and nonhuman primates varies with age such that both production and consumption are relatively low when young, begin to accelerate at some age, then at an inflection point they begin to decelerate, and finally reach an asymptote (Kaplan, 1994; Kramer, 2005b; Gurven and Kaplan, 2006). This maturation process was modeled using seven nonlinear mathematical equations that approximate these age-related behavioral properties of children's maturation as consumers and producers (SOM Appendix 1.1; SOM Figs. 1, 2, and 4). The models were fit to empirical production and consumption data (see below) using maximum likelihood, which were converted to Akaike Information Criterion (AIC) values to select the best fit model. For both production and consumption, the best-candidate model was a 4-parameter logistic, which accounts for the increase in variance (SOM Figs. 1 and 2). This general 4-parameter function is used to model production [1] and consumption [2] values for an average individual at age x .

$$P_{x_j} = a_p + \frac{b_p - a_p}{1 + e^{c_p(d_p - x_j)}} \quad [1]$$

$$C_{x_j} = a_c + \frac{b_c - a_c}{1 + e^{c_c(d_c - x_j)}} \quad [2]$$

Where x_j is the age value, a is the lower asymptote, b the higher asymptote, c the rate of change of production/consumption as offspring ages, and d the inflection point (acceleration to deceleration; SOM Fig. 3). The age at which the production exceeds consumption we use to denote the duration of juvenile dependence.

This modeling approach is advantageous compared, for example, to smoothing individual values because the parameters serve as useful point estimates that can be compared across

groups, populations, or species. This facilitates future hypothesis testing, for example, about differences between males and females in terms of the rate of age-related change, or the age at which the asymptote is reached.

Because we are interested in the effects that evolutionary changes in the duration of juvenile dependence have on the net cost of overlapping dependents, we include a range of production values. The baseline empirical data used in the best-fit model are taken from ethnographic observations for a population for whom detailed time allocation data are available and net production (juvenile independence) is reached at age 15 (SOM Appendix 3). By adjusting the model's parameters (shown in red in Eq [1]), the age at net production can be expressed as a continuous term that varies from age 10 (early juvenile independence) to age 20 (late juvenile independence; SOM Fig. 5). This range is intended to simulate the evolutionary transition from a juvenile who covers a greater proportion of their consumption and is self-supporting at a fairly young age to one who contributes little to their own consumption and is dependent until older ages. This range is conservative in not including an ape-like pattern where juveniles become autonomous feeders at weaning, when production and consumption functions become equivalent. This occurs around the age of 5, although chimpanzee juveniles continue to forage in the company of their mother until about age 10, which is the youngest age at net production that we include in parameter values. While here we use ages 10–20 as the duration of juvenile dependence, this range can be easily varied in future studies for other research questions and other species.

The same age-specific C schedule is used in all models to highlight the effect that changes in children's production have on net value. We believe this is reasonable because age-specific consumption for children growing up in traditional, subsistence-level societies is

expected to be closely related to changes in body size and not to vary appreciably above or below resources needed for growth and survival.

From the fitted age-specific values, a child's net value at age x can then be computed as $P_{xj} - C_{xj}$ (SOM Appendix 1.2; SOM Fig. 4). This value may be positive (net producers) or negative (net consumers) depending on an individual's age and how much they produce relative to their consumption. The age at which net value exceeds 0 indicates the age of juvenile independence.

Accounting for evolutionary changes in dispersal age

Dispersal age is added to the model because when offspring leave their natal group (a mother and her nulliparous offspring) may occur independently of the age at net production (Kaplan, 1994; Lee and Kramer, 2002). Dispersal has a parameter range of age 14–20, 14 corresponding to mean age at first birth in chimpanzees and 20 to mean age at first birth in modern ethnographic populations (Kaplan et al., 2000:158). Dispersal is included in the model by considering an offspring's coresidence in its natal unit up until he or she leaves and stops both consuming from it and contributing to it. The probability of staying becomes the complement of dispersal as follows:

$$P(S_{x_j}) = 1 - \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x_j - \mu)^2}{2\sigma^2}} dx_j \quad [3]$$

Where x_j is individual age x at time j and μ is the mean age at dispersal (at the population level, ranging between age 14 and 20), and σ is the standard deviation surrounding the mean.

Here we chose a standard deviation of two years. We then calculate $P(D)$, the probability of dispersing at a given age, by scaling Eq [3] to the appropriate mean and standard deviation and integrating to x_j (SOM Fig. 7).

Grouping these terms together, the net value of an individual offspring adjusted for the probability of being coresident at a given age, x_j is (SOM Fig. 8):

$$Net_{x_j} = P(S_{x_j}) \cdot \left(\mathbb{E}[P]_{x_j} - C_{x_j} \right) \quad [4]$$

Accounting for maternal fertility

We also need to account for the biological limits to a mother's fecundity. To avoid making an explicit assumption about maternal age at first birth, we begin with a mother who gives birth at $t = 0$ and can reproduce for 20 years. This time span is included in the model as an indicator function (1,0). If t exceeds 20 years, no new births occur; otherwise females reproduce at their scheduled birth interval (described in next section).

$$I = \text{indicator function denoting } p(\text{birth}) \text{ at year } t; \quad I = \begin{cases} 1 & \text{if } t \leq 20 \\ 0 & \text{if } t > 20 \end{cases}$$

Accounting for evolutionary changes in birth intervals and number of dependents

Most cost of offspring analyses only consider the age-specific cost or cumulative cost of a

single offspring. Because this may bias the cost of children from a mother's perspective, the method developed here accounts for the human pattern of raising multi-aged children by integrating the adjusted net values of individual children at each year in a mother's reproductive career. This can be visualized as a stepped matrix, similar to a lexis diagram, across t —the years after a mother gives birth to her first child. A simple illustration of this process is given in SOM Table 2 where children are born at an average birth interval and census out at independence. We vary birth intervals from 6 to 3 years. Six years is equivalent to a chimpanzee-like birth interval, and three years to the mean for modern human natural fertility populations (Alvarez, 2000). These are expressed as discrete values to facilitate graphing and interpreting the model.

Mapping the force of dependence across all life history combinations

Aggregating these components, given a k^{th} birth interval, the net cost can be calculated as the sum of the age-specific net values for all children at age x_j at each i^{th} year after a mother's first birth. This yearly calculation can be computed across a mother's reproductive career between the years after first birth t_0 until some year t_i . Thus, the force of dependence is the summation of all children's dispersal-adjusted net value for each year in a mother's reproductive career.

$$\text{Force of Dependence}_k = \sum_{i=0}^i \text{Net}_{x_j} \quad [5]$$

Where t = year after birth $i : \{0, 1, 2, \dots, 20\}$; $k = BI: \{6, \dots, 3\}$

Model parameters are perturbed within their specified ranges across 100 interpolated values for both juvenile dependence (age 10–20) and dispersal (age 14–20), and four birth interval values (6, 5, 4, and 3 years). This simulation produced a total of 40,000 unique sets of parameter combinations (100:100:4), with net cost mapped onto a 3-dimensional response surface (see SOM Fig. 9 for an interactive rotational display).² While the individual effects of these variables may seem intuitive, how these life history traits behave in concert when multiple offspring of different ages are considered is unknown. Our emphasis is on determining the relative influences of these variables on the force of dependence.

Caveats and limitations

Although other life history variables have directionally changed during hominin evolution, we focus on those variables that impact the cost of children. While child mortality is not specifically parameterized (because the length of a mother’s reproductive career is controlled), surviving fertility is implied through the birth interval term. At a 6 to 5-year birth interval, a mother who completes a 20-year reproductive career produces four surviving children, at a 4-year birth interval, five surviving children, and at 3-year birth-interval, seven surviving children. At premodern mortality levels (Gurven and Kaplan, 2007), surviving fertility would be equivalent to the reproductive output in the model’s 6 to 5-year birth interval. As an approximation, in a population of hunter-gatherer women, if 40% survive to age 45 (Gurven and Kaplan, 2007), surviving fertility of four children for women completing their reproductive career would generate the relatively low population growth characteristic of pre-Neolithic populations (Gignoux et al., 2011).

The model can incorporate other production and consumption values and be reconfigured under different life history assumptions. For example, for simplicity and not to add assumptions about past behaviors, we do not distinguish between male and female dispersal ages. However, for future research questions focused on modeling sex differences, sex can be incorporated as an added parameter. Although all possible ethnographic adjustments cannot be made here, as a rule of thumb any modification that would increase the negative net value ($P_{xi}-C_{xi}$) would raise the force of dependence. Likewise, any modification that would decrease net value ($P_{xi}-C_{xi}$) would lower the force of dependence and demand for parental / allo-care. While other variables can be added to the model to account for specific population or species attributes, most factors affecting net cost are expected to filter through juvenile dependence, dispersal age, and birth interval.

One limitation of our model is that, while evolutionary changes in birth interval and dispersal age can be fairly securely bracketed by spanning the range in living apes and contemporary humans, consumption and production behaviors may have other possible but unobservable scenarios not represented by extant species. We attempt to compensate for this by including a broad range of juvenile independence values (ages 10–20) rather than assuming that any one modern ethnographic group is representative of the past. We also assume that if a juvenile's production exceeds their consumption and they have not yet dispersed, their surplus is shared with others. In ethnographic populations, juveniles have been shown to increase the reproductive output of their mothers through shortening birth-intervals and increasing child survivorship (Lee and Kramer, 2002; Sear and Mace, 2008; Kramer, 2009). When such cooperative behaviors emerged cannot currently be known. If juveniles do not share, their net value may be underestimated. (Depending on age of net production, this may be on the order of 0.5–1.5 hours per day). We compensate for this potential overestimate in statistical analyses by

using the maximum net cost, which considers the point in a mother's reproductive career when children are the most expensive they ever will be given other life history parameters. Our question here is not to address why humans share but, instead, to assess what combinations of life history traits could not have evolved (or were constrained) unless mothers had access to help from other adults. If adults cooperate to help other mothers, it seems reasonable that an ontogeny of cooperation would have already been in place. Evidence for this is found among many other cooperative breeding mammals (and social insects), where subadults are important helpers (Thorne, 1997; Clutton-Brock, 2002; Russell, 2004; Gilchrist and Russell, 2007), and in the comparative cognitive psychology research on the ontogeny of sharing, fairness, and theory of mind in young children (Herrmann and Tomasello, 2006; Warneken and Tomasello, 2006; Warneken et al., 2007; Tomasello and Vaish, 2013).

Results

To illustrate how the force of dependence changes across a mother's reproductive career with specific life history combinations, eight profiles are taken from the full model at the end points of the parameter range for the three life history variables (Fig. 1). The ancestral combination of life history traits is shown in the upper left box, and the most derived combination in the bottom right. Negative values indicate the net cost, or amount of time (daily hours), that mothers or others have to spend subsidizing offspring. Negative values are greatest under the most derived combination of traits when juvenile are dependent until older ages, birth intervals are short, and dispersal age is late (bottom right box). In contrast, the net cost of

overlapping young is low when juveniles reach independence at a young age, regardless of other parameter values (top row).

To statistically compare the relative effects of juvenile dependence, birth intervals, and dispersal age, we extract the greatest negative net cost from each of the 40,000 life history profiles in the full model (shown by stars for eight of the profiles in Figure 1). This represents the point at which children are most expensive during a mother's reproductive career (SOM Appendix 2). Extracting maximum net costs has several advantages. In reality, the distribution of net costs across a mother's reproductive career (for an example see SOM Fig. 11) will vary within a sample of mothers with the same average life history traits because children will differ to some extent in their age at independence, birth intervals, and dispersal age. While these variables will vary to some degree among a mother's different children, the maximum net cost should not. It could be less (if for example mortality were particularly high for any one mother) but will not be greater under a set of life history parameters. Thus, if the model's net cost values are biased, they will be in the direction of overemphasizing the cost of children to mothers and others.

Figure 2 gives the composite of the maximum net cost as three dimensional surfaces illustrating the effects of juvenile dependence and dispersal age at 6- to 3-year birth intervals (see SOM Fig. 10 for an interactive rotational display). This figure illustrates, for example, that the maximum net cost is 6 hours/day across all values of dispersal age for a mother who gives birth at 3-year birth intervals (Figure 2, bottom right panel) and her juvenile offspring become self-sufficient at age 10. Maximum net cost is also standardized as the difference between the ancestral state of early juvenile independence, long birth intervals, and early age at dispersal and the derived pattern of later independence, shorter birth intervals, and later ages at dispersal

(Table 1). These standardized values indicate the magnitude of change in the force of dependence from the ancestral state and allow us to compare the relative effects of each trait on net cost.

Results are summarized by four major findings. Our first finding is that *early modifications of the ancestral pattern of long birth intervals, singleton dependents, and independence at weaning do not generate large net costs*. When juveniles are independent at a relatively young age, the maximum net cost of raising overlapping children is less than 5 hours/day across the range of decreasing birth interval and increasing dispersal ages (Fig. 1, top row). The net cost is even less throughout much of the rest of a mother's reproductive career, suggesting that early in the transition to modern life histories, supporting older juveniles, who self-provision a considerable portion of their needs, may not have required help from outside maternal-offspring groups. While it cannot be known how much time mothers in the past spent foraging or providing food and childcare to dependents, we can bracket the evolutionary changes in maternal investment to some extent with comparative data from living populations. Great ape adults spend on the order of 5 hours/day feeding (Hunt, 1989; Doran, 1997). Some portion of this supports the energetic cost of lactation, but great ape mothers do not produce surpluses to feed juveniles. In contrast, in modern ethnographic populations, women 20–50 years of age on average spend 8.93 hours (std=2.04, $n=15$ groups) in productive work (7.56 hours) and childcare (1.38 hours; see SOM Fig. 12 for detail).³

Our second finding is that the *force of dependence is least sensitive to changes in dispersal age* compared to other life history traits that affect the cost of offspring. The marginal value of dispersal while holding other variables constant is 9.36 compared to changes in juvenile dependence or birth intervals (marginal values = 66.32 and 21.32, respectively; Table 1). These

effects are notable in Figure 2, where at a given birth interval and dependence level the increased cost is negligible for juveniles to stay in their natal group. In contrast, the force of dependence is most sensitive to age of juvenile independence.

Our third finding is evidence of *an interaction that substantially increases the force of dependence under modern life history conditions* (Figs. 2 bottom right panel and 3). For every year offspring independence occurs at a later age, the maximum net cost increases a half-hour per day (slope = -0.51; Table 2) when dispersal is early and birth intervals are long. However, at a 3-year birth interval, maximum net cost increases by an hour per day (slope = -1.00; Table 2). Thus, although age at independence has an important influence on net cost, its impact is pronounced only when interactions with other life history variables are strongest—when age at dispersal is late, birth intervals are short (4–3 years), and mothers have high levels of surviving fertility (5–7 surviving children; Fig. 3, left to right). The importance of this interaction is that early evolutionary changes in life history traits have the least impact on the cost to raise children, while more derived changes result in substantial increases.

This interaction implies our fourth result. The force of dependence is greatest and *pressure to recruit adult help most pronounced under more derived life history conditions* of late juvenile independence when the interaction is strongest at 3-year birth intervals (Figs. 2 and 3, bottom right, Fig. 1, bottom row columns 3 and 4). We suggest that for much of our evolutionary history, surviving fertility for a women completing her reproductive career would be about four offspring (see previous discussion on child mortality in caveats and limitations). This rate is equivalent to a 5–6 year birth interval in the model. In this interval range, help outside the maternal-offspring unit may not be necessary to support more ancestral life history changes.

Discussion

If cooperative breeding is ancient and appears in *Homo erectus*-grade hominins (Hrdy, 2009; van Schaik and Burkart, 2010), it likely developed under not yet fully modern phenotypic expressions. Although birth interval length, dispersal age, and juvenile dependence directly affect parental investment, the impact of their changes has not been previously explored in its evolutionary context. Our goal is to develop a model to predict those life history transitions where selective pressure would have been strongest for cooperative childrearing.

The force of dependence model simulates an economic problem that would have arisen during the hominization of birth intervals, juvenile dependence, and dispersal ages. We take this approach to establish the underlying costs, constraints, and interactions among life history variables that directly affect parental investment. In future research these can be incorporated into evolutionary models that consider fitness tradeoffs of alternative strategies. The model generates two central predictions related to cooperative breeding. First, the net cost of these early changes in life history traits can be feasibly supported by maternal-offspring groups. Second, pressure to recruit adult reproductive cooperation follows under more derived conditions of late dispersal, late juvenile dependence, short birth intervals, and higher levels of surviving fertility.

Because modern human juveniles are subsidized, emphasis has focused on the demands of children driving the need for mothers to recruit adult help. In other primates, help needed to carry young is one explanation for biparental care (Clutton-Brock and Harvey, 1977; Kleiman, 1977). We reconsider this perspective for two reasons. First, while juveniles are often characterized as being expensive, existing cost of children analyses evaluate the net or

cumulative cost of a single dependent (Gurven and Kaplan, 2006). In contrast, the force of dependence model integrates the demographic and economic effects of overlapping offspring across a mother's reproductive career. This permits evaluation of potential sources of help both within and exogenous to mother-offspring dyads. Second, juvenility has undergone significant hominization. To account for this we start with an ancestral juvenile who is self-sufficient at younger ages and then model effects on net cost as juveniles move toward lower levels of self-provisioning and later ages at independence and dispersal. The greatest increase in the cost of children from the ancestral state occurs when both birth intervals are short and juvenile independence is late (Table 1), a life history combination that the model suggests is not sustainable unless mothers could rely on the help of other adults. However, early life history changes generate costs that are supportable by a mother and her juvenile offspring.

Cooperation within maternal-offspring groups

Among nonhuman species, helpers are most often sexually mature kin. Consequently, explanations for cooperative breeding have focused on the indirect and direct costs and benefits for a mature individual to delay dispersal or forego reproduction to help raise another's young (Emlen, 1995; Kokko et al., 2001; Hatchwell, 2009; Sparkman et al., 2011). Attention has not been given to the importance of juvenile help in the evolution of cooperative breeding for several likely reasons. First, because juveniles do not forgo or delay their own reproduction by helping, they present less of an evolutionary puzzle than explaining why sexually mature individuals help. Second, although juveniles cooperate in some species (Ratnieks and Anderson, 1999; Clutton-Brock et al., 2000; Clutton-Brock, 2002; Crespi et al., 2009; Uematsu et al., 2010),

their help is often sporadic and unreliable (Emlen, 1990). Because juveniles are not fully grown and are less efficient foragers than adults, the marginal value of their help may be of little added benefit to their mothers' fitness. Further, since nonhuman juveniles are independent foragers, diverting time and energy from their own survival to help may incur transient costs (Russell et al., 2003) and increased mortality risks (Crespi et al., 2009). Human juveniles, however, are less constrained as helpers both because they are partially subsidized and because the human subsistence niche offers novel helping opportunities that may positively affect the marginal value of juvenile cooperation (Kramer, 2011).

A self-sufficient chimpanzee spends about 5 hrs/day foraging (Hunt, 1989; Doran, 1997), equivalent to the time that the model's early net-producer spends working. A hominin juvenile who would otherwise work hard to support himself may incur no added time cost to cooperate while also leveraging his nonreproductive status into a potential indirect fitness benefit. Human subsistence is distinguished by large foraging ranges and a broad diversity of plant, animal, and aquatic resources, most of which require processing and use of technology to access. While juveniles may be provided with difficult-to-acquire resources, they are also able to produce other easy-to-acquire resources (fruits, nuts, berries, fire wood, water, small game) that are often shared with others (Blurton Jones et al., 1994; Hawkes et al., 1995; Kramer, 2005b, 2011; Tucker and Young, 2005; Crittenden et al., 2013). Juveniles today often specialize in tasks at which they are competent but that have greater opportunity costs for older, stronger, and more skilled individuals (Kramer, 2005b, 2011). Further, because juveniles are less efficient than adults as long-distance bipeds (DeJaeger et al., 2001; Ridley and Olds, 2008), juveniles tend to perform subsistence tasks, such as food processing, that occur in or close to residential camps, while adults forage long-distance (Lee, 1979). Task specialization, the age division of labor, and

juvenile cooperation, while often overlooked in humans, are well documented in other cooperative breeders (Jarvis, 1981; Alexander et al., 1991; Thorne, 1997; Ratnieks and Anderson, 1999; Clutton-Brock et al., 2000; Clutton-Brock, 2002; Crespi et al., 2009; Uematsu et al., 2010).

Model implications to the past and fossil hominins

Much of the discussion about the evolution of cooperative breeding in humans has centered on adults, particularly the help of grandmothers and fathers (Hawkes et al., 1998; Kaplan et al., 2000; Marlowe, 2003; Flinn et al., 2008; Hill and Hurtado, 2009; van Noordwijk et al., 2013). However, in considering selection pressures from the past forward, we also have to explain what happens to juvenile subsistence effort. Juveniles transition from an ancestry of being self-supporting to a modern pattern of greater dependence. We do not expect this transition to occur rapidly since it is linked to many other traits, nor to affect all juveniles the same. Juvenility is long life stage, lasting 10 or more years during which most growth and development is achieved. Consequently, younger and older juveniles should have very different abilities as foragers and helpers. Well-dated dental formation and eruption evidence suggest that the human pattern of juvenility is slow to develop, beginning with *Homo erectus*, but not assuming a fully modern form until early *Homo sapiens* (Smith and Tompkins, 1995; Dean et al., 2001; Dean, 2006; Smith et al., 2010). Early childhood (age 3–6) is also hypothesized to emerge as a distinct developmental phase in *Homo erectus* grade species (Thorne, 1997; Clutton-Brock, 2002; Russell, 2004; Gilchrist and Russell, 2007; Bogin, 2009; Bermúdez de Castro et al., 2010). The importance of this life stage is that these weaned but very young children likely produced

little. Model results add to this discussion by showing that older juveniles can self-provision much of their own and their younger sibling's dependence under a range of life history conditions. If so, intergenerational cooperation between a mother and her juveniles may be a critical but understudied initial step in the evolution of human cooperative childrearing.

Evolutionary changes in growth and development have generated considerable interest in recent years. While generalized changes in hominin growth trajectories are well-recognized, early *Homo*, archaic *Homo* (i.e., *H. ergaster*, *H. erectus*, *H. heidelbergensis*), and early anatomically modern humans now appear far more morphologically diverse than previously thought (Gunz et al., 2009; Hammer et al., 2011). Ontogenetic differences may help to account for some of this variation. Previous research has shown that how quickly juveniles approach and asymptote (specifically the rate of change, C , and the inflection point, D , in the production function; SOM Fig. 3) at adult production levels is particularly sensitive to local food ecology and whether task difficulty is related to skill or strength (Kramer, 2005b). As hominins expanded into more and varied environments, juvenile roles also likely became more variable in foraging ranges, burden loads, mobility patterns, and group composition. These variables condition age-specific selective pressures on achieving adult levels of locomotor efficiency. These pressures have behavioral implications to the age-division of labor, juvenile dependence, and cooperation (Kramer, 2011), and may be reflected in diverse growth rates, differences in the onset of growth spurts, and age patterning discrepancies in lower limb proportionality between juveniles and adults.

Conclusions

Model results show that the demands and opportunities for cooperative breeding are sensitive to assumptions made about other life history traits. The cooperative breeding literature often supposes a modern life history as the selective background for its evolution. However, in species where life history traits that directly affect parental care have also undergone considerable modification, this assumption may mislead interpretations of causal relationships and selective pressures. This is particularly relevant for humans because juvenile dependence, late maturity, and short birth intervals are derived since the last common ancestor and likely do not reach their modern forms until the terminal Pleistocene among *Homo sapiens* (Dean, 2006). Model results indicate that these traits interact and affect demands for parental care that in some combinations incur little cost and in others a substantial cost. If hominin cooperative breeding is ancient, its evolution should be considered against this changing life history landscape.

Acknowledgements

Data collection was funded by the National Institutes of Health (AG19044-01), the Center for the Economics and Demographics of Aging, and the National Science Foundation (0349963, 0964031). Institutional support was received by Harvard University. We are grateful to Alan Rogers (University of Utah), JHE editor Mark Teaford, and the anonymous reviewers for their helpful comments. We also thank Richard Wrangham's lab group and Tuesday lunch attendees for vetting and debating earlier versions of our analysis.

Footnotes

¹ Because both breeding females and nonbreeders (pre- and post-reproductives) share in caring for and provisioning young, humans have been described as having aspects of both communal and cooperative breeders (Lukas and Clutton-Brock, 2012). Because the definition of cooperative breeding is debated, often heatedly, and we wish to avoid focusing on semantic and typological issues, we use cooperative breeding in its most general sense as a reproductive and social system in which nonparental members of the social group help to support offspring (Russell, 2004).

² We include the life history combination of juveniles dispersing before independence because in many ethnographic populations, young people may marry before they reach the asymptote in production or become net producers (Kaplan, 1997). This also occurs among some animals. Great apes do not achieve adult levels of foraging efficiency until late in juvenility or early adulthood, long after weaning and feeding independence from their mothers.

³ Other analyses show that of the daylight hours mothers spend in economic activities, about one third are net production. In addition, she may spend up to several hours per day in childcare depending on the age of her children. Further, a mother's net production increases and plateaus early in her reproductive career (Kramer, 2009).

References

- Alexander, R. D., Noonan, K. M., Crespi, B. J., 1991. The evolution of eusociality. In: Sherman, P. W., Jarvis, J. U. M., Alexander, R. D. (Eds.), *The Biology of the Naked Mole Rat*. Princeton University Press, Princeton, pp. 3–44.
- Alvarez, H. P., 2000. Grandmother hypothesis and primate life histories. *Am. J. Phys. Anthropol.* 113:435–450.
- Bermúdez de Castro, J. M., Martínón-Torres, M., Prado, L., Gómez-Robles, A., Rosell, J., López-Polín, L., Arsuaga, J. L., Carbonell, E., 2010. New immature hominin fossil from European Lower Pleistocene shows the earliest evidence of a modern human dental development pattern. *Proc. Natl. Acad. Sci.* 107:11739–11744.
- Blurton Jones, N., Hawkes, K., Draper, P., 1994. Foraging returns of !Kung adults and children: why didn't !Kung children forage? *J. Anthropol. Res.* 50:217–248.
- Bogin, B., 2009. Childhood, adolescence, and longevity: A multilevel model of the evolution of reserve capacity in human life history. *Am. J. Hum. Biol.* 21:567–577.
- Burkart, J. M., Hrdy, S. B., van Schaik, C., 2009. Cooperative breeding and human cognitive evolution. *Evol. Anthropol.* 18:175–186.
- Clutton-Brock, T. H., 2002. Breeding together: kin selection and mutualism in cooperative vertebrates. *Science* 296:69–72.
- Clutton-Brock, T. H., Brotherton, P. N. M., O'Riain, M. J., Griffin, A. S., Gaynor, D., Sharpe, L., Kansky, R., Manser, M. B., McIlrath, G. M., 2000. Individual contributions to babysitting in a cooperative mongoose, *Suricata suricatta*. *Proc. R. Soc. B* 267:301–305.
- Clutton-Brock, T. H., Harvey, P. H., 1977. Primate ecology and social organization. *J. Zool.*

183:1–39.

Creps, B., 2014. The insectan apes. *Human Nature* 25:6–27.

Crespi, B. J., Summers, K., Dorus, S., 2009. Genomic sister-disorders of neurodevelopment: an evolutionary approach. *Evol. Appl.* 2:81–100.

Crittenden, A. N., Conklin-Brittain, N. L., Zes, D. A., Schoeninger, M. J., Marlowe, F. W., 2013. Juvenile foraging among the Hadza: implications for human life history. *Evol. Hum. Behav.* 34:299–304.

Dean, M. C., 2006. Tooth microstructure tracks the pace of human life-history evolution. *Proc. R. Soc. B* 273:2799–2808.

Dean, M. C., Leakey, M. G., Reid, D., Schrenk, F., Schwartz, G. T., Stringer, C., Walker, A., 2001. Dental growth processes in teeth distinguish modern humans from *Homo erectus* and earlier hominins. *Nature* 414:628–631.

DeJaeger, D., Willems, P. A., Heglund, N. C., 2001. The energy cost of walking in children. *Euro. J. Physiol.* 441, 538–543.

Doran, D., 1997. Influence of seasonality on activity patterns, feeding behavior, ranging and grouping patterns in Tai chimpanzees. *Int. J. Primatol.* 18:183–206.

Emery Thompson, M., 2013. Reproductive ecology of female chimpanzees. *Am. J. Primatol.* 75:222–227.

Emlen, S. T., 1990. White-fronted Bee-eaters: helping in a colonially nesting species. In: Stacey, P. B., Koenig, W. D. (Eds.), *Cooperative Breeding in Birds*. Cambridge University Press, Cambridge, pp. 487–526.

Emlen, S. T., 1995. An evolutionary theory of the family. *Proc. Natl. Acad. Sci.* 92:8092–8099.

Flinn, M. V., Quinlan, R. J., Coe, K., Ward, C. V., 2008. Evolution of the human family. In:

- Salmon, C. A., Shackelford, T. K. (Eds.), Family Relationships. An Evolutionary Perspective. Oxford University Press, Oxford, pp. 16–38.
- Foster, K. R., Ratnieks, F. L. W., 2005. A new eusocial vertebrate? Trends Ecol. Evol. 20:363–364.
- Gignoux, C. R., Henn, B. M., Mountain, J. L., 2011. Rapid, global demographic expansions after the origins of agriculture. Proc. R. Acad. Sci. 108:6044–6049.
- Gilchrist, J. S., Russell, A. F., 2007. Who cares? Individual contributions to pup care by breeders vs non-breeders in the cooperatively breeding banded mongoose (*Mungos mungo*). Behav. Ecol. Sociobiol. 61:1053–1060.
- Gunz, P., Bookstein, F. L., Mitteroecker, P., Stadlmayr, A., Seidler, H., Weber, G. W., 2009. Early modern human diversity suggests subdivided population structure and a complex out-of-Africa scenario. Proc. Natl. Acad. Sci. 106:6094–6098.
- Gurven, M., Kaplan, H., 2006. Determinants of time allocation across the lifespan. Hum. Nature 17:1–49.
- Gurven, M., Kaplan, H., 2007. Longevity among hunter-gatherers: a cross-cultural examination. Popul. Develop. Rev. 33:321–365.
- Gurven, M., Walker, R., 2006. Energetic demand of multiple dependents and the evolution of slow human growth. Proc. R. Soc. B 273, 835–841.
- Hammer, M. F., Woerner, A. E., Mendez, F. L., Watkins, J. C., Wall, J. D., 2011. Genetic evidence for archaic admixture in Africa. Proc. Natl. Acad. Sci. 108:15123–15128.
- Hatchwell, B. J., 2009. The evolution of cooperative breeding in birds: kinship, dispersal and life history. Phil. Trans. R. Soc. B 364:3217–3227.
- Hawkes, K., O'Connell, J., Blurton Jones, N., 1995. Hadza children's foraging: juvenile

- dependency, social arrangements, and mobility among hunter-gatherers. *Curr. Anthropol.* 36:688–700.
- Hawkes, K., O'Connell, J., Blurton Jones, N., Alvarez, H., Charnov, E., 1998. Grandmothering, menopause and the evolution of human life histories. *Proc. Natl. Acad. Sci.* 95:1336–1339.
- Hawkes, K., Paine, R. R., 2006. The evolution of human life histories. School of American Research Advanced Seminar Series. School of American Research, Santa Fe.
- Herrmann, E., Tomasello, M., 2006. Apes' and children's understanding of cooperative and competitive motives in a communicative situation. *Develop. Sci.* 9:518–529.
- Hill, K., Hurtado, A. M., 2009. Cooperative breeding in South American hunter-gatherers. *Proc. R. Soc. B* 276:3863–3870.
- Hrdy, S. B., 2005a. Comes the child before the man: how cooperative breeding and prolonged postweaning dependence shaped human potential. In: Hewlett, B., Lamb, M. (Eds.), *Hunter Gatherer Childhoods. Evolutionary, Developmental and Cultural Perspectives*. Transaction Publishers, New Brunswick, pp. 65–91.
- Hrdy, S. B., 2005b. Evolutionary context of human development: the cooperative breeding model. In: Carter, C. S., Ahnert, L. (Eds.), *Attachment and Bonding: A New Synthesis*. MIT Press in cooperation with Dahlem University Press, Cambridge.
- Hrdy, S. B., 2007. Evolutionary context of human development: the cooperative breeding model. In: Salmon, C. A., Shackelford, T. K. (Eds.), *Family Relationships. An Evolutionary Perspective*. Oxford University Press, Oxford, pp. 39–68.
- Hrdy, S. B., 2009. *Mothers and Others*. Belknap Press/Harvard University Press, Cambridge, MA.

- Humphrey, L. T., 2010. Weaning behavior in human evolution. *Semin. Cell Dev. Biol.* 21:453–461.
- Hunt, K. D., 1989. Positional behavior in *Pan troglodytes* at the Mahale Mountains and the Gombe Stream National Parks, Tanzania. Ph.D. Dissertation, University of Michigan.
- Jarvis, J. U. M., 1981. Eusociality in a mammal: cooperative breeding in naked mole-rat colonies. *Science* 212:571–573.
- Kaplan, H., 1994. Evolutionary and wealth flows theories of fertility: empirical tests and new models. *Popul. Dev. Rev.* 20:753–791.
- Kaplan, H., 1997. The evolution of the human life course. In: K.W. Wachter, C.E. Finch (Eds.): *Between Zeus and the Salmon: The Biodemography of Longevity*, National Academy of Sciences, Washington D.C., pp. 175–211.
- Kaplan, H., Hill, K., Lancaster, J. B., Hurtado, A. M., 2000. A theory of human life history evolution: diet, intelligence, and longevity. *Evol. Anthropol.* 9:156–185.
- Kelley, J., Schwartz, G. T., 2012. Life-history inference in the early hominins *Australopithecus* and *Paranthropus*. *Int. J. Primatol.* 33:1332–1363.
- Kennedy, G. E., 2005. From the ape's dilemma to the weanling's dilemma: early weaning and its evolutionary context. *J. Hum. Evol.* 48:123–145.
- Kleiman, D. G., 1977. Monogamy in mammals. *Q. Rev. Biol.* 52:39–69.
- Knott, C., 2001. Female reproductive ecology of the apes. In: Ellison, P. T. (Ed.), *Reproductive Ecology and Human Evolution*. Aldine de Gruyter, Hawthorn, New York, pp. 429–463.
- Kokko, H., Johnstone, R. A., Clutton-Brock, T. H., 2001. The evolution of cooperative breeding through group augmentation. *Proc. R. Soc. B* 268:187–196.
- Kramer, K. L., 2005a. Children's help and the pace of reproduction: cooperative breeding in

- humans. *Evol. Anthropol.* 14:224–237.
- Kramer, K. L., 2005b. *Maya Children: Helpers at the Farm*. Harvard University Press, Cambridge, MA.
- Kramer, K. L., 2009. Does it take a family to raise a child? In: Bentley, G., Mace, R. (Eds.), *Substitute Parents. Biological and Social Perspectives on Alloparenting in Human Societies*. Berghahn Books, New York, pp. 77–99.
- Kramer, K. L., 2010. Cooperative breeding and its significance to the demographic success of humans. *A. Rev. Anthropol.* 39:414–436.
- Kramer, K. L., 2011. The evolution of human parental care and recruitment of juvenile help. *Trends Ecol. Evol.* 26:533–540.
- Lee, R. B., 1979. *The !Kung San: Men, Women and Work in a Foraging Society*. Cambridge University Press, Cambridge.
- Lee, R. D., Kramer, K. L., 2002. Children's economic roles in the Maya family life cycle: Cain, Caldwell and Chayanov revisited. *Popul. Dev. Rev.* 28:475–499.
- Lukas, D., Clutton-Brock, T., 2012. Cooperative breeding and monogamy in mammalian societies. *Proc. Roy. Soc. B* 259:2151–2156.
- Marlowe, F., 2003. A critical period for provisioning by Hadza men: implications for pair bonding. *Evol. Hum. Behav.* 24:217–229.
- Ratnieks, F. L. W., Anderson, C., 1999. Task partitioning in insect societies. *Insectes Sociaux* 47:95–108.
- Ridley, K., Olds, T. S., 2008. Assigning energy costs to activities in children: a review and synthesis. *Med. Sci. Sports Exerc.* 40:1439–1446.
- Russell, A. F., 2004. Mammals: comparisons and contrasts. In: Koenig, W. D., Dickinson, J. I.

- (Eds.), *Ecology and Evolution of Cooperative Breeding Birds*. Cambridge University Press, Cambridge, pp. 210–227.
- Russell, A. F., Lummaa, V., 2009. Maternal effects in cooperative breeders: from hymenopterans to humans. *Proc. R. Soc. B* 364:1143–1167.
- Russell, A. F., Sharpe, L. L., Brotherton, P. N. M., Clutton-Brock, T. H., 2003. Cost minimization by helpers in cooperative vertebrates. *Proc. Natl. Acad. Sci.* 100:3333–3338.
- Sear, R., Mace, R., 2008. Who keeps children alive? A review of the effects of kin on child survival. *Evol. Hum. Behav.* 29:1–18.
- Sellen, D., 2006. Lactation, complementary feeding, and human life history. In: Hawkes, K., Paine, R. R. (Eds.), *The Evolution of Human Life History*. School of American Research Press, Santa Fe, pp. 155–196.
- Smith, H. B., Tompkins, R. L., 1995. Toward a life history of the Hominidae. *A. Rev. Anthropol.* 24:257–279.
- Smith, T., Tafforeau, P., Reid, D., Pouech, J., Lazzari, V., Zermeno, J., Guatelli-Steinberg, D., Olejniczak, A., Hoffman, A., Radovic, J., Masrour, M., Toussaint, M., Stringer, C., Hublin, J.-J., 2010. Dental evidence for ontogenetic differences between modern humans and Neanderthal. *Proc. Natl. Acad. Sci.* 107:20923–20928.
- Sparkman, A. M., Adams, J. R., Steury, T. D., Waits, L. P., Murray, D. L., 2011. Direct fitness benefits of delayed dispersal in the cooperatively breeding red wolf (*Canis rufus*). *Behav. Ecol.* 22:199–205.
- Strassman, B. I., Kurapati, N. T., 2010. Are humans cooperative breeders? Most studies of natural fertility populations do not support the grandmother hypothesis. *Behav. Brain Sci.*

33:35–38.

Thorne, B. L., 1997. Evolution of eusociality in termites. *A. Rev. Ecol. Syst.* 28:27–54.

Tomasello, M., Vaish, A., 2013. Origins of human cooperation and morality. *A. Rev. Psychol.* 64:631–255.

Tucker, B., Young, A., 2005. Growing up Mikea. Children's time allocation and tuber foraging in southwest Madagascar. In: Hewlett, B., Lamb, M. (Eds.), *Hunter-Gatherer Children*. Transaction Publishers, Somerset, NJ, pp. 147–171.

Uematsu, K., Kutsukake, M., Fukatsu, T., Shimada, M., Shibao, H., 2010. Altruistic colony defense by menopausal female insects. *Curr. Biol.* 20:1182–1186.

van Noordwijk, M. A., Kuzawa, C. W., van Schaik, C. P., 2013. The evolution of the pattern of human lactation: a comparative perspective. *Evol. Anthropol.* 22:202–212.

van Schaik, C., Burkart, J. M., 2010. Mind the gap. Cooperative breeding and the evolution of our unique features. In: Kappler, P. M., Silk, J. B. (Eds.), *Mind the Gap*. Springer, Heidelberg, pp. 477–496.

Warneken, F., Hare, B., Melis, A., Hanus, D., Tomasello, M., 2007. Spontaneous altruism by chimpanzees and young children. *PLoS Biol.* 5:1414–1420.

Warneken, F., Tomasello, M., 2006. Altruistic helping in human infants and young chimpanzees. *Science* 311:1301–1303.

Figure Captions

Figure 1. Net cost of overlapping offspring across a mother's reproductive career (years after first birth). Eight profiles from the full force of dependence model (SOM Fig. 9) are taken from the end points of the parameter range for age at independence (Juv Indep), dispersal age (Disp), and birth intervals (IBI). Negative values indicate the time (daily hours) mothers or others have to spend subsidizing offspring. The stars show the point of maximum net cost. The ancestral combination of life history traits is shown in the upper left box, the most derived conditions in the bottom right.

Figure 2. Maximum net cost of overlapping offspring stratified by 6- to 3-year intervals (IBI). Surfaces show the influence of age at juvenile independence (y-axis), dispersal age (x-axis), and birth intervals, and are the composite of the maximum net cost points from each of the 40,000 profiles in the full model (SOM Fig. 9). Color gradients follow the z-axis, where maximum net cost increases from red to blue. See SOM Figure 10 for an interactive rotational model display.

Figure 3. Linear combinations of the maximum net cost as a function of dispersal age and age at juvenile independence across 6- to 3-year birth intervals (IBI). The top row illustrates the effect of a later dispersal age (14–20) at early (age 10) and late (age 20) age of juvenile independence. The bottom row shows the effect of increasing age at juvenile independence (10–20) at early (age 14) and late (age 20) dispersal age. Slopes estimates given in Table 2.

Table 1

Change in cost of offspring (maximum net cost) from the ancestral state of early juvenile independence (age 10), early dispersal (age 14), and a 6-year birth interval^a

Sensitivity				
Juvenile Independence		Dispersal		IBI
		Early (Age 14)	Late (Age 20)	
Early		0.00	0.00	6 yrs
Late		-5.05	-6.48	
Early		-0.21	-0.21	5 yrs
Late		-6.30	-8.30	
Early		-1.22	-1.22	4 yrs
Late		-8.63	-11.08	
Early		-2.38	-2.39	3 yrs
Late		-12.36	-15.73	
Elasticity				
Juvenile Independence		Dispersal		IBI
		Early (Age 14)	Late (Age 20)	
Early		0%	0%	6 yrs
Late		158%	202%	
Early		6%	6%	5 yrs
Late		197%	259%	
Early		38%	38%	4 yrs
Late		270%	346%	

Early	74%	74%	3 yrs
Late	386%	491%	

^a Table values give the difference between the maximum net cost at the end points of the parameter range for juvenile independence (from age 10 to age 20) and dispersal (age 14 to age 20), and from a 6-year to a 5-, 4-, and 3-year birth interval. These end points are represented at the corner of the layers in Figure 2. Sensitivity is the difference in raw values and elasticity is sensitivity expressed as a percentage of the first value. The marginal value, or total effect of each variable holding the others constant, is calculated as the sum of the difference of sums in the parameter end points shown in this table. Marginal value for juvenile independence = 66.32, birth interval = 21.32, and dispersal = 9.26.

Table 2

The daily hourly increase in investment needed to support offspring (maximum net cost) for every year dispersal age and age of juvenile independence increase while holding the other variable constant at its maximum and minimum value, and stratified by birth interval (IBI)^a

Dispersal		Juvenile Independence		IBI
Early net prod	Late net prod	Early dispersal	Late dispersal	
(Age 10)	(Age 20)	(Age 14)	(Age 20)	
0.00	.24	.51	.65	6 yrs
0.00	.33	.61	.81	5 yrs
0.00	.41	.74	.99	4 yrs
0.00	.56	1.00	1.33	3 yrs

^a Values are slopes derived from the linear combinations in Figure 3 (SOM Appendix 2). Because the original maximum net costs are negative, slope values are multiplied by -1 to express more intuitively as the daily hourly change from one life history condition to another.

Figure 1

Figure 2

Figure 3