Drained coastal peatlands: A potential nitrogen source to marine ecosystems under prolonged drought and heavy storm events—A microcosm experiment

Hongjun Wang, Curtis J. Richardson, Mengchi Ho and Neal Flanagan

Duke University Wetland Center, Nicholas School of the Environment, Box 90333, Duke University,

Durham, North Carolina 27708 USA

Corresponding author: Hongjun Wang, E-mail:hw93@duke.edu, Tel: 919-613-8009

Abstract

Over the past several decades there has been a massive increase in coastal eutrophication, which is often caused by increased runoff input of nitrogen from landscape alterations. Peatlands, covering 3% of land area, have stored about 12-21% of global soil organic nitrogen (12-20 Pg N) around rivers, lakes and coasts over millennia and are now often drained and farmed. Their huge nitrogen pools may be released by intensified climate driven hydrologic events— prolonged droughts followed by heavy storms—and later transported to marine ecosystems. In this study, we collected peat monoliths from drained, natural, and restored coastal peatlands in the Southeastern U.S., and conducted a microcosm experiment simulating coupled prolonged-drought and storm events to (1) test whether storms could trigger a pulse of nitrogen export from drought-stressed peatlands and (2) assess how differentially hydrologic managements through shifting plant communities affect nitrogen export by combining an experiment of nitrogen release from litter.

During the drought phase, we observed a significant temporal variation in net nitrogen mineralization rate (NMR). NMR spiked in the third month and then decreased rapidly. This pattern indicates that drought duration significantly affects nitrogen mineralization in peat. NMR in the drained site reached up to $490 \pm 110 \text{ kg ha}^{-1} \text{ yr}^{-1}$, about 5 times higher than in the restored site. After the 14-month drought phase, we simulated a heavy storm by bringing peat monoliths to saturation. In the discharge waters, concentrations of total dissolved nitrogen in the monoliths from the drained site ($72.7 \pm 16.3 \text{ mg L}^{-1}$) was about ten times as high as from the restored site. Our results indicate that previously drained peatlands under prolonged drought are a potent source of nitrogen export. Moreover, drought-induced plant community shifts to herbaceous plants substantial raise nitrogen release with lasting effects by altering litter quality in peatlands.

Keywords: Drought, eutrophication, nitrogen mineralization, peatlands, plant community, storm

1. Introduction

Massive volumes of chemical nitrogen production via the Haber-Bosch process have been applied to the landscape following the industrial revolution, causing many natural ecosystems which previously acted as nitrogen sinks to become nitrogen sources to downstream waters after saturation (Aber, 1992; Bragazza et al., 2006; Lamers et al., 2000). Overloading of nitrogen is a leading cause of impairment of coastal marine ecosystems around the world (Boesch et al., 2001; Compton et al., 2011; Diaz and Rosenberg, 2008; Paerl and Piehler, 2008). Currently, most studies focus on anthropogenic nitrogen sources, like fertilizer applied in farmlands and fossil-fuel derived atmospheric deposition (Cui et al., 2013; Liu et al., 2013; Paerl, 1997; Paerl, 2009; Robert, 2008). However, chronically-accumulated organic nitrogen in natural ecosystems can potentially be quickly released and become a substantial source of nitrogen pollution, exacerbating degradation of aquatic ecosystems when environmental conditions, such as climatic warming and drought, change to favor nitrogen mineralization. Studies have shown large leaching losses of nitrogen from forests (Fang et al., 2009; Perakis and Hedin, 2002). But there is another important natural ecosystem—peatland that stores not only 1/3 of global soil carbon but also 12-21% of global soil organic nitrogen (12-20 Pg N) (Joosten, 2010; Limpens et al., 2006). Some peatlands have already lost their ability to absorb nitrogen due to an overload of nitrogen (Lamers et al., 2000; Qualls and Richardson, 2003). Moreover, most peatlands are located around rivers, lakes and coasts, where they may be a potentially significant nitrogen source to the adjacent aquatic ecosystems. To-date, the internal cycle of nitrogen, which is sensitive to climate change, is still poorly understood in peatlands, although its impact on adjacent downstream ecosystems may be substantial (Vassiljev and Blinova, 2012; Verhoeven et al., 1988; Wray and Bayley, 2008).

1 Soil nitrogen in peat is highly concentrated and is about 10–40 times higher than
2 concentrations found in mineral soils (Soper and Osbon, 1922). Generally, nitrogen
3 mineralization is slow under waterlogged conditions and almost all mineralized nitrogen is
4 recycled within peatlands (Lamers et al., 2000). However, if water level drops, making more
5 oxygen available in the soil, nitrogen mineralization could be accelerated, and huge amounts of
6 available nitrogen can be released (Qualls and Richardson, 2003). For example, nitrogen
7 mineralization rate found in Histosols in New York was more than $500 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ (Guthrie
8 and Duxbury, 1978), which is higher than the amount of chemical nitrogen fertilizer used in
9 intensive farming (Ju et al., 2009). Such potentially high available nitrogen in soil is the main
10 reason why large portions of peatlands were drained and reclaimed as farmlands for the past two
11 hundred years (Richardson, 1983, 2008; Soper and Osbon, 1922). In Estonia, export of nitrogen
12 from drained peatlands was more than 1.5-fold higher than from farmlands and was the major
13 source of nitrogen export to the surface water (Vassiljev and Blinova, 2012).

14 Both climatic and human disturbance have been substantially lowering peatland water
15 levels and already have driven more than 11% of global peatlands into a degraded state (IPCC,
16 2007; Joosten, 2010). The altered hydrological conditions not only directly alter the rate of
17 nitrogen mineralization by changing the availability of oxygen but also the composition of
18 organic nitrogen for mineralization by gradually changing the dominant plant communities that
19 may even have longer-lasting effects on the nitrogen cycle (Laiho et al., 2003). Importantly,
20 drought durations are projected to increase in the future (IPCC, 2007), which could substantially
21 increase the available nitrogen pool in peatlands. Therefore, large quantities of nitrogen in
22 peatlands would likely be released to freshwater bodies during heavy storms and subsequently

enter coastal estuarine and marine ecosystems with negative consequences (Boesch et al., 2001; Diaz and Rosenberg, 2008; Paerl and Piehler, 2008).

In this study, we collected peat monoliths from drained, natural and restored coastal peatlands in the Southeastern U.S and conducted a microcosm experiment. We simulated prolonged drought and heavy storms in lab to test 1) whether intensive drought and storms can cause increased export of nitrogen from drought-stressed peatlands and 2) how drought duration and long-term hydrologic conditions (drainage and restoration versus natural hydrologic regimes) affect this process. We compared these outputs with measured regional estimates of nitrogen losses from peatlands from the same area.

2. Material and methods

2.1 Study sites

Along the south Atlantic coast, peatlands cover millions of hectares and store over 20% of peat in the continental USA (Richardson, 2012; Soper and Osbon, 1922). These temperate and subtropical peatlands, called pocosins, are generally dominated by shrubs and trees. Our study area is located at Pocosin Lakes National Wildlife Refuge (PLNWR) in North Carolina and is about 8 km away from estuarine and marine wetlands. This site has a warm, humid climate with an average annual temperature of 16.8 °C. Annual precipitation is around 1230 mm and about 800 mm is lost to evapotranspiration (ET). Generally, the lowest and highest water levels occur in summer and winter, respectively. Throughout the year, groundwater levels rarely rise above ground surface and often reach 20 cm below the surface, but fall to more than 100 cm in depth in the summer(Wang et al., 2015).

Most of the pocosins were drained for farming between 1920's and 1940's (McMullan Jr, 1983). Currently, there are a series of distinct hydrologic units (800 m × 1600 m cells) divided

by ditches that have been used by PLNWR to restore the water level since the 1990s. In this study natural, drained, and restored sites were selected for treatments. The natural site experiences natural hydrologic conditions. The water level ranges from 0–60 cm below the ground surface during the winter and greater than 100 cm during the summer. Mature canopy trees—including pond pine (*Pinus serotina* Michx.), loblolly bay (*Gordonia lasianthus* (L.) Ellis), fetterbush lyonia (*Lyonia lucida* (Lam.) K. Koch), and swamp bay (*Persea palustris* (Raf.) Sarg.)—cover about 80% of the natural site. The water level in the drained site is mostly below 50 cm depth and greater than 150 cm in summer. About 80% of the ground area in the drained site is covered by western brackenfern (*Pteridium aquilinum* (L.) Kuhn). In the restored site, the water level is 20–30 cm below ground surface and native shrubs are dominant species, including large gallberry (*Ilex coriacea* (Pursh) Chapm.), inkberry (*Ilex glabra* (L.) A. Gray), fetterbush lyonia, honeycup (*Zenobia pulverulenta* (W.Bartram ex. Willd.)Pollard), and laurel greenbrier (*Smilax laurifolia* L.).

2.2 Microcosm experiment

We collected triplicate peat monoliths from the natural, drained, and restored pocosins in January 2011. In the field, peat monoliths (32 cm in diameter) were excavated in 10-cm increments to a depth of 40 cm and was transferred to a top-open plastic incubator (30-cm diameter, 37.5-cm depth). All vegetation, fresh litter, live roots and rhizomes (>1 mm diameter) were removed. We incubated these monoliths at a constant temperature of 25 °C in the lab creating a gradual loss of soil water by evaporation to simulate a prolonged drought. After a 14-month drought, we rewetted these monoliths to a saturated state to simulate a heavy-storm event (160 mm).

During the drought phase, we determined net nitrogen mineralization rate (NMR) that was calculated from the changes of total inorganic nitrogen ($\text{NH}_4^{+}\text{-N}$ and $\text{NO}_x\text{-N}$) in the monolith as a function of time and area. Soil cores (3.5-cm diameter) were collected from the monoliths on 30th, 60th, 91st, 152nd and 415th d. The core was sliced into four segments (0–5, 5–15, 15–25, and 25–38 cm). Each segment was mixed and divided into three subsamples in which one was extracted using 2 M KCl and filtered by a 0.22- μm nylon filter (Micron Separations INC, MA, USA) on the first day, one was dried by oven at 60 °C for 24 h, and the last was stored in a 4 °C cold room. Soil pH was measured with a Beckman 360 pH meter, using a soil:water ratio of 1:10. We measured soil water content for each fresh sample, and all soil properties were calculated based on dry weight. On the 4th day of rewetting, we collected the leached water from the incubators and measured pH, total inorganic nitrogen and total organic nitrogen.

2.3 Potential release of nitrogen from litter

To test whether the historic hydrological conditions have potential long-lasting effects on nitrogen release due to changing plant communities with different quality of litter, we collected senesced leaves and small branches from dominant species in the natural, drained, and restored sites in 2011 and conducted a water-extraction experiment. The leaves and branches were air dried first and then put in an oven at 60 °C for 48 h. About 5 g of dry sample was ground and passed through a 250 μm (No. 60) sieve. We put 0.25–0.30 g of sample into 15 ml deionized water, mixed them by a shaker for 48 h at 25 °C. The suspension liquid was filtered through a glass fiber filter (0.5 μm , GE Water & Process Technologies, USA) and stored at 4 °C in a refrigerator for analysis of total dissolved nitrogen (TDN) and dissolved organic carbon (DOC).

2.4 Chemical analysis

DOC in the discharged water and water extraction were both determined as the difference between total C and inorganic C by a total C analyzer (Shimadzu 5000A, Kyoto, Japan). Inorganic nitrogen (NH_4^+ and NO_x^- ($\text{NO}_3^- + \text{NO}_2^-$)) in the KCl extraction and discharged water was measured colorimetrically using a flow-injection analyzer (Lachat QuikChem 8000, Wisconsin, USA). Total dissolved nitrogen (TDN) in the discharged water and water extraction was determined by a total organic carbon analyzer with a total nitrogen measuring unit (Shimadzu TOC-L, Kyoto, Japan). Dissolved organic nitrogen (DON) was calculated based on the difference between total dissolved nitrogen and total inorganic nitrogen. Total carbon and nitrogen from the dried and milled soil samples were directly analyzed with combustion CN soil analyzer equipped with a TCD detector (ThermoQuest Flash EA1112, Milan, Italy).

2.5 Statistical analysis

We tested the parameter differences among sites by one-way ANOVA using SAS 9.3. Results were considered significant at the 95% confidence ($\alpha=0.05$). Standard error was calculated for means and error bars.

3. Results

3.1 Effects of drought

During the 14-month drought experiment, soil moisture (SM) gradually decreased, and finally about 70 to 80% of water in soil was lost in all peat monoliths (Fig. 1a). SM was always higher in the monoliths from the restored

Figure 1. Temporal variation of soil moisture (a), soil pH (b), $\text{NO}_3^- \text{N}$ (c) and $\text{NH}_4^+ \text{N}$ (d) during the drought phase in the peat monoliths from the natural, drained, and restored sites in the pocosins of coastal North Carolina.

site than those from other sites. Peat monoliths from the natural site showed higher soil pH than those from the drained and the restored sites (Fig. 1b). Along with the increasing drought stress, soil pH responded differently among peat monoliths from the different sites. At the natural site, soil pH increased first and then gradually decreased. At the restored site, it increased initially, then sharply decreased during days 60–150 and slowly increased after 150 days. At the drained site, it decreased first and gradually increased afterwards. We observed the highest extractable NO_x^- -N (Fig. 1c) in the peat monoliths from the drained site and the highest extractable NH_4^+ -N from the natural site (Fig. 1d). The concentration of NH_4^+ -N was much higher than NO_x^- -N in all peat monoliths, and the ratios of NH_4^+ -N to NO_x^- -N were 11.2 ± 5.4 , 3.6 ± 1.1 and 2.1 ± 0.3 in the natural, restored, and drained sites, respectively. Peat monoliths from the restored site always contained the lowest extractable NO_x^- -N and NH_4^+ -N (Fig. 1c, d). Both extractable NO_x^- -N and NH_4^+ -N decreased in the second month of drought, then quickly increased in the third month, and later increased slowly in all peat monoliths except that NO_x^- -N did not change in the drained site after 5 months (Fig. 1c,d). Based on the temporal changes of extractable inorganic nitrogen (NO_x^- -N + NH_4^+ -N), net nitrogen

Figure 2. Net N mineralization during the drought phase in the peat monoliths from the natural, drained, and restored sites in the pocosins of coastal North Carolina.

mineralization rates (NMR) were calculated. We found similar temporal variations of NMR from all sites (Fig. 2). NMR was negative at the beginning, spiked in the third month, and significantly decreased afterwards. The increased inorganic nitrogen in the third month represented 23%, 27%

and 77% of the total mineralized nitrogen in the monoliths from restored, natural, and drained sites during 14-month drought period. In the drained site, NMR reached $490 \pm 110 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ in the third month, which was about 5 times higher than that in the restored site.

3.2 Effects of rewetting

We found high concentrations of total dissolved nitrogen ($\text{DON} + \text{NO}_x^- \text{-N} + \text{NH}_4^+ \text{-N}$) in the discharge water— 72.7 ± 16.3 , 17.7 ± 7.1 , and $7.0 \pm 1.4 \text{ mg L}^{-1}$ —in peat monoliths from the drained, natural, and restored sites, respectively (Fig. 3). Peat monoliths from the drained site also released the highest DON and $\text{NO}_x^- \text{-N}$, which were about 20 and 5 times as high as those from the restored site. DON represented about 68% of total dissolved nitrogen in the peat monolith from the drained site and 36% from the restored site (Fig. 3). Concentrations of $\text{NH}_4^+ \text{-N}$ in the discharged water from all peat monoliths were similar, although these

Figure 3. The concentrations of dissolved organic nitrogen (DON), $\text{NH}_4^+ \text{-N}$ and $\text{NO}_x^- \text{-N}$ in the discharged water from the peat monoliths from the natural, drained, and restored sites in the pocosins of coastal North Carolina during the rewetting phase.

Figure 4. The ratio of dissolved organic carbon (DOC) to dissolved organic nitrogen (DON) in the discharge water from the peat monoliths from the natural, drained and restored sites in the pocosins of coastal North Carolina. Different letters indicate a significant difference

peat monoliths contained different amount of $\text{NH}_4^+\text{-N}$ (Fig. 1d and Fig.3). The ratio of DOC to DON was 11 times higher in the restored site (42.8 ± 7.8) than in the drained site (Fig. 4).

3.3 Potential release of nitrogen from litter

Total nitrogen (TN) in leaves of bracken fern in the drained site (about 2% of dry biomass) was two times as high as that in leaves from shrub/tree in the natural and restored sites. Other than leaves, woody materials (branches, cones and eventually tree trunks) were major sources of litter in the natural and restored sites. All small branches we collected contained much lower TN ranging from 0.25% to 0.61%.

Based on the water-extraction experiment, the leaves of bracken fern ($0.84\pm0.16 \text{ mg}$

$\text{N g}^{-1} \text{ dw}$) released much higher total dissolved nitrogen compared to the leaves ($0.17\pm0.02 \text{ mg N g}^{-1} \text{ dw}$) and branches ($0.08\pm0.01 \text{ mg N g}^{-1} \text{ dw}$) from the shrubs/trees in the natural and restored sites (Fig. 5a). Consistent with the rewetting experiment, the water extraction from the leaves of bracken fern in the drained site showed a lower ratio of DOC to DON (Fig. 5b) than leaves of shrubs/trees in the natural and restored sites.

4. Discussion

All plant-free peat monoliths in this experiment were incubated under uniform temperatures. Therefore, any changes in nitrogen mineralization resulted from the simulated drought and pre-existing (>20 years) hydrological conditions. Most previously studied drought

Figure 5. TDN—Total dissolved nitrogen (a) and the ratio of dissolved organic carbon (DOC) to TDN (b) from senescent leaves and shrub branches in the natural, drained and restored sites in pocosins of coastal North Carolina. Different letters indicate a significant difference

1 incubation experiments were relatively short-term manipulations, between several days and 1–2
2 months (Estop-Aragonés and Blodau, 2012; Fierer and Schimel, 2002; Venterink et al., 2002). In
3 this experiment, we employed a 14-month drought, which is a rare but realistic event in our field
4 situation, thus conducive to detecting an effect of long-term drought durations (Wang et al. 2015).
5 The strong temporal variations of NMR (Fig.2) implied that drought duration has substantial
6 effects on the pool of available nitrogen in peatlands. At the beginning of the drought, both NO_x^-
7 $-\text{N}$ and NH_4^+-N decreased (Fig. 1c,d), resulting in negative NMRs. Venterink et al (2002) also
8 found negative NMR during the first 12 days of drought manipulation (Venterink et al., 2002).
9 The negative NMRs implied that the growth of microbes in our study area was still limited by
10 nitrogen and the available inorganic nitrogen was rapidly assimilated into microbial biomass
11 during short-term drought, and this also indicated that short-term drought may only have minor
12 impacts on nitrogen loss in N-limited peatlands. However, NMR sharply increased during the
13 third month (Fig.2). Although such spikes were of short duration, the mineralized nitrogen had
14 major contributions to the accumulated inorganic nitrogen for the 14- month drought period.
15 Therefore, if drought lasts longer, often over 2 months in pocosin (Richardson, 2012; Richardson
16 and McCarthy, 1994), substantial nitrogen will be quickly released. The further reduction of soil
17 moisture possibly limited nitrogen mineralization, resulting in much lower NMR afterward this
18 period. Similar to what other studies found (Dominik and Jörg, 2007; Frank et al., 2014; Laine et
19 al., 2013), ammonium was the dominant inorganic nitrogen and gradually accumulated in peat
20 during drought (Figs. 1c,d). Although the peat soils in the monoliths were under aerobic
21 conditions, nitrification rates were low except in drained and natural sites during the third month.
22 Low pH (Fig. 1b) in our experiment might have restricted nitrification activity (Bridgham et al.,
23 2001), and hence the internal cycling of inorganic N in peatlands largely led to NH_4^+-N buildups.

As expected, heavy storm simulations triggered substantial nitrogen loss from the peat monoliths after a prolonged drought. The previously drained peat released the highest nitrogen. Importantly our results are consistent with field observation in pocosins in 1999 (Shelby et al., 2005). In 1999 they experienced a very dry late winter, spring and summer followed by an unprecedented series of heavy storms (555 mm in September and October). Total nitrogen export from a forested subwatershed in the pocosins was 10 kg ha^{-1} during the storm, which is nearly as high as total loss of nitrogen for an average year. The total nitrogen concentration observed in a nearby drained forested site were abnormally high (12.3 mg L^{-1}) compared to other undrained pocosin sites ($0.9\text{-}1.6 \text{ mg L}^{-1}$) (Chescheir et al., 2003; Shelby et al., 2005). We found similar results when we compared the monoliths from the drained, natural and restored sites, the monoliths from the restored and natural sites had much lower risk of TDN loss even after a 14-month drought. Such differences between the drained and restored sites suggest that 1) long-term hydrological conditions change the internal nitrogen cycle with long-lasting impacts in peatlands and 2) drainage/long-term drought makes peatlands more fragile and sensitive to further drought than saturated peatlands. Based on the hydrology and water quality data spanning 25 years (1976-2000) in more than 100 forest sites around the pocosin region (Chescheir et al., 2003), the drained organic soil sites with high hydraulic conductivities released the highest dissolved inorganic nitrogen — $10.1 \text{ kg ha}^{-1} \text{ yr}^{-1}$ which was about 15-50 times higher than the unditched natural organic soil site. Compared to the natural and restored sites, higher total nitrogen in senescent leaves of herbaceous plants from the drained sites indicates that recent litter can supply more nitrogen to the drained peatlands and plant community over there can assimilate more nitrogen. Our water extraction experiment further demonstrated that herb and fern litter in a drained peatland can also release more nitrogen (Fig. 5a). Herbaceous species in drained

1 peatlands can assimilate more inorganic nitrogen temporally, however, their litter with low ratio
2 of carbon to nitrogen is easy to be decomposed and releases more inorganic nitrogen not only to
3 downstream ecosystem during storm but also to the old peat in deeper layers to stimulate the
4 mineralization of nitrogen historically stored. Therefore, drainage/drought-induced plant
5 community shifts to more herbaceous species change the quality of litter, which may
6 substantially affect the nitrogen transformation processes in peatlands.

7 Additionally, all peat monoliths contained much more $\text{NH}_4^+\text{-N}$ than $\text{NO}_x^-\text{-N}$ at the end of
8 the drought phase experiment (Fig. 1c, d). However, in the discharge water, the concentration of
9 $\text{NH}_4^+\text{-N}$ was lower than $\text{NO}_x^-\text{-N}$ in the drained site and similar in the natural and restored sites
10 (Fig. 3). Therefore, a large portion of $\text{NH}_4^+\text{-N}$ remains bound to cation exchange sites on the soil
11 complex, which suggests storms can only flush out a small amount of $\text{NH}_4^+\text{-N}$ to the
12 downstream. In line with other studies (Bragazza and Limpens, 2004; Frank et al., 2014; Qualls
13 and Richardson, 2003), DON was the dominant form of nitrogen in the discharge water (Fig. 3).
14 The monoliths from the drained site showed unexpectedly high DON concentration (Fig. 3),
15 which suggest drainage not only enhances the release of nitrate but also DON. Peat in the
16 drained site contained higher inorganic nitrogen by increased nitrogen mineralization, which
17 possibly promote microbial dissolution of organic compounds and hence increase the production
18 of DON like amino acids (released during the hydrolysis of proteins) (Kalbitz and Geyer, 2002;
19 Qualls, 2000). In our study, the ratio of DOC to DON was lower in the drained and natural sites
20 than in the restored site, which suggest a preferential release of DON (Kalbitz and Geyer, 2002)
21 from the natural and drained sites (Figs. 3, 4). This was consistent with our water extraction
22 experiment in which leaves from the drained site also showed a lower ratio of DOC/TDN.

1 In conclusion, our findings showed that prolonged drought followed by heavy storms leads
2 to a high risk of nitrogen export from peatlands, especially from drained/degraded peatlands, to
3 downstream coastal waters. We also found that drainage/drought-induced plant community shifts
4 change the litter quality with substantial effects on the internal nitrogen cycle, and thus may
5 have long-lasting impacts on the response of nitrogen transformations to climate induced drought
6 changes. Our results also indicate that restoring drained/degraded peatlands by raising water
7 levels is conducive to conserving nitrogen. However, water levels in drained or degraded
8 peatlands should be increased gradually during restoration to leave enough time for nitrogen
9 immobilization and uptake by adapted microbe and plants in wetter conditions, otherwise,
10 enormous amounts of nitrogen may be quickly released from drained peatlands after complete
11 rewetting or a heavy storm.

12 **Acknowledgments**

13 We thank Hans W. Paerl at University of North Carolina for valuable comments. We thank
14 Wes Willis for the lab measurements, and Randy Neighbarger for technical editing. Thanks also
15 go to Sara Ward and Dave Kitts, USFWS, for helping with site selections and field assistance.
16 Coastal Carolina/ Southeastern Virginia Strategic Habitat Conservation Team, U.S. Fish and
17 Wildlife Service–Region 4, US DOE Office of Science, Terrestrial Ecosystem Sciences, under
18 grant award DE-SC0012272, The Nature Conservancy of North Carolina and the Duke
19 University Wetland Center Endowment provided financial support.

20 **References**

21 Aber, J., 1992. Nitrogen cycling and nitrogen saturation in temperate forest ecosystems. Trends
22 in ecology & evolution 7, 220-224.

1 Boesch, D., Burreson, E., Dennison, W., Houde, E., Kemp, M., 2001. Factors in the decline of
2 coastal ecosystems. *Science* 293, 1589-1591.

3 Bragazza, L., Freeman, C., Jones, T., Rydin, H., Limpens, J., Fenner, N., Ellis, T., Gerdol, R.,
4 Hájek, M., Hájek, T., Iacumin, P., Kutnar, L., Tahvanainen, T., Toberman, H., 2006.
5 Atmospheric nitrogen deposition promotes carbon loss from peat bogs. *Proceedings of the*
6 *National Academy of Sciences of the United States of America* 103, 19386-19389.

7 Bragazza, L., Limpens, J., 2004. Dissolved organic nitrogen dominates in European bogs under
8 increasing atmospheric N deposition. *Global Biogeochemical Cycles*, 18, GB4018,
9 doi:10.1029/2004GB002267.

10 Bridgham, S.D., Updegraff, K., Pastor, J., 2001. A Comparison of Nutrient Availability Indices
11 Along an Ombrotrophic–Minerotrophic Gradient in Minnesota Wetlands. *Soil Science*
12 *Society of America Journal* 65, 259-269.

13 Chescheir, G.M., Lebo, M.E., Amatya, D.M., Hughes, J., Gilliam, J.W., Skaggs, R.W.,
14 Herrmann, R.B., 2003. Hydrology and water quality of forested lands in eastern North
15 Carolina. North Carolina State University, Raleigh, N.C., p. 79.

16 Compton, J., Harrison, J., Dennis, R., Greaver, T., Hill, B., Jordan, S., Walker, H., Campbell, H.,
17 2011. Ecosystem services altered by human changes in the nitrogen cycle: a new
18 perspective for US decision making. *Ecology letters* 14, 804-815.

19 Cui, S.H., Shi, Y.L., Groffman, P.M., Schlesinger, W.H., Zhu, Y.G., 2013. Centennial-scale
20 analysis of the creation and fate of reactive nitrogen in China (1910-2010). *Proceedings of*
21 *the National Academy of Sciences of the United States of America* 110, 2052-2057.

22 Diaz, R., Rosenberg, R., 2008. Spreading dead zones and consequences for marine ecosystems.
23 *Science (New York, N.Y.)* 321, 926-929.

1 Dominik, Z., Jörg, G., 2007. The mobilisation of phosphorus, organic carbon and ammonium in
2 the initial stage of fen rewetting (a case study from NE Germany). *Biogeochemistry* 85,
3 141-151.

4 Estop-Aragonés, C., Blodau, C., 2012. Effects of experimental drying intensity and duration on
5 respiration and methane production recovery in fen peat incubations. *Soil Biology and*
6 *Biochemistry* 47, 1-9.

7 Fang, Y.T., Zhu, W.X., Gundersen, P., Mo, J.M., Zhou, G.Y., Yoh, M., 2009. Large Loss of
8 Dissolved Organic Nitrogen from Nitrogen-Saturated Forests in Subtropical China.
9 *Ecosystems* 12, 33-45.

10 Fierer, N., Schimel, J.P., 2002. Effects of drying-rewetting frequency on soil carbon and nitrogen
11 transformations. *Soil Biology & Biochemistry* 34, 777-787.

12 Frank, S., Tiemeyer, B., Gelbrecht, J., Freibauer, A., 2014. High soil solution carbon and
13 nitrogen concentrations in a drained Atlantic bog are reduced to natural levels by 10 years
14 of rewetting. *Biogeosciences* 11, 2309-2324.

15 Guthrie, T., Duxbury, J., 1978. Nitrogen mineralization and denitrification in organic soils. *Soil*
16 *Science Society of America Journal* 42, 908-912.

17 IPCC, 2007. *Climate Change 2007: The Physical Science Basis. Contribution of Working Group*
18 *I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change.*
19 *Cambridge University Press, Cambridge and New York.*

20 Joosten, H., 2010. *The Global Peatland CO₂ Picture. Peatland status and drainage related*
21 *emissions in all countries of the world, Ede, Netherlands, p. 36.*

22 Ju, X.T., Xing, G.X., Chen, X.P., Zhang, S.L., Zhang, L.J., Liu, X.J., Cui, Z.L., Yin, B., Christie,
23 P., Zhu, Z.L., Zhang, F.S., 2009. Reducing environmental risk by improving N

management in intensive Chinese agricultural systems. Proceedings of the National Academy of Sciences of the United States of America 106, 3041-3046.

Kalbitz, K., Geyer, S., 2002. Different effects of peat degradation on dissolved organic carbon and nitrogen. Organic Geochemistry 33, 319-326.

Laiho, R., Vasander, H., Penttilä, T., Laine, J., 2003. Dynamics of plant-mediated organic matter and nutrient cycling following water-level drawdown in boreal peatlands. Global Biogeochemical Cycles 17 1053, doi:10.1029/2002GB002015.

Laine, M.P.P., Strömmer, R., Arvola, L., 2013. Nitrogen Release in Pristine and Drained Peat Profiles in Response to Water Table Fluctuations: A Mesocosm Experiment. Applied and Environmental Soil Science 2013, 7.

Lamers, L.P.M., Bobbink, R., Roelofs, J.G.M., 2000. Natural nitrogen filter fails in polluted raised bogs. Global Change Biology 6, 583-586.

Limpens, J., Heijmans, Monique M., Berendse, F., 2006. The nitrogen cycle in boreal peatlands, In: Wieder, R.K., Vitt, Dale H. (Eds.), Boreal Peatland Ecosystems. Springer Verlag, Berlin, pp. 195-230.

Liu, X.J., Zhang, Y., Han, W.X., Tang, A.H., Shen, J.L., Cui, Z.L., Vitousek, P., Erisman, J.W., Goulding, K., Christie, P., Fangmeier, A., Zhang, F.S., 2013. Enhanced nitrogen deposition over China. Nature 494, 459-462.

McMullan Jr. P.S. 1983. Land-clearing trends on the Albemarle-Pamlico Peninsula 300 years of development in a wetlands region. McMullan Consulting, Durham, NC, USA. p. 132.

Paerl, H.W., 1997. Coastal eutrophication and harmful algal blooms: Importance of atmospheric deposition and groundwater as "new" nitrogen and other nutrient sources. Limnology and Oceanography 42, 1154-1165.

1 Paerl, H.W., 2009. Controlling eutrophication along the freshwater–marine continuum: dual
2 nutrient (N and P) reductions are essential. *Estuaries and Coasts* 32, 593-601.

3 Paerl, H.W., Piehler, M.F., 2008. Nitrogen and Marine Eutrophication, In: Capone, D.G.,
4 Mulholland, M., Carpenter, E. (Eds.), *Nitrogen in the Marine Environment*. Academic
5 Press, Orlando, pp. 529-567.

6 Perakis, S., Hedin, L., 2002. Nitrogen loss from unpolluted South American forests mainly via
7 dissolved organic compounds. *Nature* 415, 416-419.

8 Qualls, R.G., 2000. Comparison of the behavior of soluble organic and inorganic nutrients in
9 forest soils. *Forest Ecology and Management* 138, 29-50.

10 Qualls, R.G., Richardson, C.J., 2003. Factors controlling concentration, export, and
11 decomposition of dissolved organic nutrients in the Everglades of Florida. *Biogeochemistry*
12 62, 197-229.

13 Richardson, C.J., 1983. Pocosins: vanishing wastelands or valuable wetlands? *BioScience* 33,
14 626-633.

15 Richardson, C.J., 2008. *The Everglades Experiments*. Springer, New York, doi 10, 978-970.

16 Richardson, C.J., 2012. Pocosins: Evergreen shrub bogs of the southeast, In: Batzer, D.B., A.
17 (Ed.), *Wetland Habitats of North America*. University of California Press, pp. 189-202.

18 Richardson, C.J., McCarthy, E.J., 1994. Effect of Land-Development and Forest Management on
19 Hydrologic Response in Southeastern Coastal Wetlands - a Review. *Wetlands* 14, 56-71.

20 Robert, W.H., 2008. Coastal nitrogen pollution: A review of sources and trends globally and
21 regionally. *Harmful Algae* 8, 14-20.

- 1 Shelby, J., Chescheir, G., Skaggs, R., Amatya, D., 2005. Hydrologic and water-quality response
2 of forested and agricultural lands during the 1999 extreme weather conditions in eastern
3 North Carolina. *Transactions of the Asae* 48, 2179-2188.
- 4 Soper, E.K., Osbon, C.C., 1922. The occurrence and uses of peat in the United States. Ohio State
5 University, Washington, pp.207.
- 6 Vassiljev, A., Blinova, I., 2012. The influence of drained peat soils on diffuse nitrogen pollution
7 of surface water. *Hydrology research* 43, 352-358.
- 8 Venterink, H.O., Davidsson, T.E., Kiehl, K., Leonardson, L., 2002. Impact of drying and re-
9 wetting on N, P and K dynamics in a wetland soil. *Plant and Soil*, 243, 119-130.
- 10 Verhoeven, J.T.A., Kooijman, A.M., Wirdum, G.V., 1988. Mineralization of N and P along a
11 Trophic Gradient in a Freshwater Mire. *Biogeochemistry* 6, 31-43.
- 12 Wang, H., Richardson, C.J., Ho, M., 2015. Dual controls on carbon loss during drought in
13 peatlands. *Nature Climate Change* 5, 584-587.
- 14 Wray, H.E., Bayley, S.E., 2008. Nitrogen dynamics in floating and non-floating peatlands in the
15 Western Boreal Plain. *Canadian Journal of Soil Science* 88, 697-708.