SHORT COMMUNICATIONS

Journal of Arid Environments

Geographic variation in the thermal biology of a widespread Sonoran Desert arachnid,

***Centruroides sculpturatus* (Arachnida: Scorpiones)**

Michael M. Webber¹ and Robert W. Bryson Jr^{2*}

¹School of Life Sciences, University of Nevada, Las Vegas,
4505 S. Maryland Parkway, Las Vegas, Nevada 89154-4004 USA

²Department of Biology & Burke Museum of Natural History and Culture, University of
Washington, Box 351800, Seattle, WA 98195-1800, USA.

*Corresponding author: brysonjr@uw.edu

Telephone: 206-616-5564, Fax: 206-685-3039

Running head: Variation in thermal biology of *Centruroides sculpturatus*

Keywords: Thermal preference, geographic variation, scorpions

Abstract

Environmental temperatures can significantly influence the behavior and physiology of terrestrial ectotherms. Small-bodied terrestrial ectotherms can moderate their body temperatures behaviorally via thermoregulation; however, favorable thermal refuges may be limited across heterogeneous landscapes. In such cases, differences in the thermal environment may generate variation in preferred body temperatures among disparate populations. We tested whether geographic variation in preferred body temperatures existed for the Arizona bark scorpion *Centruroides sculpturatus*, an arachnid widely distributed across the Sonoran Desert. We predicted that geographic variation in thermal preference would exist between populations from a xeric, low-elevation site in western Arizona (Quartzsite) and a cooler, high-elevation site in eastern Arizona (Pinaleño Mountains). We found that scorpions from the Pinaleño Mountains were smaller in body size and exhibited significantly warmer diurnal body temperatures compared to scorpions from Quartzsite. However, no significant difference was detected in the preferred nocturnal temperatures of scorpions from either locality. Scorpions from the Pinaleño Mountains may have shifted their thermoregulatory patterns and selected warmer temperatures during the day as a way to maximize their exposure to favorable temperatures. Less time with mean temperatures within the preferred range of body temperatures may have also lowered metabolic rates of *C. sculpturatus* in the Pinaleño Mountains, resulting in their smaller female body sizes. The ability of *C. sculpturatus* to adapt to different thermal regimes may reflect variation in a life history trait which has allowed them to thrive and expand their distribution across the American Southwest.

Introduction

For terrestrial ectotherms, body temperatures can strongly influence physiological functions and life history traits (e.g., metabolic rate, locomotor performance, digestive rate, and adult body size), and the environmental conditions of habitats selected by these organisms can directly influence their fitness (Blouin-Demers & Weatherhead, 2001; Lourdais *et al.* 2004; Ji *et al.* 2007; Wapstra *et al.* 2010). To maintain favorable body temperatures, small-bodied terrestrial ectotherms thermoregulate by actively seeking out microhabitats with favorable thermal profiles (Halliday & Adler, 2002). Geographic variation can exist in the availability of favorable thermal refuges (Sears, 2005), particularly for species with widespread distributions across heterogeneous landscapes characterized by latitudinal and elevational variation in climate. Because terrestrial ectotherms primarily alter their body temperatures through behavioral means (Bartholomew, 1964), differences in the ambient temperature and availability of thermal refugia across landscapes may lead to intraspecific variation in preferred body temperature.

In this study, we investigated the preferred body temperatures of two populations of Arizona bark scorpion (*Centruroides sculpturatus*). *Centruroides sculpturatus* inhabit rocky terrain and riparian habitats across the Sonoran Desert from Sonora, Mexico north to the western edge of the Mohave Desert in Nevada and Utah, USA (Gantenbien *et al.* 2001; Valdez-Cruz *et al.* 2004; Rowe & Rowe 2008). In Arizona, these scorpions are particularly abundant across an array of habitats ranging from low-elevation xeric desert in the western portion of the state to cooler, high-elevation forested uplands in the east (Spence & Corman, 2005). The topographic complexity across Arizona has created a heterogeneous thermal environment for species that inhabit this region (Lowe 1961). We therefore predicted that we would observe variation in thermal preferences between populations of *C. sculpturatus* from low and high-elevation sites on

opposite ends of the state. More specifically, we hypothesized that populations of *C. sculpturatus* from the warmer, xeric low-elevation regions of western Arizona would exhibit significantly different body temperatures compared to *C. sculpturatus* populations from cooler, higher elevation regions in eastern Arizona.

Materials & Methods

We collected a total of 55 mature gravid female *C. sculpturatus* from two sites in Arizona using ultraviolet-light detection. Twenty-eight mature gravid females were collected in 2011 and 2012 near the outskirts of Quartzite in La Paz County in southwestern Arizona. The town of Quartzite sits at 266 m and is characterized by typical Sonoran Desert vegetation, including saguaro (*Carnegiea gigantea*), creosote (*Larrea tridentata*), cholla (*Cylindropuntia* spp.), and ocotillo (*Fouquieria splendens*). *Centruroides sculpturatus* were collected from under fallen vegetation and along rocky outcrops among boulder-strewn hills. On 13 April 2012, we obtained 27 mature gravid female *C. sculpturatus* from the Pinaleno Mountains in Graham County in southeastern Arizona. The Pinaleno Mountains range in elevation from 1,150 m at the northern base of the range to 3,267 m at the summit (McLaughlin 1993). Five ecological communities span the range, including semi-desert grassland at the lowest elevations, Madrean pine-oak woodland at mid-elevations, and subalpine mixed-conifer forest at the highest elevations (Coronado Planning Partnership 2008). Within the Pinaleno Mountains, *C. sculpturatus* were collected off of fractured vertical rock cuts at 1,854 m in lower Madrean pine-oak woodland near the transition to juniper-oak woodlands.

After collection, we brought all specimens to the laboratory at the University of Nevada, Las Vegas where they were individually weighed and then placed in separate plastic containers

(15.0 x 9.0 cm) with gravel substrates and maintained at temperatures of 19–29 °C. To assess thermoregulatory behavior, we recorded body temperatures of each individual within a novel thermal gradient designed for small ectotherms (Webber & Bryson, 2012). The gradient consisted of a rectangular glass enclosure 71.0 cm (length) x 15.2 cm (width) x 15.2 cm (height) with a gravel substrate. We created a linear gradient ranging from 23 to 50 °C that increased approximately 3.5 °C every 10.2 cm. These temperatures span the range of preferred body temperatures in a variety of scorpions (Warburg and Polis 1990), including the striped bark scorpion *Centruroides vittatus* (36–38 °C; Carlson and Rowe 2009), as well as upper lethal temperatures (45–47 °C; Hadley 1990). We fed all scorpions and allowed them to acclimate for 7 days before placing them in the thermal gradient. Using forceps, we placed each scorpion at the midpoint of each enclosure and allowed them to acclimate 12 hours prior to data collection. Following the acclimation period, we checked mesosomal (abdomen) body temperature (T_b) of each individual every two hours during a 24-hour period using a non-contact infrared thermometer (Model 42505, Extech Instruments, Nashua, New Hampshire, USA). We placed all scorpions on a 13 (ambient) L: 11 D cycle, which corresponded to natural summer conditions at the time of capture. The mean humidity of the room containing the gradient was 40%. We cleaned all enclosures with soap and water and replaced the gravel between trials. We calculated the mean T_b for each female using all temperature measurements taken during the 24-hour trial period (Hertz et al, 1993; Blouin-Demers and Weatherhead, 2001). To compare the preferred T_b of female *C. sculpturatus* from Quartzsite and the Pinaleno Mountains during the day (when scorpions exhibit reduced activity) and at night (when scorpions are primarily active), we divided the 24-hr trial period into two components: diurnal (0800–1800 hrs) and nocturnal (2000–0600 hrs). We calculated the mean diurnal and mean nocturnal T_b for females from each locality, and

compared these values using a repeated-measures analysis of covariance (repeated-measures ANCOVA). This method allowed us to compare the T_b of female scorpions, while controlling for differences in body size (carapace length, mm) among females.

Results

There was a significant interaction between locality and body size (carapace length, mm), in which females from Quartzsite ($\bar{X} = 5.75 \text{ mm} \pm 0.29 \text{ mm}$, $n = 28$) were larger than females from the Pinaleno Mountains ($\bar{X} = 5.43 \text{ mm} \pm 0.22 \text{ mm}$, $n = 27$; ANCOVA, $F_{(1,51)} = 5.98$, $P = 0.02$; Table 1). The mean diurnal T_b of female *C. sculpturatus* from Quartzsite, Arizona ($n = 28$), was $42.5^\circ\text{C} \pm 4.74^\circ\text{C}$, and the mean diurnal T_b of female *C. sculpturatus* from the Pinaleno Mountains ($n = 27$) was $43.2^\circ\text{C} \pm 2.15^\circ\text{C}$. After controlling for differences in body size among females, *C. sculpturatus* from the Pinaleno Mountains exhibited significantly higher diurnal T_b than females from Quartzsite (repeated-measures ANCOVA, $F_{(1,51)} = 5.59$, $P = 0.02$; Table 2). Conversely, there was no significant difference in the selected nocturnal T_b of scorpions from Quartzsite ($\bar{X} = 40.2^\circ\text{C} \pm 3.22^\circ\text{C}$; $n = 28$) and scorpions from the Pinaleno Mountains ($\bar{X} = 41.3^\circ\text{C} \pm 4.18^\circ\text{C}$; ANCOVA, $F_{(1,51)} = 2.53$, $P = 0.12$; Table 2). Additionally, there were no significant diel shifts in the preferred diurnal and nocturnal T_b of female *C. sculpturatus* from Quartzsite or the Pinaleno Mountains (Wilk's Lambda = $\lambda = 0.99$, $F_{(1,51)} = 0.33$, $P = 0.57$). Mean preferred T_b was $4.5\text{--}7.2^\circ\text{C}$ $36\text{--}38$ higher than reported for *C. vittatus* (Carlson and Rowe 2009).

Discussion

Environmental temperatures can significantly influence the behavior and physiology of terrestrial ectotherms (Blouin-Demers & Weatherhead, 2001; Lourdais *et al.* 2004; Ji *et al.* 2007;

Wapstra *et al.* 2010). However, these organisms can exert control over their body temperatures through alterations in thermoregulatory behaviors. The thermoregulatory patterns of ectotherms can be greatly influenced by the availability of suitable thermal refuges (Blouin-Demers & Weatherhead, 2002), and heterogeneity among available thermal refuges across a landscape can lead to localized variation in the temperatures selected by organisms (Freidenburg & Skelly 2004), particularly among populations of widely distributed species.

We found differences in thermal preference between populations of *C. sculpturatus* from low and high-elevation sites. We additionally found differences in body size between these two populations. These differences are likely related to variation in the climate between the two localities. For example, the average high and low air temperature in May is 28 °C and 12 °C at the base of the Pinaleno Mountains (Fort Grant, 1493 m elevation), compared to 36 °C and 18 °C at Quartzite (Cooperative Climatological Data Summaries, Western Regional Climate Center). Although temperatures of exposed rock inhabited by *C. sculpturatus* at both sites can exceed 40 °C (pers. obs.), the Pinaleno Mountains exhibit much cooler temperatures at night than at Quartzsite. This temperature difference translates to less time with mean temperatures within the preferred range of body temperatures for *C. sculpturatus* in the Pinaleno Mountains as compared to the much warmer Quartzsite locality. Less time with mean temperatures within the preferred range of body temperatures may lead to lower metabolic rates of *C. sculpturatus* in the Pinaleno Mountains, resulting in their smaller female body sizes. Although speculative, this inference can be tested in future studies by comparing female body sizes and metabolic rates from high and low-elevation sites across the distribution of this species.

Centruroides sculpturatus from the Pinaleno Mountains selected significantly higher diurnal body temperatures compared to females from Quartzsite. At night, however, female *C.*

sculpturatus from both Quartzsite and the Pinaleno Mountains selected similar body temperatures. The higher diurnal body temperatures selected by *C. sculpturatus* from the Pinaleno Mountains likely enables them to survive in a thermally challenging, high-elevation environment. The Pinaleno Mountains collecting site is higher in elevation than Quartzsite and the temperatures after sunset are typically much colder and drop quicker than at Quartzsite. In the laboratory, female *C. sculpturatus* from the Pinaleno Mountains may have shifted their thermoregulatory patterns and selected warmer temperatures during the day as a way to maximize their exposure to favorable temperatures. In the wild, this might translate to a nocturnal behavior in which high-elevation scorpions “overshoot” thermal optima early in the night that then allows them to sustain nocturnal body temperatures within the preferred range of body temperature of 41.3– 43.2 °C as the nighttime temperature quickly drops.

The thermoregulatory behaviors of ectotherms can influence a suite of life history variables. Examining intraspecific geographic variation in these patterns may therefore improve our understanding of mechanisms that contribute to the diversity in life history characteristics within widely distributed species. We discovered intraspecific variation in the thermal preference of *C. sculpturatus*, a widely distributed Sonoran Desert arachnid. The ability of *C. sculpturatus* to survive in arid lower desert and in cool high-elevation mountains appears linked to different thermoregulatory behaviors, and variability in this life history trait has likely allowed this species to expand and maintain its wide distribution across the American Southwest.

Acknowledgements

We would like to thank G. Webber Jr., M. Irick, R. Rivera, and D. Hartman for assistance in the field, and C. J. Battey and two anonymous reviewers for reviewing earlier drafts of the

manuscript. This work was partially supported by a Doctoral Dissertation Improvement Grant (IOS1209779) from the U.S. National Science Foundation, and by grants from the Graduate and Professional Student Association of the University of Nevada, Las Vegas, to M. M. Webber.

References

- Bartholomew, G.A. (1964) The roles of physiology and behaviour in the maintenance of homeostasis in the desert environment. *Symposia of the Society for Experimental Biology*, **18**: 7–29.
- Blouin-Demers, G. & Weatherhead, P.J. (2001) Thermal ecology of black rat snakes (*Elaphe obsoleta*) in a thermally challenging environment. *Ecology*, **82**, 3025–3045.
- Blouin-Demers, G. & Weatherhead, P.J. (2002). Habitat-specific behavioural thermoregulation by black rat snakes (*Elaphe obsoleta obsoleta*). *Oikos*, **97**: 59–68.
- Carlson, B.E. & Rowe, M.P. (2009) Temperature and desiccation effects on the antipredator behavior of *Centruroides vittatus* (Scorpiones: Buthidae). *Journal of Arachnology*, **37**, 321–330.
- Cooperative Climatological Data Summaries, Western Regional Climate Center (2014) Web. 11 November 2014. <http://www.wrcc.dri.edu/summary/Climsmaz.html>
- Coronado Planning Partnership. 2008. State of the Coronado National Forest: An Assessment and Recommendations for the 21st Century. Sky Island Alliance, Tucson, AZ.
- Ewing, H. (1928) The scorpions of the western part of the United States. *Proceedings of the United States National Museum*, **73**, 27–30.
- Freidenburg, K.L. & Skelly, D.K. (2004) Microgeographical variation in thermal preference by an amphibian. *Ecology Letters*, **7**: 369–373.

- Halliday, T., & Adler, K. (2002) Firefly encyclopedia of reptiles and amphibians. Toronto: Firefly Books.
- Hertz, P.E., Huey, R.B. & Stevenson, R.D. (1993) Evaluating temperature regulation by field-active ectotherms: the fallacy of the inappropriate question. *American Naturalist*, **142**: 796–818.
- Ji, X., Lin, C.X., Lin, L.H., Qiu, Q.B. & Du, Y. (2007) Evolution of viviparity in warm-climate lizards: an experimental test of the maternal manipulation hypothesis. *Journal of Evolutionary Biology*, **20**, 1037–1045.
- Lourdais, O., Shine, R., Bonnet, X., Guillon, M. & Naulleau, G. (2004) Climate affects embryonic development in a viviparous snake, *Vipera aspis*. *Oikos*, **104**, 551–560.
- Lowe, C.H. (1961) Biotic communities in the sub-Mogollon region of the inland Southwest. *Journal Arizona Academy Science*, **42**:40–49.
- McLaughlin, S.P. (1993) Additions to the flora of the Pinaleno Mountains, Arizona. *Journal of the Arizona-Nevada Academy of Science*, **27**: 5–32.
- Rowe, A.H. & Rowe, M.P. (2008) Physiological resistance of grasshopper mice *Onychomys* spp. to Arizona bark scorpion *Centruroides exilicauda* venom. *Toxicon*, **52**, 597–605.
- Russell, F.E. & Madon, M.B. (1984) Introduction of the scorpion *Centruroides exilicauda* into California and its public health significance. *Toxicon*, **22**: 658–664.
- Sears, M.W. (2005) Geographic variation in the life history of the sagebrush lizard: the role of thermal constraints on activity. *Oecologia*, **143**: 25–36.
- Spence, J.R. & Corman, T.E. (2005) *Geography, Climate and Habitats of Arizona* in The Arizona Breeding Bird Atlas. (T.E. & Wise-Gervais, C., eds.) UNM Press.

- Wapstra, E., Uller, T., While, G.M., Olsson, M. & Shine, R. (2010) Giving offspring a head start in life: field and experimental evidence for selection on maternal basking behaviour in lizards. *Journal of Evolutionary Biology*, **23**, 651–657.
- Warburg, M.R. & Polis, G.A. (1990) Behavioral responses, rhythms, and activity patterns. Pp. 112–122. In *The Biology of the Scorpions* (G.A. Polis, ed.). Stanford University Press, Stanford, California.
- Hadley, N.F. (1990). Environmental physiology. Pp. 158–1666. In *The Biology of the Scorpions* (G.A. Polis, ed.). Stanford University Press, Stanford, California.
- Webber, M.M. & Bryson Jr., R.W. (2012) A novel thermal gradient design for small-bodied ectotherms. *Euscorpius*, **140**, 1–6.

Table 1. Comparison of body size (carapace length, mm) between female *Centruroides sculpturatus* from a low-elevation site (Quartzite, n = 28) and high-elevation site (Pinaleño Mountains, n = 27) in Arizona.

Covariate	<i>df</i>	MS	F	<i>P</i>
Locality	1	100.3	5.12	0.02
Body size	1	0.13	0.008	0.93
Locality * Body size	1	103.0	5.98	0.02
Error	51	17.2	—	—

Table 2. Comparison of diurnal and nocturnal body temperatures (°C) between female *Centruroides sculpturatus* from a low-elevation site (Quartzite, n = 28) and high-elevation site (Pinaleño Mountains, n = 27) in Arizona.

Covariate	<i>df</i>	MS	F	<i>P</i>
Diurnal Body Temperature				
Locality	1	68.7	5.59	0.02
Body size	1	10.3	0.84	0.36
Locality * Body size	1	68.1	5.54	0.02
Error	51	12.3	—	—
Nocturnal Body Temperature				
Locality	1	34.5	2.53	0.12
Body size	1	7.30	0.53	0.47
Locality * Body size	1	37.2	2.73	0.11
Error	51	13.7	—	—