**Soils mediate the impact of fine woody debris on invasive and native grasses as
whole trees are mechanically shredded into firebreaks in piñon-juniper woodlands**

Zachary T. Aanderud^{1*}, Donald R. Schoolmaster², Deborah Rigby¹, Jordon Bybee¹, Tayte
Campbell¹, and Bruce Roundy¹

1
2
3
4 24 Keywords: *Bromus tectorum*, mastication, microbial biomass, phosphorus,
5
6 *Pseudoroegneria spicata*, shredding
7
8

9 26

10
11 *Corresponding author: Zachary T. Aanderud
12
13

14 28 Department of Plant and Wildlife Sciences
15

16 4105 LSB, Provo, UT 84602
17

18
19 30 Phone: (801) 422-4220, email: zachary_aanderud@byu.edu
20
21
22

23
24 32 Author contributions: ZTA, DR, JB, and BR designed the study. ZTA, DR, JB, and TC
25
26 conducted the experiments. ZTA, DRS, TC, and BR analyzed and interpreted the data,
27

28
29 34 and all authors helped write and review the manuscript. ZTA agrees to be accountable for
30
31 all aspects of the work in ensuring that questions related to the accuracy or integrity of
32

33
34 36 any part of the work are appropriately investigated and resolved.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

ABSTRACT To stem wildfires, trees are being mechanically shredded into firebreaks with the resulting fine woody debris (FWD) potentially exerting immense control over soil and plants. We linked FWD-induced changes in microbial activity and nutrient availability to the frequency of *Bromus tectorum* and three native, perennial grasses across 31 piñon-juniper woodlands, UT, USA. Using a series of mixed models, we found that FWD increased the frequency of three of the four grasses by at least 12%. Deep, as opposed to shallow, soils mediated frequencies following FWD additions but only partially explained the variation in *Bromus* and *Pseudoroegneria spicata*. Although fertile areas associated with tree-islands elicited no response, FWD-induced increases in nitrogen mineralization in deep soils (15-17 cm) caused the frequency of the exotic and *Pseudoroegneria* to rise. Higher phosphorus availability in FWD-covered surface soils (0-2 cm) had no impact on grasses. FWD altered deep soil respiration, and deep and shallow microbial biomass structuring *Pseudoroegneria* frequencies, suggesting that microorganism themselves regulated *Pseudoroegneria*. The positive effects of FWD on grass frequencies intensified over time for natives but diminished for *Bromus*. Our results demonstrate that microorganisms in deeper soils helped mediate species-specific responses to disturbance both facilitating exotic invasion and promoting native establishment.

1. Introduction

Terrestrial environments are exposed to a diverse array of disturbances that alter ecosystem characteristics and induce microbial-mediated biological changes belowground. Disturbances that alter the quantity and quality of carbon (C) resources entering environments, in particular, have the potential to exert immense control over soil processes (Zechmeister-Boltenstern et al., 2015). Wildfires, tree-fall, typhoons and/or anthropogenic activities related to timber harvesting (Tinker and Knight, 2000; Murphy et al., 2008) result in the addition of dead and downed wood that promotes soil C storage (Klopatek, 2002), stimulates and depresses the fluxes of trace gases to the atmosphere (Hafner and Groffman, 2005), and releases new sources of energy for trophic web interactions (Norden et al., 2004). Further, many of the effects of woody debris on soils may extend aboveground.

Linking potential shifts in microbial activity to plant performance is extremely difficult, but clearer relationships may arise following disturbances. Disturbances often lead to dramatic shifts in aboveground plant community composition, especially when promoting the invasion and dominance of exotic species (MacDougall and Turkington, 2005). Following woody additions, the cover of exotic understory species increases in many forested ecosystems (e.g., tropical and coniferous forests; Kerns et al., 2006; Murphy et al., 2008) in part due to sunlight reaching the once-shaded understory as downed wood opens gaps in the overstory canopy. Woody additions may also alter the belowground environment and impact plants. In surface soils, woody additions generally lead to the immobilization of inorganic nitrogen (N) as microorganisms decompose copious recalcitrant wood substrates (Laiho and Prescott, 2004). However, in the absence

1
2
3
4 78 of an overstory utilizing N, N may accumulate and be exploited by germinating and
5
6 emerging seeds, and established understory plants. Deeper in soils, the resulting dead root
7
8
9 80 system from canopy species may decompose and stimulate N mineralization and
10
11 availability (Ehrenfeld et al., 1997) aiding in seedling recruitment. Besides N dynamics,
12
13
14 82 dead and downed wood may alter other aspects of the soil environment favoring plant
15
16 invasion. For example, woody debris may cause microorganisms to: 1) elevate soil water
17
18
19 84 availability as decomposition and the production of extracellular polymeric substances
20
21 modify surface debris and/or soil structure (Or et al., 2007; Young et al., 2013); 2)
22
23
24 86 increase the release rates of essential plant macronutrients, such as phosphorus (P;
25
26 Richardson and Simpson, 2011); and 3) change in abundance with species specializing in
27
28
29 88 degrading woody substrates becoming dominant, and growth promoting rhizobacteria
30
31 (Hayat et al., 2010) and mycorrhizae (Vandenkoornhuyse et al., 2003) becoming rare.
32
33
34 90 However, the extent that these disturbance-induced shifts extend deeper into soils away
35
36 from woody materials or translate into plant responses remains relatively unclear.
37
38
39 92 The action of shredding semi-arid trees, coupled with one of the most prolific plant
40
41 invasions occurring in any ecosystem, offers an unprecedented opportunity to understand
42
43
44 94 the links between fine woody debris (FWD)-induced changes belowground and native
45
46 and exotic plant responses aboveground. The annual exotic grass, *Bromus tectorum* (L.),
47
48
49 96 is a prolific invader that degrades piñon-juniper woodlands by replacing native grass
50
51 species and creating monocultures of continuous fuels leading to catastrophic wildfires
52
53
54 98 (Brooks et al., 2004). To stem wildfires and create firebreaks in piñon-juniper woodlands
55
56 (Owen et al., 2009), thousands of hectares of trees are being mechanically shredded into
57
58 100 FWD and translocated to the soil surface. Most FWD is being deposited in place on
59
60
61
62
63
64
65

existing “tree-islands of fertility” in contrast to interspaces containing grasses, forbs, and biological soil crusts. The mosaics of islands of fertility and interspaces are chiefly responsible for differences in microbial activity prior to FWD additions (Aanderud et al., 2008) and may serve to augment the effects of FWD on belowground processes. Soils beneath trees are enriched with C and other essential elements (e.g., N, P, and calcium) due to roots acquiring nutrients from interspace soils and depositing them on the soil surface via litter inputs. Even under this enrichment, microbial activity in tree-islands is low due to the low quality of pine needle litter, and this limitation may be enhanced as more recalcitrant C sources from FWD are added to surfaces. Also, N transformations can be higher beneath tree-islands than interspace soils (Schade and Hobbie, 2005), and the addition of woody C sources, such as lignin, may only serve to immobilize N and reduce N release (Bates et al., 2002).

In this study, we evaluated the extent that soils mediated the responses of exotic and native grass species to FWD. In a replicated field experiment, where FWD was manipulated across 31 piñon-juniper woodlands, we linked FWD-induced changes in microbial activity and nutrient availability to shifts in the frequency of the invasive *B. tectorum* and three native, perennial grasses in the Great Basin Desert and Colorado Plateau, UT, USA. The three perennial species included: *Elymus elymoides* (Raf.) Swezey, *Poa secunda* (J. Presl), and *Pseudoroegneria spicata* (Pursh) Á. Löve. To understand the ecological consequences of a single disturbance type occurring at different times across 30+ ecosystems, we used a series of mixed models. Specifically, we developed models to: 1) describe the effects of shredding whole *Juniperus osteosperma* (Torr.) and the subsequent addition of FWD on N mineralization, P availability,

124 microbial respiration rates, microbial biomass, and dissolved organic C (DOC) in surface
 and subsurface soils; and, 2) relate these changes to grass species distribution. We also
 126 examined these potential links across tree-island soils covered in FWD and relatively
 bare interspace soils. The perennial grass species we selected are often influenced by *B.*
 128 *tectorum* invasions (Mangla et al. 2011) and represent dominant grasses in piñon-juniper
 woodlands. Thus, by examining these natives in tandem with *B. tectorum*, our results
 130 may offer a more comprehensive understanding into the consequences of shredding on
 species invasion. We hypothesized that: 1) the effects of FWD on shallow soils,
 132 immediately below the disturbance in particular, will explain some of the variation in the
 frequency of all grasses; 2) tree-island microsites will influence FWD-induced links
 134 between soils and grasses; and 3) changes in the availability of essential plant nutrients
 (i.e., N and P) and not microbial activity will be the prominent processes dictating grass
 136 response following FWD additions.

138 **2. Materials and methods**

2.1 Woodland sites

140 We conducted our study across 31 piñon-juniper woodland sites in the Great Basin
 Desert, Utah, USA. Dominant tree species included *Juniperus osteosperma*, and *Pinus*
 142 *edulis* (Engelm.) along with the shrubs *Artemisia tridentata* Nutt. (ssp. *wyomingensis*
 Beetle and Young, *tridentata*, and *vaseyana* [Rydb.] Beetle). We selected woodland sites
 144 with at least 15-45% tree cover estimated using images from the National Agricultural
 Imagery Program at 1 m spatial resolution (USDA-FSA-APFO Aerial Photography Field
 146 Office, Salt Lake City, UT). To ensure that the sites were indicative of semi-arid soil in

the western USA, we measured the following physiochemical characteristics: pH from soil pastes (Thermo Orion pH meter model 410, Thermo Scientific Orion, Beverly, MA); electrical conductivity from 1:1 w/v extracts (Beckman Conductivity Bridge model RC-16C, Beckman Coulter Inc., Brea, CA); soil %C and %N with dry combustion on a C to N analyzer (LECO TruSpec CN Determinator, LECO Corporation, St. Joseph, MI); and soil texture (% sand, silt, and clay) with a hydrometer. Basic soils characteristics were based on three soil samples from a combination of at least one surface and subsurface soil sample to capture the overall environment at each site. The mean annual temperature and precipitation in woodlands was based on a thirty-year period (1982-2012) from PRISM climate datasets (<http://www.prism.oregonstate.edu>, PRISM Climate Group, 2004).

2.2 FWD manipulation and soil sampling

To investigate the effects of FWD on soils, we created three, randomly-selected FWD addition and three control plots (33 × 30 m) in each site. In the addition plots, whole *J. osteosperma* trees were shredded with a toothed drum (Cline et al., 2010), and the resulting woody material was deposited on the soil surface 1-8 years prior to sampling (2003-2010), which occurred over a 2-year period during the summer months (April-August, 2011-2012). Our FWD fits best within the definition of fine down and deadwood employed by researchers in multiple other ecosystems (Norden et al., 2004; Muller-Using and Bartsch, 2009). Following whole-tree shredding, the dead and downed materials fall under the classification of FWD (diameter < 10 cm) and do contain classic FWD components (e.g., needles, twigs, small branches). These materials also consist of extremely woody components (e.g., logs and large branches) generally reserved for FWD

(diameter > 10 cm; Muller-Using and Bartsch, 2009) that are only reduced in size but not different in chemistry. Specifically, the diameter of FWD ranged from 0.1-7.5 cm (Young et al., 2014) and FWD was piled on top of tree-islands.

To begin to understand the effects of FWD on links between soils and grasses, we sampled soils in *J. osteosperma* tree-island microsites at the surface (0-2 cm depth) where seeds may germinate and emerge, and subsurface (15-17 cm depth) where establishing root systems may interact with nutrients and microorganisms. Roots of many dominant grasses in piñon-juniper woodlands, including our four study species, span these depths and are known to actively acquire resources from shallow and deep soils (James et al., 2008). The three microsite treatments included soils from interspaces (interspace), at the edge of the tree canopy or edge of woody addition piles (edge), and underneath the tree canopy or beneath the woody debris (canopy) approximately one-third the distance from the trunk of the tree to the canopy edge. In the FWD addition plots, microsites were identified using the shredded trunk of the shredded tree. We removed the soils from the three FWD addition and control plots with a corer (5 cm diameter $\times$ 10 cm length) and combined the three resulting subsamples from each treatment to create a composite sample for all microsite and depth combinations (2 treatments, FWD and control $\times$ 3 microsites $\times$ 2 depths $\times$ 31 study sites = 372 samples). The two depths were taken from different cores and were independent of each other. Soils were passed through a 2 mm sieve to remove rock fragments, stored at 4°C, and used in soil laboratory incubations within 5 days of field sampling. To standardize sampling, we evaluated woody debris piles and standing trees with canopies approximately 3 m in diameter. Last, the FWD

1
2
3
4 192 treatment and control plots were located as close to each other as possible within each
5
6 site, with the maximum distance between plots no more than 2 km.
7
8

9 194
10
11 2.3 *Grass species frequency*
12
13

14 196 To identify grass species responses to FWD, we evaluated the frequencies of exotic *B.*
15
16 *tectorum*, and three native perennial grasses *E. elymoides*, *P. secunda*, and *P. spicata* in
17
18 198 tree-island microsites following debris additions. Specifically, we measured the
19
20 occurrence of each grass across five randomly selected 30-m transects across each of the
21
22 three FWD addition and control plots using the line-point method described by Roundy et
23
24 200 al. (Roundy et al., 2014b; 2 treatments, FWD addition and control $\times$ 3 plots $\times$ 5 transects
25
26 $\times$ 31 study sites = 930 transects). Using Python version 2.7.3 (Python Software
27
28 Foundation, DE, <http://www.python.org>), first, we classified the 60 points along each
29
30 204 transect into microsites (i.e., interspace, canopy edge, and tree canopy) based on the
31
32 presence of tree canopies and woody debris. Second, we calculated the percent frequency
33
34 206 of species as the number of times a species occurred in a given microsite divided by the
35
36 total number of times that microsite occurred along each transect $\times$ 100. Third, we
37
38 208 calculated the mean species frequency present for each microsite from the five transects
39
40 in a given plot, and the mean species frequency in the three plots for the FWD and
41
42 control treatments for each woodland.
43
44
45
46
47
48 210
49
50
51
52

53 212 2.4 *Net N mineralization and P availability*
54
55

56 We measured N transformation rates and estimated soil P availability to identify the
57
58 214 influence of FWD on soil nutrients. In all soils, net N mineralization rates (mg N-NH_4^+
59
60
61
62
63
64
65

kg soil⁻¹ day⁻¹) were measured as the change in NH₄⁺ and NO₃⁻ in laboratory microcosms of surface and subsurface soils (Schaeffer et al., 2007). Microcosms consisted of 30 g of soil that was weighed into a specimen cup (155 mL), brought to a constant gravimetric water content (GWC) to simulate moist but not waterlogged conditions that semi-arid microorganisms may experience (0.3 g H₂O g dry soil⁻¹), and placed in a quart-sized Mason jar (volume = 935 mL). Jars had a gas-tight septum in the lid and contained 10 mL of water to keep samples from drying out during the incubation. Microcosms were incubated over 10 days at 30°C. We measured NH₄⁺ and NO₃⁻ at time zero and after 10 days in 0.5 M K₂SO₄ extractions (1:2 w/v) colorimetrically (SpectraMax Plus 384, Molecular Devices Corporation, Sunnyside, CA). We used the methods outlined by James et al. (James et al., 2008) for NH₄⁺ and NO₃⁻. An estimate of soil P (mg P kg soil⁻¹) availability was evaluated at time zero by extracting soils with 0.5 M NaHCO₃ (1:20 w/v) following the Olsen Sodium Bicarbonate method (Olsen et al., 1954) and measured colorimetrically.

2.5 Microbial biomass, soil respiration, and dissolved organic C

To evaluate the effects of FWD on microbial activity and potential links to grass frequencies, we evaluated microbial biomass, soil respiration, and DOC. Active soil microbial biomass (mg C kg soil⁻¹) was evaluated using a substrate-induced respiration (SIR) technique. Briefly, 10 g of soil was weighed into a specimen cup with a gas-tight septum in the lid (155 mL) and 1.1 mL of a 4.0 M sucrose solution was added to provide 4 mg sucrose g soil⁻¹ and bring the soils to a constant GWC (0.3 g H₂O g dry soil⁻¹). We measured the difference between the initial and maximum respiration (24 hours after

sucrose addition) and calculated biomass using the equation: $SMB \text{ (mg C kg soil}^{-1}\text{)} = (40.04 \times \Delta CO_2) + 0.37$ (Bailey et al., 2002), where ΔCO_2 is the change in respiration rates in units of $\mu\text{L CO}_2 \text{ g}^{-1} \text{ h}^{-1}$ caused by the addition of the sucrose solution. We also measure soil respiration rates ($\text{mg CO}_2 \text{ kg soil}^{-1} \text{ day}^{-1}$) during the first 24 hours in the same microcosms used to measure N mineralization. We extracted 10 mL of air from the microcosms with a 10 mL gas-tight syringe; measured CO_2 concentration (ppm) using an EGM-4 IRGA (PP Systems, Amesbury, MA), and converted CO_2 concentrations to mg, using the Ideal Gas Law. For DOC (mg C kg soil^{-1}), soils were extracted via a distilled water extraction (1:2 w/v), passed through a $0.45 \mu\text{m}$ nylon filter, and measured on a TOC-N analyzer (Shimadzu, Columbia, MD).

2.6 Model Set 1: FWD and soil characteristics

We evaluated links between FWD-induced changes in microbial activity and nutrient availability to shifts in grass frequency with a series of mixed models to account for site as a random effect (Bolker et al., 2009). First, we created models estimating the effect of FWD additions on five soil characteristics in shallow and deep soils with the *lmer* function of the lme4 package (Bates et al., 2014) in R (R Core Team, 2015). Models were of the form,

$$y_i \sim N\left(\alpha_{j[i]} + \sum_k \beta_k x_{ki}, \sigma_y^2\right) \quad \text{equation 1}$$

and included site location, j , as a random effect, $\alpha_j \sim N(\mu_\alpha, \sigma_\alpha^2)$, and tree-island

microsites as a covariate, x . All soil data used as inputs into the models were based on differences between FWD and control treatments by woodland sites and allowed for the

1
2
3
4 260 inclusion of time since FWD addition as a covariate. While using this differencing
5
6 method, if there was no effect of FWD treatment on a measured response (i.e., $y_{\text{FWD}} =$
7
8 y_{control}), then the response variable $y' = \log_{10}(y_{\text{FWD}}) - \log_{10}(y_{\text{control}}) = 0$. Alternatively,
9 262 values of $y' > 0$ indicated that FWD caused an increase relative to the control, and $y' < 0$
10
11 indicated that the FWD treatment caused a decrease compared to the control. For the
12
13
14 264 model, we estimated the effects of FWD on soils as model intercepts and determined
15
16 significant differences based on 95% confidence intervals. Thus, a significant effect of
17
18 266 the FWD treatment was indicated if the confidence interval of the intercept did not
19
20
21 include zero.
22
23 268

270 2.7 Model Set 2: The effects of FWD on grasses and the potential mediation of soils

271
272 Next, we fit a second set of models, one for each species estimating the total effect of
273
274 FWD additions on the change in grass frequency as a function of all five soil
275
276 characteristics from both depths representing hypothesized explanatory variables. We
277
278 followed logic of tests of mediation (Shrout and Bolger, 2002). Three possible
279
280 meaningful outcomes identifying the effects of FWD on links between species
281
282 frequencies and soil characteristics were possible (Fig. 1). The outcomes included: 1) the
283
284 effects of FWD on species frequencies were entirely mediated through changes in soil
285
286 variables due to woody additions (Fig. 1a); 2) the effects of FWD on species frequencies
287
288 were partially explained by FWD and soil variables independently, suggesting that other
289
290 unmeasured effects of the FWD treatment impacted species frequencies as well (Fig. 1b);
291
292 and 3) the FWD treatment alone explained the variation in grass frequencies (Fig. 1c).
293
294 We estimated the potential for soils to mediate this relationship by comparing the change

in intercepts between grass frequency models including and excluding soil variables. If belowground characteristics explained any of the variation in grass frequencies, then the intercept from the model, including soil variables, should be closer to zero than the intercept from the model excluding soil variables. Further, the 95% confidence intervals should also decrease in size from models that exclude to those that include soil variables. As with the first model, these models were based on FWD and control treatments by woodland sites with time since FWD addition as a covariate difference and site location as a random effect. For more explanation of the logic test, equations, and modeling process see Supplemental Model Explanation (Supplemental Materials).

Fig. 1

2.8 Model Set 3: Links between FWD-induced changes in soils and grasses

Last, we generated a third set of mixed models, similar to the second set, one for each grass species to suggest specific links between FWD-induced changes in individual soil variables and grass frequencies. The resulting links between model parameters can

roughly be interpreted as the percent change in the response variable given a 1% change in the explanatory variable. All models used the same differenced data, covariates, and random effects. The *p*-values for the effects of soil variables on grass frequency were calculated using likelihood ratio tests with the *lmer* function of the lme4 package in R. The magnitude and directionality of links were identified between soils and grasses and all models were combined into one figure to help interpret differences among grass species. For more explanation of the mixed models see Supplemental Model Explanation (Supplemental Materials).

3. Results

3.1 Woodland site characteristics

Soils were sandy to clay loams with neutral pH and low salt concentrations, but varied in total % C and % N. Across the 31 woodland sites, soil texture was dominated by sand (mean and $\pm$ standard error of the mean (SEM), $45\% \pm 2.5$) and silt ($32\% \pm 1.5$; Supplemental Table 1). All woodland soils possessed a neutral pH (7.0 ± 0.08) and low non-saline ECs (0.64 ± 0.06 dS m⁻¹). The % C varied from $0.63 (\pm 0.05)$ to $9.4 (\pm 2.2)$ and total % N varied from $0.08 (\pm 0.01)$ to $0.26 (\pm 0.05)$. Sites experienced mesic temperatures with mean annual temperatures of $9.7^\circ\text{C} \pm 0.27$, and a semi-arid climate with mean annual precipitation of $366 \text{ mm} \pm 10$. The elevation of the sites ranged from 1637-2298 m.a.s.l. (Supplemental Table 1).

3.2 Model Set 1: FWD influenced *P* availability, DOC, and *C* mineralization

The addition of FWD stimulated soil P availability and DOC in canopy surface soils and depressed microbial activity in deep interspace soils (Fig. 2). Woody debris increased the availability of P at least 2.2-fold in canopy compared to interspace and edge soils (95% confidence interval of the intercept did not include zero). The addition of FWD also elevated the concentration of DOC, but only in canopy soils. As for activity, C mineralization was 4.2% and 9.5% lower in deep interspace than edge and canopy soils respectively. For means and $\pm$ SEM from the FWD and control treatments, see Supplemental Table 2.

Fig. 2

3.3. *Model Set 2: Soils mediated the effects of FWD on exotic and native grasses*

The addition of FWD influenced the frequency of *B. tectorum*, and two of the three native perennial grasses, *P. spicata* and *E. elymoides*, but soils only mediated the frequencies of *B. tectorum* and *P. spicata*. Based on the intercepts estimating the experimental effect of FWD additions alone (models including woodland sites as a random effect and time since FWD addition as a covariate, i.e., designated as site in Fig. 3), FWD increased the frequency of the exotic by 14% and the natives by at least 12% (Fig. 3a, 3c, 3d). Alternatively, *P. secunda* was not sensitive to FWD, because its 95% confidence interval overlapped zero (Fig. 3b). Further, due to the reduction in intercept between the two models excluding and including soil variables (i.e., site versus site + soil, Fig. 3), the effects of FWD on soil characteristics reduced or explained 25% and 41% of the variation in *B. tectorum* and *P. spicata* frequencies, respectively (Fig. 3a, 3c). Thus, the exotic and native followed the second outcome from the logic tests with both FWD and the effects of woody additions on soil variables partially mediating their frequencies. Alternatively, the inclusion of the effect of FWD on soils actually increased the uncertainty in predicting the frequency of *E. elymoides*, because the intercept increased following the inclusion of soils (Fig. 3d). Thus, *E. elymoides* followed the third outcome and its frequency was entirely influenced by FWD not mediated by soils. Including time since shredding as a covariate reduced the variation in intercepts and was retained in the models. However, the inclusion of tree-island microsites into the models did not explain any of the variation in grass frequencies or improve the fit of any model and was dropped as a covariate.

Fig. 3

3.4 Model Set 3: Specific links between FWD-induced changes in soils and grasses

FWD-induced changes in microbial activity in deeper soils, (i.e., soil respiration, net N mineralization, and microbial biomass) and P availability in shallow soils, influenced the exotic and two perennial grasses. Although, FWD additions influenced only soil respiration, P availability, and DOC within tree-island microsites (Fig. 2), when microsite was dropped from the model, two new FWD-driven changes were evident (Fig. 4). These links were implied due to the detection of significant effects of N mineralization and

microbial biomass on *B. tectorum* and *P. spicata* responses. Specifically, N mineralization significantly influenced the frequency of *B. tectorum* ($p = 0.02$) and *P. spicata* ($p = 0.01$) following FWD additions. Based on the mixed models, a 1% change in N mineralization induced the frequency of *B. tectorum* and *P. spicata* to increase by 0.36% and 0.23% respectively. Also, microbial biomass in deep soils marginally depressed the frequency of *P. spicata* ($p = 0.07$), while biomass in shallow soils marginally stimulated this same species ($p = 0.06$). In deep soils, FWD reduced soil respiration, and this reduction resulted in a marginal decline in *P. spicata* ($p = 0.07$), indicating a positive relationship or linkage between this soil variable and grass. The availability of P in shallow soils influenced one native, but unlike N mineralization, the increase in P availability following FWD additions actually negatively impacted *P. secunda*. Overall, the frequency of this native grass was insensitive to FWD, but soils with higher P availability reduced *P. secunda* ($p = 0.03$) frequency. Although not statistically significant itself, the intercept for time since shredding was negative for *B. tectorum* ($-0.15, p = 0.17$) and positive for both perennials, *P. spicata* ($0.17, p = 0.20$) and *P. secunda* ($0.17, p = 0.16$), demonstrating significant links to soil processes.

386 **Fig. 4**

388 **4. Discussion**

390 The effects of woody debris on ecosystems are often highlighted in coniferous and deciduous temperate forests where debris may constitute a substantial portion of total C and strongly regulate ecosystem processes (Pregitzer and Euskirchen, 2004). Recently, however, there is an unparalleled addition of woody debris occurring in piñon-juniper woodlands for which there is no ecological precedent. The management practice of shredding piñon-juniper trees has often aided in the regrowth of native grasses in general

(Bates et al., 2007; Rose and Eddleman, 1994), and may also stimulate invasion (Owen et al., 2009). We found that shredding increased the frequency of only three of the four dominant grasses that we studied. The number of years since the disturbance occurred intensified the effects of shredding on the native grasses but diminished the effects on the exotic. In general, links between soils and grass species occurred predominantly in deeper soils (i.e., soil respiration, net N mineralization, and microbial biomass). However, FWD-induced shifts in soils explained up to 41% of species responses of only two species, *B. tectorum* and *P. spicata*. Unexpectedly, fertile areas associated with tree-islands did not structure the frequency of any grass species. Our results demonstrate the importance of plant-soil-microbial interactions in mediating the ecological effects of a disturbance, and highlight that these interactions may occur deeper in the soil profile away from the initial disturbance.

4.1 Grass-specific response to FWD and soils

Contrary to our hypothesis, we found that the effects of FWD on soils did not necessarily influence the frequencies of all four grasses but may simultaneously facilitate exotic invasion and promote native grass establishment. For example, for exotic *B. tectorum* and native *P. spicata*, the contribution of our measured soil nutrient and microbial activity variables explained 25% and 41% of their frequency, respectively. However, *E. elymoides* frequency was not directly related to any of our measured soil variables, and *P. secunda*, although insensitive to woody addition, was influenced by P availability. The remaining variation in *B. tectorum*, *P. spicata*, and *P. secunda* and all the variation in *E. elymoides* frequencies were most likely controlled by other variables, such as external

climate drivers like intra- and inter-annual precipitation variability and ecohydrological shifts caused by FWD and tree removal. Rainfall inputs and soil moisture dynamics often drive grass seedling survival in arid and semi-arid ecosystems (Abbott and Roundy, 2003), with relatively small changes in the timing or the amount of rainfall translating into consequence for plant population dynamics and species coexistence (Peters et al., 2014). Also, FWD and tree removal increases water availability for remaining plant species by decreasing rainfall interception and the amount of water lost due to evapotranspiration fluxes (Roundy et al., 2014a). For example, *J. osteosperma* removal increased the number of wet or relatively moist days (> -1.5 MPa) by at least a month to a soil depth of 30 cm (Young et al., 2013).

4.2 Deep instead of shallow soils influenced grasses

Unexpectedly, our results suggest that the effects of FWD on biology may extend deep into the soil profile, potentially deeper than the effects of woody materials on soil chemistry that extend only a few centimeters (Spears and Lajtha, 2004). We hypothesized that surface and not subsurface soils would dominant FWD-induced links between soil processes and species. Surface soils contain germinating grasses that are in direct contact with shredded materials. Thus, if soils mediated grass responses to FWD, we believed we would see these effects chiefly in surface soils. But two-thirds of the microbial-mediated processes influencing grasses occurred in deeper soils. The importance of deeper soil may have stemmed from FWD constituents leaching through the soil profile, but the decomposition of shredded piñon and juniper root systems most likely aided in altering processes. Root systems in these woodlands constitute 25-40% of the total tree biomass

(Miller et al., 1990) and are composed of mostly fine roots at our deep sampling depth (12-17 cm; Peek et al., 2006). Fine roots are relatively more labile than woody C substrates. Thus, over the last 1-8 years since shredding, the decaying root system, and not FWD, may have influenced our significant, deep soil processes (i.e., net N mineralization, respiration, and microbial biomass) to a greater extent. Unfortunately, we did not experimentally separate FWD from root effects on nutrient availability. Additionally, the reason for the limited links in surface soils may stem from the effects of FWD being masked by the relatively thick layer of litter and duff (partly decomposed organic matter or unrecognizable plant forms) existing between the woody additions and the soil mineral surface. *Juniper osteosperma* trees generate immense amounts of litter and duff beneath their canopies and these materials may continue to influence soil microorganisms more than FWD even following the disturbance.

4.3 Time since disturbance and not microsite structured grass responses

Surprisingly, tree-island microsites failed to improve the fit of any our models. We hypothesized and expected fertile areas (i.e., canopy and edge microsites) to increase grass species frequency following FWD additions. Shredded materials in manipulative studies have increased grass seedling emergence, establishment, and survival by adding more nutrients to already relatively fertile tree-island soils and trapping moisture in the spring (Young et al., 2013). Due to our post-hoc design, we may have missed the importance of microsites in structuring plant responses. We measured all existing grass species and not just the new individuals emerging since shredding. The number of years since shredding consistently reduced the uncertainty surrounding the FWD-induced soil

1
2
3
4 464 effects on grass frequencies, but the way that time influenced grasses differed between
5
6 the annual and perennials. The negative time intercept for *B. tectorum* suggested that the
7
8 effects of shredding were diminishing with time, while a positive time intercept for *P.*
9 466
10 *spicata* and *P. secunda* suggested that the effects of FWD were intensifying. Our results
11
12 support other findings where an increase of native plants, especially perennial plants,
13
14 468 started to occur a couple of years after treatment and continued to increase in frequency
15
16 with each additional year (Rose and Eddleman, 1994; Bates et al., 2007). FWD requires
17
18 470 decades to decompose, but possibly the changes in the annual *B. tectorum* are realized
19
20 relatively quickly (i.e., the first few years) due the species being an opportunistic
21
22 competitor and a prolific seed producer. Alternatively, the impact of FWD on perennial
23
24 472 grasses may require multiple years to become apparent as certain native species benefit or
25
26 contend with FWD-induced changes in soils to survive and produce seed.
27
28
29 474
30
31
32
33
34 476

35 36 4.4 Deep soil N but not P stimulated exotic and native grasses 37

38 478 Consistent with our hypothesis, net N mineralization, instead of soil respiration or
39
40 microbial biomass, dictated the highest number of grass species responses following
41
42 FWD. In deep soils, higher N availability stimulated the frequencies of both the exotic *B.*
43 480
44 *tectorum* and *P. spicata*. In N-limited ecosystems, like piñon-juniper woodlands,
45
46 inorganic N pools, especially in surface soils, are highlighted as an integral factor
47
48 482 influencing grass species success (James et al., 2008). For example, during invasion, *B.*
49
50 *tectorum* reduced soil inorganic N pools in the top 5 cm of soil by as much as 50%,
51
52 indicating the potential importance of N to the success of this exotic (Rimer and Evans,
53 484
54 2006). However, both of these species are known to utilize deeper soil N resources with
55
56
57
58 486
59
60
61
62
63
64
65

1
2
3
4 the native perennial actually acquiring similar levels of inorganic N from shallow and
5
6
7 488 deep N pools (James et al., 2008). Our results suggest that these deeper soil resources
8
9 altered by FWD enhanced the survival and/or growth of the exotic and native alike by
10
11
12 490 possibly providing inorganic N to new establishing grasses and offering more N to
13
14 already established perennial roots. As stated previously, the new N was most likely
15
16
17 492 derived from organic matter being translocated within the profile and/or the decaying root
18
19 system of the shredded tree.
20

21 494 Unexpectedly, higher levels of P availability associated with FWD elicited no
22
23 response among grasses and actually depressed the frequency of native *P. secunda*.
24
25
26 496 Generally, soil P availability is low in arid and semi-arid environments and potentially
27
28 limits the growth of grasses (Caldwell et al., 1987). Three of our four grasses were
29
30
31 498 relatively insensitive to changes in P availability. We find it difficult to explain why
32
33 higher levels of a limiting nutrient depressed plant frequency, but it may stem from
34
35
36 500 differences in grass species adaptations to low nutrient environments. Piñon-juniper
37
38 woodlands and grass species often establish on calcareous soils that strongly dictate the
39
40
41 502 adsorption and precipitation of P (Caldwell et al., 1991). In these soils, P is most often
42
43 bound by carbonates and free calcium, and grass species differ in their ability to mobilize
44
45
46 504 P via root exudates (e.g., phosphatase and inorganic acids; Marschner et al., 2006). If *P.*
47
48 *secunda* was adapted to effectively sequester relatively immobile P and the addition of
49
50
51 506 FWD increased the availability of P, then this native may have lost its competitive
52
53 advantage over other grass species and decreased in frequency. Alternatively, to help
54
55
56 508 sequester P many grasses, including *Poa* species, are facultatively mycorrhizal
57
58 (Vandenkoornhuyse et al., 2003) and mycorrhizal colonization in grasses tends to decline
59
60
61
62
63
64
65

1
2
3
4 510 following P fertilization (Eom et al., 1999). Thus, following higher P concentrations
5
6 induced by FWD, *P. secunda* may have lost the competitive benefits associated with its
7
8
9 512 mycorrhizal mutualism and decreased in frequency.

13 14 514 4.5 Microbial activity and native *P. spicata*

15
16 The depression of deep respiration following our FWD additions was associated with a
17
18
19 516 reduction in native *P. spicata* frequency. The addition of woody debris necessarily
20
21 coincides with changes in microbial metabolism. In the short-term (i.e., days to year),
22
23
24 518 metabolism is enhanced by a flush of labile C substrates (e.g., leaves, bark, and twigs)
25
26 entering soils, while in the long-term (i.e., years to decade), soil respiration may be
27
28
29 520 depressed by the more recalcitrant C sources, such as lignin (Dehlin et al., 2006). The
30
31 direct effect of soil respiration on plants is unclear. However, the lower respiration rates
32
33
34 522 under FWD additions may signal a decline in other processes associated with
35
36 microorganisms. For example, a decline in microbial activity may be related to a
37
38
39 524 depression in the release rates of essential plant micronutrients (Ranjbar and Jalali, 2012),
40
41 or a reduction in soil water holding capacity and retention due to less extracellular
42
43
44 526 polymeric substances being produced from metabolizing bacteria (Or et al., 2007). Thus,
45
46 other abiotic and biotic processes, indirectly associated with lower levels of respiration,
47
48
49 528 may have negatively influenced *P. spicata*.

50
51 As for microbial biomass, increases in biomass caused divergent effects on *P. spicata*
52
53 530 by depth. In surface soils, microbial biomass, such as growth promoting rhizobacteria,
54
55 possibly aided the establishment of seedlings. Bacterial species, such as Actinobacteria,
56
57
58 532 known to promoted seedling emergence and growth (Hayat et al., 2010), are common in
59
60
61
62
63
64
65

the rhizospheric soils of semi-arid grasses (Chen et al., 2015). Alternatively, in deeper soils, microbial biomass reduced *P. spicata* frequency. Changes in biomass are often accompanied by shifts in microbial community composition, which may influence grasses (Eom et al., 1999; Vandenkoornhuyse et al., 2003). For example, if FWD altered the mycorrhizal species diversity and richness in soils and the beneficial relationship between fungi and *P. spicata*, then *P. spicata* may be negatively impacted by FWD.

4.6 FWD and grass species advice to land managers

As land managers attempt to stem wildfires, the practice of shredding whole trees and the FWD it produces stimulates the establishment and persistence of exotic and native grass species alike. Thus, shredding may reduce the biomass of standing fuels (i.e., standing trees) but as grasses establish, the resulting biomass may act as a bridge to help fires jump breaks. The effects of FWD on grasses continued to intensify with time for perennials, but not for the exotic grass, suggesting that to gauge the effects of shredding on invasion or native grass establishment ecosystems should be monitored multiple times through time. Soils were important in understanding the effects of FWD on grass frequencies, especially in deep soils away from the layer of FWD on the surface. Deep soil processes are often overlooked in linking belowground and aboveground responses, but to be able to predict the changes in grass species after shredding, land managers should measure nutrient availability through the soil profile. Deep increases in net N mineralization, but not P availability, facilitated the increase of *B. tectorum* and *P. spicata* after shredding, suggesting that N and not P limit grass frequencies. Additionally, soil respiration and microbial biomass also influenced grass frequencies. But to more

1
2
3
4 556 fully explain these relationships, future studies need to link other microbial-mediated
5
6 processes (e.g., micronutrients, soil water relations) and specific bacterial and fungal
7
8
9 558 species (e.g., growth promoting bacteria and mycorrhizae) to grass response. Ultimately,
10
11 microbial activity and nutrients only partially explained the frequency of three of the four
12
13
14 560 grasses that we studied demonstrating that multiple other variables, like physiochemical
15
16 soil properties and external climate drivers, have the potential to also mediate the impact
17
18
19 562 of FWD on grasses.
20
21
22

23 564 **Acknowledgements**

25
26 We thank the US Joint Fire Sciences Program (JFSP Project ID:10-1-01-1) for funding
27
28
29 566 our research, and the USDI Bureau of Land Management, USDA Forest Service, and
30
31 Division of Wildlife Resources in Utah for implementation of treatments and allowing
32
33
34 568 this research on public and state lands.
35
36
37
38
39 570
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure Legends

Fig. 1 Three possible outcomes from logical mediation tests identifying the effects of FWD on links between species frequencies and soil characteristics. The tests included: 1) the effect of FWD on species frequencies was mediated through changes in soil variables due to woody additions, 2) both the FWD treatment and the effects of woody additions on soil variables partially mediated the variation in species frequencies, and 3) the FWD treatment alone explained the variation in grass frequencies.

Fig. 2 Fine woody debris (FWD) additions stimulated soil phosphorus (P) availability and dissolved organic carbon (DOC) in canopy surface soils, but depressed soil respiration in deep interspace soils. Microsite treatments included soils in barren plant interspaces (interspace), at the edge of the tree canopy or edge of FWD piles (edge), and underneath the tree canopy or beneath FWD (canopy). Values are intercepts, μ_{α} , with attending 95% confidence intervals from mixed models estimating the effects of FWD on five soil characteristics. Significant variables are indicated with an asterisk (i.e., 95% confidence interval of the intercept did not include zero). All data used in the model were based on differences between FWD and control treatments by woodland location ($n = 31$).

Fig. 3 The exotic annual, *Bromus tectorum*, and two native perennial grasses *Elymus elymoides* and *Pseudoroegneria spicata* increased in frequency following FWD, and soil characteristics explained the variation in species responses. For each of the four species, the intercepts, μ_{α} , from two different mixed models are presented, one that included

woodland sites as a random effect and time since FWD addition as a covariate (site), and another that included location and time but also soil variables (site + soil, $n = 31$). If soil variables explained any of the variation in grass frequencies, then the model intercept including soil variables should be closer to zero than the intercept excluding soil variables. Significant intercepts (i.e., 95% confidence interval of the intercept did not include zero) and reductions in the model intercept including soils are indicated with an asterisk.

Fig. 4 Changes in soils due to FWD partially mediated the effects of FWD addition on the exotic *B. tectorum* and native *P. spicata* and *Poa secunda*. The mixed model identifies links between soil variables and species frequency that were influenced by FWD ($n = 31$). Values are estimated unstandardized parameters. The model includes woodland locations as a random effect, and time since treatment or shredding as a covariate. The links between parameters (arrows) and corresponding values can be roughly interpreted as the percent change in the response variable (the one at the head of the arrow) given a 1% change in the explanatory variable (the one at the tail of the arrow). Solid arrows between soil variables and grass frequencies indicate effects that were significant ($p < 0.05$) and dashed arrows indicate marginally significant effects ($0.05 < p < 0.07$) between variables and grasses.

References

- 618 Aanderud ZT, Shuldman MI, Drenovsky RE, Richards JH (2008) Shrub-interspace
619 dynamics alter relationships between microbial community composition and
620 belowground ecosystem characteristics. *Soil Biol Biochem* 40:2206-2216 doi:
621 10.1016/j.soilbio.2008.04.008
- 622 Abbott LB, Roundy BA (2003) Available water influences field germination and
623 recruitment of seeded grasses. *J Range Manage* 56:56-64. doi: 10.2307/4003882
- 624 Bailey VL, Peacock AD, Smith JL, Bolton H (2002) Relationships between soil
625 microbial biomass determined by chloroform fumigation-extraction, substrate-
626 induced respiration, and phospholipid fatty acid analysis. *Soil Biol Biochem* 34:1385-
627 1389 doi: 10.1016/s0038-0717(02)00070-6
- 628 Bates D, Maechler M, Bolker B, Walker S. 2014, lme4: Linear mixed-effects models
629 using Eigen and S4_. R package version 1.1-7, [http://CRAN.R-](http://CRAN.R-project.org/package=lme4)
630 [project.org/package=lme4](http://CRAN.R-project.org/package=lme4)>
- 631 Bates JD, Miller RE, Svejcar T (2007) Long-term vegetation dynamics in a cut western
632 juniper woodland. *West N Am Naturalist* 67:549-561 doi: 10.3398/1527-
633 0904(2007)67[549:lvdia]2.0.co;2
- 634 Bates JD, Svejcar TJ, Miller RF (2002) Effects of Juniper cutting on nitrogen
635 mineralization. *J Arid Environ* 51:221-234 doi: 10.1006/jare.2001.0943
- 636 Bolker BM, Brooks ME, Clark CJ, Geange SW, Poulsen JR, Stevens MHH, White JS
637 (2009) Generalized linear mixed models: a practical guide for ecology and evolution.
638 *TREE* 24:127-135 doi:10.1016/j.tree.2008.10.008

Brooks ML, D'Antonio CM, Richardson DM, Grace JB, Keeley JE, DiTomaso JM, et al.
 (2004) Effects of invasive alien plants on fire regimes. *Bioscience* 54:677-688

Caldwell MM, Manwaring JH, Jackson RB (1991) Exploitation of phosphate from fertile
 soils microsites by three Great Basin perennials when in competition. *Funct Ecol*
 5:757-764 doi: 10.2307/2389538

Caldwell MM, Richards JH, Manwaring JH, Eissenstat DM (1987) Rapid shifts in
 phosphate acquisition show direct competition between neighboring plants. *Nature*
 327:615-616 doi: 10.1038/327615a0

Chen DM, Lan ZC, Hu SJ, Bai YF (2015) Effects of nitrogen enrichment on
 belowground communities in grasslands: Relative role of soil nitrogen availability vs.
 soil acidification . *Soil Biol Biochem* 89: 99-108 doi: 10.1016/j.soilbio.2015.06.028

Cline NL, Roundy BA, Pierson FB, Kormos P, Williams CJ (2010) Hydrologic response
 to mechanical shredding in a juniper woodland. *Rangeland Ecol Manag* 63:467-477
 doi: 10.2111/rem-d-09-00196.1

Dehlin H, Nilsson MC, Wardle DA (2006) Aboveground and belowground responses to
 quality and heterogeneity of organic inputs to the boreal forest. *Oecologia* 150:108-
 118 doi: 10.1007/s00442-006-0501-5

Ehrenfeld JG, Parsons WFJ, Han XG, Parmelee RW, Zhu WX (1997) Live and dead
 roots in forest soil horizons: Contrasting effects on nitrogen dynamics. *Ecology*
 78:348-362

Eom AH, Hartnett DC, Wilson GWT, Figge DAH (1999) The effect of fire, mowing and
 fertilizer amendment on arbuscular mycorrhizas in tallgrass prairie. *Am Midl Nat*
 142:55-70 doi: 10.1674/0003-0031

662

Hafner SD, Groffman PM (2005) Soil nitrogen cycling under litter and coarse woody debris in a mixed forest in New York State. *Soil Biol Biochem* 37:2159-2162 doi: 10.1016/j.soilbio.2005.03.006

666 Hayat R, Ali S, Amara U, Khalid R, Ahmed I (2010) Soil beneficial bacteria and their role in plant growth promotion: a review. *Ann Microbio* 60:579-598 doi: 10.1007/s13213-010-0117-1

James JJ, Davies KW, Sheley RL, Aanderud ZT (2008) Linking nitrogen partitioning and species abundance to invasion resistance in the Great Basin. *Oecologia* 156:637-648 doi: 10.1007/s00442-008-1015-0

672 Kerns BK, Thies WG, Niwa CG (2006) Season and severity of prescribed burn in ponderosa pine forests: Implications for understory native and exotic plants. *Ecoscience* 13:44-55 doi: 10.2980/1195-6860(2006)13[44:sasopb]2.0.co;2

Klopatek JM (2002) Belowground carbon pools and processes in different age stands of Douglas-fir. *Tree Physiol* 22:197-204

Laiho R, Prescott CE (2004) Decay and nutrient dynamics of coarse woody debris in northern coniferous forests: a synthesis. *Canadian J Forest Res-Revue Canadienne De Recherche Forestiere* 34:763-777. doi: 10.1139/x03-241

680 MacDougall AS, Turkington R (2005) Are invasive species the drivers or passengers of change in degraded ecosystems? *Ecology* 86:42-55 doi: 10.1890/04-0669

682 Mangla S, Sheley RL, James JJ, Radosevich SR (2011) Intra and interspecific competition among invasive and native species during early stages of plant growth. *Plant Ecol* 212:531-542 doi: 10.1007/s11258-011-9909-z

- Marschner P, Solaiman Z, Rengel Z (2006) Rhizosphere properties of Poaceae genotypes under P-limiting conditions. *Plant Soil* 283:11-24 doi: 10.1007/s11104-005-8295-5
- Miller PM, Eddleman LE, Kramer S (1990) Allocation patterns of carbon and minerals in juvenile and small-adult *Juniperus occidentalis*. *Forest Sci* 36:734-747
- Muller-Using S, Bartsch N (2009) Decay dynamic of coarse and fine woody debris of a beech (*Fagus sylvatica* L.) forest in Central Germany. *Eur J Forest Res* 128:287-296 doi: 10.1007/s10342-009-0264-8
- Murphy HT, Metcalfe DJ, Bradford MG, Ford AF, Galway KE, Sydes TA, et al. (2008) Recruitment dynamics of invasive species in rainforest habitats following Cyclone Larry. *Austral Ecol* 33:495-502 doi: 10.1111/j.1442-9993.2008.01904.x
- Norden B, Ryberg M, Gotmark F, Olausson B (2004) Relative importance of coarse and fine woody debris for the diversity of wood-inhabiting fungi in temperate broadleaf forests. *Biol Conserv* 117:1-10 doi: 10.1016/s0006-3207(03)00235-0
- Olsen SR, Cole CV, Watanabe FS, Dean LA (1954) Estimation of available phosphorous in soils by extraction with sodium bicarbonate. *USDA Cir. No. 939*
- Or D, Phutane S, Dechesne A (2007) Extracellular polymeric substances affecting pore-scale hydrologic conditions for bacterial activity in unsaturated soils. *Vadose Zone J* 6:298-305 doi: 10.2136/vzj2006.0080
- Owen SM, Sieg CH, Gehring CA, Bowker MA (2009) Above- and belowground responses to tree thinning depend on the treatment of tree debris. *Forest Ecol Manag* 259:71-80 doi: 10.1016/j.foreco.2009.09.044

- 1
2
3
4 706 Peek MS, Leffler AJ, Hipps L, Ivans S, Ryel RJ, Caldwell MM (2006) Root turnover and
5
6 relocation in the soil profile in response to seasonal soil water variation in a natural
7
8
9 708 stand of Utah juniper (*Juniperus osteosperma*). Tree Physiol 26:1469-1476
10
11 Peters DPC, Yao J, Browning D, Rango A (2014) Mechanisms of grass response in
12
13
14 710 grasslands and shrublands during dry or wet periods. Oecologia 174:1323-1334 doi:
15
16 10.1007/s00442-013-2837-y
17
18
19 712 Pregitzer KS, Euskirchen ES (2004) Carbon cycling and storage in world forests: biome
20
21 patterns related to forest age. Glob Change Biol 10:2052-2077 doi: 10.1111/j.1365-
22
23
24 714 2486.2004.00866.x
25
26 R core team (2015) A language and environment for statistical computing. R Foundation
27
28
29 716 for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, URL
30
31 <http://www.R-project.org>.
32
33
34 718 Ranjbar F, Jalali M (2012) Calcium, magnesium, sodium, and potassium release during
35
36 decomposition of some organic residues. Commun Soil Sci Plan 43:645-659 doi:
37
38 720 10.1080/00103624.2012.644005
39
40
41 Richardson AE, Simpson RJ (2011) Soil microorganisms mediating phosphorus
42
43 722 availability update on microbial phosphorus. Plant Physiol 156:989-996 doi: 10.
44
45 1104/ pp.111.175448
46
47
48 724 Rimer RL, Evans RD (2006) Invasion of downy brome (*Bromus tectorum* L.) causes
49
50 rapid changes in the nitrogen cycle. Am Midl Nat 156:252-258
51
52
53 726 Rose JA, Eddleman LE (1994) Ponderosa pine and understory growth following western
54
55 juniper removal. Northwest Sci 68:79-85
56
57
58
59
60
61
62
63
64
65

- 728 Roundy BA, Miller RF, Tausch RJ, Young K, Hulet A, Rau B, Jessop B, Chambers JC,
 730 Eggett D (2014a) Understory cover responses to piñon-juniper treatments across tree
 dominance gradients in the Great Basin. *Rangeland Ecol Manag* 67:482-494 doi:
 10.2111/REM-D-13-00018.1
- 732 Roundy BA, Young K, Cline N, Hulet A, Miller RF, Tausch RJ, Chambers JC, Rau B
 (2014b) Piñon-juniper reduction increases soil water availability of the resource
 734 growth pool. *Rangeland Ecol Manag* 67: 495-505
- Schade JD, Hobbie SE (2005) Spatial and temporal variation in islands of fertility in the
 736 Sonoran Desert. *Biogeochemistry* 73:541-553 doi: 10.1007/s10533-004-1718-1
- Schaeffer SM, Billings SA, Evans RD (2007) Laboratory incubations reveal potential
 738 responses of soil nitrogen cycling to changes in soil C and N availability in Mojave
 Desert soils exposed to elevated atmospheric CO₂. *Glob Change Biol* 13:854-865 doi:
 740 10.1111/j.1365-2486.2007.01324.x
- Shrout PE, Bolger N (2002) Mediation in experimental and nonexperimental studies:
 742 New procedures and recommendations. *Psychol Methods* 7:422-445 doi:
 10.1037//1082-989x.7.4.422
- 744 Spears JDH, Lajtha K (2004) The imprint of coarse woody debris on soil chemistry in the
 western Oregon Cascades. *Biogeochemistry* 71:163-175 doi: 10.1007/s10533-004-
 746 6395-6
- Tinker DB, Knight DH (2000) Coarse woody debris following fire and logging in
 748 Wyoming lodgepole pine forests. *Ecosystems* 3:472-483 doi:
 10.1007/s100210000041

- 1
- 2
- 3
- 4 750 Vandenkoornhuyse P, Ridgway KP, Watson IJ, Fitter AH, Young JPW (2003) Co-
- 5
- 6 existing grass species have distinctive arbuscular mycorrhizal communities. *Mol Ecol*
- 7
- 8 752 12:3085-3095 doi: 10.1046/j.1365-294X.2003.01967.x
- 9
- 10
- 11 Young KR, Roundy BA, Eggett DL (2013) Tree reduction and debris from mastication of
- 12
- 13 754 Utah juniper alter the soil climate in sagebrush steppe. *Forest Ecol Manag* 310:777-
- 14
- 15 785 doi: 10.1016/j.foreco.2013.09.024
- 16
- 17
- 18 756 Young KR, Roundy BA, Eggett DL (2014) Mechanical mastication of Utah juniper
- 19
- 20 encroaching sagebrush steppe increases inorganic soils N. *Appl Environ Soil Sci*
- 21
- 22 758 doi.org/10.1155/2014/632757
- 23
- 24
- 25 Zechmeister-Boltenstern S, Keiblinger KM, Mooshammer M, Peñuelas J, Richter A,
- 26
- 27 760 Sardans J, Wanek W (2015) The application of ecological stoichiometry to plant-
- 28
- 29 microbial-soil organic matter transformations. *Ecol Monogr* 85:133-155 doi:
- 30
- 31 762 10.1890/14-0777.1.sm
- 32
- 33
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
- 47
- 48
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60
- 61
- 62
- 63
- 64
- 65