Countdown to 2015: A decade of tracking progress for maternal, newborn and child survival

Cesar G. Victora^{1*}, Jennifer Harris Requejo², Aluisio J.D. Barros¹, Peter Berman³, Zulfiqar Bhutta⁴, Ties Boerma⁵, Mickey Chopra⁶, Andres de Francisco², Bernadette Daelmans⁵, Elizabeth Hazel⁷, Joy Lawn⁸, Blerta Maliqi⁵, Jennifer Bryce⁷

1. Federal University of Pelotas, 96001 Pelotas, Brazil
2. Partnership for Maternal, Newborn & Child Health, 1211 Geneva, Switzerland
3. Harvard School of Public Health, Boston, MA, USA
4. Center for Global Child Health, Hospital for Sick Children, Toronto, ON, Canada; Centre of Excellence in Women and Child Health, Aga Khan University, Karachi 74800, Pakistan
5. World Health Organization, 1211 Geneva, Switzerland
6. United Nation's Children's Fund, New York, NY, USA
7. Institute for International Programs, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, USA
8. London School of Hygiene and Tropical Medicine, London, UK

*Corresponding Author:

Cesar G. Victora, Federal University of Pelotas, 96001 Pelotas, Brazil; cvictora@gmail.com

Authors' contributions

CV, JR and JB wrote the first draft of the review. All authors contributed to revisions and writing.

Role of the funding source

Our work was supported by the Bill & Melinda Gates Foundation, the World Bank, and the Governments of Australia, Canada, Norway, Sweden, the USA, and the UK. The funders had no role in the conceptualization of the paper or in the material presented.

ACKNOWLEDGEMENTS:

Countdown would like to thank the following: Lois Park, Colleen Murray, Agbessi Amouzou, Liliana Carvajal, Courtney NG, Neha Singh, Giovanni Franca, Fernando Wehrmeister, Amani Syam, Matthews Matthai, Jo Borghi, Robin Gorna, Adam Deixel, Corrina Moucheraud, Helen Owen, , Robert Black, Li Liu

ABSTRACT

Conceived in 2003 and born in 2005 with the launch of its first report and country profiles, the Countdown to 2015 for maternal, newborn and child survival has reached its originally proposed lifespan. Major reductions in the deaths of mothers and children have occurred since Countdown's inception, even though most of the 75 priority countries failed to achieve Millennium Development Goals 4 and 5. The coverage of life-saving interventions tracked in Countdown increased steadily over time, but wide inequalities persist between and within countries. Key drivers of coverage such as financing, human resources, commodities and conducive health policies also showed important, yet insufficient increases. As a multi-stakeholder initiative of over 40 academic, international, bilateral and civil society institutions, Countdown was successful in monitoring progress and raising the visibility of the health of mothers, newborns and children. Lessons learned from this initiative have direct bearing on monitoring progress during the Sustainable Development Goals era.

INTRODUCTION

Ten years after Countdown to 2015 for Maternal, Newborn and Child Survival (Countdown) was launched, the world has become a healthier place for children and women. The Millennium Development Goals (MDG) sparked those involved in the 2003 Lancet Child Survival Series to propose the Countdown initiative, who pledged to hold regular conferences for “*ensuring that there is an overall mechanism for improving accountability, re-energising commitment, and recognizing accomplishments in child survival*”.¹

In response to this call to action, Countdown launched its first report in 2005,² which was followed by six others (2008,³ 2010,⁴ 2012,⁵ 2013,⁶ 2014⁷). The final report is being released this week, October 17, in Mexico City.⁸ From its original focus on child survival, Countdown expanded to track progress on reproductive, maternal, newborn and child health indicators relevant to MDGs 4 (reduce child mortality) and 5 (improve maternal health). The heart of the Countdown reports are the two-page country profiles (www.countdown2015mnch.org) that summarize the most recent data on intervention coverage, maternal and child mortality, and nutrition. The country profiles also highlight socioeconomic inequalities in intervention coverage, and two of the main drivers of coverage: health systems and policies, and financing.

Insert Panel 1 here

Countdown evolved in many ways during the past decade. The partnership grew from 11 to 43 institutional stakeholders, and the number of countries monitored from 60 to 75, comprising over 95% of global deaths of mothers and children. As the scope shifted beyond child survival, and in response to new evidence, the number of indicators tracked expanded from 35 to 73. By including new, proven interventions in its profiles even before data were available for many countries, Countdown has contributed to the identification of data gaps and arguably helped raise the visibility of such new interventions. Countdown recognized the importance of engaging at country level, and from 2012 embarked on a set of case studies aimed at understanding how countries have achieved progress (webannex A). Countdown is also the primary source of coverage information for the Commission on Information and Accountability for Women’s and Children’s Health (http://www.who.int/woman_child_accountability/about/coia/en/) and the Independent Expert Review Group (iERG)⁹ reports.

Insert Panel 2 here

In the decade since Countdown’s inception, the number of reports on specific RMNCH issues has grown rapidly. Countdown’s particular niche has been its action-oriented focus on intervention coverage, and the user-friendly synthesis of information in the country profiles. Our principles have not changed over this period¹: focus on monitoring the coverage of evidence-based, cost-effective interventions; maintain a country orientation; and build upon existing goals and monitoring efforts. Countdown’s commitment to these principles and its products have helped increase the global visibility of RMNCH, and has plausibly contributed to

reversing the slow rate of progress in reducing child, newborn, and maternal mortality in the 1990s that was documented in early public health series in *The Lancet*^{1, 10-12}

As the MDG era draws to a close, the global public health and development communities must take stock of progress over the past 15 years, and get ready for the Sustainable Development Goal era (<https://sustainabledevelopment.un.org/topics/sustainabledevelopmentgoals>). We assess changes in the evidence and in data availability over this period, and their implications for program managers and decision makers. We conclude by turning a critical lens on the SDG framework and future accountability efforts, drawing on Countdown's monitoring experience.

METHODS

We have reported previously on Countdown's data sources and methods,^{13, 14} and these are described in detail in Webannex B. Most coverage, equity and nutrition data come from standardized population based surveys (Demographic and Health Surveys – DHS, Multiple Indicator Cluster Surveys - MICS, or other nationally representative surveys that meet data quality standards). Mortality data are provided by United Nations sources.

Intervention coverage was monitored for the 75 Countdown countries with data from 2009 to 2015. Coverage trends are derived from countries that had at least two data points, one from between 2000-2008, and the second from 2009-2014. The “composite coverage index” (CCI) is a weighted average of eight interventions along the continuum of care, all of which have been available in most countries for at least a decade.^{15, 16} Results on socioeconomic inequalities are based on wealth quintiles calculated on the basis of household assets.¹⁶ Trends in CCI were estimated using 47 Countdown countries with at least two surveys (either DHS or MICS) where CCI was available, with survey years ranging from 1994 to 2014, using a multilevel model (see Webannex B).

Health systems and policy indicators are derived from global databases maintained by the WHO and other UN organizations, and financing data are abstracted from datasets maintained by the Organization for Economic Co-operation and Development Assistance Committee.

The 2015 Countdown Results

In this section, we summarize findings for 2015 with an emphasis on time trends; the full results are available in the 2015 report (www.countdown2015mnch.org).

Based on modeled estimates, the maternal mortality ratio has decreased by around 45% over the past two decades, and the number of annual maternal deaths has dropped from around 523,000 to 289,000.¹⁷ Although mortality reductions appear to have accelerated – three - quarters of Countdown countries reduced maternal mortality faster in 2000-2013 than in the 1990s¹⁴ – very few Countdown countries will achieve MDG 5. A 2009 analysis of maternal deaths worldwide found over half of these were due to haemorrhage, hypertensive disorders, and sepsis – causes that are preventable through the provision of quality antenatal, delivery and postnatal care.¹⁸ Recognition of the role of contraception in reducing maternal and newborn deaths has led to a major increase in resources for family planning programs¹⁹. Evidence of the importance of reaching adolescents with family planning and nutrition programs for improving birth outcomes, as well as their own health, has also resulted in increased attention to this population group^{20, 21}.

The growing recognition of stillbirths as a public health problem - 2.6 million annual third trimester stillbirths, with 1.2 million of these occurring in the intrapartum period – led Countdown to report on the stillbirth rate since 2010. Unfortunately, global visibility for stillbirths may remain limited in the SDG era given the absence of a specific target.²²

The global under-five mortality rate has dropped by 53% since 1990, from 91 deaths per 1,000 live births to 43 in 2015.²³ And, the global annual rate of reduction has steeply accelerated over time, suggesting that more progress can be expected in the coming years. In 2000, the leading causes of the 10.8 million under-five deaths were neonatal conditions (33%), diarrhea (22%), pneumonia (21%), malaria (9%) and HIV (3%).²⁴ Estimates for 2015 indicate 5.9 million deaths a year²³ with a major shift in the causes of death: preterm birth complications cause 18% of all under-five deaths, and along with other neonatal causes represent 45% of all deaths. Pneumonia (16%), diarrhea (9%), malaria (5%) and AIDS (1%) deaths have declined in relative terms, and even more so in absolute terms.²⁵ The growing concentration of deaths in the newborn period and improved understanding about causes of newborn deaths has sparked the scale up of long-existing interventions and the development of new ones.

The past 10 years have also witnessed a growing understanding of the role of nutrition in mortality and human development.^{26, 27} Sub-optimal nutrition - including fetal growth restriction, stunting, wasting, and deficiencies of vitamin A and zinc along with suboptimum breastfeeding--is an underlying cause of 45% of all deaths of children under five years of age.²⁶

Progress in the Countdown countries in reducing maternal and under-five mortality since 1990 is summarized in Table 1. One third of the 75 countries achieved the annual rate of reduction of 4.4% or higher in under-five mortality that was needed to reach MDG4 in 2015, but only six countries have achieved the annual rate of reduction of 5.5% or higher in maternal mortality required by MDG5. The successful countries are flagged in Table 1. Four countries – Cambodia, Nepal, Rwanda, and Eritrea – have achieved both MDG 4 and 5.

Table 1 also shows another measure of progress, based on whether each country has met the mortality thresholds originally set for inclusion as a Countdown country. There are two such thresholds. The first group of 60 countries was selected in 2005 for monitoring by Countdown based on their high levels of under-five mortality – either an under-five mortality rate ≥ 90 per 1,000 live births, or an absolute annual number of deaths $\geq 50,000$. Among these 60 countries, 28 have now “graduated” from Countdown by reducing child mortality below this threshold. In 2008, Countdown broadened its scope, by including eight additional countries with maternal mortality ratios of > 550 per 100,000 live births, or a ratio of > 200 plus an absolute annual number of maternal deaths ≥ 750 . Of the eight, seven have now graduated by reducing maternal mortality below this threshold. The remaining seven priority countries were added after 2008, to maintain consistency with the priority low-income countries listed in the Global Strategy for Women’s and Children’s Health, and to include South Sudan after it was formed in 2012.

Alternative estimates for maternal²⁸ and child²⁹ mortality are available from the Institute of Health Metrics and Evaluation. Although their estimates of levels and trends for specific countries may differ from those presented in Table 1, the overall conclusions are the same: a small minority of LMICs will reach either MDG 4 or 5.

Insert Table 1 around here

In the next four sections, we summarize recent results from Countdown's four streams of technical work: coverage, equity, health systems and policies, and financing. We also highlight progress in data availability and some of the most important data gaps that remain in each of these areas.

Figure 1 shows coverage across the continuum of care for selected high-impact interventions, and the service contacts through which these and other interventions can be delivered. The bars show the median coverage reported by all Countdown countries with data available in 2009 or later; the grey dots show the values for individual countries. Coverage levels are particularly low around the time of birth, and for case-management interventions for childhood illnesses. The list of coverage indicators tracked in Countdown has evolved over time. For example, indicators for rotavirus vaccine and pneumococcal conjugate vaccine were added in 2014 because of increased data availability following rapid policy adoption by countries. We no longer track the indicator of antibiotic treatment for childhood pneumonia, because validation studies have shown that this cannot be measured accurately in household surveys.³⁰ We have retained the indicator on oral rehydration therapy (ORS and/or increased fluids and continued feeding) to allow the examination of trends. However, WHO/UNICEF guidelines now recommend ORS and zinc, and in the future it will be important to track coverage for both. In 2015, 37 countries had coverage data on zinc treatment of diarrhea. The median coverage was 1%, with a high of 28% in Malawi. Results for the full set of Countdown coverage indicators are available in Webannex C, and Webannex D provides information on HIV and c-section indicators.

Insert Figure 1 about here

Data availability (Figure 1) has improved for all indicators measured through household surveys. For example, the number of countries with information on postnatal care visits for babies increased from 5 during the period 2000-2006³ to 35 in 2009-2014. The rapid expansion of household surveys in the wake of the MDGs led to real progress in ensuring that all countries have recent, high-quality data to guide their programs and policies^{31 32}

Our results continue to show unacceptably low coverage for most of these interventions and service contacts, with enormous ranges among countries around the global median values. Even interventions for the prevention of malaria, which have shown greater accelerations in coverage than any other indicators in recent years,³³ are not reaching their life-saving potential. In endemic countries with available data, only 23 percent of women report receipt of malaria prevention during pregnancy, and only 38 percent of children under five were reported to be sleeping under an insecticide-treated net. Treatment interventions are still reaching fewer than 40 percent of children with malaria or diarrhea, and only 54 percent of children with symptoms of pneumonia are taken outside the home for care. Immunizations continue to be an exception, with median coverage generally above 85 percent, although inter-country variation is still marked.

Time trends in intervention coverage were featured in an earlier publication³³ and are updated in Table 2. Three broad patterns are evident. First, the three malaria interventions that started below 40% in the earlier period showed substantial increases over time. HIV interventions are not shown in Table 2 because baseline data were not available due to methodological changes,

but the prevention of mother to child transmissions with anti-retrovirals reached 53% in the most recent period with a range of 1->95% across countries with data. Second, some interventions already showed high coverage around 2000, and increased modestly in absolute terms, partly because there was limited scope for increase; these include antenatal care (at least one visit), improved drinking water sources, and the three vaccines. Nevertheless, a substantial proportion of the gap was closed for these interventions. The third group includes all other interventions, which had coverage below 60% up to 2009 and increased by 10 percent points or less: family planning, four or more antenatal care visits skilled birth attendance, improved sanitation facilities, exclusive breastfeeding and all the case management interventions against diarrhea and pneumonia. These patterns suggest that rapid coverage increases are possible when interventions are prioritized and generously funded as in the case of the malaria or HIV indicators. On the other hand, there was very limited progress for interventions that require service contacts along the continuum of care and 24/7 service availability, particularly during pregnancy and childbirth, and for the management of childhood diarrhea and pneumonia.

Insert Table 2 about here

Further improvement is needed urgently in the quantity and quality of data for monitoring coverage. For example, there is no guarantee that women and children who report a service contact actually receive the full complement of interventions that should be delivered during that contact. Reported coverage for antenatal, delivery and postnatal care therefore represent “best case scenarios” for actual coverage of interventions that should be provided during those contacts. New Countdown secondary analyses of patterns of antenatal care in seven countries show widely varying patterns of drop-off between the first antenatal care visit and subsequent visits, with women reporting more visits having a higher probability of receiving recommended interventions.⁸ Redoubled efforts are needed to ensure contact with health services results in the delivery of all recommended life-saving interventions. New approaches are also needed to link household surveys and health facility survey data, to generate essential data on service quality that are currently lacking (see Webannex E).

The equity dimension was noticeably absent from the MDGs.³⁴ Since its first report, Countdown has provided original analyses of inequalities in intervention coverage by wealth, sex of the child, place of residence and other social determinants. These analyses consistently show systematic pro-rich inequalities for virtually all coverage indicators.³⁵ The gaps are wider for interventions, such as antenatal care (four or more visits) and skilled birth attendance, that require access to fixed health facilities, compared to interventions such as immunization that can be delivered at community level.³⁵ Our analyses have also shown that countries that made rapid progress in coverage were those that effectively reached the poorest families.³⁶

The growing availability of surveys allows disaggregated analyses of trends in coverage for the poorest and richest quintiles of mothers and children, across a large number of countries. Figure 2 shows these results for the composite coverage index in 47 Countdown countries with available information from multiple surveys. Globally, CCI coverage increased for both the richest and poorest quintiles, but the increase was steeper among the poorest (1.0 percent

point per year, 95% CI 0.8-1.1) than for the richest (0.3 percent point; 95% CI 0.2-0.4)(Figure 3a). The rich-poor gap declined from 28 percent points in 2000 to 19 points by 2014. This indicates an increase in coverage equity in both absolute and relative terms (both trends with $P<0.001$)(Figure 3b).

Insert Figure 2 about here

Whereas the coverage gap between rich and poor mothers and children persists, inequalities are decreasing, at least for the eight long-standing interventions that are part of the CCI.

Similar to coverage, data availability for equity analyses has improved, but there is still much scope for progress. Repeated surveys using consistent measurement of equity stratifiers, such as wealth, gender, residence or ethnicity, are required to identify priority groups and track subnational progress over time.

Supportive policy environments and functional health systems with adequate human resources are prerequisites for high and equitable coverage. The number of policy and systems indicators tracked in Countdown has increased from five in 2005 to 11 in this year's report, including four measures of systems that are critical to effective service delivery for women and children. These indicators are consistent with international frameworks^{37 38}. Although further work is needed to develop comparable metrics for implementation strength at national and sub-national levels, Countdown is finalizing a set of tools for describing relevant policies and aspects of RMNCH programme implementation across countries and over time.

Notable progress in adopting supportive policies has occurred across the Countdown countries (Webannex F shows adoption rates for 10 essential policies based on a 2013-2014 WHO survey). Figure 3 shows that the number of countries that adopted each of six policies for which trend data are available increased markedly between 2008 and 2014. However, more progress is needed, particularly for policies that are lagging such as maternity protection in accordance with Convention 183 of the International Labour Organisation (including maternity leave and employment protection during pregnancy and the postnatal period) and the International Code of Marketing for Breastmilk Substitutes.

Insert Figure 3 about here

Increased investment in information systems has resulted in an expansion of available data on skilled health professionals.³⁹ Three-quarters of Countdown countries remain below 22.8 physicians, nurses and midwives per 10,000 population, a threshold WHO considers necessary to achieve relatively high coverage of essential health interventions³⁹. The most recent estimates show a median density of skilled health professionals of 10.2 per 10,000 population in the Countdown countries, ranging from 1.6 in Madagascar and Niger to 142 per 10,000 population in Uzbekistan.

Countdown has contributed to substantial increases in data availability on policies by helping shape the contents of the WHO's biannual policy survey. More work is needed to overcome the limitations of using a survey-based approach, to ensure more timely and accurate information, and to explore associations between policies and changes in coverage levels. Efforts are under way to assess the feasibility of collecting data on selected tracer indicators of the quality of care (Webannex E).

The establishment of the MDG framework led to a major upswing in political prioritization of women's and children's health,⁴⁰ and official development assistance (ODA) surged after the MDG summit in 2000⁴¹. Although the growth in ODA slowed following the economic crisis in 2007-2008, trends from 2003 to 2012 in Countdown countries show a tripling of ODA to MNCH (from \$2 to \$6 billion). Most of the investment was for general health care (including health systems support), reproductive health, malaria programs, and immunization programs.⁴¹ During this same time period, ODA for projects that mention newborns grew exponentially, from \$33 million to \$1 billion, reflecting the growing visibility of newborn health, yet far too little given that neonatal causes represent almost half of under-five deaths. A comparison of the present estimates with those produced by the Institute of Health Metrics and Evaluation, the Partnership for Maternal Newborn and Child Health and United Nations Population Fund is available in the full Countdown report.⁸ Although these funding increases and the spike in high level commitments to RMNCH between 2010 and 2015 are encouraging, more is needed. In particular, better targeting of resources to countries with the greatest need could help narrow equity gaps and help achieve the next set of goals.

Each of the Countdown case studies include a detailed analysis of trends in financial flows to RMNCH, and illustrate the complexity of the funding environment. The Peru and Ethiopia case studies, for example, both showed rapid recent growth in RMNCH expenditures⁴². Yet the two countries used different resources and financing mechanisms to fund their programs. Peru, an upper-middle income country, relied mostly on domestic funding, while Ethiopia, a low income country, relied heavily on external funding. Both countries have high levels of out-of-pocket spending, which must be addressed to make healthcare more affordable to disadvantaged population groups.

What about the future? The general consensus across the panoply of resource tracking efforts is that ODA expenditures for RMNCH are increasing.^{41, 43} For domestic spending there is insufficient data on RMNCH expenditure to estimate trends. The work of the Lancet Global Commission on Investing in Health, which emphasizes the centrality of RMNCH for achieving global development, and the World Bank's recently announced Global Financing Facility for RMNCH, are signs that investments in RMNCH will continue to grow.^{44 45} However, the multiplicity of goals and targets in the SDG framework could detract from a sustained and accelerated focus on RMNCH, leaving many donor-dependent countries short of funds. The most important future strategies may be those that shore up the contribution of national sources to RMNCH, using mechanisms such as the establishment of concrete accountability procedures that strengthen the collaboration between citizens and governments and build more sustainable and efficient funding for RMNCH at local levels.

MONITORING AND ACCOUNTABILITY: HOW CAN THE CD EXPERIENCE INFORM THE SDGs

The MDGs encouraged global political consensus, provided a focus for advocacy and visibility, improved the targeting and flow of aid, and strengthened monitoring of process and outcome indicators.⁴⁰ Within the context of the health MDGs, Countdown played a unique role as a multi-stakeholder initiative. Countdown's success in monitoring and accountability at country level for MDGs 4 and 5^{46, 47} was complemented by other initiatives, such as the Independent Expert Review Group,⁹ that relied heavily on our data to push for global level accountability. Our experience has inspired others. For example, advocates for the Non-Communicable Diseases Countdown 2025 wrote: "*lessons from Countdown to 2015 include the*

importance of collaboration and inclusiveness; adaptation of global targets to the national situation; regular measurement; transparent review and publication of progress on priority interventions and outcomes; strong engagement of academia and civil society; regular reports based on fairly simple summaries of country progress; and adequate resources. Countdown to 2015, while retaining a core of basic information, has evolved to include detailed country reports and shows the value of a strong and independent partnership for global health.”⁴⁸

Leaders of global initiatives developed to track physical activity (PC Hallal, personal communication) and nutrition (L Haddad, personal communication) also acknowledge that Countdown has inspired their work.

This final Countdown report focuses on time trends over the past 15 years. Intervention coverage increased for most interventions, and particularly for those that received substantial donor investment such as interventions against malaria and HIV.^{7, 14} The gap in coverage between rich and poor seems to be narrowing, at least for interventions that have been available in low-income countries for many years. More countries are adopting RMNCH policies, and financing for the health of mothers, newborns and children has increased. Each of these statements about general trends, however, must be qualified. Cost-effective interventions are still failing to reach a large proportion of those who need them. Socioeconomic inequities in coverage are still rampant within virtually every Countdown country. Essential policies and sufficient and equitably distributed human resources and commodities are still lacking in most countries. And in spite of increased funding, there is still a huge shortfall of reliable resources for RMNCH.

The post-2015 Sustainable Development Goals (SDGs) are unquestionably much broader and more complex than the MDGs. Their adoption will bring about substantive challenges to monitoring and accountability. To conclude, we address how lessons learned from the Countdown process may be relevant to the SDG era.

The MDGs were launched in 2000 with a baseline of 1990, a peculiar choice implying that signatory countries were held accountable for trends during the preceding 10 years. In addition, the absence of real time data on births and deaths led to complex modeling procedures for estimating concurrent and past maternal and child mortality. Reliance on these methods meant that historical trends were revised every time new data became available. Baseline values therefore kept changing, and countries seemed to be aiming at moving targets. Fortunately, the SDG starting date is fixed in 2015, but substantial investments in data collection are urgently needed to provide valid and precise baseline levels.

Whereas child mortality estimates are based on actual data from censuses, surveys or civil registration, maternal mortality is modeled for most LMICs on the basis of predictor variables such as gross domestic product per capita, fertility, HIV prevalence and coverage of skilled birth attendants¹⁷. There are important problems with the acceptability of modeled estimates at country level, yet very few countries have invested in large-scale surveys or registration systems to actually measure maternal mortality,^{49 50}

In summary, monitoring during the MDG era was largely dependent on modeled mortality estimates, coverage measurements derived from infrequent household surveys, and ad-hoc systems for tracking policies, health system measures and funding flows. In addition, the MDG framework has been largely criticized for its neglect of equity. If the international community is serious about achieving the SDGs, it must invest now in improvements and innovations in

measurement.⁵¹ The area of measuring effective coverage and the quality of preventive and curative interventions deserves special attention.

There are other aspects of the SDGs that the Countdown experience suggests are likely to be problematic. For example, MDGs 4 and 5 were clearly defined, with quantitative goals for relative mortality reductions. The corresponding SDGs (3.1 and 3.2) propose absolute targets of 25 per thousand live births at national level for under-five mortality, 12 per thousand live births for neonatal mortality, and a global maternal mortality ratio of 70 per 100,000 live births. According to 2013 estimates, six CD countries already have U5MRs below 25, and 11 have MMRs below 70 (Table 1). What type of progress – if any – should these countries be aiming at? Based upon the Countdown experience of interacting with countries regarding progress towards global goals, we believe that the provisional SDG targets must be urgently revisited.

It is also important for targets to be realistic. The fact that only 25 and 6 of the 75 Countdown countries will likely reach MDGs 4 and 5, respectively, suggests that the original targets were too ambitious in light of the observed progress in terms of financial flows and health systems factors such as human resources and policies.

The Countdown experience taught us the importance of identifying evidence-based interventions, and tracking coverage change over time on a country-by-country basis. Although monitoring 73 coverage indicators is a huge task, it provides specific feedback on what needs to be improved, and where. This level of detail is essential, but we also felt a need for a summary measure – the composite coverage index. We found this to be robust, stable, and highly associated with measures of mortality.⁷ The CCI provides a promising approach to the measurement of universal health coverage, and includes a focus on equity, two pillars that lie at the heart of the health SDG.

The large number of SDG targets – 169 at some stage – has brought about substantial criticism.^{52, 53} Although work is under way to trim down this list, the Countdown experience shows that an initially small list of goals and core indicators can rapidly expand over time, as new interventions become available, and as different interest groups lobby – often with strong justification – for inclusion of additional indicators. A rigorous technical process must be in place for ensuring the validity and reliability of these new indicators, for making sure monitoring efforts stay coordinated and focused, and to minimize the reporting burden on countries.

Countdown found its niche in a plethora of global initiatives: a focus on intervention coverage at country level. Our focus does not overlook other drivers of women's and children's health, such as social or environmental determinants.⁵⁴ We recognize that biomedical interventions are one of the most important pathways through which distal, social determinants affect health. During the SDG era, tracking progress in social and environmental determinants and understanding how these impact health and development will be just as important as tracking progress in coverage or health status, and will likely face similar data availability challenges as were faced by Countdown.

Other aspects of the Countdown experience may be relevant to similar initiatives in the SDG era. Involving multiple stakeholders is essential for ensuring that data lead to action. Retaining scientific independence in the midst of stakeholders with different interests and agendas was not always a smooth process, but Countdown managed to achieve and maintain consensus on indicators that should be monitored and disseminated. Positive pressure from stakeholders led

to our expansion from child survival in 2003 to the full RMNCH continuum of care. Balancing focus with breadth is not easy, and will likely be even harder in the SDG era, both within the health goal and across all 17 SDGs.⁵⁵ Countdown engaged with civil society at the international level, as several of its stakeholders represent such constituencies. Engagement at national level, however, was mostly limited to countries where case studies were carried out and disseminated. Nevertheless, several of the key Countdown stakeholders led activities at national level that were consistent with our mandate and harmonized with our main messages.

A Viewpoint in this issue of The Lancet explores future directions for Countdown. Regardless of what lies ahead, our experience in the past ten years has established the importance and feasibility of a multistakeholder initiative, with independence and a strong technical component, in accelerating progress for the world's mothers and children. The auspicious launch of a Global Financing Facility⁴⁴ in support of the Every Woman Every Child initiative highlights the need for continued, independent monitoring and accountability in the future.

Panels

Panels: Lancet article

Panel 1. Key messages
<ul style="list-style-type: none">• Countdown to 2015 is a unique example of a multi-stakeholder initiative of academic, international, bilateral and civil society institutions focused on monitoring progress in the health of the world's women and children• Over a decade, regular reports from Countdown showed important progress by many high burden countries in increasing coverage with cost-effective, evidence-based health interventions, and in preventing deaths of mothers and children• Yet, substantial, reliable investments are required in order to achieve global targets and reduce inequalities between and within countries• Lessons learned from Countdown can inform global monitoring and accountability in the era of Sustainable Development Goals

Figures

Figure 1. Coverage of interventions varies across the continuum of care

* Countries where at least 75% of the population is at risk of malaria and where a substantial proportion (50% or more) of malaria cases is due to *Plasmodium falciparum* (n= 44) or where 50–74% of the population is at risk of malaria and where a substantial proportion (50% or more) of malaria cases is due to *P. falciparum* (n= 8)

Source: Immunization rates, WHO and UNICEF; postnatal visit for mother, Saving Newborn Lives analysis of Demographic and Health Surveys; improved water and sanitation, WHO and UNICEF Joint Monitoring Programme for Water Supply and Sanitation; all other indicators, UNICEF global database, July 2015, based on Demographic and Health Surveys, Multiple Indicator Cluster Surveys and other national surveys; does not include 2014–2015 Rwanda data.

Figure 2. (a) Global trends in the composite coverage index at national level and for the poorest and richest 20% of the national samples in 47 countries. (b) Global trends in absolute (slope index of inequality) and relative (concentration index) inequalities in the composite coverage index. Shaded areas show the 95% confidence bands for the regression lines.

Figure 3. Trends in the adoption of selected tracer policies at macro and micro policy level, 2008-2014 (68 Countdown countries, %)

Tables

Table 1. Countdown countries and 'graduation' status based on original entry criteria

Country	Year entered CD	Selection criteria, why entered	U5MR	% annual reduction (1990-2015)	Number of deaths (thousands)	Share of U5 deaths occurring neonatally	MMR	% annual reduction (1990-2015)	Number of maternal deaths	Country graduated from CD (yes/no)
			2015		2015		2013		2013	
Afghanistan	2005	U5MR≥90 and ≥50,000	91.1	2.7	94261	38.4	400	4.7	4200	no
Angola	2005	U5MR≥90 and ≥50,000	156.9	1.5	169310	31.4	460	4.9	4400	no
Azerbaijan	2005	U5MR≥90	31.7	4.4	7206	59.2	26	3.6	43	yes
Bangladesh	2005	≥50,000 child deaths	37.6	5.4	119326	62.3	170	5.0	5200	no
Benin	2005	U5MR≥90	99.5	2.4	37092	32.2	340	2.4	1300	no
Bolivia	2008	MMR>200 + ≥750 maternal deaths	38.4	4.7	9415	51.2	200	4.0	550	yes
Botswana	2005	U5MR≥90	43.6	0.9	2488	51	170	3.1	83	yes
Brazil	2005	≥50,000 child deaths	16.4	5.2	52415	54.6	69	2.4	2100	no
Burkina Faso	2005	U5MR≥90 and ≥50,000	88.6	3.3	60477	30.3	400	2.9	2800	no
Burundi	2005	U5MR≥90 and ≥50,000	81.7	3	36970	35.8	740	2.3	3400	yes
Cambodia	2005	U5MR≥90 and ≥50,000	28.7	5.6	10257	51.5	170	8.1	670	yes
Cameroon	2005	U5MR≥90 and ≥50,000	87.9	1.8	71348	29.5	590	0.9	4900	no
Central African Republic	2005	U5MR≥90	130.1	1.2	21029	33.3	880	1.3	1400	no
Chad	2005	U5MR≥90 and ≥50,000	138.7	1.7	82728	28.8	980	2.3	5800	no
China	2005	≥50,000 child deaths	10.7	6.5	181574	51.5	32	4.7	5900	no
Comoros	2011	low income	73.5	2.1	1897	46.9	350	2.6	90	
Congo	2005	U5MR≥90	45	2.9	7269	40.6	410	2.1	690	yes
Congo, Democratic Republic of	2005	U5MR≥90 and ≥50,000	98.3	2.6	304558	30.9	730	1.5	21000	no

Côte d'Ivoire	2005	U5MR≥90 and ≥50,000	92.6	2	75393	41.7	720	0.1	5300	no
Djibouti	2005	U5MR≥90	65.3	2.4	1429	51.6	230	2.4	55	yes
Egypt	2005	≥50,000 child deaths	24	5.1	65775	54.5	45	4.1	860	no
Equatorial Guinea	2005	U5MR≥90	94.1	2.8	2655	35.6	290	7.0	79	no
Eritrea	2008	MMR>200 + ≥750 materna deaths	46.5	4.7	7764	39.4	380	6.2	880	no
Ethiopia	2005	U5MR≥90 and ≥50,000	59.2	5	184186	47.5	420	5.0	13000	no
Gabon	2005	U5MR≥90	50.8	2.4	2579	46.3	240	2.0	130	yes
Gambia	2005	U5MR≥90	68.9	3.6	5540	44.6	430	2.1	340	yes
Ghana	2005	U5MR≥90 and ≥50,000	61.6	2.9	54061	47	380	2.9	3100	no
Guatemala	2008	MMR>200 + ≥750 materna deaths	29.1	4.1	12858	46.2	140	2.8	660	yes
Guinea	2005	U5MR≥90 and ≥50,000	93.7	3.7	42073	34	650	2.2	2800	no
Guinea-Bissau	2005	U5MR≥90	92.5	3.6	5883	44	560	2.2	360	no
Haiti	2005	U5MR≥90	69	3	17841	36.6	380	2.4	1000	yes
India	2005	U5MR≥90 and ≥50,000	47.7	3.9	1200998	57.9	190	4.5	50000	no
Indonesia	2005	≥50,000 child deaths	27.2	4.5	147162	50.2	190	3.5	8800	no
Iraq	2005	U5MR≥90 and ≥50,000	32	2.1	38682	58.1	67	2.0	710	yes
Kenya	2005	U5MR≥90 and ≥50,000	49.4	2.9	74429	45.3	400	0.8	6300	no
Korea, Democratic People's Republic of	2008	MMR>200 + ≥750 materna deaths	24.9	2.2	9271	54.9	87	-0.1	310	yes
Kyrgyzstan	2011	low income	21.3	4.5	3644	54.5	75	0.5	110	
Lao People's Democratic Rep	2008	MMR>550	66.7	3.6	11613	44.9	220	6.8	400	yes
Lesotho	2008	MMR>550	90.2	-0.1	5570	36.7	490	1.7	280	yes
Liberia	2005	U5MR≥90	69.9	5.2	10509	34.7	640	2.8	980	yes
Madagascar	2005	U5MR≥90 and ≥50,000	49.6	4.7	40075	40.4	440	2.3	3500	yes
Malawi	2005	U5MR≥90 and ≥50,000	64	5.3	40048	34.3	510	3.2	3400	yes
Mali	2005	U5MR≥90 and ≥50,000	114.7	3.2	82710	33.2	550	3.1	4000	no

Mauritania	2005	U5MR≥90	84.7	1.3	11050	42.5	320	2.9	430	no
Mexico	2005	>50,000 child deaths	13.2	5	31278	53.1	49	2.5	1100	yes
Morocco	2008	MMR>200 + ≥750 maternal deaths	27.6	4.3	19759	64.3	120	4.1	880	yes
Mozambique	2005	U5MR≥90 and ≥50,000	78.5	4.5	82387	35	480	4.3	4800	no
Myanmar	2005	U5MR≥90 and ≥50,000	50	3.2	46284	52.5	200	4.5	1900	yes
Nepal	2005	U5MR≥90 and ≥50,000	35.8	5.5	19900	61.6	190	6.0	1100	yes
Niger	2005	U5MR≥90 and ≥50,000	95.5	4.9	87967	29	630	2.0	5600	no
Nigeria	2005	U5MR≥90 and ≥50,000	108.8	2.7	750111	32	560	3.1	40000	no
Pakistan	2005	U5MR≥90 and ≥50,000	81.1	2.1	431568	56.7	170	3.6	7900	yes
Papua New Guinea	2005	U5MR≥90	57.3	1.8	11963	42.9	220	3.3	460	yes
Peru	2008	MMR>200 + ≥750 maternal deaths	16.9	6.2	10483	48.7	89	4.4	530	yes
Philippines	2005	≥50,000 child deaths	28	2.9	65613	45.1	120	-0.6	3000	no
Rwanda	2005	U5MR≥90 and ≥50,000	41.7	5.2	14207	44.3	320	6.1	1300	yes
Sao Tome and Principe	2011	low income	47.3	3.4	297	36.7	210	2.8	14	
Senegal	2005	U5MR≥90 and ≥50,000	47.2	4.4	27059	44.6	320	2.2	1700	yes
Sierra Leone	2005	U5MR≥90 and ≥50,000	120.4	3.1	26466	28.8	1100	3.3	2400	no
Solomon Islands	2011	low income	28.1	1.4	470	43.2	130	3.8	23	
Somalia	2005	U5MR≥90 and ≥50,000	136.8	1.1	60537	29.2	850	1.8	3900	no
South Africa	2005	≥50,000 child deaths	40.5	1.6	41930	26.6	140	0.4	1500	yes
Sudan	2005	U5MR≥90 and ≥50,000	70.1	2.4	89488	43.1	360	3.8	4600	no
S Sudan	2012	new, high burden country	92.6	4	39487	43.1	730	3.0	3000	no
Swaziland*	2005	U5MR≥90	60.7	0.8	2221	23.4	310	2.5	120	yes
Tajikistan	2005	U5MR≥90	44.8	3.5	11799	46.6	44	1.9	120	yes
Tanzania, United Republic of	2005	U5MR≥90 and ≥50,000	48.7	4.9	98180	39.3	410	3.5	7900	no
Togo	2005	U5MR≥90	78.4	2.5	19512	34.3	450	1.6	1100	yes

Turkmenistan	2005	U5MR≥90	51.4	2.3	5868	44	61	0.3	68	yes
Uganda	2005	U5MR≥90 and ≥50,000	54.6	4.9	85291	34.9	360	3.2	5900	no
Uzbekistan	2011	low income	39.1	2.4	26205	52.1	36	2.6	220	
Viet Nam	2011	low income	21.7	3.4	34191	52.4	49	4.4	690	
Yemen	2005	U5MR≥90 and ≥50,000	41.9	4.4	34351	53.1	270	2.3	2100	yes
Zambia	2005	U5MR≥90 and ≥50,000	64	4.4	38990	33.8	280	3.1	1800	yes
Zimbabwe	2005	U5MR≥90 and ≥50,000	70.7	0.3	38087	33.7	470	0.4	2100	yes

Note: 2004 is the baseline year during which Countdown assessed countries for monitoring on the basis of U5MR (U5MR≥90 or ≥50,000 deaths). In 2008, eight additional countries were included on the basis of MMR using 2005 data (MMR>550 or MMR>200 and ≥750 maternal deaths). In 2012, six additional low income countries were included to reconcile the Countdown and Global Strategy country lists: Comoros, Kyrgyzstan, Sao Tome and Principe, Solomon Islands. The 2005 report includes the U5MR but not the absolute number of deaths.

Table 2: Changes in national coverage of Countdown interventions from 2000-2008 to 2009-2014 (%) using most recent data in each period, and by percentage of the gap to 100% coverage closed between the two periods.

Indicator for which trend data are available	No. of countries with data	Median coverage (%)		Change (ppts)	Proportion of gap closed
		2000-2008	2009-2014		
Hib3 immunization ^A	13	84	95	11	69
DTP3 immunization ^A	74	77	88	11	47
First dose Measles immunization ^A	71	76	85	9	38
First line antimalarial treatment ^B	21	8	43	35	38
Antenatal care (≥ 1 visit)	63	85	90	6	36
Children sleeping under ITNs ^B	38	16	40	24	29
Vitamin A supplementation (2 doses)	47	86	90	4	29
Improved drinking water sources ^C	73	73	79	6	22
Demand for fam. planning satisfied	43	54	64	10	21
Skilled attendant at delivery	66	55	65	9	21
IPTp for malaria during pregnancy ^B	26	7	25	18	19
Exclusive breastfeeding (<6 months)	58	33	41	9	13
Careseeking for pneumonia	57	48	54	6	12
Antenatal care (4+ visits)	44	50	56	6	12
Oral rehydration salts (ORS) treatment	58	30	38	8	11
Oral rehydration therapy (ORT) with continued feeding	49	42	48	6	10
Improved sanitation facilities ^C	73	38	42	4	6
Zinc	6	2	2	1	1

Note: If more than one survey was conducted in a period, the most recent was used. These two tables include only indicators for which trend data are available in the data sets shared by UNICEF to date.

^A 2004 data used for first time period and 2012 data used for second time period.

^B Analysis includes countries where at least 75% of the population is at risk of malaria and where a substantial proportion (50% or more) of malaria cases is due to *Plasmodium falciparum* (n = 44) or where 50–74% of the population is at risk of malaria and where a substantial proportion (50% or more) of malaria cases is due to *P. falciparum* (n = 8).

^C Includes 2015 data.

Source: Immunization rates, WHO and UNICEF; postnatal visit for mother, Saving Newborn Lives analysis of Demographic and Health Surveys; improved water and sanitation, WHO and UNICEF Joint Monitoring Programme for Water Supply and Sanitation; all other indicators, UNICEF global database, July 2015, based on Demographic and Health Surveys, Multiple Indicator Cluster Surveys and other national surveys; does not include 2014-2015 Rwanda data.

REFERENCES

1. Claeson M, Gillespie D, Mshinda H, Troedsson H, Victora CG. Knowledge into action for child survival. *Lancet*. 2003; **362**(9380): 323-7.
2. Countdown to 2015. Tracking progress in child survival: the 2005 report. New York: Unicef; 2005.
3. Countdown to 2015. Tracking progress in maternal, neonatal and child survival: The 2008 report. New York: UNICEF; 2008.
4. Countdown to 2015. Countdown to 2015 decade report (2000–2010): taking stock of maternal, newborn and child survival. New York: Unicef; 2010.
5. Countdown to 2015. Building a future for women and children: the 2012 report. Washington, DC: World Health Organization and UNICEF; 2012.
6. Countdown to 2015. Countdown to 2015: accountability for maternal, newborn and child survival, the 2013 update. Geneva: WHO; 2013.
7. Countdown to 2015. Countdown to 2015: fulfilling the health agenda for women and children. Geneva: WHO; 2014.
8. Saad-Haddad G, DeJong J, Terreri N, et al. Patterns and Determinants of Antenatal Care Utilization: Analysis of National Survey Data in Seven Countdown Countries. (in press).
9. Independent Expert Review Group. iERG report; 2015.
10. Black RE AL, Bhutta ZA, Caulfield L, de Onis M, Ezzati et al. . maternal and child undernutrition: global and regional exposures and health consequences. *Lancet*. 2008; **371**(9608): 243-60.
11. Ronsmans C, Graham WJ, on behalf of the Lancet Maternal Survival Series steering group. Maternal mortality: who, when, where and why. *Lancet*. 2006; **368**(9542): 1189-200.
12. Lawn JE CS, Zupan J for the Lancet Neonatal Survival Steering Team. 4 million neonatal deaths: When? Where? Why? *Lancet*. 2005; **365**(9462): 891-900.
13. Requejo J, Victora C, Bryce J, Scientific Review Group of Countdown t. Data resource profile: countdown to 2015: maternal, newborn and child survival. *Int J Epidemiol*. 2014; **43**(2): 586-96.
14. Requejo JH, Bryce J, Barros AJ, Berman P, Bhutta Z, Chopra M, et al. Countdown to 2015 and beyond: fulfilling the health agenda for women and children. *Lancet*. 2015; **385**(9966): 466-76.
15. Boerma JT, Bryce J, Kinu Y, Axelson H, Victora CG. Mind the gap: equity and trends in coverage of maternal, newborn, and child health services in 54 Countdown countries. *Lancet*. 2008; **371**(9620): 1259-67.
16. Barros A, Victora C. Measuring coverage in maternal and child health: determining and interpreting inequalities in coverage of maternal, newborn and child health interventions. *PLoS Medicine*. 2013; (in press).
17. Maternal mortality estimation inter-agency group (World Health Organization UNCSF, United Nations Population Fund, United Nations Population Division and World Bank). Trends in maternal mortality: 1990 to 2013. . Geneva: World Health Organization; 2014.
18. Say L, Chou D, Gemmill A, Tunçalp O, Moller AB, Daniels J, et al. Global causes of maternal death: a WHO systematic analysis. *Lancet Glob Health*. 2014; **2**(6): e323-33.
19. Ahmed S, Q. Li, L. Liu, and others. Maternal deaths averted by contraceptive use: An analysis of 172 countries. *Lancet*. 2012; **380**(9837): 111-25.
20. Dean SV LZ, Imam AM, Bhutta ZA. Preconception care: closing the gap in the continuum of care to accelerate improvements in maternal, newborn and child health. *Reprod Health*. 2014; **11 Suppl 3**:S.

21. Mason E C-MV, Baltag V, Christiansen C, Lassi ZS, Bhutta ZA. Preconception care: advancing from 'important to do and can be done' to 'is being done and is making a difference'. *Reprod Health*. 2014; **11 Suppl 3:S8**
22. Lawn JE, Blencowe H, Pattinson R, Cousens S, Kumar R, Ibiebele I, et al. Stillbirths: Where? When? Why? How to make the data count? *Lancet*. 2011; **377**(9775): 1448-63.
23. You D, Hug L, Ejdemyr S, et al. Global, regional, and national levels and trends in under-5 mortality between 1990 and 2015, with scenario-based projections to 2030: a systematic analysis by the UN Inter-agency Group for Child Mortality Estimation. . *Lancet*. 2015.
24. Black RE, Morris SS, Bryce J. Where and why are 10 million children dying every year? *Lancet*. 2003; **361**(9376): 2226-34.
25. Liu L, Oza S, Hogan D, Perin J, Rudan I, Lawn JE, et al. Global, regional, and national causes of child mortality in 2000-13, with projections to inform post-2015 priorities: an updated systematic analysis. *Lancet*. 2015; **385**(9966): 430-40.
26. Black RE, Victora CG, Walker SP, Bhutta ZA, Christian P, de Onis M, et al. Maternal and child undernutrition and overweight in low-income and middle-income countries. *Lancet*. 2013; **382**(9890): 427-51.
27. World Nutrition Report; 2014.
28. Kassebaum NJ, Bertozzi-Villa A, Coggeshall MS, Shackelford KA, Steiner C, Heuton KR, et al. Global, regional, and national levels and causes of maternal mortality during 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *The Lancet*. **384**(9947): 980-1004.
29. Wang H, Liddell CA, Coates MM, Mooney MD, Levitz CE, Schumacher AE, et al. Global, regional, and national levels of neonatal, infant, and under-5 mortality during 1990-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet*. 2014; **384**(9947): 957-79.
30. Hazir T, Begum K, El Arifeen S, Khan AM, Huque MH, Kazmi N, et al. Measuring coverage in MNCH: a prospective validation study in Pakistan and Bangladesh on measuring correct treatment of childhood pneumonia. *PLoS Med*. 2013; **10**(5): e1001422.
31. Requejo JH NH, Bryc J. Measuring coverage in MNCH: challenges and opportunities in the selection of coverage indicators for global monitoring. *PLoS Med*. 2013; **10**(5): e1001416.
32. Fund UNsC. Beyond Averages: Learning from the MDGs. New York: UNICEF; 2015.
33. Walker N, Yenokyan G, Friberg IK, Bryce J. Patterns in coverage of maternal, newborn, and child health interventions: projections of neonatal and under-5 mortality to 2035. *Lancet*. 2013; **382**(9897): 1029-38.
34. Victora CG, Wagstaff A, Schellenberg JA, Gwatkin D, Claeson M, Habicht JP. Applying an equity lens to child health and mortality: more of the same is not enough. *Lancet*. 2003; **362**(9379): 233-41.
35. Barros AJ, Ronsmans C, Axelson H, Loaiza E, Bertoldi AD, Franca GV, et al. Equity in maternal, newborn, and child health interventions in Countdown to 2015: a retrospective review of survey data from 54 countries. *Lancet*. 2012; **379**(9822): 1225-33.
36. Victora CG, Barros AJ, Axelson H, Bhutta ZA, Chopra M, Franca GV, et al. How changes in coverage affect equity in maternal and child health interventions in 35 Countdown to 2015 countries: an analysis of national surveys. *Lancet*. 2012; **380**: 1149-56.
37. Organization WH. Everybody's business: Strengthening health systems to improve health outcomes. Geneva: WHO; 2007.
38. Partnership for Maternal NaCH, WHO. A multisectoral policy compendium for RMNCH. Geneva: The Partnership for Maternal, Newborn and Child Health; 2014.
39. Organization GHWAaWH. A Universal Truth: No Health Without a Workforce. Geneva: Global Health Workforce Alliance and World Health Organization; 2013.
40. Waage J, Banerji R, Campbell O, Chirwa E, Collender G, Dieltiens V, et al. The Millennium Development Goals: a cross-sectoral analysis and principles for goal setting after

- 2015 Lancet and London International Development Centre Commission. Lancet. 2010; **376**(9745): 991-1023.
41. Arregoces L DF, Pitt C et al. . Countdown to 2015:changes in official development assistance to reproductive, maternal, newborn and child health, an assessment of progress between 2003 and 2012. Lancet Global Health. 2015; **3**: e410-21.
 42. Huicho L., Segura E., Niño de Guzman J., et al. Placing child health and nutrition at the centre of the antipoverty political agenda: the Peruvian experience, 2000-2013. 2015.
 43. Dieleman JL, Graves C, Johnson E, Templin T, Birger M, Hamavid H, et al. Sources and Focus of Health Development Assistance, 1990-2014. JAMA. 2015; **313**(23): 2359-68.
 44. Desalegn H, Solberg E, Kim JY. The Global Financing Facility: country investments for every woman, adolescent and child. Lancet. 2015; **386**: 105-6.
 45. Jamison DT SL, Alleyne G, Arrow K, Berkley S, Binagwaho A, Bustreo F, Evans D, Feachem R, Frenk J, Ghosh G, Goldie SJ, Guo Y, Gupta S, Horton R, Kruk M, Mahmoud A, Mohohlo LK, Ncube M, Pablos-Mendez A, Reddy K S, Saxenian H, Soucat A, Ulltveit-Moe K, Yarney G. Global Health 2035: a world converging within a generation Lancet. 2015; **382**(9908): 1898-955.
 46. Godwin P, Misra S. Report of the external review of the accountability work for women's and children's health,. New York: United Nations (Every Woman Every Child); 2014.
 47. Commission on information and accountability for Women's and Children's Health. Keeping promises, measuring results. Final report of the Commission. Geneva: World Health Organization; 2011.
 48. Beaglehole R, Bonita R, Ezzati M, Alleyne G, Dain K, Kishore SP, et al. NCD Countdown 2025: accountability for the 25 × 25 NCD mortality reduction target. The Lancet. **384**(9938): 105-7.
 49. El Arifeen S, Hill K, Ahsan KZ, Jamil K, Nahar Q, Streatfield PK. Maternal mortality in Bangladesh: a Countdown to 2015 country case study. Lancet. 2014; **384**(9951): 1366-74.
 50. Abouzahr C. New estimates of maternal mortality and how to interpret them: choice or confusion? Reprod Health Matters. 2011; **19**(37): 117-28.
 51. Handley K, Boerma T, Victora C, Evans TG. An inflection point for country health data. Lancet Glob Health. 2015.
 52. Stokstad E. United Nations. Sustainable goals from U.N. under fire. Science. 2015; **347**(6223): 702-3.
 53. Editorial. The 169 commandments. The Economist. 2015; (March 28).
 54. World Health Organization. Commission on Social Determinants of Health. Closing the gap in a generation: Health equity through action on the social determinants of health. Geneva: WHO; 2008.
 55. Schmidt H, Gostin LO, Emanuel EJ. Public health, universal health coverage, and Sustainable Development Goals: can they coexist? Lancet. 2015; **386**(9996): 928-30.