1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19

Genes, Epigenetics and miRNA Regulation in the Placenta

Daniel Vaiman

Institut Cochin, INSERM U1016, CNRS UMR8104, Université Paris-Descartes
24, rue du Faubourg St-Jacques
75014, Paris, FRANCE

Correspondence;
Daniel Vaiman
Tel; 33 1 44412301
Fax; 33 1 44412302
Email; daniel.vaiman@inserm.fr

Key words; placenta, epigenetics, preeclampsia, micro-RNA, DNA methylation
15 pages
3480 words

l.20 Abstract.

21 This text reviews briefly the context in which epigenetics regulate gene expression in trophoblast
22 development and function. It is an attempt to focus on a limited number of recent papers that,
23 according to the author, shed new light on placental development, and constitute possible trails for
24 improving knowledge and women follow-up in pathological pregnancies.

25

II26 Introduction.

27

28 One of the earliest events of embryo differentiation in Eutherian mammals is when two major
 29 lineages differentiate into two different types of stem cells, embryonic stem cells (ESCs), on the one
 30 hand and trophoblast stem cells (TSCs), on the other hand. ESCs will conduct to the development of
 31 the different tissues of the embryo, while TSC cells are at the origin of the placenta. Clearly, early
 32 defects in the development of the placenta are susceptible to lead to functional defects, either by
 33 altering the materno-foetal immunological dialogue, the placental structure, or the placental
 34 function. These alterations will later be the cause of placental disorders, very common in humans,
 35 such as preeclampsia or Intra-Uterine Growth Restriction. Gene expression defects are considered as
 36 major culprits at the base of such diseases. Epigenetic mechanisms, that regulate gene expression
 37 without altering the DNA sequence, are now increasingly thought are important actors in the onset
 38 and severity of placental diseases in humans. This text will shortly recapitulate the major genes
 39 involved in trophoblast differentiation, then it will present briefly the different epigenetic
 40 machineries active in the cell; finally, the function of gene methylation and regulation of gene
 41 expression through the action of small RNAs will be exposed.

II42 Genes of trophoblast differentiation

431. Embryonic versus Trophoblast stem cells

44 The blastocyst stage is the first visible stage following fertilization where two morphologically
 45 different types of cells are evident. The periphery of the blastocyst is made of trophectoderm cells,
 46 and a cellular mass (inner cell mass, ICM) is located at one side in the blastocyste cavity. The
 47 trophectoderm cells near the ICM are the 'polar trophectoderm', while the opposite side is the
 48 'mural trophectoderm'. In mice, the first gene associated with trophectoderm development, the
 49 transcription factor *Cdx2* (*Drosophila* caudal-type homeobox 2) was identified in 2005 [1]. In the
 50 study, the authors demonstrate in mice the existence of a balance between *Oct3/Oct4* and *Cdx2*, the
 51 former being positioned very high in the hierarchy of pluripotency, and able to block the expression
 52 and activity of the later, the reciprocal repression being also present. Besides *Cdx2*, another
 53 transcription factor belonging to the T-box family of proteins, *Eomesodermin* (Eomes) is requested
 54 for trophoblast differentiation and proliferation (as well as for mesoderm formation) [2]. In 2007,
 55 Yagi and coworkers identified the transcription factor *Tead4* (TEA-domain family member 4) as a
 56 trophectoderm determinant acting upstream of *Cdx2* [3]. When a mouse embryo develops without
 57 *Tead4*, it dies before implantation after failure to form the blastocoele (the cavity filling the
 58 blastocyst).

592. The network of early placental genes

60 Most knowledge on early development originates from the study of mouse embryos and
 61 depends on transcription factors, most of them being highly conserved in Eutherian mammals,
 62 suggesting that at least partly, what has been discovered in mice may be widely applicable to humans
 63 despite possible differences in the gene cascades involved [4]. In mice, mural trophectoderm
 64 differentiate into primary trophoblast giant cells (TGCs). Other trophoblast giant cells (secondary
 65 TGCs) emerge from the polar trophectoderm sometimes through a transient spongiotrophoblast
 66 stage of differentiation. TGCs will constitute an interface with the maternal deciduas and will become
 67 polyploid through endoreplication, under the control of the polyploidy-regulating gene *Geminin* [5].
 68 Each of these placenta cell lineages depends on a specific subset of genes as summarized by Maltepe
 69 and coworkers in a 2010 review [6]. Amongst those, it is important to note the transcription factor
 70 *GCM1*, a major gene of the cytotrophoblasts lineage, leading to the development of the placental

syncytiotrophoblast, which allows the utero-placental exchanges in mice. The complex *HIF1-ARNT*, which is the major sensor of hypoxia in mammalian cells, plays also an essential role in spongiotrophoblast development (the cells composing the junctional zone of the mouse placenta, which is the closest to the maternal structure). Knocking-out *HIF1* in mice leads to the death of 10% of the embryos between days E6.5 and E7.5 [7]. These mice are not able to form extra-embryonic and embryonic cavities. To note as well, the importance of imprinted genes at least at later stages of development and placental growth, these genes playing a crucial role in adjusting the trade-off between mother and fetus [8]. In the recent years, in addition to the identification of crucial genes for trophoblast development, the importance of epigenetic marks has been revealed, with resistant methylated genes that prevent transdifferentiation of ES cells into TS cells [9], and the demonstration of the existence of genes able to modify the epigenetic landscape and playing a crucial role in trophoblast differentiation, such as *PRDM14* and *ARID3a*, acting respectively on DNA demethylation, and Histone acetylation [10, 11].

Genetics versus epigenetics of trophoblast differentiation

1. Definition of epigenetics

Epigenetics may be defined as the study of mechanisms regulating gene expression, that are heritable through cell division, but not involving changes in the nucleotide sequence. From an operational point of view, epigenetics works through the action of a battery of cellular machineries that regulate genome function. Epigenetics is involved in various processes of development, physiological homeostasis, and adaption to external stresses (be them physical, chemical or of biochemical/biological origins). The epigenetic marks that tag the genes as expressed or repressed are inheritable through cell divisions, such as in the mitosis processes, but also in a limited number of cases, through the meiosis process. This is the basis of 'transgenerational inheritance', when without exposure, an epigenetic mark is transmitted more than three generations after the exposure. For instance, exposure of female rats to the endocrine disrupter vinclozolin, leads to anomalies of sperm number and function (apoptosis, movement) even up to the fourth generation [12]. Mechanisms for this type of transmission remain elusive, albeit it has recently been shown that some histone marks survive the multiple erasure mechanisms of male meiosis [13] and could carry an epigenetic message; another possible explanation is given by the stock of stable mi-RNA that are present in the cells and especially in the sperm [14]. These miRNAs, may be stabilized by a specific methylation by Dnmt2 (*Trdm1*), since Dnmt2 invalidated mice apparently lose the ability to transmit at least some epigenetic memories in the case of paramutations [15].

2. The epigenetic machinery

Epigenetic marks are either apposed on the chromatin (on histones, or directly on the DNA), while epigenetic modulators may be stored in the cytoplasm of the cells such as miRNAs. It is not possible here to describe all the elements of the epigenetic machinery, and the reader can analyze numerous excellent reviews on these subjects. Among these reviews, the recent paper by Zhang and Pradhan [16], gives a comprehensive and recent vision of the various mechanisms, from DNA methylation, hydroxymethylation and demethylation to the list of enzymes able to post-translationally modify histones.

A highly simplified list of these different actors is presented as figure 1.

114 Normal epigenetic marks in the placenta

115 The actual profiling of histone marks in the placenta is not well known today, possibly because
 116 histone marks on chromatin may not be very stable. However, a recent paper analyzed extensively
 117 the placental methylome by an approach combining the use of the 450K Illumina methylation array
 118 and MethylC seq approaches [17]. A recent review listing the different technical approaches available
 119 for quantifying DNA methylation is given in Calicchio et al (2013), [18].

1201. The epigenetic status of the normal placenta

121a. The placental methylome

122 Many papers deal with placental DNA methylation, and it is not possible to analyze
 123 exhaustively the literature in this paper. Many recent reviews can be consulted on the subject such
 124 as for instance [19]. Amongst the most recent global studies, the paper of Schroeder and co-workers
 125 appear as a landmark in the field, since it compares systematically results from the Illumina 450K
 126 methylation array and MethylC-seq, in parallel with evaluating gene expression by RNA_seq [17]..
 127 The authors first show that the Illumina and MethylC-seq approaches resulted in a very similar
 128 description of the methylation profiles along the chromosomes. Another important and innovative
 129 result was the demonstration that the methylation profile is stable in the placenta during the three
 130 trimesters of pregnancy. This was unsuspected, since from an expression point of view, placentas
 131 from first, second and third trimester are quite different [20]. Before publication of the Schroeder
 132 paper, it was known that placental methylation is lower on average than in other tissues (see for
 133 instance Table 1 in [21]). The Schroeder and coworker 2013 paper revealed that the methylation is
 134 distributed in specific domains, and some of them called 'Partially Methylated Domains' are absent
 135 from adult tissues.

136b. Links between DNA methylation and gene expression

137 More and more studies tend to negate the idea that high methylation is associated simply with
 138 gene repression. The Schroeder study showed that high methylation in the gene body is rather
 139 associated to high expression levels, while the methylation inside gene promoters can be associated
 140 either to repression or increase of gene expression. To note, however, most studies attempt to
 141 correlate methylation analyses with transcriptome data. Classical transcriptome analysis measures
 142 the steady-state of mRNAs that are present in the cell at a specific moment. Therefore, the more
 143 stable mRNA may have a low expression and have a high concentration inside the cell. By contrast,
 144 short-lived mRNA will not accumulate inside the cell even if they have a very active transcription [22].
 145 This could more accurately be tackled by translatoome approaches that prior to microarray analysis,
 146 purifies polysomes with their associated mRNA being translated, and thus be indicative of the genes
 147 that are translated in proteins at a given moment [23]. To have an instant vision of the expression
 148 level of all the mRNAs is much more challenging, and may be approached rather by ChIP-seq
 149 approaches using antibodies against RNA polymerase II. For the moment such methods have not
 150 been applied to the placenta. Nevertheless, there is a clear association of the expression of several
 151 developmental genes (such as *DLX5/DLX6*) and specific methylation patterns in the placenta [17]. The
 152 mechanistic link between expression and methylation at the gene level is not very frequent in the
 153 scientific literature.

154 Abnormal environmental conditions during development may alter gene methylation. Among
 155 examples, prenatal stress imposed to mothers has been shown to modulate the methylation of the
 156 *Hmox1* promoter [24]. Preeclampsia (cause or consequence) is associated with alterations of the
 157 methylation of various genes, such as *TIMP3* [25], *RassF1* and *SERPINB5* [26], *SERPINA3* [27, 28]
 158 *Cullin7* and *Cullin 4b* [29], *11βHSD* [30], *HLA-G* [31], and several imprinted genes [32]. In 2012, it has

been shown that *SERPINA3* expression is regulated by the transcription factor *ZBTB7B*, depending upon the state of methylation of the *SERPINA3* promoter. *ZBTB7B* is also involved in regulation of extracellular matrix genes [33]. The ZBTB family of genes has been associated to highly expressed/highly methylated gene promoters [34, 35]. Overall, high-methylation in a gene is classically associated with a low level of transcription. However, many recent results show that exceptions to this paradigm are numerous.

Regulation of gene expression by microRNA

1. Definition of miRNAs, number in the mammalian genome

MiRNAs are small RNA molecules (~22 nucleotides) synthesized from the genome by the regular cell machinery either from bona fide genes (and transcribed in pri-mi-RNA), or from introns of host genes. Drosha and Pasha (DGCR8) proteins are involved in the handling of the pri-miRNA and its transformation into pre-miRNA. These are exported towards the cytoplasm through the action of Exportin 5, then the Dicer complex will eliminate the loop, and the dimer will be associated to the RISC complex (RNA-induced Silencing Complex), composed of proteins of the Argonaute family, and of a helicase able to dissociate the two RNA strands. The one that is kept in the RISC complex will hybridize to target mRNA inducing degradation or blocking protein synthesis. The processes are summarized in Treiber (2012) [36].

2. MiRNAs functions in trophoblast

In trophoblast cells, as in other context, there is an increasing corpus of data showing that miRNAs play important physiological and developmental roles, consistently with the estimation that they potentially regulate 60% of the genes in eukaryotic genomes [37]. Actually, for several physiological characteristics of trophoblast biology (immunity, invasion, proliferation, vasodilation), it has been shown that specific miRNAs are effective actors. An important issue for trophoblast tissue and especially in humans, is the existence of placental miRNAs that are specific of primates and mainly organized into two clusters, one on chromosome 19 (54 miRNAs) and one on chromosome 14 (34 miRNAs) [38, 39].

a. immunity

Trophoblasts express specific HLA molecules that differ from the ones of other cells of the body. Instead of HLA A and B, which are highly polymorphic, they express HLA-C and HLA-G (the former being moderately polymorphic, while the latter is quasi-monomorphic [40]). It has recently been shown that HLA-G expression is down-regulated by miR-148a and miR-152 [41, 42]. A very interesting recent study suggests a miRNA basis for the communication between trophoblast cells and uterine Natural Killer (uNK), the major actors of the early materno-foetal immunological dialogue. The authors showed that miR-517a-3p down-regulate PRKG1 (Protein Kinase, CGMP-Dependent, Regulatory, Type I) in Bewo cells (a choriocarcinoma cell line similar to villous trophoblast and able to fuse in culture under AMPc induction by forskolin). Exosomes from the Bewo cells were able to fuse with maternal NK cells, and the content in miRNAs induced a decrease of PRKG1 in the NK cells [43]. PRKG1 is considered as an important modulator of nitric oxide (NO) metabolism, NO being a major molecule in the regulation of the vascular system function [44]. Therefore trophoblast cells appear able to communicate with the maternal immune cells through miRNAs production, and possibly to modulate their function.

b. invasion

Trophoblast invasion is also affected by several miRNAs, often through the alteration of the Nodal pathway. Nodal is a member of the TGF β family that interacts with the ALK7 receptor (Activin receptor-like kinase 7). Both are expressed in villous and extra-villous trophoblast, and their overexpression decreases cell migration and invasion, while silencing these genes has the opposite effect [45]. Several miRNAs target genes of the Nodal pathway (miR-373a-5p, miR-675, miR-195, miR-376c), and may have a specific impact on trophoblast physiology [46-49]. Other regulators of invasion are targeted by miRNAs (*SMAD2* by miR-18a, *FOXA1* by miR-20a, *S1PR* by miR125b-1-3p, *CXCL12* by miR135b, *cMYC* by miR-34a, *MMP9* by miR-204, [50-56]). A recent study using a BAC encompassing the C19MC cluster, the team of Yoel Sadovsky showed that miRNAs encoded by this cluster target genes involved in trophoblast movements, while miR-519d, located inside this cluster, regulates invasion of Extra Villous Trophoblasts by targeting CXCL6, NR4A2 and FOXL2 [57].

c. proliferation

Trophoblast proliferate massively at the first trimester, and in humans, extravillous trophoblast will plug the maternal spiral arteries of the endometrium [58], the local hypoxia being probably important for the proliferation [59]. Again, several genes involved in proliferation are regulated by mi-RNA; this is the case for *LIF*, *CCND1*, *TTN*, *MYC/ERK*, *IRAKM/NKIRAS*, respectively targeted by miR-141, miR-155, miR-144, MiR-145, 377, let7a, and finally miR-155* [60-64].

d. vasodilation

Placentation in humans, especially the specific invasion of maternal arteries aims at providing the growing fetus with a stable hemosphere especially in terms of oxygen pressure. Indeed, while trophoblasts are well adapted to low oxygen pressure, they are very sensitive to steep variations in oxygen pressure, leading many researchers to mimic preeclampsia through a hypoxia-reoxygenation protocols, rather than a mere exposure to hypoxia [65-68]. These variations are associated to an increase in oxidative and nitrosative stress that are major actors in preeclampsia [69, 70]. In eukaryotic cells, it is clear that miR-210 plays the role of an important sensor of hypoxia, and its deregulation has been recurrently reported, see for instance [71-76]. Recently in a mouse model of preeclampsia, we also reported abnormal miR-210 regulation [77]. Amongst the various molecules important for vasodilation, NO and H₂S play prominent roles. It has recently been shown that miR-21 down-regulate CSE (Cystathione lyase), a protein involved in H₂S synthesis [78].

3. Defining miRNA targets, some examples

Compared to other epigenetic regulations, the mode of action of miRNAs is much more understood in molecular terms. In addition, it is possible to define specifically and experimentally if a given gene is a direct target for a miRNA. The idea is to clone the presumed miRNA target of the gene (often contained into the 3' Untranslated Region) in front of the luciferase coding sequence in a reporter vector. Then cells are transfected with this construction and either siRNA mimicking the miRNA or with a control siRNA. If the miRNA targets the gene, then it will lead to a reduction of the luciferase activity, thus validating the target. Following are some examples in the context of placenta and placental diseases.

a. miR-1324 in preeclampsia

In 2013, Oudejans and coworkers identified polymorphisms on the gene *INO80B* as associated with an increased risk of preeclampsia. Interestingly the susceptible allele (rs34174194) is located in the 3'UTR of the gene and modifies a binding site for the miRNA miR-1324. The common variant is highly conserved throughout mammalian species. By luciferase assays, the authors

demonstrated that the repression of the gene is only marginal in the mutant, thus leading to an increased risk of preeclampsia [79].

b. miR-675 and placental development

Another example is given by miR-675, a micro-RNA which is encoded by a highly conserved sequence present in the imprinted non-coding gene H19 [80]. In mice miR-675 synthesis is prevented during the first 11 days of gestation by binding with the HuR protein, which decreases during pregnancy. Then, miR-675 is released, and targets the *Igf1R* mRNA, encoding the active receptor of Igf2, as well as NMO1 (Nodal Modulator 1).

c. miR-34a and placental diseases

MiR-34 is frequently associated with cancer [81]. It is also strongly expressed in placental cells, especially at the end of pregnancy [82], while the pri-miR appears strongly expressed in early placentas [83]. We could show that its expression is correlated to methylation of its promoter, that this methylation is considerably increased by hypoxia. Functionally, it has been recently shown by Sun and coworkers that miR-34a inhibits cell migration and invasion by directly targeting the c-MYC proto-oncogene [51].

4. Sets of preeclampsia-associated deregulated miRNAs

One of the major interests of miRNAs is that they have a double hat. They regulate directly genes by repressing their expression, and are thus important in physiology and development, but they are also very stable in the plasma (at least several days) [84, 85], and constitute exciting potential biomarkers for a large number of human diseases; among those there are more than 20 articles exploring preeclampsia and circulating miRNAs, for instance [86-92]. For the moment, however, no consensual list of miRNAs emerges from these screening experiments; possibly, a better classification of the patients in term of preeclampsia severity, combination with IUGR, HELLP syndrome, eclampsia, or considering the ethnical background, could be very helpful to classify the various miRNA profiles and sort them more efficiently.

VI. Conclusion

This short review aims at presenting the role of the epigenetic machinery, in the context of trophoblast differentiation and normal/pathological placental development. The histone code is accessible through technically challenging and costly methods, such as in particular ChIP-seq experiments, using antibodies directed against the different histone variants. While these approaches have a great potential for the better understanding of placental development, they appear difficult to set up in the context of placental diseases. DNA methylation is now very easily implemented in humans using high-throughput microarrays that interrogate simultaneously and with a high quantitative precision more than 450,000 CpG in the genome. However, besides the use of demethylated maspin (SERPINB5) in the maternal blood [93], such epigenetic marks are not envisaged as systematically used in the context of placental diseases. An additional problem is that the link between DNA methylation and gene expression down-regulation is far from obvious [35]. Micro-RNA are probably the most promising biomarkers, cumulating the advantage of a good knowledge of their mode of action, high stability in the plasma, and the advantage of being easily detectable in the women plasma. For the moment, no clear subset of miRNAs appears as characteristic of preeclampsia; this may be due to the heterogeneity of the disease, and possibly a better clinical classification of the samples could drive significant progress in the field. This would imply analyzing a much higher of human samples thoroughly characterized, which has not been done today. A more distant additional use of small RNA could be therapeutic, since siRNA maybe used to correct specifically anomalies of gene expression. Attempts to address small vesicles encompassing

290 siRNA or therapeutic proteins towards the syncytiotrophoblast could be an interesting approach in
291 the future.
292

Figures

<ul style="list-style-type: none"> ▶ DNA methylation <ul style="list-style-type: none"> ▶ DNA methyltransferases <ul style="list-style-type: none"> ▶ Maintenance: DNMT1 ▶ De novo methylation: DNMT3A, DNMT3B ▶ Regulatory: DNMT3L ▶ Acting on RNA targets: DNMT2 ▶ Demethylases: <ul style="list-style-type: none"> ▶ TET1, TET2, TET3 ▶ Gadd45 ▶ Targets: CpG islands ▶ Stability ▶ Correlation with gene expression ? 	<ul style="list-style-type: none"> ▶ Histone code <ul style="list-style-type: none"> ▶ >70 post-translational modifications on N-ter amino-acids of H3 and H4 ▶ A complex code <ul style="list-style-type: none"> ▶ H3K4Me3 open ▶ H3K27Me3 closed ▶ H3/H4 Acet open ▶ Numerous enzymes involved <ul style="list-style-type: none"> ▶ HAT ▶ HDAC ▶ Swi-Snf ▶ P300/CEBP ▶ LSD1 ▶ Documented modifications by cellular processes/stresses 	<ul style="list-style-type: none"> □ MiRNA synthesis and regulation <ul style="list-style-type: none"> □ Targeting hundred of genes □ Powerful identification tools now available □ Quantification easy by qRT-PCR □ Biological targets can be identified □ Molecular mechanisms at least partially elucidated <ul style="list-style-type: none"> ■ Drosha ■ Dicer ■ RISC complex ■ Argonaute proteins
--	--	--

Figure 1: A short and non-exhaustive list of the mechanisms and genes involved in modulating the epigenetic anlagen in mammalian cells. To this table could be added genes involved in hydroxymethylation, an issue that has not been studied in the context of trophoblast cells.

References

- [1] Niwa H, Toyooka Y, Shimosato D, Strumpf D, Takahashi K, Yagi R and Rossant J. Interaction between Oct3/4 and Cdx2 determines trophectoderm differentiation. *Cell*. 2005;123(5):917-29.
- [2] Russ AP, Wattler S, Colledge WH, Aparicio SA, Carlton MB, Pearce JJ, Barton SC, Surani MA, Ryan K, Nehls MC, Wilson V and Evans MJ. Eomesodermin is required for mouse trophoblast development and mesoderm formation. *Nature*. 2000;404(6773):95-9.
- [3] Yagi R, Kohn MJ, Karavanova I, Kaneko KJ, Vullhorst D, DePamphilis ML and Buonanno A. Transcription factor TEAD4 specifies the trophectoderm lineage at the beginning of mammalian development. *Development*. 2007;134(21):3827-36.
- [4] Niakan KK and Eggan K. Analysis of human embryos from zygote to blastocyst reveals distinct gene expression patterns relative to the mouse. *Dev Biol*. 2013;375(1):54-64.
- [5] Gonzalez MA, Tachibana KE, Adams DJ, van der Weyden L, Hemberger M, Coleman N, Bradley A and Laskey RA. Geminin is essential to prevent endoreduplication and to form pluripotent cells during mammalian development. *Genes Dev*. 2006;20(14):1880-4.
- [6] Maltepe E, Bakardjiev AI and Fisher SJ. The placenta: transcriptional, epigenetic, and physiological integration during development. *J Clin Invest*. 2010;120(4):1016-25.
- [7] Covello KL, Kehler J, Yu H, Gordan JD, Arsham AM, Hu CJ, Labosky PA, Simon MC and Keith B. HIF-2 α regulates Oct-4: effects of hypoxia on stem cell function, embryonic development, and tumor growth. *Genes Dev*. 2006;20(5):557-70.
- [8] Constancia M, Kelsey G and Reik W. Resourceful imprinting. *Nature*. 2004;432(7013):53-7.
- [9] Cambuli F, Murray A, Dean W, Dudzinska D, Krueger F, Andrews S, Senner CE, Cook SJ and Hemberger M. Epigenetic memory of the first cell fate decision prevents complete ES cell reprogramming into trophoblast. *Nat Commun*. 2014;5:5538.
- [10] Burton A, Muller J, Tu S, Padilla-Longoria P, Guccione E and Torres-Padilla ME. Single-cell profiling of epigenetic modifiers identifies PRDM14 as an inducer of cell fate in the mammalian embryo. *Cell Rep*. 2013;5(3):687-701.
- [11] Rhee C, Lee BK, Beck S, Anjum A, Cook KR, Popowski M, Tucker HO and Kim J. Arid3a is essential to execution of the first cell fate decision via direct embryonic and extraembryonic transcriptional regulation. *Genes Dev*. 2014;28(20):2219-32.
- [12] Anway MD, Cupp AS, Uzumcu M and Skinner MK. Epigenetic transgenerational actions of endocrine disruptors and male fertility. *Science*. 2005;308(5727):1466-9.
- [13] van de Werken C, van der Heijden GW, Eleveld C, Teeuwssen M, Albert M, Baarends WM, Laven JS, Peters AH and Baart EB. Paternal heterochromatin formation in human embryos is H3K9/HP1 directed and primed by sperm-derived histone modifications. *Nat Commun*. 2015;5:5868.
- [14] Rassoulzadegan M, Grandjean V, Gounon P, Vincent S, Gillot I and Cuzin F. RNA-mediated non-mendelian inheritance of an epigenetic change in the mouse. *Nature*. 2006;441(7092):469-74.
- [15] Kiani J, Grandjean V, Liebers R, Tuorto F, Ghanbarian H, Lyko F, Cuzin F and Rassoulzadegan M. RNA-mediated epigenetic heredity requires the cytosine methyltransferase Dnmt2. *PLoS Genet*. 2013;9(5):e1003498.
- [16] Zhang G and Pradhan S. Mammalian epigenetic mechanisms. *IUBMB Life*. 2014;66(4):240-56.
- [17] Schroeder DI, Blair JD, Lott P, Yu HO, Hong D, Crary F, Ashwood P, Walker C, Korf I, Robinson WP and LaSalle JM. The human placenta methylome. *Proc Natl Acad Sci U S A*. 2013;110(15):6037-42.
- [18] Calicchio R, Doridot L, Miralles F, Mehats C and Vaiman D. DNA methylation, an epigenetic mode of gene expression regulation in reproductive science. *Curr Pharm Des*. 2013;20(11):1726-50.
- [19] Bianco-Miotto T, Mayne BT, Buckberry S, Breen J, Rodriguez Lopez CM and Roberts CT. Recent progress towards understanding the role of DNA methylation in human placental development. *Reproduction*. 2016;152(1):R23-30.
- [20] Winn VD, Haimov-Kochman R, Paquet AC, Yang YJ, Madhusudhan MS, Gormley M, Feng KT, Bernlohr DA, McDonagh S, Pereira L, Sali A and Fisher SJ. Gene expression profiling of the human

- maternal-fetal interface reveals dramatic changes between midgestation and term. *Endocrinology*. 2007;148(3):1059-79.
- [21] Fuke C, Shimabukuro M, Petronis A, Sugimoto J, Oda T, Miura K, Miyazaki T, Ogura C, Okazaki Y and Jinno Y. Age related changes in 5-methylcytosine content in human peripheral leukocytes and placentas: an HPLC-based study. *Ann Hum Genet*. 2004;68(Pt 3):196-204.
- [22] Honkela A, Peltonen J, Topa H, Charapitsa I, Matarese F, Grote K, Stunnenberg HG, Reid G, Lawrence ND and Rattray M. Genome-wide modeling of transcription kinetics reveals patterns of RNA production delays. *Proc Natl Acad Sci U S A*. 2015;112(42):13115-20.
- [23] Zuccotti P and Modelska A. Studying the Translatome with Polysome Profiling. *Methods Mol Biol*. 2016;1358:59-69.
- [24] Solano ME, Kowal MK, O'Rourke GE, Horst AK, Modest K, Plosch T, Barikbin R, Remus CC, Berger RG, Jago C, Ho H, Sass G, Parker VJ, Lydon JP, DeMayo FJ, Hecher K, Karimi K and Arck PC. Progesterone and HMOX-1 promote fetal growth by CD8+ T cell modulation. *J Clin Invest*. 2015;125(4):1726-38.
- [25] Yuen RK, Penaherrera MS, von Dadelszen P, McFadden DE and Robinson WP. DNA methylation profiling of human placentas reveals promoter hypomethylation of multiple genes in early-onset preeclampsia. *Eur J Hum Genet*. 2010;18(9):1006-12.
- [26] Robins JC, Marsit CJ, Padbury JF and Sharma SS. Endocrine disruptors, environmental oxygen, epigenetics and pregnancy. *Front Biosci (Elite Ed)*. 2011;3:690-700.
- [27] Chelbi ST, Mondon F, Jammes H, Buffat C, Mignot TM, Tost J, Busato F, Gut I, Rebourcet R, Laissue P, Tsatsaris V, Goffinet F, Rigourd V, Carbonne B, Ferre F and Vaiman D. Expressional and epigenetic alterations of placental serine protease inhibitors: SERPINA3 is a potential marker of preeclampsia. *Hypertension*. 2007;49(1):76-83.
- [28] Chelbi ST, Wilson ML, Veillard AC, Ingles SA, Zhang J, Mondon F, Gascoin-Lachambre G, Doridot L, Mignot TM, Rebourcet R, Carbonne B, Concordet JP, Barbaux S and Vaiman D. Genetic and epigenetic mechanisms collaborate to control SERPINA3 expression and its association with placental diseases. *Hum Mol Genet*. 2012;21(9):1968-78.
- [29] Gascoin-Lachambre G, Buffat C, Rebourcet R, Chelbi ST, Rigourd V, Mondon F, Mignot TM, Legras E, Simeoni U, Vaiman D and Barbaux S. Cullins in human intra-uterine growth restriction: expressional and epigenetic alterations. *Placenta*. 2010;31(2):151-7.
- [30] Causevic M and Mohaupt M. 11beta-Hydroxysteroid dehydrogenase type 2 in pregnancy and preeclampsia. *Mol Aspects Med*. 2007;28(2):220-6.
- [31] Chen S, Zhao G, Miao H, Tang R, Song Y, Hu Y, Wang Z and Hou Y. MicroRNA-494 inhibits the growth and angiogenesis-regulating potential of mesenchymal stem cells. *FEBS Lett*. 2015;589(6):710-7.
- [32] Bourque DK, Avila L, Penaherrera M, von Dadelszen P and Robinson WP. Decreased placental methylation at the H19/IGF2 imprinting control region is associated with normotensive intrauterine growth restriction but not preeclampsia. *Placenta*. 2010;31(3):197-202.
- [33] Widom RL, Lee JY, Joseph C, Gordon-Froome I and Korn JH. The hckrox gene family regulates multiple extracellular matrix genes. *Matrix Biol*. 2001;20(7):451-62.
- [34] Sasai N, Nakao M and Defossez PA. Sequence-specific recognition of methylated DNA by human zinc-finger proteins. *Nucleic Acids Res*. 2010;38(15):5015-22.
- [35] Spruijt CG and Vermeulen M. DNA methylation: old dog, new tricks? *Nat Struct Mol Biol*. 2014;21(11):949-54.
- [36] Treiber T, Treiber N and Meister G. Regulation of microRNA biogenesis and function. *Thromb Haemost*. 2012;107(4):605-10.
- [37] Friedman RC, Farh KK, Burge CB and Bartel DP. Most mammalian mRNAs are conserved targets of microRNAs. *Genome Res*. 2009;19(1):92-105.
- [38] Donker RB, Mouillet JF, Chu T, Hubel CA, Stolz DB, Morelli AE and Sadovsky Y. The expression profile of C19MC microRNAs in primary human trophoblast cells and exosomes. *Mol Hum Reprod*. 2012;18(8):417-24.

- 401 [39] Morales-Prieto DM, Chaiwangyen W, Ospina-Prieto S, Schneider U, Herrmann J, Gruhn B and
 402 Markert UR. MicroRNA expression profiles of trophoblastic cells. *Placenta*. 2012;33(9):725-34.
- 403 [40] Kirszenbaum M, Moreau P, Gluckman E, Dausset J and Carosella E. An alternatively spliced form
 404 of HLA-G mRNA in human trophoblasts and evidence for the presence of HLA-G transcript in adult
 405 lymphocytes. *Proc Natl Acad Sci U S A*. 1994;91(10):4209-13.
- 406 [41] Zhang C, Li Q, Ren N, Li C, Wang X, Xie M, Gao Z, Pan Z, Zhao C, Ren C and Yang W. Placental miR-
 407 106a approximately 363 cluster is dysregulated in preeclamptic placenta. *Placenta*. 2014;36(2):250-2.
- 408 [42] Manaster I, Goldman-Wohl D, Greenfield C, Nachmani D, Tsukerman P, Hamani Y, Yagel S and
 409 Mandelboim O. MiRNA-mediated control of HLA-G expression and function. *PLoS One*.
 410 2012;7(3):e33395.
- 411 [43] Kambe S, Yoshitake H, Yuge K, Ishida Y, Ali MM, Takizawa T, Kuwata T, Ohkuchi A, Matsubara S,
 412 Suzuki M, Takeshita T and Saito S. Human exosomal placenta-associated miR-517a-3p modulates the
 413 expression of PRKG1 mRNA in Jurkat cells. *Biol Reprod*. 2014;91(5):129.
- 414 [44] Birschmann I and Walter U. Physiology and pathophysiology of vascular signaling controlled by
 415 guanosine 3',5'-cyclic monophosphate-dependent protein kinase. *Acta Biochim Pol*. 2004;51(2):397-
 416 404.
- 417 [45] Nadeem L, Munir S, Fu G, Dunk C, Baczyk D, Caniggia I, Lye S and Peng C. Nodal signals through
 418 activin receptor-like kinase 7 to inhibit trophoblast migration and invasion: implication in the
 419 pathogenesis of preeclampsia. *Am J Pathol*. 2011;178(3):1177-89.
- 420 [46] Gao WL, Liu M, Yang Y, Yang H, Liao Q, Bai Y, Li YX, Li D, Peng C and Wang YL. The imprinted H19
 421 gene regulates human placental trophoblast cell proliferation via encoding miR-675 that targets
 422 Nodal Modulator 1 (NOMO1). *RNA Biol*. 2012;9(7):1002-10.
- 423 [47] Fu G, Ye G, Nadeem L, Ji L, Manchanda T, Wang Y, Zhao Y, Qiao J, Wang YL, Lye S, Yang BB and
 424 Peng C. MicroRNA-376c impairs transforming growth factor-beta and nodal signaling to promote
 425 trophoblast cell proliferation and invasion. *Hypertension*. 2013;61(4):864-72.
- 426 [48] Bai Y, Yang W, Yang HX, Liao Q, Ye G, Fu G, Ji L, Xu P, Wang H, Li YX, Peng C and Wang YL.
 427 Downregulated miR-195 detected in preeclamptic placenta affects trophoblast cell invasion via
 428 modulating ActRIIA expression. *PLoS One*. 2012;7(6):e38875.
- 429 [49] Luo L, Ye G, Nadeem L, Fu G, Yang BB, Honarparvar E, Dunk C, Lye S and Peng C. MicroRNA-378a-
 430 5p promotes trophoblast cell survival, migration and invasion by targeting Nodal. *J Cell Sci*.
 431 2012;125(Pt 13):3124-32.
- 432 [50] Yu Y, Wang L, Liu T and Guan H. MicroRNA-204 suppresses trophoblast-like cell invasion by
 433 targeting matrix metalloproteinase-9. *Biochem Biophys Res Commun*. 2015;463(3):285-91.
- 434 [51] Gu Y, Zhang X, Yang Q, Wang J, He Y, Sun Z and Zhang H. Aberrant Placental Villus Expression of
 435 miR-486-3p and miR-3074-5p in Recurrent Miscarriage Patients and Uterine Expression of These
 436 MicroRNAs during Early Pregnancy in Mice. *Gynecol Obstet Invest*. 2015.
- 437 [52] Pang RT, Leung CO, Ye TM, Liu W, Chiu PC, Lam KK, Lee KF and Yeung WS. MicroRNA-34a
 438 suppresses invasion through downregulation of Notch1 and Jagged1 in cervical carcinoma and
 439 choriocarcinoma cells. *Carcinogenesis*. 2010;31(6):1037-44.
- 440 [53] Tamaru S, Mizuno Y, Tochigi H, Kajihara T, Okazaki Y, Okagaki R, Kamei Y, Ishihara O and Itakura
 441 A. MicroRNA-135b suppresses extravillous trophoblast-derived HTR-8/SVneo cell invasion by directly
 442 down regulating CXCL12 under low oxygen conditions. *Biochem Biophys Res Commun*.
 443 2015;461(2):421-6.
- 444 [54] Anton L, Olarerin-George AO, Hogenesch JB and Elovitz MA. Placental expression of miR-517a/b
 445 and miR-517c contributes to trophoblast dysfunction and preeclampsia. *PLoS One*.
 446 2015;10(3):e0122707.
- 447 [55] Wang Y, Zhang Y, Wang H, Wang J, Pan Z and Luo S. Aberrantly up-regulated miR-20a in pre-
 448 eclampsic placenta compromised the proliferative and invasive behaviors of trophoblast cells by
 449 targeting forkhead box protein A1. *Int J Biol Sci*. 2014;10(9):973-82.
- 450 [56] Xu P, Zhao Y, Liu M, Wang Y, Wang H, Li YX, Zhu X, Yao Y, Qiao J, Ji L and Wang YL. Variations of
 451 microRNAs in human placentas and plasma from preeclamptic pregnancy. *Hypertension*.
 452 2014;63(6):1276-84.

- [57] Xie L, Mouillet JF, Chu T, Parks WT, Sadovsky E, Knofler M and Sadovsky Y. C19MC microRNAs regulate the migration of human trophoblasts. *Endocrinology*. 2014;155(12):4975-85.
- [58] Carbillon L, Challier JC, Alouini S, Uzan M and Uzan S. Uteroplacental circulation development: Doppler assessment and clinical importance. *Placenta*. 2001;22(10):795-9.
- [59] Genbacev O. To proliferate or to divide - to be or not to be. *Early Pregnancy*. 2001;5(1):63-4.
- [60] Morales-Prieto DM, Schleussner E and Markert UR. Reduction in miR-141 is induced by leukemia inhibitory factor and inhibits proliferation in choriocarcinoma cell line JEG-3. *Am J Reprod Immunol*. 2011;66 Suppl 1:57-62.
- [61] Dai Y, Qiu Z, Diao Z, Shen L, Xue P, Sun H and Hu Y. MicroRNA-155 inhibits proliferation and migration of human extravillous trophoblast derived HTR-8/SVneo cells via down-regulating cyclin D1. *Placenta*. 2012;33(10):824-9.
- [62] Liang Y, Lin Q, Luo F, Wu W, Yang T and Wan S. Requirement of miR-144 in CsA induced proliferation and invasion of human trophoblast cells by targeting titin. *J Cell Biochem*. 2014;115(4):690-6.
- [63] Farrokhnia F, Aplin JD, Westwood M and Forbes K. MicroRNA regulation of mitogenic signaling networks in the human placenta. *J Biol Chem*. 2014;289(44):30404-16.
- [64] Xue P, Zheng M, Diao Z, Shen L, Liu M, Gong P, Sun H and Hu Y. miR-155* mediates suppressive effect of PTEN 3'-untranslated region on AP-1/NF-kappaB pathway in HTR-8/SVneo cells. *Placenta*. 2013;34(8):650-6.
- [65] Yang Z, Bai B, Luo X, Xiao X, Liu X, Ding Y, Zhang H, Gao L, Li J and Qi H. Downregulated Kruppel-like factor 8 is involved in decreased trophoblast invasion under hypoxia-reoxygenation conditions. *Reprod Sci*. 2014;21(1):72-81.
- [66] Leach RE, Kilburn BA, Petkova A, Romero R and Armant DR. Diminished survival of human cytotrophoblast cells exposed to hypoxia/reoxygenation injury and associated reduction of heparin-binding epidermal growth factor-like growth factor. *Am J Obstet Gynecol*. 2008;198(4):471 e1-7; discussion e7-8.
- [67] Belkacemi L, Bainbridge SA, Dickinson MA, Smith GN and Graham CH. Glyceryl trinitrate inhibits hypoxia/reoxygenation-induced apoptosis in the syncytiotrophoblast of the human placenta: therapeutic implications for preeclampsia. *Am J Pathol*. 2007;170(3):909-20.
- [68] Kurlak LO, Mistry HD, Cindrova-Davies T, Burton GJ and Pipkin FB. Human placental renin-angiotensin system in normotensive and pre-eclamptic pregnancies at high altitude and after acute hypoxia-reoxygenation insult. *J Physiol*. 2015.
- [69] Cotechini T, Komisarenko M, Sperou A, Macdonald-Goodfellow S, Adams MA and Graham CH. Inflammation in rat pregnancy inhibits spiral artery remodeling leading to fetal growth restriction and features of preeclampsia. *J Exp Med*. 2014;211(1):165-79.
- [70] Myatt L, Eis AL, Brockman DE, Greer IA and Lyall F. Endothelial nitric oxide synthase in placental villous tissue from normal, pre-eclamptic and intrauterine growth restricted pregnancies. *Hum Reprod*. 1997;12(1):167-72.
- [71] Mayor-Lynn K, Toloubeydokhti T, Cruz AC and Chegini N. Expression profile of microRNAs and mRNAs in human placentas from pregnancies complicated by preeclampsia and preterm labor. *Reprod Sci*. 2011;18(1):46-56.
- [72] Ishibashi O, Ohkuchi A, Ali MM, Kurashina R, Luo SS, Ishikawa T, Takizawa T, Hirashima C, Takahashi K, Migita M, Ishikawa G, Yoneyama K, Asakura H, Izumi A, Matsubara S and Takeshita T. Hydroxysteroid (17-beta) dehydrogenase 1 is dysregulated by miR-210 and miR-518c that are aberrantly expressed in preeclamptic placentas: a novel marker for predicting preeclampsia. *Hypertension*. 2012;59(2):265-73.
- [73] Muralimanoharan S, Maloyan A, Mele J, Guo C, Myatt LG and Myatt L. MIR-210 modulates mitochondrial respiration in placenta with preeclampsia. *Placenta*. 2012;33(10):816-23.
- [74] Lee DC, Romero R, Kim JS, Tarca AL, Montenegro D, Pineles BL, Kim E, Lee J, Kim SY, Draghici S, Mittal P, Kusanovic JP, Chaiworapongsa T, Hassan SS and Kim CJ. miR-210 targets iron-sulfur cluster scaffold homologue in human trophoblast cell lines: siderosis of interstitial trophoblasts as a novel

504 pathology of preterm preeclampsia and small-for-gestational-age pregnancies. *Am J Pathol.*
505 2011;179(2):590-602.

506 [75] Anton L, Oларerin-George AO, Schwartz N, Srinivas S, Bastek J, Hogenesch JB and Elovitz MA.
507 miR-210 inhibits trophoblast invasion and is a serum biomarker for preeclampsia. *Am J Pathol.*
508 2013;183(5):1437-45.

509 [76] Enquobahrie DA, Abetew DF, Sorensen TK, Willoughby D, Chidambaram K and Williams MA.
510 Placental microRNA expression in pregnancies complicated by preeclampsia. *Am J Obstet Gynecol.*
511 2011;204(2):178 e12-21.

512 [77] Doridot L, Chatre L, Ducat A, Vilotte JL, Lombes A, Mehats C, Barbaux S, Calicchio R, Ricchetti M
513 and Vaiman D. Nitroso-Redox Balance and Mitochondrial Homeostasis Are Regulated by STOX1, a
514 Pre-Eclampsia-Associated Gene. *Antioxid Redox Signal.* 2014;21(6):819-34.

515 [78] Cindrova-Davies T, Herrera EA, Niu Y, Kingdom J, Giussani DA and Burton GJ. Reduced
516 cystathionine gamma-lyase and increased miR-21 expression are associated with increased vascular
517 resistance in growth-restricted pregnancies: hydrogen sulfide as a placental vasodilator. *Am J Pathol.*
518 2013;182(4):1448-58.

519 [79] Oudejans CB, Michel OJ, Janssen R, Habets R, Poutsma A, Sistermans EA, Weiss MM, Incarnato
520 D, Oliviero S, Kleiverda G, Van Dijk M and Arngimsson R. Susceptibility allele-specific loss of miR-
521 1324-mediated silencing of the INO80B chromatin-assembly complex gene in pre-eclampsia. *Hum*
522 *Mol Genet.* 2015;24(1):118-27.

523 [80] Keniry A, Oxley D, Monnier P, Kyba M, Dandolo L, Smits G and Reik W. The H19 lincRNA is a
524 developmental reservoir of miR-675 that suppresses growth and Igf1r. *Nat Cell Biol.* 2012;14(7):659-
525 65.

526 [81] Ghandadi M and Sahebkar A. MicroRNA-34a and its target genes: Key factors in cancer multidrug
527 resistance. *Curr Pharm Des.* 2015.

528 [82] Gu Y, Sun J, Groome LJ and Wang Y. Differential miRNA expression profiles between the first and
529 third trimester human placentas. *Am J Physiol Endocrinol Metab.* 2013;304(8):E836-43.

530 [83] Doridot L, Houry D, Gaillard H, Chelbi ST, Barbaux S and Vaiman D. miR-34a expression,
531 epigenetic regulation, and function in human placental diseases. *Epigenetics.* 2014;9(1).

532 [84] Mitchell PS, Parkin RK, Kroh EM, Fritz BR, Wyman SK, Pogosova-Agadjanyan EL, Peterson A,
533 Noteboom J, O'Briant KC, Allen A, Lin DW, Urban N, Drescher CW, Knudsen BS, Stirewalt DL,
534 Gentleman R, Vessella RL, Nelson PS, Martin DB and Tewari M. Circulating microRNAs as stable
535 blood-based markers for cancer detection. *Proc Natl Acad Sci U S A.* 2008;105(30):10513-8.

536 [85] Chen X, Ba Y, Ma L, Cai X, Yin Y, Wang K, Guo J, Zhang Y, Chen J, Guo X, Li Q, Li X, Wang W, Wang
537 J, Jiang X, Xiang Y, Xu C, Zheng P, Zhang J, Li R, Zhang H, Shang X, Gong T, Ning G, Zen K and Zhang CY.
538 Characterization of microRNAs in serum: a novel class of biomarkers for diagnosis of cancer and
539 other diseases. *Cell Res.* 2008;18(10):997-1006.

540 [86] Li Q, Long A, Jiang L, Cai L, Xie LI, Gu J, Chen X and Tan L. Quantification of preeclampsia-related
541 microRNAs in maternal serum. *Biomed Rep.* 2015;3(6):792-6.

542 [87] Yang S, Li H, Ge Q, Guo L and Chen F. Deregulated microRNA species in the plasma and placenta
543 of patients with preeclampsia. *Mol Med Rep.* 2015;12(1):527-34.

544 [88] Yang Q, Lu J, Wang S, Li H, Ge Q and Lu Z. Application of next-generation sequencing technology
545 to profile the circulating microRNAs in the serum of preeclampsia versus normal pregnant women.
546 *Clin Chim Acta.* 2011;412(23-24):2167-73.

547 [89] Fu G, Brkic J, Hayder H and Peng C. MicroRNAs in Human Placental Development and Pregnancy
548 Complications. *Int J Mol Sci.* 2013;14(3):5519-44.

549 [90] Gunel T, Zeybek YG, Akcakaya P, Kalelioglu I, Benian A, Ermis H and Aydinli K. Serum microRNA
550 expression in pregnancies with preeclampsia. *Genet Mol Res.* 2011;10(4):4034-40.

551 [91] Choudhury M and Friedman JE. Epigenetics and microRNAs in preeclampsia. *Clin Exp Hypertens.*
552 2012;34(5):334-41.

553 [92] Luque A, Farwati A, Crovetto F, Crispi F, Figueras F, Gratacos E and Aran JM. Usefulness of
554 circulating microRNAs for the prediction of early preeclampsia at first-trimester of pregnancy. *Sci*
555 *Rep.* 2014;4:4882.

556 [93] Chim SS, Tong YK, Chiu RW, Lau TK, Leung TN, Chan LY, Oudejans CB, Ding C and Lo YM.
557 Detection of the placental epigenetic signature of the maspin gene in maternal plasma. Proc Natl
558 Acad Sci U S A. 2005;102(41):14753-8.
559
560