Weathering of pyrogenic organic matter induces fungal oxidative enzyme response in single culture inoculation experiments

Christy Gibson ^a, Timothy D. Berry ^a, Ruzhen Wang ^{a,b}, Julie A. Spencer ^c, Cliff T. Johnston ^d,
Yong Jiang ^b, Jeffrey A. Bird ^{e,f}, Timothy R. Filley ^{a,d*}

^a *Department of Earth, Atmospheric, and Planetary Sciences and the Purdue Climate Change
Research Center, Purdue University, West Lafayette, IN 47907, USA*

^b *Soil Chemistry Research Group, Institute of Applied Ecology, Chinese Academy of Sciences,
Shenyang 110016, China*

^c *Department of Chemistry, Johns Hopkins University, Baltimore, MD, 21218, USA*

^d *Department of Agronomy, Purdue University, West Lafayette, IN 47907, USA*

^e *School of Earth and Environmental Sciences, Queens College, City University of New York,
Flushing, NY 11367, USA*

^f *Earth and Environmental Sciences PhD Program, The Graduate Center, City University of New
York, New York, NY 10016, USA*

* Corresponding author. Tel.: 1-765-494-658, E-mail address: filley@purdue.edu (T.R. Filley)

Keywords: ¹³C, pyrogenic organic matter, fungal decomposition, enzyme, phenol oxidase,
peroxidase, priming

Abbreviations: Fourier transform infrared spectroscopy (FTIR), Diffuse reflectance Fourier transform infrared spectroscopy (DR-FTIR), Nuclear magnetic resonance spectroscopy (NMR), Malt extract agar (MEA), Pyrogenic organic matter (PYOM), X-ray photoelectron spectroscopy (XPS), Highly Condensed Aromatic Carbon (HCAC)

ABSTRACT

The addition of pyrogenic organic matter (PyOM), the aromatic carbon-rich product of the incomplete combustion of plant biomass or fossil fuels, to soil can influence the rate of microbial metabolism of native soil carbon. The interaction of soil heterotrophs with PyOM may be governed by the surficial chemical and physical properties of PyOM that evolve with environmental exposure. We present results of a 36-day laboratory incubation investigating the interaction of a common white-rot fungus, *Trametes versicolor*, with three forms of ^{13}C -enriched (2.08 atom% ^{13}C) PyOM derived from *Pinus ponderosa* (450 °C): one freshly produced, and two artificially weathered (254 nm, UV light-water treatment and water-leaching alone). Analysis (FTIR, XPS) of the UV-weathered PyOM showed increased aliphatic C–H content and oxidation of aromatic carbon relative to both the original and water-leached PyOM. The addition of both weathered forms of PyOM stimulated (positively primed) fungal respiration of the growth media, while the unaltered PyOM mildly inhibited (negatively primed) respiration. Artificial weathering resulted in higher oxidative (laccase and peroxidase) enzyme activity than unaltered PyOM, possibly the result of a diminished capacity to bind reactive substrates and extracellular enzymes after weathering. However, and contrary to expectations, simple water-leached weathering resulted in a relatively higher enzyme activity and respiration than UV-weathering. The ^{13}C content of respired CO_2 indicated negligible fungal oxidation of PyOM for all treatments, demonstrating the overall low microbial reactivity of this high temperature PyOM. The increased enzymatic and positive priming response of *T. versicolor* to weathered PyOM highlights the importance of weathering-induced chemistry in controlling PyOM–microbe–soil carbon interactions.

1. Introduction

Pyrogenic organic matter (PyOM), the solid, aromatic, carbon-rich product of the incomplete combustion of plant biomass or fossil fuels, can exert many important influences on terrestrial ecosystem dynamics through its alteration of the chemical, physical, and microbiological properties of soil (Hammes and Schmidt, 2009; Lehman et al., 2009; Hart and Luckai, 2013). PyOM can make up a significant proportion (~5–45%) of soil organic carbon (SOC) (Glaser et al., 1998; Schmidt et al., 1999; Skjemstad et al., 1996, 2002) with estimates as high as 89% of SOC in some desert and urban soils (Hamilton and Hartnett, 2013). The mean residence time (MRT) of biomass-derived PyOM C in soil is much higher than its unpyrolyzed source material (Santos et al., 2012; Maestrini et al., 2013). Six-month laboratory soil mesocosms of ponderosa pine-derived PyOM showed microbial decay losses of ~0.4% compared with 30% C loss of its parent pine material (Santos et al., 2012).

MRTs of PyOM are typically on the centennial scale although a broader range, ~90–1616 years, has been reported (Singh et al., 2012). This “recalcitrance” imparted to biomass carbon is based upon an enhanced resistance to microbial decay derived from thermodynamic, structural, and inhibitory controls (i.e., high hydrophobicity, high aromaticity and condensation, and potentially toxic gaseous or extractive chemicals) (Hammes et al., 2008). The potential for enhanced C sequestration by PyOM in soil, in addition to positive influences to cation exchange capacity, pH, and enhanced microbial nutrient cycling, is one of the asserted positive benefits for the intentional production and addition of PyOM or biochar to agricultural soils, although a beneficial response to soil has not been consistently observed (Lehman et al., 2006; McHenry, 2009; Laird et al., 2010a,b; Kauffman et al., 2014). The impact that increased PyOM input to

surface soils will have in forest ecosystems experiencing greater frequency and intensity of forest fires is still being debated (Adams, 2013).

As stable as PyOM seems to be under natural soil conditions, it is, however, susceptible to environmental oxidative alteration by both biotic and abiotic factors (Kawamoto et al., 2005; Cheng et al., 2006; Zimmerman, 2010; LeCroy et al., 2013). PyOM may be susceptible to oxidization in the dark under aqueous conditions (Zimmerman, 2010), by interaction with ozone (Kawamoto et al., 2005), and on contact with air (Cohen-Ofri et al., 2006), demonstrating that PyOM can be progressively weathered by natural abiotic processes in soil or while entrained in the atmosphere (Cheng et al., 2006). Generation of an oxidative rind (LeCroy et al., 2013) can influence its wettability, solubility, and porosity and thus the ability of microbes to both inhabit and utilize it as a carbon source (Hockaday et al., 2006; Lehman et al., 2011).

It is estimated that over 7.6 Tg of particulate PyOM is emitted into the atmosphere globally with 42% contributed from open biomass burning (Bond et al., 2013). Recent studies, however, have shown that global emission values may be underestimated by a factor of 3 (Randles et al., 2013) and that rural emission contribution to this estimate is very low (Santos et al., 2014). PyOM particles can persist in the atmosphere on a time scale of days to weeks and can travel long distances prior to deposition (EPA, 2010; Santos et al., 2014) as is evidenced by the influence of PyOM on the albedo and subsequent melting of arctic glaciers (Clarke and Noone, 1985). It is unknown exactly how much PyOM is redeposited from the atmosphere and is incorporated in soil but estimates place it between < 1% to 17% of SOC (Hamilton and Hartnett, 2013; Santos et al., 2014; Lehndorff et al., 2015). Photolytic oxidation of PyOM, either at the surface of the soil or once entrained as particulate matter in the atmosphere, could be extensive, particularly where UV exposure is high (Khalizov et al., 2009). Once oxidized to produce

soluble PyOM, fractions of this material may further react photolytically or, once in the soil or aquatic system, be metabolized by microbes or bind to particle surfaces (Kuhlbusch, 1998; Hockaday et al., 2006; Fujii et al., 2009). However, current estimates of PyOM MRT do not consider the potential effects of photochemical oxidation or environmental weathering on its fate.

The addition of PyOM to soil can influence the stability of native soil carbon by increasing (positive priming) or decreasing (negative priming) rates of microbial metabolism (Hamer et al., 2004; Liang et al., 2010; Zimmerman et al., 2011; Maestrini et al., 2014). However, the magnitude and direction of the priming effect can vary with soil and PyOM type, addition rate, incubation conditions, and length of study (e.g., Zimmerman et al., 2011; Santos et al., 2012; Woolf and Lehman, 2012). In soil incubations, positive priming effects are often observed in the initial stages of the experiment which are subsequently followed by a trend to negative or neutral priming. This trend has been attributed to an initial co-metabolism of labile fractions within PyOM and SOC (Hamer et al., 2004; Cheng et al., 2009; Jones et al., 2011; Keith et al., 2011; Zimmerman et al., 2011; Maestrini et al., 2014). Negative priming effects have been attributed to physical and chemical protection mechanisms such as sorption of dissolved organic matter to PyOM surfaces, or accumulation of the more recalcitrant portion of the PyOM after the labile materials are decomposed (Bell and Worrall, 2011; Jones et al., 2011, Zimmerman et al., 2011; Singh et al., 2012; Farrell et al., 2013). More recently, the potential for PyOM to disrupt intracellular signaling of specific microbial groups has been posited as a phenomenon contributing to negative priming effects (Masiello et al., 2013).

Field and laboratory PyOM incubations have recently demonstrated that a wide range of soil microbe groups have the ability to assimilate and oxidize PyOM in soil (Santos et al., 2012;

Farrell et al., 2013; Gomez et al., 2014). Saprotrophic wood decay fungi are considered to have an enhanced capacity to degrade many forms of highly condensed aromatic carbon (HCAC) given the complement of extracellular phenoloxidases and peroxidases they produce. In particular, many forms of white-rot decay fungi, such as *Trametes versicolor*, are known to degrade and take up many forms of HCAC including manufactured fullerol nanocarbon, PAHs, and coal products (Hofrichter et al., 1999; Marquez-Rocha et al., 2000; Leonowicz et al., 2001; Andersson et al., 2003; Schreiner et al., 2009). Recent decay studies of highly condensed aromatic nanocarbon materials (e.g., carbon nanotubes, fullerenes) indicate that surface oxidation, as would be induced through chemical, biological, or photolytic oxidation (i.e. weathering that adds hydroxyl or carboxyl groups) greatly increases the metabolic response of saprophytic soil fungi (Schreiner et al., 2009; Berry et al., 2014). Knowledge of how specific microbial groups, like saprotrophic fungi or other potential degraders, respond to the presence of different weathering states of highly condensed aromatic carbon may be an important key to understanding both the short and long term ecological impact of PyOM in soils.

We present the results of a 36-day laboratory single culture incubation study investigating the interaction of a common saprotrophic white-rot fungus, *T. versicolor*, with artificially weathered (both 254 nm light in a water slurry and water leached) and unaltered 450 °C PyOM produced from ¹³C-enriched (3 atom% ¹³C) *Pinus ponderosa*. We hypothesized that the photolytically weathered PyOM will enhance the C mineralization of the malt extract agar (MEA) growth media (i.e. positive priming) compared with untreated PyOM. Similarly, we also expected greater enzymatic oxidation of the photolytically weathered PyOM resulting in the production of measurable ¹³C-labeled CO₂. In contrast, we expect that PyOM that is unaltered or simply weathered through water leaching will exhibit little inductive or primary reactivity given

its higher degree of condensed aromatic surface chemistry, lower surface oxidation, and decreased solubility of low molecular components. We quantified the decomposition of culture growth media and PyOM by monitoring the isotopic CO₂ efflux and carbon isotopic composition, and the influence of PyOM on fungal enzymatic response by measuring the activity of the oxidative enzymes laccase and peroxidase.

2. Materials and Methods

2.1. Production of PyOM

Detailed description of the methods to grow Ponderosa pine, *P. ponderosa*, saplings under ¹³C-CO₂ atmosphere and the subsequent pyrolysis were given in Bird and Torn (2006) and Santos et al. (2012). Briefly, *P. ponderosa* saplings were grown in a chamber with a pulsed ¹³CO₂ atmosphere, at the University of California, Davis. Stems were then pyrolyzed at a constant temperature of 450 °C for 5 h under N₂ with the resultant PyOM characterized using solid state ¹³C NMR, and elemental and stable isotope analysis (Santos et al., 2012). The stable carbon isotopic composition was measured to be 848‰ (2.3 atom% ¹³C), and the elemental abundance was 77.9% C, 3.4% H and 14.4% O (Santos et al., 2012). Additional detailed chemical characterization of Ponderosa pine PyOM and precursor wood, including DR-FTIR and solid state NMR, has been reported elsewhere (Chatterjee et al., 2012). Briefly, solid state ¹³C nuclear magnetic resonance (NMR) and diffuse reflectance Fourier transform infrared spectroscopy (DR-FTIR) of the source pine and PyOM showed with pyrolysis the accumulation of heat resistant aliphatic domains and fused or stacked aromatic rings and diminished polysaccharide-derived oxygenated aliphatic and carboxyl groups (Chatterjee et al., 2012). Aryl

or alkene C contributed 93% of the total acquired signal as determined by direct polarization (DP) ^{13}C NMR demonstrating that PyOM used in this study is dominated by aromatic C.

2.2. Artificial weathering of PyOM

Weathering effects were simulated by reacting the fresh, untreated PyOM (PyOMF) in an aqueous slurry for 4 weeks either in the dark (PyOMW), or under UV (254 nm) light (PyOMUV). Briefly, to produce PyOMUV, the original fresh untreated PyOMF (particle size ≤ 2 μm) was loaded as 30 mg subsamples into 4 separate 16 ml quartz vials (New England Ultraviolet Company, Branford, CT) containing 10 ml of ultra-pure water. The vials were then sealed and attached to a gently rotating carousel (5 rpm) with continuous exposure to two fixed low-pressure mercury lamps, which emit light at 254 nm, for 4 weeks in a procedure modified from Skjemstad et al. (1996). The intensity of incident UV light from these lamps was $\sim 1.5 \times 10^9$ Einstein cm^2/s (Chu et al., 1998; Deihl et al., 2002). Foil-covered vials were also prepared and used as the dark reaction comparisons, simulating only water exposure and leaching. The UV treatment applied to PyOM in this experiment was identical to Chu et al. (1998) and is calculated to produce $\sim 1879 \text{ W}/\text{cm}^2$ of energy over the four weeks. This amount of energy is equivalent to ~ 11 days of solar radiance at mid latitudes ($39^\circ 16' 22 \text{ N}$) at 2944 m above sea level (McKenna and Andreas, 1997).

Prior to isolation of the weathered PyOM, after 4 weeks of treatment, a 3 ml headspace sample was removed from the vials and injected into pre-evacuated exetainer vials (Labco Ltd., Ceredigion, UK) to determine the extent of oxidation of PyOM to CO_2 and to assess the ^{13}C -label content. The contents of the vials were then combined within treatment, filtered to $0.22 \mu\text{m}$, using a cellulose acetate filter and air dried to recover the weathered PyOM.

2.3. Analysis of PyOM surface chemistry

2.3.1. Fourier-transform infrared (FTIR)

FTIR spectra of the untreated and weathered PyOM was obtained using a Nicolet 6700 FTIR spectrometer (Thermo Fisher, Waltham, MA). A Smart Diffuse Reflectance accessory (Thermo Fisher, Waltham, MA) was used in the sample bench. The samples were analyzed by mixing 10 mg of PyOM with 190 mg of spectral grade KBr (Pike Technologies, Madison, WI) then grinding in an agate vial for 30 s using a Wig L Bug mixer (Dentsply Rinn, York, PA). The mixtures were then packed into a 6 mm diameter sample cup and the mixture surface was flattened with a metal spatula prior to analysis. Diffuse Reflectance FTIR (DR-FTIR) spectra were collected with 64 individual scans which were signal averaged using 4 cm^{-1} optical resolution in the region from $4000\text{--}580\text{ cm}^{-1}$. Sample spectra were normalized by a background spectrum of KBr using the standard Kubelka Monk transformation in OMNIC Series Software Version 8.2 (Thermo Fisher, Waltham, MA). Spectra were further analyzed by investigating the peak areas associated with O–H stretching ($3600\text{--}3300\text{ cm}^{-1}$), aromatic C–H stretch ($3000\text{--}3100\text{ cm}^{-1}$), aliphatic C–H stretch ($2850\text{--}3000\text{ cm}^{-1}$), C=O stretch ($1700\text{--}1725\text{ cm}^{-1}$), C=O stretching/C=C aromatic stretch ($1400\text{--}1600\text{ cm}^{-1}$), and C–O stretch ($1210\text{--}1320\text{ cm}^{-1}$) (Cheng et al., 2008; Keiluweit et al., 2010; Demyan et al., 2012). As a semi-quantitative measure, the relative contributions of peak area intensities were calculated as outlined in Demyan et al. (2012) and Harvey et al. (2012). Briefly, a local peak area associated with each functional group was selected by establishing a lower and upper boundary and drawing a local baseline within these boundaries to designate integration limits based on group frequencies reported by Cheng et al. (2008), Keiluweit et al. (2010) and Demyan et al. (2012). The peaks within the established

integration limits were then baseline corrected and integrated to obtain peak areas using Grams/32 AI Version 6.0 (Galactic Software, Salem, NH). Ratios of the resultant peak areas were then calculated to determine spectral difference amongst the different types of PyOM. Although FTIR is a bulk analysis, the chemical characteristics and subsequent measured differences with treatment may be attributed to surface chemical phenomenon reported in previous studies (Harvey et al., 2012).

2.3.2. X-ray photoelectron spectroscopy (XPS)

XPS was used to obtain an average surface carbon oxidation assessment of the particles as this method penetrates the top ~10 nm of the surface. For the analysis, 30 mg was sampled from the three initial pools of PyOMF. Approximately 3 mg of PyOM was dusted onto copper tape until the tape was fully covered. Samples were analyzed in a PHI 5600 XPS system using Mg K α X-rays (1253.6 eV, 15 kV, 300 W) and a high-energy electron energy analyzer, operating at a constant pass energy of 5.85 eV. XP spectra were processed using commercially available software (CasaXPS version 2.3.16, CASA Software, Ltd., Teignmouth, UK) and integration of the relevant peak areas was used for quantification. Each spectrum was given a Shirley background and energy adjusted to 284.6 eV in the C (1s) region.

2.4. Fungal inoculation experiments

To evaluate the response of the white-rot fungus *T. versicolor* (strain MAD-697-R) to untreated and artificially weathered ^{13}C -labeled PyOM, single culture inoculation experiments were established in malt extract agar (MEA), a growth media produced from malted barley with an isotopic composition of -23.8‰ $\delta^{13}\text{C}$. The large difference between $\delta^{13}\text{C}$ values in the growth

media and the PyOM allows for distinction of the two sources in evolved CO₂ and thus is used to determine relative microbial mineralization of source. *T. versicolor* was obtained from the Forest Pathology and Wood Microbiology collection at the University of Minnesota, St. Paul, Minnesota. Decay studies were performed in 12 ml borosilicate exetainer vials (Labco Ltd., Ceredigion, UK) with hard plastic screw cap fitted with glass fiber filter (GF/F) paper to allow for gaseous exchange or septum to permit headspace CO₂ accumulation. PyOM particles were mixed into 0.75 ml of a 2% MEA at a loading of 5% of the MEA C. This was poured onto a 1.5 ml, non-nutritive base layer of 3% agar precharged into the vial. The denser base media was included to prevent large PyOM from settling at the bottom of the containers during the incubation period. MEA without PyOM additions was also inoculated and prepared as controls. Microcosms (except for no-fungi controls) were then inoculated with a 1 cm diameter plug taken from the 2% MEA stock plate and incubated for 4 days in the dark at 25 °C without headspace analysis to allow fungal colonization. Headspace monitoring began after day 4 and continued for 36 days. The headspace of each vial was sampled for CO₂ on days 4, 23, 30 and 36 after 1 h accumulation periods. This was done by replacing the GF/F-containing caps with caps fitted with Teflon coated rubber septa, flushing with moistened CO₂ free air prior to accumulation time, and removing 3 ml of headspace from each vial after accumulation of gases. The sample was then injected into pre-evacuated exetainer vials (Labco Ltd., Ceredigion, UK).

2.5. Stable isotope trace gas analysis

To distinguish between growth media or ¹³C-labeled PyOM in the inoculation experiments, the stable isotope content of evolved CO₂ was determined using a Sercon Cryoprep TG2 interfaced to a Sercon 20-22 Isotope Ratio Mass Spectrometer (IRMS) (Sercon, Crewe,

U.K.). Isotopic ratios are expressed in standard delta notation (δ) and in units of per mil (‰) relative to the Vienna Pee Dee Belemnite (VBDB) standard. The CO₂ carbon source was calculated using a two component mixing model as follows:

(Equation 1 and 2)

Total C respired was calculated using the estimated cumulative C mineralized at each sample time point. This method assumed that the rate of CO₂ evolved for each subsampled time point was representative of the rate of CO₂ evolved during the period of time elapsed since the previous sample point. Cumulative CO₂ was calculated using the following equation:

(Equation 3)

where C_i is cumulative C mineralized until the current time point (i) of the CO₂ analysis, n is the number of total time points in this experiment, R_i is respiration rate of the current time point, T_i is the time interval between i point and the following time point ($i+1$). Total C respired was calculated by using the sum of the cumulative CO₂ for all sample points.

2.5. Enzyme activity assays

To determine the oxidative enzyme response of *T. versicolor* to the control and PyOM treatments, a 2, 2'-azino-bis-(3-ethylbenzothiazoline-6-sulphonate) (ABTS) assay was performed on enzymes extracted via micro-centrifugation of growth media, modified from the methods described in Heinonsalo et al. (2011) and Berry et al. (2014). To recover enzymes from the MEA layer, 4 mm square plugs of media plus fungi were extracted from each tube of which replicates within the same treatment were combined and homogenized. Three equal volume subsamples of the homogenate were added to a micro-centrifuge tube with 0.45 μ m cellulose acetate membrane filters (Costar Spin-X, Corning Inc., NY) and then centrifuged at room temperature and 15,700 g

for 30 min. Replication, therefore, for the enzyme assays was at the analytical level, not the experimental treatment level.

The reaction solution consisted of 0.5 mM MnSO₄, 1.0 mM ABTS, 5 µl of H₂O₂ (added after measuring laccase activity) and 50 mM sodium malonate buffer at pH 6 (Sigma-Aldrich, St. Louis, MO). The supernatant from each micro-centrifuge tube was added (~200 µl) to the reaction solution and the absorbance read at 420 nm after 20 s. After the initial measurement, H₂O₂ was added and absorbance was measured after 20 s. Enzyme activity was quantified against a standard curve prepared by reacting purified laccase of known activity from *T. versicolor* with reaction solution. Peroxidase activity was determined 20 s after the addition of H₂O₂ and was calculated by subtraction of laccase activity from total activity. All activities are reported in mU/mL of supernatant.

2.7. Statistical methods

Each inoculation treatment was prepared in triplicate and standard errors of means for FTIR and stable isotope measurements were calculated from the individual analysis of each replicate. Enzyme assays were done on combined inoculation treatments. Means were compared using a Two-way ANOVA and Tukey's HSD test at $p < 0.05$ using SAS JMP 10 (SAS Institute, Cary, NC).

3. Results

3.1. ¹³C Isotopic evidence for weathering-induced oxidation of PyOM

Stable isotope analysis of sample headspace after 4 weeks of artificial weathering showed highly ¹³C-enriched CO₂ values for PyOMUV (~306‰), but only mildly ¹³C enriched values for

PyOMW ($\sim -5.0\text{‰}$) compared to the head space blank $\delta^{13}\text{C}$ CO_2 value (-11‰) (Table 1). The UV weathering-derived CO_2 , however, was significantly depleted with respect to the reported $\delta^{13}\text{C}$ value of the PyOMF at $\sim 840\text{‰}$ (Santos et al., 2012). The enriched PyOMUV head space indicates UV promoted oxidation of the PyOM surface, but because the head space and water used in the experiment were not flushed of background CO_2 , the stable isotope value was diluted by more than half the anticipated $\sim 840\text{‰}$ CO_2 value. These values clearly indicate that UV weathering (PyOMUV) resulted in surface oxidation while water leaching (PyOMW) liberated only small amounts of ^{13}C -labeled CO_2 .

3.2. Chemical analysis of weathered and fresh PyOM

XPS spectra resolved into %C and %O for each treatment revealed a minor decrease in surface C and a concomitant increase in surface O for the UV light PyOMUV treatment (Table 2). These changes resulted in a slight shift in the elemental O:C ratio from 0.148 (PyOMF) to 0.177 (PyOMUV) indicating a slight increase in surficial oxygen content through photolysis. PYOMW showed minimal change in the XPS analysis.

The FTIR analysis of the PyOM treatments exhibited trends consistent with changes indicated in the XPS analysis. The sum of the combined peak areas of O–H stretching vibrations associated with free hydroxyl groups (3630 , 3535 and 3490 cm^{-1}) decreased for both PyOMUV and PyOMW with respect to PyOMF, with the largest decrease observed in the UV light treatment (Fig. 1a, Table 3). This change was driven primarily by 3630 cm^{-1} as there was no significant change in peak intensity at this wavelength among the treatments (Fig. 1). The aromatic C–H and C=C stretching vibrations (3050 cm^{-1} and 1600 cm^{-1}) decreased in PyOMUV with respect to PyOMF (Fig. 1b,c) while only the C=C stretching vibration (1600 cm^{-1})

decreased in PyOMW (Fig. 1b,c). For PyOMUV, the aliphatic C–H (2965, 2910 and 2870 cm^{-1}) and the carbonyl C=O (1705 cm^{-1}) stretching vibrations increased relative to PyOMF and PyOMW (Fig. 1b,c). This same region decreased in PyOMW relative to PyOMF. The region from 1274 cm^{-1} (C–O stretching vibration and OH plane deformation) exhibited a decrease for both weathering treatments, but with the PyOMUV treatment exhibiting a large (ca. 50%), reduction relative to the initial PyOMF.

The relative changes of functional groups attributed to oxidized carbon (carbonyl, carboxylate), aliphatic carbon (C–H), aromatic C=C and C=C–H, and OH stretching vibrations (those highly susceptible to photochemical oxidation) are used herein to elucidate how artificial weathering changes surface chemical properties that could influence interactions with fungi (Table 4). The ratios of carbonyl C=O stretching vibration (1705 cm^{-1}) to aryl C=C vibration (1600 cm^{-1}) versus aryl C–H stretching vibration (3050 cm^{-1}) to aliphatic C–H stretching vibration (2965, 2910 and 2870 cm^{-1}) shows that the PyOMUV samples become relatively enriched in oxidized carbonyl-C (C=O) over aryl-C while simultaneously trending toward a chemistry relatively enriched in aliphatic C at the expense of aromatic C–H (Table 4). Additionally, the relative contribution of the carbonyl (1705 cm^{-1}) to carboxylate (1435 cm^{-1}) C–O band increased significantly in the PyOMUV samples compared to both PyOMF and PyOMW (Table 4). Relative changes of the C–O to broad OH stretching vibrations (Table 4) exhibit a dramatic reduction concomitant with the shift to aliphatic carbon driven by a large loss in the C–O stretch associated with aromatic hydroxyl groups. Together, these results reflect the combined effects of photo-oxidation and water leaching of soluble carboxylic groups.

3.3. Total carbon respired and carbon source partitioning

Over the 36 day incubation with *T. versicolor* the cumulative production of CO₂ ranged from a low of 16.6 mg C/g MEA C for the MEA + PyOMF treatment to a high of 260 mg C/g MEA C for the MEA + PyOMW treatment (Fig. 2). The addition of PyOMF to MEA resulted in a marginally significant ($\alpha = 0.10$) reduction in CO₂ efflux with respect to the MEA-fungi control. Addition of both weathered PyOM treatments increased the production of CO₂ by over 2× for PyOMUV and 5× for PyOMW, respectively.

Across all four treatments, the average $\delta^{13}\text{C}$ values of respired CO₂ ranged from -23.5 (± 0.8), -22.1 (± 1.1), -23.9 (± 0.5), and -24.5 (± 1.5) for the 4, 23, 30 and 36 day samplings, respectively. Despite the large differences in the concentration in CO₂ efflux among the treatments, the $\delta^{13}\text{C}$ values of the CO₂ showed no difference among the treatments or between the weathered and control experiments.

3.3. Enzyme activity

Laccase activity ranged from 50.8 (control) to 265.8 mU/mL media (PyOMW) while peroxidase ranged from 55.3 (control) to 386.4 mU/mL media (PyOMW) (Fig. 3). The addition of PyOMUV and PyOMW resulted in a significant increase in both laccase and peroxidase activity above the MEA-fungi control ($\alpha = 0.05$) which had the lowest values of both enzymes. The addition of PyOMF resulted in only a moderate increase ($\alpha = 0.10$) in laccase and no increase in peroxidase activity. As with total C respired, the addition of PyOMW resulted in the most significant effects on enzyme activity.

4. Discussion

4.1. UV-treatment induced PyOM surface oxidation consistent with natural weathering

Weathering by UV exposure and water leaching induced significant changes to the surface chemistry of PyOM particles (PyOMUV) used in this incubation (Table 4). Overall, these chemical changes are consistent with expected surface alteration of highly condensed aromatic carbon substances by photo-oxidation (Chen et al., 2010; Bitter et al., 2014). Aromatic carbon readily absorbs UV light, and in the presence of oxygen and water leads to the oxidation and cleavage of the C=C bonds resulting in the formation of hydroxyl, carbonyl, and/or carboxylic groups (Bertilsson and Stefan, 1998; Wu et al., 2001). Observed changes in our current study included increased surface O content (Table 2) and a general increase in carbonyl and aliphatic chemical groups at the expense of aromatic ones (Table 3). The changes in FTIR signatures for the leached weathering treatment (PyOMW) (Fig. 1, Tables 3 and 4), which were limited to minor decreases in aromatic (3050 and 1600 cm^{-1}), minor increases in carboxyl stretches (1435 cm^{-1}), and a decrease in the broad hydroxyl distribution (3630 , 3535 and 3490 cm^{-1}) may be attributed to redistribution of more mobile moieties from within the PyOM structure to the surface without significant oxidation. These chemical alterations are also consistent with other investigations of artificial and natural aging of PyOM, which showed an increase in surficial carbonyl and carboxylic acid functional groups (Yao et al., 2010; LeCroy et al., 2013; Liu et al., 2013).

Chemical weathering of PyOM also involves a decrease in the degree of condensation of the aromatic ring structure. Weathered PyOM that has undergone extensive surface oxidation were shown to consist of less than ten fused aromatic rings substituted with carboxyl groups and one or two hydroxyl groups compared to greater than ten for their un-weathered precursors (Mao et al., 2012). In our study, higher intensity of both aliphatic and aryl C-H peak areas in PyOMUV is likely the result of a number additive processes including the leaching and

photochemical decarboxylation of carboxylic groups, the reabsorption of aliphatic compounds mobilized from the interior of the PyOM particles, and the photochemical breaking of the aryl C=C bonds (Cheng et al., 2006).

While most chemical characterization studies of PyOM aging in soils have involved the recovery of 6 month to 100 year PyOM, short-term exposure of PyOM in soils where microbial interactions may be the dominant control on chemical change have also resulted in increased surface O and carboxyl groups (Hockaday et al., 2007; Nguyen et al., 2009; LeCroy et al., 2013) as we have also seen here. Surficial changes in soil aging, however, have an added ambiguity in that it is nearly impossible to distinguish microbial oxidation from sorption of oxidized dissolved organic matter in the soil.

4.2. Artificial weathering of PyOM induces fungal enzymatic response

We found a significant increase in both laccase and peroxidase enzyme activities with the addition of PyOMUV and PyOMW ($\alpha = 0.05$). However, PyOMF resulted in only a marginal increase in laccase activity ($\alpha = 0.10$) compared to the no-PyOM control (Fig. 3). In contrast to our original hypothesis, PyOMW exhibited a significantly greater response in enzyme activity than PyOMUV ($P < 0.001$). These findings, in conjunction with the chemical characterization data, demonstrate that surface PyOM chemistry and degree of condensation plays a significant role in microbial response to PyOM (D'Annibale et al., 2005; Schreiner et al., 2008; Arun et al., 2011; Berry et al., 2014).

The distinct chemical differences observed for PyOMF, PyOMUV and PyOMW (Fig. 1, Tables 3 and 4) should influence a variety of physical properties such as inorganic and organic sorptive and binding capacity, and surface charge, which could be a controlling factor in the

distinct enzymatic and mineralization responses observed (Kasozi et al., 2010; Lehman et al., 2011). These chemical differences may even influence the binding and thus the activity of extracellular enzymes. Cho and Bailey (1978) observed decreased apparent glucose oxidase and gluconolactonase activity with the addition of char-based activated carbon to a maltose solution inoculated with the enzymes. Similarly, a study by Zhang and Yu (2000) demonstrated that activated C decreased dye decolorization and laccase activity when added to an azo dye inoculated with *T. versicolor*. In the context of the present study, such findings indicate that our untreated PyOMF may have inhibited the enzymatic activity of *T. versicolor* explaining the marginal suppression of MEA C mineralization.

In general, our results for PyOMUV and PyOMW in comparison to PyOMF are consistent with previous work on the white-rot fungal response to single wall carbon nanotubes (CNT), highly condensed aromatic carbon structures with different degrees of surface oxidation (Berry et al., 2014). These authors found that the two white rot fungi *T. versicolor* and *Phlebia tremellosa* both increased the activity of laccase and peroxidase enzymes in single culture media when CNT, with just 3% carboxyl content, were added to the growth media. However, these same fungi exhibited no response when pure, unfunctionalized CNT were added, leading to the hypothesis that surface carboxyl functional groups caused an increase in degradative enzyme activity and thus oxidative aging would be a critical factor for environmental microbial degradation. This response may be governed by the sorptive capacity of PyOM and other HCAC (Kasozi et al., 2010; Masiello et al., 2013). In some cases, this could lead to the disruption of extracellular signaling of the fungus, as demonstrated by Masiello et al. (2013) who observed reduced gene expression by gram negative bacteria as a result of sorption of stimulatory compounds on the surface of PyOM. This has implications for the weathered (leached and photo-

treated) analogs of PyOMF as increased weathering may lower capacity to bind key regulatory compounds. These results suggest that substrate and enzyme sorption, occlusion of enzyme active sites, and binding of regulatory compounds may be involved in the inhibition of substrate mineralization in PyOMF and to a lesser degree weathered PyOM. The overall suppressive effect of PyOMF is in keeping with strong adsorption of both extracellular enzymes and nutrients which should limit decomposition of both media and PyOM; similar to the interpretation of past mesocosm studies where activated char was shown to bind enzyme active sites and prevented dye decolorization (Zhang et al., 2000).

It is not immediately clear, however, what mechanisms would result in the seemingly non-linear relationship between surface chemical change, highest for PyOMUV, and fungal reactivity response, highest for PyOMW. Our original hypothesis stated that, although this particular 450 °C PyOM has been shown to have very low lability in soil (Santos et al., 2012), the oxidized PyOMUV would exhibit the highest fungal enzyme activity, PyOM decomposition, and MEA decomposition as this material would leach lignin-like signaling compounds, like veratryl alcohol (Dekker et al., 2001), to induce oxidative enzyme production (Schreiner et al., 2009; Berry et al., 2014). Interestingly, although PyOMW exhibited surface chemical changes far less than PyOMUV, as exhibited by the XPS, FTIR, and ¹³CO₂ analyses (Tables 1, 2, and 3 and Fig. 1), it induced the greatest enzyme response and MEA priming. Taken together, the relative response between PyOMF and PyOMUV can be reasonably explained by suppressive surface binding effects of PyOMF and an amplifying reactivity effect from PyOMUV because of potential inducer compounds; a hypothesis consistent with earlier studies (Berry et al., 2014). That explanation, unfortunately, runs counter to the relative reactivity observed between PyOMUV and PyOMW and their assumed chemical differences. One possible explanation that

could make sense of the three treatment responses might be gathered from the FTIR data. The increase in aryl CH and aliphatic CH stretching vibrations in PyOMUV relative to PyOMF (Fig. 1, Tables 3 and 4) suggests that the PyOMUV surface may be relatively more hydrophobic than PyOMW and thus could have potentially greater surface sorptive properties for enzymes or labile MEA compounds. Without further chemical characterization, it is not possible to unequivocally determine relative controls.

4.3. Weathered PyOM induces fungal priming but is not simultaneously degraded

Similar to the differential response of the enzyme activity upon addition of fresh and weathered PyOM, *T. versicolor* exhibited a distinct capacity to mineralize the MEA growth media C based on PyOM type (Fig. 2). We anticipated that total C mineralized would increase with degree of simulated weathering in the order MEA < PyOMF < PyOMW < PyOMUV, and that each PyOM treatment would induce some degree of MEA decomposition. However, the greatest yield in respired CO₂ was in the PyOMW treatment with the overall relative rank as PyOMF ≤ MEA < PyOMUV < PyOMW. In fact, PyOMF mildly suppressed ($\alpha = 0.1$) MEA mineralization with respect to the no-PyOM MEA-fungi control. Surprisingly, and contrary to our hypotheses, there was no PyOM ¹³CO₂ mineralized during the incubation process, at least after the first week of the experiment when CO₂ accumulation began. Therefore, the main effect of the weathered PyOM was to induce, or positively prime (Kuzyakov et al., 2009) MEA decomposition while the main influence of the PyOMF was to mildly suppress (or negatively prime) MEA decomposition. Certainly, under the appropriate conditions, *T. versicolor*, and many white-rot fungi in general, have the capacity to mineralize highly condensed polyaromatic carbon materials, which has been shown in similar laboratory and soil inoculations (Zhang and Yu,

2000; Schreiner et al., 2008; Arun et al 2011). Fungal decay studies using *T. versicolor* and highly condensed aromatic compounds have also shown that carbon mineralization potential decreases with the degree of ring condensation (Andersson et al., 2003; Atagana et al., 2009). Andersson et al. (2003) also observed a suppression of PAH degradation after wood addition. With respect to large, highly condensed aromatic particles like PyOM, prior field and laboratory soil mesocosm studies, including one by Santos et al. (2012) that investigated the exact PyOMF materials used in the present study, demonstrated PyOM oxidation to CO₂ and incorporation into both fungal and bacterial PLFAs (Maestrini et al., 2014). Furthermore, Santos et al. (2012) showed that uptake of PyOM C into fungal PLFA was lower (> 20 mol%) than in bacterial PLFA while the reverse was observed for soil-derived C. A low preference for PyOM by fungi is in agreement with the results demonstrated herein that showed no PyOM-¹³C in the headspace and indicates either a preference for MEA C, relatively lower substrate use efficiency, the inability of fungi to penetrate PyOM pore spaces to access labile PyOM C, or the need for complex microbial consortia (Boonchan et al., 2000; Jin et al., 2010; Zimmerman et al., 2011).

A portion of the resilience of PyOM in soil has been ascribed, in part, to enhancement of physical protection mechanisms within soil aggregates (Hamer et al., 2004; Zimmerman et al., 2011) or coating by a protective rind (Brodowski et al., 2006). Kasozi et al. (2010) showed that compounds containing hydroxyl and carboxyl groups can be progressively sorbed to PyOM surfaces over time and that PyOM can sorb up to 0.1% of its weight in natural organic matter (NOM) and about 1% of low molecular weight phenolic compounds. While soil aggregation is not possible in the present study, it suggests that there is potential for PyOM suspended in a nutrient rich growth media, to be protected from mineralization by essentially being coated in a protective layer that provides a barrier between the PyOM surface and extracellular enzymes.

Previous PyOM-soil mesocosm studies exhibit a positive priming effect at the early stage of the incubation (Hamer et al., 2004; Zimmerman et al., 2010, 2011; Jones et al., 2011; Cross and Sohi, 2011). Proposed mechanisms for this phenomenon include co-metabolism through extracellular enzyme production, microbial community behavior, i.e., copiotrophs versus oligotrophs, and sorption of carbon sources to PyOM (Hamer et al., 2004; Zimmerman et al., 2011; Maestrini et al., 2014). However, our study shows that no PyOM C was mineralized as CO₂, suggesting that these labile components are either inaccessible to fungi, or play a different role in the mineralization process when abundant metabolizable carbon is present. It is important to note that we allowed the fungi to incubate with the media and PyOM while the hyphae initially grew before sampling the headspace. It is possible that all the rapidly metabolizable PyOM C was consumed in this short time frame. While our study focused on one species of fungi, these results suggest possibly a less-significant role of fungi in PyOM decomposition; a finding consistent with recently published studies from soil-PyOM incubations. Using ¹³C-labeled PyOM to trace metabolic uptake into phospholipid fatty acids both Santos et al. (2012) and Farrell et al. (2013) demonstrated that gram positive bacteria were able to incorporate more PyOM C in their cells than either gram negative bacteria or fungi and they also showed that fungi had a greater preference for wood. It is also possible that if our experiments used PyOM produced over range of production temperature we would observe very different results with respect to fungal accessibility of the weathered and fresh PyOM as has been demonstrated with soil incubations (Lou et al., 2011 Zimmerman et al., 2011).

5. Conclusions

This study provides the first evidence for the direct role of surface oxidation in the response of white-rot fungi to pyrogenic organic matter in a decomposition study. Addition of weathered PyOM resulted in increased mineralization of malt extract media as indicated by increases in both CO₂ evolved and enzyme activity, serving as a microbial inducer rather than an available C source. While this study focused on one species of fungi, these results may shed light on the role of fungi in PyOM studies involving soil and sediments and help to explain why microbial priming results vary extensively among experiments. It could be that fungi are more involved in the positive priming of SOC in the presence of PyOM than bacteria. While this study did not measure the sorptive character of the different PyOM used, our results demonstrate that over time, weathering of PyOM might aid in the sequestration of C by increased binding of organic compounds with increased weathering by altering surface charge. However, further study is needed to confirm that sorption to PyOM surface is a major factor in the response of enzymatic activity and total C mineralized as well as confirming whether these fungi exhibit strategies that select for SOC over PyOM in priming studies. The increased enzymatic and positive priming response of *T. versicolor* to weathered PyOM highlights the importance of weathering-induced chemistry in controlling PyOM–microbe–soil carbon interactions.

Acknowledgements

The authors would like to thank the reviewers for their comments and suggestions, which were invaluable while completing this manuscript. The authors also acknowledge support from the National Science Foundation under award BIO-1127253 and thank Dr. Howard Fairbrother for assistance with XPS analysis and characterization of PyOM materials used in this study, Dr.

Gnanasiri S. Premachandra for assistance with DRIFTS analysis of PyOM, and Dr. Robert Blanchette for providing fungal cultures.

Associate Editor—**Myrna Simpson**

References

Adams, M.A., 2013. Mega-fires, tipping points and ecosystem services: Managing forests and woodlands in an uncertain future. *Forest Ecology and Management* 294, 250-261.

Andersson, B.E., Lundstedt, S., Tornberg, K., Schnürer, Y., Öberg, L.G., Mattiasson, B., 2003. Incomplete degradation of polycyclic aromatic hydrocarbons in soil inoculated with wood-rotting fungi and their effect on the indigenous soil bacteria. *Environmental Toxicology and Chemistry* 22, 1238-1243.

Arun, A., Eyini, M., 2011. Comparative studies on lignin and polycyclic aromatic hydrocarbons degradation by basidiomycetes fungi. *Bioresource Technology* 102, 8063-8070.

Atagana, H.I., 2009. Biodegradation of PAHs by fungi in contaminated-soil containing cadmium and nickel ions. *African Journal of Biotechnology* 8, 5780-5789.

Bell, M.J., Worrall, F., 2011. Charcoal addition to soils in NE England: a carbon sink with environmental co-benefits? *Science of the Total Environment* 409, 1704-1714.

Berry, T.D., Filley, T.R., Blanchette, R.A., 2014. Oxidative enzymatic response of white-rot fungi to single-walled carbon nanotubes. *Environmental Pollution* 193, 197-204.

Bertilsson, S., Stefan, L.J., 1998. Photochemically produced carboxylic acids as substrates for freshwater bacterioplankton. *Limnology and Oceanography* 43, 885-895.

- 549 Bird, J.A., Torn, M.S., 2006. Fine roots vs. needles: a comparison of ^{13}C and ^{15}N dynamics in a
550 ponderosa pine forest soil. *Biogeochemistry* 79, 361-382.
- 551 Bitter, J.L., Yang, J., Milani, S.B., Jafvert, C.T., Fairbrother, D.H., 2014. Transformations of
552 oxidized multiwalled carbon nanotubes exposed to UVC, 254 nm irradiation.
553 *Environmental Science: Nano* 1, 324-337.
- 554 Bond, T.C., Doherty, S.J., Fahey, D.W., Forster, P.M., Berntsen, T., DeAngelo, B.J., Flanner,
555 M.G., Ghan, S., Kärcher, B., Koch, D., 2013. Bounding the role of black carbon in the
556 climate system: A scientific assessment. *Journal of Geophysical Research: Atmospheres*
557 118, 5380–5552.
- 558 Boonchan, S., Britz M.L., Stanley G.A., 2000. Degradation and mineralization of high-
559 molecular-weight polycyclic aromatic hydrocarbons by defined fungal-bacterial
560 cocultures. *Applied and Environmental Microbiology* 66, 1007-1019.
- 561 Brodowski, S., John, B., Flessa, H., Amelung, W., 2006. Aggregate-occluded black carbon in
562 soil. *European Journal of Soil Science* 57, 539-546.
- 563 Chatterjee, S., Santos, F., Abiven, S., Itin, B., Stark, R., Bird, J., 2012. Elucidating the chemical
564 structure of pyrogenic organic matter by combining magnetic resonance, mid-infrared
565 spectroscopy and mass spectrometry. *Organic Geochemistry* 51, 35-44.
- 566 Chen, C.Y., Jafvert, C.T., 2010. Photoreactivity of carboxylated single-walled carbon nanotubes
567 in sunlight: reactive oxygen species production in water. *Environmental Science and*
568 *Technology* 44, 6674-6679.
- 569 Cheng, C.H., Lehmann, J., Thies, J.E., Burton, S.D., Engelhard, M.H., 2006. Oxidation of black
570 carbon by biotic and abiotic processes. *Organic Geochemistry* 37, 1477-1488.

- 571 Cheng, C.H., Lehmann, J., 2009. Ageing of black carbon along a temperature gradient.
572 Chemosphere 75, 1021-1027.
- 573 Cho, Y.K., Bailey, J.E., 1978. Immobilization of enzymes on activated carbon: properties of
574 immobilized glucoamylase, glucose oxidase, and gluconolactonase. Biotechnology and
575 Bioengineering 20, 1651-1665.
- 576 Chu, W., Jafvert, C. T., Diehl, C.A., Marley, K., Larson, R.A., 1998. Phototransformations of
577 polychlorobiphenyls in Brij 58 micellar solutions. Environmental Science and
578 Technology 32, 1989-1993.
- 579 Clarke, A.D., Noone, K.J., 1985. Soot in the Arctic snowpack: A cause for perturbations in
580 radiative transfer. Atmospheric Environment 19, 2045-2053.
- 581 Cohen-Ofri, I., Weiner, L., Boaretto, E., Mintz, G., Weiner, S., 2006. Modern and fossil
582 charcoal: aspects of structure and diagenesis. Journal of Archaeological Science 33, 428-
583 439.
- 584 Cross, A., Sohi, S.P., 2011. The priming potential of biochar products in relation to labile carbon
585 contents and soil organic matter status. Soil Biology and Biochemistry 43, 2127-2134.
- 586 D'Annibale, A., Ricci, M., Leonardi, V., Quarantino, D., Mincione, E., Petruccioli, M., 2005.
587 Degradation of aromatic hydrocarbons by white-rot fungi in a historically contaminated
588 soil. Biotechnology and Bioengineering 90, 723-731.
- 589 Demyan, M.S., Rasche, F., Schulz, E., Breulmann, M., Müller, T., Cadisch, G., 2012. Use of
590 specific peaks obtained by diffuse reflectance Fourier transform mid-infrared
591 spectroscopy to study the composition of organic matter in a Haplic Chernozem.
592 European Journal of Soil Science 63, 189-199.

- 593 Environmental Protection Agency, 2012. Report to Congress on Black Carbon. EPA Publication
594 450/R-12-001
- 595 Diehl, C.A., Jafvert, C.T., Marley, K.A., Larson, R.A., 2002. Surfactant-assisted UV-photolysis
596 of nitroarenes. *Chemosphere* 46, 553-560.
- 597 Farrell, M., Kuhn, T.K., Macdonald, L.M., Maddern, T.M., Murphy, D.V., Hall, P.A., Singh, B.,
598 Baumann, K., Krull, E.S., Baldock, J.A., 2013. Microbial utilisation of biochar-derived
599 carbon. *Science of the Total Environment* 465, 288–297.
- 600 Fujii, K., Uemura, M., Hayakawa, C., Funakawa, S., Kosaki, T., Ohta, S., 2009. Fluxes of
601 dissolved organic carbon in two tropical forest ecosystems of East Kalimantan,
602 Indonesia. *Geoderma* 152, 127-136.
- 603 Glaser, B., Haumaier, L., Guggenberger, G., Zech, W., 1998. Black carbon in soils: the use of
604 benzenecarboxylic acids as specific markers. *Organic Geochemistry* 29, 811-819.
- 605 Gomez, J.D., Deneff, K., Stewart, C.E., Zheng, J., Cotrufo, M.F., 2014. Biochar addition rate
606 influences soil microbial abundance and activity in temperate soils. *European Journal of*
607 *Soil Science* 65, 28-39.
- 608 Hamer, U., Marschner, B., Brodowski, S., Amelung, W., 2004. Interactive priming of black
609 carbon and glucose mineralisation. *Organic Geochemistry* 35, 823-830.
- 610 Hamilton, G.A., Hartnett, H.E., 2013. Soot black carbon concentration and isotopic composition
611 in soils from an arid urban ecosystem. *Organic Geochemistry* 59, 87–94.
- 612 Hammes, K., Schmidt, M.W.I., 2009. Changes of biochar in soil. In: Lehmann, J., Joseph, S.
613 (Eds), *Biochar for Environmental Management: Science and Technology*. Earth Scan,
614 London, pp. 169–182.

- 615 Hammes, K., Smernik, R.J., Skjemstad, J.O., Schmidt, M.W., 2008. Characterisation and
616 evaluation of reference materials for black carbon analysis using elemental composition,
617 colour, BET surface area and ^{13}C NMR spectroscopy. *Applied Geochemistry* 23, 2113-
618 2122.
- 619 Hart, S., Luckai, N., 2013. Review: Charcoal function and management in boreal ecosystems.
620 *Journal of Applied Ecology* 50, 1197-1206.
- 621 Harvey, O.R., Herbert, B.E., Kuo, L.J., Louchouart, P., 2012. Generalized two-dimensional
622 perturbation correlation infrared spectroscopy reveals mechanisms for the development
623 of surface charge and recalcitrance in plant-derived biochars. *Environmental Science and*
624 *Technology* 46, 10641-10650.
- 625 Heinonsalo, J., Kabiersch, G, Niemi, RM, Simpanen, S, 2012. Filter centrifugation as a sampling
626 method for miniaturization of extracellular fungal enzyme activity measurements in solid
627 media. *Fungal Ecology* 5, 261-269.
- 628 Hockaday, W.C., Grannas, A.M., Kim, S., Hatcher, P.G., 2006. Direct molecular evidence for
629 the degradation and mobility of black carbon in soils from ultrahigh-resolution mass
630 spectral analysis of dissolved organic matter from a fire-impacted forest soil. *Organic*
631 *Geochemistry* 37, 501-510.
- 632 Hofrichter, M., Vares, T., Kalsi, M., Galkin, S., Scheibner, K., Fritsche, W., Hatakka, A., 1999.
633 Production of manganese peroxidase and organic acids and mineralization of ^{14}C -labelled
634 lignin (^{14}C -DHP) during solid-state fermentation of wheat straw with the white rot fungus
635 *Nematoloma frowardii*. *Applied and Environmental Microbiology* 65, 1864-1870.

- 636 Jones, D.L., Murphy, D.V., Khalid, M., Ahmad, W., Edwards-Jones, G., DeLuca, T.H., 2011.
637 Short-term biochar-induced increase in soil CO₂ release is both biotically and abiotically
638 mediated. *Soil Biology and Biochemistry* 43, 1723-1731.
- 639 Kauffman, N., Dumortier, J., Hayes, D.J., Brown, R.C., Laird, D.A., 2014. Producing energy
640 while sequestering carbon? The relationship between biochar and agricultural
641 productivity. *Biomass and Bioenergy* 63, 167-176.
- 642 Kasozi, G.N., Zimmerman, A.R., Nkedi-Kizza, P., Gao, B., 2010. Catechol and humic acid
643 sorption onto a range of laboratory-produced black carbons (biochars). *Environmental*
644 *Science and Technology* 44, 6189-6195.
- 645 Kawamoto, K., Ishimaru, K., Imamura, Y., 2005. Reactivity of wood charcoal with ozone.
646 *Journal of Wood Science* 51, 66-72.
- 647 Keiluweit, M., Nico, P.S., Johnson, M.G., Kleber, M., 2010. Dynamic molecular structure of
648 plant biomass-derived black carbon (biochar). *Environmental Science and Technology*
649 44, 1247-1253.
- 650 Keith, A., Singh, B., Singh, B.P., 2011. Interactive priming of biochar and labile organic matter
651 mineralization in a smectite-rich soil. *Environmental Science and Technology* 45, 9611-
652 9618.
- 653 Khalizov, A.F., Xue, H., Wang, L., Zheng, J., Zhang, R., 2009. Enhanced light absorption and
654 scattering by carbon soot aerosol internally mixed with sulfuric acid. *The Journal of*
655 *Physical Chemistry* 113, 1066-1074.
- 656 Kuhlbusch, T.A., 1998. Enhanced: Black carbon and the carbon cycle. *Science* 280, 1903-1904.

- 657 Kuzyakov, Y., Subbotina, I., Chen, H., Bogomolova, I., Xu, X., 2009. Black carbon
658 decomposition and incorporation into soil microbial biomass estimated by ^{14}C labeling.
659 Soil Biology and Biochemistry 41, 210-219.
- 660 Laird, D.A., Fleming, P., Davis, D.D., Horton, R., Wang, B., Karlen, D.L., 2010. Impact of
661 biochar amendments on the quality of a typical Midwestern agricultural soil. Geoderma
662 158, 443-449.
- 663 Laird, D., Fleming, P., Wang, B., Horton, R., Karlen, D., 2010. Biochar impact on nutrient
664 leaching from a Midwestern agricultural soil. Geoderma 158, 436-442.
- 665 LeCroy, C., Masiello, C.A., Rudgers, J.A., Hockaday, W.C., Silberg, J.J., 2013. Nitrogen,
666 biochar, and mycorrhizae: Alteration of the symbiosis and oxidation of the char surface.
667 Soil Biology and Biochemistry 58, 248-254.
- 668 Lehmann, J., Gaunt, J., Rondon, M., 2006. Bio-char sequestration in terrestrial ecosystems—a
669 review. Mitigation and Adaptation Strategies for Global Change 11, 395-419.
- 670 Lehmann, J., Czimczik, C., Laird, D., Sohi, S., 2009. Stability of biochar in soil, In: Lehmann, J.,
671 Joseph, S. (Eds), Biochar for Environmental Management: Science and Technology.
672 Earth Scan, London, pp. 183-206.
- 673 Lehmann, J., Rillig, M.C., Thies, J., Masiello, C.A., Hockaday, W.C., Crowley, D., 2011.
674 Biochar effects on soil biota—a review. Soil Biology and Biochemistry 43, 1812-1836.
- 675 Lehdorff, E., Wolf, M., Litt, T., Brauer, A., Amelung, W., 2015. 15,000 years of black carbon
676 deposition—a post-glacial fire record from maar lake sediments (Germany). Quaternary
677 Science Reviews 110, 15-22.
- 678 Leonowicz, A., Cho, N.S., Luterek, J., 2001. Fungal laccase: properties and activity on lignin.
679 Journal of Basic Microbiology 41, 185-227.

- 680 Liang, B., Lehmann, J., Sohi, S., Thies, J., O'Neill, B., Trujillo, L., Gaunt, J., Solomon, D.,
681 Grossman, J., Neves, E., Luizão, F., 2010. Black carbon affects the cycling of non-black
682 carbon in soil. *Organic Geochemistry* 41, 206-213.
- 683 Liu, Z., Demisie, W., Zhang, M., 2013. Simulated degradation of biochar and its potential
684 environmental implications. *Environmental Pollution* 179, 146-152.
- 685 Maestrini, B, Nannipieri, P, Abiven, S, 2014. A meta-analysis on pyrogenic organic matter
686 induced priming effect. *GCB Bioenergy* 7, 577-590.
- 687 Mao, J.D., Johnson, R.L., Lehmann, J., Olk, D.C., Neves, E.G., Thompson, M.L., Schmidt-Rohr,
688 K., 2012. Abundant and stable char residues in soils: implications for soil fertility and
689 carbon sequestration. *Environmental Science and Technology* 46, 9571-9576.
- 690 Márquez-Rocha, F.J., Hernández-Rodríguez, V.Z., Vázquez-Duhalt, R., 2000. Biodegradation of
691 soil-adsorbed polycyclic aromatic hydrocarbons by the white rot fungus *Pleurotus*
692 *ostreatus*. *Biotechnology Letters* 22, 469-472.
- 693 Masiello, CA, Chen, Y, Gao, X, Liu, S, 2013. Biochar and microbial signaling: production
694 conditions determine effects on microbial communication. *Environmental Science and*
695 *Technology* 47, 11496-11503.
- 696 McHenry, M.P., 2009. Agricultural bio-char production, renewable energy generation and farm
697 carbon sequestration in Western Australia: Certainty, uncertainty and risk. *Agriculture,*
698 *Ecosystems and Environment* 129, 1-7.
- 699 McKenna, E., Andreas, A., 1997. South Park Mountain Data: South Park, Colorado (Data);
700 NREL Report No. DA-5500-56521.<http://dx.doi.org/10.5439/1052562>
- 701 Nguyen, B.T., Lehmann, J., Kinyangi, J., Smernik, R., Riha, S.J., Engelhard, M.H., 2009. Long-
702 term black carbon dynamics in cultivated soil. *Biogeochemistry* 92, 163-176.

- 703 Norwood, M.J., Louchouart, P., Kuo, L.J., Harvey, O.R., 2013. Characterization and
704 biodegradation of water-soluble biomarkers and organic carbon extracted from low
705 temperature chars. *Organic Geochemistry* 56, 111-119.
- 706 Randles, C.A., Colarco, P.R., Silva, A.D., 2013. Direct and semi-direct aerosol effects in the
707 NASA GEOS-5 AGCM: aerosol-climate interactions due to prognostic versus prescribed
708 aerosols. *Journal of Geophysical Research: Atmospheres* 118, 149-169.
- 709 Santos, F., Torn, M., Bird, J., 2012. Biological degradation of pyrogenic organic matter in
710 temperate forest soils. *Soil Biology and Biochemistry* 51, 115-124.
- 711 Santos, F., Fraser, M.P., Bird, J.A., 2014. Atmospheric black carbon deposition and
712 characterization of biomass burning tracers in a northern temperate forest. *Atmospheric*
713 *Environment* 95, 383-390.
- 714 Schmidt, M.W.I., Skjemstad, J.O., Gehrt, E., Kögel-Knabner, I., 1999. Charred organic carbon in
715 German chernozemic soils. *European Journal of Soil Science* 50, 351-365.
- 716 Schreiner, K.M., Filley, T.R., Blanchette, R.A., Bowen, B.B., Bolskar, R.D., Hockaday, W.C.,
717 Masiello, C.A., Raebiger, J.W., 2009. White-rot basidiomycete-mediated decomposition
718 of C₆₀ fullerol. *Environmental Science and Technology* 43, 3162-3168.
- 719 Singh, BP, Cowie, AL, Smernik, RJ, 2012. Biochar carbon stability in a clayey soil as a function
720 of feedstock and pyrolysis temperature. *Environmental Science and Technology* 46,
721 11770-11778.
- 722 Singh, N., Abiven, S., Maestrini, B., Bird, J., Torn, M., Schmidt, M., 2014. Transformation and
723 stabilization of pyrogenic organic matter in a temperate forest field experiment. *Global*
724 *Change Biology* 20, 1629-1642.

- 725 Skjemstad, J.O., Clarke, P., Taylor, J.A., Oades, J.M., McClure, S.G., 1996. The chemistry and
726 nature of protected carbon in soil. *Soil Research* 34, 251-271.
- 727 Skjemstad, J.O., Reicosky, D.C., Wilts, A.R., McGowan, J.A., 2002. Charcoal carbon in US
728 agricultural soils. *Soil Science Society of America Journal* 66, 1249-1255.
- 729 Woolf, D., Lehmann, J., 2012. Modelling the long-term response to positive and negative
730 priming of soil organic carbon by black carbon. *Biogeochemistry* 111, 83-95.
- 731 Wu, C., Liu, X., Wei, D., Fan, J., Wang, L., 2001. Photosonochemical degradation of phenol in
732 water. *Water Research* 35, 3927-3933.
- 733 Yao, F.X., Arbestain, M.C., Virgel, S., Blanco, F., Arostegui, J., Maciá-Agulló, J.A., Macías, F.,
734 2010. Simulated geochemical weathering of a mineral ash-rich biochar in a modified
735 Soxhlet reactor. *Chemosphere* 80, 724-732.
- 736 Zhang, F., Yu, J., 2000. Decolourisation of Acid Violet 7 with complex pellets of white rot
737 fungus and activated carbon. *Bioprocess Engineering* 23, 295-301.
- 738 Zimmerman, A.R., 2010. Abiotic and microbial oxidation of laboratory-produced black carbon
739 (biochar). *Environmental Science and Technology* 44, 1295-1301.
- 740 Zimmerman, A.R., Gao, B., Ahn, M.Y., 2011. Positive and negative carbon mineralization
741 priming effects among a variety of biochar-amended soils. *Soil Biology and*
742 *Biochemistry* 43, 1169-1179.
- 743
- 744
- 745
- 746
- 747
- 748

Table Captions

Table 1.

Mean $\delta^{13}\text{C}$ (‰) values of headspace CO_2 with standard errors ($n = 5$) after four weeks of artificial weathering in a water slurry in sealed quartz vials while exposed to UV-light (PyOMUV) and in the dark (PyOMW).

Table 2.

X-ray photoelectron spectroscopy (XPS) analysis resolved into C and O elemental % for fresh pyrogenic organic matter (PyOMF) and artificially weathered forms of PyOM treated for four weeks in a water slurry either exposed to UV light (PyOMUV) or in the dark (PyOMW). Measurements were made prior to addition to fungal inoculation experiments.

Table 3.

FTIR band assignments and peak area for fresh pyrogenic organic matter (PyOMF) and artificially weathered forms of PyOM treated for four weeks in a water slurry either exposed to UV light (PyOMUV) or in the dark (PyOMW). Measurements were made prior to addition to fungal inoculation experiments. The mean and standard error ($n = 3$) of each of the PyOMF, PyOMW, and PyOMUV treatments is shown.

Table 4.

Ratios of FTIR peak areas for of aryl C–H stretching vibration to aliphatic C–H stretching vibration ($3050/(2965, 2910, 2870 \text{ cm}^{-1})$), carbonyl C=O stretching vibration to aryl C=C vibration ($1705/1600 \text{ cm}^{-1}$), C–O stretching to OH stretching ($1274/3630, 3535, 3490 \text{ cm}^{-1}$),

carbonyl to carboxylate ($1705/1435\text{ cm}^{-1}$), and aryl C=C stretching vibration to aliphatic C-H stretching vibration ($1600/2965, 2910, 2870\text{ cm}^{-1}$) for PyOMF, PyOMW, and PyOMUV treatments. The mean and standard error ($n = 3$) of each of the PyOMF, PyOMW, and PyOMUV treatments is shown. Measurements were made prior to addition to fungal inoculation experiments.

Figure captions

Fig. 1. FTIR spectra of fresh pyrogenic organic matter (PyOMF) and artificially weathered forms treated for four weeks in a water slurry either exposed to UV light (PyOMUV) or in the dark (PyOMW) in the: (a) O-H stretching region ($3600\text{--}3300\text{ cm}^{-1}$), (b) C-H stretching region ($3100\text{--}2820\text{ cm}^{-1}$), and (c) the lower frequency fingerprint region ($1710\text{--}1200\text{ cm}^{-1}$).

Fig. 2. Total C respired, normalized per gram of malt media agar (MEA) C, over 36 days for unamended MEA (Control) and MEA amended with fresh pyrogenic organic matter (PyOMF) and artificially weathered forms treated for four weeks in a water slurry either exposed to UV light (PyOMUV) or in the dark (PyOMW). Treatments that do not share the same letters are significantly different from each other ($\alpha = 0.05$).

Fig. 3. Laccase and peroxidase enzyme activity from *T. versicolor* inoculated microcosms ($n = 3$) of malt extract agar (MEA) (Control) and MEA containing either fresh pyrogenic organic matter (PyOMF) or artificially weathered forms treated for four weeks in a water slurry either exposed

794 to UV light (PyOMUV) or in the dark (PyOMW). Treatments which do not share letters are
795 significantly different for laccase (lower case) and peroxidase (upper case) activity ($\alpha = 0.05$).

Figure 1.

Figure 2.

Figure 3.

Table 1.

Treatment	CO ₂ (μmole)			δ ¹³ C (‰)
PyOMF F	-	±	-	840 ± 0.20
PyOMW	0.13	±	0.01	-5.81 ± 3.68
PyOMUV	1.36	±	0.10	305.7 ± 16.0

F This value was obtained from elemental analysis of untreated PyOM (Santos et al., 2012).

Table 2

Element	PyOMF	PyOMW	PyOMUV
%C	85.6	84.8	83.2
%O	12.7	12.8	14.7

Table 3

Wavenumber (cm ⁻¹)	Assignment	PyOMF Area	PyOMW Area	PyOMUV Area
3630, 3535, 3490	Free O-H stretching vibration	4.5±0.9	3.7±0.3	3.3±0.3
3050	aromatic C–H stretching vibration	1.4±0.3	1.1±0.1	1.6±0.2
2965, 2910, 2870	aliphatic C–H stretching vibration	1.3±0.1	1.1±0.0	2.2±0.3
1705	carbonyl stretching vibration of carboxylic groups	3.2±0.2	2.8±0.2	5.0±0.9
1600	aromatic C=C vibration, C=O stretching vibration	14.7±0.9	11.7±0.8	9.1±0.9
1435	aromatic skeletal vibration with C–H in plane deformation or C–O stretching vibration from carbonates	8.0±0.4	6.2±0.9	5.2±0.4
1274	C–O stretching vibration, OH plane deformation	37±4	30±7	20±3

Table 4.

Band Assignments	Ratio of Wavenumbers (cm ⁻¹)	PyOMF Ratio Values	PyOMW Ratio Values	PyOMUV Ratio Values
Carbonyl C=O/Ar C=C	1705/1600	0.2±0.0	0.3±0.0	0.6±0.1
C–O/OH	1274/3630,3535,3490	8.5±0.7	8±1	6.2±0.9
Carbonyl C=O/Carboxylate C–O	1705/1435	0.4±0.0	0.50±0.0	1.0±0.1
Ar C–H/AL–CH	3050/2965,2910,2870	1.1±0.1	1.0±0.0	0.7±0.0
Ar C=C /AL C–H	1600/2965,2910,2870	11.6±0.9	10.2±0.3	4.1±0.5