Are Chinese consumers at risk due to exposure to metals in crayfish?

A bioaccessibility-adjusted probabilistic risk assessment

Qian Peng^{a,#}, Luís M. Nunes^{b,#}, Ben K. Greenfield^c, Fei Dang^d, Huan Zhong^{a,e,*}

^a Nanjing University, School of Environment, State Key Laboratory of Pollution Control and Resource Reuse, Nanjing, Jiangsu Province, People's Republic of China

^b University of Algarve, Civil Engineering Research and Innovation for Sustainability Center, Faro, Portugal

^c Environmental Health Sciences Division, School of Public Health, University of California, Berkeley, USA

^d State Key Laboratory of Soil Environment and Pollution Remediation, Institute of Soil Science, Chinese Academy of Sciences, Nanjing, Jiangsu Province, People's Republic of China

^e Environmental and Life Sciences Program, Trent University, Peterborough, Ontario, Canada

Abbreviations: HQ, hazard quotient; HI, hazard index; ILTR, increased lifetime risk; FAO, Food and Agriculture Organization; NBSC, National Bureau of Statistics of China; USITC, United States International Trade Commission; MLYR, middle and lower reaches of the Yangtze River; ICP-MS, inductively coupled plasma-mass spectroscopy; CRM, certified reference material; USEPA, United States Environmental Protection Agency; RfD, reference dose; EDI, estimated daily intake; CSF, cancer slope factor; IR, ingestion rate; BW, body weight; ED, exposure duration; EF, exposure frequency; AT, averaging time; NCRPM, National Council on Radiation Protection and Measurements; MCS, Monte Carlo Simulation; *pd*, probability distribution; USFDA, United States Food and Drug Administration.

These authors contributed equally.

* Corresponding author at: School of Environment, Nanjing University, State Key Laboratory of Pollution Control and Resource Reuse, 163 Xian Lin Da Dao, Nanjing, 210023, China. Tel.: +86 25 89680136. E-mail address: zhonghuan@nju.edu.cn

Abstract:

Freshwater crayfish, the world's third largest crustacean species, has been reported to accumulate high levels of metals, while the current knowledge of potential risk associated with crayfish consumption lags behind that of finfish. We provide the first estimate of human health risk associated with crayfish (*Procambarus clarkii*) consumption in China, the world's largest producer and consumer of crayfish. We performed Monte Carlo Simulation on a standard risk model parameterized with local data on metal concentrations, bioaccessibility (ϕ), crayfish consumption rate, and consumer body mass. Bioaccessibility of metals in crayfish was found to be variable (68–95%) and metal-specific, suggesting a potential influence of metal bioaccessibility on effective metal intake. However, sensitivity analysis suggested risk of metals via crayfish consumption was predominantly explained by consumption rate (explaining > 92% of total risk estimate variability), rather than metals concentration, bioaccessibility, or body mass. Mean metal concentrations (As, Cd, Cu, Ni, Pb, Se and Zn) in surveyed crayfish samples from 12 provinces in China conformed to national safety standards. However, risk calculation of ϕ -modified hazard quotient (HQ) and hazard index (HI) suggested that crayfish metals may pose a health risk for very high rate consumers, with a HI of over 24 for the highest rate consumers. Additionally, the ϕ -modified increased lifetime risk (ILTR) for carcinogenic effects due to the presence of As was above the acceptable level (10^{-5}) for both the median (ILTR = 2.5×10^{-5}) and 90th percentile (ILTR = 1.8×10^{-4}), highlighting the relatively high risk of As in crayfish. Our results suggest a need to consider crayfish when assessing human dietary exposure to metals and associated health risks, especially for high crayfish-consuming populations, such as in China, USA and Sweden.

Keywords: Metal, Crayfish, Human Health Risk Assessment, Uncertainty, Sensitivity

1. Introduction

Freshwater crayfish (Decapoda: Parastacidae and Astacidae) are a significant worldwide aquatic food (FAO, 2014; NBSC, 2013), with a long history of human consumption (Patoka et al., 2014). In some European countries, long-standing cultural traditions are still linked with crayfish consumption, especially in Scandinavia (Taugbøl and Skurdal, 1999). In the USA, crayfish used as food is popular in Wisconsin, Minnesota, Arkansas, Mississippi, and Texas, and especially popular in Louisiana, where more than 90 percent of US crayfish are processed (Holdich, 1993; USITC, 2003). In China, crayfish have been widely used as food since the 1980s (Mu et al., 2007).

Crayfish are also widely reported to accumulate high concentrations of metals (Alcorlo et al., 2006; Anderson et al., 1997a, b; Kouba et al., 2010; Kuklina et al., 2014). Due to their omnivory as well as necrophagia and life spans up to six years, crayfish bioaccumulate multiple toxins and heavy metals in their shells and flesh. Hazards associated with crayfish consumption have been reported, including Haff disease (Xie et al., 2010; Zhang et al., 2012), human paragonimiasis (Lane et al., 2009), *Vibrio* infections (Anda et al., 2001; Bean et al., 1998; Kay et al., 2012), intoxication by natural or man-made toxins (Vasconcelos, 1999; Vasconcelos et al., 2001), and effects resulting from exposure to organic contaminants (Gewurtz et al., 2000; Levengood and Schaeffer, 2011; Schilderman et al., 1999). Multiple studies have demonstrated heavy metal contamination in crayfish in the USA (Mason et al., 2000; Besser et al., 2007; Hothem et al., 2007; Moss et al., 2010), Italy (Bellante et al., 2015; Bruno et al., 2006), Spain (Devesa et al., 2002; Alcorlo et al., 2006; Vioque-Fernandez et al., 2007) and China (Huang, 2009; Wu et al., 2010; Peng et al., 2015). However, most of the previous studies were based on localized sampling,

84 especially in contaminated areas. Levels of metal contamination in crayfish intended
85 for human consumption at regional or national scales have rarely been reported.

86 China is the world's largest producer and consumer of crayfish (FAO, 2014; Mu
87 et al., 2007; NBSC, 2013). Of the more than 500,000 tons of reported crayfish
88 production and consumption annually worldwide (McClain and Romaine, 2007),
89 China alone is responsible for 529,000 tons per year, exporting about 26,000 tons,
90 mainly into the USA and Europe (Mu et al., 2007; NBSC, 2013; Schuler et al., 2000).
91 The majority of crayfish production is used for human consumption within China (Mu
92 et al., 2007; Qiu et al., 2012; Yue et al., 2009). Dietary metal exposure has also been
93 shown to pose health risk to consumers in China. In particular, arsenic (As) exposure
94 from consumption of rice and other foods poses incremental lifetime cancer risk
95 increases to 1 to 2 per 1,000 Chinese individuals (Li et al., 2011). Given the above
96 mentioned hazards and wide population exposure, the health risk of crayfish
97 consumption in China warrants study. Among the few studies focused on health risk
98 due to metal exposure via crayfish consumption (Schuler et al., 2000; Wu et al., 2010),
99 a small-scale survey in Shanghai City showed that lead (Pb) intake due to crayfish
100 consumption could pose risk to human health (Wu et al., 2010). However, data are
101 limited at regional or national scales in China on metal concentrations and dietary
102 bioavailability in crayfish, as well as dietary consumption rates.

103 When assessing metal exposure via diet, total metal burden often provides a
104 conservative (i.e., high) estimate of exposure due to the fact that only a fraction of
105 metal is bioavailable (Oomen et al., 2002; Versantvoort et al., 2005). In vitro
106 bioaccessibility tests, validated against in vivo studies (Li et al., 2014a; Li et al.,
107 2014b; Schroder et al., 2004), provide a reliable method to assess metal bioavailability,
108 which aids in dietary exposure and risk assessment (Amiard et al., 2008).

We report on the first large-scale assessment of health risk due to heavy metals in crayfish in China. Our study includes regions throughout China, but focuses on the main region of crayfish consumption, the middle and lower reaches of the Yangtze River delta area. Cooked red swamp crayfish (*Procambarus clarkii*) were collected from restaurants in 23 cities in China, and determined for metal concentrations (As, Cd, Cu, Ni, Pb, Se and Zn). Effective metal intake (i.e., metal bioaccessibility) was estimated using an in vitro digestion model and considered in a probabilistic risk calculation. Risk indices include increased lifetime cancer risk (ILTR) for As, and the hazard quotient (HQ) for all examined metals. Finally, sensitivity analysis was performed to compare the relative importance of metal concentration, bioaccessibility, ingestion rate, and consumer body weight for risk via crayfish consumption.

2. Materials and methods

2.1 Crayfish sampling

We obtained 210 red swamp crayfish (*Procambarus clarkii*) samples from 23 cities (12 provinces, Supplemental Information Figure S1), in which crayfish consumption has been frequently reported. Samples were obtained in September, 2012, during the peak season of crayfish consumption (i.e., June to September, Peng et al., 2015). Among the selected cities, 15 were located in the middle and lower reaches of the Yangtze River (MLYR), China's main crayfish producing and consuming area (Mu et al., 2007), and 8 were located in other regions of the country (Figure S1). A single site (restaurant) was sampled for crayfish in each city, except Nanjing, for which three separate restaurants were sampled.

Exposure was assessed as consumed, rather than from contamination of the raw commodity. Therefore, all samples were of cooked crayfish intended for human

consumption, obtained from local restaurants. Crayfish were cooked by either stewing or frying. Cooking methods had no significant effect on metal (except for Zn) bioaccessibility (nested ANOVA, site nested within cooking, $p > 0.05$; Supplemental Information Table S1), and thus the effects of cooking methods on the risk of metals were not considered in this study. We did not distinguish farm-raised versus wild-caught crayfish in this study. However, the majority of crayfish consumed in China are farm-raised (Mu et al., 2007; Yue et al., 2009). All samples were vacuum-packed, stored on ice, and transported to the lab within 3 days. After tail length determination (5.4 ± 0.8 cm), tail muscle (edible part of crayfish) was dissected, weighed (3.0 ± 1.1 g), rinsed with ultrapure water and freeze-dried. Samples were ball-milled to a fine powder and stored at -20°C until further analysis.

2.2 Determining metal concentrations in crayfish tail muscle

Subsamples of crayfish tail muscle were digested in concentrated trace metal grade nitric acid (HNO_3), and then analyzed for As, Cd, Cu, Ni, Pb, Se and Zn by inductively coupled plasma-mass spectroscopy (ICP-MS, Perkin Elmer/NexION300). The instrument detection limits were 0.08, 0.06, 0.43, 0.07, 0.48, 0.50 and 0.63 ng L^{-1} for Cd, Cu, Ni, Pb, As, Se and Zn, respectively. Quality control for digestion and ICP-MS analysis included blanks, certified reference material (CRM; scallop tissue, GBW10024), and duplicate measurement ($\text{RSD} < 10\%$). Instrumental quality control included calibration checks and running a standard after every 30 samples. Additionally, internal standards were used to correct the sensitivity drift and matrix effects (Long and Martin, 1989): Bi, Ge, In and Sc were spiked into the digested samples ($1 \text{ } \mu\text{g L}^{-1}$) before analysis. CRM (3 replicates) was digested and analyzed with each batch of samples. The recoveries (mean $\pm$ SD) of As, Cd, Cu, Ni, Pb, Se and

Zn in the CRM were $98 \pm 8\%$, $97 \pm 6\%$, $99 \pm 11\%$, $103 \pm 11\%$, $96 \pm 15\%$, $102 \pm 10\%$ and $97 \pm 11\%$, respectively.

2.3 Bioaccessibility of metals in crayfish tail muscle

Metal bioaccessibility was measured in 24 samples from 8 restaurants in MLYR, including Suqian, Suzhou, Anqing, Hangzhou, Anji, and Xinyang cities, and 2 restaurants in Nanjing City (see Supplemental Information Table S2). In this study, freeze-dried tissues were used for assessing metal bioaccessibility. Our preliminary experiments (using 10 crayfish individuals from 5 sampling sites) indicated that freeze-drying had insignificant effects on either concentration or bioaccessibility of metals. Dietary metal bioaccessibility was estimated by the *in vitro* digestion model developed by Versantvoort et al. (2005), simulating the digestion processes in the human mouth, stomach and small intestine. Briefly, digestion juices, including saliva, gastric juice, duodenal juice and bile juice, were prepared artificially and heated to $37 \pm 2^\circ\text{C}$ before use (details described in Supplemental Information Table S3). Crayfish subsamples were mixed with saliva ($\text{pH} = 6.8$) and incubated for 5 min, followed by addition of gastric juice ($\text{pH} = 1.3$) for 2 h (mixed $\text{pH} = 2\text{--}3$), and then, simultaneous administration of duodenal juice ($\text{pH} = 8.1$), bile ($\text{pH} = 8.2$) and 1 M NaHCO_3 solution for another 2 h. The final pH in the mixture was 6.5–7. The sample-to-fluid ratio was 1:95 (g mL^{-1}) at a volume ratio of 3 saliva:6 gastric juice:6 duodenal juice:3 bile:1 NaHCO_3 for the digestion juice (Oomen et al., 2003). A sample-to-fluid ratio of 1:95 instead of 1:98 (Oomen et al., 2003) was used in this study to facilitate calculating solution volumes. However, minor changes in sample-to-fluid ratios have insignificant effects on quantified metal bioaccessibility (Hamel et al., 1998; Oomen et al., 2003). Mixing was conducted at 55 rpm at 37°C , followed by centrifugation

(1640×g, 5 min) to separate the supernatants and pellets. Scallop tissue (GBW10024) was used as a CRM for quality control.

Both supernatants and residues were digested in concentrated HNO₃ and analyzed for metals by ICP-MS as described in Section 2.2. Bioaccessibility (% extractability) was defined as 100×(metal in supernatant)/(metal in tail muscle before extraction). The recoveries of metal extraction, i.e., 100×(metal in supernatant + metal remaining in the residue)/(metal in tail muscle before extraction), exhibited ranges of 89–112%, 91–109%, 86–107%, 93–112%, 91–107% and 93–113% for As, Cd, Cu, Ni, Se and Zn respectively.

2.4 Ingestion rates

We based rates of crayfish ingestion by humans on a previously published questionnaire survey performed in Nanjing City, a representative city in MLYR (Peng et al., 2015). Survey answers were validated by asking some logically identical questions with different wording and cross-checking for inconsistencies among the answers (Iarossi, 2006). Description of the survey method and study population may be found in Peng et al. (2015). Ingestion rates for survey participants were calculated based on the product of (1) the amount of crayfish consumed per meal (g/meal), and (2) the frequency of crayfish consumption during the peak season (meal/week). Ingestion rates during the peak season were averaged and reported as daily intakes. The sampled population included both consumers and non-consumers (ca. 11%) of crayfish.

2.5 Health risk assessment and uncertainty analysis

We evaluated human health risks of exposure to metals in crayfish tail muscle,

employing the methodology used for determining exposure levels to environmental contaminants that are likely to be associated with adverse health effects (USEPA, 1989). We employed the reference dose method to evaluate hazard of noncarcinogenic effects. In this method, estimated daily intake is compared to a reference dose (RfD), with an acceptably low lifetime risk of deleterious effect, considered protective to broader population groups, including sensitive populations (USEPA, 1989, 2015a, b).

We estimated hazard quotients (HQ; unitless) to evaluate hazard of non-carcinogenic effects; HQ are obtained by dividing the estimated mean daily intake for a specific metal i (EDI_i ; mg/kg/d) by the RfD (mg/kg/d):

$$HQ_i = EDI_i / RfD_i \quad (1)$$

A hazard index is then obtained by summing the HQ for the different metals:

$$HI = \sum_i HQ_i \quad (2)$$

For As, being a carcinogenic metal, the increased lifetime risk of cancer (ILTR) is also estimated, corresponding to the increased probability of developing cancer due to the quantified exposure:

$$ILTR_j = EDI_j \times CSF_j \quad (3)$$

CSF_j stands for cancer slope factor, which is a plausible upper-bound estimate of the probability of a response per unit intake of a chemical over a lifetime (USEPA, 1989). The acceptable carcinogenic risk is usually considered to be between 1 per million and 1 per hundred thousand (Hunter and Fewtrell, 2001). The RfD and CSF parameters in the risk models (1) through (3) were set at values provided by USEPA (2015a, b) and ATDSR (2007), compiled in Table 1. <Table 1>

Estimated daily intake (EDI) from crayfish consumption was obtained by multiplying daily consumption rate by corresponding metal concentrations in the

foodstuff, adjusted by the bioaccessibility (ϕ_i), considered equal regardless of the cooking method, as justified above:

$$EDI_i = (C_i \times \phi_i \times IR \times ED \times EF) / (BW \times AT) \quad (4)$$

where C_i = concentration of metal i in crayfish (mg/kg); IR = ingestion rate (kg/d); BW = body weight (kg); ED = exposure duration (years); EF = exposure frequency (days/year); and AT = averaging time, set at 75 years $\times$ 365 days/year.

Quantitative uncertainty analysis was employed to obtain a population estimation of the range of risks encountered (NCRPM, 1996). Sensitivity analysis was further performed to identify the parameters which had the strongest impact on the model outcomes (Hamby, 1994), being therefore the best candidates for further scientific assessment, as well as the best indicators for decision making and public communication.

Monte Carlo Simulation (MCS) was used to perform quantitative sensitivity and uncertainty analysis (Metropolis and Ulam, 1949; Thompson and Graham, 1996), employing as inputs probability distributions based on empirical data. Specifically, the propagation of uncertainty through the risk equations (1) through (4) was simulated. Statistical distributions were derived for human body weight, ingestion rate, crayfish metal concentration, and bioavailability. The non-parametric Kolmogorov-Smirnov test ($\alpha = 0.01$) was used to compare each empirical statistical distribution with the sample probability distribution (pd). Only Normal, Lognormal and Weibull distributions were tested, as they are the most frequently used in exposure and risk characterization studies (Apt, 1976; Caldwell et al., 2008). Body weight (BW) per age and sex was separated in yearly classes from birth until age 20, then every ten years until age 80. Above 80, a single age class was considered (Li et al., 2011). The pd for BW was obtained by attributing to each age class the calculated BW and

percentage of the national population. The remaining variables were obtained specifically for the study and therefore their *pd* were obtained directly from the sampled data. MCS were performed in Crystal Ball© (Oracle, Redwood City, CA, USA), using Latin hypercube sampling. Individual parameter contributions to variance in the outcomes (noncancer hazard index and increased lifetime cancer risk) were estimated by taking the square of the spearman rank correlation coefficient between each input variable and each outcome, and normalizing to 100%. The maximum number of trials that Crystal Ball ran before stopping the simulation was set at 500,000.

3. Results and discussion

3.1 Metal concentrations in crayfish

Observed As, Cd, Cu, Ni, Pb, Se and Zn concentrations (wet weight, mean $\pm$ SD) were 253.3 ± 142.4 $\mu\text{g/kg}$, 29.8 ± 39.2 $\mu\text{g/kg}$, 9.2 ± 5.8 mg/kg , 171.2 ± 141.2 $\mu\text{g/kg}$, 23.3 ± 29.5 $\mu\text{g/kg}$, 294.4 ± 154.1 $\mu\text{g/kg}$ and 20.1 ± 6.3 mg/kg , respectively (N = 210, tail muscle samples). Concentrations varied widely among cities and large variations in metal concentrations were observed for each metal (Table 2). It has been widely reported that effects of cooking on metal concentrations in fish and shellfish are minimal (Perelló et al., 2008; Kalogeropoulos et al., 2012), and thus the reported crayfish metal levels should not be significantly affected by cooking. Metal concentrations for As, Cd and Pb in all samples were below Chinese national safety standards (GB2762-2012; SAPRC, 2012); Cu, Ni, Se and Zn are not listed in the national standards. Crayfish metal concentrations found in our study were generally within the ranges of those reported in Chinese freshwater fish (Cheung et al., 2008), although As concentrations (253 ± 142 $\mu\text{g/kg}$) were higher than the freshwater fish

from the MLYR region (17 ± 9 $\mu\text{g/kg}$ for 469 fish samples, Yi et al., 2011). Kouba et al. (2011) summarized global literature on Cu, Zn, Ni, Cd, and Pb in crayfish tail muscle on a dry weight basis. We converted concentration ranges in our study from wet to dry weight assuming 75% moisture content (Peng et al., 2015), to facilitate comparison to this range of concentrations observed globally. Concentration ranges were comparable for Cu, which were 7.2–157.6 mg/kg dry weight in our study versus 2.3–110.8 mg/kg in Kouba et al. (2011). Concentration ranges were also comparable for Ni (0.06–3.8 our study versus 0.3–3.8 mg/kg globally) and Zn (40.4–196 versus 4.0–95.6 mg/kg). However, our study concentration ranges were generally lower than global concentration ranges for Cd (0.003–1.71 versus 0.015–4.3 mg/kg) and Pb (<0.001–1.01 versus 0.015–11.9 mg/kg). The relatively low concentrations of Cd and Pb may be explained by the fact that crayfish is mostly farm-raised in China (Mu et al., 2007; Yue et al., 2009). In particular, aquaculture management practices in China, including short cultivation duration and rapid growth rates (Mu et al., 2007), may result in growth dilution. This could be partly responsible for the relatively low metal levels (Zhang et al., 2010). <Table 2>

Compared to total As in our study crayfish (mean = 253 $\mu\text{g/kg}$ wet; 1012 $\mu\text{g/kg}$ dry weight), As concentrations in crayfish varied widely among regions sampled globally. Muscle As concentrations were lower in crayfish muscle obtained from two remote freshwater streams in Maryland, USA (60 and 120 $\mu\text{g/kg}$ dry weight; data extracted from Figure 9 in Mason et al., 2000) and from *Pacifastacus leniusculus* and *Procambarus clarkii* collected from California, USA streams with historic gold mining influence (240 to 680 $\mu\text{g/kg}$ dry weight; Hothem et al., 2007). In contrast, total As was higher in freshwater crayfish (*Cherax destructor* Clark) muscle from western Victoria, Australia sites with historic coal or gold mining (1700 to 2600 $\mu\text{g/kg}$ dry

weight; Williams et al., 2009) and whole crayfish (*P. clarkii*) obtained from a creek below a severe breach of a mine tailings reservoir in Doñana National Park, Andalusia, Spain (1600 to 8500 µg/kg dry weight; Devesa et al., 2002). Concentrations were also generally higher in crayfish (*P. clarkii*) muscle from two agriculturally impacted lakes in Sicily, Italy (median = 1820; range < 200 to 4030 µg/kg dry weight; Bellante et al., 2015).

3.2 Metal bioaccessibility in crayfish

Across all measured samples, mean bioaccessibility (% extractability) of the examined trace elements in crayfish was $77.6\% \pm 16.5$, $78.7\% \pm 6.1$, $82.6\% \pm 12.6$, $89.2\% \pm 17.1$, $92.5\% \pm 2.9$, and $95.4\% \pm 6.2$ for Zn, Cd, Se, Ni, Cu and As, respectively (Figure 1; Supplemental Information Table S2). Lead bioaccessibility was not determined because extracted Pb levels were too low to be precisely measured. <Figure 1>

Metal bioaccessibility in cooked aquatic food has been reported in only a few studies (He et al., 2010; He and Wang, 2013; Houlbreque et al., 2011; Metian et al., 2009). Bioaccessibility of metals in this study were generally in agreement with those reported in steamed seafood tissue (47.8%–85.9% for clams, He and Wang, 2013; 52.2%–84.3% for seabass, He et al., 2010), but higher than those in cooked mussels (34%–63%, Metian et al., 2009). Cooking was reported to decrease metal bioaccessibility in mussels (Houlbreque et al., 2011), gastropods (Amiard et al., 2008), and fish (He et al., 2010). In most previous surveys, metal bioaccessibility was quantified in uncooked shellfish, which may not be realistic for human consumption of cooked shellfish. The explicit inclusion of bioaccessibility of cooked samples in exposure estimates should reduce the uncertainty and bias in risk assessment. All

bioaccessibility data for a specific metal was treated as coming from a single population, and the propagation of this uncertainty was assessed through the MCS.

3.3 Ingestion rates

Peng et al. (2015, Table 4) estimated consumption of crayfish in China and in the middle and lower reaches of the Yangtze River (MLYR). According to these estimates, crayfish consumption outside MLYR was only about 4% of the total national consumption. This observation is consistent with previous studies (Qiao and Jiang, 2010; Shen, 2010; Wang, 2010), and reports that MLYR is the main producing and consuming area of crayfish in China and globally (FAO, 2014; Mu et al., 2007; NBSC, 2013). In MLYR, crayfish consumption rates could reach more than 100 metric tons per day per city (Mu et al., 2007) during the peak season (i.e., June to September; Peng et al., 2015). Therefore, crayfish consumption in MLYR would represent the majority of human exposure in China, as well as a conservative (i.e., high) estimate of individual exposure rates for crayfish consumers in China. For these reasons, we estimate crayfish ingestion rate (IR) using survey data from Nanjing (Peng et al., 2015), as a proxy of a typical city in MLYR.

The ingestion rate (IR) in the Nanjing survey data (Peng et al., 2015) is highly skewed and varies dramatically among individuals (Figure 2). The skewness is illustrated by median IR (8.8 g/d/person), well below mean (27.6 g/d/person) and 90th percentile (93.3 g/d/person). <Figure 2>

To accommodate such variability in the estimate, the IR frequency distribution was introduced in the probabilistic modelling of risk as a custom *pd*. Specifically, ingestion rate was modeled by drawing repeated samples from the individual survey results. Given the strong seasonal variation in consumption and resulting intermittent

exposure, the calculation of the risk from intermittent exposures was calculated as the sum of short-term exposures, by using the averaging procedure included in equation (1), with EF equal to 120 d/year. Using data reported in Table 4 and Table 5 of Peng et al. (2015), the mean consumption rate from the Nanjing survey in the peak season (27.6 g/d/person) is approximately 6 times of that of Hubei Province (4.7 g/d/person) and 14 times of the MLYR as a whole (2.0 g/d/person), the latter values obtained from China population annual fishery statistics (NBSC, 2013; Peng et al., 2015). The discrepancy likely results from the fact that most crayfish consumption occurs during the 120 day harvesting season, when the survey was conducted. Thus, the Nanjing survey results employed in our probabilistic risk calculations represent a conservative estimate of exposure during the peak harvesting season.

3.4 Health risk due to exposure to metals by consuming crayfish in MLYR

As discussed above (Section 3.3) and documented in previous studies (Mu et al., 2007; Peng et al., 2015; Qiao and Jiang, 2010; Shen, 2010; Wang, 2010), crayfish consumption in MLYR represents the highest potential health hazard for crayfish consumption in China. Therefore, the following risk calculation and discussion focuses on MLYR, which could be interpreted as a conservative estimate of risk associated with crayfish consumption in China. The estimated exposure for MLYR residents was expressed by probability distributions derived from data presented in Table 3. Correlations between model parameters were weak as indicated by correlation coefficients always lower than 0.45 ($p < 0.05$), indicating that parameter dependencies have little impact on the overall risk results (USEPA, 2001). <Table 3>

Sensitivity analysis results indicated that greater than 92% of the variation in calculated risk was due to ingestion rate, with a small portion (< 7%) attributable to

arsenic concentrations. All other parameters (including body weight, bioaccessibility, and other metals concentrations) explained < 0.5% individually and < 1.5% in combination (Table 4). The unimportance of bioaccessibility (< 0.1%) is somewhat surprising, and likely stems from high bioaccessibility coefficients and a limited range. In light of this low sensitivity, any uncertainty in bioaccessibility would be expected to have a trivial effect on risk estimates. The overwhelming importance of crayfish consumption rate stems from the wide range in consumption rates among individuals sampled. High variability in seafood consumption rates has been well documented across individuals and populations, and as a result local and subpopulation-specific consumption rate estimates are useful prior to generating seafood consumption guidance (e.g., Allen et al., 1996; Sechena et al., 2003; SFEI and CDHS, 2000; USEPA, 2000; USEPA, 2011). Our results, in combination with these findings, suggest that consumption rate estimates would be the primary monitoring priority for obtaining better estimates of risk due to dietary metals exposure from crayfish. <Table 4>.

MCS-derived median (50th percentile) and P90 (90th percentile) HQ values were well below 1.0, indicating low health hazard per metal (Table 1). The highest HQ was found for As (P90 = 0.40), followed by Cu (P90 = 0.11). P90 for the combined HI was 0.57, indicating that the highly exposed individuals are exposed to intakes close to the maximum admissible. Although less than 1.0, these HI values may still be important for crayfish consumers because crayfish is not the only significant source of metal exposure (e.g., Li et al., 2011). In the consumption survey (Peng et al., 2015), 0.5% (i.e., 3 out of 501 respondents) of the surveyed individuals (high consumption population) reported consuming over 466 g/d of crayfish muscle, resulting in a HI of over 24 (see range in Table 1). Although high-end estimates of consumption rates are

likely subject to greater measurement uncertainty (Sechena et al., 2003), our results do suggest that some individuals in Nanjing City (and MLYR generally) may be exposed to metals at hazardous levels from crayfish consumption.

The relative contribution of each metal to our estimated HI differed from that for freshwater fish of the MLYR (Yi et al., 2011). For our MCS P90 results, arsenic contributed 70% to total HI, followed by Cu (18%), Zn (5%) and Se (4%), with a trivial ($< 0.001\%$) contribution due to Pb (Table 1). In contrast, for freshwater fish, Pb explained 36.5% of total HI, Cd 30.2%, Cu 11.4%, and Zn 6.3% (Yi et al., 2011). These differences result from higher As concentrations in crayfish than in freshwater fish in China (discussed above). The moderately high contribution of Cu to dietary exposure risk, together with prior studies (e.g., Wang et al., 2001), indicates the importance of considering Cu contamination in aquatic food and associated food safety issues in China.

The median and 90th percentile of ILTR from As were estimated to be 2.5 per 100,000 and 18 per 100,000, respectively. Both are greater than typical acceptable risk thresholds (i.e., above 1 per 100,000; USEPA, 2000). A previous study calculated a total ILTR due to As intake in the Chinese diet as 177 per 100,000 (Li et al., 2011). Employing this rate, for typical versus high rate consumers in the MLYR, crayfish could contribute to approximately 1.4–9.6% of the total risk due to dietary As exposure. Of course these results do not address the combined risks from additional carcinogenic chemicals that may be present in seafood (e.g., halogenated organic compounds; USEPA, 2000), nor the potential for underestimation of risk due to intermittent exposures (Felter et al., 2011; Halmes et al., 2000).

It should also be noted that variations in As speciation in crayfish (e.g., mainly as dimethylarsinate, Williams et al., 2009) could play a role in As bioavailability,

toxicity, and resulting risk. Our results demonstrated that As bioaccessibility was generally high across samples (i.e., $95.4 \pm 6.2\%$, Supplemental Information Table S2), suggesting that potential variations in As speciation in these samples may play a minor role in As bioaccessibility. However, because we did not determine As species, and inorganic arsenic is more toxic than organic arsenic (ATSDR 2007), our risk interpretations are conservative, and As speciation in crayfish prepared for human consumption warrants further investigation.

4. Conclusion

This study provided the first large-scale survey and probabilistic risk assessment of metals in crayfish consumed in China, the world's largest producer and consumer of crayfish. Metal concentrations in crayfish were generally low to moderate, with concentrations of Cd and Pb on the low end of observed global concentrations. An exception was As, which exhibited higher concentrations than finfish in China (Yi et al., 2011) and crayfish from USA streams (Mason et al., 2000; Hothem et al., 2007). Metal bioaccessibility in crayfish, determined by an in vitro digestion model, were a priori, and based on previous studies, considered to provide a more realistic risk assessment. Results found variable bioaccessibility among different metals, which was considered in our risk calculations. Risk models showed, however, little sensitivity to bioaccessibility coefficients, likely due to high values and limited ranges. The calculated median and 90th percentile value of HQ and HI (bioaccessibility-modified) suggested low health hazard due to most heavy metals in crayfish, indicating that contamination by Cu, Cd, Pb, Ni, Se, and Zn in crayfish are unlikely to pose a health hazard for noncarcinogenic effects to the general population in China.

The risk for carcinogenic effects due to the presence of As was above the acceptable level of 1 per 100,000, for both the median ($ILTR = 2.5 \times 10^{-5}$) and 90th percentile ($ILTR = 1.8 \times 10^{-4}$). Consequently, As contamination in crayfish may pose a health risk for high rate consumers in regions of China where crayfish consumption is popular (e.g., MLYR). Considering the already high reported human dietary exposure to As in China, particularly due to rice consumption (Li et al., 2011), we may expect that supplementary sources of As intake for high rate crayfish consumers may pose a meaningful risk increase. There are two caveats to this interpretation. First, we did not determine the proportions of the more toxic inorganic As species, which have been quite variable in the few studies where they were measured in crayfish (Devesa et al., 2002; Williams et al., 2009). Second, risk calculations in this study were mainly based on data from MLYR, which is the main area of crayfish consumption in China. Thus, conclusions about risk drawn in this study are conservative and should be extrapolated with caution to other areas in China, with generally lower crayfish consumption.

Results from the stochastic simulation emphasized the dominant role of consumption rate in controlling risk of metals via crayfish consumption in China. For instance, risk calculation demonstrated that crayfish metals may pose a health risk for high rate consumers (e.g., in MLYR), highlighting the importance of considering ingestion rates when assessing risk of crayfish metals. Therefore, guidance for crayfish consumption in China (especially in MLYR cities) and other world regions with high crayfish consumption may help reduce potential hazard, especially for high consumption individuals. To mitigate the potential risk associated with eating crayfish, and further delineate exposure for high rate consumers and sensitive populations, additional consumption surveys for crayfish and other freshwater fish in China are also warranted. For example, this survey about consumption rates was conducted in

urban areas, which may differ from rural populations. Probabilistic risk assessments may also be warranted for crayfish consumers in other crayfish consuming regions such as Scandinavian countries and the southeastern United States, especially for As which varies widely among regions. Finally, in addition to metals, contamination of crayfish by organic contaminants may pose an added risk to crayfish consumers. These issues may affect the overall risk associated with crayfish consumption, warranting further investigation in future studies.

Acknowledgements

The authors thank Dr. Douglas Evans for his kind help and two anonymous reviewers for providing very constructive comments to the manuscript. Financial support was provided to HZ by the National Natural Science Foundation of China (41273087). LN was supported by European Union Marie Curie Actions, Grant FP People 2010 “IRSES Electroacross” and BG by the SAGE-IGERT Fellowship (US National Science Foundation).

References

- Alcorlo P, Otero M, Crehuet M, Baltanas A, Montes C. The use of the red swamp crayfish (*Procambarus clarkii*, Girard) as indicator of the bioavailability of heavy metals in environmental monitoring in the River Guadiamar (SW, Spain). *Sci Total Environ* 2006; 366:380–90.
- Allen MJ, Velez PV, Diehl DW, McFadden SE, Kelsh M. Demographic variability in seafood consumption rates among recreational anglers of Santa Monica Bay, California, in 1991-1992. *Fish B-Noaa* 1996; 94:597–610.
- Amiard JC, Amiard-Triquet C, Charbonnier L, Mesnil A, Rainbow PS, Wang WX. Bioaccessibility of essential and non-essential metals in commercial shellfish from Western Europe and Asia. *Food Chem Toxicol* 2008; 46:2010–22.
- Anda P, Del Pozo JS, Garcia JMD, Escudero R, Pena FJG, Velasco MCL, Sellek RE, Chillaron MRJ, Serrano LPS, Navarro JFM. Waterborne outbreak of tularemia associated with crayfish fishing. *Emerg Infect Dis* 2001; 7:575–82.
- Anderson MB, Preslan JE, Jolibois L, Bollinger JE, George WJ. Bioaccumulation of lead nitrate in red swamp crayfish (*Procambarus clarkii*). *J Hazard Mater* 1997a; 54:15–29.
- Anderson MB, Reddy P, Preslan JE, Fingerman M, Bollinger J, Jolibois L, Maheshwarudu G, George WJ. Metal accumulation in crayfish, *Procambarus clarkii*, exposed to a petroleum-contaminated bayou in Louisiana. *Ecotox Environ Safe* 1997b; 37:267–72.
- Apt KE. Applicability of the Weibull distribution function to atmospheric radioactivity data. *Atmos Environ* 1976; 10(9):777–81.
- ATSDR. Toxicological profile for Arsenic. Atlanta, GA: Agency for Toxic Substances and Disease Registry. United States Department of Health and

Human Services; 2007. <http://www.atsdr.cdc.gov/toxprofiles/index.asp> [accessed
2 December 2015]

Bean NH, Maloney EK, Potter ME, Korazemo P, Ray B, Taylor JP, Seigler S,
Snowden J. Crayfish: A newly recognized vehicle for vibrio infections.
Epidemiol Infect 1998; 121:269–73.

Bellante A, Maccarone V, Buscaino G, Buffa G, Filiciotto F, Traina A, Del Core M,
Mazzola S, Sprovieri M. Trace element concentrations in red swamp crayfish
(*Procambarus clarkii*) and surface sediments in Lake Preola and Gorgi Tondi
natural reserve, SW Sicily. Environ Monit Assess 2015; 187:404.

Besser JM, Brumbaugh WG, May TW, Schmitt CJ. Biomonitoring of lead, zinc, and
cadmium in streams draining lead-mining and non-mining areas, southeast
Missouri, USA. Environ Monit Assess 2007; 129:227–41.

Bruno G, Volpe MG, De Luise G, Paolucci M. Detection of heavy metals in farmed
Cherax destructor. B Fr Pêche Piscic 2006; 380-381:1341–9.

Buchet JP, Lison D, Ruggeri M, Foa V, Elia G. Assessment of exposure to inorganic
arsenic, a human carcinogen, due to the consumption of seafood. Arch Toxicol
1996; 70:773–8.

Caldwell DJ, Mastrocco F, Hutchinson TH, Laenge R, Heijerick D, Janssen C,
Anderson PD, Sumpter JP. Derivation of an aquatic predicted no-effect
concentration for the synthetic hormone, 17 alpha-ethinyl estradiol. Environ Sci
Technol 2008; 42(19):7046–54.

Cheung KC, Leung HM, Wong MH. Metal concentrations of common freshwater and
marine fish from the Pearl River Delta, South China. Arch Environ Con Tox
2008; 54:705–15.

Devesa, V, Suner, MA, Lai, VWM, Granchinho, SCR, Martinez, JM, Velez, D,

547 Cullen, WR, Montoro, R. Determination of arsenic species in a freshwater
 548 crustacean *Procambarus clarkii*. Appl Organomet Chem 2002; 16:123–32.

549 FAO. FAO Yearbook: Fishery and Aquaculture Statistics. Rome, Italy: Food and
 550 Agriculture Organization; 2014.

551 Felter SP, Conolly RB, Bercu JP, Bolger PM, Boobis AR, Bos PMJ, Carthew P,
 552 Doerr NG, Goodman JJ, Harrouk WA, Kirkland DJ, Lau SS, Llewellyn GC,
 553 Preston RJ, Schoeny R, Schnatter AR, Tritscher A, Van Velsen F, Williams GM.
 554 A proposed framework for assessing risk from less-than-lifetime exposures to
 555 carcinogens. Crit Rev Toxicol 2011; 41:507–44.

556 Gewurtz SB, Lazar R, Haffner GD. Comparison of polycyclic aromatic hydrocarbon
 557 and polychlorinated biphenyl dynamics in benthic invertebrates of Lake Erie,
 558 USA. Environ Toxicol Chem 2000; 19:2943–50.

559 Halmes NC, Roberts SM, Tolson JK, Portier CJ. Reevaluating cancer risk estimates
 560 for short-term exposure scenarios. Toxicol Sci 2000; 58:32–42.

561 Hamby D. A review of techniques for parameter sensitivity analysis of environmental
 562 models. Environ Monit Assess 1994; 32:135–54.

563 Hamel SC, Buckley B, Liou PJ. Bioaccessibility of metals in soils for different liquid
 564 to solid ratios in synthetic gastric fluid. Environ Sci Technol 1998; 32:358–62

565 He M, Ke CH, Wang WX. Effects of cooking and subcellular distribution on the
 566 bioaccessibility of trace elements in two marine fish species. J Arg Food Chem
 567 2010; 58:3517–23.

568 He M, Wang WX. Bioaccessibility of 12 trace elements in marine molluscs. Food
 569 Chem Toxicol 2013; 55:627–36.

570 Holdich DM. A review of astaciculture: freshwater crayfish farming. Aquat Living
 571 Resour 1993; 6(04):307–17.

572 Hothem RL, Bergen DR, Bauer ML, Crayon JJ, Meckstroth AM. Mercury and trace
 573 elements in crayfish from Northern California. *B Environ Contam Tox* 2007;
 574 79:628–32.

575 Houlbreque F, Herve-Fernandez P, Teyssie JL, Oberhaensli F, Boisson F, Jeffree R.
 576 Cooking makes cadmium contained in Chilean mussels less bioaccessible to
 577 humans. *Food Chem* 2011; 126:917–21.

578 Huang X. The contamination evaluation of the five heavy metals enriched in
 579 *Procambarus clarkii* and the water in Anhui. Academic Dissertation. Anhui
 580 Agricultural University; 2009. [In Chinese].

581 Hunter P, Fewtrell L. Acceptable risk. In: Fewtrell L, Bartram J (Eds.): World Health
 582 Organization. *Water Quality: Guidelines, Standards and Health*. London,
 583 England: IWA Publishing; 2001; 207–27.

584 Iarossi G. The power of survey design: A user's guide for managing surveys,
 585 interpreting results, and influencing respondents. Washington, DC: The World
 586 Bank; 2006.

587 Kalogeropoulos N, Karavoltos S, Sakellari A, Avramidou S, Dassenakis M, Scoullou
 588 M. Heavy metals in raw, fried and grilled Mediterranean finfish and shellfish.
 589 *Food Chem Toxicol* 2012; 50(10):3702–8.

590 Kay MK, Cartwright EJ, MacEachern D, McCullough J, Barzilay E, Mintz E, Duchin
 591 JS, MacDonald K, Turnsek M, Tarr C. *Vibrio mimicus* infection associated with
 592 crayfish consumption, Spokane, Washington, 2010. *J Food Protect* 2012;
 593 75:762–4.

594 Kouba A, Buric M, Kozak P. Bioaccumulation and effects of heavy metals in crayfish:
 595 A review. *Water Air Soil Poll* 2010; 211:5–16.

596 Kuklina I, Kouba A, Buric M, Horka I, Duris Z, Kozak P. Accumulation of heavy

metals in crayfish and fish from selected Czech reservoirs. *BioMed Res Int* 2014;
 306103, 9 pages. <http://dx.doi.org/10.1155/2014/306103> [accessed 2 December
 2015]

Lane MA, Barsanti MC, Santos CA, Yeung M, Lubner SJ, Weil GJ. Human
 paragonimiasis in North America following ingestion of raw crayfish. *Clin Infect
 Dis* 2009; 49:E55–61.

Levengood JM, Schaeffer DJ. Polycyclic aromatic hydrocarbons in fish and crayfish
 from the calumet region of Southwestern Lake Michigan. *Ecotoxicology* 2011;
 20:1411-21.

Li G, Sun GX, Williams PN, Nunes L, Zhu YG. Inorganic arsenic in Chinese food
 and its cancer risk. *Environ Int* 2011; 37:1219–25.

Li HB, Cui XY, Li K, Li J, Juhasz AL, Ma, LQ. Assessment of in vitro lead
 bioaccessibility in house dust and its relationship to in vivo lead relative
 bioavailability. *Environ Sci Technol* 2014a; 48:8548–55.

Li HB, Li J, Juhasz AL, Ma LQ. Correlation of in vivo relative bioavailability to in
 vitro bioaccessibility for arsenic in household dust from China and its
 implication for human exposure assessment. *Environ Sci Technol* 2014b;
 48:13652–9.

Long SE, Martin TD. Determination of Trace Elements in Waters and Wastes by
 Inductively Coupled Plasma-Mass Spectrometry: Test Method 200.8[M]. US
 Environmental Protection Agency, Office of Research and Development,
 Environmental Monitoring and Systems Laboratory; 1989.

Mason, RP, Laporte JM, Andres S. Factors controlling the bioaccumulation of
 mercury, methylmercury, arsenic, selenium, and cadmium by freshwater
 invertebrates and fish. *Arch Environ Con Tox* 2000; 38:283–97.

622 McClain WR, Romaine RP. *Procambarus clarkii* (Girard, 1852). Cultured Aquatic
623 Species Information Program. In FAO Fisheries and Aquaculture Department
624 [Online]. Rome, Italy: Food and Agriculture Organization; 2007.
625 http://www.fao.org/fishery/culturedspecies/Procambarus_clarkii/en#tcNA00D9
626 [accessed 2 December 2015]

627 Metian M, Charbonnier L, Oberhaensli F, Bustamante P, Jeffree R, Amiard JC,
628 Warnau M. Assessment of metal, metalloid, and radionuclide bioaccessibility
629 from mussels to human consumers, using centrifugation and simulated digestion
630 methods coupled with radiotracer techniques. *Ecotox Environ Safe* 2009;
631 72:1499–502.

632 Metropolis N, Ulam S. The monte carlo method. *J Am Stat Assoc* 1949;
633 44(247):335-41.

634 Moss JC, Hardaway CJ, Richert JC, Sneddon J. Determination of cadmium copper,
635 iron, nickel, lead and zinc in crawfish [*Procambrus clarkii*] by inductively
636 coupled plasma optical emission spectrometry: A study over the 2009 season in
637 Southwest Louisiana. *Microchem J* 2010; 95:5–10.

638 Mu F, Cheng Y, Wu X. Distribution and industrial development of crayfish in the
639 world. *J Shanghai Fish Univ* 2007; 16(1):64–72 [In Chinese].

640 NBSC. China Statistical Yearbook. National Bureau of Statistics of China; 2013.
641 <http://www.stats.gov.cn/tjsj/ndsj/2013/indexeh.htm> [accessed 2 December 2015]

642 NCRPM. A guide for uncertainty analysis in dose and risk assessment related to
643 environmental contamination. Bethesda, MD, USA: National Council on
644 Radiation Protection and Measurements; 1996.

645 Oomen AG, Hack A, Minekus M, Zeijdner E, Cornelis C, Schoeters G, Verstraete W,
646 Van de Wiele T, Wragg J, Rempelberg CJM, Sips A, Van Wijnen JH.

647 Comparison of five in vitro digestion models to study the bioaccessibility of soil
648 contaminants. *Environ Sci Technol* 2002; 36:3326–34.

649 Oomen AG, Rompelberg CJM, Bruil MA, Dobbe CJG, Pereboom D, Sips A.
650 Development of an in vitro digestion model for estimating the bioaccessibility of
651 soil contaminants. *Arch Environ Con Tox* 2003; 44:281–7.

652 Patoka J, Fisakova MN, Kalous L, Skrdla P, Kuca M. Earliest evidence for human
653 consumption of crayfish. *Crustaceana* 2014; 87:1578–85.

654 Peng Q, Greenfield BK, Dang F, Zhong H. Human exposure to methylmercury from
655 crayfish (*Procambarus clarkii*) in China. *Environ Geochem Hlth* 2015; 1–13.

656 Perelló G, Martí-Cid R, Llobet JM, Domingo JL. Effects of various cooking processes
657 on the concentrations of arsenic, cadmium, mercury, and lead in foods. *J Agr*
658 *Food Chem* 2008; 56(23):11262–9.

659 Qiao G, Jiang S. Report on development of the crayfish industry in Jiangsu Province.
660 *Fish Guide to be Rich* 2010; 13:24-6. [In Chinese].

661 Qiu Y, Strid A, Bignert A, Zhu Z, Zhao J, Athanasiadou M, Bergman Å. Chlorinated
662 and brominated organic contaminants in fish from Shanghai markets: A case
663 study of human exposure. *Chemosphere* (2012); 89(4):458–66.

664 Richert JC, Sneddon J. Determination of inorganics and organics in crawfish. *Appl*
665 *Spectrosc Rev* 2008; 43:49–65.

666 Schilderman P, Moonen EJC, Maas LM, Welle I, Kleinjans JCS. Use of crayfish in
667 biomonitoring studies of environmental pollution of the river Meuse. *Ecotox*
668 *Environ Safe* 1999; 44:241-52.

669 Schroder L, Basta NT, Casteel SW, Evans TJ, Payton ME, Si J. Validation of the in
670 vitro gastrointestinal (IVG) method to estimate relative bioavailable lead in
671 contaminated soils. *J Environ Qual* 2004; 33:513–21.

672 Schuler LJ, Howell JP, Heagler MG. Mercury concentrations in Louisiana and
673 Chinese crayfish. *B Environ Contam Tox* 2000; 64:27–32.

674 Sechena R, Liao S, Lorenzana R, Nakano C, Polissar N, Fenske R. Asian American
675 and Pacific Islander seafood consumption — a community-based study in King
676 County, Washington. *J Expo Anal Environ Epidemiol* 2003; 13:256–66.

677 SFEI and CDHS. San Francisco Bay Seafood Consumption Study. Richmond, CA:
678 San Francisco Estuary Institute and California Department of Health Services;
679 2000.

680 Shen Y. Report on the development of the crayfish industry in Jiangsu Province.
681 *China Fish* 2010; 6:14–6. [In Chinese].

682 SAPRC. Maximum Levels of Contaminants in Foods. GB2762-2012. Beijing, China:
683 Standardization Administration of the People’s Republic of China; 2012.

684 Tan S, Zhao L, Guan P. Research on the acute effect of 5 kinds of fishery drugs on
685 crayfish *Procambarus clarkii*. *J Anhui Agr Sci* 2010; 38(33):18852–4. [In
686 Chinese].

687 Taugbøl T, Skurdal J. The future of native crayfish in Europe: How to make the best
688 of a bad situation? p. 271-9 In: Gheraardi F, Holdich DM (Eds.): *Crayfish in*
689 *Europe as alien species: How to make the best of a bad situation*; 1999.

690 Thompson KM, Graham JD. Going beyond the single number: Using probabilistic
691 risk assessment to improve risk management. *Hum Ecol Risk Assess* 1996;
692 2:1008–34.

693 USEPA. Risk Assessment Guidance for Superfund Volume I Human Health
694 Evaluation Manual (Part A) Interim Final. Washington, DC: United States
695 Environmental Protection Agency; 1989.

696 USEPA. Guidance for assessing chemical contaminant data for use in fish advisories.

697 Volume 1. Fish sampling and analysis. 3rd edition. Washington, DC: United
 698 States Environmental Protection Agency; 2000.

699 USEPA. Risk Assessment Guidance for Superfund: Volume III-Part A, Process for
 700 Conducting Probabilistic Risk Assessment. Washington, DC: United States
 701 Environmental Protection Agency; 2001.

702 USEPA. Exposure Factors Handbook 2011 Edition. Washington, DC: National Center
 703 for Environmental Assessment, United States Environmental Protection Agency;
 704 2011.

705 USEPA. Regional Screening Table [Online]. Washington, DC: United States
 706 Environmental Protection Agency; 2015a.
 707 <http://www2.epa.gov/risk/regional-screening-table> [accessed 2 December 2015]

708 USEPA. Integrated Risk Information System (IRIS) 2015b. Washington, DC: United
 709 States Environmental Protection Agency; 2015b. <http://www.epa.gov/iris/>
 710 [accessed 2 December 2015]

711 USFDA. Guidance Documents for Trace Elements in Seafood. Washington, DC:
 712 Center for Food Safety and Applied Nutrition, United States Food and Drug
 713 Administration; 1993.

714 USITC. Crawfish tail meat from China. Investigation No. 731-TA-752 (Review).
 715 Publication 3614. Washington, DC: United States International Trade
 716 Commission; 2003.

717 Vasconcelos V, Oliveira S, Teles FO. Impact of a toxic and a non-toxic strain of
 718 *Microcystis aeruginosa* on the crayfish *Procambarus clarkii*. Toxicon 2001;
 719 39(10):1461–70.

720 Vasconcelos VM. Cyanobacterial toxins in Portugal: Effects on aquatic animals and
 721 risk for human health. Braz J Med Biol Res 1999; 32:249–54.

722 Versantvoort CHM, Oomen AG, Van de Kamp E, Rompelberg CJM, Sips A.
 723 Applicability of an in vitro digestion model in assessing the bioaccessibility of
 724 mycotoxins from food. Food Chem Toxicol 2005; 43:31–40.

725 Vioque-Fernandez A, Alves de Almeida E, Ballesteros J, Garcia-Barrera T,
 726 Gomez-Ariza JL, Lopez-Barea J. Doñana National Park survey using crayfish
 727 (*Procambarus clarkii*) as bioindicator: Esterase inhibition and pollutant levels.
 728 Toxicol Lett 2007; 168:260–8.

729 Wang, WX, Yang Y, Guo X, He M, Guo F, Ke C. Copper and zinc contamination in
 730 oysters: subcellular distribution and detoxification. Environ Toxicol Chem 2011;
 731 30(8):1767–74.

732 Wang Z. Report on the development of crayfish industry in Hubei Province. China
 733 Fish 2010; 10:14-5. [In Chinese].

734 Williams G, West JM, Koch I, Reimer KJ, Snow ET. Arsenic speciation in the
 735 freshwater crayfish, *Cherax destructor* Clark. Sci Total Environ
 736 2009; 407(8):2650–8.

737 Wu C, Liu H, Fang Y, Lu D, Xing Z, Yuan W, Duan S, Qin L. Survey on the content
 738 and intake of lead and cadmium in crayfish from market in Shanghai. J Environ
 739 Occup Med 2010; 27(11):650–6. [In Chinese].

740 Xie W, Chen K, Zhu X, Nie X, Zheng G, Pan D, Wang S. Evaluation on heavy metal
 741 contents in water and fishes collected from the waterway in the Pearl River Delta,
 742 South China. J Agro- Environ Sci 2010; 29:1917–23. [In Chinese].

743 Yi Y, Yang Z, Zhang S. Ecological risk assessment of heavy metals in sediment and
 744 human health risk assessment of heavy metals in fishes in the middle and lower
 745 reaches of the Yangtze River Basin. Environ Pollut 2011; 159:2575–85.

746 Yue CF, Wang TT, Wang YF, Peng Y. Effect of combined photoperiod, water calcium

747 concentration and pH on survival, growth, and moulting of juvenile crayfish
748 (*Procambarus clarkii*) cultured under laboratory conditions. Aquac
749 Res 2009; 40(11):1243–50.

750 Zhang B, Yang G, Yu X, Mao H, Xing C, Liu J. Haff disease after eating crayfish in
751 East China. Internal Med 2012; 51:487–9.

752 Zhang H, Feng X, Larssen T, Qiu G, Vogt RD. In inland China, rice, rather than fish,
753 is the major pathway for methylmercury exposure. Environ Health Persp 2010;
754 118(9): 1183–8.

755

Table 1. Estimated oral reference doses (RfD, mg/kg/d) and cancer slope factor (CSF, (mg/kg/d)⁻¹) for metals (USEPA, 2015a, b), daily intake for crayfish consuming populations in China (EDI, mg/kg/d), hazard quotient (HQ, unitless), hazard index (HI, unitless), and increased lifetime cancer risk (ILTR, unitless).

		Metal						
		As	Cd	Cu	Ni	Pb	Se	Zn
CSF	RfD	3×10 ⁻⁴	1×10 ⁻³	0.04	0.02	0.02 ^a	0.005 ^a	0.3
CSF		1.5	NA	NA	NA	NA	NA	NA
EDI	P50	1.7×10 ⁻⁵	9.9×10 ⁻⁷	5.8×10 ⁻⁴	8.3×10 ⁻⁶	5.7×10 ⁻⁹	1.8×10 ⁻⁵	1.2×10 ⁻³
	P90	1.2×10 ⁻⁴	1.1×10 ⁻⁵	4.2×10 ⁻³	7.5×10 ⁻⁵	1.2×10 ⁻⁷	1.2×10 ⁻⁴	7.8×10 ⁻³
	Range	5.5×10 ⁻³	1.2×10 ⁻³	0.19	3.6×10 ⁻³	8.6×10 ⁻⁶	4.8×10 ⁻³	0.26
HQ	P50	0.057	9.9×10 ⁻⁴	0.015	4.2×10 ⁻⁴	2.8×10 ⁻⁷	3.5×10 ⁻³	4.1×10 ⁻³
	P90	0.40	1.1×10 ⁻²	0.105	3.7×10 ⁻³	6.2×10 ⁻⁶	0.024	0.029
	Range	18.3	1.2	4.7	0.18	4.8×10 ⁻⁴	0.97	0.87
HQ/HI (p90)		0.70	0.02	0.18	0.01	1.1×10 ⁻⁵	0.04	0.05
HI	P50				0.089			
	P90				0.572			
	Range				24			
ILTR	P50	2.5×10 ⁻⁵						
	P90	1.8×10 ⁻⁴			NA			
	Range	8.2×10 ⁻³						

NA: not applicable.

^a The value corresponds to the Minimal Risk Level.

Table 2. Trace element concentrations (wet weight) in tail muscle of cooked crayfish (*Procambarus clarkii*) sampled in restaurants in China.Mean $\pm$ standard deviation (SD).

City	Province	Sample size	As $\mu\text{g/kg}$	Cd $\mu\text{g/kg}$	Cu mg/kg	Ni $\mu\text{g/kg}$	Pb $\mu\text{g/kg}$	Se $\mu\text{g/kg}$	Zn mg/kg
Nanjing	Jiangsu	26	259.8 ± 121.0	43.8 ± 35.0	11.1 ± 8.2	211.1 ± 89.9	18.0 ± 8.4	254.6 ± 82.8	18.2 ± 3.4
Suqian	Jiangsu	9	450.3 ± 125.5	46.8 ± 32.6	16.1 ± 7.0	274.1 ± 209.5	44.5 ± 49.5	173.1 ± 83.7	17.0 ± 2.2
Lianyungang	Jiangsu	9	174.7 ± 39.3	44.8 ± 58.4	13.1 ± 5.0	199.3 ± 123.3	16.0 ± 10.1	169.7 ± 61.8	14.6 ± 2.8
Suzhou	Jiangsu	9	237.1 ± 93.3	41.1 ± 19.9	6.1 ± 1.6	74.5 ± 39.3	21.7 ± 16.1	229.7 ± 94.8	16.4 ± 1.6
Huai'an	Jiangsu	9	300.5 ± 112.6	30.0 ± 37.0	8.7 ± 3.8	109.1 ± 35.8	31.3 ± 20.9	167.3 ± 51.7	16.1 ± 4.6
Yangzhou ^a	Jiangsu	2	237.9 ± 13.0	20.2 ± 0.1	3.7 ± 1.7	98.8 ± 18.0	ND ^b	159.2 ± 31.2	23.2 ± 0.4
Yiyang	Hunan	9	211.2 ± 42.1	30.0 ± 1.7	6.0 ± 1.4	62.5 ± 69.2	1.7 ± 3.8	614.2 ± 80.4	22.1 ± 2.1
Bengbu	Anhui	9	147.4 ± 45.3	25.3 ± 0.7	7.9 ± 2.4	64.5 ± 14.3	12.3 ± 19.4	236.8 ± 93.3	23.5 ± 2.8
Suzhou	Anhui	9	128.8 ± 17.9	24.7 ± 3.3	14.2 ± 5.5	272.0 ± 148.4	49.1 ± 27.8	234.9 ± 36.0	21.3 ± 2.0

Chuzhou	Anhui	9	274.6 ± 67.3	33.8 ± 4.3	11.4 ± 5.8	268.5 ± 45.9	16.4 ± 25.5	328.4 ± 112.2	30.6 ± 3.5
Anqing ^a	Anhui	6	95.5 ± 29.4	45.5 ± 49.4	16.7 ± 8.7	101.4 ± 35.3	19.6 ± 30.9	205.7 ± 31.4	26.0 ± 4.0
Wuhan	Hubei	9	412.2 ± 155.3	38.2 ± 2.0	5.5 ± 1.3	337.3 ± 72.8	ND ^b	222.1 ± 41.1	36.9 ± 5.4
Huang gang	Hubei	9	331.2 ± 105.2	37.0 ± 11.3	12.7 ± 5.4	181.6 ± 56.2	3.8 ± 11.3	215.3 ± 34.8	28.1 ± 2.7
Hangzhou	Zhejiang	9	215.2 ± 55.5	2.8 ± 1.7	6.1 ± 2.4	76.5 ± 54.3	74.8 ± 74.0	331.7 ± 150.6	17.3 ± 2.3
Anji	Zhejiang	9	105.1 ± 16.3	2.8 ± 0.4	5.7 ± 1.8	65.2 ± 16.0	30.4 ± 6.1	408.6 ± 93.3	17.4 ± 0.8
Zigong ^a	Sichuang	5	168.8 ± 83.7	38.1 ± 7.0	3.5 ± 1.2	73.8 ± 43.3	41.1 ± 62.3	249.5 ± 106.5	27.5 ± 1.7
Anyang	Henan	9	552.4 ± 186.8	16.9 ± 10.3	8.7 ± 3.2	164.8 ± 49.5	41.1 ± 65.1	203.9 ± 62.3	13.6 ± 1.4
Xinyang	Henan	9	113.0 ± 37.3	1.2 ± 0.4	6.2 ± 2.4	71.9 ± 17.1	19.8 ± 7.3	241.4 ± 18.9	16.3 ± 1.6
Beijing	Beijing	9	193.8 ± 33.7	3.1 ± 0.9	3.4 ± 1.1	173.8 ± 80.5	34.3 ± 7.3	274.0 ± 104.2	20.2 ± 1.1
ZaoZhuang	Shandong	9	249.0 ± 92.0	12.6 ± 6.6	4.4 ± 2.4	293.4 ± 182.9	21.5 ± 3.6	703.0 ± 215.2	14.4 ± 0.8
Haerbin	Heilongjiang	9	335.1 ± 90.2	32.0 ± 15.9	11.7 ± 4.2	93.8 ± 43.5	13.3 ± 11.3	206.4 ± 39.7	15.1 ± 2.4
Xi'an	Shanxi	9	284.6 ± 148.3	29.4 ± 48.9	8.9 ± 2.0	375.4 ± 307.7	12.6 ± 11.5	258.4 ± 98.8	16.7 ± 3.6

Guiyang	Guizhou	9	234.0 ± 105.7	60.9 ± 137.3	12.2 ± 4.8	91.5 ± 133.3	11.0 ± 6.7	385.1 ± 56.1	20.0 ± 3.6
Mean	-	210	253.3 ± 142.4	29.8 ± 39.2	9.2 ± 5.8	171.2 ± 141.2	23.3 ± 29.5	294.4 ± 154.1	20.1 ± 6.3
Safety standards	-	-	5000 ^{c,d}	500 ^c	- ^e	- ^e	500 ^c	- ^e	- ^e

^a Fewer crayfish individuals were obtained due to limited sample availability.

^b ND = not detected; individual samples that were ND were set at quantification limit (0.15 µg/kg) for generating means.

^c Referring to GB2762-2012 (SAPRC, 2012).

^d Assuming that 10% of As is inorganic As (Buchet et al., 1996; USFDA, 1993).

^e Cu, Zn, Ni and Se were not listed in the new national standard (SAPRC, 2012).

Table 3. Parameters used in the risk calculation.

Variable	Definition	Statistical distribution	Origin of information
IR (g/day)	Ingestion rate ^a	Custom distribution: see text	Peng et al., 2015
ED (year)	Exposure duration	Constant = 75	Life expectancy (NBSC, 2013c)
EF (day)	Exposure frequency	Constant = 120	Duration of peak season (Peng et al., 2015)
BW	Body weight	Custom distribution: see text	Li et al., 2011
AT	Averaging time	Constant = 75 year×365 days/year	USEPA, 1989
C _i (mg/kg)	Metal concentration	Lognormal	This study (Table 2)
φ _i (%) ^b	Bioaccessibility (in vitro extraction model)	As N (95.4; 6.2) Cd N (78.7;6.1) Cu N (92.5;2.9) Ni N (89.2;17.1) Pb 100 ^c Se N (82.6;12.6) Zn LN (77.6;16.5)	This study (Figure 1; Table S2)

N: Normal distribution (mean; SD), truncated to values > 0 when needed; LN:

Lognormal distribution (mean; SD)

^a Ingestion rates were averaged and reported as daily intakes (more details in Peng et al., 2015).

^b Outliers were removed (more details in Table S2).

^c Lead bioaccessibility was not determined because extracted Pb levels were too low to be precisely measured.

Table 4. Model sensitivity to input uncertainty. Contribution to variance (%).

Parameters	Risk model	
	Noncancer hazard index (HI)	Increased lifetime cancer risk (ILTR)
IR	95.3	92.5
As concentration	3.4	6.5
BW woman	0.5	0.4
BW man	0.4	0.4
Cu concentration	0.3	-
φ_{As}	< 0.1	0.1
Cd, Ni, Pb, Se, Zn concentration	< 0.1	-
φ_{Cd} , φ_{Cu} , φ_{Ni} , φ_{Se} , φ_{Zn}	< 0.1	-

BW: body weight

Figure captions

Figure 1. Element bioaccessibility (ϕ_i) using the in vitro extraction model described by Versantvoort et al. (2005). Mean $\pm$ standard deviation (SD).

Figure 2. Histogram of ingestion rate (IR) during peak season, based on survey results described in Peng et al. (2015).

Figure 1

Figure 2

Q: Are consumers at risk due to exposure to metals in crayfish? **A:** Arsenic poses health risk, especially for high rate consumers.