1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29

Preliminary analysis of beak stable isotopes ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$) stock
variation of neon flying squid, *Ommastrephes bartramii*, in the North
Pacific Ocean

Zhou Fang^{1,5,4}, Katherine Thompson⁴, Yue Jin^{1,5}, Xinjun Chen^{1,2,3,5},
Yong Chen^{4,5}

- 1. College of Marine Sciences, Shanghai Ocean University, Shanghai 201306, China;
- 2 National Engineering Research Center for Oceanic Fisheries, Shanghai Ocean University, Shanghai 201306, China;
- 3. The Key Laboratory of Sustainable Exploitation of Oceanic Fisheries Resources, Ministry of Education, Shanghai Ocean University, Shanghai 201306, China
- 4. School of Marine Sciences, University of Maine, Orono, Maine 04469, USA
- 5. Collaborative Innovation Center for Distant-water Fisheries, Shanghai 201306, China

Tel.:021-61900306; Fax:021-61900304
E-mail: xjchen@shou.edu.cn

Abstract

Stable isotopes (^{13}C and ^{15}N) have been confirmed as a useful tool for understanding trophic position and related dietary variation in squid. In this study, we performed isotopic analysis on *Ommastrephes bartramii* beaks to examine differences between the eastern and western stocks in the North Pacific Ocean. Isotopic values of the upper beak (UB) and the lower beak (LB) were also compared. A generalized additive model (GAM) was used to select variables that explain stock variation. Isotopic signatures ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$) were significantly different between the two stocks whereas there was no difference in C/N ratios. All isotopic values were significantly different between the UB and LB. Trophic niche width was distinct between two stocks with little overlap. The $\delta^{13}\text{C}$ signature increased with latitude and mantle length (ML) with greater variability in the eastern stock. None of the variables could explain the variation in $\delta^{13}\text{C}$ values for the western stock. The $\delta^{15}\text{N}$ signature increased rapidly with ML in the eastern stock, whereas $\delta^{15}\text{N}$ gradually increased with latitude and ML in the western stock. The isotopic variations can be successfully explained by different migration patterns and feeding behaviors of the two stocks. The $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of the UB were lower than those of the LB, possibly due to variable chitin/protein ratios at different developmental stages. Future studies should account for a greater sample size and the functionality of the UB.

Keywords: Beak; stable isotope; ontogeny; stock variation; *Ommastrephes bartramii*; GAM

Introduction

Cephalopods, especially ommastrephid squids, play an important role in the marine food web by connecting the higher and lower trophic levels, acting as both predator and prey (Rodhouse and Nigmatullin, 1996; Navarro et al., 2013). Commercial oceanic (ommastrephid) squid landings increased drastically over the past two decades, from approximately 2.0 million metric tons in 1990s to over 4.3 million metric tons in 2010 (FAO, 2013). The neon flying squid, *Ommastrephes bartramii*, which can reach large body sizes and yield high economic value, is widely distributed in temperate and subtropical waters of the Pacific, Indian and Atlantic Oceans, but is commercially exploited only in the North Pacific Ocean with some Asian countries (Bower and Ichii 2005; Chen et al., 2008; Jereb and Roper, 2010).

Some studies have been discussed about the population structure of *O. bartramii* based on mantle length distribution and rates of infection by helminth parasites (Murata, 1990; Bower and Margolis, 1991; Nagasawa et al., 1998; Yatsu et al., 1998). It is generally accepted that this species is believed to consist of two main cohorts, the autumn cohort that is mainly distributed east of 170°E (Eastern stock) with fishing season from May to June and the winter-spring cohort located west of 170°E (Western stock) with fishing season from July to November (Bower and Ichii, 2005). These two cohorts have been confirmed as two different stocks with major genetic differentiation, although they are overlapping geographically in some areas (Katugin, 2002; Bower and Ichii, 2005). Geographic separation of the two cohorts causes them to experience very different oceanographic conditions, which leads to asynchronous growth patterns

and varied abundances (Chen and Chiu, 2003; Chen, 2010).

Oceanic squid, tend to migrate long distances in north-south direction during their lifetime and may be exposed to different environments in each life phase (Yatsu et al., 1997; Murata and Nakamura, 1998; Ichii et al., 2004). Optimum oceanographic environment and food richness are the two main reasons cause the migration (Ichii et al. 2009). Dietary change is also closely related with the squid growth, especially adult squid (Watanabe et al., 2004; 2008) Feeding behavior tends to be the dominant factor for the migration with the body size increase in adult squid. With voracious feeding habits and a fast metabolism, squid effectively contribute to energy exchange in the marine food web (Clarke, 1996; Piatkowski et al., 2001; Boyle and Rodhouse 2008). Examining their feeding habits is important to elucidating the trophic role of squid in the marine ecosystem. Stomach content analysis is a conventional approach that directly examines feeding habits (Rodhouse and Nigmatullin, 1996; Watanabe et al., 2004; Cherel and Hobson, 2005). This time-consuming method with some obvious disadvantages was replaced by some indirect approach (Ruiz-Cooley et al., 2006; Jackson et al., 2007). Stable isotope analysis (SIA), specifically analysis of carbon and nitrogen isotopes (^{13}C and ^{15}N), has been successfully used in several studies of squid feeding habits over the last decade (Takai et al., 2000; Ruiz-Cooley et al., 2004, 2006; Lorrain et al., 2011; Argüelles et al., 2012; Miller et al., 2013). The trophic position of a species can be identified by quantifying the nitrogen isotope ratios $\delta^{15}\text{N}$ ($^{15}\text{N}/^{14}\text{N}$) (McCutchan et al., 2003; Vanderklift and Ponsard, 2003). Carbon isotope ratios $\delta^{13}\text{C}$ ($^{13}\text{C}/^{12}\text{C}$) reflects variability in primary production and

trophic dynamics along migration routes (Rubenstein and Hobson, 2004). Stable isotope ratios ($\delta^{15}\text{N}$ and $\delta^{13}\text{C}$) of animal tissues provide a record of historic diet composition (Rau et al., 1983; Hobson and Welch, 1992; 1994). Therefore, it is possible to obtain relatively long-term feeding information, which is difficult to do with conventional method (Onthank, 2013).

Different tissues have been analyzed for isotopic signatures $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$, the most common of which is muscle due to its easy availability (Ruiz-Cooley et al., 2004, 2006; Stowasser et al., 2006; Parry, 2008; Cherel et al., 2009a; 2009b; Hunsicker et al., 2010; Argüelles et al., 2012). Unlike archival tissues, muscle is continuously regenerating with fast cell metabolism and protein synthesis (Cherel et al., 2009b; Moltschaniwskyj and Carter, 2010), so historic nutrient records are not retained (Onthank, 2013). Additionally, muscle has high lipid content yielding negative $\delta^{13}\text{C}$ values, which alter the absolute $\delta^{13}\text{C}$ value and confound interpretation of foraging behavior dynamics (DeNiro and Epstein, 1977; Focken and Becker, 1998; Logan et al., 2008; Ruiz-Cooley et al., 2011). As a main feeding organ, the beak has very low lipid content and it has been verified that the beak has a slightly higher ^{13}C content but lower ^{15}N content compared to lipid-free muscle (Cherel and Hobson, 2005; Hobson and Cherel, 2006; Cherel et al., 2009b). The growth of beak is deposited with continuous layers of cells (Raya et al., 2010), so different parts of beak can represent different life stages (Cherel et al., 2009b; Guerra et al., 2010). Thus, beak is confirmed as a useful archival tissues that can reflect long-term ecological and life history information to examine habitat and trophic variation (Cherel and Hobson,

2005; Hobson and Cherel, 2006; Ruiz-Cooley et al., 2006; Hunsicker et al., 2010; Ruiz-Cooley et al., 2013; Xavier et al., 2014).

In this study, we used quantitative beak stable isotopic data ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$) to estimate the different trophic level of *O. bartramii* between two stocks with pertinent dietary-induced migration patterns. We also used a generalized additive model (GAM) to analyze the main factors influencing isotopic variation in different life stages. The isotopic difference between upper and lower beaks was also analyzed. This study improves our understanding of the stock variation pattern in feeding behavior and tests different migration routes of two *O. bartramii* stocks in the North Pacific Ocean.

Materials and Methods

A total of 60 random squid samples were obtained from the Chinese commercial jigging vessel *F/V Jinhai 827* operated on two main fishing grounds (152°E-174°W and 38°-45°N) in the North Pacific Ocean from May to November in 2010 (Fig.1). Samples were immediately frozen at -18°C for laboratory work. Samples from different areas were treated as different stocks based on age back-calculation analysis, in which eastern stock and western stock belong to autumn cohort and winter-spring cohort, respectively (Table 1 and 2)(Jin et al., 2015; Fang et al., 2016).

Dorsal mantle length (ML) was measured to the nearest 1 mm in the laboratory. The maturity stage was also determined according to Lipinski and Underhill (1995). The upper (UB) and lower beaks (LB) were dissected from the buccal mass, washed with fresh water and stored in 75% ethyl alcohol, which does not change the isotopic

composition of tissues and prevents beaks from dehydrating (Hobson et al., 1997).

Before isotopic analysis, beaks were cleaned with MilliQ water for at least 5 min and dried in a freeze dryer (Alpha 1-4 LD Plus, Christ, Germany) for 24 h. Each beak sample was then ground to a fine powder with a Mixer Mill MM 400 (Retsch GmbH, Haan, Germany) for 1.5 min. Approximately 1.50 ± 0.05 mg of each beak powder sample was then weighed in an aluminum tray using an electronic scale (BSA223S, Sartorius, Göttingen, Germany). Samples were analyzed with an ISOPRIME 100 isotope ratio mass spectrometer (Isoprime Corporation, Cheadle, UK) and a vario ISOTOPE cube elemental analyzer (Elementar Analysensysteme GmbH, Hanau, Germany) in the Key Laboratory of Sustainable Exploitation of Oceanic Fisheries Resources, Ministry of Education, at Shanghai Ocean University.

The standard reference material for carbon was Vienna Pee Dee Belemnite (V-PDB) and for nitrogen was atmospheric nitrogen (N_2). USGS 24 (-16.049‰ V-PDB) was used as primary standard for ^{13}C and USGS 26 (53.7‰ V- N_2) was used to quantify ^{15}N . To assess the associated errors within and between runs, repeated analyses of internal laboratory standards (-26.98‰ vPDB and 5.96‰ v N_2) were performed every 10 samples. The analytical precision was less than 0.1‰ both for carbon and nitrogen. The $\delta^{13}C$ and $\delta^{15}N$ of samples are expressed in standard notation as the following (Ruiz-Cooley et al., 2006):

$$\delta X(\text{‰}) = \left(\frac{R_{\text{sample}}}{R_{\text{standard}}} - 1 \right) \times 1000$$

where R_{sample} and R_{standard} are the ratios of $^{13}C/^{12}C$ or $^{15}N/^{14}N$ in the sample and standards, respectively. δX is the measure of the ratio of heavy to light isotope ratio in

the sample.

All the isotopic values had constant variance and conformed to normal distribution. We used an independent two-group t-test to analyze the $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ variation between the two stocks. Differences of isotopic values between upper and lower beaks were also evaluated with a paired t-test. Convex hull of carbon–nitrogen biplot was also constructed to evaluate the trophic niche width for two stocks. Variable geographic ranges of the two stocks may cause isotopic variation, so the distance to the shelf break (DSB) was calculated for each sample. We used R to calculate DSB with function “dist2Line” in the package of “geosphere”, which calculates the minimum distance between a specific point to the nearest continental shelf border along a great circle (shortest distance on a sphere) (Hijmans, 2014). It is possible that the relationships between two dependent variables ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$) and three independent variables (latitude (LAT), ML, DSB) are nonlinear and multivariate (Argüelles et al., 2012), thus generalized additive models (GAM) were used to analyze isotopic variation in different situation (Wood, 2006; Argüelles et al., 2012). The best fitted model was the one with the lowest AIC (Akaike information criterion) value (Akaike, 1974). Relationships were developed for the upper and lower beaks and isotopic values ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$) were compared between stocks.

Results

Isotopic variation between two stocks

Isotopic values of western stock were higher than those of eastern stock (Table 1

and 2). The independent two-group t-test results indicated that ML was significantly different between the two geographical stocks ($P < 0.05$; Table 3). Isotopic values were also significantly different between stocks ($P < 0.05$; Table 3). There was no difference between the C/N ratio of the UB and LB ($P > 0.05$; Table 3). Isotopic values of UB were also lower than those of LB (Table 1 and 2) and these values were all significantly different between the two stocks ($P < 0.05$; Table 3).

The considerable overlapped trophic niche width of two stocks was observed from the biplot (Fig. 2A). Western stock had more concentrated trophic niche width with higher $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values (Fig. 2A). In the eastern stock, the trophic niche width was changed substantially from low $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ to high $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values (Fig. 2A). This transformation was also companied with the ML increase.

Considering the disparity of trophic niche width of eastern stock for different body size, two subgroups were established to evaluate the variation of eastern stock in different sizes. It is obvious that eastern large squids (ML > 350mm, mean ML 418.1mm) have similar trophic level (mean $\pm$ SD Upper $\delta^{13}\text{C}$: -18.2 ± 0.2 , $\delta^{15}\text{N}$: 8.5 ± 0.5 ; Lower $\delta^{13}\text{C}$: -17.8 ± 0.1 , $\delta^{15}\text{N}$: 9.1 ± 0.6) with those of western stock (mean ML 274.4mm; Upper $\delta^{13}\text{C}$: -18.0 ± 0.3 , $\delta^{15}\text{N}$: 8.8 ± 0.8 ; Lower $\delta^{13}\text{C}$: -17.8 ± 0.2 , $\delta^{15}\text{N}$: 9.1 ± 0.7) (Fig. 2B). The eastern small squids (ML < 350mm, mean ML 262.8mm) presented a very distinct trophic level with a low value (Upper $\delta^{13}\text{C}$: -18.6 ± 0.2 ; $\delta^{15}\text{N}$: 6.0 ± 0.9 ; Lower $\delta^{13}\text{C}$: -18.3 ± 0.3 ; $\delta^{15}\text{N}$: 6.5 ± 0.8) compared to large sized groups (Fig. 2B).

Isotopic variation among different ML groups

The $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values varied between ML groups (Fig. 3). For a given stock, the $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values increased with ML by 0.5 – 0.8‰ for $\delta^{13}\text{C}$ and 3 – 3.5‰ for $\delta^{15}\text{N}$ in the eastern stock (Fig. 3). In the western stock, $\delta^{15}\text{N}$ values also changed by 1.5 – 2‰ with increasing ML, but $\delta^{13}\text{C}$ values changed little between ML groups (Fig. 3). The isotopic values in LB were slightly higher than those in UB (Fig. 3).

$\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values increased rapidly from ML 250 – 300 mm to 350 – 400 mm in the eastern stock (Fig. 3). Isotope values plateaued in ML 350 – 400 mm and > 400 mm for the eastern stock. In the western stock, there was a gradual increase in $\delta^{15}\text{N}$ values from ML > 200 mm to 250 – 300 mm, whereas values leveled off in ML 300 – 350 mm and > 350 mm (Fig. 3).

GAMs selection and simulation

ML was identified as the single most important variable in all GAMs (Table 4). There was no difference between UB and LB in both the stocks for the variables identified as significant in the GAMs (Table 4). The GAM models explained from 23.8% to 91.6% of the variation in $\delta^{13}\text{C}$ or $\delta^{15}\text{N}$ values (Table 5).

In the eastern stock, latitude and ML were the two main effects for $\delta^{13}\text{C}$ values of UB and LB (Fig. 4). The $\delta^{13}\text{C}$ values decreased with latitude and increased with ML with little variability (Fig. 4). ML was also the only variable that had significant effects on $\delta^{15}\text{N}$ values and $\delta^{15}\text{N}$ increased with ML (Figs. 4C and 4F). An isotopic plateau could be seen between ML 350 and 400mm (Fig. 4).

In the western stock, all of the three isotope variables failed to explain $\delta^{13}\text{C}$ values in the GAMs for both UB and LB (Table 5). The $\delta^{15}\text{N}$ values increased linearly with latitude. A similar result was observed for $\delta^{15}\text{N}$ with $\delta^{15}\text{N}$ values gradually increasing with ML and then reaching a plateau between ML 300 and 350 mm (Fig. 5).

Discussion

Intraspecific variation is usually affected by both biotic factors (e.g., intrinsic genetic structure) and abiotic factors (e.g., environmental conditions) (Pecl et al., 2004; Sandoval-Castellanos et al., 2007). Variation between stocks may be due to asynchronous growth, disparity in migration routes and differences in diet composition, which has been shown in other ommastrephid squids (Crespi-Abril and Baron, 2012; Liu et al., 2013; Arbuckle and Wormuth, 2014). Trophic niche width of the two stocks was separate except eastern large individuals (Fig. 2). Different feeding habitat of the two areas is the main reason to impact this disparity. Watanabe et al. (2008) indicated that winter-spring cohort mainly prey micronektonic squid (*Watasenia scintillans*) and Japanese anchovy (*Engraulis japonicas*) when stay in the subarctic frontal zone (SAFZ) and prepare southward migration during the late autumn and early spring (Fig. 1). The migration of prey species was also influenced by the variation of Kuroshio–Oyashio (Takahashi et al., 2001). Adults of autumn cohort mainly fed on the transitional-water myctophid (*Symbolophorus californiensis*), squid (*Onychoteuthis borealijaponica*) and subtropical water myctophids (*Ceratoscopelus warmingii*), squid (*Gonatus berryi* and *Berryteuthis anonychus*)

(Watanabe et al., 2004), which has similar trophic level with the preys for western stock. However, juveniles fed on zooplankton, planktonic crustaceans (Euphausiids and amphipods) actively (Watanabe et al., 2004), which has lower trophic level than fish and squid. And this latitude is also the suitable habitat for the male squids of eastern stock always staying in the spawning/nursery ground (subtropical frontal zone, STFZ). This is the main reason why the feeding niche width distinct with little overlap for two stocks, as well as the different size for eastern stock (Fig. 2).

Marine phytoplankton $\delta^{13}\text{C}$ decreases from the equator poleward at different rates between the southern and northern hemispheres (Rau et al., 1982; Takai et al., 2000). Lower $\delta^{13}\text{C}$ values at higher latitudes in the eastern stock suggest that variation of phytoplankton $\delta^{13}\text{C}$ could be reflected in squid muscle (Fig. 3a, Takai et al., 2000). Latitudinal variation in the western stock failed to explain $\delta^{13}\text{C}$ variability, even though the western stock has a greater latitudinal range than the eastern stock (Table 1). The $\delta^{13}\text{C}$ signature of the western stock may also be affected by proximity to shore, since the western stock is closer to the shelf break than the eastern stock (Fig. 1). There seem to be more factors involved in $\delta^{13}\text{C}$ dynamics of the western stock, which should be considered in future analysis.

ML is another factor influencing $\delta^{13}\text{C}$ values (Fig. 3A). Oceanic squid often experience multiple environments along large-scale migration throughout their lifetime. All of the individuals less than 300 mm ML are males in this study. Females and males share the same habitat until they reach ~25 cm ML. During gametogenesis, females in the autumn cohort migrate north to the transition zone (TZ) and then to the

SAFZ, while males remain in the spawning/nursery ground (STFZ) (Fig. 1) (Ichii et al., 2009). Once mature, females migrate south and meet males for breeding in the STFZ (Chen and Chiu, 2003; Ichii et al., 2009). This explains the sharp decrease of $\delta^{13}\text{C}$ in small squid and rapid increase of $\delta^{13}\text{C}$ in large squid, due to higher Chl-a concentration at the TZ and SAFZ compared to the less productive STFZ (Seki et al., 2002; Ichii et al., 2004). Increasing $\delta^{13}\text{C}$ values of squid at ML 350 – 400 mm and over 400 mm can be explained by the autumn cohort squid of this size swimming south to the STFZ, thus experiencing lower Chl-a concentration and decreasing lower $\delta^{13}\text{C}$ levels throughout their migration route (Figs. 3A, 4B, 4E).

ML had no effect on $\delta^{13}\text{C}$ variation of the western stock. The $\delta^{13}\text{C}$ values of the western stock were higher than eastern stock values for every ML group (Fig. 3A). The winter-spring cohort (western stock in this study) inhabits an area with relatively low Chl-a concentration in their early life stages and then migrates north to an area of high productivity (Ichii et al., 2009). Thus, this stock tends to consist of small-sized individuals. The earlier that these squid mature, the longer they inhabit the more productive area. DSB is also a potential factor contributing to $\delta^{13}\text{C}$ differences between stocks, although it was not significant in the GAM. A larger sample size is needed to explore its effect in future analysis.

Nitrogen isotopes (^{15}N) have been shown to reflect latitudinal variation in phytoplankton isotopes (Wada and Hattori, 1990). Maximum nitrate concentrations are easily reached at relatively low levels compared with other nutrient saturation concentrations, although more nitrate rises to the euphotic layer by vertical water

293 mixing at high latitudes (Takai et al., 2000). The majority of squid from the western
294 stock in this study were caught in the nitrate-rich water off Kuril Island where
295 significant isotope fractionation results in the low $\delta^{15}\text{N}$ values of this area (Fig.1,
296 Takai et al., 2000). Isotope fractionation is also reduced by fast utilization of inorganic
297 nitrogen compounds, so the phytoplankton often has lower $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ at higher
298 latitudes than at lower latitudes. However, the negative relationship between latitude
299 and isotopic variation is weak and may deteriorate with trophic variation, which
300 differed by 1.2‰ in $\delta^{15}\text{N}$ compared to 0.05‰ in $\delta^{13}\text{C}$ among size groups (Fig. 3B).
301 The diet of the winter-spring cohort changes from zooplankton to fish and squid as
302 they grow along their northern migration (Watanabe et al., 2004). This may explain
303 why $\delta^{15}\text{N}$ values increased with latitude in the western stock. The autumn cohort also
304 migrates north while maturing into adult squid (Ichii et al., 2009). The eastern stock
305 may have similar $\delta^{15}\text{N}$ variation, but it is possible that latitude was not significant in
306 this analysis due to low sample size.

307 The $\delta^{15}\text{N}$ signature tends to increase abruptly with ML in the eastern stock (Figs.
308 3B, 4C, 4F). The rate of $\delta^{15}\text{N}$ increase is about 1‰ between ML groups except for
309 between ML groups D and E (ML 350 – 400 mm and ML > 400 mm) (Figs. 3B). The
310 overall $\delta^{15}\text{N}$ difference was approximately 3‰, which represents nearly a trophic
311 level (Mingawa and Wada, 1986; Petersen and Fry, 1987). During the northern
312 migration of autumn cohort females from STFZ to SAFZ, the main prey items are fish
313 and squid based on stomach content analyses (Watanabe et al., 2004; Bower and Ichii
314 2005). Meanwhile, males still living in the STFZ prey on lower trophic species such

as euphausiids and amphipods. Therefore, sexual differences and mantle size variation in feeding behavior lead to trophic variation between size classes. This may also explain similar patterns in the western stock. A $\delta^{15}\text{N}$ plateau was also observed in both stocks at different MLs, which is discussed in a previous study of the same species caught off Hawaii (Parry, 2008). The leveling off of $\delta^{15}\text{N}$ values is observed in squid undergoing reproductive maturation. In this phase, somatic growth yields to reproductive growth, resulting in less nitrogen incorporated into other tissues (Parry, 2008). This pattern results from squid reproductive maturation and energy partitioning rather than migration dynamics.

Interestingly, an increasing trend continued after the plateau in $\delta^{15}\text{N}$ values for both stocks (Figs. 4 and 5). Parry (2008) analyzed the muscle of *O. bartramii* in waters off Hawaii and observed $\delta^{15}\text{N}$ values did not change after 350 to 400mm ML. This suggested that *O. bartramii* reached peak (maybe maximum) $\delta^{15}\text{N}$ concentrations when values became constant (Parry, 2008). In this study we analyzed beaks rather than muscle and reported low $\delta^{15}\text{N}$ values in the beak, which mainly consists of chitin, compared to 3-4‰ $\delta^{15}\text{N}$ in muscle (Schimmelmann and DeNiro, 1986; Hobson and Cherel, 2006; Cherel et al., 2009b). Mean maximum $\delta^{15}\text{N}$ values of muscle are nearly 15-16‰ (Parry, 2008), whereas mean maximum $\delta^{15}\text{N}$ values of the beak are about 11-12‰. Chitin tends to have lower $\delta^{15}\text{N}$ content than protein content, since protein is the main chemical structure in the pigmented part of the beak (Miserez et al., 2008). Beak pigmentation is related to the maturation process and occurs over a very short period of the squid life cycle (García, 2003). The maturation process is accompanied

by protein enrichment due to rapid beak pigmentation, leading to the $\delta^{15}\text{N}$ values in larger squid reaching a potential threshold.

We also compared the isotopic signatures between UB and LB in this study. $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values were significantly different between UB and LB and the UB exhibited lower values than the LB in both stocks (Fig. 3). The opposite scenario was observed in a previous study (Hobson and Cherel, 2006). Differences in chitin content between the UB and LB may be the most important factor influencing isotopic signatures. The UB size is larger than the LB's, so smaller individuals have more chitin structure resulting in a higher proportion of enriched $\delta^{15}\text{N}$ pigmented parts in the LB than the UB. This is reversed for larger squid with a darker, more pigmented UB. Early beak development may also have an impact on the isotopic signature in the UB and LB (Boletzky, 2007; Uchikawa et al., 2009). While the beak is an effective archival tissue to explore isotopic variation representing different feeding habitats and trophic levels, the LB is the best structure for such analysis (Cherel and Hobson, 2005; Hobson and Cherel, 2006; Ruiz-Cooley et al., 2006; Cherel et al., 2009b). In order to study the rare or deep-sea species, undigested beaks from stomachs of larger fish or marine mammals is the only material for the related investigation. The UB and LB ratio has been found to vary for some species (Xavier et al., 2011). Therefore, the UB should also be analyzed to evaluate functionality in order to comprehensively understand squid ecology.

In conclusion, we analyzed $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ in the beaks of two *O. bartramii* stocks, and compared isotopic values between the UB and LB. The $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values

were significantly different between the two stocks except for C/N and UB and LB significantly differed in all isotopic signatures. Both $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values increased drastically with size in the eastern stock, while $\delta^{15}\text{N}$ only slightly increased and $\delta^{13}\text{C}$ varied minimally in the western stock. Trophic niche width was distinct between two stocks with little overlap. GAMs selected latitude and ML as significant variables in explaining $\delta^{13}\text{C}$ variation, whereas ML was the only significant factor for $\delta^{15}\text{N}$ in the eastern stock. In the western stock, none of the variables explained the $\delta^{13}\text{C}$ variation, while latitude and ML were both significant for $\delta^{15}\text{N}$. $\delta^{15}\text{N}$ leveled off in both stocks, which may relate to squid consuming the highest trophic levels of prey after reaching reproductive maturity. Stable isotope values of the UB were less than those of the LB, possibly due to different proportions of chitin between the UB and LB. Future studies might focus on increasing sample size, analyzing different structures and identifying the function of the UB.

Acknowledgements

The support of the scientific surveys by commercial jigging vessel *F/V Jinhai 827* is gratefully acknowledged. This work was funded by the National Science Foundation of China (NSFC41306127 and NSFC41276156), National Science Foundation of Shanghai (13ZR1419700), the innovation Program of Shanghai Municipal Education Commission (13YZ091), State 863 projects (2012AA092303), the Funding Program for Outstanding Dissertations in Shanghai Ocean University and

Shanghai Leading Academic Discipline Project (Fisheries Discipline). The involvement of Y. Chen was supported by SHOU International Center for Marine Studies and Shanghai 1000 Talent Program.

References

- Akaike, H., 1974. A new look at the statistical model identification. Automatic Control. IEEE. Transactions. 19, 716-723.
- Argüelles, J., Lorrain, A., Cherel, Y., Graco, M., Tafur, R., Alegre, A., Espinoza, P., Taípe, A., Ayón, P., Bertrand, A., 2012. Tracking habitat and resource use for the jumbo squid *Dosidicus gigas*: a stable isotope analysis in the Northern Humboldt Current System. Mar. biol. 159, 2105-2116.
- Arbuckle, N.S.M., Wormuth, J.H., 2014. Trace elemental patterns in Humboldt squid statoliths from three geographic regions. Hydrobiologia. 725, 115-123.
- Boletzky, S.V., 2007. Origin of the lower jaw in cephalopods: a biting issue. Paläontologische Zeitschrift. 81, 328-333.
- Bower, J.R., Ichii, T., 2005. The red flying squid (*Ommastrephes bartramii*): a review of recent research and the fishery in Japan. Fish. Res. 76, 39-55.
- Bower, S.M., Margolis, L., 1991. Potential use of helminth parasites in stock identification of flying squid, *Ommastrephes bartrami*, in North Pacific waters. Can. J. Zool. 69, 1124-1126.

403 Boyle, P., Rodhouse, P., 2008. Cephalopods: ecology and fisheries. John Wiley &
404 Sons.

405 Chen, C.S., Chiu, T.S., 2003. Variations of life history parameters in two geographical
406 groups of the neon flying squid, *Ommastrephes bartramii*, from the North Pacific.
407 Fish. Res. 63, 349–366.

408 Chen, C.S., 2010. Abundance trends of two neon flying squid (*Ommastrephes*
409 *bartramii*) stocks in the North Pacific. ICES. J. Mar. Sci. 67, 1336-1345.

410 Chen, X.J., Liu, B.L., Chen, Y. 2008. A review of the development of Chinese distant-
411 water squid jigging fisheries. Fish. Res. 89, 211-221.

412 Cherel, Y., Hobson, K.A., 2005. Stable isotopes, beaks and predators: a new tool to
413 study the trophic ecology of cephalopods, including giant and colossal squids.
414 Proc. R. Soc. Lond. B. 272, 1601-1607.

415 Cherel, Y., Ridoux, V., Spitz, J., Richard, P., 2009a. Stable isotopes document the
416 trophic structure of a deep-sea cephalopod assemblage including giant octopod
417 and giant squid. Biol. Let. 5, 364-367.

418 Cherel, Y., Fontaine, C., Jackson, G.D., Jackson, C.H., Richard, P., 2009b. Tissue,
419 ontogenetic and sex-related difference in $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of the oceanic
420 squid *Todarodes fillipovae* (Cephalopoda: Ommastrephidae). Mar. Biol. 156, 699-
421 708.

422 Clarke, M.R., 1996. The role of cephalopods in the world's oceans: general
423 conclusions and the future. Philos. Trans. R. Soc. Lond. B. 351, 977–1112.

424 Crespi-Abril, A.C., Baron, P.J., 2012. Revision of the population structuring of *Illex*

425 *argentinus* (Castellanos, 1960) and a new interpretation based on modelling the
 426 spatio-temporal environmental suitability for spawning and nursery. Fish. Oceano.
 427 21, 199-214.

428 DeNiro, M.J., Epstein, S., 1977. Mechanism of carbon isotope fractionation
 429 associated with lipid synthesis. Science. 197, 261–263.

430 Fang, Z., Liu, B.L., Li, J.H., Su, H., Chen, X.J., 2014. Stock identification of neon
 431 flying squid (*Ommastrephes bartramii*) in the North Pacific Ocean on the basis of
 432 beak and statolith morphology. Sci. Mar. 78, 239-248.

433 Fang, Z., Li, J.H., Katherine, T., Hu, F.F., Chen, X.J., Liu, B.L., Chen, Y., 2016. Age,
 434 growth and population structure of neon flying squid (*Ommastrephes bartramii*)
 435 in the North Pacific Ocean based on beak microstructure. Fish. Bull. 114, 34-44.

436 FAO. 2013. Fishery and Aquaculture Statistics. 2011. FAO yearbook, Rome, 10-11.

437 Focken, U., Becker, K., 1998. Metabolic fractionation of stable carbon isotopes:
 438 implications of different proximate compositions for studies of the aquatic food
 439 webs using $\delta^{13}\text{C}$ data. Oecologia. 115, 337–343.

440 García, V.H., 2003. Growth and pigmentation process of the beaks of *Todaropsis*
 441 *ebblanae* (Cephalopoda: Ommastrephidae). Berliner. Palaobiol. Abh. 3, 131-140.

442 Guerra, Á., Rodríguez-Navarro, A.B., González, Á.F., Romanek, C.S., Álvarez-Lloret,
 443 P., Pierce, G.J., 2010. Life-history traits of the giant squid *Architeuthis dux*
 444 revealed from stable isotope signatures recorded in beaks. ICES. J. Mar. Sci. 67,
 445 1425-1431.

446 Hijmans, R.J., 2014. geosphere: Spherical Trigonometry. R package version 1.3-11.

447 <http://CRAN.R-project.org/package=geosphere>

448 Hobson, K.A., Welch, H.E., 1992. Determination of trophic relationships within a
449 high Arctic marine food web using $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ analysis. Mar. Ecol. Prog. Ser.
450 84, 9–18.

451 Hobson, K.A., Welch, H.E., 1994. Cannibalism and trophic structure in a high Arctic
452 lake: insights from stable-isotope analysis. Can. J. Fish. Aquat. Sci. 52, 1195–
453 1201.

454 Hobson, K.A., Gibbs, H.L., Gloutney, M.L., 1997. Preservation of blood and tissue
455 samples for stable-carbon and stable-nitrogen isotope analysis. Can. J. Zool. 75,
456 1720–1723.

457 Hobson, K.A., Cherel, Y., 2006. Isotopic reconstruction of marine food webs using
458 cephalopod beaks new insight from captive raised *Sepia officinalis*. Can. J. Zool.
459 84, 766–770.

460 Hunsicker, M.E., Essington, T.E., Aydin, K.Y., Ishida, B., 2010. Predatory role of the
461 commander squid *Berryteuthis magister* in the eastern Bering Sea: insights from
462 stable isotopes and food habits. Mar. Ecol. Prog. Ser. 415, 91–108.

463 Ichii, T., Mahapatra, K., Sakai, M., Inagake, D., Okada, Y., 2004. Differing body size
464 between the autumn and the winter–spring cohorts of neon flying squid
465 (*Ommastrephes bartramii*) related to the oceanographic regime in the North
466 Pacific: a hypothesis. Fish. Oceano. 13, 295–309.

467 Ichii, T., Mahapatra, K., Sakai, M., Okada, Y., 2009. Life history of the neon flying
468 squid: effect of the oceanographic regime in the North Pacific Ocean. Mar. Ecol.

469 Prog. Ser. 378, 1–11.

470 Jackson, G.D., Bustamante, P., Cherel, Y., Fulton, A., Grist, E.P.M., Jackson, C.H.,
 471 Nichols, P.D., Pethybridge, H., Phillips, K., Ward, R.D., Xavier, J.C., 2007.
 472 Applying new tools to cephalopod trophic dynamics and ecology: perspectives
 473 from the Southern Ocean Cephalopod Workshop, 2–3 February 2006. Rev. Fish.
 474 Biol. Fish. 17,79–99.

475 Jereb, P., Roper, C.F.E., 2010. Cephalopods of the world. An annotated and illustrated
 476 catalogue of cephalopod species known to date. vol.2. Myopsid and Oegopsid
 477 Squids, FAO Species Catalogue for Fishery Purposes, Rome, p. 269

478 Jin, Y., Fang, Z., Li, Y.K., Chen, X.J., Hu, H.L., 2015. Growth characteristics analysis
 479 of *Ommastrephes bartramii* beaks in eastern North Pacific. Marine. Fishery. 37,
 480 101-106. (In Chinese with English abstract)

481 Katugin, O.N., 2002. Patterns of genetic variability and population structure in the
 482 North Pacific squids *Ommastrephes bartramii*, *Todarodes pacificus*, and
 483 *Berryteuthis magister*. Bull. Mar. Sci. 71, 383-420.

484 Lipinski, M.R., Underhill, L.G., 1995. Sexual maturation in squid: quantum or
 485 continuum? S. Afr. J. Mar. Sci. 15, 207-223.

486 Liu, B.L., Chen, X.J., Yi, Q., 2013. A comparison of fishery biology of jumbo flying
 487 squid, *Dosidicus gigas* outside three Exclusive Economic Zones in the Eastern
 488 Pacific Ocean. Chin. J. Oceanol. Limnol. 31, 523-533.

489 Logan, J.M., Jardine, T.D., Miller, T.J., Bunn, S.E., Cunjak, R.A., Lutcavage, M.E.,
 490 2008. Lipid corrections in carbon and nitrogen stable isotope analyses:

491 comparison of chemical extraction and modelling methods. J. Anim. Ecol. 77,
492 838-846.

493 Lorrain, A., Arguelles, J., Alegre, A., Bertrand, A., Munaron, J.M., Richard, P., Cherel,
494 Y., 2011. Sequential isotopic signature along gladius highlights contrasted
495 individual foraging strategies of Jumbo Squid (*Dosidicus gigas*). PLoS. One. 6,
496 e22194.

497 McCutchan, J.H., Lewis, W.M., Kendall, C., McGrath, C.C., 2003. Variation in
498 trophic shift for stable isotope ratios of carbon, nitrogen, and sulphur. Oikos. 102,
499 378–390.

500 Miller, T.W., Bosley, K.L., Shibata, J., Brodeur, R.D., Omori, K., Emmett, R., 2013.
501 Contribution of prey to Humboldt squid *Dosidicus gigas* in the northern California
502 Current, revealed by stable isotope analyses. Mar. Ecol. Prog. Ser. 477, 123-134.

503 Mingawa, M., Wada, E., 1986. Stepwise enrichment of ^{15}N along food chains: further
504 evidence and the relation between $\delta^{15}\text{N}$ and animal age. Geochm. Cosmochim.
505 Acta. 48, 1135–1140.

506 Miserez, A., Schneberk, T., Sun, C., Zok, F.W., Waite, J.H., 2008. The transition from
507 stiff to compliant materials in squid beaks. Science. 319, 1816-1819.

508 Moltschaniwskyj, N.A., Carter, C.G., 2010. Protein synthesis, degradation, and
509 retention: mechanisms of indeterminate growth in cephalopods. Physiol. Biochem.
510 Zool. 83, 997-1008.

511 Murata, M., 1990. Oceanic resources of squids. Mar. Behav. Phys. 18, 19–71.

512 Murata, M., Nakamura, Y., 1998. Seasonal migration and diel vertical migration of the

513 neon flying squid, *Ommastrephes bartramii*, in the North Pacific. In: Okutani, T.
 514 (Ed.), Contributed Papers to International Symposium on Large Pelagic Squids.
 515 Japan Marine Fishery Resources Research Center, Tokyo, pp. 13–30
 516 Nagasawa, K., Mori, J., Okamura, H., 1998. Parasites as biological tags of stocks of
 517 neon flying squid (*Ommastrephes bartramii*) in the North Pacific Ocean, In:
 518 Okutani, T. (1998). Contributed papers to International Symposium on Large
 519 Pelagic Squids, July 18-19, 1996, for JAMARC's 25th anniversary of its
 520 foundation, pp 49-64.
 521 Navarro, J., Coll, M., Somes, C., Olson, R.J., 2013. Trophic niche of squids: Insights
 522 from isotopic data in marine systems worldwide. Deep. Sea. Res. Topi. Stud.
 523 Oceano. 95, 93-102.
 524 Onthank, K.L., 2013. Exploring the life histories of cephalopods using stable isotope
 525 analysis of an archival tissue (Doctoral dissertation, Washington State University).
 526 Parry, M., 2008. Trophic variation with length in two ommastrephid squids,
 527 *Ommastrephes bartramii* and *Sthenoteuthis oualaniensis*. Mar. Biol. 153, 249-256.
 528 Pecl, G.T., Moltschaniwskyj, N.A., Tracey, S.R., Jordan, A.R., 2004. Inter-annual
 529 plasticity of squid life history and population structure: ecological and
 530 management implications. Oecologia. 139, 515-524.
 531 Petersen, B.J., Fry, B., 1987. Stable isotopes in ecosystem studies. Annu. Rev. Ecol.
 532 Syst. 18, 293–320.
 533 Piatkowski, U., Pierce, G.J., Morais da Cunha, M., 2001. Impact of cephalopods in
 534 the food chain and their interaction with the environment and fisheries: an

535 overview. Fish. Res. 52, 5-10.

536 Rau, G.H., Sweeney, R.E., Kaplan, I.R., 1982. Plankton $^{13}\text{C}/^{12}\text{C}$ ratio changes with
537 latitude: Differences between northern and southern oceans. Deep-Sea. Res.
538 29,1035-1039.

539 Rau, G.H., Mearns, A.J., Young, D.R., Olson, R.J., Schafer, H.A., Kaplan, I.R., 1983.
540 Animal $^{13}\text{C}/^{12}\text{C}$ correlates with trophic level in pelagic food webs. Ecology. 64,
541 1314–1318.

542 Raya, C., Bartolomé, A., García-Santamaría, M.T., Pascual-Alayón, P., Almansa, E.,
543 2010. Age estimation obtained from analysis of octopus (*Octopus vulgaris* Cuvier,
544 1797) beaks: Improvements and comparisons. Fish. Res. 106, 171-176.

545 Rodhouse, P.G., Nigmatullin, C.M., 1996. Role as consumers. Phil. Trans. R. Soc.
546 Lond. B. 351,1003–1022.

547 Rubenstein, D.R., Hobson, K.A., 2004. From bird to butterflies: animal movements
548 patterns and stable isotopes. Trends. Ecol. Evol. 19, 256–263

549 Ruiz-Cooley, R.I., Gendron, D., Aguñiga, S., Mesnick, S., Carriquiry, J.D., 2004.
550 Trophic relationships between sperm whales and jumbo squid using stable
551 isotopes of C and N. Mar. Ecol. Progr. Ser. 277, 275-283.

552 Ruiz-Cooley, R.I., Markaida, U., Gendron, D., Aguiniga, S., 2006. Stable isotopes in
553 jumbo squid (*Dosidicus gigas*) beaks to estimate its trophic position: comparison
554 between stomach contents and stable isotopes. J. Mar. Biol. Ass. U. K., 86, 437-
555 445.

556 Ruiz-Cooley, R.I., Garcia, K.Y., Hetherington, E.D., 2011. Effects of lipid removal

557 and preservatives on carbon and nitrogen stable isotope ratios of squid tissues:
 558 Implications for ecological studies. J. Exp. Mar. Biol. Ecol. 407, 101-107.

559 Ruiz-Cooley, R.I., Ballance, L.T., McCarthy, M.D., 2013. Range expansion of the
 560 jumbo squid in the NE Pacific: $\delta^{15}\text{N}$ decrypts multiple origins, migration and
 561 habitat use. PloS. One. 8, e59651.

562 Sandoval-Castellanos, E., Uribe-Alcocer, M., Díaz-Jaimes, P., 2010. Population
 563 genetic structure of the Humboldt squid (*Dosidicus gigas* d'Orbigny, 1835)
 564 inferred by mitochondrial DNA analysis. J. Exp. Mar. Biol. Ecol. 385, 73-78.

565 Schimmelmann, A., DeNiro, M.J., 1986. Stable isotopic studies on chitin. III. The
 566 D/H and $^{18}\text{O}/^{16}\text{O}$ ratios in arthropod chitin. Geochim. Cosmochim. Acta. 50, 1485-
 567 1496.

568 Seki, M.P., Polovina, J.J., Kobayashi, D.R., Bidigare, R.R., Mitchum, G.T., 2002. An
 569 oceanographic characterization of swordfish (*Xiphias gladius*) longline fishing
 570 grounds in the springtime subtropical North Pacific. Fish. Oceano. 11, 251-266.

571 Stowasser, G., Pierce, G.J., Moffat, C.F., Collins, M.A., Forsythe, J.W., 2006.
 572 Experimental study on the effect of diet on fatty acid and stable isotope profiles of
 573 the squid *Lolligun culabrevis*. J. Exp. Mar. Biol. Ecol. 333, 97-114.

574 Takahashi, M., Watanabe, Y., Kinoshita, T., Watanabe, C., 2001. Growth of larval and
 575 early juvenile Japanese anchovy, *Engraulis japonicus*, in the Kuroshio-Oyashio
 576 transition region. Fish. Oceano. 10, 235-247.

577 Takai, N., Onaka, S., Ikeda, Y., Yatsu, A., Kidokoro, H., Sakamoto, W., 2000.
 578 Geographical variations in carbon and nitrogen stable isotope ratios in squid. J.

579 Mar. Biol. Ass. U. K. 80, 675-684.

580 Uchikawa, K., Sakai, M., Wakabayashi, T., Ichii, T., 2009. The relationship between
581 paralarval feeding and morphological changes in the proboscis and beaks of the
582 neon flying squid *Ommastrephes bartramii*. Fish. Sci. 75, 317-323.

583 Vanderklift, A., Ponsard, S., 2003. Sources of variation in consumer diet $\delta^{15}\text{N}$
584 enrichment: a meta-analysis. Oecologia. 136,169–182

585 Wada, E., Hattori, A., 1990. Nitrogen in the sea: forms, abundance, and rate processes.
586 CRC press.

587 Watanabe, H., Kubodera, T., Ichii, T., Kawahara, S., 2004. Feeding habits of neon
588 flying squid *Ommastrephes bartramii* in the transitional region of the central
589 North Pacific. Mar. Ecol. Progr. Ser. 266, 173-184.

590 Watanabe, H., Kubodera, T., Ichii, T., Sakai, M., Moku, M., Seitou, M., 2008. Diet
591 and sexual maturation of the neon flying squid *Ommastrephes bartramii* during
592 autumn and spring in the Kuroshio–Oyashio transition region. J. Mar. Biol. Ass. U.
593 K. 88, 381-389.

594 Wood, S., 2006. Generalized additive models: an introduction with R. CRC press.

595 Xavier, J.C., Phillips, R.A., Cherel, Y., 2011. Cephalopods in marine predator diet
596 assessments: why identifying upper and lower beaks is important. ICES. J. Mar.
597 Sci. 68, 1857-1864.

598 Xavier, J.C., Allcock, A.L., Cherel, Y., Lipinski, M.R., Pierce, G.J., Rodhouse, P.G.,
599 Rosa, R., Shea, E.K., Strugnell, J.M., Vidal, E.A.G., Villanueva, R., Ziegler, A.
600 2014. Future challenges in cephalopod research. J. Mar. Biol. Ass. U. K.

DOI:10.1017/S0025315414000782.

Yatsu, A., Midorikawa, S., Shimada, T., Uozumi, Y., 1997. Age and growth of the
neon flying squid, *Ommastrephes bartrami*, in the North Pacific Ocean. Fish. Res.
29, 257–270

Yatsu, A., Tanaka, H., Mori, J., 1998. Population structure of the neon flying squid,
Ommastrephes bartramii, in the North Pacific. In: Okutani T. (ed), Contributed
Papers to International Symposium on Large Pelagic Squids. Japan Marine
Fishery Resources Research Center, Tokyo, pp 31–48.

628

629 Table 1. Summary information of sampled eastern stock of *Ommastrephes bartramii*
630 in North Pacific Ocean (max: maximum; min: minimum; ave: average; std: standard
631 deviation)

ID	Catch date (yy/mm/dd)	Coordinate	ML(mm)	Sex	Maturity stage	Upper beak			Lower beak		
						¹³ C (‰)	¹⁵ N (‰)	¹³ C/ ¹⁵ N	¹³ C (‰)	¹⁵ N (‰)	¹³ C/ ¹⁵ N
2	2010/6/9	172°17'W, 39°52'N	454	♀	3	-18.22	9.06	3.79	-17.71	9.59	3.26
75	2010/5/30	175°34'E, 38°45'N	285	♀	1	-18.82	6.52	4.09	-18.39	6.76	3.48
103	2010/6/20	175°18'W, 39°57'N	253	♂	2	-18.82	5.28	4.26	-18.72	6.24	3.58
136	2010/6/20	175°18'W, 39°57'N	244	♀	1	-18.70	5.95	4.37	-18.61	6.17	3.62
139	2010/6/20	175°18'W, 39°57'N	248	♂	1	-18.67	7.07	4.14	-18.32	7.35	3.45
161	2010/5/18	175°22'E, 38°49'N	403	♀	2	-18.32	8.32	4.28	-17.88	9.35	3.41
164	2010/5/18	175°22'E, 38°49'N	363	♀	2	-18.06	8.65	4.10	-17.70	9.07	3.51
165	2010/5/18	175°22'E, 38°49'N	293	♀	1	-18.90	6.70	4.34	-18.58	6.55	3.61
179	2010/5/24	175°24'E, 39°03'N	346	♀	2	-18.18	7.82	4.07	-17.93	8.75	3.40
182	2010/5/16	176°15'E, 38°37'N	418	♀	2	-18.51	7.66	4.09	-18.13	8.37	3.20
194	2010/6/10	171°58'W, 39°52'N	446	♀	3	-18.43	8.21	3.99	-17.92	8.58	3.34
218	2010/5/18	175°22'E, 38°49'N	435	♀	2	-18.10	8.12	4.01	-17.74	8.72	3.29
223	2010/5/18	175°22'E, 38°49'N	208	♀	1	-18.40	4.81	4.10	-17.97	5.41	3.54
270	2010/5/10	171°43'E, 39°13'N	473	♀	3	-18.01	10.31	3.26	-17.13	10.35	3.08
274	2010/5/21	175°14'E, 38°51'N	325	♀	2	-18.35	7.79	4.18	-17.85	8.31	3.48
281	2010/5/21	175°14'E, 38°51'N	461	♀	3	-17.84	9.03	3.94	-17.68	9.13	3.30
291	2010/5/19	175°17'E, 38°50'N	305	♀	2	-18.62	6.56	3.26	-17.95	6.72	3.11
346	2010/6/17	172°53'W, 40°18'N	376	♀	2	-18.27	8.52	3.99	-17.92	9.29	3.37
385	2010/5/27	170°01'E, 38°22'N	352	♀	2	-18.14	7.82	4.22	-17.91	8.13	3.58
404	2010/6/8	172°06'W, 39°46'N	319	♀	1	-18.83	5.86	4.30	-18.31	6.85	3.11
415	2010/6/20	175°18'W, 39°57'N	250	♂	1	-18.57	5.70	4.13	-18.15	6.88	3.05
417	2010/6/25	176°28'W, 39°47'N	255	♂	1	-18.82	5.39	4.24	-18.55	5.87	3.47
418	2010/6/20	175°18'W, 39°57'N	265	♀	1	-18.81	6.39	3.22	-18.28	6.67	3.10
421	2010/6/26	175°38'W, 39°37'N	244	♂	1	-18.39	5.65	4.22	-18.35	6.52	3.47
424	2010/6/20	175°18'W, 39°57'N	220	♂	1	-18.29	5.83	4.08	-18.26	6.52	3.50
428	2010/6/20	175°18'W, 39°50'N	212	♂	2	-18.70	4.65	3.99	-18.38	5.58	3.31
430	2010/6/27	176°48'W, 39°57'N	232	♂	1	-18.86	5.05	4.18	-18.81	6.03	3.62
432	2010/6/20	175°18'W, 39°47'N	270	♂	2	-18.80	5.42	4.06	-18.58	6.04	3.41
441	2010/6/29	176°28'W, 39°57'N	230	♂	2	-18.79	5.57	4.14	-18.50	5.44	3.53
448	2010/6/20	175°18'W, 39°57'N	253	♀	1	-18.65	6.46	3.19	-18.34	6.49	3.11
max	-	-	473	-	-	-17.84	10.31	4.37	-17.13	10.35	3.62
min	-	-	208	-	-	-18.90	4.65	3.19	-18.81	5.41	3.05
ave	-	-	315	-	-	-18.50	6.87	4.01	-18.15	7.39	3.38
std	-	-	84	-	-	0.30	1.48	0.33	0.38	1.43	0.18

632

633

634

635 Table 2. Summary information of sampled western stock of *Ommastrephes bartramii*

636 in North Pacific Ocean (max: maximum; min: minimum; ave: average; std: standard

637 deviation)

ID	Catch date (yy/mm/dd)	Coordinate	ML(mm)	Sex	Maturity stage	Upper beak			Lower beak		
						¹³ C (‰)	¹⁵ N (‰)	¹³ C/ ¹⁵ N	¹³ C (‰)	¹⁵ N (‰)	¹³ C/ ¹⁵ N
9	2010/7/9	157°47'E, 38°55'N	187	♀	1	-17.34	7.86	4.02	-17.43	8.00	3.45
156	2010/9/13	156°27'E, 43°39'N	285	♀	2	-17.53	9.51	4.09	-17.29	10.22	3.25
234	2010/8/2	157°12'E, 41°02'N	178	♂	1	-18.14	7.35	4.08	-18.11	7.59	3.54
237	2010/8/2	157°12'E, 41°02'N	163	♂	1	-18.11	8.17	3.63	-18.06	8.62	3.20
324	2010/11/13	149°52'E, 40°15'N	341	♂	3	-17.92	8.95	3.96	-17.76	8.98	3.33
520	2010/10/3	153°41'E, 41°41'N	293	♂	2	-18.35	8.96	3.93	-17.97	9.60	3.37
536	2010/8/10	156°59'E, 41°35'N	304	♂	3	-17.86	9.08	4.02	-17.59	9.30	3.34
541	2010/8/10	156°59'E, 41°35'N	215	♂	1	-18.09	8.35	3.99	-18.11	8.69	3.52
654	2010/7/19	134°25'E, 39°53'N	211	♀	2	-17.85	8.19	4.16	-17.89	8.50	3.59
683	2010/7/19	134°25'E, 39°53'N	194	♀	2	-17.99	7.51	4.15	-17.73	8.39	3.40
854	2010/8/15	154°14'E, 41°49'N	274	♂	3	-17.75	8.34	4.19	-17.60	8.63	3.46
963	2010/10/23	154°38'E, 42°09'N	374	♀	2	-18.55	8.79	4.10	-18.23	9.12	3.40
1004	2010/9/22	154°05'E, 43°26'N	361	♀	2	-17.89	8.87	4.00	-17.63	9.50	3.33
1009	2010/9/22	154°05'E, 43°26'N	381	♀	2	-17.82	10.11	3.19	-17.57	9.72	3.07
1046	2010/10/21	155°43'E, 41°50'N	351	♀	2	-18.32	9.05	3.90	-17.78	9.49	3.07
1101	2010/8/17	155°36'E, 43°29'N	247	♀	2	-18.08	9.27	4.02	-17.92	9.46	3.37
1132	2010/10/10	156°09'E, 42°58'N	326	♀	2	-18.06	9.79	3.81	-17.82	9.65	3.28
1150	2010/7/20	153°42'E, 39°36'N	201	♀	2	-18.34	7.07	3.56	-17.95	7.31	3.30
1167	2010/7/20	153°42'E, 39°36'N	207	♂	1	-18.28	8.38	3.94	-18.35	8.54	3.36
1298	2010/10/16	157°06'E, 43°17'N	242	♂	1	-17.93	8.94	3.89	-17.66	9.51	3.33
1329	2010/9/11	156°28'E, 43°19'N	267	♀	2	-18.12	9.62	3.73	-17.97	9.72	3.20
1425	2010/8/7	154°53'E, 42°30'N	313	♀	2	-17.84	8.52	3.31	-17.53	8.92	3.11
1440	2010/8/7	154°53'E, 42°30'N	256	♂	2	-17.79	8.88	3.75	-17.41	8.82	3.35
1679	2010/10/12	156°47'E, 42°35'N	236	♀	2	-18.04	8.96	4.03	-17.85	9.26	3.54
1703	2010/7/6	157°46'E, 39°08'N	225	♂	1	-18.15	8.37	4.14	-17.89	9.16	3.41
1817	2010/9/19	158°35'E, 44°27'N	261	♂	2	-18.17	9.22	4.07	-17.98	9.49	3.21
1854	2010/11/5	154°07'E, 41°57'N	421	♀	3	-18.52	9.05	4.06	-18.17	9.69	3.23
1855	2010/11/5	154°07'E, 41°57'N	405	♀	3	-18.15	10.22	3.70	-17.72	10.13	3.18
1874	2010/11/5	154°07'E, 41°57'N	281	♂	3	-18.10	9.64	3.98	-17.89	9.41	3.39
1878	2010/7/13	153°36'E, 39°24'N	231	♀	2	-17.99	8.53	3.72	-17.63	8.49	3.38
max	-	-	421	-	-	-17.34	10.22	4.19	-17.29	10.22	3.59
min	-	-	163	-	-	-18.55	7.07	3.19	-18.35	7.31	3.07
ave	-	-	274	-	-	-18.04	8.79	3.90	-17.82	9.06	3.33
std	-	-	71	-	-	0.26	0.76	0.24	0.26	0.69	0.13

638

Table 3. Analysis of isotopic values of western and eastern North Pacific Ocean *Ommastrephes bartramii* stocks (t-test results)

Group	Variable	<i>df</i>	<i>t</i>	<i>P</i>
Stock	ML	58	2.01	0.05
	CU	58	-6.30	<0.01
	NU	58	-6.31	<0.01
	C/NU	58	1.38	0.17 ^{ns}
	CL	58	-4.02	<0.01
	NL	58	-5.78	<0.01
	C/NL	58	1.08	0.28 ^{ns}
UB and LB	CE	29	-9.89	<0.01
	NE	29	-7.71	<0.01
	C/NE	29	14.10	<0.01
	CW	29	-7.83	<0.01
	NW	29	-4.97	<0.01
	C/NW	29	17.00	<0.01

Abbreviations: ML=Mantle length, CU=Carbon isotope value of upper beak, NU=Nitrogen isotope value of upper beak, C/NU= Carbon/Nitrogen value of upper beak, CL=Carbon isotope value of lower beak, NL=Nitrogen isotope value of lower beak, C/NL= Carbon/Nitrogen value of lower beak, CE=Carbon isotope value of eastern stock, NE=Nitrogen isotope value of eastern stock, C/NE= Carbon/Nitrogen value of eastern stock, CW=Carbon isotope value of western stock, NW=Carbon isotope value of western stock, C/NW= Carbon/Nitrogen value of western stock
ns= non-significant

Table 4. The AIC values for the selection of General Additive Model (GAM) for *Ommastrephes bartramii* isotopic analysis

RV	AO	LAT	DSB	ML	LAT+DSB	LAT+ML	DSB+ML	ALL
$\delta^{13}\text{C}$	UE	10.983	6.334	-14.325	7.669	<u>-19.932</u>	-12.378	-19.823
	LE	14.924	9.068	-6.809	7.730	<u>-11.827</u>	-4.819	-6.257
	UW	8.076	6.213	7.705	7.636	8.617	<u>3.134</u>	5.027
	LW	6.610	5.317	5.751	7.194	6.881	<u>5.200</u>	7.150
$\delta^{15}\text{N}$	UE	102.491	96.104	<u>55.705</u>	95.754	57.623	57.304	58.836
	LE	100.444	94.158	<u>46.486</u>	95.518	48.115	47.486	49.243
	UW	55.235	52.596	46.658	52.819	<u>44.287</u>	49.110	46.031
	LW	47.909	48.668	42.425	46.922	<u>37.307</u>	44.511	37.700

Abbreviations: RV=Response variable, AO=Analyzed object, LAT=Latitude, DSB=Distance to shelf break, ML= Mantle length, ALL=All the three independent variables, UE=Upper beak of eastern stock, LE=Lower beak of eastern stock, UW=Upper beak of western stock, LW=Lower beak of western stock

Underlines are the lowest AIC values which the best model chosen

Table 5. Statistical outputs of GAMs in $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ for *Ommastrephes bartramii*

Variable	Sample part	Source	e.d.f	<i>F</i>	<i>P</i>	% explained
$\delta^{13}\text{C}$	UE	LAT	1.00	5.07	0.03	81.6
		ML	6.28	9.99	<0.01	
	LE	LAT	1.00	5.50	0.029	83.0
		ML	8.25	7.19	<0.01	
	UW	ML	1.51	1.70	0.20 ^{ns}	32.8
	LW	ML	1.83	0.87	0.44 ^{ns}	23.8
$\delta^{15}\text{N}$	UE	ML	2.16	69.82	<0.01	84.7
	LE	ML	5.76	36.33	<0.01	91.6
	UW	LAT	1.00	9.11	<0.01	66.1
		ML	4.30	8.24	<0.01	
	LW	LAT	1.00	10.82	<0.01	70.4
		ML	3.38	6.35	<0.01	

Abbreviations: IV=Isotopic variable, RV=Response variable, LAT=Latitude, DSB=Distance to shelf break, ML= Mantle length, ALL=All the three independent variables, UE=Upper beak of eastern stock, LE=Lower beak of eastern stock, UW=Upper beak of western stock, LW=Lower beak of western stock; ns= non-significant

Figure legends

Fig. 1. Sampling stations of *Ommastrephes bartramii*, and descriptions of Subarctic frontal zone (SAFZ), Transition zone (TZ) and Subtropical frontal zone (STFZ) in the North Pacific Ocean. Western (solid dots) and eastern (triangles) stock represent different stocks. Vertical line in 170°E (longdash) represents the geographical boundary of different stocks.

Fig. 2. Biplot of convex hull carbon-nitrogen isotopes (A), and mean $\pm$ standard deviation isotopic values (B) for upper and lower beaks of western and eastern stocks. Eastern large individuals mean ML>350mm; Eastern small individuals mean ML<350mm.

Fig. 3. Isotopic variations of $\delta^{13}\text{C}$ (A) and $\delta^{15}\text{N}$ (B) for upper and lower beaks of western and eastern stocks. Stick in boxplot means the range of isotopic values, upper counter means the 25% quartile, lower counter means the 75% quartile, solid line in the box mean the medium value.

Fig. 4. General Additive Models (GAMs) for significant variables of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of the eastern stocks for upper (A-C) and upper (D-F) beaks. Black solid line is the mean value, grey filled area is the 95% confident interval.

Fig. 5. General Additive Models (GAMs) for significant variables of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of the western stocks for upper (A-B) and upper (C-D) beaks. Black solid line is the mean value, grey filled area is the 95% confident interval.

778

779

780 Fig.1

781

782

783

784

785

786

787

788

789

790

791

792

793

794

A

B

Fig. 2

807
808
809
810

811
812
813

A

814
815
816
817

Fig. 3

Fig. 4

846

847

848 Fig. 5