1 **Corn Residue, Tillage, and Nitrogen Rate Effects on Soil Properties**

2 María B. Villamil*, Joseph Little, and Emerson D. Nafziger

3 Department of Crop Sciences, University of Illinois, 1102 S. Goodwin Ave., Urbana, IL 61821.

4 *Corresponding author:

5 María Villamil, Department of Crop Sciences, University of Illinois, 1102 S. Goodwin Ave.,

6 Urbana, IL 61801. Phone: +1 (217) 333 4690. villamil@illinois.edu

7 **Abbreviations:** PR, penetration resistance; BD, bulk density; WAS, water aggregate stability;

8 SOC_s, soil organic carbon stocks; SOC_c, soil organic carbon concentration.

Abstract

Harvesting corn (*Zea mays* L.) residue for use as a biofuel feedstock may alter important chemical and physical properties of soils. Long-term residue removal, linked with tillage and fertilizer practices, could lower soil organic carbon (SOC), as well as increase soil compaction and susceptibility to erosion. A study initiated in 2006 included three levels of residue removal (none, partial, or full removal), two tillage treatments (no-till or chisel plow), and two N application rates (134 and 268 kg N/ha). These experiments were established in continuous corn (CC) at four Illinois sites, with treatments remaining in the same plots each year. In the spring at the beginning of the eighth growing season, we measured total carbon stocks (SOC), water aggregate stability (WAS), soil bulk density (BD), and penetration resistance (PR). Results showed that with all crop residue retained in the field, SOC stocks 13% lower under chisel tillage than in no-tilled plots, but removal of some or all of the residue lowered the SOC levels of no-tilled plots to those of tilled plots. On average for the studied depths, no-till plots had 5% and 39% higher BD and PR respectively than tilled plots, and residue removal significantly increased PR under no-till. Regardless of tillage treatment, the highest WAS values were found without residue removal at the lower N rate and with partial removal at the higher N rate. The higher N rate slightly lowered the BD under partial removal of residue for both tillage treatments at both studied depths. These results indicate that crop residue harvest results in only modest changes in soil under continuous corn system, and that these changes are likely smaller than are the effects of tillage and N fertilizer use. But as long as residue amount and tillage practices are sufficient to limit losses of soil by erosion to acceptable levels, we believe that corn residue represents a viable feedstock for a sustainable bioenergy industry in the U.S. Midwest.

1. Introduction

1 With growing interest in biofuel alternatives to petroleum, corn residue has become a
2 leading candidate feedstock for lignocellulosic ethanol production. The United States
3 Department of Energy (DOE) considers residue—the above-ground stalks, leaves, and vegetative
4 residues that remain after grain harvest—a feasible option because of its wide availability and
5 low cost (Perlack et al., 2011). In the United States, farmers planted 38.6 million hectares of corn
6 in 2013; estimates of annually harvestable residue range from 58.3 million to 100 million metric
7 tons (USDA-NASS 2013; Graham et al., 2007; Gallagher and Baumes, 2012). While few
8 lignocellulosic ethanol refineries have begun operations in the Midwest with corn residue as the
9 sole feedstock, the scale of residue harvest could increase in the region as refineries attempt to
10 meet federal mandates for production, currently set to reach 61 billion liters by 2022 (Energy
11 Independence and Security Act, 2007).

12 The commercial use of residue may provide supplemental income from the sale of baled
13 residue, reduced incidence of corn diseases, and better seed placement and germination in fields
14 where corn residue is removed. Pathogens that cause stalk rot and leaf blights overwinter on corn
15 refuse, so clearing a field of residue can help reduce the incidence of disease the following
16 season (White, 1999). Corn residue may also have an autotoxic effect on seedlings that slows
17 development (Singh, et al., 2010). Furthermore, cooler soil temperatures persist in the spring for
18 longer periods of time under residue cover, and with warmer soils, corn seedlings may grow
19 faster in soils without residue cover (Swan et al., 1987).

20 Despite the agronomic benefits of removing residue, the long-term sustainability of
21 residue harvest is unclear. Over 13 years of continuous corn, SOC remained unchanged under
22 residue harvest in no-till plots, but increased by 14% when residue was returned to no-till plots
23 (Clapp et al., 2000). Using the natural abundance of the ¹³C isotope, these researchers also

1 reported that the average half-life of relic SOC was 122 years in plots where residue remained
2 and high N rates (200 kg N ha⁻¹) were used; this half-life was lowered to 71 years in the absence
3 of N fertilizer, and was further lowered, to 49 years, by the combination of residue harvest and
4 no N added. In contrast, several studies have shown that N fertilizer and residue harvest had no
5 effect on SOC within the same tillage treatment (Reicosky et al., 2002; Wilts et al., 2004; Hooker
6 et al., 2005) yet no-till plots had more SOC stock compared to tilled plots with no residue harvest
7 (Hooker et al., 2005).

8 Tillage could increase the rate of residue decomposition by mechanically breaking down
9 residues to increase the surface area of residue exposed to soil microbes and by incorporating
10 residue into the soil where microbes and moisture are more plentiful. This might contribute to
11 faster additions to the SOC pool, but could also disrupt soil macroaggregates that protect relic
12 SOC, thereby increasing the loss of relic SOC (Hammerbeck et al., 2012). The conversion of
13 virgin land to farmland has been estimated to result in the loss of 50% of the original SOC during
14 the first 25 years of cultivation in temperate climates (Matson et al., 1997). However, after the
15 initial conversion to farmland, SOC stocks may reach a new equilibrium and resist further
16 decline (Buyanovsky, et al., 1994; Plante and Parton, 2007). Several studies have shown tillage
17 to have little effect on SOC, though tillage can change the distribution of SOC in the soil profile
18 (Dolan et al., 2006; Blanco-Canqui and Lal, 2008).

19 Fertilizer N could affect SOC by increasing plant growth, thereby increasing the amount
20 of biomass returned to the soil and by providing the substrate needed for microorganisms to
21 consume materials with high C:N ratios (Gregorich et al., 1996). Several studies have reported
22 increased SOC in continuous corn with N application rates ranging from 120 kg N/ha to 280 kg
23 N/ha (Varvel, 1994; Studdert and Echeverria, 2000; Liebig et al., 2002; Jagadamma et al., 2008).

1 Compared against the SOC level measured before a 13-year experiment was initiated, SOC did
2 not decrease only under both N fertilization and residue return (Allmaras et al., 2004). There is
3 some disagreement over the role of N fertilizers in SOC storage, as other researchers have
4 posited that high concentrations of N could enable microorganisms to more readily consume
5 relic SOC (Mulvaney et al., 2009).

6 In addition to their role as the primary source of C inputs, crop residues and their
7 management have significant impacts on soil physical properties including bulk density (BD)
8 root penetration resistance (PR), and water aggregate stability (WAS). Observing lower BD
9 under higher nitrogen fertilizer rates, Jagadamma et al. (2008) suggested that by inducing greater
10 biomass production, which in turn increases the porosity of the soil in the root zone, N fertilizers
11 may indirectly contribute to lowering BD. Wander and Bollero (1999) observed that BD and PR
12 were significantly higher under no-till compared to conventional-till systems. Retention of
13 residues generally increases the number and stability of soil aggregates which in turn promotes
14 SOC stabilization (Buyanovsky et al., 1994; Paustian et al., 1997). In fields where a quarter or
15 more of residue was removed, soil macroaggregates were reduced by 40% (Blanco-Canqui and
16 Lal, 2009). Moebius-Clune et al. (2008) found that tillage had more effect than residue harvest
17 on studied soil properties, with aggregation significantly less stable under residue harvest than
18 under residue return in a no-till system.

19 Our objective of this study was to determine the effects of residue removal in
20 combination with tillage and N rate on soil SOC stocks, WAS, BD and PR in order to identify
21 potential impacts of long term use of these management practices on soil quality parameters that
22 may influence future productivity.

2. Materials and Methods

2.1 Field sites

Field experiments were established in the fall of 2005 following uniformly-cropped corn at four University of Illinois Crop Sciences research centers located near DeKalb (41°55′ N, 88°45′ W), Monmouth (40°54′ N, 90°38′ W), Perry (39°46′ N, 90°44′ W), and Urbana, IL (40°6′ N, 88°12′ W). Soil types were Flanagan silt loam (fine, smectitic, mesic Aquic Argiudolls) at DeKalb and Urbana, Muscatine silt loam (fine-silty, mixed, superactive, mesic Aquic Argiudolls) at Monmouth, and Clarksdale silt loam (fine, smectitic, mesic Udollic Endoaqualfs) at Perry. Full description of each of these soil orders can be found at the official USDA- NRCS query website (Soil Survey Staff, 2014). Briefly, all three soil series are silt loam soils with a slope of about 2%, formed in thick loess over till. Flanagan series consists of dark colored, somewhat poorly drained soils, developed in 1-1.5m of loess over loam till under prairie vegetation. Permeability is moderate and surface runoff is slow to medium. Muscatine series consist of dark colored, very deep soils developed in loess 2-3 m thick over till under prairie vegetation. Muscatine soils are somewhat poorly drained with moderate permeability and low surface runoff potential. Clarksdale soils are very deep, somewhat poorly drained soils formed in 1-3 m loess over till under mixed prairie and forest vegetation. Mean annual precipitation ranges from 890 to 914 mm, and the mean annual temperature from 10.6 to 11.1°C (Illinois State Water Survey, 2010).

2.2 Experimental layout and field methods

The experimental design was a split-split plot arrangement in a randomized complete block with four replications, and all treatments remained in the same place each year. Main plots consisted of one of three levels of corn residue removal (full, partial, and none) in the fall following harvest. Then, subplots were tilled with a chisel plow or similar implement or left undisturbed. In

1 the spring, tilled plots were leveled with secondary tillage before planting, and no-till plots were
2 slot-planted with no tillage of any kind. Rates of N fertilizer (134 and 268 kg N/ha) were
3 assigned to sub-subplots, and applied as 28% urea-ammonium nitrate (UAN) solution either at
4 planting or before the crop reached the 5-leaf stage. Sub-subplots were eight, 76-cm rows (6.1 m)
5 wide by 9.1 to 13.1 m long, depending on location.

6 Residue removal and tillage treatments were implemented in the fall of 2005, and N fertilizer
7 treatments commenced in 2006, the first year of cropping. Full removal of residue involved
8 chopping the residue with a rotary mower at a height of 5 cm, raking the residue into windrows,
9 which were removed. Partial removal of residue was achieved by raising the setting of the rake
10 to a height of 7 cm and running it through unchopped residue to produce windrows about half the
11 size of those from plots with full removal of residue; much of the residue remaining was
12 segments of lower stalk still attached at the root. These smaller windrows were removed and the
13 remaining residue was chopped with a rotary mower at a height of 5 cm. In plots with no residue
14 removed, residue was chopped with a rotary mower at a height of 5 cm. The intent was to
15 simulate full and partial removal of residue using the same methods and equipment that
16 producers might use. Averaged across locations, years, tillage systems, and N rates it was
17 estimated that 7.6, 4.2, and 0.8 Mg/ha of residue remained with no, partial, and full removal of
18 residue, respectively (Coulter and Nafziger, 2008, adjusted for subsequent yields).

19 The chisel plow tillage system used in this study involved chisel plowing to a depth of 25 cm in
20 the fall after residue removal, followed by field cultivating to a depth of 9 cm just before
21 planting. Each year, corn was planted 4 to 5 cm deep at 81,500 seeds/ha with a no-till planter in
22 late April. No-till corn was planted approximately 15 cm to the side of the rows from the
23 previous year. Weeds were controlled with preemergence and postemergence herbicides, which

varied with location and year. Plots were also hand weeded to control escaped weeds. Soil P, K, and pH levels were maintained according to recommendations from the Illinois Agronomy Handbook (Fernandez and Hoefft, 2009).

2.2 Soil sampling and lab procedures

Soil sampling was conducted in the spring of 2013, at the beginning of the eighth crop year of the experiment. Penetration resistance (PR, kPa) was recorded with a Field Scout SC 900 Soil Compaction Meter (Spectrum Technologies, Plainfield, IL) with a cone basal area of 1.28 cm² and a cone angle of 30°. Three subsamples, each consisting of five sub-subsamples, were recorded at each plot and the results averaged at the selected depths of 0-15, 15-30-cm in order to get one measurement per plot per depth considered. Gravimetric water content was determined simultaneously and later used as a covariate in the statistical analysis of PR. For determination of WAS and SOC, three soil cores were taken in each sub-subplot with hand-held probes 20-mm in diameter. Samples were subdivided into depths of 0-15 and 15-30 cm, then cores from each depth were composited. Samples were air-dried and sieved through <2mm, then three subsamples of the 1- 2-mm soil fraction were used to determine WAS (%) with an Eijkelkamp wet sieving apparatus (Eijkelkamp Agrisearch Equipment, Netherlands) using the procedure developed by Kemper and Rosenau (1986). Soil organic carbon concentration (SOCc, %) was obtained using loss on ignition (Davis, 1974) and the results adjusted according to equations developed by Konen et al. (2002) for Illinois soils.

Soil bulk density (BD, Mg/m³) was determined using the core method (Blake and Hartge, 1986) from two subsamples taken within the 0-15 and 15-30-cm depths with a slide-hammer probe with a 5 cm core diameter in each plot. Bulk density values were used to convert SOCc to a

1 basis of weight per unit area, or SOC stocks (SOCs), in Mg/ha (Ellert and Bettany, 1995).

2 Though values of SOC_c are reported in Tables for interested readers the discussion will focus on
3 SOC_cs. This consideration relates to the findings of Goidts et al. (2009), who reported that in
4 croplands, SOC concentration usually contributes more than bulk density to the variability of
5 SOC stocks.

6 2.3 Statistical analysis

7 Data was analyzed using the MIXED procedure of SAS software version 9.4 (SAS Institute Inc.,
8 2013). Residue removal, tillage, and N rates were considered fixed effects, while replicates
9 (blocks within sites) and sites were considered random effects. Significance of random effects
10 was calculated with a Wald Z test statistic using the COVTEST option in the MIXED procedure.
11 Depth (D) was analyzed using a repeated measures technique with a variance-covariance
12 structure selected on the basis of the lowest Akaike's Information Criteria (Littell et al., 2006).
13 Pre-planned estimates were used for mean comparison purposes setting the probability of Type I
14 error or alpha level (α) at 0.05. The CORR procedure of SAS was used to evaluate the relationship
15 between SOC_cs and WAS. Statistical model and SAS codes are available upon request from the
16 authors.

18 3. Results

19 We did not find that residue removal affected any of the parameters measured on its own, but it
20 interacted with tillage treatments (RxT, Table 1) in the responses observed for SOC_cs, SOC_c, and
21 PR (Table 1) while showing an interaction with N rate on BD (RxN, Table 1) and WAS, though
22 the latter was expressed differently at the successive depths (RxNx_D, Table 1). Tillage options

also had a different response on the surface versus the deeper soil layer in the responses measured for SOC_s, SOC_c, and BD (Tx_D, Table 1) whereas N rate interacted with depth on its effects on PR (Nx_D, Table 1). The triple way interaction effects of TxNx_D as well as the four-way interaction of RxTxNx_D among treatments were not detected in the present study (Table 1). Among residue removal levels, SOC_s were significantly lower under chisel tillage than in no-tilled plots with no residue removal, but residue removal under no-till lowered the SOC_s to those found in tilled plots (Table 2). Soil PR, a measurement that mimics the pressure that roots must exert to move through soil at a given depth, on the other hand, showed a clear response to residue removal that depended upon tillage option considered yet the magnitude of the difference in PR between tillage options was consistent among residue treatments (Table 2). Soil PR was consistently lower in chisel tilled plots than in no-tilled plots regardless of residue removal rates, and residue removal significantly increased PR under no-till but had no effect under chisel tilled management (Table 2).

The SOC_s stocks were greater with all residue retained and dropped more in the surface soil (0 to 15 cm) than in the 15-30 cm layer with partial and full residue removal (Table 3).

Soil physical properties of BD and PR were greater in the deeper layer compared to those in the surface soil (Table 2 to 4). Tillage decreased PR, by about 39% (Table 4) and it reduced SOC_s partially by decreasing BD (Tables 1 and 4). The effect of tillage on BD and thus on SOC_s was greater at the 0-15 cm depth than at the 15-30 cm depth, but the T x D interaction was also significant for SOC_c (Tables 1 and 4), so the change in BD with tillage did not fully account for the change in SOC_s.

Nitrogen rate by itself did not affect any of the measured parameters, but interacted with residue removal for BD, with a slightly lower BD value from the lower to the higher N rate with partial

removal, but no change with N rate at either the full or no removal (Tables 1 and 5). Nitrogen rate interacted with depth for PR with a slight increase in this property with the higher N rate; from 1102 kPa to 1208 kPa at the higher N rate. The only significant effect of any treatment or treatment combination for WAS was the RxNx_D interaction (Table 6). Higher WAS values were found with no residue removal at the lower N rate and with partial residue removal at the higher N rate than in the other residue-N rate combinations yet these effects were confined to the surface 0-15cm soil layer (Table 6).

4. Discussion

Under conventional tillage, crop residues and soil aggregates are mechanically disturbed and exposed to soil microorganisms which stimulates decomposition and can lead to reductions in SOC stocks as previously protected C mineralizes (Balesdent et al., 2000). In contrast, no-till soils are undisturbed and crop residues accumulate on the soil surface protecting pre-existing and stable soil organic matter. In this regard, our results show larger SOC stocks within the top soil layer, reflecting the stratification associated with topsoil addition of residues and nutrients, especially under conservation tillage systems (Wilhelm et al., 2004; Dolan et al., 2006). Similar results were reported by Zuber et al. (2015) when comparing tillage and crop rotation effects on soil properties of typical Illinois soils. The authors found a 7% reduction in SOC stocks after 15 years of chisel-till compared to no-tilled plots and under different crop rotations.

In conditions of no residue removal, our results show that SOC stocks were 13.4% greater under no-till than in chisel tilled plots (Table 2). However, partial or full removal of residues under no-till management, lower the SOC stocks to the levels found under chisel-till. These results partially agrees

1 with the results of other long-term, continuous-corn studies, where plots with full residue present
2 had higher SOC stocks than did plots with residue removed (Wilts et al., 2004; Liska et al.,
3 2014). On the other hand, our results show that SOC_s change abruptly under no-till conditions
4 with removal of residues yet SOC_s do not change with removal of residues under chisel-till. We
5 hypothesize that the full removal of residue in these highly fertile soils allows for a faster
6 turnover of the soil carbon due to the higher temperature and reduced water content that
7 commonly limits carbon cycling in this environments particularly under no-till conditions
8 (Needelman et al. 1999). Yet this might be a temporary effect which indicates the need to further
9 evaluate these trends to see if they are continuing in the same direction.

10 Unlike studies that reported increased SOC stocks with increasing N rates in continuous corn
11 production (Varvel, 1994; Studdert and Echeverria, 2000; Liebig et al., 2002; Jagadamma et al.,
12 2008) we found no evidence of N fertilizer rate effect on SOC_s averaged across residue and
13 tillage treatments. These results are in agreement with the findings of Russell et al. (2009), and
14 Brown et al. (2014). Likely, increased SOC decay rates with more available N was offset by the
15 larger amount of biomass produced at higher N rates (Kirkby et al., 2013; Neff et al., 2002).

16 The formation and stability of soil aggregates is often related to the amount of SOC_s and WAS
17 has been reported to increase with a corresponding increase in SOC_s; in our study, SOC_s were
18 positively correlated with WAS ($r = 0.38$, $P = <.0001$), as expected. The explanations of SOC
19 responses to N rate put forward by both Kirkby et al., (2013) and by Neff et al. (2002) may also
20 apply to our results on soil aggregate stability. Water aggregate stability determination uses the
21 soil macro aggregates fraction within 2-1 mm; these aggregates are organized around plant
22 fragments and represent a transient C pool (Buyanovsky et al., 1994). Our results lend support to
23 the idea that this transient pool is more sensitive to management practices such as residue

1 removal and N rates than are total SOC stocks, at least over a period of several years. The drop
2 in WAS with residue removal at the low N rate and the increase in WAS with residue removal at
3 the high N rate reflects this interplay between residue amount and N supply; without residue
4 removal, residue breakdown may be constrained by N supply, while at full removal, N rate has
5 little effect (Table 6).

6 Bulk density was 8% greater under no-till (1.50 Mg/m^3) than under chisel-till management (1.38
7 Mg/m^3) within the top 0-15cm soil layer yet the difference was only 2% within the deeper 15-30
8 cm layer (Table 4). The increased BD under no-till has been reported in a number of other
9 studies, and is ascribed to the lack of mechanical fracturing of the soil under NT (Balesdent et
10 al., 2000). These results are consistent with the observation that residues are less dense and much
11 more elastic than soils, and so tend to resist compaction to a greater extent (Kladviko, 1994);
12 without residue interspersed throughout the surface layer, soil under no-till would tend to be
13 more easily compacted. Tillage, on the other hand, disrupts surface compaction and increases, at
14 least temporarily, the porosity of the soil (Kay and VandenBygaart, 2002), thereby lowering BD
15 and PR compared to no-till (Cassel et al., 1995; Blanco-Canqui et al., 2006; Moebius-Clune et
16 al., 2007). In this regard, our results show that soil PR was 39% lower under chisel-till
17 management compared to no-tilled plots over both depths considered regardless residue removal
18 levels. Still, the PR and BD values we found were substantially less than the theoretical root-
19 restricting levels of 1.75 Mg/m^3 for BD and of 2000 kPa for PR in silt loam and silty clay loam
20 soils (Hamblin, 1985), so we would not expect any of these treatments to represent a cause for
21 concern regarding root growth and crop production.

22 The soil conditions created with partial residue removal and high N rate could hasten turnover of
23 corn residue, increasing aggregation, WAS, and soil porosity, resulting in a lower BD. Yet since

1 this effect was so small we consider it to be of no practical consequence. This may also be said
2 of the finding that, when averaged across residue and tillage treatments, soil PR within the
3 deeper soil layer was increased from 1102 kPa to 1208kPa at the higher N rate; these levels of
4 PR recorded values are far from the 2000 to 4000 kPa threshold range suggested as restrictive for
5 crop production and root growth (Hamblin, 1985).

6 **5. Conclusions**

7 Overall, our results showed that residue removal, tillage, and N fertilization in continuous corn
8 production systems influence soil properties in ways that are closely linked to the role and levels
9 of residue left in the field and what we think may be their rate of decomposition. The levels of
10 corn yield play an important role since in these productive soils of Illinois, with annual corn
11 yield averages of 10 Mg/ha or more, removing residue increases yield regardless of tillage, but
12 more under no-till than when tilled (Coulter and Nafziger, 2008; unpublished data of the authors
13 since 2007). While our results show the combination of no-till and no residue removal has the
14 highest SOC stocks and WAS in the surface soil, partial or full removal of residues in these
15 systems will lower the SOC levels to those of tilled plots and significantly increase the PR of
16 the soil. No till plots also had the highest BD within the top 30 cm regardless of the level of
17 residue. But even after seven years with residue and tillage treatments in place, we found no
18 consistent degradation of the studied soil properties in response to partial or full removal. Soils at
19 all locations are relatively high in organic matter, which is a key both to their productivity and,
20 we believe, to the fact that they are buffered against effects of removing carbon in the form of
21 residue, hence limiting to some extent the feedstock needed to maintain SOC levels. In any case, we
22 find no reason to believe that removal of at least some of the residue each year in a continuous
23 corn cropping system represents any threat to its sustainability, as long as care is taken to maintain

conditions to prevent loss by erosion. Likewise, within these systems, tillage does not seem to represent a threat to soil quality, even when combined with removal of residue in these high residue systems. However a more in depth and long-term evaluation of these systems is needed to verify this claim.

Acknowledgements

This research was supported in part by the North Central Regional Sun Grant Center at South Dakota State University through a grant provided by the U.S. Department of Energy Office of Biomass Programs #802 DOE Sub SDSU 3TP146, and through a grant from the Illinois Fertilizer Research & Education Council, Project #335. We acknowledge with gratitude the work of Lyle Paul, Eric Adee, Martin Johnson, Mike Vose, Mike Kleiss, Jeff Warren, and Gevan Behnke in plot care and sample collection and analyses.

References

- Allmaras, R.R., Linden, D.R., Clapp, C.E., 2004. Corn-residue transformations into root and soil carbon as related to nitrogen, tillage, and stover management. *Soil Sci. Soc. Am. J.* 68: 1366-1375.
- Balesdent, J., Chenu, C., Balabane, M., 2000. Relationship of soil organic matter dynamics to physical protection and tillage. *Soil Till. Res.* 53: 215-230.
- Blake, G.R., Hartge, K.H., 1986. Bulk density. In: A. Klute, ed. *Methods of soil analysis. Part I: Physical and mineralogical methods*. 2nd ed. ASA and SSSA, Madison, WI. p. 363-375.
- Blanco-Canqui, H., Lal, R., Post, W.M., Izaurralde, R.C., Owens, L.B., 2006. Corn stover impacts on near-surface soil properties of no-till corn in Ohio. *Soil Sci. Soc. Am. J.* 70:266-278.
- Blanco-Canqui, H., Lal, R., 2008. No-tillage and soil-profile carbon sequestration: An on-farm assessment. *Soil Sci. Soc. Am. J.* 72:693-701.
- Blanco-Canqui, H., Lal, R., 2009. Corn stover removal for expanded uses reduces soil fertility and structural stability. *Soil Sci. Soc. Am. J.* 73:418-426.
- Brown, K.H., Bach, E.M., Drijber, R.A., Hofmockel, K.S., Jeske, E.S., Sawyer, J.E., Castellano, M.J., 2014. A long-term nitrogen fertilizer gradient has little effect on soil organic matter in a high-intensity maize production system. *Glob. Change Biol.* 20:1339-1350.

- 1 Buyanovsky, G.A., Aslam, M., Wagner, G.H., 1994. Carbon turnover in soil physical fractions.
2 Soil Sci. Soc. Am. J. 58, 1167–1173.
- 3 Cassel, D.K., Raczowski, C.W., Denton, H.P., 1995. Tillage effects on corn production and soil
4 physical conditions. Soil Sci. Soc. Am. J. 59:1436-1443.
- 5 Clapp, C.E., Allmaras, R.R., Layese, M.F., Linden, D.R., Dowdy, R.H., 2000. Soil organic
6 carbon and ^{13}C -abundance as related to tillage, crop residue, and nitrogen fertilizer under
7 continuous corn in Minnesota. Soil Till. Res. 55:127-142.
- 8 Coulter, J.A., Nafziger, E.D., 2008. Continuous corn response to residue management and
9 nitrogen fertilization. Agron. J. 100: 1774-1780.
- 10 Davies, B.E., 1974. Loss-on-ignition as an estimate of soil organic matter. Soil Sci. Soc. Am.
11 Proc. 38, 150-151.
- 12 Dolan, M.S., Clapp, C.E., Allmaras, R.R., Baker, J.M., Molina, J.A.E., 2006. Soil organic carbon
13 and nitrogen in a Minnesota soil as related to tillage, residue, and nitrogen management.
14 Soil Till. Res. 89:221-231.
- 15 Ellert, B.H., Bettany, J.R., 1995. Calculation of organic matter and nutrients stored in soils under
16 contrasting management regimes. Can. J. Soil Sci. 75, 529-538.
- 17 Energy Independence and Security Act. 2007. Pub. L. No. 110-140. 110th Congr., 1st Sess.
- 18 Fernandez, F., Hoeft R., 2009. Managing soil pH and crop nutrients. In: E.D. Nafziger (ed.) Ch.
19 8. Illinois Agronomy Handbook. 24th ed. University of Illinois Extension. IL. 224 p.
- 20 Gallagher, P.W., Baumes, H., 2012. Biomass supply from corn residues: Estimates and critical
21 review of procedures. Agricultural Economic Report No. 847. USDA Office of the Chief
22 Economist.
- 23 Graham, R.L., Nelson, R., Sheehan, J., Perlack, R.D., Wright, L.L., 2007. Current and potential
24 U.S. corn stover supplies. Agron. J. 99:1-11.
- 25 Gregorich, E.G., Ellert, B.H., Drury, C.F., Liang, B.C., 1996. Fertilization effects on soil organic
26 matter turnover and corn residue C storage. Soil Sci. Soc. Am. J. 60: 472-476.
- 27 Goidts E., Wesemael, B.v., Van Oost, K., 2009. Driving forces of soil organic carbon evolution
28 at the landscape and regional scale using data from a stratified soil monitoring. Glob.
29 Change Bio. 15:2981-3000.
- 30 Hamblin, A.P., 1985. The influence of soil structure on water movement, crop root growth, and
31 water uptake. Adv. Agron. 38:95-158.
- 32 Hammerbeck, A.L., Stetson, S.J., Osborne, S.L., Schumacher, T.E., Pikul Jr., J.L., 2012. Corn
33 residue removal impact on soil aggregates in a no-till corn/soybean rotation. Soil Sci.
34 Soc. Am. J. 76:1390-1398.
- 35 Hooker, B.A., Morris, T.F., Peters, R., Cardon, Z.G., 2005. Long-term effects of tillage and corn
36 stalk return on soil carbon dynamics. Soil Sci. Soc. Am. J. 69:188-196.
- 37 Illinois State Water Survey, 2010. Illinois climate normals. Illinois State Water Survey.
38 <http://www.isws.illinois.edu/atmos/statecli/newnormals/newnormals.htm#stationlist>
39 (accessed 5 Jan 2015).

- 1 Jagadamma, S., Lal, R., Hoef, R.G., Nafziger, E.D., Ade, E.A., 2008. Nitrogen fertilization and
2 cropping system impacts on soil properties and their relationship to crop yield in the
3 central Corn Belt, USA. *Soil Till. Res.* 98:120-129.
- 4 Kay, B.D., VandenBygaart, A.J., 2002. Conservation tillage and depth stratification of porosity
5 and soil organic matter. *Soil Till. Res.* 66:107-118.
- 6 Kemper, W.D., Rosenau, R.C., 1986. Aggregate stability and size distribution. In: A. Klute, (ed.)
7 *Methods of soil analysis. Part I: Physical and mineralogical methods*, 2nd ed. ASA and
8 SSSA, Madison, WI. p. 425-442.
- 9 Kirkby, C.A., Richardson, A.E., Wade, L.J., Batten, G.D., Blanchard, C., Kirkegaard, J.A., 2013.
10 Carbon-nutrient stoichiometry to increase soil carbon sequestration. *Soil Biol. Biochem.*
11 60:77-86.
- 12 Kladvko, E.J., 1994. Residue effects on soil physical properties. In: P.W. Unger (ed.) *Managing*
13 *agricultural residues*. CRC Press, Boca Raton, FL. p. 123-141.
- 14 Konen, M.E., Jacobs, P.M., Lee Burras, C., Talaga, B.J., Mason, J.A., 2002. Equations for
15 predicting soil organic carbon using loss-on-ignition for north central U.S. soils. *Soil Sci.*
16 *Soc. Am. J.* 66:1878-1881.
- 17 Liebig, M.A., Varvel, G.E., Doran, J.W., Wienhold, B.J., 2002. Crop sequence and nitrogen
18 fertilization effects on soil properties in the western Corn Belt. *Soil Sci. Soc. Am. J.*
19 66(2), 596-601.
- 20 Liska, A.J., Yang, H., Milner, M., Goddard, S., Blanco-Canqui, H., Letlon, M.P., Fang, X.X.,
21 Zhu, H., Suyker, A.E., 2014. Biofuels from crop residue can reduce soil carbon and
22 increase CO₂ emissions. *Nat. Clim. Change* 4: 398-401.
- 23 Littell, R.C., Milliken, G.A., Stroup, W.W., Wolfinger, R.D., Schabenberger, O., 2006. *SAS for*
24 *mixed models*. 2nd ed. SAS Inst., Cary, NC.
- 25 Matson, P.A., Parton, W.J., Power, A.G., and Swift, M.J. 1997. Agricultural intensification and
26 ecosystem properties. *Science* 277: 504-509.
- 27 Moebius-Clune, B.N., van Es, H.M., Idowu, O.J., Schindelbeck, R.R., Moebius-Clune, D.J.,
28 Wolfe, D.W., Abawi, G.S., Thies, J.E., Gugino, B.K., Lucey, R., 2008. Long-term effects
29 of harvesting maize stover and tillage on soil quality. *Soil Sci. Soc. Am. J.* 72(4), 960-
30 969.
- 31 Mulvaney, R.L., Khan, S.A., Ellsworth, T.R., 2009. Synthetic nitrogen fertilizers deplete soil
32 nitrogen: A global dilemma for sustainable cereal production. *J. Environ. Qual.* 38:2295-
33 2314.
- 34 Neff, J.C., Townsend, A.R., Gleixner, G., Lehman, S.J., Turnbull, J., Bowman, W.D., 2002.
35 Variable effects of nitrogen additions on the stability and turnover of soil carbon. *Nature*
36 419:915-917.
- 37 Paustian, K., Collins, H.P., Paul, E.A., 1997. Management controls on soil carbon. p. 15–49. In
38 E.A. Paul et al. (ed.) *Soil organic matter in temperate agroecosystems*. CRC Press, Boca
39 Raton, FL.

- 1 Perlack, R.D., Eaton, L.M., Turhollow Jr., A.F., Langholtz, M.H., Brandt, C.C., Downing, M.E.,
2 et al., 2011. U.S. billion-ton update: Biomass supply for a bioenergy and bioproducts
3 industry. U.S. Dep. Of Energy, USDA, Oak Ridge National Laboratory, Oak Ridge, TN.
- 4 Plante, A.L., Parton, W.J., 2007. The dynamics of soil organic matter and nutrient cycling. In:
5 E.A. Paul (ed.) Soil microbiology, ecology, and biochemistry, 3rd ed. Academic Press,
6 Boston, MA. p. 433-467.
- 7 Reicosky, D.C., Evans, S.D., Cambardella, C.A., Allmaras, R.R., Wilts, A.R., Huggins, D.R.,
8 2002. Continuous corn with moldboard tillage: Residue and fertility effects on soil
9 carbon. J. Soil Water Conserv. 57:277-284.
- 10 Russell, A.E., Cambardella, C.A., Laird, D.A., Jaynes, D.B., Meek, D.W., 2009. Nitrogen
11 fertilization effects on soil carbon balances in Midwestern U.S. agricultural systems.
12 Ecol. Applic. 19:1102-1113.
- 13 SAS Institute. 2013. SAS Software v. 9.4 SAS Institute Inc., Cary, NC.
- 14 Singh, N.B., Singh, A., Singh, D., 2010. Autotoxicity of maize and its mitigation by plant growth
15 promoting rhizobacterium *Paenibacillus polymyxa*. Allelopathy J. 25, 195-204.
- 16 Soil Survey Staff. 2014. Official soil series descriptions. Query facility. USDA Natural
17 Resources Conservation Service. <https://soilseries.sc.egov.usda.gov/osdname.asp>
18 (Accessed 6 June 2014).
- 19 Studdert, G.A., Echeverria, H.E., 2000. Crop rotations and nitrogen fertilization to manage soil
20 organic carbon dynamics. Soil Sci. Soc. Am. J. 64:1496-1503.
- 21 Swan, J.B., Schneider, E.C., Moncrief, J.F., Paulson, W.H., Peterson, A.E., 1987. Estimating
22 corn growth, yield, and grain moisture from air growing degree days and residue cover.
23 Agron. J. 79(1), 53-60.
- 24 USDA-NASS. 2013. National corn acres planted. USDA National Agriculture Statistics Service.
25 [http://quickstats.nass.usda.gov/results/08638011-B478-3942-B44F-](http://quickstats.nass.usda.gov/results/08638011-B478-3942-B44F-FA130ADDE283?pivot=short_desc)
26 [FA130ADDE283?pivot=short_desc](http://quickstats.nass.usda.gov/results/08638011-B478-3942-B44F-FA130ADDE283?pivot=short_desc) (Accessed 6 June 2014).
- 27 Varvel, G.E., 1994. Rotation and nitrogen fertilization effects on changes in soil carbon and
28 nitrogen. Agron. J. 86:319-325.
- 29 Wander, M.M., Bollero, G.A., 1999. Soil quality assessment of tillage impacts in Illinois. Soil
30 Sci. Soc. Am. J. 63:961-971.
- 31 White, D.G. (ed.). 1999. Compendium of corn diseases. 3rd ed. Am. Phytopathol. Soc. Press, St.
32 Paul, MN.
- 33 Wilhelm, W.W., Johnson, J.M.F, Hatfield, J.L., Voorhees, W.B., Linden, D.R., 2004. Crop and
34 soil productivity response to corn residue removal: A literature review. Agron. J. 96:1-17.
- 35 Wilts, A.R., Reicosky, D.C., Allmaras, R.R., Clapp, C.E., 2004. Long-term corn residue effects:
36 Harvest alternatives, soil carbon turnover, and root-derived carbon. Soil Sci. Soc. Am. J.
37 68: 1342-1351.
- 38 Zuber, S.M., Behnke, G., Nafziger, E.D., Villamil, M.B., 2015. Crop rotation and tillage effects
39 on soil physical and chemical properties in Illinois. Agron. J. In press.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15

Tillage	Depth cm	SOCs Mg/ha	SOCc %	WAS %	BD Mg/m ³	PR kPa
No-till	0-15	49.6 a	2.20 a	79.4	1.50 b	900
Chisel tilled		45.4 c	2.18 a	81.0	1.38 c	449
No-till	15-30	48.3 b	2.10 b	80.7	1.52 a	1380
Chisel tilled		47.5 b	2.11 b	81.9	1.49 b	948
No-till		49.0	2.15	80.8	1.51	1140 a
Chisel tilled		46.5	2.14	81.5	1.44	699 b

Table 1. Probability values and numerator degrees of freedom (df) associated with the different sources of variation in the analysis of variance for soil organic carbon stock (SOCs, Mg/ha), and concentration (SOCc, %), water aggregate stability (WAS, %), soil bulk density (BD, Mg/m³), and penetration resistance (PR, kPa), measured across four sites after seven cropping seasons following initiation of the experiment.

	df	SOCs	SOCc	WAS	BD	PR
Source of variation						
Residue Removal (R)	2	0.3997	0.3269	0.6292	0.7408	0.1194
Tillage (T)	1	0.0076	0.7894	0.3604	<.0001	<.0001

R x T	2	0.0021	0.0005	0.9602	0.8705	0.0105
N rates (N)	1	0.4385	0.4651	0.2041	0.7858	0.1221
R x N	2	0.5481	0.1536	0.1771	0.0428	0.9943
T x N	1	0.8677	0.3252	0.2184	0.1570	0.8921
R x T x N	2	0.9653	0.9444	0.3639	0.2816	0.3528
Depth (D)	1	0.1748	<.0001	0.2027	<.0001	<.0001
R x D	2	0.0561	<.0001	0.3796	0.1222	0.3003
T x D	1	<.0001	0.0281	0.7891	<.0001	0.6203
R x T x D	2	0.1556	0.2410	0.8105	0.1705	0.6573
N x D	1	0.439	0.1344	0.2893	0.8481	0.0221
R x N x D	2	0.1491	0.4138	0.0026	0.2846	0.1660
T x N x D	1	0.847	0.8407	0.7948	0.8782	0.8037
R x T x N x D	2	0.5037	0.9528	0.1561	0.5490	0.9479

Table 2. Interaction effects of residue removal and tillage on total soil C stock (SOCs), soil C concentration (SOCc), water aggregate stability (WAS), bulk density (BD), and soil penetration resistance (PR) in the top 30 cm of soil, across four Illinois sites following seven cropping seasons with treatments in place. Means within a column with the same lower case letter are not significantly different at $\alpha=0.05$; no letters indicate no significant differences were detected.

Residue removal	Tillage	SOCs Mg/ha		SOCc %		WAS %	BD Mg/m ³	PR kPa	
None	No-till	52.1	a	2.29	a	80.9	1.51	1002.3	b
	Chisel tilled	45.1	b	2.08	cd	82.6	1.44	724.5	c
Partial	No-till	48.1	b	2.12	bcd	79.0	1.51	1167.5	a
	Chisel tilled	46.5	b	2.15	bc	80.8	1.44	697.5	c
Full	No-till	46.7	b	2.04	d	80.4	1.52	1249.4	a
	Chisel tilled	47.7	b	2.20	ab	81.2	1.44	674.5	c