Title: Long-term agricultural management maximizing hay production can significantly reduce belowground C storage

Author names and affiliations: LENKA SOCHOROVÁ^{a, b *}, JAN JANSÁ^b, ERIK VERBRUGGEN^c, MICHAL HEJCMAN^a, JÜRGEN SCHELLBERG^d, E. TOBY KIERS^e, NANCY COLLINS JOHNSON^f

^aDepartment of Ecology, Faculty of Environmental Sciences, Czech University of Life Sciences, Kamýcká 1176, Prague 6 CZ-165 21, the Czech Republic

^b Institute of Microbiology, Academy of Sciences of the Czech Republic, Vídeňská 1083, Prague 4 CZ-142 00, the Czech Republic

^cDepartment of Plant and Vegetation Ecology, University of Antwerp, Universiteitsplein 1, Antwerpen BE-2610, Belgium

^d Institute of Crop Science and Resource Conservation, University of Bonn, Katzenburgweg 5, Bonn D-53115, Germany

^e Department of Ecological Science, Faculty of Earth and Life Sciences, Vrije Universiteit, De Boelelaan 1085-1087, Amsterdam NL-1081HV, the Netherlands

^f School of Earth Sciences and Environmental Sustainability and Department of Biological Sciences, Northern Arizona University, PO Box: 5694, Flagstaff AZ 86011, the USA

Corresponding author: Lenka Sochorová, Telephone: +420 224 383 781, Fax: +420 296 442 384, e-mail: lsochorova@fzp.czu.cz

Abstract

Liming and fertilization of grasslands have been used for centuries to sustain hay production. Besides improving hay yields, these practices induce compositional shifts in plant and soil microbial communities, including symbiotic arbuscular mycorrhizal (AM) fungi. However, in spite of increasing interest in soil carbon (C) sequestration to offset anthropogenic CO₂ emissions, little is known about the long-term effects of these agronomic interventions on soil C stocks. We examined how plants, AM fungi, and soil C respond to more than seven decades of annual applications of lime, mineral nitrogen (N), and mineral phosphorus (P) to test the hypotheses that (1) management practices increasing aboveground plant production decrease C allocation to roots, AM fungi and the soil; and (2) the relative availability of N and P predicts belowground C allocation in a consistent manner. Our study was conducted at the Rengen Grassland Experiment, established in 1941. Lime combined with N increased hay yields and promoted development of AM fungal hyphae in soil, while reducing relative C allocation to roots. Simultaneous enrichment of soil with lime, N, and P further boosted hay production, promoted grasses and suppressed other plant functional groups. This treatment also decreased soil organic C and strongly suppressed AM fungi in the soil, although the response to P varied among different AM fungal taxa. Our results indicate that agricultural management practices aimed at maximization of hay production may, in the long run, significantly (-20%) reduce belowground C storage. This is a great concern with respect to the intended use of grasslands as anthropogenic CO₂ sinks because the fertilization-induced decrease in soil C stocks can partly or fully negate the C sequestration potential of the grassland ecosystems as a whole.

Keywords: arbuscular mycorrhiza, nitrogen, phosphorus, lime, soil carbon, long term ecological research

1. Introduction

Grasslands cover about 40% of the terrestrial surface of the planet and about half of this area is currently used in agricultural production (Lal, 2007). Because soils contain more organic carbon (C) than both the atmosphere and biota combined and because grassland soils generally have much larger C storage capacity than annually cropped soils, grasslands are extremely important for global soil C sequestration (Conant *et al.*, 2001; Lal, 2013). Long-term productivity of grasslands is influenced by many factors including climate, geographic location, and management practices, and particularly by the level of fertilization (Sala *et al.*, 1988; Conant *et al.*, 2001). For sustainable hay production, soil organic matter is critical because it supplies plants with nutrients, helps to reduce soil erosion, and increases cation exchange and water holding capacity (Schulten & Schnitzer, 1997; Daynes *et al.*, 2013). Consequently, management practices that restore and/or maintain high levels of soil organic matter are desirable. This in combination with recent interest in soil C sequestration, as a means to offset rising CO₂ levels in the atmosphere, has led to a keen interest in optimizing grassland management for C storage (Conant & Paustian, 2002; Lal, 2008). An earlier review of 115 studies from around the world demonstrated that grasslands fertilization, especially with nitrogen (N), tends to be associated with increased soil C (Conant *et al.*, 2001; van Groenigen *et al.*, 2006), suggesting that fertilization is potentially an important driver of soil C sequestration. However, the relationship is difficult to predict because it may critically depend on the type of fertilizer, and is likely to be nonlinear (Wander *et al.*, 1994). One reason for this is that belowground C allocation by the plants and subsequent C transformations within soils likely depend on plant and microbial responses to relative, not absolute nutrient limitations (Cleveland & Liptzin, 2007; Kaiser *et al.*, 2014).

The soils and roots of plants in natural and managed grasslands are abundantly colonized by arbuscular mycorrhizal (AM) fungi. These fungi, belonging to the phylum Glomeromycota, form symbiotic relationships with plants that are often mutually beneficial for both partners. Plant hosts supply carbohydrates in exchange for fungal provisioning of mineral nutrients, especially phosphorus (P) and zinc, some of which would otherwise be unavailable for plants (Allen, 1991; Jansa *et al.*, 2003). Besides facilitating nutrient uptake, AM fungi can provide

benefits to colonized plants, such as resistance against pathogens and drought tolerance (Newsham *et al.*, 1995; Smith *et al.*, 2010). High densities of AM fungal hyphae in soil form underground networks that contribute to creation and stability of soil aggregates and reduce soil losses through erosion (Chaudhary *et al.*, 2009). Because many of these features are considered beneficial for the sustainability of agricultural production, there is a great interest in increasing mycorrhizal symbioses through agricultural management (Gianinazzi *et al.*, 2010; Fitter *et al.*, 2011).

Mycorrhizal fungi are key components of grassland C sequestration (Treseder & Allen, 2000; Zhu & Miller, 2003; Johnson *et al.*, 2006; Wilson *et al.*, 2009). Hyphae of AM fungi may comprise a large fraction (estimates go up to a half) of microbial biomass in grassland soil (Olsson *et al.*, 1999), and therefore their responses to fertilization are highly relevant to soil C dynamics. Although AM fungi are expected to rapidly channel and immobilize a significant portion of plant-derived C through the soil, there is an on-going debate about the importance of this C highway in the buildup of stable soil organic matter and soil C sequestration (Staddon *et al.*, 2003; Phillips *et al.*, 2013; Averill *et al.*, 2014; Balasooriya *et al.*, 2014). Density of AM fungal hyphae in the soil was previously found to be positively correlated with the responses of soil organic C to long-term N enrichment and burning in a North American tall-grass prairie, while application of fungicides suppressed AM fungal hyphae and reduced soil organic carbon (Wilson *et al.*, 2009). Responses of AM fungi to N fertilization have been shown to vary with availability of soil P. For example, N enrichment of P deficient grasslands often increases AM fungal biomass but N enrichment of P rich grasslands often decreases it (Johnson *et al.*, 2003; Wilson *et al.*, 2009; Liu *et al.*, 2012).

The functional equilibrium model can help explain why N enrichment may either increase or decrease AM fungal biomass. This conceptual model states that plants should preferentially allocate photosynthates to structures that acquire the most limiting resources; plants growing in infertile soil should allocate more to roots and mycorrhizas and plants growing in nutrient rich soil should allocate more to shoots and leaves (Brouwer, 1983; Johnson *et al.*, 2003). Increasing soil N availability in P-limited soils exacerbates P limitation and enhances the importance of symbiotically acquired P; thus plants are expected to increase allocation of photosynthates belowground to AM fungi. In contrast, N enrichment of P-rich soils is likely to

remove limitation by any belowground resource and, in accordance with the functional equilibrium model, plants are expected to shift allocation of photosynthates aboveground to shoots and away from AM fungi (Johnson, 2010).

Changes in nutrient availabilities and consequent shifts in C allocation by plants can have large effects on AM fungal communities (Egerton-Warburton *et al.*, 2001, 2007; Liu *et al.*, 2012). Application of fertilizers systematically increases plant productivity, but also reduces plant diversity (Silvertown *et al.*, 2006; Hejerman *et al.*, 2007; Semelová *et al.*, 2008). Generally, the most discussed nutrients in relation to the management of grasslands are N, P, and calcium (Ca), of which the last mentioned is most often provided in carbonate form (lime) not to primarily correct for plant nutrient deficiencies, but to reverse soil acidification, developing mainly due to application of N fertilizers (Johnston *et al.*, 1986). Soil acidification is recognized as one of the main limits of agricultural production and provision of ecosystem services on a wide range of soils (e.g., Scott *et al.*, 2000). The effect of liming on AM fungi is not easy to predict, in some cases the application of Ca promoted AM fungal colonization of roots (Johnson *et al.*, 2005), whereas in others it had no effect (Kennedy *et al.*, 2005). Recent unconstrained monitoring of AM fungal communities in a number of agricultural soils (including grasslands) in Switzerland pointed to a particular importance of soil pH in structuring indigenous AM fungal communities (Jansa *et al.*, 2014), which provides some mechanistic explanation on how could induced soil pH shifts (due to liming of acidic soils) affect the indigenous AM fungi. The effects of fertilization on individual AM fungal taxa are highly context-dependent and are often confounded with the effects of soil disturbance and crop rotation (Jansa *et al.*, 2006 and references therein). Individual taxa of AM fungi differ in their responses to fertilization; some species increase and others decrease in abundance in response to nutrient enrichment (Douds & Schenck, 1990; Johnson, 1993; Egerton-Warburton *et al.*, 2001).

The aim of this study was to analyze the responses of soil C and AM fungi to 70 years of annual applications of lime and mineral fertilizers at one of the oldest agricultural experiments worldwide – the Rengen Grassland Experiment. Our goal was to test the ability of the functional equilibrium model to predict responses of soil C stocks to grassland inputs. We examined the abundance of AM fungi inside and outside plant roots and used molecular techniques to determine the abundance of a few individual AM fungal taxa. This allowed us to test the

hypothesis that lime and fertilizers would reduce plant allocation of photosynthates to belowground structures including the mycorrhizas. We predicted that the differences of lime and fertilizer inputs among treatments would have consequences for the long-term buildup of soil C stocks, co-incident with the effects on AM fungi: N enrichment will increase allocation to AM fungi due to increasing relative demand for limiting P, while fertilization with both N and P will decrease allocation to AM fungi, consequently reducing the soil C sequestration.

2. Material and methods

2.1. Study site description and experimental design

This study was conducted at the Rengen Grassland Experiment established in 1941 by the University of Bonn in low productivity *Nardetum* grassland in the Eifel Mountains, Germany (50°13' N, 6°51' E). The site is 475 m above sea level, average annual precipitation is 811 mm and the mean annual temperature is 6.9°C. Soil type is a pseudogley. The experimental plots (3 m × 5 m) received annual amendments of lime (in the form of Ca(OH)₂, obtained by combination of CaO with water) and fertilizers for more than seven decades, reaching levels considered sufficient for sustainable hay production in that area (Wendland *et al.*, 2012). We sampled four treatment combinations (average inputs in kg ha⁻¹ yr⁻¹): Ca (715 Ca), CaN (100 N, 752 Ca), CaNP (100 N, 35 P, 936 Ca), and a control (unfertilized plots). The different amounts of Ca in the different treatments were due to the forms of applied fertilizers (such as Ca₃PO₄ in the CaNP treatment, which were not counterbalanced by reducing lime inputs in the same treatment as compared to the other limed treatments). The experimental plots are arranged in two blocks containing each five randomly arranged replicates of the Ca, CaN and CaNP treatments, totaling 30 experimental units with ten replicate plots per treatment. No differences in the underlying geology or soil properties were detected between the two blocks when comparing concentrations of available P, total N, organic C, mobile Ca in the soil and the soil pH (analyses not shown), justifying pooling the replicates across the two blocks. Since the original field experiment did not contain non-fertilized controls, ten control plots were designated as late as 1998 on adjacent grassland strip below one of the blocks. This grassland strip has never been fertilized or grazed, it has been subjected to 2 hay cuts since 1941, but for historical reasons the plots are not spatially randomized with the other treatments. However, no significant soil gradient within soil properties has been found in the experimental area either between the blocks (see above) or along the axis perpendicular to the block arrangement, except the soil pH, which showed a significant shift with the distance from the control plots, driven by the strong difference in soil pH between the controls and the other (limed) plots (see Table 1). These analyses justified inclusion of the control plots to the comparison with the other soil treatments even if they were not perfectly randomized with the other treatments in the experimental field.

Further details of the site and the experimental design have been published previously (Schellberg *et al.*, 1999; Hejzman *et al.*, 2007).

2.2. Sampling procedure and plant analysis

Experimental plots were cut twice a year, first in early June and again in mid-October. Aboveground biomass for laboratory analyses was collected along a strip 1.25 m wide and 5 m long and weighted. Subsamples of 500 g (fresh weight) from each of the plots were dried at 60°C for 48 h to determine the dry biomass (hay yield). The percentage of surface cover by individual vascular plant species was estimated visually in twenty plots (i.e., one experimental block and half of the control plots) on 20 June 2011 according to a regional plant species list (Rothmahler *et al.*, 2000), considering three plant functional groups (grasses, herbs (i.e., dicots without legumes) and legumes). The nomenclature of plant species was based on Kubát *et al.* (2002). To eliminate edge effect, relevés were taken in the center of each plot, considering an area of 1.8 m × 3.2 m.

Three types of belowground samples were collected: (1) composite soil+root samples, (2) root standing biomass, and (3) new root growth. All samples were transported to the Czech University of Life Sciences in Prague and frozen within 24 hours of collection.

(1) On 20 October 2011, eight small diameter cores (2.5 cm diameter × 10 cm depth) were collected across the central area of each plot and mixed into one composite sample. Roots recovered from a portion of each composite sample were analyzed for AM fungal colonization and the remaining soil from the composite samples was analyzed for extraradical (soil-borne) AM fungal hyphae and other soil properties as described below.

(2) At the same time, one large (5 cm diameter) core was taken from the center of each plot to measure the standing root biomass. The deepest possible core was taken and the resulting core depths varied from 10 to 15 cm. Each core was weighed and its exact length was recorded. A subsample of each core was used to estimate gravimetric soil moisture by weighing the soil sample before and after drying at 70°C for 72h. The rest of the core was used to extract root biomass (see below). Each hole created through this sampling was immediately filled with granitic river sand.

(3) In mid June 2012, the area was re-sampled using the same corer and sampling depth as above (point 2) to estimate new root production during eight months of root growth.

Roots were removed from the large core samples by soaking them in soapy water for 5 to 10 min, and then rinsed with tap water through a 1mm sieve. Roots were also carefully removed from the new growth (sand) cores by rinsing with tap water through a 1 mm sieve. Cleaned roots were weighed, dried at 80°C for 48 h and weighed again. Standing root biomass in the topsoil (10 cm depth) per ha was calculated by using estimated root biomass ($\times 2$) and the soil core volume. Root-to-shoot biomass ratio was then calculated as the ratio of dry weight of roots (t ha^{-1}) and annual yield (t ha^{-1} , average value from years 2008 through 2012).

2.3. Determination of AM fungal colonization and extraradical hyphal length

The extent of root colonization by AM fungi was determined in the roots from the composite soil samples. Roots were soaked in 5% KOH at room temperature for two days, and stained using the ink and vinegar technique (Vierheilig *et al.*, 1998). Stained roots were examined for mycorrhizal colonization using the magnified intersection method, scoring 100 intersections per sample (McGonigle *et al.*, 1990) at a magnification of 200 $\times$. Then we calculated percentages of root length colonized by arbuscules, coils, vesicles, mycorrhizal hyphae, and the fraction of root length devoid of any mycorrhizal structures.

Composite soil samples were thawed and a 5 g subsample was used for determination of extraradical hyphal length density following the protocol described in Rillig *et al.* (1999), with minor modifications. In short, soil was suspended in a solution containing 112 ml of deionized water and 420 mg of $\text{Na}(\text{PO}_4)_6$, shaken for 30 s, allowed to settle for 30 min, and decanted into a 38 μm sieve. All material on the sieve was suspended in 200 ml of deionized water, shaken, and 3 ml was transferred onto a membrane filter with a pore size of 0.45 μm (GN-6 25mm; Pall Life science, Mijdrecht, the Netherlands), placed in a vacuum manifold. Material on the filter was stained with 0.02% trypan blue in lactoglycerol for 5 min, washed with excess water and the filter was then mounted on microscopic slides. AM hyphae were distinguished from other hyphae according to descriptions in Mosse (1959): “dark-to light-blue stained aseptate hyphae with characteristic unilateral angular projections”. Intersects in 30 random grids were counted

using a compound microscope at 200× magnification. Hyphal length was calculated per g soil dry weight by estimating soil water loss after drying in an oven at 80 °C for four days.

2.4. Real-time PCR

A subsample of each composite soil sample was dried at 80°C for 48 h, homogenized, and sieved through a 2 mm sieve. The soil DNA was extracted from 500 mg (dry weight) samples using the Macherey-Nagel NucleoSpin Soil DNA kit (Macherey-Nagel GmbH & Co., Düren, Germany) following the standard protocol and stored at –20°C until use. Abundances of six AM fungal taxa were estimated using quantitative real-time polymerase chain reaction (qPCR) with specific AM fungal primers and hydrolysis (TaqMan) probes, using an AB StepOnePlus instrument (Applied Biosystems, Foster City CA, USA.). We employed taxon-specific markers targeting the nuclear large ribosomal subunit (nLSU) genes (Wagg *et al.*, 2011; Thonar *et al.*, 2012) for *Cetraspora pellucida*, *Claroideoglomus claroideum*, *Funneliformis mosseae*, *Diversispora celata*, *Gigaspora margarita*, and *Rhizophagus irregularis*. Additionally, for detection of *Rhizophagus* sp., we also used a marker targeting the mitochondrial large ribosomal subunit (mtLSU) gene (Kiers *et al.*, 2011; Couillerot *et al.*, 2012). Twenty billion copies of an internal standard (linearized plasmid carrying a fragment of cassava mosaic virus, GenBank accession AJ427910) were added into each sample before extraction, and quantified after the extraction, to estimate the DNA extraction efficiency and DNA extract quality, namely absence of PCR inhibitors (Thonar *et al.*, 2012; Janoušková *et al.*, 2015). Each qPCR reaction mixture contained 2 µl template (DNA extracts diluted 1:10), 0.4 µl forward primer (25 µM), 0.4 µl reverse primer (25 µM), 0.1 µl hydrolysis probe (25 µM), 4 µl of 5× HOT FIREPol® PROBE qPCR Mix Plus (ROX) (Solis BioDyne, Tartu, Estonia), and 13.1 µl PCR-grade water.

Cycling conditions were as follows: initial DNA denaturation and DNA polymerase activation at 95°C for 15 min, then 50 cycles each with denaturation at 95°C for 10 s, annealing conditions specified in Table S1 (electronic supplement), and elongation at 72°C for 10 s.

2.5. Soil properties

To measure soil pH, 30 g of dry soil was suspended in 30 ml deionized water for 20 minutes, centrifuged at 3044 g for 20 minutes and the supernatants (soil extracts) were filtered through a

paper filter to remove floating particles. The pH was determined using a combination pH electrode G-P Combo w/RJ (Corning, USA). The concentration of mobile Ca^{2+} ions in the supernatants brought to pH 7.0 ± 0.2 (using 1% HCl or KOH) was measured using a Ca-selective electrode (type 20-35, Monokrystal, Turnov, Czech Republic) and corrected for acidic error using the calibration of the electrode response with solutions of known Ca concentrations at different pH values. Soil organic C and total N concentrations were analyzed using a CN analyzer (Flash EA 2000, ThermoFisher Scientific, Waltham MA, USA). Soil available P was determined spectrophotometrically in aqueous soil extracts (3 g dry soil and 30 ml of water) after shaking for 25 hours, subsequent centrifugation at 3044 g for 20 min and filtration through a 0.22 μm filter (CME, Carl Roth, Karlsruhe, Germany). This approach to measure P availability was chosen because of large differences in soil pH between the different soil application treatments. Soil pH can namely largely affect the estimates of P availability in soils when assessed by various extraction methods (Demaria *et al.*, 2005).

2.6. Bulk density and soil carbon stocks

Bulk density was determined from the large soil cores (5 cm diameter). The entire core was weighted fresh, and a half of the core was then taken, weighted, dried at 70°C for 3 days and weighted again. The ratio of fresh to dry weights of the soil subsample was used to calculate dry weight of the entire soil core of known dimensions. In order to determine the amount of organic C per ha, the volumetric organic C content of the topsoil was calculated by using gravimetric organic C concentrations measured on the composite soil samples (cores with 2.5 cm diameter, dried and sieved to pass 2 mm sieve) and the soil density assessed on the large soil cores, and scaling up to $1 \times 10^6 \text{L}$ of soil (1 ha surface, 10 cm depth). Both gravimetric and volumetric C concentrations were used to compare the differently treated soils whereas the volumetric C concentration was then used to establish a relationship with the extraradical hyphal length density. This is because a gravimetric approach would artificially amplify the differences due to the well-established fact that more organic soils have a lower bulk density (Aggelides *et al.*, 2000; Celik *et al.*, 2004; Hati *et al.*, 2007).

2.7. Statistical analyses

Prior to analysis, root colonization data (percentage values + 1) were square root-arcsine transformed (Zar, 1999). To evaluate differences between treatments, one-way ANOVA in STATISTICA 8.0 (Statsoft, Tulsa, OK, USA) and an additional post-hoc test using the least significant difference F-test ($p = 0.05$) was used. Control plots were considered as the fourth treatment in addition to the Ca, CaN, and CaNP for statistical comparisons, given the apparent absence of underlying soil gradients on the experimental grounds (see above). Redundancy Analysis (RDA) and Canonical Correspondence Analysis (CCA) in the CANOCO 4.5 program (terBraak & Šmilauer, 2002) were used to examine the relationships among AM fungal variables, soil properties, and hay yields. Further, a Monte Carlo permutation test with 499 permutations was used to reveal if the tested explanatory environmental variables (soil C, available soil P, total soil N, available soil Ca, soil pH, and hay productivity at the plot level) had a significant impact on the AM fungal community composition and abundance. Results of RDA and CCA were visualized in an ordination diagram constructed by the CanoDraw program. Simple linear regression was used to examine the relationships between the volumetric organic C content of the topsoil, density of AM fungal hyphae in soil and hay production.

3. Results

3.1. Soil properties

Long-term fertilization and liming caused large changes in soil chemical properties across the different management treatments (Tables 1 and 2). Plant available P in soil was an order of magnitude higher in P fertilized plots compared to the other treatments (Table 1). Long-term liming, with or without fertilizer additions, considerably increased soil pH. It increased with liming and fertilizer from 5.32 in control plots to 7.14 in CaNP plots (Table 1, Table S2). The fertilization had a profound effect on the total N and organic C concentrations in the soil, but neither of the practices had significantly changed mobile Ca levels. Soil bulk density varied between individual samples from 0.83 to 1.18 kg L⁻¹, and was significantly lower in the unfertilized control treatment as compared to the CaNP treatment (Table 2, Table S2).

3.2. Plant responses to lime and fertilizers

Hay yield was significantly increased by application of lime and the N and P fertilizers ($F = 118.8$, $p < 0.001$; Fig. 1a). The standing biomass of roots in October 2011 tended to have lower values in the CaN treatment as compared to the other treatments, yet the differences remained insignificant ($F = 2.06$, $p = 0.123$; Fig. 1b, Table S3). Root biomass in June 2012 showed similar pattern as in October 2011, with no significant differences between treatments ($F = 1.58$, $p = 0.21$), but the values were about 3 times lower than those in October 2011 (data not shown). Root-to-shoot biomass ratio was higher in the unfertilized control and in the limed treatments than in the CaN and CaNP treatments ($F = 13.8$, $p < 0.001$; Fig. 1c). Plant species richness decreased with the complexity of soil amendments, with significantly fewer species being found in the CaNP plots than the other treatments ($F = 11.1$, $p < 0.001$; Fig. 1d); relative cover of herbs tended to decrease ($F = 3.16$, $p = 0.0535$; Fig. 1e), grass cover increased ($F = 5.22$, $p = 0.011$; Fig. 1f), and cover of legumes decreased ($F = 5.53$, $p = 0.015$, data not shown) with increasing complexity of the soil amendments.

3.3. Mycorrhizal responses to soil amendments

The percentage of root length colonized by any AM fungal structures varied significantly among

the treatments ($F = 3.18$, $p = 0.036$; Fig. 2a) and was lower in the CaNP treatment than in the unfertilized control. The colonization of roots in the CaN treatment showed similarly high values as in the unfertilized control plots (Fig. 2a, Table S4). A similar picture was observed when comparing the impact of the management practices on the rates of root colonization by specific AM fungal structures (Fig. 2b – d), but only the coils (Fig. 2d) showed statistically significant ($F = 3.64$, $p = 0.022$) decrease in abundance in the CaNP as compared to the other treatments. The density of extraradical AM fungal hyphae in the soil was significantly higher in the CaN plots as compared to the control and the CaNP treatments, with the latter having the lowest hyphal density among all the management treatments ($F = 5.91$, $p = 0.002$; Fig. 2e). According to the CCA (Monte Carlo permutation test $p = 0.018$; Fig. S1), root colonization and density of AM fungal hyphae in the soil were negatively related to available soil P and hay yield and positively related to the N and C concentrations in the soil.

The qPCR profiling of individual AM fungal taxa abundance in soil yielded positive detection between 30% (*Claroideoglossum*) and nil (*Cetranspora*) of the analyzed samples (details not shown). There were either no effects of soil management on the abundance of the tested AM fungal taxa or the abundance in the limed and/or fertilized treatment was higher than in the control plots (Fig. S1). Additionally, there was an obvious difference in the response to P fertilizer between *Claroideoglossum*, whose abundance decreased in response to P fertilization as compared to N fertilization only, and *Diversispora*, which was not significantly affected by P fertilization (Fig. S2 a, b). Multivariate analysis (Fig. 3) showed that the abundances of the tested AM fungal taxa were significantly affected by soil properties (Monte Carlo permutation test $p = 0.02$) and mostly driven by soil amendments, positively correlating with pH (*Rhizophagus*, *Claroideoglossum*, *Diversispora*), yields and/or soil P levels (*Funnelformis*).

3.4. Soil C stocks

The amount of soil organic C in the top 10 cm was significantly increased by liming without mineral fertilizer amendment, and significantly decreased by simultaneous addition of lime, N and P as compared to the unfertilized controls ($F = 12.7$, $p < 0.001$; Fig. 2f). There was also a strong decrease in the soil C/N ratio with increasing complexity of amendments (Table 2). Soil C/N ratios were higher in the control and limed treatments as compared to the other treatments,

with the values being still significantly higher in the CaN than in the CaNP treatments (Table 2). Volumetric organic C content of topsoil had a negative relationship with hay yields ($R^2 = 0.21$, $p = 0.003$) and a positive relationship with the density of AM fungal hyphae in the soil ($R^2 = 0.22$, $p = 0.002$; Fig. 4), whereas it showed no significant relationship with any of the AM fungal abundance estimates inside the roots (analyses not shown).

4. Discussion

Our results show that seven decades of yield-promoting measures, particularly N and P fertilization, dramatically reduced soil C stocks in a temperate grassland. This reduction in the top 10 cm is about 6 tons C ha⁻¹ when comparing the fully fertilized treatment with the unmanaged control. There is even a higher reduction of 11 tons C ha⁻¹ when the fully fertilized treatment is contrasted against the lime-only treated plots. This latter comparison represents a reduction of 20.4% of the C stocks in the topsoil of the limed-only plots as compared to CaNP treatment, which is more than half of the figure to be expected due to land use change from grassland to cropland (-36%, Poeplau *et al.*, 2011). The C loss from the topsoil of the CaNP treatment over the last 70 years outweighed the C contained in the increased annual hay yield by 6- and 12-fold when compared to the control with no inputs and to the limed treatment, respectively. This means that increased hay yields over 6 to 12 subsequent years are just compensated by reduction of belowground C stocks within the top 10 cm. This comparison takes into account average hay yields over the last fifteen years, assuming 45% C content of hay. Due to lack of records, we are unable to draw a full C balance of the system over the entire 70 years duration of the experiment or for soil depths greater than 10 cm. Nevertheless, the magnitude of the observed effect calls for a very careful consideration of the design of grassland management practices that aim at both maximizing hay yield and enhancing belowground C sequestration. There is an urgent need to better understand the mechanisms that control the accrual of soil C stocks in agricultural systems and, based on this information, to design highly productive agrosystems with high C sequestration potential.

Photosynthesis is the source of the vast majority of C in grassland soils, and thus one would assume that the greater the productivity, the greater the C remaining in the system. There are, however, many complicating factors: the processes of decomposition and mineralization determine the ultimate fate of soil organic C; and the stoichiometry of C, N, and P within the system has a strong influence on the complex interactions involved in these processes. Organic C enters the soil as root or shoot debris (necromass), root exudates, AM fungal biomass (trophically completely dependent on host plants) or AM fungal necromass. Thereafter, organic C enters the belowground food web involving soil animals, fungi and prokaryotes. At every

414 trophic step, a portion of organic C is lost as CO₂ and a part is stabilized in the soil over shorter
415 or longer periods of time. As a result, realized soil C stocks depend on plant productivity (i.e.,
416 input quantity), biomass partitioning between roots and shoots, degree of association of plants
417 with AM fungi, degradability of plant and/or AM fungal necromass (i.e., input quality), activity
418 of soil microbes, and competition among plants, AM fungi, and soil saprobes for essential
419 resources such as N and P, as well as fluctuations of environmental properties such as
420 temperature and moisture (Cotrufo *et al.*, 2013). Theoretically, plants and associated AM fungi
421 could be seen as competitors of soil saprobes for available P and/or N thus suppressing their
422 decomposition activity (Leifheit *et al.*, 2015). Alternatively, AM fungi and fine roots could
423 prime soil saprotrophic life with fresh C from aboveground (Murphy *et al.*, 2015), thus
424 increasing activity of soil saprobes and eventually reducing the soil C stocks. Below, we will
425 describe three non-mutually exclusive mechanisms that may explain the decline of soil C that
426 was associated with P fertilization in our study: 1) allocation to roots and AM symbioses, 2)
427 degradability of litter inputs, and 3) stoichiometric control of soil microbial activity.

429 *4.1. Allocation to roots and AM symbioses*

430 The functional equilibrium model predicts that plant allocation to roots will decrease when the
431 plants are not limited by mineral nutrients (Brouwer, 1983; Nielsen *et al.*, 2001; Hermans *et al.*,
432 2006). Here we showed that the root-to-shoot biomass ratio significantly decreased in the CaN
433 and CaNP treatments, corroborating the idea that plants allocate biomass to the organs required
434 to gain access to the most limited resources. Enrichment with N or with N and P together
435 increased relative allocation of biomass/C aboveground to shoots and reduced allocation to roots,
436 indicating greater limitation of plants' growth by photosynthesis than by nutrients in those soil
437 management treatments (Fig. 1). The functional equilibrium model applies to plant allocation to
438 AM symbioses as well as roots (Johnson *et al.*, 2003; Johnson *et al.*, 2013). It is interesting to
439 note that in our study allocation to roots was reduced in both the CaN and CaNP treatments,
440 while allocation to AM fungi (soil hyphae) was only reduced in the CaNP treatment (Fig. 2e).
441 This strongly suggests that AM symbioses are critical for the P nutrition while roots are critical
442 for the N nutrition of plants.

The opposite relationships of hay yield and AM hyphae with soil C suggests that plant allocation to shoots (hay) and away from AM symbioses may have a large (and possibly a causal, AM-mediated) impact on soil C storage (Fig. 4). The positive correlation between soil C and AM fungal hyphae could arise from increased C inputs via hyphal biomass and necromass as well as increased C storage through enhanced soil aggregation and hence protection against degradation (Zhu & Miller, 2003; Rillig, 2004; Leifheit *et al.*, 2015), as well as from lower microbial activity in the unfertilized soil due to low P availability for the microbes, which could further be enhanced by direct competition for resources like available P between AM fungi and other soil biota (van Groenigen *et al.*, 2006; Welc *et al.*, 2010). Other studies have also reported a positive correlation between AM fungal abundance and soil C levels (Treseder & Allen, 2000; Johnson *et al.*, 2006; Wilson *et al.*, 2009). Future research is needed to elucidate the mechanisms generating these correlations. The degree to which AM fungi and their associated microbes increase soil C stocks though direct input of recalcitrant forms of C versus their influence on soil aggregate structure (and protection from decomposition) is currently unknown. It is important to recognize that in some systems, AM fungi might actually reduce soil C levels by stimulating (nutrient rich) organic matter decomposition by other soil microbes (Hodge *et al.*, 2001; Cheng *et al.*, 2012). This idea is also supported by the view that most C allocated to the AM fungi flushes through the ecosystem and is respired back to the atmosphere within few days (Johnson *et al.*, 2002; Staddon *et al.*, 2003; Lendenmann *et al.*, 2011). Depending on the relative importance of either of these processes (increased C inputs and stabilization or stimulation of microbial decomposition), the effects of AM fungi on C stocks may be positive, negative or neutral (Verbruggen *et al.*, 2013). From our correlative findings we can simply conclude that AM fungi thrive in more organic soil. Indeed, some common organic compounds (i.e., chitin or cellulose) may significantly affect the presence and abundance of AM fungi in soil (Gryndler *et al.*, 2003; Joner & Jakobsen, 1995).

The community composition of plants and AM fungi was strongly influenced by the different fertilizer treatments. Apart from stimulating mycorrhizal fungi in general (Fig. S1), application of lime together with N increased the abundance of AM fungal taxa *Claroideoglossum* and *Diversispora* (Fig. S2). In contrast, P enrichment drove a sharp decrease of extraradical AM hyphae in the soil, although the specific response of the individual fungi to P application

differed, with some taxa such as *Diversispora celata*, not responding at all to addition of P (Fig. S2). This may suggest that plants either selectively down-regulate certain fungal taxa in response to high P, or that high P-levels directly inhibit some AM fungal taxa whereas others are indifferent to elevated P levels. This interpretation is supported by another fertilization experiment that shows the relative abundance of *Diversispora* spp. increases while that of *Glomus* spp. decrease with N and P fertilization (Liu et al. 2015). It should be noted that in our study the observed changes in the fungal community composition could only roughly explain changes in AM fungal hyphal density in soil because the method used to screen for the AM fungal taxa abundances misses many taxa for which the quantitative molecular markers are not yet available. This was the case of the control treatment that showed a complete absence of both *Diversispora* and *Claroideoglomus*, although other AM fungi were obviously present in great abundance (Fig. S2). Our measured abundances of certain AM taxa, although correct, do not signal lower activity of AM fungi in general, but only of the particular focal species (Jansa et al., 2014; Ohsowski et al., 2014). Finally, reduced mycorrhization in the CaNP treatment could have arisen because of selection of a plant community with reduced mycorrhizal dependency over the course of the experiment. The shift from a diverse herb-rich community to a less diverse grass-dominated community (Fig. 1) can also be expected to generate many changes that could generate strong feedbacks on soil C dynamics other than mycorrhizal abundance.

4.2. Degradability of litter inputs

Litter quality is likely to vary across the different fertilizer treatments and this may have a strong influence on decomposition. Previous research at the Rengen Grassland Experiment has shown that P concentration in harvested forage from the CaNP plots was about twice as high as in the other plots, with most of the other parameters remaining the same (Schellberg et al., 1999; Hejerman et al., 2010). Concentration of N in the harvested forage from the CaN plots was higher than in the CaNP plots, but only slightly and insignificantly elevated as compared to control and the limed-only plots (Hejerman et al., 2010). This resulted in much lower N/P ratio of the harvested forage in the CaNP treatment (4.8) than in the other three treatments (where the mean ranged between 13.6 and 17.7, Hejerman et al., 2010). The P-rich biomass in the CaNP treatment is likely to be degraded faster than if P in the biomass is low (Cornelissen & Thompson, 1997;

Freschet *et al.*, 2013), resulting in more C to be removed from the P-rich organic inputs than if the inputs are of low quality.

In addition to the direct influence of chemical fertilizers on forage quality, the indirect effect through altered plant community composition is also likely to be important. We observed the highest soil C storage in the treatments with the highest plant diversity, the unfertilized control and the limed plots (Fig. 1), in agreement with other observations from the Jena and Wageningen biodiversity experiments, where plant diversity correlated positively with belowground C storage (Steinbeiss *et al.*, 2008; Cong *et al.*, 2014). In our case this correlation could be explained either through production of more degradation-recalcitrant litter by the plants adapted to nutrient-limited environments (Pastor & Cohen, 1997), or through changes in the composition and/or activity of soil microbial communities (including the AM fungi) in response to the differential management. Indeed, it has recently been demonstrated that manipulation of the soil microbial diversity could have dramatic consequences on multiple ecosystem functions including litter decomposition, nutrient cycling and C sequestration (Wagg *et al.*, 2014).

4.3. Stoichiometric control of soil microbial activity

Decomposer microorganisms require a certain balance of essential nutrients and the most limiting nutrient may control the decomposition process (Sinsabaugh *et al.*, 2013). The processes of mineralization and immobilization are well known to be sensitive to the C/N ratio of the substrate and surrounding environment; however, C/N/P ratios can also be very important (Elser *et al.*, 2000). Many grassland systems, particularly those on highly weathered or carbonate rich soils are P limited and N enrichment of these systems may exacerbate P limitation and immobilize organic substrates. For instance, in a laboratory based decomposition study, Craine *et al.* (2007) found that increasing mineral N levels reduces litter decomposition, potentially leading to higher soil C accumulation. Also, van Groenigen *et al.* (2006) found a strong positive effect of N availability on build-up of the soil C reservoir. Looking at actual C stocks in grasslands, Fornara & Tilman (2012) and Fornara *et al.*, (2013) report significantly higher C under only-N fertilization in studies lasting for 27 and 19 years, respectively. In the latter study, simultaneous addition of N and P entirely negated this effect. These observations were more recently confirmed by both laboratory and field studies, indicating that both C mineralization and

gaseous N emissions are promoted by P additions to soil (He & Dijkstra, 2015; Fisk *et al.*, 2015). This broadly agrees with our results: when P limitation is removed, soil C storage decreases. However, in our work the N-only addition did not significantly increase soil C compared to the controls, even after 70 years of fertilization (Fig. 2f). Yet it tended (not significantly) to decrease standing root biomass and showed a tendency to increase the abundance of AM fungal hyphae in soil, indicating the possible interconnections between AM fungal biomass/activity and the buildup of soil C stocks.

Concluding remarks

The Rengen Grassland Experiment provides an unprecedented opportunity to examine how long-term management influences plant and microbial communities and ecosystem processes such as C sequestration. Our study provides correlative evidence that AM symbioses contribute to soil C stocks and that this C pool may be diminished if AM fungal biomass is reduced in response to P fertilization. Future experimental work is needed to more thoroughly understand the mechanisms causing these correlations. Isotopes can be used to measure C pools and track C fluxes through plants, mycorrhizas and decomposer food-webs. Also, stoichiometric models should be tested to elucidate the relative importance of C/N/P ratios in soil C dynamics. Although we acknowledge a number of limitations of our study, such as lack of records on C stocks in deeper soil horizons and limited data on the total AM fungal biomass and community composition, our results convey a very clear message - that designing future agroecosystems to simultaneously maximize aboveground yields and maximize belowground C storage may be a difficult or even an impossible goal to achieve.

Acknowledgments

LS was supported by a grant IGA 20134275 and 20144219 from the Czech University of Life Sciences in Prague. JJ was supported by the Fellowship J.E. Purkyně, Czech Ministry of Education, Youth and Sports (LK11224), and the long-term development program RVO 61388971. EV was funded by NWO and Freie Universität Berlin. ETK was funded by the ERC starting grant and NWO (Vidi and Meervoud grants). MH was supported by project GAČR 505/12/1390. NCJ was funded through a research fellowship from the Fulbright Commission of the Czech Republic, the United States National Science Foundation (DEB-0842327), and the United States Department of Agriculture (NIFA 2011-67009-30002). The maintenance of the Rengen Grassland Experiment was partly funded by the Faculty of Agriculture of the University of Bonn, Germany. JS also acknowledges funding from the German Research Foundation (DFG), project number SCHE 549/2-1. Constructive comments by two anonymous reviewers are gratefully acknowledged.

References

- Aggelides, S.M., Londra, P.A., 2000. Effects of compost produced from town wastes and sewage sludge on the physical properties of a loamy and a clay soil. *Bioresource Technology* 71, 253 – 259.
- Allen, M.F., 1991. *The ecology of mycorrhizae*. Cambridge University Press, New York. USA.
- Averill, C., Turner, B.L, Finzi, A.C., 2014. Mycorrhiza-mediated competition between plants and decomposers drives soil carbon storage. *Nature* 505, 543-545.
- Balasoorya, W.K., Denef, K., Huygens, D., Boeckx, P., 2014. Translocation and turnover of rhizodeposit carbon within soil microbial communities of an extensive grassland ecosystem. *Plant Soil* 376, 61-73.
- Brouwer, R., 1983. Functional equilibrium: sense or non-sense? *Netherlands Journal of Agriculture Science* 31, 335–348.
- Celik, I., Ortas, I., Kilic, S., 2004. Effects of compost, mycorrhiza, manure and fertilizer on some physical properties of a Chromoxerert soil. *Soil & Tillage Research* 78, 59 – 67.
- Chaudhary, V.B., Bowker, M.A., O'Dell, T.E., Grace, J.B., Redman, A.E., Rillig, M.C., Johnson, N.C., 2009. Untangling the biological contributions to soil stability in semiarid shrublands. *Ecological Applications* 19, 110–122.
- Cheng, L., Booker, F.L., Tu, C. *et al.*, 2012. Arbuscular mycorrhizal fungi increase organic carbon decomposition under elevated CO₂. *Science* 337, 1084–1087.
- Cleveland, C.C., Liptzin, D., 2007. C: N: P stoichiometry in soil: is there a “Redfield ratio” for the microbial biomass? *Biogeochemistry* 85, 235-252.
- Conant, R.T., Paustian, K., Elliott, E.T., 2001. Grassland management and conservation into grassland: effects on soil carbon. *Ecological Applications* 11, 343–355.
- Conant, R.T., Paustian, K., 2002. Potential soil carbon sequestration in overgrazed grassland ecosystems. *Global Biogeochemical Cycles* 135, 90-114.
- Cong, W.F., van Ruijven, J., Mommer, L., De Deyn, G.B., Berendse, F., Hoffland, E., 2014. Plant species richness promotes soil carbon and nitrogen stock in grassland without legumes. *Journal of Ecology* 102, 1163 – 1170.
- Cornelissen, J.H.C., Thompson, K., 1997. Functional leaf attributes predict litter decomposition rate in herbaceous plants. *New Phytologist* 135, 109-114.

- Cotrufo, M.F., Wallenstein, M.D., Boot, C.M., Denef, K., Paul, E., 2013. The Microbial Efficiency-Matrix Stabilization (MEMS) framework integrates plant litter decomposition with soil organic matter stabilization: do labile plant inputs form stable soil organic matter? *Global Change Biology* 19, 988-995.
- Couillerot, O., Ramírez-Trujillo, A., Walker, V., *et al.*, 2012. Comparison of prominent *Azospirillum* strains in *Azospirillum-Pseudomonas-Glomus* consortia for promotion of maize growth. *Applied Microbiology and Biotechnology* 97, 4639–4649.
- Craine, J.M., Morrow, C., Fierer, N., 2007. Microbial nitrogen limitation increases decomposition. *Ecology* 88, 2105–2113.
- Daynes, C.N., Field, D.J., Saleeba, J.A., Cole, M.A., McGee, P.A., 2013. Development and stabilisation of soil structure via interactions between organic matter, arbuscular mycorrhizal fungi and plant roots. *Soil Biology and Biochemistry* 57, 683-694.
- Demaria, P., Flisch, R., Frossard, E., Sinaj, S., 2005. Exchangeability of phosphate extracted by four chemical methods. *Journal of Plant Nutrition and Soil Science* 168, 89 – 93.
- Douds, D.D., Schenck, N.C., 1990. Increased sporulation of vesicular-arbuscular mycorrhizal fungi by manipulation of nutrient regimens. *Applied and Environmental Microbiology* 56, 413–418.
- Elser, J.J., Fagan, W.F., Denno, R.F., *et al.* 2000. Nutritional constraints in terrestrial and freshwater food webs. *Nature* 408, 578-580.
- Egerton-Warburton, L.M., Graham, R.C., Allen, E.B., Allen, M.F., 2001. Reconstruction of the historical changes in mycorrhizal fungal communities under anthropogenic nitrogen deposition. *Proceeding of the Royal Society B-Biological Sciences* 268, 2479-2484.
- Egerton-Warburton, L.M., Johnson, N.C., Allen, E.B., 2007. Mycorrhizal community dynamics following nitrogen fertilization: a cross-site test in five grasslands. *Ecological Monographs* 77, 527–544.
- Fisk, M., Santangelo, S., Minick, K., 2015. Carbon mineralization is promoted by phosphorus and reduced by nitrogen addition in the organic horizon of northern hardwood forests. *Soil Biology & Biochemistry* 81, 212-218.
- Fitter, A.H., Helgason, T., Hodge, A., 2011. Nutritional exchanges in the arbuscular mycorrhizal symbiosis: Implications for sustainable agriculture. *Fungal Biology Reviews* 25, 68-72.

630 Fornara, D.A., Banin, L., Crawley, M.J., 2013. Multi-nutrient vs. nitrogen-only effects on carbon
631 sequestration in grassland soils. *Global Change Biology* 19, 3848–3857.

632 Fornara, D.A., Tilman, D., 2012. Soil carbon sequestration in prairie grasslands increased by
633 chronic nitrogen addition. *Ecology* 93, 2030–2036.

634 Freschet, G.T., Cornwell, W.K., Wardle, D.A., *et al.* 2013. Linking litter decomposition of
635 above-and below-ground organs to plant–soil feedbacks worldwide. *Journal of Ecology* 101,
636 943-952.

637 Gianinazzi, S., 2010. Agroecology: the key role of arbuscular mycorrhizas in ecosystem services.
638 *Mycorrhiza* 20, 519–530.

639 Gryndler, M., Jansa, J., Hršelová, H., Chvátalová, I., Vosátka, M., 2003. Chitin stimulates
640 development and sporulation of arbuscular mycorrhizal fungi. *Applied Soil Ecology* 22, 283
641 – 287.

642 He, M., Dijkstra, F.A., 2015. Phosphorus addition enhances loss of nitrogen in a phosphorus-
643 poor soil. *Soil Biology & Biochemistry* 82, 99-106.

644 Hejerman, M., Klauisová, M., Schellberg, J., Honsová, D., 2007. The Rengen Grassland
645 Experiment: Plant species composition after 64 years of fertilizer application. *Agriculture,*
646 *Ecosystems and Environment* 122, 259–266.

647 Hati, K.M., Swarup, A., Dwivedi, A.K., Misra, A.K., Bandyopadhyay, K.K., 2007. Changes in
648 soil physical properties and organic carbon status at the topsoil horizon of a vertisol of
649 central India after 28 years of continuous cropping, fertilization and manuring. *Agriculture,*
650 *Ecosystems and Environment* 119, 127 – 134.

651 Hejerman, M., Szaková, J., Schellber, J., Tlustoš, P., 2010. The Rengen Grassland Experiment:
652 relationship between soil and biomass chemical properties, amount of elements applied, and
653 their uptake. *Plant Soil* 333, 163 – 179.

654 Hermans, C., Hammond, J.P., White, P.J., Verbruggen, N., 2006. How do plants respond to
655 nutrient shortage by biomass allocation? *Trends in Plant Science* 11, 610-617.

656 Hodge, A., Campbell, C.D., Fitter, A.H., 2001. An arbuscular mycorrhizal fungus accelerates
657 decomposition and acquires nitrogen directly from organic material. *Nature* 413, 297–299.

658 Janoušková, M., Püschel, D., Hujšlová, M., Slavíková, R., Jansa, J., 2015. Quantification of
 659 arbuscular mycorrhizal fungal DNA in roots: how important is material preservation?
 660 Mycorrhiza, in press, doi 10.1007/s00572-014-0602-7.

661 Jansa, J., Mozafar, A., Frossard, E., 2003. Long-distance transport of P and Zn through the
 662 hyphae of an arbuscular mycorrhizal fungus in symbioses with maize. *Agronomie* 23, 481–
 663 588.

664 Jansa, J., Wiemken, A., Frossard, E., 2006. The effects of agricultural practices on arbuscular
 665 mycorrhizal fungi. In: E Frossard, WEH Blum, BP Warkentin (eds.) *Function of Soils for*
 666 *Human Societies and the Environment*. Geological Society, London, Special Publications
 667 266, 89–115.

668 Jansa, J., Erb, A., Oberholzer, H.R., Šmilauer, P., Egli, S., 2014. Soil and geography are more
 669 important determinants of indigenous arbuscular mycorrhizal communities than
 670 management practices in Swiss agricultural soils. *Molecular Ecology* 23, 2118–2135.

671 Johnson, D., Leake, J.R., Ostle, N., Ineson, P., Read, D.J., 2002. In situ (CO₂)-C-13 pulse-
 672 labelling of upland grassland demonstrates a rapid pathway of carbon flux from arbuscular
 673 mycorrhizal mycelia to the soil. *New Phytologist* 153, 327–334.

674 Johnson, D., Leake, J.R., Read, D.J., 2005. Liming and nitrogen fertilization affects phosphatase
 675 activities, microbial biomass and mycorrhizal colonization in upland grassland. *Plant and*
 676 *Soil* 271, 15–164.

677 Johnson, D., Leake, J.R., Read, D.J., 2006. Role of arbuscular mycorrhizal fungi in carbon and
 678 nutrient cycling in grassland. In: Gadd GM (ed.) *Fungi in Biogeochemical Cycles*, PP. 129–
 679 150, Cambridge University Press.

680 Johnson, N.C., 1993. Can fertilization of soil select less mutualistic mycorrhizae? *Ecological*
 681 *Applications* 3, 749–757.

682 Johnson, N.C., 2010. Tansley Review: Resource stoichiometry elucidates the structure and
 683 function of arbuscular mycorrhizas across scales. *New Phytologist* 185, 631–647.

684 Johnson, N.C., Rowland, D.L., Corkidi, L., Egerton-Warburton, L.M., Allen, E.B., 2003.
 685 Nitrogen enrichment alters mycorrhizal allocation at five mesic to semiarid grasslands.
 686 *Ecology* 7, 1895–1908.

687 Johnson, N.C., Angelard, C., Sanders, I.R., Kiers, E.T., 2013. Predicting community and
688 ecosystem outcomes of mycorrhizal responses to global change. *Ecology Letters* 16, 140–
689 153.

690 Johnston, A.E., Goulding, K.W.T., Poulton, P.R. (1986) Soil acidification during more than 100
691 years under permanent grassland and woodland at Rothamsted. *Soil Use and Management* 2,
692 3-10.

693 Joner, E.J., Jakobsen, I., 1995. Growth and extracellular phosphatase activity of arbuscular
694 mycorrhizal hyphae as influenced by soil organic matter. *Soil Biology and Biochemistry* 27,
695 1153 – 1159.

696 Kaiser, C., Franklin, O., Dieckmann, U., Richter, A., 2014. Microbial community dynamics
697 alleviate stoichiometric constraints during litter decay. *Ecology Letters* 17, 680-690.

698 Kennedy, N., Connolly, J., Clipson, N., 2005. Impact of lime, nitrogen and plant species on
699 fungal community structure in grassland microcosms. *Environmental Microbiology* 6, 780-
700 788.

701 Kiers, E.T., Duhamel, M., Beesetty, Y., *et al.* 2011. Reciprocal Rewards Stabilize Cooperation in
702 the Mycorrhizal Symbiosis. *Science* 333, 880-882.

703 Kubát, K., Hrouda, L., Chrtěk, J., Kaplan, Z., Kirschner, J., Štěpánek, J., Zázvorka, J., 2002. Klíč
704 ke květeně České republiky. Academia, Praha.

705 Lal, R., 2007. Anthropogenic influences on world soils and implications to global food security.
706 *Advances in Agronomy* 93, 69-93.

707 Lal, R., 2008. Carbon sequestration. *Philosophical Transactions of the Royal Society B:*
708 *Biological Sciences* 363, 815-830.

709 Lal, R., 2013. Intensive agriculture and the soil carbon pool. *Journal of Crop Improvement* 27,
710 735-751.

711 Lendenmann, M., Thonar, C., Barnard, R.L., Salmon, Y., Werner, R.A., Frossard, E., Jansa, J.,
712 2011. Symbiont identity matters: carbon and phosphorus fluxes between *Medicago*
713 *truncatula* and different arbuscular mycorrhizal fungi. *Mycorrhiza* 21, 689-702.

714 Leifheit, E.F., Verbruggen, E., Rillig, M.C., 2015. Arbuscular mycorrhizal fungi reduce
715 decomposition of woody plant litter while increasing soil aggregation. *Soil Biology &*
716 *Biochemistry* 81, 323-328.

717 Liu, Y., Shi, G., Mao, L. *et al.*, 2012. Direct and indirect influence of 8 yr of nitrogen and
 718 phosphorus fertilization on Glomeromycota in an alpine meadow ecosystem. *New*
 719 *Phytologist* 194, 523–535.

720 Liu, Y., Johnson, N.C., Mao, L. *et al.* 2015. Phylogenetic structure of arbuscular mycorrhizal
 721 community shifts in response to increasing soil fertility. *Soil Biology & Biochemistry* 89,
 722 196-205.

723 McGonigle, T.P., Miller, M.H., Evans, D.G., Fairchild, G.L., Swan, J.A., 1990. A new method
 724 which gives an objective measure of colonization of roots by vesicular-arbuscular
 725 mycorrhizal fungi. *New Phytologist* 115, 495–501.

726 Mosse, B., 1959. Observations on the extra-matrical mycelium of a vesicular arbuscular
 727 endophyte. *Transactions British Mycological Society* 4, 439 – 448.

728 Murphy, C.J., Baggs, E.M., Morley, N., Wall, D.P., Paterson, E., 2015. Rhizosphere priming can
 729 promote mobilisation of N-rich compounds from soil organic matter. *Soil Biology &*
 730 *Biochemistry* 81, 236-243.

731 Newsham, K., Fitter, A., Watkinson, A.R., 1995. Multifunctionality and biodiversity in
 732 arbuscular mycorrhizas. *Trends in Ecology and Evolution* 10, 407–411.

733 Nielsen, K.L., Eshel, A., Lynch, J.P., 2001. The effect of phosphorus availability on the carbon
 734 economy of contrasting common bean (*Phaseolus vulgaris* L.) genotypes. *Journal of*
 735 *Experimental Botany* 52, 329-339.

736 Ohsowski, B.M., Zaitsoff, P.D., Öpik, M., Hart, M.M., 2014. Where the wild things are: looking
 737 for uncultured Glomeromycota. *New Phytologist* 204, 171–179.

738 Olsson, P.A., Thingstrup, I., Jakobsen, I., Baath, E., 1999. Estimation of the biomass of
 739 arbuscular mycorrhizal fungi in a linseed field. *Soil Biology and Biochemistry* 31, 1879–
 740 1887.

741 Pastor, J., Cohen, Y., 1997. Herbivores, the functional diversity of plants species, and the cycling
 742 of nutrients in ecosystems. *Theoretical Population Biology* 51, 165-179.

743 Phillips, R.P., Brzostek, E., Midgley, M.G., 2013. The mycorrhizal-associated nutrient
 744 economy: a new framework for predicting carbon-nutrient couplings in temperate forests.
 745 *New Phytologist* 199, 41-51.

746 Poeplau, C., Don, A., Vesterdal, L., Leifeld, J., van Wesemael, B., Schumacher, J., Gensior, A.,
 747 2011. Temporal dynamics of soil organic carbon after land-use change in the temperate zone
 748 – carbon response functions as a model approach. *Global Change Biology* 17, 2415–2427.
 749 Rillig, M.C., 2004. Arbuscular mycorrhizae and terrestrial ecosystem processes. *Ecology Letters*
 750 7, 740–754.
 751 Rillig, M.C., Field, C.B., Allen, M.F., 1999. Soil biota responses to long-term atmospheric CO₂
 752 enrichment in two California annual grasslands. *Oecologia* 119, 572–577.
 753 Rothmaler, W., Jager, E.J., Werner, K., 2000. Exkursionsflora von Deutschland, Bd. 3,
 754 Gefäßpflanzen Atlasband, Heidelberg, Spektrum Akademischer Verlag.
 755 Sala, O.E., Parton, W.J., Joyce, L.A., Lauenroth, W.K., 1988. Primary production of the central
 756 grassland region of the United States. *Ecology* 69, 40–45.
 757 Schellberg, J., Mösel, B.M., Kühbauch, W., Rademacher, I.F., 1999. Long-term effects of
 758 fertilizer on soil nutrient concentrations, yield, forage quality and floristic composition of a
 759 hay meadow in the Eifel mountains, Germany. *Grass and Forage Science* 54, 195–207.
 760 Schulten, H.R., Schnitzer, M., 1997. Chemical model structures for soil organic matter and
 761 soils. *Soil Science* 162, 115–130.
 762 Scott, B.J., Ridley, A.M., Conyers, M.K. (2000) Management of soil acidity in long-term
 763 pastures of south-eastern Australia: a review. *Australian Journal of Experimental*
 764 *Agriculture* 40, 1173–1198.
 765 Semelová, V., Hejcman, M., Palvů, V., Vacek, S., Podrázký, V., 2008. The Grass Garden in the
 766 Giant Mts. (Czech Republic): Residual effect of long-term fertilization after 62 years.
 767 *Agriculture, Ecosystems and Environment* 123, 337–342.
 768 Sinsabaugh, R.L., Manzoni, S., Moorhead, D.L., Richter, A., 2013. Carbon use efficiency of
 769 microbial communities: stoichiometry, methodology and modelling. *Ecology Letters* 16,
 770 930–939.
 771 Silvertown, J., Poulton, P., Johnston, E., Edwards, G., Heard, M., Biss, P.M., 2006. The Park
 772 Grass Experiment 1856–2006: its contribution to ecology. *Journal of Ecology* 94, 801–814.
 773 Smith, S.F., Facelli, E., Pope, S., Smith, F.A., 2010. Plant performance in stressful environments:
 774 interpreting new and established knowledge of the roles of arbuscular mycorrhizas. *Plant*
 775 *Soil* 326, 3–20.

776 Staddon, P.L., Ramsey, C.B., Ostle, N., Ineson, P., Fitter, A.H., 2003. Rapid turnover of hyphae
777 of mycorrhizal fungi determined by AMS microanalysis of C-14. *Science* 300, 1138 –
778 1140.

779 StatSoft, 1995. Statistica for Windows. StatSoft. Tulsa. OK. USA.

780 Steinbeiss, S., Bessler, H., Engels, C. *et al.*, 2008. Plant diversity positively affects short-term
781 soil carbon storage in experimental grasslands. *Global Change Biology* 14, 2937-2949.

782 ter Braak, C.J.F, Šmilauer, P., 2002. CANOCO Reference Manual and CanoDraw for Windows
783 User's Guide: Software for Canonical Community Ordination (Version 4.5). Microcomputer
784 Power. Ithaca. NY. USA.

785 Thonar, C., Erb, A., Jansa, J., 2012. Real-time PCR to quantify composition of arbuscular
786 mycorrhizal fungal communities – marker design, verification, calibration and field
787 validation. *Molecular Ecology Resources* 12, 219–232.

788 Treseder, K.K., Allen, M.F., 2000. Mycorrhizal fungi have a potential role in soil carbon storage
789 under elevated CO₂ and nitrogen deposition. *New Phytologist* 147, 189-200.

790 van Groenigen, K., Six, J., Hungate, B.A., de Graaff, M., van Breemen, N., van Kessel, C., 2006.
791 Element interactions limit soil carbon storage. *Proceedings of the National Academy of*
792 *Sciences of the United States of America* 103, 6571 – 6574.

793 Verbruggen, E., Veresoglou, S.D., Anderson, I.C., Caruso, T., Hammer, E.C., Kohler, J., Rillig,
794 M.C., 2013. Arbuscular mycorrhizal fungi – short-term liability but long-term benefits for
795 soil carbon storage? *New Phytologist* 197, 366–368.

796 Vierheilig, H., Coughlan, A.P., Wyss, U., Piché, Y., 1998. Ink and vinegar, a simple staining
797 technique for arbuscular-mycorrhizal fungi. *Applied and Environmental Microbiology* 64,
798 5004–5007.

799 Wagg, C., Jansa, J., Stadler, M., Schmid, B., van der Heijden, M.G.A., 2011. Mycorrhizal fungal
800 identity and diversity relaxes plant-plant competition. *Ecology* 92, 1303–1313.

801 Wagg, C., Bender, S.F., Widmer, F., van der Heijden, M.G.A., 2014. Soil biodiversity and soil
802 community composition determine ecosystem multifunctionality. *Proceedings of the*
803 *National Academy of Sciences of the United States of America* 111, 5266-5270.

804 Wander, M.M., Traina, S.J., Stinner, B.R., Peters, S.E., 1994. Organic and conventional
805 management effects on biologically active soil organic matter pools. *Soil Science Society of*
806 *America Journal* 58, 1130-1139.

807 Welc, M., Ravnskov, S., Kieliszewska-Rokicka, B., Larsen, J., 2010. Suppression of other soil
808 microorganisms by mycelium of arbuscular mycorrhizal fungi in root-free soil. *Soil Biology*
809 *and Biochemistry* 42, 1534–1540.

810 Wendland, M., Diepolder, M., Capriel, P., 2012. Leitfaden für die Düngung von Acker- und
811 Grünland. Institut für Agrarökologie. Ökologischen Landbau und Bodenschutz. Freising-
812 Weihenstephan.

813 Wilson, G.W.T., Rice, C.W., Rillig, M.C., Springer, A., Hartnett, D.C., 2009. Soil aggregation
814 and carbon sequestration are tightly correlated with the abundance of arbuscular mycorrhizal
815 fungi: results form long-term field experiments. *Ecology Letters* 12, 452–461.

816 Zar, J.H., 1999. Biostatistical analysis. Pearson Education. India.

817 Zhu, Y.G., Miller, R.M., 2003. Carbon cycling by arbuscular mycorrhizal fungi in soil-plant
818 systems. *Trends in Plant Science* 8, 407-409.

819

Figure legends:

Fig. 1. Annual hay yield (a), root standing biomass (b), root-to-shoot biomass ratio (c), plant species richness (d), cover of herbs (e), and cover of grasses (f) as affected by the management treatments (Control – no amendments, Ca – lime added, CaN – lime and nitrogen added, CaNP – limed and nitrogen and phosphorus fertilizers added). Bars represent means +1 standard errors (n = 10 for a-c and n = 5 for d-f). Different letters indicate significant differences between treatment means (least significant difference F-test, $p < 0.05$). Biomass of roots has been estimated only for the topsoil (0-10 cm depth) in October 2011, and upscaled to 1 ha surface to compare with hay yields. Average yields of seasons 2008 – 2012 are shown.

Fig. 2. Extent of root length colonized by any of the arbuscular mycorrhizal (AM) fungal structures (a), and the fraction of root length colonized specifically by arbuscules (b), vesicles (c), and AM fungal coils (d). The density of the extraradical mycelium (ERM) of AM fungi in the soil (e) and the concentration of organic carbon (C) in the topsoil (f) as affected by the management treatments. Bars represent means +1 standard errors (n=10). Different letters indicate significant differences between treatment means (least significant difference F-test, $p < 0.05$). The percentage values were arcsin-square root transformed for the statistical comparisons so as not to violate ANOVA assumptions. Soil C stocks have been estimated only for the topsoil (0-10 cm depth) and upscaled to 1 ha surface.

Fig. 3. Ordination diagram showing results of redundancy analysis of abundance of individual arbuscular mycorrhizal fungal taxa in soil in relation to soil chemical properties and hay yield. *Rhizophagus irregularis* abundance was assessed with quantitative PCR marker targeting mitochondrial large ribosomal subunit (LSU) gene whereas the other AM fungal taxa were assessed with markers for nuclear LSU genes. Environmental variables: Soil N and C – total nitrogen and organic carbon concentrations in the soil, respectively. Soil P and Ca – available phosphorus and mobile calcium concentrations in the soil, respectively. Yield – hay productivity per year (average of 2008-2012 seasons). First and second axes explain 62.8% and 25.1% of the mycorrhizal fungal community and environmental data relationship, respectively.

850 **Fig. 4.** Negative relationship between volumetric soil carbon (C) concentration and hay yields (a)
851 and a positive relationship between the volumetric soil C concentration and the length density of
852 extraradical mycelium (ERM, log transformed) of the arbuscular mycorrhizal fungi in the soil
853 (b). Every dot represents a separate experimental plot. Both regression lines $p < 0.01$.
854

Table 1. Soil chemical properties (0 – 10 cm depth) of the Rengen Grassland Experiment. Soil data measured on samples collected in 2011 are shown. Mean values (n=10) are shown of soil pH, available soil phosphorus (P), total soil nitrogen (N) and soil mobile calcium (Ca) concentrations. Different letters within a column indicate significant differences between treatment means according to the least significant difference post-hoc test following significant ANOVA (*F* - and *p* – values are shown in italics).

Treatment	pH	P (mg kg ⁻¹)	N (g kg ⁻¹)	Ca (mg kg ⁻¹)
control	5.32c	0.307b	3.79bc	70.5
Ca	6.96a	0.663b	4.11a	138.3
CaN	7.06ab	0.452b	4.03ab	121.9
CaNP	7.14b	9.947a	3.57c	103.7
<i>F value</i>	<i>211.34</i>	<i>92.98</i>	<i>6.26</i>	<i>2.40</i>
<i>p-value</i>	<i><0.001</i>	<i><0.001</i>	<i>0.002</i>	<i>0.084</i>

Table 2. Gravimetric and volumetric carbon (C) concentrations of the soils, soil bulk density, and the soil C/N ratios as affected by the different management treatments. Mean values (n=10, $\pm$ standard errors) are shown. Different letters within a column indicate significant differences between treatment means according to the least significant difference post-hoc test following significant ANOVA (F - and p – values are shown in italics).

Treatment	Gravimetric C concentration (g kg ⁻¹)	Soil bulk density (kg l ⁻¹)	Volumetric C concentration (g l ⁻¹)	Soil C/N ratio
Control	51.7 ($\pm$ 0.6) ab	0.97 ($\pm$ 0.02) b	50.3 ($\pm$ 0.9) b	13.6 ($\pm$ 0.09) a
Ca	55.5 ($\pm$ 1.9) a	1.00 ($\pm$ 0.03) ab	55.4 ($\pm$ 1.7) a	13.5 ($\pm$ 0.20) a
CaN	49.0 ($\pm$ 1.9) b	1.03 ($\pm$ 0.02) ab	50.5 ($\pm$ 1.8) ab	12.1 ($\pm$ 0.14) b
CaNP	41.5 ($\pm$ 1.9) c	1.06 ($\pm$ 0.02) a	44.1 ($\pm$ 2.3) c	11.6 ($\pm$ 0.23) c
<i>F value</i>	<i>12.7</i>	<i>2.96</i>	<i>6.84</i>	<i>33.97</i>
<i>p-value</i>	<i><0.001</i>	<i>0.045</i>	<i><0.001</i>	<i><0.001</i>

Figure 2

Figure 3

Figure 4