The surface-atmosphere exchange of carbon dioxide, water, and sensible heat across a dryland wheat-fallow rotation

Elizabeth S.K. Vick^a, Paul C. Stoy^{a*}, Angela C.-I. Tang^a, Tobias Gerken^{a,b}

^a Department of Land Resources and Environmental Sciences, Montana State University, Bozeman, MT 59717, USA

^b Department of Meteorology, The Pennsylvania State University, University Park, PA 16802, USA

* Corresponding author

Phone: +1 406 600 3577

Fax: +1 406 994 3933

Email: paul.stoy@montana.edu

Abstract

Summer fallow – the practice of keeping a field out of production during the growing season – is a common practice in dryland wheat (*Triticum aestivum* L.) cropping systems to conserve soil water resources. Fallow also depletes soil carbon stocks and thereby soil quality. The area of fallow has decreased by tens of millions of hectares since the 1970s in the northern North American Great Plains as producers have recognized that avoiding fallow usually confers both an economic and soil conservation benefit. Observed summertime cooling across parts of this region has coincided with fallow reduction, suggesting that the role of fallow in atmospheric processes needs to be ascertained. We measured carbon dioxide, latent heat, and sensible heat flux across a winter wheat – spring wheat – fallow sequence in Montana, USA to determine the effects of dryland crop management on ecosystem carbon resources and energy partitioning at the surface-atmosphere interface. Winter wheat and spring wheat fields were carbon sinks ($F_c = -203 \pm 52$ g

C-CO₂ m⁻² and -107 ± 29 g C-CO₂ m⁻²), respectively, during the April to September study period, but the fallow field was a carbon source of 135 ± 73 g C-CO₂ m⁻². Evapotranspiration in the wheat crops was over 100 mm greater than the 275 ± 39 mm observed in the fallow field during the study period. Modeled maximum daily atmospheric boundary layer height was on average 210 m higher and up to 900 m higher in fallow compared to the spring wheat field with more crossings of the modeled atmospheric boundary layer and lifted condensation level, suggesting that regional studies of the effects of fallow on near-surface temperature and moisture are necessary to understand the effects of fallow reduction on regional climate dynamics. Results demonstrate that fallow has a detrimental impact to soil carbon resources yet is less water intensive, with consequences for regional climate via its impacts on atmospheric boundary layer development and global climate via its carbon metabolism.

Keywords

Atmospheric boundary layer; Carbon dioxide flux; Eddy covariance; Evapotranspiration; Fallow; Sensible heat flux; Wheat

1. Introduction

Wheat (*Triticum aestivum* L.) provides more than 20% of the calories and protein for the global population (Hawkesford et al., 2013) and nearly 220 Mha of wheat was harvested globally in 2013 (FAO, 2013). Wheat therefore plays a central role in not only global food production, but also in the global exchange of water, energy, and climate-relevant trace gases like carbon dioxide between the land surface and the atmosphere (West and Marland, 2002; Buyanovsky and Wagner, 1998).

Our understanding of the interaction between wheat cropping systems and the atmosphere remains incomplete. Wheat has a very low canopy resistance to water vapor transport during its

main growth period (Bonan, 2008), and models tend to accurately simulate latent heat flux (LE , see Table 1 for a list of abbreviations), and sensible heat flux (H) during these periods (Ingwersen et al., 2011). Aspects of the seasonal timing of crop development including ripening (Ingwersen et al., 2011) and management decisions like harvesting (Sus et al., 2010) on surface-atmosphere fluxes continue to challenge ecosystem models. Management practices including crop rotations have been identified as important contributors to carbon metabolism at the field scale (Béziat et al., 2009; Schmidt et al., 2012), but have largely been studied in winter wheat crops and/or mesic cropping systems to date (e.g. Billesbach et al., 2014; Moureaux et al., 2008). Surface-atmosphere exchange in rotations common to dryland cropping systems including spring wheat and chemical fallow (hereafter ‘fallow’) have not been measured.

Fallow is a common management practice in the dryland wheat-growing regions of the northern North American Great Plains to conserve water for subsequent crops (Lubowski et al., 2006). Fallow however also increases erosion (Wischmeier, 1959) and soil carbon loss (Cihacek and Ulmer, 1995), and fallow-small grain management strategies are not considered sustainable from the soil conservation perspective (Merrill et al., 1999). Management practices are changing. The area of fallow in the Prairie Provinces of Canada has decreased from over 15 Mha in the 1970s to under 2 Mha at the present (Figure 1) as producers have realized that the water-savings benefit of fallow is outweighed by the economic losses of not planting (Dhuyvetter et al., 1996). The area under fallow in the United States has likewise decreased from 16 Mha to 6 Mha across the same time frame (Lubowski et al., 2006), largely in the northern Great Plains and other areas of the semiarid West (Figure 1). Despite the decreasing trend in fallow area across the North American northern Great Plains, fallow remains common in many regions including major land resource area (MLRA) 52 in north-central Montana – the largest wheat-growing region in the state – where some 40% of agricultural lands may remain in fallow in any given year. In contrast, fallow has been reduced in northeastern Montana (MLRA 53) by hundreds of kha over the past

decade (Long et al., 2014, 2013) as producers have adopted continuous cropping or alternate cropping practices (Burgess et al., 2012; Miller et al., 2003, 2002).

The widespread decline of fallow in agricultural areas of the Canadian Prairie Provinces (Figure 1) has coincided with a summertime cooling trend since the 1970s (Betts et al., 2013a, 2013b; Gameda et al., 2007; Mahmood et al., 2014). Extreme temperature events now occur less frequently than in the recent past, maximum summer temperatures have decreased by *ca.* 2 °C, relative humidity has increased by some 7% (Betts et al., 2013b), and summer precipitation has increased by an average of 10 mm/decade across parts of the Canadian Prairie Provinces (Gameda et al., 2007). A remarkable 6 W m⁻² summer cooling has been observed (Betts et al., 2013a); for reference, anthropogenic greenhouse gasses are responsible for a *ca.* 2.5 W m⁻² warming globally since the dawn of the Industrial Era (IPCC, 2007). These climate benefits have only occurred during the growing season; fall, winter, and spring temperatures have followed global trends (Betts et al., 2013b).

Studies suggest that these climate changes are the result of land management, specifically the trend away from leaving fields fallow during summer (Betts et al., 2013a; Gameda et al., 2007; Mahmood et al., 2014). These large-scale declines in fallow area have resulted in less sensible heat and more water vapor entering the atmosphere and a moister, shallower atmospheric boundary layer (*ABL*) and lower lifted condensation level (*LCL*) with an increased probability of convective precipitation (Betts et al., 2013b; Gameda et al., 2007). In other words, the observed regional climate cooling is broadly consistent with the effects of fallow avoidance on climate processes.

Using a conceptual model, Gameda et al. (2007) suggested that fallow supports an *ABL* height (*h*) on the order of 2-3 km, but that *h* is only some 0.5 km over crops. These estimates have not been supported by observations or models, in part because the impacts of field-scale management changes on *H* and *LE*, which help determine *h*, have not been studied to date in

wheat-fallow rotations. Doing so is an important step in understanding the role of human management on climate processes in the North American Northern Great Plains.

Cropping systems that include wheat also influence land surface biogeochemistry. Many studies to date have used the eddy covariance technique to find that wheat tends to be a strong carbon sink at the field scale during the growing season, but is infrequently a net annual carbon sink. For example, Anthoni et al. (2004) found that a central European winter wheat field was a sink for atmospheric carbon dioxide during the growing season, but a source of carbon annually, with a net biome productivity of 45 to 105 g C m⁻² y⁻¹, indicating a loss of C to the atmosphere when the end products of the wheat crop were accounted for. Crop rotations are also important for carbon metabolism. Béziat et al. (2009) measured the carbon balance of a three-crop rotation and found that the net biome productivity was -161 +/- 66 g C m⁻² y⁻¹ in winter wheat, but a maize crop at the same site a year earlier was a carbon source of 372 +/- 78 g C m⁻² y⁻¹ due in part to residue from previous crops. These observations emphasize the importance of studying crop rotations for a comprehensive understanding of the impacts of agriculture on surface-atmosphere exchange.

Here, we use the eddy covariance technique to measure the surface-atmosphere exchange of carbon dioxide (F_c), LE , and H across a dryland winter wheat - spring wheat - fallow sequence in central Montana, USA. We quantify differences in surface fluxes during the April – September growing period and use models of h and the height of the lifted condensation level (h_{LCL}) driven by observed surface-atmosphere energy flux and near-surface air temperature and humidity to help us understand the physical mechanisms that underlie the observed effects of agricultural management on ABL , LCL , and regional climate processes (Gameda et al., 2007). We additionally discuss the impacts of wheat and fallow on carbon and water resources to provide information to producers for land management decisions.

2. Materials and Methods

2.1 Site Description

Eddy covariance and micrometeorological measurements were made in two fields in central Montana, USA. Measurements were made on the first field during 2013 when it was planted with winter wheat, and during 2014 when it was planted with spring wheat. Measurements in the second field were made during 2014 when it was held in fallow.

The first field was planted with winter wheat in autumn, 2012 following a year of chemical fallow. In March 2013, a 3 m tower with eddy covariance and micrometeorological instrumentation was installed in this field at 46° 59' 41.1" N, 109° 36' 49.5" W (Figure 2). Eddy covariance instrumentation was mounted at 1.8 m with a fetch of 200 m from the nearest field edge to the west. The winter wheat crop was harvested on August 7, 2013 and instruments were left on the tower until September 2013, when they were removed for calibration and reinstalled. The tower was moved to accommodate the planting of spring wheat on May 5, 2014 and it measured spring wheat during the 2014 growing season. It was removed in September 2014 following the spring wheat harvest on August 18, 2014.

A second tower of identical height and instrumentation height was installed during spring, 2014, at 46° 59' 44.8" N, 109° 37' 46.3" W in a no-till chemical fallow field that was planted with spring wheat the previous year. Chemical treatments were applied to discourage weedy growth during the study period, with the exception of the immediate vicinity of the tower to avoid damaging sensors. The fallow field was two fields to the west of the first (wheat) field, and the fallow field tower was 1.2 km from the wheat field tower (Figure 2). The minimum fetch was 200 m to the southwest. For the purposes of this study, measurements from all fields are studied for the April-September period - hereafter the 'study period' - that encompasses the main growing season.

Mean annual temperature and precipitation were 6.6 °C and 388 mm, respectively, in Moccasin, MT where the Montana State University Central Agricultural Research Center is located some 25

km from the study fields. The slope of both fields was less than 0.5 degrees on Judith and Danvers clay loams.

2.2 Measurements

2.2.1 Meteorological Measurements

Micrometeorological variables were measured at both towers as summarized in Table 2. Canopy heights and snow depths were measured using a SR50A-L sonic depth sensor (Campbell Scientific Inc., Logan, UT, USA) at the wheat tower. Incident shortwave (SW_{in}), outgoing shortwave, incident longwave, outgoing longwave radiation, and thereby net radiation (R_n) were measured at both towers using NR01 four-component net radiometers (Hukseflux, Delft, The Netherlands) mounted at 1.8 m above the soil surface. Air temperature (T_a) and relative humidity were measured using HMP45C temperature and relative humidity probes (Vaisala, Vantaa, Finland) at 2 m. Soil heat flux (G) was measured using self-calibrating HFP01 heat flux plates (Hukseflux) at 5 cm below the soil surface. Soil moisture was measured at 5 cm and 10 cm using two CS616 time domain reflectometer sensors (Campbell Scientific) in the wheat field and two CS650 sensors (Campbell Scientific) in the fallow field. Soil temperature was measured at multiple depths using copper-constantan thermocouples in the wheat field and the CS650 sensors in the fallow field. Measurements were made every minute and half-hour averages were stored using CR3000 and CR1000 data loggers (Campbell Scientific).

2.2.2 Turbulent Flux Measurements

F_c and LE were measured using the eddy covariance technique. This involved the coupling of CSAT-3 sonic anemometers (Campbell Scientific) with LI-7200 CO₂/H₂O enclosed infrared gas analyzers (LiCor, Lincoln, NE) installed 1.8 m above the ground surface on both towers. H was measured using the CSAT-3 sonic anemometers. Data were recorded at 10 Hz on CR3000 data loggers (Campbell Scientific) and processed into half-hourly flux sums using EddyPro (LiCor). Eddy covariance processing steps involved double rotation rather than the planar fit method given the rapid changes in wheat canopy height (Kaimal and Finnegan, 1994), covariance maximization

for time lag detection (Mauder and Foken, 2004), and plausibility ranges for the of five standard deviations from the mean for vertical velocity and 3.5 standard deviations from the mean for CO₂ and H₂O concentrations. Negative values refer to fluxes from the atmosphere to the surface following the micrometeorological convention.

Surface-atmosphere fluxes are often underestimated by eddy covariance measurements, especially during periods of insufficient turbulent exchange that usually occur at night (Aubinet et al., 2000; Falge et al., 2001a; Gu et al., 2005; Papale et al., 2006; Reichstein et al., 2005). A friction velocity (u^*) filter following Reichstein et al. (2005) was applied to the data set to identify and filter periods of insufficient turbulence. F_c measurements made during night, defined as periods for which the solar zenith angle exceeded 90°, were binned into six T_a classes for three month periods and further binned into twenty u^* classes. The u^* filter was chosen to be the value at which the mean value of the F_c observations in a given u^* class first exceeded 95% of the remaining F_c observations at higher values of u^* . u^* threshold values determined using this approach exhibited trivial differences (not shown) from a statistical approach in which the u^* threshold was determined as the u^* class at which the F_c observations were not statistically different than F_c observations at higher u^* as determined by a one-sided t-test following Papale (2012). Nighttime F_c observations that occurred under insufficient u^* , and F_c observations that exceeded logical upper bounds of 20 $\mu\text{mol m}^{-2} \text{s}^{-1}$ and lower bounds of -35 $\mu\text{mol m}^{-2} \text{s}^{-1}$ determined using probability distribution functions were excluded from the data record.

2.3. Gapfilling and Data Processing

Missing data occurred due to power outages and equipment malfunction, as well as damage to wires by animals. Missing SW_{in} , T_a , relative humidity, and soil temperature data were gapfilled using linear regression from the neighboring tower if data were available. If data from the neighboring tower were also missing, data from the Moccasin Soil Climate Analysis Network (SCAN) site, located 25 km from the study fields, were used in the gapfilling procedure.

2.3.1 Gapfilling Soil Heat Flux and Net Radiation

Missing G observations were gapfilled by establishing a linear relationship with R_n on a daily basis, relating the resulting slope and intercept parameters with canopy height, and gapfilling using regression parameters that varied as a function of canopy height. If measured R_n data were not available from either tower, a linear relationship for each day between tower R_n and SW_{in} from the Moccasin SCAN site was established and used for gapfilling R_n , which in turn was used to gapfill G .

2.3.2 Gapfilling Carbon Dioxide Flux

Our gapfilling model for F_c needed to simultaneously account for photosynthetic and respiratory processes due to rapid changes in canopy biomass during the growth period and for non-vegetated conditions before plant growth and after harvest. To gapfill F_c , a rectangular hyperbolic model for ecosystem production and the respiration model of Lloyd and Taylor (1994) were combined:

$$F_c = -\frac{\alpha\beta SW_{in}}{\alpha SW_{in} + \beta} + R_{10} \exp\left(E_0 \left(\frac{1}{T_{10} - T_t} - \frac{1}{T_a - T_t}\right)\right) \quad (1)$$

where α is the initial response of the light response curve ($\mu\text{mol CO}_2 \text{ J}^{-1}$), β is gross ecosystem productivity (GEP) at light saturation ($\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$), R_{10} ($\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$) is ecosystem respiration (RE) at a T_a of 10 °C (T_{10}), E_0 is the activation energy parameter (K), and the reference temperature T_t is set to 227.13 K following Falge et al. (2001b). Parameters were fit using least squares regression for observations within a seven-day moving window of F_c , SW_{in} , and T_{air} observations. Parameter sets during periods for which the parameter estimation routine did not converge to an optimal solution or for which observations were not available were estimated using the previous day's values for the case of small gaps of two days or less, and using linear interpolation of parameter values from preceding and subsequent periods for large gaps of more than two days.

2.3.3 Gapfilling Latent Heat Flux

Evapotranspiration (ET) was calculated as LE divided by the latent heat of vaporization with units set to equal mm per half hour eddy covariance measurement period. Missing LE (and thereby ET)

observations, including those that exceeded logical bounds of -100 W m^{-2} and 600 W m^{-2} , were gapfilled using the Priestley-Taylor model (Priestley and Taylor, 1972) adjusted to include an uncertainty term, ε_{PT} , that functions as an intercept parameter:

$$ET_0 = \frac{1}{\lambda} \frac{s(R_n - G)}{s + \gamma} \alpha_{PT} + \varepsilon_{PT}. \quad (2)$$

Here, λ is the latent heat of vaporization, s is the slope of the saturation vapor pressure-temperature relationship, γ is the psychrometric constant, and α_{PT} is the Priestley-Taylor coefficient. α_{PT} and ε_{PT} were determined for the winter wheat, spring wheat, and fallow fields individually for each day of the study period using linear regression against measured ET . α_{PT} and ε_{PT} for days during which insufficient data were available were estimated using the previous day's parameters for the case of small gaps of two days or less, and linear interpolation from the parameters of preceding and subsequent periods for large gaps of more than two days.

2.3.4 Gapfilling Sensible Heat Flux

Missing H values, including those that exceeded logical bounds of -100 W m^{-2} and 500 W m^{-2} , were gapfilled by fitting a daily linear regression between observed H and R_n . The linear model included a slope and an intercept parameter similar to equation 2, and missing H values were filled using the results of this regression. As with the other flux gapfilling routines, parameters which were unable to be calculated due to missing data were estimated using the previous day's parameters for the case of small gaps of two days or less, and linear interpolation from the parameters of preceding and subsequent periods for large gaps of more than two days.

2.4 Uncertainty Analysis

Uncertainty in eddy covariance observations is often on the order of 10-15% (e.g. Goulden et al., 1997), and varies as a function of flux magnitude (Richardson et al., 2008, 2006). The total uncertainty in eddy covariance measurements is a function of observational uncertainty (Moncrieff et al., 1996), gapfilling uncertainty (Falge et al., 2001a, 2001b), and spatial

uncertainty due to the assumption that a spatially variable flux can be expressed as an average on a square meter basis (Oren et al., 2006).

We applied the approach of Richardson et al. (2008) to estimate the random uncertainty of F_c , LE , and H . Briefly, data from the same half hour period of consecutive days were screened for similar micrometeorological conditions defined as those for which R_n differed by less than 75 $W\ m^{-2}$, air temperature differed by less than 3 °C, and average wind speed differed by less than 1 $m\ s^{-1}$ under the assumption that fluxes measured during these conditions on consecutive days should be similar and any differences are related to the random error of eddy covariance measurements. Differences in F_c , LE , and H identified using this approach were found to be linearly related to the mean flux (Richardson et al., 2008, 2006), the model for which is taken to be the random uncertainty of the flux measurements. Random uncertainty of the seasonal sums of F_c , LE , and H is taken to be the mean absolute value of these fluxes multiplied by the percent random uncertainty following Stoy et al. (2006a, 2006b).

Random uncertainty was propagated through the gapfilling routines following the recommendations of Motulsky and Ransnas (1987) by perturbing the input flux observations with a random value drawn from a normal distribution multiplied by the random uncertainty calculated using the Richardson et al. (2008) routine. This procedure was repeated 100 times for each day of observations, and 100 parameter sets for the gapfilling model were subsequently fit for each day using least squares regression. Each of the corresponding parameter sets for the case of F_c were tested for convergence of the optimization routine, and all parameter sets were adjusted for missing values using the routines for large and small gaps described above. Gapfilling uncertainty was then determined as the standard deviation of the April – September seasonal flux sum. Total uncertainty for each flux over the study periods was calculated by summing the variance attributable to random uncertainty and that attributable to gapfilling parameter uncertainty - noting the propagation of random error into the estimate of gapfilling uncertainty - in order to obtain a conservative value of total flux uncertainty for each field for each study period.

2.5 Energy Balance

The surface-atmosphere energy balance is rarely closed using the eddy covariance and micrometeorological measurements of R_n , G , LE , and H at a single tower (Franssen et al., 2010; Stoy et al., 2013; Wilson et al., 2002), although 26 of the 173 tower sites studied by Stoy et al. (2013) had an energy balance closure greater than 100%. Energy balance closure for the winter wheat, spring wheat, and fallow fields was calculated as the percent of the sum of available energy during the April – September study periods (i.e. $R_n - G$) that is realized as the measured sum of sensible and latent heat fluxes ($H + \lambda E$), excluding periods for which turbulent fluxes were gapfilled.

2.6 Atmospheric Boundary Layer Height Model

We used surface-atmosphere exchange observations and a simple one-dimensional *ABL* model to test the conceptual model of Gameda et al. (2007), which assumes that fallow results in h on the order of 2-3 km and vegetated surfaces result in h on the order of 500 m in the Canadian Prairies. h was modeled for each of the management practices following Luyssaert et al. (2014):

$$\Delta h = (H_v \Delta t) / (\rho_a c_p \gamma_v) \quad (3)$$

where Δh is the change in h in meters per unit time (Δt), ρ_a is the air density in kg m^{-3} , c_p is the specific heat capacity in $\text{J kg}^{-1} \text{K}^{-1}$, γ_v is the virtual temperature inversion strength in K m^{-1} set to 0.003 K m^{-1} , and H_v is the virtual heat flux in W m^{-2} obtained by measured H and LE :

$$H_v = H + 0.07LE. \quad (4)$$

Equation 3 was calculated for each day during the study periods and h was set to 150 m at the beginning of each day, also following Luyssaert et al. (2014). We focus our comparisons on model predictions of maximum daily h for the different agricultural treatments.

Modeled h is compared to the modeled height of the lifted condensation level (h_{LCL}), which was calculated by dry-adiabatically raising an air parcel with near-surface temperature and moisture to its saturation point approximated by (Stull, 2012):

$$T_{LCL} = \frac{2840}{3.5 \ln(T_a) - \ln\left(\frac{Pr}{0.622+r}\right) - 7.108} + 55 \quad (5)$$

with atmospheric pressure at the surface (P) in kPa, the mixing ratio r , and T_a in K. We focus our discussion of h and h_{LCL} on the 2014 growing season when spring wheat was measured by the first tower and fallow was measured by the second tower.

3. Results

3.1 Prevailing weather conditions

On average, 2014 was cooler, cloudier, and wetter than 2013. Mean annual T_a during 2013 was 7.0 °C and mean April-September T_a was 14.3 °C. Mean annual T_a (6.5°C) and April-September mean T_a (13.7 °C) were about 0.5 °C cooler in 2014 than 2013. Annual cumulative SW_{in} was 3% greater in 2013 (5290 MJ m⁻² year⁻¹) than in 2014 (5140 MJ m⁻² year⁻¹), and the April-September cumulative incident SW_{in} was 5% greater in 2013. Cumulative precipitation measured at the Moccasin SCAN site was 452 mm in 2013 and 503 mm in 2014. The study period cumulative precipitation was 360 mm in 2013 and 391 mm in 2014, 162 mm of which fell during a large precipitation event during Aug. 21-24 shortly after the harvest of the spring wheat field on Aug. 18, 2014.

3.2 Plant Height Growth

Canopy height was near zero after the 2012 fallow growing season that preceded winter wheat planting. Measureable winter wheat growth began in early May 2013 (Figure 3). The winter wheat crop reached its maximum height of 0.64 m on July 9, 2013. The canopy was reduced to stubble measured at 0.2 m after harvest on August 7, 2013. With the exception of variation due to snow cover, the plant canopy remained at 0.2 m during the 2013/2014 winter, and was effectively 0 m after spring wheat planting on May 2, 2014. Measurable spring wheat crop growth began on May 26 and reached 0.83 m on July 25, *ca.* two weeks after winter wheat reached its maximum

height in 2013. Canopy height was likewise reduced to 0.2 m stubble after harvest on Aug. 18, 2014 (Figure 3).

3.3 u^* Thresholds and Data Availability

The Reichstein et al. (2005) algorithm selects a unique u^* threshold for three month periods, which correspond to April through June and July through September during the study period. The identified u^* thresholds were 0.077 m s^{-1} for 2013 winter wheat, 0.048 m s^{-1} for 2014 spring wheat, and 0.073 m s^{-1} for the fallow field during the April through June period, and 0.054 m s^{-1} for 2013 winter wheat, 0.092 m s^{-1} for 2014 spring wheat, and 0.057 m s^{-1} for the fallow field during the July through September period. Turbulent flux observations taken during periods where measured u^* was less than these values during nighttime were removed from the observational record and filled using the gapfilling procedures described in sections 2.3.2-2.3.4. The percent of missing data for each treatment that resulted from the u^* filter and logical filters, as well as power system and instrumental failures, is described in Table 3. Uncertainty in flux sums that result from missing data is propagated through the data gapfilling routines to result in monthly and seasonal uncertainty estimates as described in section 2.4.

3.4 Carbon Dioxide Fluxes

The cumulative sum of F_c in units of carbon in CO_2 (C- CO_2) in all fields during the study period is shown in Figure 4A. The 2013 winter wheat field became a net sink of carbon (since April 1, 2013) around May 8, 2013, and the 2014 spring wheat field became a net sink of carbon (since April 1, 2014) around May 23, 2014. The winter wheat field had a cumulative C- CO_2 uptake of $-208 \pm 53 \text{ g C m}^{-2}$ during the 2013 April – September study period (expressed as the mean of the gapfilling model simulations plus or minus one standard deviation of random plus gapfilling uncertainty), and the spring wheat field took up $-107 \pm 29 \text{ g C m}^{-2}$ during the 2014 study period (Figure 4B). The fallow field lost $135 \pm 73 \text{ g C-}\text{CO}_2$ to the atmosphere during the 2014 study period with two brief periods of net C- CO_2 uptake of *ca.* $10\text{-}20 \text{ g C m}^{-2}$ corresponding to periods of weedy growth near the tower during late June and weedy growth following the large rain event

after the spring wheat harvest in August. The larger uncertainty for F_c in the fallow field is attributable to greater uncertainty in net respiratory fluxes calculated by the Richardson et al. (2008) algorithm, and the larger uncertainty in F_c at the winter wheat field *versus* the spring wheat field is due in part to the larger uncertainties attributable to larger average flux magnitude.

Maximum monthly F_c occurred at the winter wheat field in June at -186 ± 47 g C-CO₂ month⁻¹ (Figure 4B) and in July at the spring wheat field at -113 ± 30 g C-CO₂ month⁻¹, which is of greater magnitude than F_c in June at the spring wheat field at -100 ± 27 g C-CO₂ month⁻¹ as identified by a two-sided t-test ($p < 0.05$, Figure 4B). Monthly F_c from the fallow field peaked in July at 39 ± 22 g C-CO₂ month⁻¹.

3.5 Evapotranspiration

Cumulative ET from the 2013 winter wheat field was 410 ± 36 mm during the study period (Figure 5A), significantly less than the 437 ± 44 mm observed for the 2014 spring wheat field during the cooler and wetter 2014 ($p < 0.05$), but only by some 6% and reflective of the differences in April-September precipitation observed between years. The rate of increase in cumulative ET in the 2014 fallow field was not as steep as that in the winter wheat and spring wheat fields, and the Apr. – Sept. total reached 275 ± 39 mm, *ca.* 2/3 of observed ET from the spring wheat field. Monthly total ET from the winter wheat field peaked in June at 130 ± 10 mm and monthly total ET from the spring wheat field peaked in July at 147 ± 10 mm (Figure 5B), mirroring the monthly patterns of F_c uptake (Figure 4B). Monthly total ET in the Fallow treatment peaked in June and July at *ca.* 69 ± 10 mm.

3.6 Sensible Heat Fluxes and Energy Balance Closure

H for all treatments before 2013 winter wheat and 2014 spring wheat growth began was similar at *ca.* 2.5 MJ m⁻² day⁻¹. After crop growth initiated, of the increase in cumulative H diverged between the spring wheat and fallow fields (Figure 6A). The winter wheat and spring wheat fields had similar totals of H of 445 ± 66 and 465 ± 67 MJ m⁻² for the respective study periods. H at the fallow field reached 759 ± 98 MJ m⁻² for the study period, an increase in H proportional to the

decrease in ET (Figures 5A and 6A). Monthly H was greater at the fallow field than the wheat fields until harvest in August, especially from the winter wheat field which ceased growth and was harvested sooner than the spring wheat field (see e.g. Figures 6B). Energy balance closure during the study period was calculated to be 86% at the winter wheat field, 105% at the spring wheat field, and 107% at the fallow field.

3.7 Modeled Atmospheric Boundary Layer Height and Lifted Condensation Level

Modeled h using surface flux data from the spring wheat and fallow fields that were both measured in 2014 (Figure 7A) demonstrates a seasonal pattern that follows the seasonal trend in H (Figure 6B), which is the largest contributor to H_v (equation 4). Maximum modeled h was 2070 m for the spring wheat field and 2440 m for the fallow field (Figure 7A), and the maximum daily difference between the two fields was over 900 m (Figure 7B). The mean difference in modeled maximum daily h between the spring wheat and fallow fields was 210 m during the April – September study period.

Differences in modeled h_{LCL} between fallow and spring wheat were minor and averaged 20 m over the study period. As a consequence, h_{LCL} modeled using data from the spring wheat and fallow fields were averaged for subsequent analyses. The mean value of maximum daily h_{LCL} was *ca.* 1600 m but exceeded 3500 m during the month before the spring wheat harvest (Figure 7A). A comparison of modeled h and h_{LCL} for both spring wheat and fallow indicates that the growing convective ABL height reaches the LCL ($h > h_{LCL}$) on average some 30 minutes earlier for fallow, and such crossings occurred during 81 days with fallow and 74 days with spring wheat (Figure 8).

4. Discussion

It is important to recall that the 2014 study period was cooler and wetter than 2013 and comparisons between the winter wheat field (measured in 2013) and the spring wheat and fallow fields (measured in 2014) are subject to these differences in prevailing weather conditions. We

first discuss differences in surface-atmosphere carbon dioxide exchange with a focus on the winter wheat and spring wheat fields – noting that the fallow field lost carbon to the atmosphere as expected – followed by an analysis of the impacts of spring wheat *versus* fallow management on the turbulent fluxes of latent and sensible heat and consequences for modeled atmospheric boundary layer development.

4.1 Carbon Dioxide Exchange

The spring wheat and fallow fields – both measured during 2014 – lost a similar amount of C-CO₂ during April and May before crop growth and during initial stages of spring wheat emergence as quantified as measureable height growth by the sonic distance sensor (Figure 2; 37 g C m⁻² for April and May for the spring wheat field and 43 g C m⁻² for April and May for the fallow field). Mean T_a (8.2 °C) and soil temperature (9.0 °C in the spring wheat field and 9.1 °C in the fallow field) were characteristically low during this period and carbon fluxes during the largely fallow period in both fields behaved in a similar manner. Following crop emergence, the spring wheat field turned into a net carbon sink (since April 1, 2014) in mid-June. Outside of brief periods of weedy growth, the consistent C source from the fallow field resulted in net April-September C-CO₂ losses (135 ± 73 g C m⁻² study period⁻¹) that were of similar magnitude to C sinks in the spring wheat field (-107 ± 29 g C m⁻² study period⁻¹).

Both wheat fields were a net sink of C-CO₂ during the study period on the order of 200 g C m⁻² for the winter wheat field and 100 g C m⁻² for the spring wheat field. It has been suggested that non-woody C₃-dominated ecosystems should have similar net ecosystem productivity due to metabolic and organism size constraints (Peichl et al., 2013), bringing into question why the winter wheat field had two-fold greater C-CO₂ uptake than the spring wheat field, especially when canopy height growth was greater in the spring wheat field than the winter wheat field (Figure 3). Potential explanations include 1) differences in climatological variables between growing seasons; 2) net C-CO₂ efflux in the spring wheat field during April and May after planting (i.e. the difference among fields during the April-September study period is a result of

planting date); or 3) respiratory losses from organic matter left from winter wheat field in the spring wheat field noting that chemical fallow preceded winter wheat. We briefly discuss each sequentially.

To test the effects of the cooler and wetter conditions on determining 2014 fluxes, we parameterized the daily F_c model using 2014 flux and micrometeorological observations and forced it with 2013 SW_{in} and T_a observations. This analysis follows the approach of Richardson et al. (2007) to interpret the role of climate *versus* biological and management responses to climate variability of surface-atmosphere fluxes. The cumulative sum of F_c during the study period using the 2014 models forced by 2013 meteorology was $-120 \text{ g C-CO}_2 \text{ m}^{-2}$, a difference of only a *ca.* $10 \text{ g C-CO}_2 \text{ m}^{-2}$ for the study period, suggesting that differences in F_c are not primarily due to differences in growing season climate.

The second explanation – differences in crop emergence timing – contributes to the differences in CO_2 uptake between the winter wheat and spring wheat crops. The spring wheat field lost *ca.* 40 g C m^{-2} during April and May before the spring wheat field became a net daily sink for C-CO_2 around May 29, 2014. Due to earlier planting and thereby crop emergence, the winter wheat field became a net C-CO_2 sink at the beginning of May. Differences in seasonal F_c even among winter wheat treatments has been attributed to differences in early season crop development (Dufranne et al., 2011).

The third explanation – that there is a greater respiratory source in the spring wheat field – can be confirmed by analyzing the parameters of the gapfilling model (eq. 1). The mean R_{10} parameter during the period of peak growth in the winter wheat field is *ca.* $2.5 \mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$, but over $3 \mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ in the spring wheat field. Using mean R_{10} and E_0 parameters in the respiratory component of equation 1 with observed T_a – noting that equation 1 is designed to gapfill F_c rather than partition it into its components (Reichstein et al., 2012; Wehr et al., 2016) - results in a modeled RE of 263 g C m^{-2} during the study period in the winter wheat field and

g C m⁻² during the study period in the spring wheat field. This difference may seem large, but using these modeled *RE* values to model *GEP* as the difference between *F_c* and *RE* results in -466 g C m⁻² during the study period in the winter wheat field and -595 g C m⁻² in the spring wheat field, a difference of about 25% that is proportional to the difference in maximum canopy height (Figure 3). Greater *GEP* in the spring wheat field results in more labile C available for *RE* (Högberg et al., 2001; Ryan and Law, 2005; Stoy et al., 2007), but differences in modeled *RE* are greater than differences in *GEP* among fields, pointing to a respiratory source in the spring wheat field that is not present in the winter wheat field. These results are consistent with the findings of Schmidt et al. (2012) and Waldo et al. (2016), who demonstrated the importance of plant residues from previous crops to the crop carbon balance. The plant residues that were left after the 2013 winter wheat harvest (e.g. Figure 3) that remained in the field during the 2014 spring wheat crop - noting that chemical fallow preceded winter wheat - cannot be excluded as a cause of the differences in C-CO₂ uptake between our winter wheat and spring wheat observations. These findings are in agreement with the findings of Peichl et al. (2013) in that differences in carbon metabolism in these non-woody C₃ systems was dominated by the impacts of land management.

4.2 Evapotranspiration and Water Management

Daily *ET* sums quantified here are similar to or greater than other studies in wheat cropping systems. Maximum daily *ET* was *ca.* 7 mm day⁻¹ for the winter wheat crop (in June) and the spring wheat crop (in July), which corresponds to low surface resistance to water by wheat (Bonan, 2008) during periods with sufficient soil moisture and the high incident radiation load of the study sites. That being said, the maximum value of α_{PT} calculated by the gapfilling models was 0.75 for winter wheat and 0.85 for spring wheat, far less than the theoretical maximum of 1.26 described by Priestley and Taylor (1972) and values found for irrigated wheat crops (e.g. 1.17-1.26; Zhang et al., 2004), suggesting limitations to the transport of water between soil and atmosphere that may be due to either stomatal closure or limited evaporation due to dry soil surface conditions. Future work should seek to identify the source of this water limitation,

perhaps using advanced algorithms for estimating the contribution of evaporation and transpiration to ET using high frequency eddy covariance data (Scanlon and Kustas, 2010; Scanlon and Sahu, 2008; Sulman et al., 2016).

There were notable differences amongst ET observed here and in other wheat cropping systems. Anthoni et al. (2004) found maximum rates of ET on the order $3\text{--}4\text{ mm day}^{-1}$ in mid-May in a winter wheat crop in Germany, far lower than the 7 mm day^{-1} maximum observed here. We can estimate SW_{in} from $PPFD$ observations at the Gebesee study site of Anthoni et al. (2004) and approximate annual SW_{in} to be on the order of $3500 - 4010\text{ MJ m}^{-2}\text{ year}^{-1}$, which is approximately $2/3$ of the annual SW_{in} received by the Judith Basin study sites. In other words, differences in maximum ET among our observations and those of Anthoni et al. (2004) are due in part to lower SW_{in} during the growing season in central Europe.

ET was predictably higher in the winter wheat and spring wheat fields than the fallow field (Figure 5A), confirming that fallowing results in soil water savings if lateral water transport and drainage losses can be assumed to be minimal during the growth period. Winter wheat ET was 410 mm during the April – September study period, which represented *ca.* 90% of the total precipitation based on the data from the Moccasin SCAN site in 2013 for the same time frame, noting that precipitation maps suggest that the study fields receive slightly more precipitation than the Moccasin SCAN site on average due to topographical influence. Total ET for the spring wheat and fallow fields during the 2014 study period was 440 mm and 315 mm , which was *ca.* 90% and 60% respectively of Moccasin SCAN precipitation observations in 2014. In other words, wheat cropping returned almost all incident precipitation to the atmosphere, while the fallow field left over 100 mm of water available for a subsequent crop, or for deep drainage.

It is likely that the no-till treatment reduced ET in the fallow field *versus* a situation in which it was tilled; Chi et al. (2016) for example demonstrated that tillage tended to increase soil evaporation. It is important to note that ET may be underestimated in the winter wheat field and overestimated in the spring wheat and fallow fields owing to imperfect energy balance closure,

but it is unclear if radiometric or turbulent flux observations are the cause of the imperfect closure and current practices suggest that turbulent flux observations should not be adjusted to match radiometric observations (Baldocchi, 2008).

The amount of water that would need to be replenished to equal the water savings conferred by fallow during the study period is on the order of 100 mm. Stubble height in our case is 0.2 m (Figure 3), and snow tended to accumulate to stubble height in the study fields. If we assume a characteristic snow density in MT to be on the order of 0.08 g cm^{-3} (Jacobson, 2010; Welch et al., 2015), the depth of water that corresponds to 0.2 m of snow of this density is 16 mm of water, an order of magnitude less than the amount of water that needs to be replenished to account for water use of the wheat fields. In other words, changes in stubble management are unlikely to make up the difference in water lost to the atmosphere among the wheat and fallow fields. If water in the soil column during the non-growing season is replenished such that soil moisture is at or near field capacity before crop growth regardless of prior agricultural treatment, the water savings benefits of fallow should be questioned.

4.3 Impacts of Surface Conditions on Atmospheric Boundary Layer & Lifted Condensation Level

Cumulative H at both wheat fields was *ca.* 450 MJ m^{-2} during the study period, with important seasonal differences due to crop development and harvest that have implications for energy inputs into the *ABL* and its diurnal growth (Figure 7A). The choice to fallow is among these management decisions with implications for *ABL* dynamics as evidenced by large differences in modeled maximum daily h between the spring wheat and fallow fields, especially during the peak crop growth period in June and July in our case (Figure 7B).

Gameda et al. (2007) studied the impacts of fallow reduction on regional climate in the Canadian Prairie Provinces and used a conceptual model to argue that fallow results in (modeled) h on the order of 2-3 km but h resulting from vegetated surfaces is on the order of 0.5 km. Their reasoning is that R_n in fallow is greater due to lower albedo and is preferentially partitioned into H , which results in greater virtual heat flux following equation 4, while vegetated surfaces

partition R_n preferentially into LE resulting in a cooler and shallower ABL . Figure 7A agrees with Gameda et al. (2007) in that the modeled h is higher as a result of observed surface fluxes from the fallow field *versus* the spring wheat field, but our results differ in magnitude. Our results suggest that the conceptual model of Gameda et al. (2007) underestimates h in wheat scenarios. It is important to note that the one-dimensional simulations presented here simply estimate the influence of a single surface (spring wheat *versus* fallow) on a modeled vertical column of the atmosphere and do not seek to simulate realistic atmospheric flows.

The intersection of the convective ABL height with the LCL is a necessary, albeit not sufficient, condition for the initiation of local convection (e.g. Juang et al., 2007; Juang et al., 2007). Maximum daily h_{LCL} differed on average by *ca.* 20 m between the two plots, which can be explained by their close proximity. Low relative humidities and hot air temperatures during the month preceding spring wheat harvest led to modeled h_{LCL} greater than 2500 m, which exceeded maximum h and suggests that convective precipitation during this period is unlikely. There was a tendency to reach h_{LCL} sooner and more often in the fallow simulation due to higher H and thus h above fallow, (Figure 8). These findings, however, need to be interpreted with caution for two reasons. (i) Differences in surface meteorology between the experimental plots are small, and (ii) h as well as h_{LCL} are calculated from local quantities, which do not account for feedbacks between the regional surface energy balance and the thermodynamic state of the atmosphere as well as non-local effects such as the development of mesoscale circulations between patches with different surface characteristics (e.g. Taylor et al., 2007).

Overall, large-scale reduction of fallow has the potential to decrease h by several hundred meters. Similarly, Betts et al. (2013b) found that there was a systematic reduction in h_{LCL} in the Canadian Prairies of approximately 200 m from 1953 to 2011, which is similar to the calculated mean change in h of 210 m, highlighting the uncertain nature of the net effect on convective development. h and h_{LCL} have more crossings in the fallow simulation yet fallow reduction is thought to result in an increase in cloud development and convective initiation. Our results

suggest that studies of the effects of fallow on near-surface temperature and humidity following Betts et al. (2013b) should be extended to the regions of fallow reduction in the U.S. to ascertain how land use changes across the entire northern North American Great Plains has impacted regional climate.

The details of the differences – and similarities – among modeled h and h_{LCL} as they impact cloud formation processes and convective precipitation over wheat growing areas has yet to be ascertained. Given the reduction in the practice of fallow over the past four decades on the order of tens of millions of hectares (Figure 1), the role of cropping systems and planting decisions needs to be further investigated to quantify the proposed role of agricultural management in the observed summertime cooling trend and increase in precipitation across parts of the northern North American Great Plains (Betts et al., 2013a, 2013b; Mahmood et al., 2014). Future research should study the impacts of management patterns in space and time in h and h_{LCL} dynamics - and measurements of h should be made or estimated from radiosonde data (e.g. Seidel et al., 2010) – to understand the consequences of land management across the northern North American Great Plains on atmospheric dynamics of importance to convective precipitation.

5. Conclusions

Greater CO₂ uptake by the winter wheat field (*ca.* -200 g C m⁻²) than the spring wheat (*ca.* -100 g C m⁻²) field during the April – September study period were largely due to the effects of ecosystem management: Carbon losses in April and May from the spring wheat field combined with a larger respiratory source during the main growth period – consistent with the respiration of residues from the previous winter wheat crop – explain most of the difference in F_c among the wheat fields. CO₂ losses from the wheat fields before plant emergence and after harvest occur at a similar rate as CO₂ losses from the fallow field, which represented a total C-CO₂ loss to the atmosphere of over 100 g C m⁻² during the April – September study period. Total April – September ET was similar between the winter wheat and spring wheat fields despite differences

in prevailing climatic conditions and crop emergence and harvest dates, suggesting that ecosystem water use efficiency was greater in the winter wheat field with greater F_c . Cropping systems management also had important implications for H and thereby ABL dynamics, the maximum daily value of which was *ca.* 200 m less in spring wheat *versus* fallow simulations, which is on the order of the reduction of h_{LCL} demonstrated by Betts et al. (2013b) from the period 1951-1991 to 1992-2011. Future research should quantify the amount of ET resulting from evaporation and transpiration to understand sources of water to the atmosphere and its interaction with plant water use and growth, and investigate the implications of widespread changes in agricultural management in the northern North American Great Plains – including alternate cropping sequences – on regional climatology.

Acknowledgements

This work was supported by the Montana Wheat and Barley Committee, the Montana University System Water Center, the USDA National Institute of Food and Agriculture Hatch project 228396, and NSF DEB 1552976 ‘The role of ecosystem management on boundary layer development and precipitation in the Northern Plains’. We would like to thank Adam Sigler, Simon Fordyce, Aiden Johnson, Robby Robertson, Dr. Lin Hua, and Angela Tang for field assistance and coordination and David Walker for advice on agricultural statistics. We would also like to thank the landowner Bob Otten and the farm operator Brandon Morris for permission, coordination, and invaluable insight. Stephanie Ewing, Perry Miller, and Rob Payn provided valuable comments on the manuscript and Clain Jones and Adam Sigler provided valuable information on soil characteristics.

References

- Anthoni, P.M., Freibauer, A., Kolle, O., Schulze, E.-D., 2004. Winter wheat carbon exchange in Thuringia, Germany. *Agric. For. Meteorol.* 121, 55–67.
- Anthoni, P.M., Knohl, A., Rebmann, C., Freibauer, A., Mund, M., Ziegler, W., Kolle, O., Schulze, E.-D., 2004. Forest and agricultural land-use-dependent CO₂ exchange in Thuringia, Germany. *Glob. Chang. Biol.* 10, 2005–2019.
- Aubinet, M., Grelle, A., Ibrom, A., Rannik, U., Moncrieff, J., Foken, T., Kowalski, A.S., Martin, P.H., Berbigier, P., Bernhofer, C., Clement, R., Elbers, J., Granier, A., Grunwald, T., Morgenstern, K., Pilegaard, K., Rebmann, C., Snijders, W., Valentini, R., Vesala, T., 2000. Estimates of the annual net carbon and water exchange of forests: The EUROFLUX methodology. *Adv. Ecol. Res.* 30, 113–175.
- Baldocchi, D.D., 2008. “Breathing” of the terrestrial biosphere: lessons learned from a global network of carbon dioxide flux measurements systems, Turner Review. *Aust. J. Bot.* 56, 1–26.
- Betts, A.K., Desjardins, R., Worth, D., 2013a. Cloud radiative forcing of the diurnal cycle climate of the Canadian Prairies. *J. Geophys. Res. Atmos.* 118, 8935–8953.
- Betts, A.K., Desjardins, R., Worth, D., Cerkowniak, D., 2013b. Impact of land use change on the diurnal cycle climate of the Canadian Prairies. *J. Geophys. Res. Atmos.* 118, 11–996.
- Béziat, P., Ceschia, E., Dedieu, G., 2009. Carbon balance of a three crop succession over two cropland sites in South West France. *Agric. For. Meteorol.* 149, 1628–1645.
- Billesbach, D.P., Berry, J.A., Seibt, U., Maseyk, K., Torn, M.S., Fischer, M.L., Abu-Naser, M., Campbell, J.E., 2014. Growing season eddy covariance measurements of carbonyl sulfide and CO₂ fluxes: COS and CO₂ relationships in Southern Great Plains winter wheat. *Agric. For. Meteorol.* 184, 48–55.
- Bonan, G.B., 2008. Forests and climate change: forcings, feedbacks, and the climate benefits of forests. *Science* (80-.). 320, 1444–1449.

637 Burgess, M.H., Miller, P.R., Jones, C.A., 2012. Pulse crops improve energy intensity and
638 productivity of cereal production in Montana, USA. *J. Sustain. Agric.* 36, 699–718.

639 Buyanovsky, G.A., Wagner, G.H., 1998. Changing role of cultivated land in the global carbon
640 cycle. *Biol Fertil Soils* 27, 242–245.

641
642 Chi, J., Waldo, S., Pressley, S., O’Keeffe, P., Huggins, D., Stöckle, C., Pan, W.L., Brooks, E.,
643 Lamb, B., 2016. Assessing carbon and water dynamics of no-till and conventional tillage
644 cropping systems in the inland Pacific Northwest US using the eddy covariance method.
645 *Agric. For. Meteorol.* 218–219, 37–49.

646 Cihacek, L.J., Ulmer, M.G., 1995. Estimated soil organic carbon losses from long-term crop-
647 fallow in the Northern Great Plains of the USA. *Soil Manag. Greenh. Eff.* 85–92.

648 Dhuyvetter, K.C., Thompson, C.R., Norwood, C.A., Halvorson, A.D., 1996. Economics of
649 dryland cropping systems in the Great Plains: A review. *J. Prod. Agric.* 9, 216–222.

650 Dufranne, D., Moureaux, C., Vancutsem, F., Bodson, B., Aubinet, M., 2011. Comparison of
651 carbon fluxes, growth and productivity of a winter wheat crop in three contrasting growing
652 seasons. *Agric. Ecosyst. Environ.* 141, 133–142.

653 Falge, E., Baldocchi, D., Olson, R., Anthoni, P., Aubinet, M., Bernhofer, C., Burba, G.,
654 Ceulemans, R., Clement, R., Dolman, H., Granier, A., Gross, P., Grunwald, T., Hollinger,
655 D., Jensen, N.O., Katul, G., Keronen, P., Kowalski, A., Lai, C.T., Law, B.E., Meyers, T.,
656 Moncrieff, J., Moors, E., Munger, J.W., Pilegaard, K., Rannik, U., Rebmann, C., Suyker, A.,
657 Tenhunen, J., Tu, K., Verma, S., Vesala, T., Wilson, K., Wofsy, S., 2001a. Gap filling
658 strategies for defensible annual sums of net ecosystem exchange. *Agric. For. Meteorol.* 107,
659 43–69.

660 Falge, E., Baldocchi, D., Olson, R., Anthoni, P., Aubinet, M., Bernhofer, C., Burba, G.,
661 Ceulemans, R., Clement, R., Dolman, H., Granier, A., Gross, P., Grunwald, T., Hollinger,
662 D., Jensen, N.O., Katul, G., Keronen, P., Kowalski, A., Lai, C.T., Law, B.E., Meyers, T.,

663 Moncrieff, J., Moors, E., Munger, J.W., Pilegaard, K., Rannik, U., Rebmann, C., Suyker, A.,
 664 Tenhunen, J., Tu, K., Verma, S., Vesala, T., Wilson, K., Wofsy, S., 2001b. Gap filling
 665 strategies for long term energy flux data sets. *Agric. For. Meteorol.* 107, 71–77.
 666 FAO, F.A.O., 2013. Statistical Yearbook 2013: World Food and Agriculture. Food Agric. Organ.
 667 United Nations, Rome 289.
 668 Franssen, H.J.H., Stöckli, R., Lehner, I., Rotenberg, E., Seneviratne, S.I., 2010. Energy balance
 669 closure of eddy-covariance data: A multisite analysis for European FLUXNET stations.
 670 *Agric. For. Meteorol.* 150, 1553–1567. doi:10.1016/j.agrformet.2010.08.005
 671 Gameda, S., Qian, B., Campbell, C.A., Desjardins, R.L., 2007. Climatic trends associated with
 672 summerfallow in the Canadian Prairies. *Agric. For. Meteorol.* 142, 170–185.
 673 Goulden, M.L., Daube, B.C., Fan, S.-M., Sutton, D.J., Bazzaz, A., Munger, J.W., Wofsy, S.C.,
 674 1997. Physiological responses of a black spruce forest to weather. *J. Geophys. Res.* 102,
 675 28987–28996.
 676 Gu, L.H., Falge, E.M., Boden, T., Baldocchi, D.D., Black, T.A., Saleska, S.R., Suni, T., Verma,
 677 S.B., Vesala, T., Wofsy, S.C., Xu, L.K., 2005. Objective threshold determination for
 678 nighttime eddy flux filtering. *Agric. For. Meteorol.* 128, 179–197.
 679 Hawkesford, M.J., Araus, J., Park, R., Calderini, D., Miralles, D., Shen, T., Zhang, J., Parry,
 680 M.A.J., 2013. Prospects of doubling global wheat yields. *Food Energy Secur.* 2, 34–48.
 681 Högberg, P., Nordgren, A., Buchmann, N., Taylor, A.F.S., Ekblad, A., Högberg, M.N., Nyberg,
 682 G., Ottoson-Löfvenius, M., Read, D.J., 2001. Large-scale forest girdling shows that current
 683 photosynthesis drives soil respiration. *Nature* 411, 789–792.
 684 Ingwersen, J., Steffens, K., Högy, P., Warrach-Sagi, K., Zhunusbayeva, D., Poltoradnev, M.,
 685 Gäbler, R., Wizemann, H.-D., Fangmeier, A., Wulfmeyer, V., Streck, T., 2011. Comparison
 686 of Noah simulations with eddy covariance and soil water measurements at a winter wheat
 687 stand. *Agric. For. Meteorol.* 151, 345–355.
 688 IPCC, 2007. Climate Change 2007: The Physical Science Basis. Contribution of Working Group

I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change.
 Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.

Jacobson, M.D., 2010. Inferring snow water equivalent for a snow-covered ground reflector using
 GPS multipath signals. *Remote Sens.* 2, 2426. doi:10.3390/rs2102426

Juang, J.-Y., Katul, G.G., Porporato, A., Stoy, P.C., Siqueira, M.S., Detto, M., Kim, H.-S., Oren,
 R., 2007. Eco-hydrological controls on summertime convective rainfall triggers. *Glob.*
Chang. Biol. 887–896.

Juang, J.-Y., Porporato, A., Stoy, P.C., Siqueira, M.B.S., Oishi, A.C., Detto, M., Kim, H.-S.,
 Katul, G.G., 2007. Hydrologic and atmospheric controls on initiation of convective
 precipitation events. *Water Resour. Res.* 43.

Kaimal, J.C., Finnegan, J.J., 1994. Atmospheric boundary layer flows: their structure and
 measurements. Oxford, New York.

Lloyd, J., Taylor, J.A., 1994. On the temperature dependence of soil respiration. *Funct. Ecol.* 8,
 315–323.

Long, J.A., Lawrence, R.L., Greenwood, M.C., Marshall, L., Miller, P.R., 2013. Object-oriented
 crop classification using multitemporal ETM+ SLC-off imagery and random forest.
GIScience Remote Sens. 50, 418–436.

Long, J.A., Lawrence, R.L., Miller, P.R., Marshall, L.A., Greenwood, M.C., 2014. Adoption of
 cropping sequences in northeast Montana: A spatio-temporal analysis. *Agric. Ecosyst.*
Environ. 197, 77–87.

Lubowski, R.N., Vesterby, M., Bucholtz, S., Baez, A., Roberts, M.J., 2006. Major uses of land in
 the United States, 2002. United States, Department of Agriculture, Economic Research
 Service.

Luyssaert, S., Jammet, M., Stoy, P.C., Estel, S., Pongratz, J., Ceschia, E., Churkina, G., Don, A.,
 Erb, K., Ferlicoq, M., Gielen, B., Grunwald, T., Houghton, R.A., Klumpp, K., Knohl, A.,
 Kolb, T., Kuemmerle, T., Laurila, T., Lohila, A., Loustau, D., McGrath, M.J., Meyfroidt, P.,

715 Moors, E.J., Naudts, K., Novick, K., Otto, J., Pilegaard, K., Pio, C.A., Rambal, S.,
 716 Rebmann, C., Ryder, J., Suyker, A.E., Varlagin, A., Wattenbach, M., Dolman, A.J., 2014.
 717 Land management and land-cover change have impacts of similar magnitude on surface
 718 temperature. *Nat. Clim. Chang.* 4, 389–393.

719 Mahmood, R., Pielke, R.A., Hubbard, K.G., Niyogi, D., Dirmeyer, P.A., McAlpine, C., Carleton,
 720 A.M., Hale, R., Gameda, S., Beltrán-Przekurat, A., 2014. Land cover changes and their
 721 biogeophysical effects on climate. *Int. J. Climatol.* 34, 929–953.

722 Mauder, M., Foken, T., 2004. Documentation and instruction manual of the eddy covariance
 723 software package TK2. Univ., Abt. Mikrometeorologie.

724 Merrill, S.D., Black, A.L., Fryrear, D.W., Saleh, A., Zobeck, T.M., Halvorson, A.D., Tanaka,
 725 D.L., 1999. Soil wind erosion hazard of spring wheat–fallow as affected by long-term
 726 climate and tillage. *Soil Sci. Soc. Am. J.* 63, 1768–1777.

727 Miller, P.R., Gan, Y., McConkey, B.G., McDonald, C.L., 2003. Pulse crops for the northern
 728 Great Plains. *Agron. J.* 95, 980–986.

729 Miller, P.R., McConkey, B.G., Clayton, G.W., Brandt, S.A., Staricka, J.A., Johnston, A.M.,
 730 Lafond, G.P., Schatz, B.G., Baltensperger, D.D., Neill, K.E., 2002. Pulse crop adaptation in
 731 the northern Great Plains. *Agron. J.* 94, 261–272.

732 Moncrieff, J.B., Malhi, Y., Leuning, R., 1996. The propagation of errors in long-term
 733 measurements of land-atmosphere fluxes of carbon and water. *Glob. Chang. Biol.* 2, 231–
 734 240.

735 Motulsky, H.J., Ransnas, L.A., 1987. Fitting curves to data using nonlinear regression: a practical
 736 and nonmathematical review. *FASEB J.* 1, 365–374.

737 Moureaux, C., Debacq, A., Hoyaux, J., Suleau, M., Tourneur, D., Vancutsem, F., Bodson, B.,
 738 Aubinet, M., 2008. Carbon balance assessment of a Belgian winter wheat crop (*Triticum*
 739 *aestivum* L.). *Glob. Chang. Biol.* 14, 1353–1366.

740 Oren, R., Hsieh, C.I., Stoy, P.C., Albertson, J.D., McCarthy, H.R., Harrell, P., Katul, G.G., 2006.
 741 Estimating the uncertainty in annual net ecosystem carbon exchange: spatial variation in
 742 turbulent fluxes and sampling errors in eddy-covariance measurements. *Glob. Chang. Biol.*
 743 12, 883–896.
 744 Papale, D., 2012. Data gap filling, in: *Eddy Covariance*. Springer, pp. 159–172.
 745 Papale, D., Reichstein, M., Aubinet, M., Canfora, E., Bernhofer, C., Kutsch, W., Longdoz, B.,
 746 Rambal, S., Valentini, R., Vesala, T., Yakir, D., 2006. Towards a standardized processing of
 747 Net Ecosystem Exchange measured with eddy covariance technique: algorithms and
 748 uncertainty estimation. *Biogeosciences* 3, 571–583.
 749 Peichl, M., Sonnentag, O., Wohlfahrt, G., Flanagan, L.B., Baldocchi, D.D., Kiely, G., Galvagno,
 750 M., Gianelle, D., Marcolla, B., Pio, C., Migliavacca, M., Jones, M.B., Saunders, M., 2013.
 751 Convergence of potential net ecosystem production among contrasting C3 grasslands. *Ecol.*
 752 *Lett.* 16, 502–512.
 753 Priestley, C.H.B., Taylor, R.J., 1972. On the assessment of surface heat flux and evaporation
 754 using large-scale parameters. *Mon. Weather Rev.* 100, 81–92.
 755 Reichstein, M., Falge, E., Baldocchi, D., Papale, D., Aubinet, M., Berbigier, P., Bernhofer, C.,
 756 Buchmann, N., Gilmanov, T.G., Granier, A., Grünwald, T., Havránková, K., Ilvesniemi, H.,
 757 Janous, D., Knohl, A., Laurila, T., Lohila, A., Loustau, D., Matteucci, G., Meyers, T.,
 758 Miglietta, F., Ourcival, J.-M., Pumpanen, J., Rambal, S., Rotenberg, E., Sanz, M.,
 759 Tenhunen, J., Seufert, G., Vaccari, F., Vesala, T., Yakier, D., Valentini, R., 2005. On the
 760 separation of net ecosystem exchange into assimilation and ecosystem respiration: review
 761 and improved algorithm. *Glob. Chang. Biol.* 11, 1424–1439.
 762 Reichstein, M., Stoy, P.C., Desai, A.R., Lasslop, G., Richardson, A.D., 2012. Partitioning of net
 763 fluxes, in: *Eddy Covariance*. Springer, pp. 263–289.
 764 Richardson, A.D., Hollinger, D.Y., Burba, G.G., Davis, K.J., Flanagan, L.B., Katul, G.G.,
 765 Munger, J.W., Ricciuto, D.M., Stoy, P.C., Suyker, A.E., Verma, S.B., Wofsy, S.C., 2006. A

766 multi-site analysis of random error in tower-based measurements of carbon and energy
767 fluxes. *Agric. For. Meteorol.* 136, 1–18.

768 Richardson, A.D., Hollinger, D.Y., Aber, J.D., Ollinger, S. V., Braswell, B.H., 2007.
769 Environmental variation is directly responsible for short- but not long-term variation in
770 forest-atmosphere carbon exchange. *Glob. Chang. Biol.* 13, 788–803. doi:10.1111/j.1365-
771 2486.2007.01330.x

772 Richardson, A.D., Mahecha, M.D., Falge, E., Kattge, J., Moffat, A.M., Papale, D., Reichstein,
773 M., Stauch, V.J., Braswell, B.H., Churkina, G., Kruijt, B., Hollinger, D.Y., 2008. Statistical
774 properties of random CO₂ flux measurement uncertainty inferred from model residuals.
775 *Agric. For. Meteorol.* 148, 38–50.

776 Ryan, M.G., Law, B.E., 2005. Interpreting, measuring, and modeling soil respiration.
777 *Biogeochemistry* 73, 3–27.

778 Scanlon, T.M., Kustas, W.P., 2010. Partitioning carbon dioxide and water vapor fluxes using
779 correlation analysis. *Agric. For. Meteorol.* 150, 89–99.

780 Scanlon, T.M., Sahu, P., 2008. On the correlation structure of water vapor and carbon dioxide in
781 the atmospheric surface layer: A basis for flux partitioning. *Water Resour. Res.* 44,
782 W10418.

783 Schmidt, M., Reichenau, T.G., Fiener, P., Schneider, K., 2012. The carbon budget of a winter
784 wheat field: An eddy covariance analysis of seasonal and inter-annual variability. *Agric.*
785 *For. Meteorol.* 165, 114–126.

786 Seidel, D.J., Ao, C.O., Li, K., 2010. Estimating climatological planetary boundary layer heights
787 from radiosonde observations: Comparison of methods and uncertainty analysis. *J.*
788 *Geophys. Res. Atmos.* 115, D16113.

789 Stoy, P., Mauder, M., Foken, T., Marcolla, B., Boegh, E., Ibrom, A., Arain, M.A., Arneth, A.,
790 Aurela, M., Bernhofer, C., Cescatti, A., Dellwik, E., Duce, P., Gianelle, D., van Gorsel, E.,
791 Kiely, G., Knohl, A., Margolis, H., McCaughey, H., Merbold, L., Montagnani, L., Papale,

792 D., Reichstein, M., Saunders, M., Serrano-Ortiz, P., Sottocornola, M., Spano, D., Vaccari,
 793 F., Varlagin, A., 2013. A data-driven analysis of energy balance closure across FLUXNET
 794 research sites: The role of landscape scale heterogeneity. *Agric. For. Meteorol.* 171, 137–
 795 152.

796 Stoy, P.C., Katul, G.G., Siqueira, M.B.S., Juang, J.-Y., McCarthy, H.R., Oishi, A.C., Uebelherr,
 797 J.M., Kim, H.-S., Oren, R., 2006a. Separating the effects of climate and vegetation on
 798 evapotranspiration along a successional chronosequence in the southeastern U.S. *Glob.*
 799 *Chang. Biol.* 12, 2115–2135.

800 Stoy, P.C., Katul, G.G., Siqueira, M.B.S., Juang, J.-Y., Novick, K.A., Uebelherr, J.M., Oren, R.,
 801 2006b. An evaluation of models for partitioning eddy covariance-measured net ecosystem
 802 exchange into photosynthesis and respiration. *Agric. For. Meteorol.* 141, 2–18.

803 Stoy, P.C., Palmroth, S., Oishi, A.C., Siqueira, M.B.S., Juang, J.-Y., Novick, K.A., Ward, E.J.,
 804 Katul, G.G., Oren, R., 2007. Are ecosystem carbon inputs and outputs coupled at short time
 805 scales? A case study from adjacent pine and hardwood forests using impulse-response
 806 analysis. *Plant, Cell Environ.* 30, 700–710.

807 Stull, R.B., 2012. An introduction to boundary layer meteorology. Springer Science & Business
 808 Media.

809 Sulman, B.N., Roman, D.T., Scanlon, T.M., Wang, L., Novick, K.A., 2016. Comparing methods
 810 for partitioning a decade of carbon dioxide and water vapor fluxes in a temperate forest.
 811 *Agric. For. Meteorol.* 226–227, 229–245.

812 Sus, O., Williams, M., Bernhofer, C., Béziat, P., Buchmann, N., Ceschia, E., Doherty, R.,
 813 Eugster, W., Grünwald, T., Kutsch, W., 2010. A linked carbon cycle and crop
 814 developmental model: Description and evaluation against measurements of carbon fluxes
 815 and carbon stocks at several European agricultural sites. *Agric. Ecosyst. Environ.* 139, 402–
 816 418.

817 Taylor, C.M., Parker, D.J., Harris, P.P., 2007. An observational case study of mesoscale

atmospheric circulations induced by soil moisture. *Geophys. Res. Lett.* 34.

Waldo, S., Chi, J., Pressley, S.N., O’Keeffe, P., Pan, W.L., Brooks, E.S., Huggins, D.R., Stöckle, C.O., Lamb, B.K., 2016. Assessing carbon dynamics at high and low rainfall agricultural sites in the inland Pacific Northwest US using the eddy covariance method. *Agric. For. Meteorol.* 218–219, 25–36.

Wehr, R., Munger, J.W., McManus, J.B., Nelson, D.D., Zahniser, M.S., Davidson, E.A., Wofsy, S.C., Saleska, S.R., 2016. Seasonality of temperate forest photosynthesis and daytime respiration. *Nature* 534, 680–683.

Welch, C.M., Stoy, P.C., Rains, F.A., Johnson, A. V, McGlynn, B.L., 2015. The impacts of mountain pine beetle disturbance on the energy balance of snow during the melt period. *Hydrol. Process.*

West, T.O., Marland, G., 2002. A synthesis of carbon sequestration, carbon emissions, and net carbon flux in agriculture: comparing tillage practices in the United States. *Agriculture, Ecosystems & Environment* 91, 217–232.

Wilson, K., Goldstein, A., Falge, E., Aubinet, M., Baldocchi, D., Berbigier, P., Bernhofer, C., Ceulemans, R., Dolman, H., Field, C., Grelle, A., Ibrom, A., Law, B.E., Kowalski, A., Meyers, T., Moncrieff, J., Monson, R., Oechel, W., Tenhunen, J., Valentini, R., Verma, S., 2002. Energy balance closure at FLUXNET sites. *Agric. For. Meteorol.* 113, 223–243.

Wischmeier, W.H., 1959. A rainfall erosion index for a universal soil-loss equation. *Soil Sci. Soc. Am. J.* 23, 246–249.

Zhang, Y., Liu, C., Yu, Q., Shen, Y., Kendy, E., Kondoh, A., Tang, C., Sun, H., 2004. Energy fluxes and the Priestley–Taylor parameter over winter wheat and maize in the North China Plain. *Hydrol. Process.* 18, 2235–2246.

Figure Legends

Figure 1. Trends in summer fallow area from 1920 until the present in Canada (red dashed line) and the United States (blue solid line) using data from Statistics Canada and the United States Department of Agriculture Economic Research Service.

Figure 2. A map of the winter wheat (WW 2013) and spring wheat (SW 2014) tower location and the fallow (2014) tower location in central Montana, USA. (Image date: 7/25/2014, Google Earth).

Figure 3. The daily maximum canopy height or snow height for the winter wheat and spring wheat rotations measured by a sonic distance sensor (Table 2). Canopy height was assumed to be near zero before winter wheat growth began in earnest in early May, 2013. Stubble left on the winter wheat field in 2013 measured *ca.* 0.2 m, and was assumed to drop to 0 m following the planting of spring wheat on May 2, 2014.

Figure 4. The cumulative (A) and monthly (B) sums of the net ecosystem exchange of carbon in CO_2 (F_c) between the winter wheat, spring wheat, and fallow fields and the atmosphere during the April-September study period. Negative values denote carbon uptake by the biosphere. Error bars represent one standard deviation from the cumulative or monthly sum.

Figure 5. The cumulative (A) and monthly (B) sums of evapotranspiration (ET) from the winter wheat, spring wheat, and fallow study fields during the the April – September study periods. Error bars represent one standard deviation from the cumulative or monthly sum.

Figure 6. The cumulative (A) and monthly (B) sums of sensible heat flux (H) from the winter wheat, spring wheat, and fallow study fields during the the April – September study periods. Error bars represent one standard deviation from the cumulative or monthly sum.

Figure 7. (A) The maximum daily height of the modeled atmospheric boundary layer (h) calculated using surface virtual heat flux measurements from the spring wheat and fallow fields and a one dimensional boundary layer model following Luyssaert et al. (2014) (equation 3) and the height of the lifted condensation level (h_{LCL}) for maximum daily air temperature modeled by lifting air parcels dry adiabatically to the saturation point temperature (equation 5) using meteorological observations from 2014 from the spring wheat and fallow study sites. Daily values were smoothed using a seven-day digital filter. (B) The difference in simulated h among the spring wheat and fallow fields. Dots represent daily maximum modeled h and lines represent the output of a seven day digital filter.

Figure 8. Histogram of times during the April – September 2014 study period when modeled daytime lifted condensation level (h_{LCL}) first exceeds the height of the atmospheric boundary layer (h). ‘None’ and ‘cloudy’ refers to days when daytime h_{LCL} never exceeds h and when h_{LCL} at sunrise was below 150 m, respectively.

Table 1. A list of abbreviations with their definitions.

Abbreviation	Definition
α	Initial slope of the light response curve
α_{PT}	Priestley-Taylor coefficient
β	Gross ecosystem productivity at light saturation
γ	Psychrometric constant
γ^*	Virtual temperature inversion strength
Δh	Change in height of the atmospheric boundary layer
ε_{PT}	Intercept term of the Priestley-Taylor-type model
λ	Latent heat of vaporization
ρ_a	Density of air
ABL	Atmospheric boundary layer
c_p	Specific heat capacity
E_0	Activation energy
ET	Evapotranspiration
F_c	Carbon dioxide flux
G	Soil heat flux
GEP	Gross ecosystem productivity
h	Height of the atmospheric boundary layer
h_{LCL}	Height of the lifting condensation level
H	Sensible heat flux
H_v	Virtual heat flux
LE	Latent heat flux
P	Atmospheric surface pressure
r	Water vapor mixing ratio
RE	Ecosystem respiration
R_n	Net radiation
R_{10}	Ecosystem respiration at an air temperature of 10 °C
s	Slope of the vapor pressure-temperature relationship at saturation
SW_{in}	Incident shortwave radiation
T_a	Air temperature
T_{10}	Air temperature of 10 °C
T_{LCL}	Air temperature at the lifting condensation level
T_t	Reference temperature (227.13 K)
u^*	Friction velocity

Table 2. Variables measured at each site (see Table 1) along with sensor type. W: Wheat (winter wheat in 2013, spring wheat in 2014), F: Fallow (2014).

Measurement	Sensor(s)	Site
SW_{in}	NR01 net radiometer	W, F
R_n	NR01 net radiometer	W, F
Canopy height	SR50 sonic distance sensor	W
T_a and relative humidity	HMP-50 temperature/relative humidity probe	W, F
G	HFP01 heat flux plate	W, F
H	CSAT-3	W, F
LE	CSAT-3 and LI-7200	W, F
F_c	CSAT-3 and LI-7200	W, F

Table 3: The percent of available turbulent flux data during the entire April-September study period for each study site. Daytime refers to periods when the solar zenith angle was less than 90°.

Site	Period	F_c	LE	H
Winter wheat (2013)	Daytime	52%	52%	53%
	Total	48%	49%	49%
Spring wheat (2014)	Daytime	48%	48%	48%
	Total	45%	45%	45%
Fallow (2014)	Daytime	44%	50%	51%
	Total	42%	49%	50%