**Agricultural land use and macroinvertebrate assemblages in lowland temporary streams of
the Willamette Valley, Oregon, USA**

William J. Gerth*, Judith Li, and Guillermo R. Giannico

Department of Fisheries and Wildlife, Oregon State University, Corvallis, Oregon 97331 USA

*Corresponding author. E-mail address: william.gerth@oregonstate.edu

ABSTRACT

Streams that dry during part of the year are common throughout the world, yet studies of the macroinvertebrate assemblages in these types of streams are rare compared to those in permanent streams; and studies that assess the effects of agriculture on temporary stream invertebrates are even rarer. We studied macroinvertebrate assemblages in lowland temporary streams of a region with high agricultural land use, the southern Willamette Valley, Oregon, USA. Overall assemblages were dominated by non-insects, and invertebrates tolerant of organic pollution. Nonetheless, these invertebrates displayed adaptations to life in temporary habitats, and as such they may be unique to temporary streams and seasonal wetlands, providing an important addition to regional biodiversity. Stream invertebrates are also important as a prey base for native fish and amphibians using these channels. Benthic invertebrate densities were higher at sites with slower water and more in-stream vegetation; to a lesser degree greater agricultural land use was associated with lower densities. Taxon richness was also negatively affected by agriculture, but this was most evident when least-disturbed and highly agricultural sites were compared. Sites in watersheds with a lower proportion of their area under agriculture (mostly west of the Willamette River) had a variety of taxa in disturbance-sensitive insect orders Ephemeroptera, Plecoptera, and Trichoptera (EPT), plus flies in the family Simuliidae present. In addition, they had greater relative abundances of 2 types of flies in the family Chironomidae. In contrast, sites in watersheds with high agricultural land use (mainly east of the Willamette River) had greater relative abundances of non-insects, including ostracods, nematodes, and oligochaete worms. In highly agricultural watersheds, when stream-bottom vegetation was abundant, it was associated with greater benthic invertebrate density, but not with higher taxon richness. Our results suggest that increasing stream-bottom vegetation could be useful when food is limiting for native vertebrates. On the other hand, reduced agricultural land use allows for the development of more diverse benthic invertebrate assemblages.

Key words: temporary streams, agricultural streams, macroinvertebrate communities, agriculture, land use, conservation, Oregon, United States.

1. Introduction

Streams that dry during part of a year are common throughout the world and naturally occur where rainfall is limited or highly seasonal, or where porous geology allows for the loss of surface flow (Dieterich and Anderson 1995, Paltridge et al. 1997, Uys and O'Keefe 1997, Meyer and Meyer 2000, Aguiar et al. 2002, Datry et al. 2014, Mazor et al. 2014). These types of streams can vary in the length and predictability of their dry phase. Some, but not all, have permanently flowing headwaters, and they may or may not be used by fish during the wet phase. The terms ephemeral, temporary, and intermittent have been applied variously to streams that do not flow continuously. Our study focused on temporary streams, defined by Dieterich and Anderson (2000) as streams without permanent headwaters, but with flow persisting for >4 consecutive months per year. Based on the changing conditions in these kinds of systems, the annual cycle can be divided into 3 hydrologic stages: a running water stage with flowing water, a pool phase with standing water, and a terrestrial stage with surface water absent (Williams and Hynes 1976, Williams 1996). Our studies were conducted towards the end of the running water stage during late winter/early spring in temporary streams draining grass seed producing lands of the Willamette Valley, Oregon, U.S.A.

Previous studies of temporary streams in the Pacific Northwest region of North America have examined fishless sites within small drainage areas in relatively steep, forested terrain (Tew 1971, Dieterich and Anderson 2000, Price et al. 2003, Banks et al. 2007, Stewart and Anderson 2008). In contrast, our study examined lowland temporary streams in a region with high agricultural land use. These streams supported native fish and amphibians during the wet season, serving as nursery grounds for some species (Colvin 2006, Colvin et al. 2009). As invertebrates are essential prey for these and other vertebrate predators, invertebrate abundance or biomass are considered an important component of habitat suitability for vertebrates (Wipfli and Gregovich 2002, Taft and Haig 2005) in these temporary habitats.

Although temporary streams are less well studied than permanent streams, several patterns have been documented. These systems often have fewer benthic invertebrate taxa during their running phase than permanent streams (Williams 1996, Williams et al. 2003, Storey and Quinn 2008, Sanchez-Montoya

et al. 2010), and taxon richness decreases as the duration of the dry period increases (Meyer and Meyer 2000, Williams 2006, Datry et al. 2014). In addition, invertebrate assemblages in streams that do not flow continuously can be quite distinct from those in permanent streams (Dieterich and Anderson 2000, Meyer and Meyer 2000, Williams 2006, Storey and Quinn 2008) with many taxa in temporary streams exhibiting traits that allow them to survive or avoid the loss of surface water (Tew 1971, Dieterich and Anderson 1995, Jacobi and Cary 1996, Williams et al. 2003). In other cases, minimal differences in richness and assemblage composition between these types of streams have been found, especially when non-permanent channels being studied do not dry in all years, or where seeps, pools, or moist stream substrates remain throughout the dry period (Del Rosario and Resh 2000, Price et al. 2003, Banks et al. 2007).

Agricultural land use has often led to degraded riparian zones, and increased water temperatures, nutrients, pesticides, and fine sediments in permanent streams (Cuffney et al. 2000, Waite and Carpenter 2000, Anderson et al. 2003, Stone et al. 2005), but these effects are less well documented for temporary streams. In addition, in agricultural zones the hydrology tends to be altered as watercourses are often channelized, ditches are dug and, sometimes, sub-surface drainage is installed to move water quickly off the land (Boag 1992, Armitage et al. 2003, Williams et al. 2003, Stone et al. 2005). Physico-chemical conditions associated with agriculture can be accompanied by reduced aquatic macroinvertebrate taxonomic richness, increased proportional abundances of tolerant invertebrates, and/or other indications of biological impairment (Rothrock et al. 1998, Cuffney et al. 2000, Anderson et al. 2003, Stone et al. 2005). However, impact levels may vary with the types of crops (e.g., annual row crops vs. permanent or orchard crops) or animals being reared and the intensity of production.

Effects of emergent or submerged vegetation in temporary streams are also poorly understood. Tew (1971) found in one temporary stream that the highest invertebrate density, and the greatest taxonomic richness and proportional abundance of non-insects were associated with the channel segments that had grass growing in them. This coincides with what others have reported for permanent streams, where habitats with aquatic plants that provide structural complexity and potential refuge during high

flows had different assemblage composition and higher macroinvertebrate richness and density than habitats without vegetation (Cogerino et al. 1995, Armitage et al. 2001, Armitage et al. 2003).

This study examined macroinvertebrate assemblages within a network of temporary streams flowing through agricultural lands in the southern Willamette Valley where annual and perennial grasses were the major crops. The objectives of our research were: 1) to characterize physico-chemical conditions and watershed land use, across a broad range of sites in lowland, temporary streams; 2) to examine macroinvertebrate density, taxon richness and assemblage composition along a gradient from intense agricultural use to minimally disturbed sites; and, 3) to determine the effect of stream-bottom vegetation on macroinvertebrate density and richness in streams draining agricultural lands.

2. Materials and Methods

2.1. Study location

This study was conducted in the Willamette Valley, Oregon, USA, a lowland area lying between the Coast Range and the Cascade Mountains where sediments from the Missoula floods were deposited between 15,000 and 13,000 years ago towards the end of the last ice age (Boag 1992). The broad valley bottom has very little slope with predominantly poorly drained, clayey soils (Hulse et al. 2002, Wigington et al. 2005). The climate is characterized by cool, wet winters and warm dry summers; most of the approximately 1,100mm of annual precipitation occurs as rain. More than 95% of the precipitation falls from September through June, with about 50% in December through February (Taylor and Bartlett 1993, Wigington et al. 2003). As a result, temporary streams in this region dry during summer and only resume flowing in October or November after the rains return. Temporary and permanent streams of the southern valley drain land that has been converted largely for agricultural use; a variety of crops are grown, but much of the land is used for grass seed production and pasture (Jackson 1993, Klock et al. 2002, Wyss et al. 2013). According to historical accounts of Euro-American settlers in the mid-19th century, the valley was swampy during the wet season, and streams meandered and spread across the valley floor without definite channels (Boag 1992, Taft and Haig 2003). Since the late 19th century, however, the land has

been ditched and tiled to facilitate drainage, altering the hydrology and habitat available in present-day streams. Despite alterations, stream networks expand considerably and predictably during the wetseason (Wigington et al. 2005) and provide seasonal habitats for aquatic fauna (Colvin et al 2009).

2.2. Field sampling and data collection

Sample and data collection took place in 3 stages for this study. During the 2002-2003 wetseason, sites in five sub-basins of the southern Willamette Valley were sampled in conjunction with a related fish and amphibian survey (Colvin 2006, Colvin et al. 2009). Because leastdisturbed sites (< 55% of watershed under agriculture) were not well represented in this initial survey, in 2008 we sampled hand-picked leastdisturbed sites and resampled some of the same sites sampled in 2002-2003 within highlyagricultural watersheds (> 85% of area under agriculture) for comparison. In addition, in spring 2006 we sampled paired sites within streams in highly agricultural watersheds to test for the local effects of stream-bottom vegetation.

Southern valley survey, and leastdisturbed vs highlyagricultural sites.—For the southern valley survey (2002-2003), we used observations from road crossings and aerial photographs to identify a group of 59 potential sampling sites on streams likely to be dry during summer. From these, we chose 18 sampling sites in independent drainages that met the following criteria: 1) channels were dry during a scouting visit in the month of August prior to sampling, 2) watershed area sufficient to maintainwater in the channels throughout the wetseason (6–9 months), 3) watersheds with little urban influence and primarily draining the flattest part of the valley (Prairie Terrace level IV ecoregion) and 4) landowners granted permission for sampling (Fig. 1A).

In 2008, to identify macroinvertebrate communities likely to occur under the best conditions in these types of streams, we collected additional invertebratesamples at 5 leastdisturbed sites and repeated sampling in 5 of the most agriculturallyinfluenced sites from the previous survey for comparison. The leastdisturbed sites had riparian vegetation and three of them were in nature preserves or park/natural areas; one stream running through private property had active wetland mitigation just beyond the riparian

zone at the sampling site. Although not a criterion for selection, all leastdisturbed sites had evidence of beaver (*Castor canadensis*) activity nearby. The 5 highlyagricultural stream sites had variable levels of riparian vegetation development (e.g., from no vegetation to corridors of shrubs or forest), and no evidence of beaver activity.

At each site, a 100m long study section was established. In 2002-2003, water quality grab samples were collected every 2 to 3 weeks throughout the wetseason (mid-December through early May; 7 sample dates per site). Physical habitat surveys were conducted once at each site in late April–early May 2003. Wetted widths were measured at 6 equally spaced transects perpendicular to the direction of flow in each study section. At 3 equally spaced points across these transects, depth and velocity were measured, and substrate was categorized as vegetated or non-vegetated. At non-vegetated sample points, the substrate particle size was noted. No physical habitat data or water quality samples were collected in association with samples taken in 2008.

For both years, benthic invertebrate samples consisting of 6 randomlylocated Surber samples (500 μ m mesh netting; sampling area = 0.09 m² each) were collected once in late March–early April, when streams were flowing, but discharge was less variable than in winter. At each collection point, stream-bottom substrate and any rooted vegetation in the sampling frame was thoroughly disturbed and water flow pushed dislodged invertebrates and stream-bottom material into the collection net. Samples collected in this manner were preserved with 95% ethanol and were held for processing in the lab.

Effects of stream-bottom vegetation in highlyagricultural sites.—To test the effect of stream-bottom vegetation on benthic invertebrates, adjacentpairs of sites with and without extensive cover of stream-bottom vegetation (henceforward called “vegetated” and “non-vegetated” sites) were picked in 4 streams of the Calapooia River sub-basin east of the Willamette River (Fig. 1B). All sampling sites had >95% of their watersheds under agricultural use. Paired sites were approximately 200–500m apart and within the same stream.

Each site was sampled once for invertebrates between the end of February and the beginning of April 2006. Paired sites were always sampled on the same day. Macroinvertebrate sampling was

conducted in the same way as in other components of the study. In late April 2006 physical habitat surveys were conducted, but measurements were made in greater detail than those in 2003. Wetted widths were measured at 11 equally spaced transects perpendicular to the direction of flow. At 5 equally spaced points on these transects, depth and velocity were measured and substrate was characterized. Water quality grab samples were collected once at the time of habitat surveys.

2.3. Sample processing and data compilation

Water quality grab samples were collected and transported in coolers to the USDA Agricultural Research Lab on the Oregon State University campus in Corvallis, Oregon, to be analyzed for pH, conductivity, suspended sediment, nitrate-N, and orthophosphate-P. Conductivity and pH were determined using meters (Hydrolab Quanta multi-parameter water quality sensor and Orion model 811 pH meter). Suspended sediment concentration was determined by weighing the dried filtrate from a known volume of sample water. Nitrate concentration was measured by flow injection analysis (Wendt 1999) and orthophosphate concentration was determined by flow injection analysis colorimetry (Prokopy 1994). For the 2002-2003 southern valley survey, data were averaged across sample dates to determine the average water quality to which benthic invertebrates were exposed during the wetseason. For the 2006 paired sites, water quality samples were only available from the end of April, so wet-seasonlong averages could not be calculated.

Wetted width, depths, and velocities measured in physical habitat surveys were averaged by site for both the southern valley survey and the pairedsite stream-bottom vegetation test. Percent vegetated stream bottom was calculated by counting the number of points on transects where vegetation was present, then dividing by the total number of points where observations were made.

All sample site coordinates were entered into a geographic information system (GIS) (ArcGIS 8.1, Esri, Redlands, CA). Areas draining to sites were delineated using hydrologic modeling with 10m digital elevation models (DEMs)(Oregon Geospatial Enterprise Office) and site coordinates as pour points. Watershed areas and slopes (from the highest points in watersheds to sample sites) were calculated

as well as stream distances from the sampling sites to downstream permanent water. Watershed land use was determined by clipping delineated watersheds from a Willamette Valley land use/land cover GIS layer (Klock et al. 2002). Land use categories provided in the land use/land cover layer were combined and simplified to “Agriculture” (primarily perennial grass but also including annual grass, orchard, berry, and row crop fields), “Forest”, “Urban”, and “Other”, and the proportion of each watershed in agricultural use (% Watershed Agriculture) was calculated.

The six benthic invertebrate samples collected at each site were combined to form a site composite sample. Composite samples were subsampled using a gridded sieve (Caton 1991) with a fixed count protocol (sensu Larsen and Herlihy 1998) and a target of identifying and enumerating 500 organisms per site. This was accomplished by taking random subsamples from each composite sample until at least 500 organisms were picked or the entire sample was processed. Aquatic invertebrates were identified to the lowest practical taxonomic resolution using dichotomous keys and taxonomic descriptions (Merritt and Cummins 1996, Wiggins 1996, Smith 2001, Thorp and Covich 2001, Stewart and Stark 2002, Santos-Flores and Dodson 2003). Aquatic insects were identified to genus with the exception of Simuliidae that were left at family, and Chironomidae and Ceratopogonidae that were identified to subfamily or tribe. Non-insects were identified to the finest practical level, varying from phylum (e.g., for nematodes) to genus (e.g., for amphipods and isopods). Samples were processed in the years in which they were collected, but all identifications were performed by the first author to maintain consistency among sample sets. For each site, invertebrate count data were converted to densities by dividing by the proportion of the composite sample examined and, then, dividing by the area of stream bottom sampled.

2.4. Statistical analyses

Southern valley survey.—Invertebrate summary metrics related to whole assemblages (total density, total taxon richness) and specific types of taxa (non-insect invertebrates and insects in the disturbance-sensitive orders Ephemeroptera, Plecoptera, and Trichoptera (EPT): % non-insects, %EPT,

and EPT taxon richness) were calculated. In addition, we calculated values of the Hilsenhoff Biotic Index (HBI) modified for use with taxa found in Oregon aquatic environments (OWEB 2001). HBI values are assemblage level, abundance weighted averages of taxon specific water quality tolerance values. HBI values range from 0 to 10, with low values indicating assemblages dominated by clean-water taxa intolerant of organic enrichment, and high values indicating greater assemblage pollution tolerance.

Relationships between 12 environmental variables and invertebrate density and total taxon richness were assessed with multiple regressions. The environmental variables used for these analyses were wetted width, channel depth, water velocity, % vegetated substrate, suspended sediment, nitrate-N, orthophosphate-P, conductivity, distance to downstream permanent water, watershed area, watershed slope, and watershed % agriculture. Prior to regression, variables were transformed appropriately. All possible 1, 2 and 3 variable models were run and models were ranked by Akaike's Information Criteria modified for use with small sample sizes (AIC_c) (Burnham and Anderson 2002). Top models were identified as all models within 2 AIC_c units of the model with the lowest AIC_c value.

Variation in assemblage composition was explored using non-metric multidimensional scaling (NMS) ordination with Bray-Curtis dissimilarity measures (PC-ORD, version 4.41, MjM Software, Gleneden Beach, Oregon). Prior to ordination, invertebrate data were $\log(x+1)$ transformed to reduce the influence of dominant taxa on the analysis. Subsequently, data were standardized by site totals to eliminate the influence of differences in total invertebrate density among sites. Dimensionality of the ordination solution was determined using the scree test (Kruskal and Wish 1978) to assess the point at which additional axes provided little improvement in fit (stress reduction). Site ordination coordinates were correlated with the same environmental variables used in the multiple regression analyses. Correlations between ordination coordinates and invertebrate variables (summary metrics and transformed taxa values in the ordination matrix) were also run. Directions and strengths of correlations for the most highly correlated environmental variables and invertebrate summary metrics ($r \geq 0.45$) were shown as vectors overlaid on the ordination graph.

Leastdisturbed vs. highlyagricultural sites.—Invertebrate metric values for leastdisturbed and highlyagricultural sites were compared statistically using t-tests after appropriate transformations. Invertebrate data from this part of the study were also combined with data from the southern valley survey to give context to the previous information. Invertebrate metric medians and ranges were compared for the least disturbed, highlyagricultural and southern valley survey sites. Assemblage data from sites sampled in 2003 and 2008 were combined in one matrix, $\log(x+1)$ transformed, and standardized by site totals; assemblage composition patterns were examined using NMS ordination with Bray-Curtis dissimilarity measures.

Effects of stream-bottom vegetation in highlyagricultural watersheds.—Site physico-chemical and watershed characteristics were summarized and compared for the paired sites with and without large amounts of rooted stream-bottom vegetation. Invertebrate density and taxon richness were the invertebrate metrics used to test for effects of stream-bottom vegetation. Paired t-tests were run to test for all differences.

3. Results

3.1 Southern valley survey

Site characteristics.—Median watershed area for the sites was 8.8km²;the largest was almost 22 km² (Table1). Sites were generally several kilometers upstream of permanent water (median distance = 2.1 km), but the furthest one was6.5 stream kilometers away from where water was present during summer. Agriculture was the major land use in allwatersheds, ranging from 55% to 98% of watershed land. Perennial grass was the highest agricultural land cover in all watersheds, but the site with the lowest proportion of its watershed in agriculture also had about a third of its agricultural land in orchard/berry fields. Row crop fields were rare; they represented only 5% of the land area in the watershed where they were most abundant. Watersheds were shallowly sloping, reflecting the regional topography.

Though sitespecific physical measurements were made in late April-early May, we observed highly variable channel depth, wetted width, and water velocity throughout the wetseason depending on

rainfall. In general, these measures were highest between mid-November and late January. After February they were lower and less variable. An example of this seasonal pattern is illustrated by the hydrograph for one of the southern valley survey sites from a year in which invertebrates were not collected (2003-2004; Fig. 2). In late April–early May 2003, about a month after invertebrate sampling, median site channel depth and width were 0.30m and 3.0m respectively (Table 1). Median water velocity was 0.14m/s, but site velocities ranged over an order of magnitude.

Generally, nitrate concentrations were high early in the wetseason and decreased through time. Five of the 18 sites had at least one reading of nitrate-N concentration above the USEPA drinking water standard of 10 mg/L. When nitrate concentrations were averaged over the wetseason, median concentration for the sites was 4.1 mg/L, but the highest nitrate concentration was almost 10 times the lowest (Table 1). Other water quality measures were also variable through time, but did not have clear seasonal trends. Averaged over the wetseason, phosphate concentrations at sites ranged over approximately one order of magnitude. Conductivity and suspended sediment were less variable among sites.

Stream bottoms at all sites were predominantly hard, compressed clay. However, sites varied in the amount of rooted, stream-bottom vegetation. Plants growing in the channels included bentgrass (*Agrostis* sp.), reed canarygrass (*Phalaris arundinacea* L.), spike rush (*Eleocharis* sp.), water plantain (*Alisma* sp.) and cattail (*Typha* sp.).

Benthic invertebrates.—We collected a median of 18 benthic taxa per site, and a total of 49 macroinvertebrate taxa valley-wide. Non-insects collected were adults and juveniles since both are aquatic and these invertebrates don't have active, terrestrial lifestages. In contrast, most insects collected were juvenile life stages, since juvenile aquatic insects in most orders (Diptera, Ephemeroptera, Plecoptera, Trichoptera, Megaloptera, and Odonata) are the ones found in water, and their winged adults are terrestrial/aerial. The exceptions among the aquatic insects are beetles and true bugs (Orders Coleoptera and Hemiptera respectively) that have aquatic juveniles; however, their winged adult forms can also be found in aquatic habitats. Median macroinvertebrate density was 2923 invertebrates per square

meter, but individual site densities varied by almost 2 orders of magnitude (Table 2). These temporary stream sites had low total taxonrichness (median = 18), generally dominated by non-insects (median = 88.7%, Table 2). Oligochaete and nematode worms were collected at all sites; ostracods were collected at 17 of the 18 sites (Table 3). Free-living flatworms (Turbellaria), 2 subfamilies/tribes of chironomids, and macrocrustaceans (isopods and amphipods) were found at over 80% of sites. Modified Hilsenhoff Biotic Index (HBI) scores (OWEB 2001) between 5.5 and 7.0 indicated that these invertebrate assemblages were dominated by organisms tolerant of organic pollution and low dissolved oxygen (Table 2).

Insects in the disturbance-sensitive orders, Ephemeroptera, Plecoptera, and Trichoptera (EPT), were uncommon. Though EPT taxonrichness ranged from 0 to 6, the median was only 1. Despite being collected at over 70% of the sites, *Limnephilus* spp. caddisflies were generally not abundant within sites. Even though individuals were not abundant and identifications were left at the genus level for analyses, four obvious *Limnephilus* morpho-species (based on case form and head capsule coloration) were present in the collections, indicating additional diversity within this genus.

Regression models.—Variation in benthic invertebrate density could be explained by 4 approximately equally good multiple regression models (within 2 ΔAIC_c units of one another). Water velocity and % vegetated substrate were each in three top models. The percent of watershed in agricultural land use (abbreviated to watershed % agriculture in the model) was represented in 2 top models. The equation for the model with all of these variables was

$$\begin{aligned} \text{Log density} = & 5.23 - (4.29 \times \text{velocity}) + (1.01 \times \arcsin \sqrt{\% \text{vegetated substrate}}) \\ & - (1.95 \times \arcsin \sqrt{\text{watershed \% agriculture}}). \end{aligned}$$

None of the independent variables in this model were highly correlated with the others (all correlation coefficients: $|r| \leq 0.35$) so multicollinearity should not be a major concern and interpretation of the signs and magnitudes of the regression coefficients is straightforward (Graham 2003). Thus, this model

indicated that sites with slower water velocities, more vegetation growing in the channel, and less agriculture in the watershed had greater densities of benthic macroinvertebrates (adjusted $R^2 = 0.56$; $p < 0.01$). As individual explanatory variables, velocity and %vegetated substrate explained 34% and 32% of the variation in invertebrate density respectively; watershed %agriculture explained 7%.

Variation in taxon richness was less satisfactorily explained by the measured physico-chemical and watershed variables. Eight regression models were about equivalent (within 2 ΔAIC_c units of one another), but the best of these only explained 6% of the variation in richness ($p = 0.17$). Taxonomic richness showed little relationship to agricultural landuse ($r = -0.20$; $p = 0.42$), or watershed area ($r = -0.26$; $p = 0.31$), which we thought could be a proxy for the duration of stream flow. The site with the lowest richness (9 taxa) had 73% of its watershed in agricultural land use, which was below the median agricultural land use for sites in the southern valley survey (Table 1). If this site was considered an outlier and was removed from the analysis, there was a stronger relationship between richness and watershed % agriculture ($r = -0.35$; $p = 0.07$), but not between richness and watershed area ($r = -0.19$; $p = 0.29$).

Community analysis.—Benthic assemblage composition gradients were best represented by a three-dimensional NMS ordination. This solution accounted for 87% of the information in the original site dissimilarity matrix with a stress of 10.0. Axes 1 and 2 explained the majority of the dissimilarity, accounting for 45% and 28% respectively (Fig. 3A).

Axis 1 represented a gradient in assemblages from those with greater relative abundances of non-insects, including ostracods, nematodes, and oligochaete worms, to those with greater numbers of total and EPT taxa, presence of Simuliidae, and greater relative abundances of 2 types of chironomids (Tanypodinae and Tanytarsini). This assemblage gradient was associated with decreasing watershed %agriculture and conductivity. Axis 1 also displayed a geographic component; most of the sites from west of the Willamette River (to the right in the ordination plot) had watersheds with somewhat less agriculture. These sites had more diverse invertebrate assemblages than sites from east of the Willamette (to the left in the ordination).

Local site conditions including stream-bottom vegetation, water velocity, and channel width were correlated with the assemblage composition gradient along Axis 2 (Fig. 3A). At sites with slower water, more vegetation growing on stream bottoms, and wider wetted channels (higher Axis 2 scores) there were higher total benthic densities and greater proportions of cyclopoid copepods, *Crangonyx* amphipods, *Caecidotea* isopods, and mites. In the opposite direction, sites with lower Axis 2 scores had higher water velocities, narrower channels, and less stream-bottom vegetation. These sites had lower benthic invertebrate densities but higher relative abundances of Orthocladiinae chironomids and *Baetis* mayflies.

Though Axis 3 also indicated a gradient in invertebrate assemblage composition, it explained little (11%) of assemblage dissimilarity, and was moderately correlated with suspended sediment ($r = 0.48$).

3.2. Leastdisturbed vs. highlyagricultural sites

Leastdisturbed and highlyagricultural sites sampled in 2008 were compared using the 6 invertebrate metrics listed in Table 2. Leastdisturbed sites had significantly greater total and EPT taxon richness (both $p < 0.01$), higher proportional abundances of EPT individuals (%EPT; $p < 0.01$) and lower proportional abundances of non-insect individuals (%non-insect; $p < 0.01$) than highlyagricultural sites. However, leastdisturbed and highlyagricultural sites did not differ in invertebrate density ($p = 0.61$), or in HBI score ($p = 0.13$).

Leastdisturbed and highlyagricultural sites had several metric values that were at opposite ends of the ranges for sites from the 2003 southern valley survey. Leastdisturbed sites had some of the highest values for %EPT and taxon (total and EPT) richness, and lowest values for %non-insects (Table 2). The additional sampling in 2008 added 4 taxa to the valley-wide taxa list—3 only from leastdisturbed sites (a mayfly, stonefly, and caddisfly) and 1 from leastdisturbed and highlyagricultural sites (a beetle).

HBI median values ranged from 6.0 at leastdisturbed sites to 6.2 and 6.4 (on a scale of 0 to 10) at other sites (Table 2). These values indicated assemblages at all sites regardless of land use were dominated by organisms tolerant of organic pollution and low dissolved oxygen (OWEB, 2001). The fact

that invertebrate densities did not differ between leastdisturbed and highlyagricultural sites indicates that the weak relationship between invertebrate density and watershed agricultural land use seen with the southern valley survey data may be spurious. Additionally even though leastdisturbed sites had higher %EPT than other sites, they were still dominated by other types of organisms. None of the leastdisturbed sites had $\geq 25\%$ EPT individuals and only 2 sites had $> 10\%$ EPT individuals.

A three-dimensional NMS ordination combining data from the 2003 southern valley survey and the 2008 samples (Fig. 3B) revealed assemblage gradients and correlations with invertebrate metrics that were largely the same as those seen in the singleyear ordination (Fig. 3A). This combined solution accounted for 91% of the information in the original site dissimilarity matrix with a stress of 10.5. Axis 1 explained 62% of the dissimilarity, and Axes 2 and 3 explained 15% and 14% respectively. Assemblages at 2008 leastdisturbed and highlyagricultural sites were distinct from one another at opposite ends of Axis 1, while overlapping with assemblages collected in the more extensive valley-wide survey. Again there was a geographic component to this ordination because we only found leastdisturbed sites west of the Willamette River and the highlyagricultural sites were east of the Willamette.

3.3. Effects of stream-bottom vegetation in highlyagricultural sites

In the 2006 pairedsite comparison, rooted vegetation covered 60% to 87% of benthic substrates at vegetated sites, whereas non-vegetated sites had 0% to 7% vegetation cover. On average, sites with vegetated substrates were 10cm shallower than corresponding non-vegetated sites within the same stream (paired t-test, $p < 0.01$), but paired sites did not differ in width ($p = 0.30$) or average velocity ($p = 0.78$). Average wetted widths for these sites ranged from 2 to 4.5 meters and average water velocity ranged from 0.04 to 0.12 m/s. Because paired sites were close together within streams, water quality and watershed land use differences were minor. All watersheds had $>95\%$ agricultural land use.

For three streams, benthic invertebrate densities werecomparatively low at non-vegetated sites (~200 to 1500 individuals per square meter) and 6–45X higher at vegetated sites (Fig. 4). These results supported findings in the southern valley survey, that local stream-bottom vegetation can be one of the

primary factors affecting benthic invertebrate density. However, in a fourth stream, where the benthic invertebrate density at the non-vegetated site was more than an order of magnitude higher than in the other streams, there was little difference between vegetated and non-vegetated sites. Consequently, density differences associated with stream bottom vegetation were only marginally significant (paired t-test; $p = 0.08$). At these pairs of highly agricultural sites, we collected a median of 11 taxa per site. As in the 2003 survey, stream-bottom vegetation did not have a significant effect on taxon richness (paired t-test, $p = 0.18$).

4. Discussion

Our extensive valley-wide survey in combination with subsequent comparisons of least disturbed and highly agricultural sites and sites with and without stream-bottom vegetation revealed a suite of invertebrate taxa adapted to the seasonal nature of Willamette Valley lowland temporary streams. Many of the taxa we found in our study sites were also found in stillwater southern Willamette Valley seasonal wetlands (Wyss et al. 2013). Most of these taxa have resting eggs or other desiccation resistant life stages that allow them to remain dormant in place during the dry season (Williams 2006). In addition, the aquatic insects have winged adults which allow them to colonize temporary habitats overland from nearby permanent or temporary aquatic habitats. Non-insect invertebrates, especially nematodes, oligochaetes, various crustaceans and flatworms, dominated benthic assemblages at our study sites, but in certain locations aquatic insects including those in the family Simuliidae and several EPT taxa could also be found. These invertebrates were affected by the physico-chemical conditions imposed by agricultural land use and by local stream-bottom vegetation. Though previous studies of invertebrates in non-permanent streams in agricultural landscapes examined relationships between invertebrates, local land use and physico-chemical and riparian characteristics (Aguiar et al. 2002, Davis et al. 2003, Stone et al. 2005), the broader view of our study illustrated the importance of testing for effects at both local and watershed scales.

Not surprisingly, we found that taxonomic richness in our temporary streams was considerably lower than at permanent streams in forested watersheds of western Oregon (Herlihy et al., 2005), and somewhat lower than at permanent Willamette Valley lowland streams sampled during summer (Li et al. 2001). Based on the literature, we also expected that taxonrichness would be related to the duration of the presence of surface water in the temporary channels we studied (Williams 1996, Meyer and Meyer 2000). Although we did not directly measure the duration of water presence, we expected that watershed size could be used as a surrogate for duration of the wet phase. Contrary to expectations, we did not find any relationship between taxon richness and watershed area. The temporal measure of wet phase duration may require a more local and empirical measure to detect its influence on invertebrate composition.

One of the factors that did appear to influence invertebrate taxonomic richness and assemblage composition was watershed agricultural land use. The relationship between the proportion of the watershed under agricultural use and macroinvertebrate assemblage composition was apparent in the valley-wide survey and with the addition of assemblage information from leastdisturbed and highlyagricultural sites. On the other hand, the relationship between taxonomic richness and agriculture was not apparent across the limited range in the proportions of watershed area under agriculture in our southern valley survey, but became obvious when leastdisturbed sites were compared with highlyagricultural sites. Because agriculture can lead to various types of habitat alteration and water quality changes (Cuffney et al. 2000, Waite and Carpenter 2000, Anderson et al. 2003, Stone et al. 2005), and the watersheds we studied had multiple landowners with varying farming practices, we were unable to narrow down particular practices or water quality parameters that had the greatest effects on aquatic invertebrates. However, identifying specific factors associated with changes in invertebrate richness and assemblage compositionand determining best management practices could be a fruitful area for future research.

We did noticethat the invertebrate gradients we found that were related to agriculture also had a geographic component. The flat part of the valley (Prairie Terrace level IV ecoregion) that is more heavily used for agriculture tends to be narrower west of the Willamette River; study sites on that side of

the Willamette are closer to the Valley Foothills, which have a greater proportion of their lands under forest cover (level IV ecoregion in the level III Willamette Valley ecoregion). Thus, compared to sites in the wider, flat part of the valley east of the Willamette, western sites were generally characterized by having a smaller proportion of the watershed area under agriculture, greater stream invertebrate taxonomic richness, and assemblages less dominated by non-insects. Such geographic differences in land use and invertebrate assemblage patterns have been noted in other studies at landscape scales (e.g., Aguiar et al. 2002).

Temporary stream sites were all characterized by invertebrate assemblages dominated by organisms considered highly tolerant of organic pollution and low dissolved oxygen (Modified HBI scores > 5.0; OWEB, 2001). Given the high degree of agricultural land use in many of the watersheds, it was not surprising that pollution-tolerant organisms would dominate invertebrate assemblages at most sites. However, even the invertebrate assemblages in least disturbed sites had high modified HBI scores. These high scores at least disturbed sites suggest that adaptations that enable invertebrates to complete their life cycles in temporary streams (e.g., dormant life stages, fast development, flexible life cycles, and/or good dispersal ability; Williams 2006) could be similar to traits that allow them to inhabit streams with seasonally poor water quality. As a result, it is likely that bioassessment criteria developed for permanent streams will give misleading results for lowland temporary streams and, therefore, new criteria should be developed. The one study that contradicts this notion that temporary and permanent streams need different bioassessment criteria is from an arid region in southern California where perennial streams are rare and both temporary and perennial streams draw from the same regional pool of taxa adapted to intermittent flow (Mazor et al. 2014). The authors of this study, however, admit that their results are not likely to be applicable to wetter regions (such as Oregon) where richer regional faunas are more likely to allow the development of distinctly different assemblages in temporary and perennial streams.

Another way to look at variation in invertebrate assemblage composition across sites with differing amounts of agricultural land use is to consider the biological traits of taxonomic groups. In particular, in our study, we noted that relative abundance of non-insects was greatest in sites with high

watershed agricultural land use; conversely, these sites had low abundance of aquatic insects. which were found in greater abundances in less disturbed sites. Two trait differences between aquatic insects and non-insects could account for this pattern. First, although relatively uniform, high modified HBI scores at all sampling sites indicate that invertebrate assemblages had high pollution tolerance, the average taxon specific HBI values differed between temporary stream insect and non-insect taxa. The average tolerance values for the aquatic taxa were 4.4 for insects and 6.8 for non-insects. This means, that as a group, aquatic insects are less likely to occur in temporary streams with degraded water quality than non-insects (OWEB 2001). In addition, as indicated before, aquatic insects have winged adults, so they have greater dispersal potential than non-insects. Therefore, in watersheds with less disturbances, there may be greater source populations of aquatic insects accounting for higher colonization rates in their temporary streams.

Despite the apparent overall disturbance tolerance of invertebrate assemblages in temporary streams like the ones in this study, lowland temporary streams of the Willamette Valley are also likely important for sustaining certain specialist taxa. We collected several EPT genera (the stoneflies *Ostrocerca* and *Podmosta*, the mayflies *Ameletus* and *Anafroptilum*-formerly *Centroptilum* (Jacobus and Wiersma 2014)-and the caddisflies in the genus *Limnephilus*) known to have species previously found in temporary streams of western Oregon (Dieterich and Anderson 2000, Banks et al 2007, Stewart and Anderson 2010). Endowed with specialized adaptations to withstand or avoid the dry phase of temporary streams, these species are considered to be more abundant in, if not exclusive to, temporary habitats. In addition, we collected a little known anomopod cladoceran, *Dumontia oregonensis*, which is also known from seasonal wetlands in the Willamette Valley (Wyss et al. 2013), but was previously only reported from a few vernal pools in southern Oregon and central California (Santos-Flores and Dodson 2003, Van Damme and Dumont 2008). Our field sampling and laboratory subsampling procedures, and level of taxonomic identification did not allow us to develop an exhaustive list of all taxa, so there may be additional taxa (particularly species) unique to these temporary streams that were not documented in our study.

Like us, Williams et al. (2003) found that temporary, anthropogenically modified streams in England, which they called “ditches”, had lower taxon richness than perennial streams and rivers in the same region, but that they supported specialist temporary water invertebrates not found elsewhere. As a result, they considered the ditches an important aquatic habitat because of their contribution to regional biodiversity. In contrast, Datry et al (2014) in their studies in several regions of Europe, North America, and New Zealand, did not note specialist taxa in rivers and streams without perennial flow and indicated that taxa in non-perennial lotic habitats were nested subsets of those found in perennial systems. This disagreement between studies indicates that regional or other environmental factors may determine whether or not specialist temporary water invertebrates occur in temporary streams. Further studies at broad spatial scales may help identify factors accounting for differences in current study results.

Although agricultural land use has been implicated in increasing water temperatures, nutrients, pesticides, and fine sediments in permanent streams (Rothrock et al. 1998, Cuffney et al. 2000, Waite and Carpenter 2000, Stone et al. 2005, Anderson et al. 2006), effects of agriculture may be less for temporary streams in the southern Willamette Valley. First, because these streams dry during summer and only flow during the cooler parts of the year, removal of riparian plants that could affect summer water temperatures would not likely cause temperatures in other seasons to rise above physiological thresholds for temporary stream invertebrates. In addition, much of the cropland in the southern Willamette Valley is used for grassseed production (Jackson 1993; Klock et al. 2002). Grasses grown for seed production complete their growth during the wet season, and seed is harvested in the dry season (Nelson et al. 2006); as a result, water withdrawal for irrigation is often unnecessary. In addition, many of the grasses grown for seed are perennial crops, so there is less soil disturbance in years after stand establishment. Consequently, these fields export less sediment and adsorbed agrochemicals to stream channels than those in which annual crops are grown. Valley-wide soil losses from water erosion in perennial rye grass fields are estimated to be 2.1 metric tons/ha/year compared to 4.5 metric tons/ha/year for annual cereal grains in this region (Steiner et al. 2006); the US national average rate of soil loss for all cropland due to water erosion is 6.1 metric tons/ha/year (USDA-NRCS 2010). Additional soil conservation (e.g., returning chopped straw

residue to fields and direct seeding/no till planting) and nutrient management strategies have been investigated, which could further protect water and habitat quality for aquatic organisms (Nelson et al. 2006, Steiner et al. 2006).

As stream-bottom vegetation management could counteract some of the effects of watershed agriculture on invertebrates, we examined the effects of different amounts of stream-bottom vegetation. Studies in permanent stream have demonstrated higher invertebrate densities and numbers of taxa associated with habitats where aquatic vegetation was present (Gregg and Rose 1985, Armitage et al. 2001, Armitage et al. 2003). In contrast, our results indicated that local stream-bottom vegetation affected invertebrate density but did not affect taxon richness. Though increasing benthic invertebrate density may be useful if food is limiting for native fish and amphibians in these temporary streams, stream-bottom vegetation does not seem to support more invertebrate taxa, particularly those characteristic of less agriculturally influenced sites.

Another factor to consider about stream-bottom vegetation is that we never observed high levels of stream bottom vegetation if a corridor of riparian trees was present at a study site. This was especially apparent in the four sets of paired sites used to test the effect of stream-bottom vegetation in highly agricultural watersheds, but was also noticeable in several sites in the southern valley survey. In the paired sites within the same streams, sites without riparian trees had extensive growths of stream-bottom vegetation, while just a few hundred meters away sites with riparian trees had very little stream-bottom vegetation. Perhaps shading from the riparian canopy prevents dense growth of stream-bottom plants. However, in the Willamette Valley, areas with riparian trees and shrubs support more wintering birds and bird species than those without woody vegetation (McComb et al. 2005), and areas with trees adjacent to bodies of water are important for amphibians that make extensive use of upland habitat (Pearl et al. 2005). Thus, it seems there may be a need to find a balance between encouraging stream-bottom vegetation for increasing invertebrate densities and maintaining habitat conditions that benefit several native vertebrate species in the riparian habitats.

Another option to enhance invertebrate assemblages could be to take some agricultural land out of production. Our results show that even modest reduction in the percentage of watershed land in agriculture could be helpful, because at sites with approximately 75% of watershed area under agricultural use, the invertebrate assemblages were similar to those at least disturbed sites. Currently, many low-lying areas in the southern Willamette Valley are farmed despite the fact that they remain wet for extended periods and consequently have lower crop yields (J. J. Steiner, USDA-ARS, personal communication). These lands with lower crop production could potentially be set aside with the aid of federal conservation programs to maximize biodiversity and minimize effects on farm incomes.

5. Conclusions

Our study found a range of invertebrate assemblages that included dominant non-insect groups characterized by drought-resistant stages that survive the dry period on site, and insects that may have resistant stages for on-site survival, but also have winged adult stages to disperse to and from other habitats. Because our results suggest that invertebrates, integral to wet-season food webs, are affected by agricultural land use, these aquatic communities should be integrated into conservation planning and agricultural management. Studies of other temporary streams in agriculturally rich regions will further our understanding of how to balance the conservation of stream biota within the context of vital agricultural landscapes.

Acknowledgements

We thank cooperating grass seed farmers, The Nature Conservancy Willow Creek Preserve staff, and the Benton County Natural Areas and Parks Department for site access. Jeff Steiner of the U.S. Department of Agriculture, Agricultural Research Service (USDA-ARS) and Mark Mellbye of the Oregon State University (OSU) Extension Service were instrumental in launching a unique cooperation between USDA-ARS researchers, grass seed producers, and OSU faculty. Continuing support was provided by Kathryn Boyer (USDA, Natural Resources Conservation Service), and Steve Griffith

(USDA-ARS). Field help was provided by Randy Colvin, Lance Wyss, Sue Reithel, Richard Van Driesche, Bill Gavin, and Rick Caskey. Machel Nelson, Don Streeter and others at the USDA-ARS analytical lab in Corvallis, OR, performed water quality analyses. Allison Evans and D. Christopher Rogers helped to confirm the identification of *Dumontia oregonensis*. Additional statistical help was provided by Alan Herlihy at OSU Department of Fisheries and Wildlife.

Funding

This work was supported by a special grant from the USDA, USDA NIFA (Conservation Effects Assessment Program Grant # 2006-51130-03705), by additional contributions from the USDA Natural Resources Conservation Service, the Oregon Seed Council and by Oregon State University.

577 **References**

- 578 Aguiar, F.C., Ferreira, M.T., Pinto, P., 2002. Relative influence of environmental variables on
579 macroinvertebrate assemblages from an Iberian basin. *J. N. Am. Benthol. Soc.* 21:43-53.
- 580 Anderson, B.S., Hunt, J.W., Phillips, B.M., Nicely, P.A., Gilbert, K.D., De Vlaming, V., Connor, V.,
581 Richard, N., Tjeerdema, R.S., 2003. Ecotoxicologic impacts of agricultural drain water in the Salinas
582 River, California, USA. *Environ. Toxicol. Chem.* 22:2375-2384.
- 583 Anderson, B.S., Phillips, B.M., Hunt, J.W., Worcester, K., Adams, M., Kapellas, N., Tjeerdema, R.S.,
584 2006. Evidence of pesticide impacts in the Santa Maria River Watershed, California, USA. *Environ.*
585 *Toxicol. Chem.* 25:1160-1170.
- 586 Armitage, P.D., Lattman, K., Kneebone, N., Harris, I., 2001. Bank profile and structure as determinants of
587 macroinvertebrate assemblages—seasonal changes and management. *Regul. River.* 17:543-556.
- 588 Armitage, P.D., Szoszkiewicz, K., Blackburn, J.H., Nesbitt, I., 2003. Ditch communities: a major
589 contributor to floodplain biodiversity. *Aquat. Conserv.* 13:165-185.
- 590 Banks, J.L., Li, J., Herlihy, A.T., 2007. Influence of clearcut logging, flow duration, and season on
591 emergent aquatic insects in headwater streams of the central Oregon Coast Range. *J. N. Am. Benthol.*
592 *Soc.* 26:620-632.
- 593 Boag, P.G., 1992. Environment and experience: settlement culture in nineteenth century Oregon.
594 University of California Press, Berkeley, California.
- 595 Burnham, K.P., Anderson, D.R., 2002. Model selection and multimodel inference: a practical
596 information-theoretic approach. 2nd edition. Springer-Verlag, New York.
- 597 Caton, L.W., 1991. Improved subsampling methods for the EPA "Rapid Bioassessment" benthic
598 protocols. *N. Am. Benthol. Soc. Bul.* 8:317-319.
- 599 Coggerino, L., Cellot, B., Bournard, M., 1995. Microhabitat diversity and associated macroinvertebrates in
600 aquatic banks of a large European river. *Hydrobiologia* 304:103-115.
- 601 Colvin, R.W., 2006. Fish and amphibian use of intermittent streams within the upper Willamette Basin,
602 Oregon. MS Thesis. Oregon State University, Corvallis, Oregon.
- 603 Colvin, R.W., Giannico, G.R., Li, J., Boyer, K.L., Gerth, W.J., 2009. Fish use of intermittent
604 watercourses draining agricultural lands in the upper Willamette River valley, Oregon. *Trans. Am. Fish.*
605 *Soc.* 138:1302-1313.
- 606 Cooper, C.M., 1993. Biological effects of agriculturally derived surface water pollutants on aquatic
607 systems--a review. *J. Environ. Qual.* 22:402-408.
- 608 Cuffney, T.F., Meador, M.R., Porter, S.D., Gurtz, M.E., 2000. Responses of physical, chemical, and
609 biological indicators of water quality to a gradient of agricultural land use in the Yakima River Basin,
610 Washington. *Environ. Monit. Assess.* 64:259-270.

611 Datry, T., Larned, S.T., Fritz, K.M., Bogan, M.T., Wood, P.J., Meyer, E.I., Santos, N.A., 2014. Broad-
612 scale patterns of invertebrate richness and community composition in temporary rivers: effects of flow
613 intermittence. *Ecography* 37:94-104.

614 Davis, S., Golladay, S.W., Vellidis, G., Pringle, C.M., 2003. Macroinvertebrate biomonitoring in
615 intermittent coastal plain streams impacted by animal agriculture. *J. Environ. Qual.* 32:1036-1043.

616 Del Rosario, R.B., Resh, V.H., 2000. Invertebrates in intermittent and permanent streams: is the
617 hyporheic zone a refuge from drying? *J. N. Am. Benthol. Soc.* 19:680-696.

618 Dieterich, M., Anderson, N.H., 1995. Life cycles and food habits of mayflies and stoneflies from
619 temporary streams in western Oregon. *Freshw. Biol.* 34:47-60.

620 Dieterich, M., Anderson, N.H., 2000. The invertebrate fauna of summer-dry streams in western Oregon.
621 *Archiv für Hydrobiol.* 147:273-295.

622 Graham, M.H. 2003. Confronting multicollinearity in ecological multiple regression. *Ecology* 84:2809-
623 2815.

624 Gregg, W.W., Rose, F.L., 1985. Influences of aquatic macrophytes on invertebrate community structure,
625 guild structure, and microdistribution in streams. *Hydrobiologia* 128:45-56.

626 Herlihy, A.T., Gerth, W.J., Li, J., Banks, J.L., 2005. Macroinvertebrate community response to natural
627 and forest harvest gradients in western Oregon headwater streams. *Freshw. Biol.* 50:905-919.

628 Hulse, D.W., Gregory, S.V., Baker J.P. (editors), 2002. Willamette River basin: trajectories of
629 environmental and ecological change. Oregon State University Press, Corvallis, Oregon.

630 Jackson, P.L., 1993. Agriculture. Pages 59-69 in P. L. Jackson, and I. J. Schlosser(editors). *Atlas of the*
631 *Pacific Northwest*. Oregon State University Press, Corvallis, Oregon.

632 Jacobi, G.Z., Cary, S.J., 1996. Winter stoneflies (Plecoptera) in seasonal habitats in New Mexico, USA. *J.*
633 *N. Am. Benthol. Soc.* 15:690-699.

634 Jacobus, L.M., Wiersma, N.A., 2014. The genera *Anafroptilum* Kluge, 2011 and *Neocloeon* Traver, 1932,
635 reinstated status, in North America, with remarks about the global composition of *Centroptilum* Eaton,
636 1869 (Ephemeroptera: Baetidae). *Zootaxa* 3814:385-391.

637 Klock, C., Smith, S., O'Neil, T., Goggans, R., Barrett, C., 2002. Willamette Valley land use/land cover
638 map. Northwest Habitat Institute, Corvallis, Oregon.

639 Kruskal, J.B., Wish, M., 1978. *Multidimensional Scaling*. Sage Publications, Beverly Hills, California.

640 Larsen, D.P., Herlihy, A.T., 1998. The dilemma of sampling streams for macroinvertebrate richness. *J. N.*
641 *Am. Benthol. Soc.* 17:359-366.

642 Li, J., Larsen, D.P., 2001. Variability in stream macroinvertebrates at multiple spatial scales. *Freshw. Biol.*
643 46:87-97.

644 Mazor, R.D., Stein, E.D., Ode, P.R., Schiff, K., 2014. Integrating intermittent streams into watershed
645 assessments: applicability of an index of biotic integrity. *Freshw. Sci.* 33:459-474.

646 McComb, B., Bilsland, C.D., Steiner, J.J., 2005. Associations of winter birds with riparian condition in
647 the lower Calapooia watershed, Oregon. Northwest Sci. 79:164-171.

648 Merritt, R.W., Cummins, K.W. (editors), 1996. An introduction to the aquatic insects of North America.
649 Kendall/Hunt Publishing Company, Dubuque, IA.

650 Meyer, A., Meyer, E.I., 2000. Discharge regime and the effect of drying on macroinvertebrate
651 communities in a temporary karst stream in East Westphalia (Germany). Aquatic Sci. 62:216-231.

652 Nelson, M.A., Griffith, S.M., Steiner, J.J., 2006. Tillage effects on nitrogen dynamics and grass seed crop
653 production in western Oregon, USA. Soil Sci. Am. J. 70:825-831.

654 OWEB, 2001. Water quality monitoring technical guidebook. Oregon Plan for Salmon and Watersheds,
655 Salem, Oregon. http://www.oregon.gov/OWEB/docs/pubs/wq_mon_guide.pdf

656 Paltridge, R.M., Dostine, P.L., Humphrey, C.L., Boulton, A.J., 1997. Macroinvertebrate recolonization
657 after re-wetting of a tropical seasonally-flowing stream (Magela Creek, Northern Territory, Australia).
658 Mar. Freshwater Res. 48:633-645.

659 Pearl, C.A., Adams, M.J., Leuthold, N., Bury, R.B., 2005. Amphibian occurrence and aquatic invaders in
660 a changing landscape: Implications for wetland mitigation in the Willamette Valley, Oregon, USA.
661 Wetlands 25:76-88.

662 Price, K., Suski, A., McGarvie, J., Beasley, B., Richardson, J.S., 2003. Communities of aquatic insects of
663 old-growth and clearcut coastal headwater streams of varying flow persistence. Can. J. Forest Res.
664 33:1416-1432.

665 Prokopy, W.R., 1994. Ammonia in surface water, wastewater. QuikChem Method 10-115-01-1-A. Lachat
666 Instruments, Milwaukee, Wisconsin.

667 Rinella, F.A., Janet, M.L., 1998. Seasonal and spatial variability of nutrients and pesticides in streams of
668 the Willamette Basin, Oregon, 1993-95. U.S. Geological Survey. Report 97-4082-C.

669 Rothrock, J.A., Barten, P.K., Ingman, G.I., 1998. Land use and aquatic biointegrity in the Blackfoot River
670 watershed, Montana. J. Am. Water Resour. Assoc. 34:565-581.

671 Sanches-Montoya, M.M., Vidal-Abarca, M.R., Suarez, M.L., 2010. Comparing the sensitivity of diverse
672 macroinvertebrate metrics to a multiple stressor gradient in Mediterranean streams and its influence on
673 the assessment of ecological status. Ecol. Indic. 10:896-904.

674 Santos-Flores, C.J., Dodson, S.I., 2003. *Dumontia oregonensis* n. fam., n. gen., n. sp., a cladoceran
675 representing a new family of 'water-fleas' (Crustacea, Anomopoda) from U.S.A., with notes on the
676 classification of the order Anomopoda. Hydrobiologia 500:145-155.

677 Smith, D.G., 2001. Pennak's freshwater invertebrates of the United States: Porifera to Crustacea, fourth
678 edition. John Wiley and Sons, New York.

679 Stewart, K.W., Anderson, N.H., 2008. The nymphs of three Nemouridae species (Plecoptera) from
680 Oregon temporary headwater streams. Trans. Am. Entomol. Soc. 134:173-183.

681 Stewart, K.W., Anderson, N.H., 2010. The life-history of *Ostrocerca dimicki* (Frison) in a short-flow,
682 summer-dry Oregon stream. *Illiesia* 6:52:57.

683 Stewart, K.W., Stark, B.P., 2002. Nymphs of North American stonefly genera. The Caddis Press,
684 Columbus, OH.

685 Steiner, J.J., Griffith, S.M., Mueller-Warrant, G.W., Whittaker, G.W., Banowetz, G.M., Elliot, L.F., 2006.
686 Conservation practices in western Oregon permanent grass seed systems: impacts of direct seeding and
687 maximum residue management on production. *Agron. J.* 98:177-186.

688 Stone, M.L., Whiles, M.R., Webber, J.A., Williard, K.W.J., Reeve, J.D., 2005. Macroinvertebrate
689 communities in agriculturally impacted southern Illinois streams: patterns with riparian vegetation, water
690 quality, and in-stream habitat quality. *J. Environ. Qual.* 34:907-917.

691 Storey, R.G., Quinn, J.M., 2008. Composition and temporal changes in macroinvertebrate communities of
692 intermittent streams in Hawke's Bay, New Zealand. *New Zeal. J. Mar. Fresh.* 42:109-125.

693 Taft, O.W., Haig, S.M., 2003. Historical wetlands in Oregon's Willamette Valley: implications for
694 restoration of winter waterbird habitat. *Wetlands* 23:51-64.

695 Taft, O.W., Haig, S.M., 2005. The value of agricultural wetlands as invertebrate resources for wintering
696 shorebirds. *Agric. Ecosyst. Environ.* 110:249-256.

697 Taylor, G.H., Bartlett, A., 1993. The climate of Oregon, climate zone 2, Willamette Valley. Oregon
698 Climate Service, Oregon State University Agricultural Research Station. Special Report 914. Corvallis,
699 Oregon.

700 Tew, M.P., 1971. The species composition and adaptations of insects in an intermittent stream in western
701 Oregon. MS Thesis. Oregon State University, Corvallis, Oregon.

702 Thorp, J.H., Covich, A.P., 2001. Ecology and classification of North American freshwater invertebrates.
703 Academic Press, San Diego, CA.

704 USDA-NRCS, 2010. Summary Report-2007 National resources inventory: soil erosion on cropland. US
705 Department of Agriculture Natural Resources Conservation Service.
706 <http://www.nrcs.usda.gov/technical/NRI/2007/nri07erosion.html>

707 Uys, M.C., O'Keefe, J.H., 1997. Simple words and fuzzy zones: early directions for temporary river
708 research in South Africa. *J. Environ. Manage.* 21:517-531.

709 Van Damme, K., Dumont, H.J., 2008. Corrections and additions to the Dumontiidae Santos-Flores &
710 Dodson, 2003 (Crustacea: Branchiopoda: Anomopoda), and implications for anomopod phylogeny.
711 *Hydrobiologia* 598:399-401.

712 Waite, I.R., Carpenter, K.D., 2000. Associations among fish assemblage structure and environmental
713 variables in Willamette Basin streams, Oregon. *Trans. Am. Fish. Soc.* 129:754-770.

714 Wendt, K., 1999. Determination of nitrate/nitrite by flow injection analysis (low flow method).
715 QuikChem Method 10-107-04-1-A. Lachat Instruments, Milwaukee, Wisconsin.

- 716 Wiggins, G.B., 1996. Larvae of the North American caddisfly genera (Trichoptera), second edition.
717 University of Toronto Press, Toronto, Ontario, Canada.
- 718 Wigington, P.J.J., Griffith, S.M., Field, J.A., Baham, J.E., Horwath, W.R., Owen, J., Davis, J.H., Rain,
719 S.C., Steiner, J.J., 2003. Nitrate removal effectiveness of a riparian buffer along a small agricultural
720 stream in western Oregon. *J. Environ. Qual.* 32:162-170.
- 721 Wigington, P.J.J., Moser, T.J., Lindeman, D.R., 2005. Stream network expansion: a riparian water quality
722 factor. *Hydrol. Proces.* 19:1715-1721.
- 723 Williams, D.D., 1996. Environmental constraints in temporary fresh waters and their consequences for the
724 insect fauna. *J. N. Am. Benthol. Soc.* 15:634-650.
- 725 Williams, D.D., 2006. *The Biology of Temporary Waters*. Oxford University Press, Oxford, UK.
- 726 Williams, D.D., Hynes, H.B.N., 1976. The ecology of temporary streams 1. The faunas of two Canadian
727 streams. *Int. Rev. gesamen Hydrobiol.* 61:761-787.
- 728 Williams, P., Whitfield, M., Biggs, J., Bray, S., Fox, G., Nicolet, P., Sear, D., 2003. Comparative
729 biodiversity of rivers, streams, ditches and ponds in an agricultural landscape in southern England. *Biol.*
730 *Cons.* 115:329-341.
- 731 Wipfli, M.S., Gregovich, D.P., 2002. Export of invertebrates and detritus from fishless headwater streams
732 in southeastern Alaska: implications for downstream salmonid production. *Freshw. Biol.* 47:957-969.
- 733 Wyss, L.A., Dugger, B.D., Herlihy, A.T., Gerth, W.J., Li, J.L., 2013. Effects of grass seed agriculture on
734 aquatic invertebrate communities inhabiting seasonal wetlands of the southern Willamette Valley,
735 Oregon. *Wetlands* 33:921-937.

Figure 1.

Locations of southern Willamette Valley temporary stream sites sampled in 2002-03 and 2008 in western Oregon (A), and paired vegetated-non-vegetated sites sampled in 2006 (B). Black lines in map A indicate the Willamette River (flowing north). Gray lines are major Willamette River tributaries; dashed lines are temporary streams (not drawn in map A). In map A, filled gray circles indicate sites sampled in the southern valley survey (2002-03); the 5 circles with central black dots are highly agricultural sites that were resampled in 2008 to compare with least-disturbed sites. Stars indicate least-disturbed sites sampled in 2008 only.

Figure 2.

Wet-season hydrograph for one of the Willamette Valley temporary stream sites sampled. Measurements from a pressure transducer in a year when invertebrate samples were not collected (2003-04). Gap in recording occurred in January while data were downloaded.

Figure 3.

NMS ordination plots of Willamette Valley temporary stream sites based on Bray-Curtis dissimilarity of benthic invertebrate assemblages. Two main axes of three-dimensional ordinations shown. A) Plot of 18 sites from the 2002-2003 southern Willamette Valley survey. Sites east (Δ) and west (O) of the Willamette River. Vectors show correlations with invertebrate metrics (black) and physico-chemical and landscape variables (gray). B) Plot of southern Willamette Valley survey sites (2002-2003) and sites from highly agricultural and least disturbed watersheds (2008). Southern valley survey sites east (Δ) and west (O) of the Willamette River, and sites from highly agricultural ($\blacktriangle$) and least disturbed ($\bullet$) watersheds.

Figure 4.

Benthic invertebrate densities at paired vegetated and non-vegetated substrate sites within 4 Willamette Valley lowland temporary streams.

Table 1.

Physico-chemical and watershed characteristics of temporary stream sites sampled in the southern Willamette Valley survey (2002-2003). Site physical measurements were taken late April or early May. Water quality data were averaged from 7 grab samples collected every 2 to 3 weeks from mid-December 2002 through early May 2003. Watershed measurements were performed with GIS. Land use data were from the Northwest Habitat Institute (Klock et al. 2002).

Variable	Median (Range)
Depth	0.30m (0.11 – 0.54m)
Velocity	0.14m/s (0.03 – 0.35m/s)
Wetted width	3.0m (1.7 – 7.3m)
Vegetated stream bottom	25% (0 – 83%)
NO ₃ -N	4.1mg/L (0.8 – 7.5mg/L)
PO ₄ -P	0.05mg/L (0.01 – 0.11mg/L)
Conductivity	170µS/cm (55 – 244µS/cm)
Suspended sediment	81mg/L (68 – 116mg/L)
Watershed area	8.8km ² (0.7 – 21.8km ²)
% Watershed agriculture	86% (55 – 98%)
Distance to downstream permanent water	2.1km (0.4 – 6.5km)
Watershed slope	1.5% (0.1 – 3.3%)

Table 2.

Benthic macroinvertebrate summary metrics for temporary stream sites from the southern Willamette Valley survey (2002-03) and least disturbed and highly agricultural sites (2008).

Variable	Southern valley survey Median (Range)	Least disturbed Median (Range)	Highly agricultural Median (Range)
Density (# / m ²)	2923 (207 – 14207)	5201 (3228 – 22471)	3465 (2393 – 13167)
%non-insect individuals	88.7 (31.1 – 99.8)	62.8 (40.9 – 86.1)	97.2 (77.3 – 100)
%EPT individuals	0.9 (0 – 4.9)	9.6 (1.3 – 24.2)	0 (0 – 0.7)
Total taxon richness	18 (9 – 26)	24 (19 – 27)	13 (9 – 15)
EPT richness	1 (0 – 6)	5 (2 – 7)	0 (0 – 1)
Modified Hilsenhoff Biotic Index	6.2 (5.5 – 7.0)	6.0 (5.2 – 6.5)	6.4 (5.9 – 7.1)

Table 3.

Most common invertebrates collected in 2003 benthic samples at temporary stream sites in southern Willamette Valley. Taxa present at over 60% of the 18 survey sites are shown.

Benthic taxa	% of sites where taxa occurred
Nematoda	100
Oligochaeta	100
Ostracoda	94.4
Orthocladinae	88.9
Tanytarsini	88.9
<i>Caecidotea</i>	88.9
<i>Crangonyx</i>	83.3
Turbellaria	83.3
Ceratopogoninae	77.8
Harpacticoida	77.8
<i>Limnephilus</i> spp.	72.2
Acari	72.2
Cyclopoida	66.7
<i>Hydra</i>	66.7
Lymnaeidae	61.1
Pisidiidae	61.1

Mean daily stage height (m)