Maximum likelihood estimation of the mixture of log-concave densities

Hao Hu^{a,*}, Yichao Wu^a, Weixin Yao^b

^aDepartment of Statistics, North Carolina State University, Raleigh, North Carolina 27695, U.S.A.

^bDepartment of Statistics, University of California, Riverside, California 92521, U.S.A.

Abstract

Finite mixture models are useful tools and can be estimated via the EM algorithm. A main drawback is the strong parametric assumption about the component densities. In this paper, a much more flexible mixture model is considered, which assumes each component density to be log-concave. Under fairly general conditions, the log-concave maximum likelihood estimator (LCMLE) exists and is consistent. Numeric examples are also made to demonstrate that the LCMLE improves the clustering results while comparing with the traditional MLE for parametric mixture models.

Keywords: Consistency, Log-concave maximum likelihood estimator (LCMLE), Mixture model.

1. Introduction

The finite mixture model (McLachlan & Peel, 2000; McNicholas & Murphy, 2008) provides a flexible methodology for both theoretical and practical analysis. It has a density of the form

$$f(x) = \sum_{j=1}^K \lambda_j g_j(x; \theta_j) \quad x \in \mathbb{R}^d, \quad (1.1)$$

5 where $\lambda_1, \dots, \lambda_K$ are the mixing proportions and $g_j(x; \theta_j)$'s are component densities. The unknown parameters in the mixture model (1.1) can be estimated by the EM algorithm, see e.g. Dempster *et al.* (1977) and McLachlan & Krishnan (2007). One major

*Correspondence to: North Carolina State University, 5109 SAS Hall, 2311 Stinson Dr, Raleigh, NC, 27695, United States

Email addresses: hhu5@ncsu.edu (Hao Hu), wu@stat.ncsu.edu (Yichao Wu), weixin.yao@ucr.edu (Weixin Yao)

drawback of the traditional mixture model (1.1) is the strong parametric assumption about the component density g_j . It is often too restrictive and the density estimation
10 may be inaccurate due to the model misspecification. Another drawback is that each model requires a specific EM algorithm based on the parametric assumption.

To relax the parametric assumption, nonparametric shape constraints are becoming increasingly popular. In this paper, we make one fairly general shape constraint for our mixture model. We assume that each component density is log-concave. A density g
15 is log-concave if $\log g$ is concave. Examples of log-concave densities include normal, Laplace, logistic, as well as gamma and beta with certain parameter constraints. Log-concave densities have lots of nice properties as described by Balabdaoui *et al.* (2009). Their nonparametric maximum likelihood estimators were studied by Dümbgen & Rufibach (2009), Cule *et al.* (2010), Cule & Samworth (2010), Chen & Samworth (2013),
20 Pal *et al.* (2007) and Dümbgen *et al.* (2011) (referred as [DSS 2011] thereafter). The convergence rates of these estimators for log-concave densities were studied by Doss & Wellner (2013) and Kim & Samworth (2014). Such estimators provide more generality and flexibility without any tuning parameters.

In our model, we assume that $\mathbf{x}_1, \dots, \mathbf{x}_n$ are independent d -dimensional random
25 variables with distribution Q_0 and the mixture density f_0 . The mixture density f_0 belongs to a given class

$$\mathcal{F} = \{f : f(x) = \sum_{j=1}^K f_j(x) = \sum_{j=1}^K \lambda_j \exp\{\phi_j(x)\}, \lambda \in \Lambda, \phi \in \Phi\}, \quad (1.2)$$

where $\lambda = (\lambda_1, \dots, \lambda_K)$, $\Lambda = \{(\lambda_1, \dots, \lambda_K) : 0 < \lambda_j < 1, \sum_{j=1}^K \lambda_j = 1\}$,
 $\phi = (\phi_1, \dots, \phi_K)$, and $\Phi = \{(\phi_1, \dots, \phi_K) : \phi_j \text{ is concave}\}$. We assume that each ϕ_j is continuous and is coercive in the sense that $\phi_j(x) \rightarrow -\infty$ as $\|x\| \rightarrow \infty$ ($j =$
30 $1, \dots, K$).

One issue for mixture models is that the likelihood might be unbounded in some cases. For example, the likelihood function for a normal mixture takes the form of
 $L(\theta|x) = \sum_{i=1}^n (\lambda g(x_i|\mu_1, \sigma_1^2) + (1-\lambda)g(x_i|\mu_2, \sigma_2^2))$, where $\theta = \{(\lambda, \mu_1, \mu_2, \sigma_1^2, \sigma_2^2) : \sigma_1^2, \sigma_2^2 > 0, \lambda \in (0, 1)\}$ and g is the density function for the standard normal distribution. When $\mu_1 = x_1$ and $\sigma_1^2 \rightarrow 0$, $L(\theta|x) \rightarrow \infty$ (see Section 3.10 of McLachlan
35

& Peel (2000) for detailed discussions). Many methods have been proposed to solve the unboundedness issue of mixture likelihood, see for example, Hathaway (1985), Chen *et al.* (2008), and Yao (2010). Note that, similar to traditional normal mixture models with unequal variances, the likelihood functions for mixtures of log-concave
40 densities are unbounded as well. Thus, similar to Hathaway (1985), we will define LCMLE on a constrained parameter space. Let $M_j(\phi) = \max_{x \in \mathbb{R}^d} \{\phi_j(x)\}$, $M_{(1)}(\phi) = \min_j \{M_j(\phi)\}$, and $M_{(K)}(\phi) = \max_j \{M_j(\phi)\}$. We further define the ratio $\mathcal{S}(\phi) = M_{(1)}(\phi)/M_{(K)}(\phi)$. Here, we borrow the idea of Hathaway (1985) by restricting our interest to a constrained subspace Φ_η such that $\Phi_\eta = \{\phi \in \Phi : |\mathcal{S}(\phi)| \geq$
45 $\eta > 0\}$ for some $\eta \in (0, 1]$. This restriction avoids estimating the case that the modes of different components differ a lot. By restricting on Φ_η , we focus our interest on $f \in \mathcal{F}_\eta$, where

$$\mathcal{F}_\eta = \{f : f(x) = \sum_{j=1}^K f_j(x) = \sum_{j=1}^K \lambda_j \exp\{\phi_j(x)\}, \lambda \in \Lambda, \phi \in \Phi_\eta\}. \quad (1.3)$$

Let Q_n be the empirical distribution of $X_1, \dots, X_n$. The (restricted) log-concave maximum likelihood estimator (LCMLE) is

$$f_n = f(\cdot | Q_n) = \operatorname{argmax}_{f \in \mathcal{F}_\eta} \int \log(f) dQ_n. \quad (1.4)$$

50 In practice, similar to Hathaway (1985), picking η can be tricky for some extreme case. If η is too small, there might be a chance that some boundary point $|\mathcal{S}(\phi)| = \eta$ maximizes the log-likelihood and the solution will depend on the choice of η . In this paper, we do not focus on the issue of choosing η . The constrained subspace Φ_η is mainly used for theoretical development. Based on our empirical experience, if we
55 start the algorithm from a reasonable initial value, such as the maximum likelihood estimate assuming all components are normal with equal variance, the unboundedness issue is very rare.

Many methods have been proposed to relax the parametric assumption of (1.1). Hunter *et al.* (2007), Bordes *et al.* (2006a), Butucea & Vandekerckhove (2014), and
60 Chee & Wang (2013) considered the extension of (1.1) by assuming all component densities are symmetric but unknown. Bordes *et al.* (2006b), Bordes & Vandekerckhove

(2010), Hohmann & Holzmam (2013), Xiang *et al.* (2014), and Ma & Yao (2015) considered the extension of (1.1) when $K = 2$ and one of the component densities is symmetric but unknown. Mixtures of log-concave densities have been studied by
65 Chang & Walther (2007), Cule *et al.* (2010) and Balabdaoui & Doss (2014). Chang & Walther (2007) provided an EM-type algorithm and demonstrated sound numerical results in the simulation study. Cule *et al.* (2010) applied the log-concave mixture model to the Wisconsin breast cancer data set. Balabdaoui & Doss (2014) considered a special case when all components have the same symmetric log-concave densities but
70 with different location parameters, and proved the $\sqrt{n}$ -consistency of their proposed M-estimators for mixing proportion as well as location parameters. Note that these models are special cases from the family of $\mathcal{F}$. Therefore, their estimators and asymptotic results cannot be applied here. For example, the mixture of normal distributions with different component means and variances belongs to $\mathcal{F}$ but does not belong to the
75 model family considered by Balabdaoui & Doss (2014).

To the best of our knowledge, none of the existing works has studied the theoretical properties of the estimator for the log-concave mixture model (1.2) under such general conditions. This paper aims to fill in this gap. We show that theoretically, the LCMLE (in the restricted subset $\mathcal{F}_\eta$) exists, and is consistent under fairly general
80 conditions. However, we want to point out that the extension of the properties of the log-concave density to mixtures of log-concave densities is not trivial. The log-density $l_n = l(\cdot|Q_n) = \log f_n$ is no longer guaranteed to be a concave function. Consequently, many nice theoretical properties stated in DSS 2011 no longer hold for our mixture model.

85 The rest of the paper is organized as follows. Section 2 introduces the basic setup, model details, and notations. Section 3 states the theoretical properties. We review the EM-type algorithm for log-concave mixture models in Section 4. Simulation and real data studies are conducted in Section 5 and 6. We end the article with a short conclusion in Section 7. The proofs and lemmas are presented in the appendix.

90 2. Log-concave maximum likelihood estimator

Let $\mathcal{Q} = \mathcal{Q}(d)$ be the family of all distributions Q on $\mathbb{R}^d$. Our goal is to maximize a log-likelihood-type functional:

$$L(\phi, \lambda, \pi, Q) = \int \log\left[\sum_{j=1}^K \lambda_j \exp\{\phi_j(x)\}\right] dQ(x) - \sum_{j=1}^K \pi_j \left(\int \exp\{\phi_j(x)\} dx - 1\right), \quad (2.1)$$

where π_j 's are Lagrange multipliers to incorporate the constraint $\int \exp\{\phi_j(x)\} dx = 1$ ($j = 1, \dots, K$). We define a profile log-likelihood:

$$L(Q) = \sup_{\phi \in \Phi, \lambda \in \Lambda, \pi} L(\phi, \lambda, \pi, Q). \quad (2.2)$$

If, for fixed Q , (ψ, λ^*, π^*) maximizes $L(\phi, \lambda, \pi, Q)$, it will automatically satisfy that:

$$\pi_j^* = E(\pi(j|x)) = \int \frac{\lambda_j^* \exp\{\psi_j(x)\}}{(\sum_{h=1}^K \lambda_h^* \exp\{\psi_h(x)\})} dQ(x); \quad (2.3)$$

$$\int \exp\{\psi_j(x)\} dx = 1 \quad (j = 1, 2, \dots, K). \quad (2.4)$$

95 Note that differing from the non-mixture setting in DSS 2011, π_j^* is not equal to 1.

To verify this, note that $\phi + c \in \Phi$ for any fixed vector of functions $\phi \in \Phi$ and arbitrary $c = (c_1, \dots, c_K)^T \in \mathbb{R}^K$, and

$$\begin{aligned} \frac{\partial L(\psi + c, \lambda, \pi, Q)}{\partial c_h} \Big|_{c=0} &= \left(\int \frac{\lambda_h \exp\{\psi_h(x)\}}{\sum_{j=1}^K \lambda_j \exp\{\psi_j(x)\}} dQ(x) - \pi_h \int e^{\psi_h(x)} dx \right) = 0, \\ \frac{\partial L(\psi, \lambda, \pi, Q)}{\partial \pi_h} &= 1 - \int \exp\{\psi_h(x)\} dx = 0. \end{aligned}$$

The maximizer (ψ, λ^*) forms the log-likelihood maximizer $l^*(x) = \log \sum_{j=1}^K \lambda_j^* e^{\psi_j(x)}$.

3. Theoretical Properties

Before we state the main theories, we first define the convex support of a distribution.

Definition For any distribution Q , let $Q(C)$ be the probability measure of the set C . The convex support of Q is the set such that:

$$csupp(Q) = \bigcap \{C : C \subseteq \mathbb{R}^d \text{ closed and convex, } Q(C) = 1\}.$$

100 The convex support is itself closed and convex with $Q(csupp(Q)) = 1$.

In the following text, we define:

$\mathcal{Q}^1 = \{Q \in \mathcal{Q} : \int \|x\| dQ < \infty\}$, (we define $\|x\|$ as Euclidean norm in our paper).

$\mathcal{Q}^0 = \{Q \in \mathcal{Q} : interior(csupp(Q)) \neq \emptyset\}$.

Theorem 1. For any $Q \in \mathcal{Q}^1 \cap \mathcal{Q}^0$, the value of $L(Q)$ is real and there exists a
105 maximizer:

$$(\psi, \lambda^*, \pi^*) = \underset{\phi \in \Phi_\eta, \lambda \in \Lambda, \pi}{\operatorname{argmax}} L(\phi, \lambda, \pi, Q) \text{ such that } \int e^{\psi_j(x)} dx = 1 \text{ for } j = 1, \dots, K.$$

Next, we establish the consistency of the estimated mixture density. In the following, we refer to the concept of convergence of distribution as converging with respect to Mallows distance D_1 : $D_1(Q, Q') = \inf_{(X, X')} E\|X - X'\|$, where Q and Q' are two distributions and the infimum is taken over all pairs of (X, X') such that $X \sim Q$
110 and $X' \sim Q'$. The convergence of Q_n to Q_0 with respect to Mallows distance, i.e. $\lim_{n \rightarrow \infty} D_1(Q_n, Q) = 0$, is equivalent to assuming that Q_n weakly converges to Q_0 , denoted by $Q_n \rightarrow^w Q$, and $\int \|x\| dQ_n(x) \rightarrow \int \|x\| dQ(x)$ as $n \rightarrow \infty$.

Theorem 2. Let Q_n be a sequence such that $\lim_{n \rightarrow \infty} D_1(Q_n, Q_0) = 0$ for some $Q_0 \in \mathcal{Q}^1 \cap \mathcal{Q}^0$. Then,

$$\lim_{n \rightarrow \infty} L(Q_n) = L(Q_0).$$

Let ϕ_{nj} 's and λ_{nj} 's be the maximizer corresponding to profile log-likelihood $L(Q_n)$, i.e. $f_n(x) = \sum \lambda_{nj} \exp\{\phi_{nj}(x)\} = f(\cdot | Q_n) \in \mathcal{F}_\eta$. For $f_0(x) = f(\cdot | Q_0) \in \mathcal{F}_\eta$, we have:

$$\lim_{n \rightarrow \infty, x \rightarrow y} f_n(x) = f_0(y) \text{ for all } y \notin \partial\{f_0 \geq 0\}, \quad (3.1)$$

$$\lim_{n \rightarrow \infty, x \rightarrow y} f_n(x) \leq f_0(y) \text{ for all } y \in \mathbb{R}^d, \quad (3.2)$$

$$\lim_{n \rightarrow \infty} \int |f_n(x) - f_0(x)| dx = 0. \quad (3.3)$$

The above theorem showed the consistency of the estimated mixture density. If we further assume that the true mixture density $f_0(x)$ is identifiable, then each estimated
115 component density and mixing proportions are also consistent. We will discuss more about the identifiability issue in Section 7.

4. EM-type algorithm

The EM algorithm for estimating log-concave mixture densities has already been developed by Chang & Walther (2007). Here we briefly summarize it. First we randomly generate initial values for the normal mixture EM-algorithm and run the normal mixture EM-algorithm until convergence, which will provide a good initial value. Then we use the outcome as the starting values for our EM-type algorithm. We assume the observed data $\mathbf{X} = (\mathbf{x}_1, \dots, \mathbf{x}_n)^T \in \mathbb{R}^{n \times d}$ to be incomplete and define the missing value $\mathbf{Z} = (\mathbf{z}_1, \dots, \mathbf{z}_n)^T$, where $\mathbf{x}_i = (x_{i,1}, \dots, x_{i,d})^T$ and $\mathbf{z}_i$ is a K -dimension vector with its j -th element given by:

$$z_{ij} = \begin{cases} 1 & \text{if } x_i \text{ belongs to } j\text{th group,} \\ 0 & \text{otherwise.} \end{cases}$$

So the complete log-likelihood is:

$$\log f(\phi, \lambda; \mathbf{X}, \mathbf{Z}) = \log \prod_{i=1}^n \prod_{j=1}^K [\lambda_j e^{\phi_j(\mathbf{x}_i)}]^{z_{ij}} = \sum_{i=1}^n \sum_{j=1}^K z_{ij} [\log \lambda_j + \phi_j(\mathbf{x}_i)],$$

In E-step, we replace z_{ij} by

$$z_{ij}^{(t+1)} = \frac{\lambda_j^{(t)} e^{\hat{\phi}_j^{(t)}(\mathbf{x}_i)}}{\sum_{h=1}^K \lambda_h^{(t)} e^{\hat{\phi}_h^{(t)}(\mathbf{x}_i)}}.$$

In M-step, first we update λ by $\lambda_j^{(t+1)} = \frac{1}{n} \sum_{i=1}^n z_{ij}^{(t+1)}$, $j = 1, \dots, K$. Then we update ϕ_j by maximizing $\sum_{i=1}^n z_{ij}^{(t+1)} \phi_j(\mathbf{x}_i)$ with respect to ϕ_j through the function called `mlelcd` in the R package `LogConcDEAD` (Cule *et al.* (2009)) and get estimator $\hat{\phi}_j^{(t+1)}$ for $j = 1, \dots, K$. The estimation of $\hat{\phi}_j$ has been studied by Walther (2002) and Rufibach (2007). Given i.i.d. data $X_1, \dots, X_n$ from distribution f , the Log-concave Maximum Likelihood Estimator (LCMLE) $\hat{f}_n$ exists uniquely and has support on the convex hull of the data (by Theorem 2 of Cule *et al.* (2010)). The log-likelihood estimator $\log \hat{f}_n$ is a piecewise linear function with knots which are a subset of $\{X_1, \dots, X_n\}$. Walther (2002) and Rufibach (2007) provided algorithms for computing $\hat{f}_n(X_i)$, $i = 1, \dots, n$. The entire log-density $\log \hat{f}_n$ can be computed by linearly interpolating between $\log \hat{f}_n(X_{(i)})$ and $\log \hat{f}_n(X_{(i+1)})$. Walther

(2002) and Rufibach (2007) also pointed out that it is natural to apply weights for an EM-type algorithm. The $z_{1j}^{(t+1)}, \dots, z_{nj}^{(t+1)}$ can be viewed as weights for $\mathbf{x}_1, \dots, \mathbf{x}_n$ when estimating the log-concave density ϕ_j in our algorithm for $j = 1, \dots, K$. The algorithm stops once the increasing increment $\ell^{(t+1)} - \ell^{(t)}$ is below 10^{-7} , where $\ell^{(t)} = \sum_{i=1}^n \log \sum_{j=1}^K \lambda_j^{(t)} \exp\{\phi_j^{(t)}(x_i)\}$.

To avoid the local maximum, we restart the algorithm 20 times and choose the result with the highest log-likelihood. As we discussed in Section 1, the unboundedness issue of the log-likelihood does happen infrequently, mostly due to an inappropriate initial. In our algorithm, we borrow the idea of restarting process in many existing EM-algorithms for parametric mixture models, e.g. Benaglia *et al.* (2009). If the log-likelihood goes to infinity in any iteration, our EM-type algorithm will be forced to restart from the beginning with a new randomly chosen initial.

5. Simulation Results

5.1. Copula procedure to generate multivariate log-concave mixtures

As we do not have a tuning issue for LCMLE, the most attractive application of LCMLE is the density estimation with dimensionality higher than 1. To generate data from a multivariate log-concave mixture model, we borrow the idea of the copula procedure from Chang & Walther (2007). For a d -dimensional log-concave mixture density, we observe n observations $\mathbf{x}_1, \dots, \mathbf{x}_n$, where $\mathbf{x}_i = (x_{i,1}, \dots, x_{i,d})^T \in \mathbb{R}^d$. To simplify our simulation, we focus on the model whose univariate marginal distributions are log-concave. We model the dependence structure with a normal copula. Suppose $(N_1, \dots, N_d)^T$ are multivariate normal with mean $\mathbf{0}$ and covariance matrix Σ . Let $F_1, \dots, F_d$ be the CDFs of the desired univariate log-concave distributions. Then,

$$\mathbf{x}_i = (x_{i,1}, \dots, x_{i,d})^T = (F_1^{-1}(\Phi(N_1)), \dots, F_d^{-1}(\Phi(N_d)))^T.$$

5.2. Significant Improvement when densities are log-concave mixtures

We first generate 500 observations from a univariate log-concave mixture model: $0.3\text{Logistic}(0, 1) + 0.7\text{Laplace}(5, 1)$ (referred as Model I). This setup is a more general form of Chang & Walther (2007), as Chang & Walther (2007) only considered the

case that one component is a location shift of the other. For the multivariate cases, we generate 500 observations based on the copula procedure as we discussed in Section 5.1 for Model II through IV, which are multivariate log-concave mixture models with dimensionality d from 2 to 4. For each model, component 1 (with probability 0.3) is generated as a joint normal distribution $N(0, \mathbf{I}_d)$; component 2 (with probability 0.7) is generated through a normal copula $N(0, 0.5\mathbf{I}_d + 0.5\mathbf{1}_d)$, where $\mathbf{I}_d$ is a $d \times d$ identity matrix and $\mathbf{1}_d$ is a $d \times d$ matrix of ones. The marginal distributions of component 2 are summarized in Table 1.

Table 1: The simulation setups of Model II - IV.

Model	d	Marginal Distribution of Component 2
II	2	$N(0, 1)$, and $Gamma(2, 1) + 2$
III	3	$N(0, 1)$, $Gamma(2, 1) - 1$, and $Beta(4, 1)$
IV	4	$N(0, 1)$, $Gamma(2, 1) + 2$, $Beta(4, 1)$, $Laplace(0, 1) + 1$

We repeat the simulation 100 times for each model. When evaluating the simulation results for mixture models, there is a well-known *label switching* issue when sorting the labels for mixture models (Stephens (2000); Yao & Lindsay (2012)). In this paper, we adopt the method of Yao (2015) to find labels by minimizing the distance between the estimated classification probabilities and the true labels over different permutations. After sorting the labels, we compute the mean square errors obtained by the log-concave EM algorithm (MSE_2) and compare them with the parametric normal EM-algorithm (MSE_1) to compare the accuracy of the estimated λ 's. As mixture models also serve as methods of classification, we compute the average misclassification number (denoted as AMN_2 for the log-concave EM-algorithm and AMN_1 for the normal EM-algorithm) among the 100 replicates. We are also interested in the difference between two classification methods. One of many possible measurements to summarize the similarity between two clusterings is the Adjusted Rand Index (ARI), which ranges from -1 to 1, see Hubert & Arabie (1985) for detailed description of ARI . In this paper, we compute the average Adjusted Rand Index ($AvARI$) among the 100

replicates.

180 We report results over the 100 replicates in Table 2. We observe significant smaller MSEs for the estimated λ obtained by log-concave mixture models. Especially for Model I and II, the mean square errors obtained by log-concave mixture model are less than half of those obtained by normal mixture model. In terms of classification, the average misclassification number among the 500 observations are significantly reduced
185 as well. The *AvARI* indicates that the classification results of the log-concave mixture model and the normal mixture model are quite different, especially for Model I and II.

Table 2: Simulation results of Model I - IV.

Model	d	AMN_1	AMN_2	MSE_1	MSE_2	$AvARI$
I	1	17.56	10.86	0.0016	0.0007	0.91
II	2	30.35	13.28	0.0085	0.0013	0.86
III	3	12.43	4.79	0.0010	0.0005	0.93
IV	4	7.97	3.21	0.0006	0.0004	0.95

To compare the classification result for each replicate, we take $d = 4$ as an example and show the clustering results in Figure 1. In Figure (1a), each point represents a single replicate from Model IV's setup. The x -axis represents the number of misclassification by the Normal mixture EM-algorithm. The y -axis represents the number of misclassification by our log-concave mixture EM-algorithm. We observe significant
190 improvement in the misclassification rates, as all the points for our 100 replicates are below the identity line.

To better illustrate the finite sample performance of the LCMLE, we pick one replicate from Model I. To compare the fitted densities with the true densities, in Figure 2,
195 we plot the true component densities in the solid lines and the fitted densities in the dashed line. Even for a finite sample size of 500, the LCMLE for the log-concave mixture model approximates the true component densities well.

Figure 1: Four-dimensional clustering result: normal mixture EM-algorithm vs log-concave mixture EM-algorithm by number of misclassifications. The solid lines represent the identity.

Figure 2: EM-type algorithm estimation for log-concave mixtures for a single replicate of Model I. Solid line represents the truth and dashed line represents the estimation results. The fitted $\hat{\lambda} = 0.3076$.

5.3. Insignificant penalty when the parametric assumptions are correct

200 We are also interested in the price that we have to pay for the flexibility while the data actually are from normal mixtures. For Model V - VIII, we generate $n = 500$

observations from a joint normal mixture distribution, in which the first component (with probability 0.4) is a d -dimensional normal distribution with mean $\mathbf{0}_d$ and covariance matrix $0.5\mathbf{I}_d + 0.5\mathbf{1}_d\mathbf{1}_d^T$, and the second component (with probability 0.6) is a d -dimensional normal distribution with mean $\boldsymbol{\mu}_d$ and the same covariance matrix, where $\mu_1 = 5$, $\boldsymbol{\mu}_2 = (3, 2)^T$, $\boldsymbol{\mu}_3 = (3, 2, 2)^T$, and $\boldsymbol{\mu}_4 = (3, 1, 3, 1)^T$. We also repeat the simulation 100 times and compare the same criteria.

From Table 3, we observe no significant penalty for applying log-concave mixture models instead of normal mixture models. The MSEs and average misclassification numbers for log-concave mixture models are either almost the same or only a little bit higher than those for the multivariate normal mixture model. The classification results of the log-concave mixture model and the normal mixture model are quite similar to each other, as we observe the *AvARI*'s are close to 1 in Table 3. This phenomena is further supported in Figure (1b), which shows the classification results for Model VIII ($d = 4$). We observe no significant difference in terms of misclassifications, as most points in Figure (1b) are around the identity line. Consequently, we conclude that the log-concave mixture model is a more flexible methodology without significant penalties if the data are actually from normal mixtures.

Table 3: Simulation results of Model V - VIII.

Model	d	AMN_1	AMN_2	MSE_1	MSE_2	$AvARI$
V	1	3.17	3.51	0.0004	0.0005	0.99
VI	2	33.95	37.12	0.0018	0.0020	0.91
VII	3	21.95	23.57	0.0008	0.0008	0.94
VIII	4	17.71	18.20	0.0018	0.0020	0.96

6. Real Data Application

To further illustrate the performance of log-concave mixture models, we apply the log-concave EM algorithm to the crab data set of Campbell & Mahon (1974), which

contains two types of crabs in the data set: 100 blue crabs and 100 orange crabs. We focus on these blue crabs, which include $n_1 = 50$ males and $n_2 = 50$ females referred to as groups G_1 and G_2 , respectively. For each crab, there are five measurements. We are only interested in two of them: RW (rear width) and BD (body depth), both in unit of mm. In Figure 3, we give the scatter plot of RW and BD .

Figure 3: Scatter plot of RW (rear width) and BD (body depth) of the Blue Crab data set.

Fitting a 2-dimensional two component log-concave mixture model results in 18 observations misclassified. Fifteen observations from G_1 are misclassified into G_2 and three observations from G_2 are misclassified into G_1 . The normal mixture model results in 20 observations misclassified in total. Two additional observations from G_1 are misclassified into G_2 .

7. Conclusion

The log-concave maximum likelihood estimator (LCMLE) provides more flexibility to estimate mixture densities, when compared to the traditional parametric mixture models. The estimation of LCMLE for log-concave mixtures can be achieved by an EM-type algorithm. The LCMLE is not sensitive to the model mis-specification

and consequently, only one implementation of the EM-type algorithm is necessary. Through simulation studies, we observed significant improvements in the sense of classification and no significant penalties when the parametric assumption is indeed
240 correct.

In this paper, we proved the existence of the LCMLE for log-concave mixture models. The consistency is also proved for the estimated mixture density. If the true mixture density is identifiable, then the estimated component densities are also identifiable. However, it is not an easy task to prove the overall identifiability for the most general
245 family of mixtures of log-concave distributions in (1.2) from a nonparametric point of view. Some restrictive conditions, such as symmetry, are needed to ensure identifiability. Hunter *et al.* (2007) and Bordes *et al.* (2006a) proved the identifiability of (1.1) if $K = 2$ and both component densities are symmetric but with different location parameters. Balabdaoui & Doss (2014) has considered a special case of (1.2), when
250 $\phi_j(x; \theta_j) = \phi(x - \theta_j)$ and ϕ is a symmetric concave function about 0, and the identifiability of (1.2) follows from Hunter *et al.* (2007) and Bordes *et al.* (2006a) when $K = 2$.

Acknowledgment

We wish to thank the Associate Editor and two reviewers for their helpful comments and suggestions that led to improvements in this paper. Hu's research is partially
255 supported by National Institutes of Health grant R01-CA149569. Wu's research is partially supported by National Institutes of Health grant R01-CA149569 and National Science Foundation grant DMS-1055210. Yao's research is supported by NSF grant DMS-1461677.

260 Appendix A: Lemmas

Lemma 1 is taken from Cule & Samworth (2010). Lemma 2 to Lemma 5 are taken from DSS 2011. Lemma 6 is the extension of Lemma 2.13 of DSS 2011.

Lemma 1. For any log-concave distribution Q with density f , there exist finite constants $B_1 = B_1(Q) > 0$ and $B_2 = B_2(Q) > 0$ such that $f(x) \leq B_1 \exp(-B_2 \|x\|)$ for all $x \in \mathbb{R}^d$.

Lemma 2. The following properties of Q are equivalent:

- (a) $\text{csupp}(Q)$ has non-empty interior.
- (b) $Q(H) < 1$ for any hyperplane $H \subset \mathbb{R}^d$.
- (c) With Leb denoting Lebesgue measure on $\mathbb{R}^d$,

$$\limsup_{\delta \downarrow 0} \{Q(A) : A \subset \mathbb{R}^d \text{ closed and convex, } \text{Leb}(A) \leq \delta\} < 1.$$

Lemma 3. Let ϕ be the function such that for any $x, y \in \text{interior}(\text{dom}(\phi))$ and $t \in (0, 1)$, if $tx + (1-t)y \in \text{interior}(\text{dom}(\phi))$, $\phi(tx + (1-t)y) \geq t\phi(x) + (1-t)\phi(y)$ and for $C \subseteq \mathbb{R}^d$, $\int_C e^{\phi(x)} dx \leq 1$. We define $D_q = \{x \in C : \phi(x) \geq q\}$. For any $r < M \leq \max_{x \in \mathbb{R}^d} \phi(x)$,

$$\text{Leb}(D_r) \leq (M-r)^d e^{-M} / \int_0^{M-r} t^d e^{-t} dt.$$

Lemma 4. Let $\bar{\phi}, \phi_1, \phi_2, \dots$ be concave functions and $\phi_n \leq \bar{\phi}$. Further we assume the set $H = \{x : \liminf_{n \rightarrow \infty} \phi_n(x) > -\infty\}$ is not empty. Then there exist a subsequence $(\phi_{n(k)})_k$ of $(\phi_n)_n$ and a function ϕ such that $H \subset \text{dom}(\phi) \stackrel{d}{=} \{\phi > -\infty\}$:

$$\begin{aligned} \lim_{k \rightarrow \infty, x \rightarrow y} \phi_{n(k)}(x) &= \phi(y) \text{ for all } y \in \text{interior}(\text{dom}(\phi)), \\ \lim_{k \rightarrow \infty, x \rightarrow y} \phi_{n(k)}(x) &\leq \phi(y) \text{ for all } y \in \mathbb{R}^d. \end{aligned}$$

Lemma 5. Suppose Q_n is a sequence converged to some distribution Q and h be a nonnegative and continuous function, then

$$\liminf_{n \rightarrow \infty} \int h dQ_n \geq \int h dQ.$$

If the stronger statement $\liminf_{n \rightarrow \infty} \int h dQ_n = \int h dQ < \infty$ holds, then for any function f such that $|f|/(1+h)$ is bounded,

$$\lim_{n \rightarrow \infty} \int f dQ_n = \int f dQ.$$

Lemma 6. A point $x \in \mathbb{R}^d$ is an interior point of C if and only if

$$h(Q, x) = \sup\{Q(E) : E \subset C, E \text{ closed and convex}, x \notin \text{interior}(E)\}/Q(C) < 1.$$

Proof For $x \notin \text{interior}(E)$ and closed and convex E , there exists a unit vector $u_j \in \mathbb{R}^d$ such that E is contained in the closed set H_C which is a subset of C :

$$C \supseteq H_C(x) = \{y \in C : u^T y \leq u^T x\} \supseteq E.$$

By the definition of $h(Q, x)$ we conclude $h(Q, x) \leq Q(H_C)/Q(C) \leq 1$. There are two cases: $E \subset H_C$ and $E = H_C(x)$. For the case $E \subset H_C$, by definition $h(Q, x) < 1$ strictly. For the case $E = H_C(x)$, as we have $x \notin \text{interior}(E)$ but $x \in H_C(x)$, we conclude $x \in \partial H_C(x)$. Now if $x \notin \text{interior}(C)$, by definition, $h(Q, x) = 1$. On the other hand, if $h(Q, x) = 1$, then $Q(H_C(x)) = Q(C)$, which leads to $C = H_C(x) = E$. Combined with $x \notin \text{interior}(H_C(x))$ we can conclude that $x \notin \text{interior}(C)$. Consequently, $x \notin \text{interior}(C) \iff h(Q, x) = 1$. Thus, $x \in \text{interior}(C) \iff h(Q, x) < 1$.

Appendix B: Proof of Theorem 1

The first thing is to prove the finiteness of the log-likelihood type function.

$L(Q)$ is the supreme of $L(\phi, \lambda, \pi, Q)$ over all $\phi \in \Phi, \lambda \in \Lambda, \lambda \in \mathbb{R}^K$. If we take a special case that $\phi_j^*(x) = -(\log \lambda_j^*) - \|x\|$, $L(\phi^*, \lambda^*, \pi, Q) = \log K - \int \|x\| dQ > -\infty$. Consequently, $L(Q) > -\infty$.

Now we show $L(Q) < \infty$. As discussed at the end of Section 2, we do restrict our interest to the ϕ such that $\int e^{\phi_j(x)} dx = 1$ for $j = 1, \dots, K$. Consequently, we define the log-density as $l(x) = \log \sum_{j=1}^K \lambda_j e^{\phi_j(x)}$ and rewrite the log-likelihood-type function as $L(l, Q) = L(\phi, \lambda, \pi, Q)$. For the convenience of the proof, we define an envelope function $\bar{\phi}(x) = \max_j \{\phi_j(x)\}$, i.e. $\bar{\phi}(x) \geq l(x)$ for every $x \in \mathbb{R}^d$. This function is continuous but not smooth on $d - 1$ dimensional boundaries. These boundaries divide the $\text{csupp}(Q)$ into K sets: $C_1, \dots, C_K$. Each set C_j is defined as $C_j = \{x \in \mathbb{R}^d : \bar{\phi}(x) = \phi_j(x)\}$. The sets $C_1, \dots, C_K$ are disjoint except on the boundaries and $\text{Leb}(C_i \cap C_j) = 0$ for every $i \neq j$. For any $x, y \in C_j$ and $t \in (0, 1)$,

290 $\bar{\phi}(tx + (1-t)y) \geq t\bar{\phi}(x) + (1-t)\bar{\phi}(y)$ and $\int_{C_j} e^{\bar{\phi}(x)} dx \leq 1$. We define $M_j(\phi)$ and $\mathcal{S}(\phi)$ as stated in Section 1. As $L(l, Q) \leq \sum_{j=1}^K Q(C_j)M_j$, $M_j > -\infty$, and the restriction $|\mathcal{S}(\phi)| \geq \eta > 0$, we focus our interest on $M_j > 0$ and the only case which we have to worry about is all M_j 's increasing to infinity. We define $D_q = \{x \in \mathbb{R}^d : \bar{\phi}(x) \geq q\}$. For any $c > 0$,

$$\begin{aligned} L(l, Q) &\leq \int \bar{\phi}(x) dQ = \int_{c\text{supp}(Q) \setminus D_{-cM_{(1)}}} \bar{\phi}(x) dQ + \int_{D_{-cM_{(1)}}} \bar{\phi}(x) dQ \\ &\leq -cM_{(1)}(1 - Q(D_{-cM_{(1)}})) + M_{(K)}Q(D_{-cM_{(1)}}) \\ &\leq (1 + c\eta) \left(Q(D_{-cM_{(1)}}) - \frac{c\eta}{c\eta + 1} \right) M_{(K)}. \end{aligned}$$

We can always find sufficient large c such that the set $D_{-cM_{(1)}}$ is a closed and convex subset of $\mathbb{R}^d$. We define the set $D_{j,q} = \{x \in C_j : \bar{\phi}(x) \geq q\} \subset C_j$. Obviously $Leb(D_{-cM_{(1)}}) = \sum_{j=1}^K Leb(D_{j,-cM_{(1)}})$. For any $c > 0$, applying Lemma 3 to set $D_{j,-cM_{(1)}}$ and letting $M = M_{(1)}$ yield $Leb(D_{j,-cM_{(1)}}) \leq (1+c)M_{(1)}^d e^{-M_{(1)}}/(d! + o(1)) \rightarrow 0$ as $M_{(1)} \rightarrow \infty$ for every $j = 1, \dots, K$. Consequently, $Leb(D_{-cM_{(1)}}) \rightarrow 0$ as $M_{(1)} \rightarrow \infty$. By our definition, $\eta \in (0, 1]$. Thus, by Lemma 2, we can find sufficiently large c and small δ such that

$$\sup\{Q(D) : D \subset \mathbb{R}^d, Leb(D) \leq \delta\} < \frac{c\eta}{c\eta + 1}.$$

Thus, $L(l, Q) \rightarrow -\infty$ as $M_{(1)} \rightarrow \infty$, which indicates that when all modes of log-concave densities increase to infinity, the log-likelihood is poorly characterized. On the other hand, $L(l, Q) \leq M_{(K)}$. These considerations show that $L(Q)$ is finite and equals the supremum of $L(l, Q)$ for suitable finite M_j 's such that $M_j \in [M_{*j}, M_j^*]$ ($j = 1, \dots, K$).

Let $\phi_{m,j}$'s and $\lambda_{m,j}$'s form a sequence $l_m(x) = \log \sum \lambda_{m,j} \exp\{\phi_{m,j}(x)\}$ such that $-\infty < L(l_m, Q) \uparrow L(Q)$ as $m \rightarrow \infty$. Next, we will prove that for every $j \in \{1, \dots, K\}$, there exists a point, say, $x_{0,j} \in \text{interior}(c\text{supp}(Q))$, such that $\liminf_{m \rightarrow \infty} \phi_{m,j}(x_{0,j}) > -\infty$.

We define $\bar{\phi}_m(x) = \max_j \{\phi_{m,j}(x)\}$, $C_{m,j} = \{x \in \mathbb{R}^d : \bar{\phi}_m(x) = \phi_{m,j}(x)\}$, and $M_{m,j} = \max_{x \in \mathbb{R}^d} \phi_{m,j}(x)$. For any $j^* \in \{1, \dots, K\}$, by picking any $x_{0,j^*} \in C_{m,j^*}$

such that $\phi_{m,j^*}(x_{0,j^*}) \in [M'_{m,j^*}, M_{m,j^*})$, where $M'_{m,j^*} = \max_{x \in \partial\{C_{m,j^*}\}} \phi_{m,j^*}(x)$,
 305 there exists a sufficient small $\epsilon \geq 0$ such that the set $E_{m,j^*} = \{x \in C_{m,j^*} : \phi_{m,j^*}(x) \geq \phi_{m,j^*}(x_{0,j^*}) + \epsilon\}$ is a closed and convex subset of C_{m,j^*} and x_{0,j^*} is not an interior point of E_{m,j^*} . Thus,

$$\begin{aligned} L(l_m, Q) &= \int l_m dQ \leq \int \bar{\phi}_m(x) dQ = \sum_{j \neq j^*} \int_{C_{m,j}} \phi_{m,j}(x) dQ + \int_{C_{m,j^*}} \phi_{m,j^*} dQ \\ &\leq \sum_{j \neq j^*} M_{m,j} Q(C_{m,j}) + \phi_{m,j^*}(x_{0,j^*}) Q(C_{m,j^*}) + (M_{m,j^*} - \phi_{m,j^*}(x_{0,j^*})) Q(E_{m,j^*}) \\ &\leq \sum_{j=1}^K \max(M_{m,j}, 0) + \phi_{m,j^*}(x_{0,j^*}) Q(C_{m,j^*}) (1 - h_{j^*}(Q, x_{0,j^*})). \end{aligned}$$

These inequalities hold for the case of $\phi_{m,j^*}(x_{0,j^*}) = M_{m,j^*}$ as well ($\epsilon = 0$ accordingly). By Lemma 6, $h_{j^*}(Q, x_{0,j^*}) < 1$. Due to the fact that M_{m,j^*} is finite, $\text{interior}(C_{m,j^*})$ is not empty. Consequently, $\liminf_{m \rightarrow \infty} Q(C_{m,j^*}) > 0$, which yields

$$\begin{aligned} \phi_{m,j^*}(x_{0,j^*}) &\geq - \frac{\sum_{j=1}^K \max(M_{m,j}, 0) - L(l_m, Q)}{Q(C_{m,j^*})(1 - h_{j^*}(Q, x_{0,j^*}))} \\ &> - \frac{\sum_{j=1}^K \max(M_j^*, 0) - L(l_1, Q)}{Q(C_{m,j^*})(1 - h_{j^*}(Q, x_{0,j^*}))} > -\infty. \end{aligned}$$

Hence, the set $H_j = \{x : \liminf_{m \rightarrow \infty} \phi_{m,j}(x) > -\infty\}$ is not empty for every $j \in \{1, \dots, K\}$. From Lemma 1 we conclude that for each ϕ_j , we can find suitable finite positive constants $a_j, b_j > 0$ such that $\phi_j(x) \leq a_j - b_j \|x\| \leq a - b \|x\|$, where $a = \max_j a_j > 0$ and $b = \min_j b_j > 0$. Then by Lemma 4, there exist a subsequence $(\phi_{1,m(k_1)})_{k_1}$ of $(\phi_{1,m})_m$ and a concave function ϕ_1 such that:

$$\begin{aligned} \lim_{k_1 \rightarrow \infty, x \rightarrow y} \phi_{1,m(k_1)}(x) &= \phi_1(y) \text{ for all } y \in \text{interior}(\text{dom}(\phi_1)), \\ \lim_{k_1 \rightarrow \infty, x \rightarrow y} \phi_{1,m(k_1)}(x) &\leq \phi_1(y) \text{ for all } y \in \mathbb{R}^d. \end{aligned}$$

If we define $\phi_1 = -\infty$ on $\mathbb{R}^d \setminus \text{dom}(\phi_1)$, then we can rewrite them as:

$$\begin{aligned} \limsup_{k_1 \rightarrow \infty} \phi_{1,m(k_1)}(x) &\leq \phi_1(x) \text{ for all } x \in \partial\{\text{dom}(\phi_1)\}, \\ \lim_{k_1 \rightarrow \infty} \phi_{1,m(k_1)}(x) &= \phi_1(x) \text{ for all } x \in \mathbb{R}^d \setminus \partial\{\text{dom}(\phi_1)\}. \end{aligned}$$

We can find a sub-subsequence in the original subsequence, which has the similar property for $\phi_{2,m(k_2)}$. Keeping doing this sequentially for all $\phi_{m,j}$'s and $\lambda_{m,j}$'s will yield the common subsequence $l_{m(k)}$ and a function $l^*(x) = \log \sum \lambda_j \exp\{\phi_j(x)\}$ such that:

$$\begin{aligned} \limsup_{k \rightarrow \infty} l_{m(k)}(x) &\leq l^*(x) \quad \text{for all } x \in \mathcal{P}, \\ \lim_{k \rightarrow \infty} l_{m(k)}(x) &= l^*(x) \quad \text{for all } x \in \mathbb{R}^d \setminus \mathcal{P}, \end{aligned}$$

where $\mathcal{P} = \cup_{j=1}^K (\partial\{\text{dom}(\phi_j)\})$ and $\text{Leb}(\mathcal{P}) = 0$. The next step is to prove that $l^*(x)$ is the maximizer. Applying Fatou's lemma to the subsequence function $l_{m(k)}(x) \leq a - b\|x\|$ yields

$$\limsup_{k \rightarrow \infty} \int l_{m(k)} dQ \leq \int l^* dQ.$$

Hence,

$$L(Q) \geq l(l^*, Q) \geq \limsup_{k \rightarrow \infty} L(l_{m(k)}, Q) = L(Q),$$

from which we conclude $L(l^*, Q) = L(Q)$. The first inequality follows by the definition of $L(Q)$. The last equality follows by the definition that $l_{m(k)}$ is a sequence that
310 maximizes $L(l_{m(k)}, Q)$ to $L(Q)$ as $k \rightarrow \infty$. Thus, it concludes the existence of the maximizer l^* , which indicates the existence of λ_j^* 's and ϕ_j^* 's.

Appendix C: Proof of Theorem 2

We proof the theorem for a subsequence of Q_n . Let $L(Q_n) \rightarrow \Gamma$. As in the proof of Theorem 1, $l_n(x) \leq a - b\|x\|$ and $\inf \phi_{n,j}(x_0) > -\infty$ for some $x_0 \in \text{interior}(\text{csupp}(Q))$. Therefore, for a subsequence of $(Q_n)_n$, there exists a function l^* such that $l_n(y), l^*(y) \leq a - b\|y\|$, and

$$\begin{aligned} \limsup_{k \rightarrow \infty} l_{n(k)}(x) &\leq l^*(x) \quad \text{for all } x \in \mathcal{P}, \\ \lim_{k \rightarrow \infty} l_{n(k)}(x) &= l^*(x) \quad \text{for all } x \in \mathbb{R}^d \setminus \mathcal{P}. \end{aligned}$$

By Skorohod's theorem, there exists a probability space with random variables $X_n \sim Q_n$, $X \sim Q$ such that $X_n \rightarrow X$ almost surely. We define a random variable

$H_n = a - b\|X_n\| - l_n(X_n) \geq 0$. Applying Fatou's lemma to H_n yields,

$$\begin{aligned}\Gamma &= \lim_{n \rightarrow \infty} \int l_n dQ_n = \lim_{n \rightarrow \infty} \int (a - b\|x\|) dQ_n - E(H_n) = a - b\gamma - \liminf_{n \rightarrow \infty} E(H_n) \\ &\leq a - b\gamma - E\left(\liminf_{n \rightarrow \infty} (H_n)\right) \leq a - b\gamma - E(a - b\|X\| - l^*(X)) \\ &= b\left(\int \|x\| dQ_0 - \gamma\right) + \int l^*(X) dQ_0 = L(l^*, Q_0) \leq L(Q_0).\end{aligned}$$

Let $l_0(x) = \log \sum \lambda_j \phi_j(x)$, i.e. λ_j 's and ϕ_j 's are the results corresponding with l_0 . In the following proof we utilize a special approximation scheme. Let $l^{(\epsilon)}(x) = \log \sum \lambda_j^{(\epsilon)} \phi_j^{(\epsilon)}(x)$, $\lambda_j^{(\epsilon)} = \lambda_j$ and $\phi_j^{(\epsilon)} = \inf_{v,c}(v_j^T x + c_j)$ such that $\|v_j\| \leq \epsilon^{-1}$ and $\phi_j(y) \leq v_j^T y + c_j$. DSS 2011 shows that the approximation $\phi_j^{(\epsilon)}$ is real valued and Lipschitz continuous with constant ϵ^{-1} . Consequently, $l^{(\epsilon)}(x)$ is also Lipschitz-continuous with constant ϵ^{-1} . Moreover, $\phi_j^{(\epsilon)} \geq \phi_j$ and $\phi_j^{(\epsilon)} \downarrow \phi_j$ pointwise as $\epsilon \downarrow 0$. Thus, $l^{(\epsilon)} \downarrow l_0$ pointwise as $\epsilon \downarrow 0$ and $l^{(1)} \geq l^{(\epsilon)} \geq l_0$ for $\epsilon \in (0, 1)$. With this approximation, it follows from Lipschitz-continuity, $\int \|x\| dQ_0 = \gamma < \infty$, and the stronger version of Lemma 5 that

$$\begin{aligned}\Gamma &= \lim_{n \rightarrow \infty} \int l_n dQ_n \geq \lim_{n \rightarrow \infty} L(l^{(\epsilon)}, Q_n) = \lim_{n \rightarrow \infty} \int l^{(\epsilon)} dQ_n - \sum \pi_j \int e^{\phi_j^{(\epsilon)}(x)} dx + 1 \\ &= \int l^{(\epsilon)} dQ_0 - \sum \pi_j \int \exp(\phi_j^{(\epsilon)}(x)) dx + 1.\end{aligned}$$

Applying monotone convergence theorem to function $l^{(1)} - l^{(\epsilon)}$ and dominated convergence theorem to $\exp\{\phi_j^{(\epsilon)}\}$'s yields, $\lim_{\epsilon \rightarrow 0+} L(l^{(\epsilon)}, Q_0) = L(l_0, Q_0)$. Hence,
315 $\Gamma \geq L(Q_0)$. Combining with $\Gamma \leq L(l^*, Q_0) \leq L(Q_0)$ yields $\Gamma = L(Q_0) = L(l^*, Q_0)$, which indicates that l^* equals the maximizer $l_0 = l(\cdot | Q_0)$ that corresponds to $L(Q_0)$.

Applying to density $f_n = \exp\{l_n\}$ and $f_0 = \exp\{l_0\}$ yields,

$$\begin{aligned}\lim_{n \rightarrow \infty, x \rightarrow y} f_n(x) &= f_0(y) \text{ for all } x \in \mathbb{R}^d \setminus \mathcal{P}, \\ \lim_{n \rightarrow \infty, x \rightarrow y} f_n(x) &\leq f_0(y) \text{ for all } y \in \mathcal{P},\end{aligned}$$

where $\mathcal{P} = \cup_{j=1}^K (\partial\{f_{0j} > 0\})$ and $\text{Leb}(\mathcal{P}) = 0$. Consequently, $(f_n)_n \rightarrow f_0$ almost everywhere with respect to Lebesgue measure. In addition, $|f_n(x)| \leq e^{a-b\|x\|}$, and $\int e^{a-b\|x\|} dx$ is finite. Applying Lebesgue's dominated convergence theorem yields,

$$\lim_{n \rightarrow \infty} \int |f_n(x) - f_0(x)| dx = 0.$$

Consequently, we claim Theorem 2 to be true for a subsequence of the original sequence $(Q_n)_n$. It remains to show it is true for the entire sequence.

320 Suppose any assertion about f_n is false, then one could replace the initial sequence $(Q_n)_n$ from the start with a subsequence such that one of the following three conditions is satisfied:

- (i) $\lim_{n \rightarrow \infty} f_n(x_n) > f_0(y)$ for some sequence $(x_n)_n$ converge to point y ;
- (ii) $\lim_{n \rightarrow \infty} f_n(x_n) < f_0(y)$ for some sequence $(x_n)_n$ converge to point y ;
- 325 (iii) $\lim_{n \rightarrow \infty} \int |f_n(x) - f_0(x)| dx > 0$.

Any of these three properties are transmitted to subsequence of $(Q_n)_n$, which would lead to a contradiction.

References

- Balabdaoui, Fadoua, & Doss, Charles R. 2014. Inference for a mixture of symmetric distributions under log-concavity. *arXiv preprint arXiv:1411.4708*.
- 330 Balabdaoui, Fadoua, Rufibach, Kaspar, & Wellner, Jon A. 2009. Limit distribution theory for maximum likelihood estimation of a log-concave density. *Annals of statistics*, **37**(3), 1299.
- Benaglia, Tatiana, Chauveau, Didier, Hunter, David, & Young, Derek. 2009. mixtools: An R package for analyzing finite mixture models. *Journal of Statistical Software*, 32(6), 1–29.
- 335 Bordes, Laurent, & Vandekerkhove, Pierre. 2010. Semiparametric two-component mixture model with a known component: an asymptotically normal estimator. *Mathematical Methods of Statistics*, **19**(1), 22–41.
- 340 Bordes, Laurent, Mottelet, Stéphane, Vandekerkhove, Pierre, *et al.* 2006a. Semiparametric estimation of a two-component mixture model. *The Annals of Statistics*, **34**(3), 1204–1232.
- Bordes, Laurent, Delmas, Céline, & Vandekerkhove, Pierre. 2006b. Semiparametric Estimation of a Two-component Mixture Model where One Component is known. *Scandinavian journal of statistics*, **33**(4), 733–752.
- 345

- Butucea, Cristina, & Vandekerckhove, Pierre. 2014. Semiparametric mixtures of symmetric distributions. *Scandinavian Journal of Statistics*, **41**(1), 227–239.
- Campbell, NA, & Mahon, RJ. 1974. A multivariate study of variation in two species of rock crab of the genus *Leptograpsus*. *Australian Journal of Zoology*, **22**(3), 417–425.
- 350 Chang, George T, & Walther, Guenther. 2007. Clustering with mixtures of log-concave distributions. *Computational Statistics & Data Analysis*, **51**(12), 6242–6251.
- Chee, Chew-Seng, & Wang, Yong. 2013. Estimation of finite mixtures with symmetric components. *Statistics and Computing*, **23**(2), 233–249.
- Chen, Jiahua, Tan, Xianming, & Zhang, Runchu. 2008. Inference for normal mixtures
355 in mean and variance. *Statistica Sinica*, **18**(2), 443.
- Chen, Yining, & Samworth, Richard J. 2013. Smoothed log-concave maximum likelihood estimation with applications. *Statist. Sinica*, **23**, 1373–1398.
- Cule, Madeleine, & Samworth, Richard. 2010. Theoretical properties of the log-concave maximum likelihood estimator of a multidimensional density. *Electronic*
360 *Journal of Statistics*, **4**, 254–270.
- Cule, Madeleine, Gramacy, Robert, & Samworth, Richard. 2009. LogConcDEAD: An R package for maximum likelihood estimation of a multivariate log-concave density. *Journal of Statistical Software*, **29**(2), 1–20.
- Cule, Madeleine, Samworth, Richard, & Stewart, Michael. 2010. Maximum likelihood
365 estimation of a multi-dimensional log-concave density. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, **72**(5), 545–607.
- Dempster, Arthur P, Laird, Nan M, & Rubin, Donald B. 1977. Maximum likelihood from incomplete data via the EM algorithm. *Journal of the Royal Statistical Society. Series B (Methodological)*, 1–38.
- 370 Doss, Charles, & Wellner, Jon A. 2013. Global rates of convergence of the MLEs of log-concave and s-concave densities. *arXiv preprint arXiv:1306.1438*.

- Dümbgen, Lutz, & Rufibach, Kaspar. 2009. Maximum likelihood estimation of a log-concave density and its distribution function: Basic properties and uniform consistency. *Bernoulli*, **15**(1), 40–68.
- 375 Dümbgen, Lutz, Samworth, Richard, & Schuhmacher, Dominic. 2011. Approximation by log-concave distributions, with applications to regression. *The Annals of Statistics*, **39**(2), 702–730.
- Hathaway, Richard J. 1985. A constrained formulation of maximum-likelihood estimation for normal mixture distributions. *The Annals of Statistics*, 795–800.
- 380 Hohmann, Daniel, & Holzmam, Hajo. 2013. Semiparametric location mixtures with distinct components. *Statistics*, **47**(2), 348–362.
- Hubert, Lawrence, & Arabie, Phipps. 1985. Comparing partitions. *Journal of classification*, **2**(1), 193–218.
- Hunter, David R, Wang, Shaoli, & Hettmansperger, Thomas P. 2007. Inference for
385 mixtures of symmetric distributions. *The Annals of Statistics*, 224–251.
- Kim, Arlene KH, & Samworth, Richard J. 2014. Global rates of convergence in log-concave density estimation. *arXiv preprint arXiv:1404.2298*.
- Ma, Yanyuan, & Yao, Weixin. 2015. Flexible estimation of a semiparametric two-component mixture model with one parametric component. *Electronic Journal of
390 Statistics*, **9**, 444–474.
- McLachlan, Geoffrey, & Krishnan, Thriyambakam. 2007. *The EM algorithm and extensions*. Vol. 382. John Wiley & Sons.
- McLachlan, Geoffrey, & Peel, David. 2000. *Finite mixture models*. John Wiley & Sons.
- 395 McNicholas, Paul David, & Murphy, Thomas Brendan. 2008. Parsimonious Gaussian mixture models. *Statistics and Computing*, **18**(3), 285–296.

- Pal, Jayanta Kumar, Woodroffe, Michael, & Meyer, Mary. 2007. Estimating a Polya Frequency Function. *Lecture Notes-Monograph Series*, 239–249.
- Rufibach, Kaspar. 2007. Computing maximum likelihood estimators of a log-concave density function. *Journal of Statistical Computation and Simulation*, **77**(7), 561–574.
- Stephens, Matthew. 2000. Dealing with label switching in mixture models. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, **62**(4), 795–809.
- Walther, Guenther. 2002. Detecting the presence of mixing with multiscale maximum likelihood. *Journal of the American Statistical Association*, **97**(458), 508–513.
- Xiang, Sijia, Yao, Weixin, & Wu, Jingjing. 2014. Minimum profile Hellinger distance estimation for a semiparametric mixture model. *Canadian Journal of Statistics*, **42**(2), 246–267.
- Yao, Weixin. 2010. A profile likelihood method for normal mixture with unequal variance. *Journal of Statistical Planning and Inference*, **140**(7), 2089–2098.
- Yao, Weixin. 2015. Label switching and its solutions for frequentist mixture models. *Journal of Statistical Computation and Simulation*, **85**(5), 1000–1012.
- Yao, Weixin, & Lindsay, Bruce G. 2012. Bayesian mixture labeling by highest posterior density. *Journal of the American Statistical Association*.