The Robust EM-type Algorithms for Log-concave Mixtures of Regression Models

Hao Hu^{a,*}, Weixin Yao^b, Yichao Wu^a

^a*Department of Statistics, North Carolina State University, Raleigh, North Carolina 27695, U.S.A.*

^b*Department of Statistics, University of California, Riverside, California 92521, U.S.A.*

Abstract

Finite mixture of regression (FMR) models can be reformulated as incomplete data problems and they can be estimated via the expectation-maximization (EM) algorithm. The main drawback is the strong parametric assumption such as FMR models with normal distributed residuals. The estimation might be biased if the model is misspecified. To relax the parametric assumption about the component error densities, a new method is proposed to estimate the mixture regression parameters by only assuming that the components have log-concave error densities but the specific parametric family is unknown.

Two EM-type algorithms for the mixtures of regression models with log-concave error densities are proposed. Numerical studies are made to compare the performance of our algorithms with the normal mixture EM algorithms. When the component error densities are not normal, the new methods have much smaller MSEs when compared with the standard normal mixture EM algorithms. When the underlying component error densities are normal, the new methods have comparable performance to the normal EM algorithm.

Keywords: EM algorithm; Log-concave Maximum Likelihood Estimator; Mixture of Regression Model; Robust regression.

*Correspondence to: North Carolina State University, 5109 SAS Hall, 2311 Stinson Dr, Raleigh, NC, 27695, United States

Email addresses: hhu5@ncsu.edu (Hao Hu), weixin.yao@ucr.edu (Weixin Yao), wu@stat.ncsu.edu (Yichao Wu)

1. Introduction

Suppose we have n subjects where the measurement of observation i is a d -dimensional vector $\mathbf{x}_i = (x_{i1}, \dots, x_{id})^T$ for $i = 1, \dots, n$. Additionally, for a fixed finite integer k , $\mathbf{x}_i$ has a k -component mixture density of $f : \mathbb{R}^d \rightarrow \mathbb{R}$:

$$f(\mathbf{x}_i; \boldsymbol{\psi}) = \sum_{j=1}^k \lambda_j g_j(\mathbf{x}_i; \boldsymbol{\theta}_j), \quad (1.1)$$

where $\boldsymbol{\theta}_j \in \Theta_j \subseteq \mathbb{R}^{q_j}$ is the parameter corresponding to the component density g_j , λ_j 's are the mixing proportions, $\lambda_j \in (0, 1)$ for $j = 1, \dots, k$, $\sum_{j=1}^k \lambda_j = 1$, and $\boldsymbol{\psi} = (\lambda_1, \dots, \lambda_{k-1}, \boldsymbol{\theta}_1^T, \dots, \boldsymbol{\theta}_k^T)^T \in \mathbb{R}^{\sum_{j=1}^k q_j + k - 1}$. Model like (1.1) is called a finite mixture model, which provides a flexible methodology when the observations are from a number of classes with unknown class indicators.

Finite mixture models are widely used in econometrics, biology, genetics, and engineering; see, e.g. Frühwirth-Schnatter (2001), Grün & Hornik (2012), and Liang (2008) and Plataniotis (2000). For this reason, there is a rich history of studying mixture models both theoretically and practically. Everitt & Hand (1981), Lindsay (1995), and McLachlan & Peel (2000) provided great summaries of the theories, algorithms, and many technical details of mixture models.

When a random variable has a finite mixture density that depends on certain covariates, we obtain a finite mixture of regression (FMR) model. Suppose we observe univariate response y_i and p -dimensional covariate $\mathbf{x}_i$, the FMR model can be written as follows:

$$f(y_i | \mathbf{x}_i; \boldsymbol{\psi}) = \sum_{j=1}^k \lambda_j g_j(y_i - \mathbf{x}_i^T \boldsymbol{\beta}_j), \quad (1.2)$$

where $\boldsymbol{\beta}_j \in \mathbb{R}^p$, $\lambda_j \in (0, 1)$, $\sum_{j=1}^k \lambda_j = 1$, $\boldsymbol{\psi} = (\lambda_1, \dots, \lambda_{k-1}, \boldsymbol{\beta}_1^T, \dots, \boldsymbol{\beta}_k^T)^T \in \mathbb{R}^{kp + k - 1}$, and g_j is a parametric distribution function, such as normal, for j -th component, $j = 1, \dots, k$.

The parametric FMR model (1.2) can be estimated through the maximum likelihood estimators. As there are usually no explicit solutions to the unknown parameters, it is natural to reformulate the likelihood as an incomplete data problem and apply the expectation-maximization (EM) algorithm for the FMR models; see, e.g. Dempster

et al. (1977) and McLachlan & Krishnan (2007). Besides estimating the parameters in the FMR models, the EM algorithms also provide the probabilities that an observation belongs to certain classes. Consequently, FMR models can also be considered as
 30 unsupervised classification methods, even though clustering might not always be the goal.

Usually people assume that the densities g_j 's belong to certain parametric families, e.g. normal distribution. These parametric assumptions are often too strong and restrictive. There exist some previous works for Model (1.1) and (1.2) with non-normal
 35 component error densities g_j 's. Galimberti & Soffritti (2014), Song *et al.* (2014), Ingrassia *et al.* (2014), Punzo & McNicholas (2014), and Yao *et al.* (2014) discussed clustering and FMR model with heavy-tailed error distributions such as t or *Laplace* distributions. Liu & Lin (2014), Lin *et al.* (2007), Lin (2010), Zeller *et al.* (2011), and Lachos *et al.* (2011) explored the finite mixture models with skewed error distributions
 40 such as skewed-normal or skewed- t distribution. Verkuilen & Smithson (2012), Ingrassia *et al.* (2015), Punzo & Ingrassia (2016), and Bartolucci & Scaccia (2005) discussed the FMR model with other specific families such as beta or exponential distribution.

These previous works adjusted some certain model misspecification. However, most of the time, we are still not sure which parametric family we should apply, say
 45 error densities from logistic distribution vs *Laplace* distribution. Moreover, the parameter estimation may still be biased if the parametric model is misspecified. Another drawback is that each model requires a specific EM algorithm based on the parametric assumption. As a result, it would be valuable to have a universal EM algorithm for all, or at least some classes of the FMR models. Possible solutions include traditional
 50 nonparametric methods, e.g. Hunter & Young (2012) and Wu & Yao (2016), to adjust the parametric model mis-specification. These traditional nonparametric methods, e.g. kernel methods, bring new difficulties in selecting the tuning parameters.

To relax the parametric assumption, nonparametric shape constraints are becoming increasingly popular. In this paper, we make one shape constraint instead of a specific
 55 parametric assumption for each component density. We assume each component density g_j to be log-concave. A density $g(x)$ is log-concave if its log-density, $\phi(x) = \log g(x)$, is concave. Examples of log-concave densities include, but are not

limited to normal, Laplace, chi-square, logistic, gamma with shape parameter greater than 1, and beta distribution with both parameters greater than 1. Log-concave densities are unimodal but unimodal densities are not necessarily log-concave. Log-concave densities have many favorable properties as described by Balabdaoui *et al.* (2009). To estimate the log-density $\phi(x)$, Dümbgen *et al.* (2011) proposed an estimator by maximizing a log-likelihood-type functional:

$$L(\phi, Q) = \int \phi dQ - \int \exp\{\phi(x)\} dx + 1, \quad (1.3)$$

where $Q \in \mathcal{Q}$, $\mathcal{Q}$ is the family of all d -dimensional distributions, $\phi \in \Phi$ and Φ is the family of all concave function. For linear regression with log-concave error density, Dümbgen *et al.* (2011) proposed an estimator by maximizing:

$$\hat{L}(\phi, \beta, Q) = \frac{1}{n} \sum_{i=1}^n \phi(y_i - \mathbf{x}_i^T \beta) - \int \exp\{\phi(x)\} dx + 1. \quad (1.4)$$

Such estimators, like the maximizers of (1.3) and (1.4), are called log-concave maximum likelihood estimators (LCMLEs), which were studied by, for example, Dümbgen & Rufibach (2009), Cule *et al.* (2010), Cule & Samworth (2010), Chen & Samworth (2013), and Dümbgen *et al.* (2011). Dümbgen *et al.* (2011) proved the existence, uniqueness, and consistency of LCMLEs for (1.3) and (1.4) under fairly general conditions. These estimators provide more generality and flexibility without any tuning parameter. For log-concave mixture models, Chang & Walther (2007) proposed a log-concave EM-type algorithm for mixture density estimation, along with the application in clustering. Hu *et al.* (2016) further proposed the LCMLE, which is the maximizer of a log-likelihood type functional, and proved the existence and consistency for the LCMLE for the log-concave mixture models. To the best of our knowledge, none of the existing works has studied the log-concave FMR models as well as their computational algorithms. This paper aims to fill in this gap.

In this paper, we adopt the idea of log-concave density estimation and combine it with the FMR models. The identifiability of the proposed model has been established by Wang *et al.* (2012), Balabdaoui & Doss (2014), and Wu & Yao (2016). We propose two EM-type algorithms, which aim at adjusting the model misspecification. The remainder of this paper is organized as follows. We introduce the basic setup, model

85 details and notations in Section 2. We propose the EM-type algorithms for the log-concave mixtures of regression models in Section 3. Simulation studies and real data analysis are conducted in Sections 4 and 5. We end the paper with a short conclusion in Section 6.

2. Mixtures of Regression Models with Log-concave Error Densities

90 In this paper, we let Z be a latent variable with $\mathbb{P}(Z = j) = \lambda_j$, where $\lambda_j \in (0, 1)$, and $\sum_{j=1}^k \lambda_j = 1$ for $j = 1, \dots, k$. While given the latent variable $Z = j$, the response y_i has a linear relationship with $\mathbf{x}_i \in \mathbb{R}^p$:

$$y_i = \mathbf{x}_i^T \boldsymbol{\beta}_j + \epsilon_j, \quad (2.1)$$

where $\boldsymbol{\beta}_j = (\beta_{0,j}, \beta_{1,j}, \dots, \beta_{p-1,j})^T \in \mathbb{R}^p$ and ϵ_j is the error term with the distribution function g_j ($j = 1, \dots, k$). We assume that each component's error distribution g_j 95 is an unknown density function with the mean 0 for $j = 1, \dots, k$. If we do not assume a zero mean for g_j , $\boldsymbol{\beta}_j$ does not contain the intercept term accordingly. To relax the traditional parametric assumption about g_j , we only assume that g_j 's are log-concave, i.e. $\log g_j$ is concave for $j = 1, \dots, k$. We define $\boldsymbol{\theta}_j = (\lambda_j, \boldsymbol{\beta}_j^T)^T$ for $j = 1, \dots, k$ and $\boldsymbol{\theta} = (\boldsymbol{\theta}_1^T, \dots, \boldsymbol{\theta}_k^T)^T$. The likelihood function for the mixture of regressions model 100 can be presented as:

$$f(y_i | \mathbf{x}_i, \boldsymbol{\theta}, \mathbf{g}) = \sum_{j=1}^k \lambda_j g_j(y_i - \mathbf{x}_i^T \boldsymbol{\beta}_j), \quad (2.2)$$

where $\boldsymbol{\theta} \in \Theta = \{\boldsymbol{\theta} = (\boldsymbol{\theta}_1^T, \dots, \boldsymbol{\theta}_k^T)^T \mid \boldsymbol{\beta}_j \in \mathbb{R}^p, \lambda_j \in (0, 1), \sum_{j=1}^k \lambda_j = 1\} \subset \mathbb{R}^{kp+k-1}$.

Let $\mathbf{y} = (y_1, \dots, y_n)^T \in \mathbb{R}^n$, $\mathbf{x}_i = (1, x_{i,1}, \dots, x_{i,p-1})^T$ and $\mathbf{X} = (\mathbf{x}_1, \dots, \mathbf{x}_n)^T \in \mathbb{R}^{n \times p}$ be the n observations for the mixture of regressions model, where $n \gg kp +$ 105 $k - 1$. In order to estimate the model (2.2), it is natural to maximize the observed log-likelihood function:

$$\ell(\boldsymbol{\theta}, \mathbf{g} | \mathbf{X}, \mathbf{y}) = \sum_{i=1}^n \log \sum_{j=1}^k \lambda_j g_j(y_i - \mathbf{x}_i^T \boldsymbol{\beta}_j), \quad (2.3)$$

where $g_j(x) = \exp\{\phi_j(x)\}$ for some unknown concave function $\phi_j(x)$.

3. The EM-type Algorithms for Log-concave FMR Models

We define the missing value $\mathbf{Z} = (\mathbf{z}_1, \dots, \mathbf{z}_n)^T \in \mathbb{R}^{n \times k}$, where $\mathbf{z}_i = (z_{i1}, \dots, z_{ik})^T$
 110 $(i = 1, \dots, n)$ is a k -dimensional indicator vector with its j -th element given by

$$z_{ij} = \begin{cases} 1 & \text{if } (\mathbf{x}_i, y_i) \text{ belongs to } j\text{th group;} \\ 0 & \text{otherwise.} \end{cases}$$

Consequently, the complete log-likelihood for equation (2.3) is:

$$\ell_c(\boldsymbol{\theta}, \mathbf{g} | \mathbf{X}, \mathbf{y}, \mathbf{Z}) = \sum_{i=1}^n \sum_{j=1}^k z_{ij} \{\log \lambda_j + \log g_j(y_i - \mathbf{x}_i^T \boldsymbol{\beta}_j)\}. \quad (3.1)$$

In the E-step, given the current estimate $\boldsymbol{\theta}^{(t)}$ and $g_j^{(t)}$'s, we need to compute

$$Q(\boldsymbol{\theta}, \mathbf{g} | \boldsymbol{\theta}^{(t)}, \mathbf{g}^{(t)}, \mathbf{X}, \mathbf{y}) = \mathbb{E}\{\ell_c(\boldsymbol{\theta}, \mathbf{g} | \mathbf{X}, \mathbf{y}, \mathbf{Z}) \mid \boldsymbol{\theta}^{(t)}, \mathbf{g}^{(t)}, \mathbf{X}, \mathbf{y}\},$$

which is equivalent to computing

$$\begin{aligned} z_{ij}^{(t+1)} &= E(Z_{ij} | \boldsymbol{\theta}^{(t)}, \mathbf{g}^{(t)}, \mathbf{X}, \mathbf{y}) = Pr(Z_{ij} = 1 | \boldsymbol{\theta}^{(t)}, \mathbf{g}^{(t)}, \mathbf{X}, \mathbf{y}) \\ &= \frac{\lambda_j^{(t)} g_j^{(t)}(y_i - \mathbf{x}_i^T \boldsymbol{\beta}_j^{(t)})}{\sum_{h=1}^k \lambda_h^{(t)} g_h^{(t)}(y_i - \mathbf{x}_i^T \boldsymbol{\beta}_h^{(t)})}. \end{aligned} \quad (3.2)$$

In M-step, we need to maximize the following Q function:

$$\begin{aligned} Q(\boldsymbol{\theta}, \mathbf{g} | \boldsymbol{\theta}^{(t)}, \mathbf{X}, \mathbf{y}) &= \sum_{i=1}^n \sum_{j=1}^k z_{ij}^{(t+1)} \{\log \lambda_j + \log g_j(y_i - \mathbf{x}_i^T \boldsymbol{\beta}_j)\} \\ &= \sum_{i=1}^n \sum_{j=1}^k z_{ij}^{(t+1)} \log \lambda_j + \sum_{i=1}^n \sum_{j=1}^k z_{ij}^{(t+1)} \log g_j(y_i - \mathbf{x}_i^T \boldsymbol{\beta}_j). \end{aligned} \quad (3.3)$$

The first part of (3.3) is maximized by $\lambda_j^{(t+1)} = \frac{1}{n} \sum_{i=1}^n z_{ij}^{(t+1)}$, $j = 1, \dots, k$. However, for the second part, there is no explicit solution for $\boldsymbol{\beta}_j$'s and g_j 's. Consequently,
 115 we propose to alternatively update g_j 's and $\boldsymbol{\beta}_j$'s to maximize the second part of (3.3).

It is also well known that MLEs can be sensitive to outliers, see e.g. Yao *et al.* (2014) and García-Escudero *et al.* (2009). To overcome this problem, we further propose a robust technique, which adopts the idea of least trimmed squares (LTS), see

e.g. Rousseeuw (1985) for a detailed description of LTS. For each algorithm, when
 120 updating λ_j 's and β_j 's in the t -th iteration ($j = 1, \dots, k$), we drop s observations
 with the least log-likelihood values. In that way, we sacrifice some efficiency to gain
 the robustness to the outliers. The number s is the trimming tuning parameter, which
 satisfies $0 < s < n/2$. In this paper, we mainly use this trimmed idea to get a stable
 125 estimate of log-concave component error densities while enjoying its robustness when
 the component error densities are highly skewed or have heavy tails. Our empirical
 experience suggests that the choice of $s = \lfloor n/40 \rfloor$ works well. Note that a larger value
 of s would make our algorithms more robust if there are outliers in the data set and the
 sample size is not too small.

Our methodology is summarized as follows. First, we apply some stochastic se-
 130 arch strategy, which will be addressed later, to create the initial value for normal mix-
 tures of regression models from function `regmixEM` in R package `mixtools`, see
 Benaglia *et al.* (2009), until convergence. We treat the outcome of the normal mix-
 ture EM algorithm as the starting values for our EM-type algorithms, i.e. $\psi^{(0)} =$
 $(\hat{\lambda}_1^{(0)}, \dots, \hat{\lambda}_k^{(0)}, \hat{\beta}_1^{(0)T}, \dots, \hat{\beta}_k^{(0)T})^T$. The normal mixture of regressions model usually
 135 provides good initial values and our proposed EM algorithm will further improve the
 estimate if the error density is not normally distributed. The initial estimated density
 g_j can be obtained by the function `mlelcd` in R package `LogConcDEAD` (Cule *et al.*,
 2009).

First, we propose the Algorithm 3.1 for the case that all components have the same
 140 error density g .

Algorithm 3.1. *The EM-type algorithm when all g_j 's are the same, i.e. $g_j \equiv g$.*

*Initialize $\psi^{(0)}$ and $z_{ij}^{(0)}$ from normal mixture EM algorithm with equal variances
 and initialize the trimmed index subset of size $n - s$, denoted by $I^{(0)}$, which has the
 $n - s$ largest log-likelihoods. Initialize $g^{(0)}$ by the function `mlelcd` through fitted
 145 residuals $y_i - \mathbf{x}_i^T \beta_j^{(0)}$ with weights $z_{ij}^{(0)}$ for $i = 1, \dots, n$, $j = 1, \dots, k$.*

In t -th iteration, it consists of the following steps.

E-step: Given $\psi^{(t)}$ and $g^{(t)}$, we calculate

$$z_{ij}^{(t+1)} = E(Z_{ij} | \mathbf{X}, \mathbf{y}, \psi^{(t)}, g^{(t)}) = \frac{\lambda_j^{(t)} g^{(t)}(y_i - \mathbf{x}_i^T \boldsymbol{\beta}_j^{(t)})}{\sum_{h=1}^k \lambda_h^{(t)} g^{(t)}(y_i - \mathbf{x}_i^T \boldsymbol{\beta}_h^{(t)})}, \quad (3.4)$$

for $i = 1, \dots, n$, $j = 1, \dots, k$.

M-step:

(A) Calculate the log-likelihood value for each observation:

$$\ell_i^{(t)} = \ell(\mathbf{x}_i, y_i | g^{(t)}, \psi^{(t)}) = \log \sum_{j=1}^k \lambda_j^{(t)} g^{(t)}(y_i - \mathbf{x}_i^T \boldsymbol{\beta}_j^{(t)}),$$

150 from $i = 1, \dots, n$. Update the trimmed index subset of size $n - s$, denoted by $I^{(t+1)}$, which has the $n - s$ largest log-likelihoods.

(B) Update λ simply through

$$\lambda_j^{(t+1)} = \frac{1}{n - s} \sum_{i \in I^{(t+1)}} z_{ij}^{(t+1)}, j = 1, \dots, k. \quad (3.5)$$

(C) Update β :

$$\tilde{\boldsymbol{\beta}}_j^{(t+1)} \leftarrow \arg \max_{\boldsymbol{\beta}_j} \sum_{i \in I^{(t+1)}} z_{ij}^{(t+1)} \log g^{(t)}(y_i - \mathbf{x}_i^T \boldsymbol{\beta}_j), \quad j = 1, \dots, k. \quad (3.6)$$

(D) Shift the intercept of $\tilde{\boldsymbol{\beta}}_j^{(t+1)}$ so that the residuals have a zero mean.

$$\hat{\boldsymbol{\beta}}_j^{(t+1)} = (\hat{\beta}_{j,0}^{(t+1)}, \tilde{\beta}_{j,1}^{(t+1)}, \dots, \tilde{\beta}_{j,p-1}^{(t+1)}),$$

where

$$\hat{\beta}_{j,0}^{(t+1)} = \tilde{\beta}_{j,0}^{(t+1)} + c_j^{(t+1)} \quad \text{with} \quad c_j^{(t+1)} = \frac{1}{n - s} \sum_{i \in I^{(t+1)}} z_{ij}^{(t+1)} (y_i - \mathbf{x}_i^T \tilde{\boldsymbol{\beta}}_j^{(t+1)}),$$

for $j = 1, \dots, k$.

155 (E) Update g by:

$$g^{(t+1)} \leftarrow \arg \max_{g \in \mathcal{G}} \sum_{i=1}^n \sum_{j=1}^k z_{ij}^{(t+1)} \log g(y_i - \mathbf{x}_i^T \hat{\boldsymbol{\beta}}_j^{(t+1)}). \quad (3.7)$$

where $\mathcal{G}$ is the family of all log-concave densities.

In (3.6), β_j is updated through the function `optim` in R. The evaluation of $\log \hat{g}^{(t)}(y_i - \mathbf{x}_i^T \beta_j^{(t+1)})$ is calculated through the function `dlcd` in R package `LogConcDEAD`. In (3.7), the error density g is updated through the function called `mlelcd` in the R package `LogConcDEAD` through kn fitted residuals $y_i - \mathbf{x}_i^T \beta_j^{(t+1)}$ with weights $z_{ij}^{(t+1)}$, $i = 1, \dots, n, j = 1, \dots, k$. The algorithm is terminated if either t_{max} of iterations has been reached, or if $\ell^{(t+1)} - \ell^{(t)} < 10^{-8}$, where $\ell^{(t)} = \sum_{i \in I^{(t)}} \ell_{(i)}^{(t)}$ is the trimmed log-likelihood.

The algorithm usually converges after 10 iterations for $p = 2$ or 3. For each iteration, the most time consuming step is the (E) step for the density updating, which usually takes about 20 seconds for a sample size of 400. Consequently, the average computational time is about 3-5 minutes.

To avoid the local maximum, we follow the similar stochastic search strategy proposed by Dümmbgen *et al.* (2013). We restart the entire algorithm 20 times. For each restart, we randomly sample $\lfloor \alpha n \rfloor$ ($\alpha \in (0, 1)$) observations k times, fit k simple linear regressions, obtain the k groups of coefficients, and treat them as the starting values of β 's in the normal EM algorithm for k components. Additionally, we generate λ_j 's from a *Uniform*(0,1) distribution, scale them so that their sum is one, and treat them as the starting values of the mixing proportions in the normal EM algorithm. We then fit a normal FMR model, obtain the estimated coefficients, and use them as the initial values for our algorithm. We repeat this procedure 20 times and select the solution with the highest trimmed likelihood to avoid getting stuck in a local maximum.

The LCMLE $\hat{g}$ has been studied by Walther (2002) and Rufibach (2007). Here, we briefly summarize the results. Given i.i.d. data $X_1, \dots, X_n$ which follow a distribution g , the Log-concave Maximum Likelihood Estimator (LCMLE) $\hat{g}$ exists uniquely and has support on the convex hull of the dataset (by Theorem 2 of Cule *et al.* (2010)). In addition, $\log \hat{g}$ is a piecewise linear function whose knots are a subset of $\{X_1, \dots, X_n\}$. Walther (2002) and Rufibach (2007) provided algorithms for computing $\hat{g}(X_i)$, $i = 1, \dots, n$. The entire log-density $\log \hat{g}$ can be computed by linear interpolation between $\log \hat{g}(X_{(i)})$ and $\log \hat{g}(X_{(i+1)})$. Walther (2002) and Rufibach (2007) also pointed out that it is natural to apply weights in the density estimation step of the

EM-type algorithms. The $z_{ij}^{(t+1)}$'s can be viewed as weights for the kn fitted residuals $y_i - \mathbf{x}_i^T \boldsymbol{\beta}_j^{(t+1)}$ ($i = 1, \dots, n, j = 1, \dots, k$), while estimating the log-concave density g for M-step 3 in our algorithms.

190 A more general case is that the components' error terms do not share a common distribution, i.e. at least one g_j is different. Consequently, we propose the following Algorithm 3.2. The main difference is that, in Algorithm 3.2, each component density g_j is estimated by the iterative residuals only from the according component class, instead of being estimated by the entire residuals from all components in Algorithm 3.1.

Algorithm 3.2. *The EM-type algorithm when g_j 's are different.*

Initialize $\boldsymbol{\psi}^{(0)}$ and $z_{ij}^{(0)}$ from normal mixture EM algorithm with unequal variances and initialize the trimmed subset of size $n - s$, denoted by $I^{(0)}$, which has the $n - s$ largest log-likelihoods. For $j \in \{1, \dots, k\}$, initialize $g_j^{(0)}$ by the function `mlelcd` through fitted residuals $y_i - \mathbf{x}_i^T \boldsymbol{\beta}_j^{(0)}$ with weights $z_{ij}^{(0)}$ for $i = 1, \dots, n$.

In t -th iteration, it consists of the following steps.

E-step: Given $\boldsymbol{\psi}^{(t)}$ and $g^{(t)}$, we calculate

$$z_{ij}^{(t+1)} = E(Z_{ij} | \mathbf{X}, \mathbf{y}, \boldsymbol{\psi}^{(t)}, g^{(t)}) = \frac{\lambda_j^{(t)} g_j^{(t)} (y_i - \mathbf{x}_i^T \boldsymbol{\beta}_j^{(t)})}{\sum_{h=1}^k \lambda_h^{(t)} g_h^{(t)} (y_i - \mathbf{x}_i^T \boldsymbol{\beta}_h^{(t)})}, \quad (3.8)$$

for $i = 1, \dots, n, j = 1, \dots, k$.

M-step:

(A) Calculate the log-likelihood value for each observation:

$$\ell_i^{(t)} = \ell(\mathbf{x}_i, y_i | \mathbf{g}^{(t)}, \boldsymbol{\psi}^{(t)}) = \log \sum_{j=1}^k \lambda_j^{(t)} g_j^{(t)} (y_i - \mathbf{x}_i^T \boldsymbol{\beta}_j^{(t)}),$$

205 for $i = 1, \dots, n$. Update the trimmed subset of size $n - s$, denoted by $I^{(t+1)}$, which has the $n - s$ largest log-likelihoods.

(B) Update λ simply through

$$\lambda_j^{(t+1)} = \frac{1}{n - s} \sum_{i \in I^{(t+1)}} z_{ij}^{(t+1)}, j = 1, \dots, k. \quad (3.9)$$

(C) Update β :

$$\tilde{\beta}_j^{(t+1)} \leftarrow \arg \max_{\beta_j} \sum_{i \in I^{(t+1)}} z_{ij}^{(t+1)} \log g_j^{(t)}(y_i - \mathbf{x}_i^T \beta_j), \quad j = 1, \dots, k. \quad (3.10)$$

(D) Shift the intercept of $\tilde{\beta}_j^{(t+1)}$ so that the residuals have a zero mean.

$$\hat{\beta}_j^{(t+1)} = (\hat{\beta}_{j,0}^{(t+1)}, \tilde{\beta}_{j,1}^{(t+1)}, \dots, \tilde{\beta}_{j,p-1}^{(t+1)}),$$

where

$$\hat{\beta}_{j,0}^{(t+1)} = \tilde{\beta}_{j,0}^{(t+1)} + c_j^{(t+1)} \quad \text{with} \quad c_j^{(t+1)} = \frac{1}{n-s} \sum_{i \in I^{(t+1)}} z_{ij}^{(t+1)} (y_i - \mathbf{x}_i^T \tilde{\beta}_j^{(t+1)}),$$

210 for $j = 1, \dots, k$.

(E) Update g_j by:

$$g_j^{(t+1)} \leftarrow \arg \max_{g_j \in \mathcal{G}} \sum_{i=1}^n z_{ij}^{(t+1)} \log g_j(y_i - \mathbf{x}_i^T \hat{\beta}_j^{(t+1)}). \quad (3.11)$$

for $j = 1, \dots, k$, where $\mathcal{G}$ is the family of all log-concave densities.

In (3.11), the j -th component density g_j is updated through the function called `mlelcd` in the R package `LogConcDEAD` through n fitted residuals $y_i - \mathbf{x}_i^T \hat{\beta}_j^{(t+1)}$ with weights $z_{ij}^{(t+1)}$, $i = 1, \dots, n$ for $j \in \{1, \dots, k\}$. The algorithm is terminated if
 215 either the maximum number of iterations t_{max} has been reached, or if $\ell^{(t+1)} - \ell^{(t)} < 10^{-8}$, where $\ell^{(t)} = \sum_{i \in I^{(t)}} \ell_i^{(t)}$ is the trimmed log-likelihood for t -th iteration.

4. Numerical Experiments

4.1. Simulation setup

220 In this section, we study the performance of our EM-type algorithms and compare them with the according EM algorithms for the normal FMR models. For the convenience purposes, in the following text and tables, we denote Algorithm 3.1 as “LCD-EM1” and compare it with the normal EM algorithm with equal variance and similar trimming techniques, denoted as “Normal-EM1”. We also denote Algorithm 3.2

225 as “LCD-EM2” and compare it with the normal EM algorithm with unequal variance and similar trimming techniques, denoted as “Normal-EM2”.

We generate data from 2-component log-concave FMR models:

$$y_i = \begin{cases} \beta_1^T \mathbf{x}_i + e_{i,1} & \text{with probability } \lambda; \\ \beta_2^T \mathbf{x}_i + e_{i,2} & \text{with probability } 1 - \lambda. \end{cases} \quad (4.1)$$

For Model I through Model VI, we set $\mathbf{x}_i = (1, x_{1,i})^T$, where $x_{1,i}$ ’s are independently generated from $Uniform(-1, 3)$. We let $\lambda = 0.3$, $\beta_1 = (\beta_{0,1}, \beta_{1,1})^T =$
 230 $(0, 2)^T$, and $\beta_2 = (\beta_{0,2}, \beta_{1,2})^T = (-2, 5)^T$. For Model VII and Model VIII, we set $\mathbf{x}_i = (1, x_{1,i}, x_{i,2})^T$, where $x_{1,i}$ ’s and $x_{i,2}$ ’s are both independently generated from $Uniform(-1, 3)$. We let $\lambda = 0.3$, $\beta_1 = (\beta_{0,1}, \beta_{1,1}, \beta_{2,1})^T = (0, 2, 1)^T$, and $\beta_2 = (\beta_{0,2}, \beta_{1,2}, \beta_{2,2})^T = (-2, 5, 3)^T$. We let $e_{i,1} = e_{i,2} \equiv e_i$, where e_i ’s are independently and identically generated based on the parametric form from Table 1. For all
 235 eight models, we generate data for a finite sample size of $n = 400$.

Table 1: The error densities for Model I to Model VIII and the summary of the according features.

	e_i ’s distribution	log-concave	symmetric
Model I	Standard Normal: $N(0, 1)$	Yes	Yes
Model II/VII	Centered Beta: $3(Beta(1, 2) - 1/3)$	Yes	No
Model III/VIII	Centered Exponential: $Exp(2) - 2$	Yes	No
Model IV	Standard Laplace: $Laplace(0, 1)$	Yes	Yes
Model V	Centered Beta: $4(Beta(0.25, 0.75) - 1/4)$	No	No
Model VI	Centered t: t_4	No	Yes

For Model IX to Model XI, we let $\mathbf{x}_i = (1, x_{1,i})^T$, where $x_{1,i}$ ’s are independently generated as $Uniform(-1, 3)$. We set $n = 400$, $\lambda = 0.4$, $\beta_1 = (\beta_{0,1}, \beta_{1,1})^T = (0, 1)^T$, and $\beta_2 = (\beta_{0,2}, \beta_{1,2})^T = (-3, 4)^T$. The component error densities are generated based on the parametric form of Model IX to Model XI in Table 2.

240 For all nine models, we repeat the simulation $N = 200$ times. We visualize the generated data of Model III and IX for a single replicate in Figure 1. For both replicates, our proposed algorithm has monotone increasing log-likelihood.

Table 2: The error densities for Model IX, X, and XI.

	$e_{i,1}$'s distribution	$e_{i,2}$'s distribution
Model IX	$N(0, 1)$	$N(0, 0.25)$
Model X	$3(\text{Beta}(1, 2) - 1/3)$	$N(0, 0.25)$
Model XI	$\frac{2}{3}\text{Laplace}(0, 1)$	$\frac{2}{3}(\text{Exp}(2) - 2)$

(a) Model III: Generated Data

(b) Model XI: Generated Data

(c) Model III: Monotone Increasing likelihood

(d) Model XI: Monotone Increasing likelihood

Figure 1: Generated data for Model III and XI (green and red line represent the true coefficients for the two components) and the monotone increasing log-likelihood value in each iteration.

It is also valuable to compare our proposed algorithm with some other parametric EM algorithms that are proposed in the literature. In this study, we select two popular ones, EM algorithm for *Laplace*-FMR models (denoted as “*Laplace*-EM”) and EM algorithm for *t*-FMR models (denoted as “*t*-EM”). We compare the same criteria in Model IV and Model VI.

There is a well-known *label switching* issue when sorting the labels for mixture

models (Stephens, 2000; Yao & Lindsay, 2009). In this paper, we adopt the method
of Yao (2015) to find the labels by minimizing the distance between the estimated
classification probabilities and the true labels over different permutations. After sorting
the labels, we compute the MSE of all parameters over the N replicates, i.e. $MSE =$
 $N^{-1} \sum_{h=1}^N (\hat{\theta}_h - \theta_0)^2$, where $\hat{\theta}_h = (\hat{\beta}_1^T, \hat{\beta}_2^T, \hat{\lambda}_h^T)^T$ is the vector of parameter estimates
of the h -th replicate and θ_0 is the true value for the vector of the parameters. As the
mixtures of regression models serve as a methodology for classification, we compute
the average of misclassification numbers (AMN) as well.

4.2. Selecting the trimming constant s and the stochastic search proportion α

One important key step for both algorithms is to select the appropriate trimming
constant s and the stochastic search proportion α . Typically s is a relatively small
positive constant. If s is too small (approaching zero), the algorithms do not have
enough robustness powers against the outliers. If s is too large, we scarify too much
on the efficiency. Choosing the trimming tuning parameters adaptively is not easy and
has long been a challenging problem. One option is to choose s using the graphical
way suggested by Neykov *et al.* (2007). In practice, we use $s = 0.01$ - 0.05 . In our
later simulation example, we choose $s = 0.025$, which means we drop 2.5% of the
observations while updating the parameter. Table 3 shows the simulation results of
 MSE 's and AMN 's over $N = 200$ replicates for Model III with different trimming size.
We observe that the trimming size between 0.01 or 0.05 is appropriate.

Table 3: Simulation results for Model III with different trimming size.

Model	Method	$\beta_{0,1}$	$\beta_{1,1}$	$\beta_{0,2}$	$\beta_{1,2}$	λ	AMN
III	LCD-EM1(s=0)	0.01900	0.05215	0.02505	0.01878	0.00500	48.15
	LCD-EM1(s=0.01)	0.01770	0.05215	0.02474	0.01534	0.00307	48.02
	LCD-EM1(s=0.025)	0.01095	0.02746	0.02039	0.01676	0.00304	47.49
	LCD-EM1 (s=0.05)	0.01010	0.01910	0.02778	0.02018	0.00359	51.60
	LCD-EM1 (s=0.10)	0.01691	0.02437	0.02855	0.03010	0.00313	51.45

On the other hand, the selection of α is not that sensitive as long as $\alpha \leq 0.5$. We
 270 take Model II as an example. In Figure 2, we plot the $R(\beta)$ vs different α 's, where
 $R(\beta) = N^{-1} \sum_{h=1}^N \|\hat{\beta}_h - \beta_0\|^2$, β_0 is the vector of true coefficient values, and $\hat{\beta}_h$
 is the estimator for replicate h . We observe that $R(\beta)$ are almost at the same level for
 $\alpha \in (0, 0.5)$. As α approaching 1, though not very frequently, the algorithm is more
 likely to get stuck in some local maximum estimator instead of the local maximum,
 275 as we are repeatedly using the same starting value. In practice, we choose $\alpha = 0.10$,
 which usually works very well.

Figure 2: Select Stochastic Search Proportion α

4.3. Simulation results

Table 4 displays the *MSEs* of parameter estimates (the value of Model IV and
 Model VI are multiplied by 10) and the average of misclassification numbers over
 280 $N = 200$ simulations for Algorithm 3.1. For the density which is not normally distri-
 buted, even is not log-concave (Model V and VI), Algorithm 3.1 demonstrates signifi-
 cant improvement over the traditional normal mixture EM algorithm in terms of much
 smaller *MSE*. Especially for Model II, III and V, many *MSEs* from LCD-EM1 are 30%
 less than those from the Normal-EM1. This phenomena is still true when we increase
 285 the dimensionality. Table 5 display the simulation results for $p = 3$. For Model VII
 and VIII, we still observe much smaller *MSEs* for LCD-EM1 when comparing with
 Normal-EM1.

Table 4: Simulation results for Model I-VI.

Model	Method	$\beta_{0,1}$	$\beta_{1,1}$	$\beta_{0,2}$	$\beta_{1,2}$	λ	AMN
I	LCD-EM1	0.03096	0.01028	0.00874	0.00380	0.00081	41.55
	Normal-EM1	0.02671	0.00992	0.00819	0.00348	0.00072	41.43
II	LCD-EM1	0.00847	0.00273	0.00273	0.00051	0.00101	26.95
	Normal-EM1	0.01310	0.00341	0.00416	0.00134	0.00671	29.68
III	LCD-EM1	0.01095	0.02746	0.02039	0.01676	0.00304	47.49
	Normal-EM1	0.14997	0.04237	0.038357	0.03090	0.00402	62.17
IV	LCD-EM1	0.03794	0.01526	0.01304	0.00475	0.00152	54.25
	Normal-EM1	0.05371	0.01558	0.01538	0.00686	0.00146	55.86
	<i>Laplace</i> -EM	0.03314	0.02184	0.01404	0.00354	0.00102	54.08
V	LCD-EM1	0.01639	0.00317	0.00695	0.00031	0.00113	33.13
	Normal-EM1	0.05458	0.01504	0.02268	0.00480	0.00121	51.66
VI	LCD-EM1	0.01021	0.02783	0.01654	0.01300	0.00245	53.45
	Normal-EM1	0.01304	0.03233	0.02011	0.01813	0.00236	54.91
	<i>t</i> -EM	0.01421	0.01367	0.00816	0.01416	0.00257	52.08

When the error density truly comes from the specific parametric family, the new algorithm still has comparable performance. The $MSEs$ of LCD-EM1 is almost the same or only slightly worse than the according parametric EM algorithms for Model I/IV/VI. Notice that in most cases, we are unsure about the component densities and which parametric EM algorithm we should apply. In that case, our proposed algorithm shows great flexibility. After fitting this EM algorithm, one could characterize the component densities quite well and can further select the appropriate parametric EM algorithm to get further precisions.

To show the performance of Algorithm 3.2, we report the result over 200 replicates and compare the same criteria as we did for (4.1). Similar phenomena (shown in Table

Table 5: Simulation results for Model VII-VIII.

Model	Method	$\beta_{0,1}$	$\beta_{1,1}$	$\beta_{2,1}$	λ
VII	LCD-EM1	0.00001	0.00046	0.00224	0.00040
	Normal-EM1	0.00018	0.00589	0.00452	0.00158
VIII	LCD-EM1	0.04092	0.00027	0.01603	0.00037
	Normal-EM1	0.19841	0.00022	0.09888	0.00040
Model	Method	$\beta_{0,2}$	$\beta_{1,2}$	$\beta_{2,2}$	AMN
VII	LCD-EM1	0.00163	0.00062	0.00051	22.32
	Normal-EM1	0.01002	0.00139	0.00247	24.38
VIII	LCD-EM1	0.00939	0.00009	0.00001	36.29
	Normal-EM1	0.03050	0.00440	0.00436	47.90

6) are observed for Algorithm 3.2. For the component that truly comes from normal distribution (Model IX and Component 2 of Model X), our proposed algorithm has
300 comparable performance to the normal EM algorithm with unequal variances and a similar trimming technique. For the components which are misspecified (Model XI and component 1 of Model X), potential improvements are gained if we apply LCD-EM2 instead of the Normal-EM2.

To further illustrate the performance of LCMLE for a single replicate, Figure 3a
305 shows the fitted LCMLE for a single replicate of Model III's simulation. The fitted error density by Algorithm 3.1 (green dashed line) approximates the true density (red solid line) well, even under a finite sample size of 400. Similar phenomena holds for Algorithm 3.2. The fitted log-concave error densities for the two components (red and green dashed lines) approximate the true densities (red and green solid line) for both
310 two components well under a finite sample size of 400 for Model XI.

Table 6: Simulation results for Model IX-XI.

Model	Method	$\beta_{0,1}$	$\beta_{1,1}$	$\beta_{0,2}$	$\beta_{1,2}$	λ	AMN
IX	LCD-EM2	0.01473	0.00700	0.00187	0.00088	0.00117	30.21
	Normal-EM2	0.01390	0.00551	0.00175	0.00081	0.00087	29.88
X	LCD-EM2	0.00543	0.00122	0.00199	0.00080	0.00084	26.97
	Normal-EM2	0.00633	0.00266	0.00183	0.00073	0.00075	27.89
XI	LCD-EM2	0.00658	0.00436	0.03836	0.00004	0.00022	49.20
	Normal-EM2	0.01943	0.01671	0.08099	0.00004	0.00185	61.28

(a) Model III: fitted LCMLE vs true density (b) Model XI: fitted LCMLEs vs true densities

Figure 3: Fitted LCMLE for Model III and XI vs the true densities.

4.4. Classification results

One important feature of the FMR model is that it serves as a tool of unsuper-
 315 vised learning. Consequently, we compare the average number of misclassifications
 (AMN's) among the 200 replicates in Table 4 and Table 6. For Model I and Model
 VII, the average misclassification numbers for our EM-type algorithms are almost the
 same or only a little bit higher than the normal EM algorithm with similar trimming
 techniques. When the models are misspecified, the average misclassification numbers
 320 obtained from log-concave FMRs are smaller than those from the normal mixture EM
 algorithm with similar trimming techniques.

To further illustrate the classification result, we show the classification results for

every replicate in Model I, III, IX and XI. In Figure 4, each point represents a single replicate. The x -axis represents the number of misclassifications by normal mixture EM algorithm. The y -axis represents the number of misclassifications by our log-concave mixture EM algorithm. We observe significant improvement in the sense of misclassification rates when the models are misspecified (in Figure 4b and 4d, the majority of points are under the identical line). When the component error densities are indeed normal, we observe no significant penalties if we apply the log-concave EM algorithm (Figure 4a and 4c).

4.5. Robustness to outliers

We artificially create Model XII, which has the setup of 4.1 with $\mathbf{x}_i = (1, x_{1,i})^T$, where $x_{1,i}$'s are independently generated as $Uniform(-1, 3)$. We set $\lambda = 0.3$, $\beta_1 = (\beta_{0,1}, \beta_{1,1})^T = (0, 2)^T$, and $\beta_2 = (\beta_{0,2}, \beta_{1,2})^T = (-1, 2)^T$. We let e_i as $Laplace(0, 1)$ and artificially replace 10 observations (2.5%) with the extreme outliers. We generated five y values at $x = -1$ from a $Uniform(-15, -10)$. We also generated another five y values at $x = 2$ from a $Uniform(20, 25)$.

Table 7: Simulation results for Model XII.

Model	Method	$\beta_{0,1}$	$\beta_{1,1}$	$\beta_{0,2}$	$\beta_{1,2}$	λ	AMN
XII	LCD-EM1(s=0)	0.12227	0.11354	0.02471	0.00086	0.00019	54.05
	LCD-EM1(s=0.01)	0.08403	0.01152	0.02666	0.00084	0.00009	53.80
	LCD-EM1 (s=0.025)	0.03180	0.00013	0.02359	0.00047	0.00001	53.35
	LCD-EM1 (s=0.05)	0.05297	0.00427	0.02595	0.00043	0.00015	53.35
	Normal-EM1 (s=0)	0.16953	0.22098	0.20755	0.17174	0.00423	66.35
	Normal-EM1 (s=0.025)	0.14019	0.04473	0.05549	0.16231	0.00459	65.35
	Normal-EM1 (s=0.05)	0.03666	0.02956	0.09437	0.02881	0.00348	64.75

We report the similar criteria in Table 7. We observe that combining log-concave EM algorithm with trimming has the best robustness against the non-normal distributed density and outliers.

Figure 4: Numbers of misclassifications: normal mixture EM algorithm vs log-concave mixture EM algorithm for mixtures of regression models. The solid lines represents the identity.

5. Data Analysis

The tone dataset (from package `mixtools`) contains 150 trials from the same musician; see Cohen (1980) for a detailed description. In each trial, a fundamental tone, which was purely determined by a stretching ratio, was first provided to the musician. Then the musician tuned the tone one octave above. The tuning ratio, which was measured as the adjusted tone divided by the fundamental tone, was recorded. The purpose

of this experiment was to demonstrate the “two musical perception theory”. We give the scatter plot of the data in Figure 5.

Figure 5: Tone data from the tone perception study of Cohen (1980).

For the entire dataset, by applying Algorithm LCD-EM1 with $k = 2$, we obtain the fitted coefficients (shown as the solid lines in Figure 5) and the fitted log-likelihood value. We refit the data with Algorithm Normal-EM1, and report the same criteria as we did for Algorithm 3.1.

To further demonstrate the prediction power of Algorithms 3.1, we apply a 10-folder cross validation to the data set. Denote the full data set as $\mathcal{D}$. We randomly partition $\mathcal{D}$ into a training set $\mathcal{R}_h$ and a testing set $\mathcal{T}_h$ with the property $\mathcal{D} = \mathcal{R}_h + \mathcal{T}_h$ for $h = 1, \dots, H$, where $H = 10$. For each folder $h \in 1, \dots, H$, we estimated the parameters $\hat{\lambda}_j^h$'s and $\hat{\beta}_j^h$'s, as well as the estimated log-concave density g^h through the training set $\mathcal{R}_h$. We then calculate the following two types of mean square errors:

- $E_1 = H^{-1} \sum_{h=1}^H \sum_{i \in \mathcal{T}_h} \sum_{j=1}^k \hat{p}_{ij}^h \{(y_i - \mathbf{x}_i^T \hat{\beta}_j^h)^2\};$
- $E_2 = H^{-1} \sum_{h=1}^H \sum_{i \in \mathcal{T}_h} \min_j \{(y_i - \mathbf{x}_i^T \hat{\beta}_j^h)^2\};$

where $\hat{p}_{ij}^h$ is the estimated probability that (x_i, y_i) is from j -th component for folder h :

$$\hat{p}_{ij}^h = \frac{\hat{\lambda}_j^h g^h(y_i - \mathbf{x}_i^T \hat{\boldsymbol{\beta}}_j^h)}{\sum_{m=1}^k \hat{\lambda}_m^h g^h(y_i - \mathbf{x}_i^T \hat{\boldsymbol{\beta}}_m^h)},$$

360 for $i \in \mathcal{T}_h$ and $j \in \{1, \dots, k\}$.

We report the same criteria based on the coefficients obtained by the Normal-EM1 algorithm. The results of fitting the log-concave FMR model and the normal FMR model are summarized in Table 8. The fitted result obtained by LCD-EM1 has a much larger log-likelihood. Additionally, Algorithm 3.1 provides much smaller mean square
 365 errors for both E_1 and E_2 , which indicates that our proposed algorithm predicts the response more precisely than the traditional normal EM algorithm.

Table 8: Estimated parameters and other characteristics of LCD-EM algorithm and Normal-EM algorithm for the tone dataset.

	LCD-EM1		Normal-EM1	
	Comp 1	Comp 2	Comp 1	Comp 2
β_0	-0.0143	1.9488	-0.0388	1.8924
β_1	0.9968	0.0263	0.9989	0.0559
λ	0.4253	0.5747	0.3256	0.6744
ℓ	170.91		158.54	
E_1	0.0039		0.0105	
E_2	0.0033		0.0041	

6. Conclusion and Discussion

This paper proposed two robust EM-type algorithms for the log-concave mixtures of regression models. These algorithms provide more flexibility, which allows a large
 370 family of error densities in the mixtures of regression models. By estimating the log-concave error density in every M-step of our algorithms, the log-concave maximum likelihood estimator corrects the model misspecification, e.g. adjusting skewness and heavy tails when the error distribution is not normal, in a nonparametric way without specifying the families of error densities.

375 Through numerical studies, our proposed algorithms have better performances than
the parametric EM algorithms whose parametric families are misspecified. We also
observe no significant penalties for applying our proposed algorithms instead of the
according EM algorithms which correctly characterize the error densities in the FMR
model.

380 Future work includes, but is not limited to the theoretical investigation of consistency
and convergence properties for the log-concave FMR models, as an extension of
Section 3 of Dümbgen *et al.* (2011). It would also be a challenging task to prove the
ascending properties for these nonparametric EM algorithm.

Acknowledgment

385 Hu's research is partially supported by National Institutes of Health grant R01-
CA149569. Yao's research is supported by NSF grant DMS-1461677. Wu's research is
partially supported by National Institutes of Health grant R01-CA149569 and National
Science Foundation grant DMS-1055210.

References

- 390 Balabdaoui, Fadoua, & Doss, Charles R. 2014. Inference for a mixture of symmetric
distributions under log-concavity. *arXiv preprint:1411.4708*.
- Balabdaoui, Fadoua, Rufibach, Kaspar, & Wellner, Jon A. 2009. Limit distribution
theory for maximum likelihood estimation of a log-concave density. *The Annals of
Statistics*, **37**(3), 1299–1331.
- 395 Bartolucci, Francesco, & Scaccia, Luisa. 2005. The use of mixtures for dealing with
non-normal regression errors. *Computational Statistics & Data Analysis*, **48**(4),
821–834.
- Benaglia, Tatiana, Chauveau, Didier, Hunter, David, & Young, Derek. 2009. **mixtools**:
An R package for analyzing finite mixture models. *Journal of Statistical Software*,
400 **32**(6), 1–29.

- Chang, George T, & Walther, Guenther. 2007. Clustering with mixtures of log-concave distributions. *Computational Statistics & Data Analysis*, **51**(12), 6242–6251.
- Chen, Yining, & Samworth, Richard J. 2013. Smoothed log-concave maximum likelihood estimation with applications. *Statistica Sinica*, **23**(3), 1373–1398.
- 405 Cohen, Elizabeth A. 1980. Inharmonic tone perception. *Unpublished Ph. D. Dissertation, Stanford University*.
- Cule, Madeleine, & Samworth, Richard. 2010. Theoretical properties of the log-concave maximum likelihood estimator of a multidimensional density. *Electronic Journal of Statistics*, **4**, 254–270.
- 410 Cule, Madeleine, Gramacy, Robert, & Samworth, Richard. 2009. LogConcDEAD: An R package for maximum likelihood estimation of a multivariate log-concave density. *Journal of Statistical Software*, **29**(2), 1–20.
- Cule, Madeleine, Samworth, Richard, & Stewart, Michael. 2010. Maximum likelihood estimation of a multi-dimensional log-concave density. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, **72**(5), 545–607.
- 415 Dempster, Arthur P, Laird, Nan M, & Rubin, Donald B. 1977. Maximum likelihood from incomplete data via the EM algorithm. *Journal of the Royal Statistical Society. Series B (Statistical Methodological)*, 1–38.
- Dümbgen, Lutz, & Rufibach, Kaspar. 2009. Maximum likelihood estimation of a log-concave density and its distribution function: Basic properties and uniform consistency. *Bernoulli*, **15**(1), 40–68.
- 420 Dümbgen, Lutz, Samworth, Richard, & Schuhmacher, Dominic. 2011. Approximation by log-concave distributions, with applications to regression. *The Annals of Statistics*, **39**(2), 702–730.
- 425 Dümbgen, Lutz, Samworth, Richard J, & Schuhmacher, Dominic. 2013. Stochastic search for semiparametric linear regression models. *Pages 78–90 of: From Probability to Statistics and Back: High-Dimensional Models and Processes—A Festschrift in Honor of Jon A. Wellner*. Institute of Mathematical Statistics.

- Everitt, Brian S, & Hand, David J. 1981. *Finite mixture distributions*. Vol. 9. Chapman
430 and Hall London.
- Frühwirth-Schnatter, Sylvia. 2001. Markov chain Monte Carlo estimation of classical
and dynamic switching and mixture models. *Journal of the American Statistical
Association*, **96**(453), 194–209.
- Galimberti, Giuliano, & Soffritti, Gabriele. 2014. A multivariate linear regression ana-
435 lysis using finite mixtures of t distributions. *Computational Statistics & Data Ana-
lysis*, **71**, 138–150.
- García-Escudero, Luis Angel, Gordaliza, Alfonso, San Martin, Roberto, Van Aelst,
Stefan, & Zamar, Ruben. 2009. Robust linear clustering. *Journal of the Royal
Statistical Society: Series B (Statistical Methodology)*, **71**(1), 301–318.
- 440 Grün, Bettina, & Hornik, Kurt. 2012. Modelling human immunodeficiency virus ribo-
nucleic acid levels with finite mixtures for censored longitudinal data. *Journal of the
Royal Statistical Society: Series C (Applied Statistics)*, **61**(2), 201–218.
- Hu, Hao, Wu, Yichao, & Yao, Weixin. 2016. Maximum likelihood estimation of the
mixture of log-concave densities. *Computational Statistics & Data Analysis*, **101**,
445 137–147.
- Hunter, David R, & Young, Derek S. 2012. Semiparametric mixtures of regressions.
Journal of Nonparametric Statistics, **24**(1), 19–38.
- Ingrassia, Salvatore, Minotti, Simona C, & Punzo, Antonio. 2014. Model-based cluste-
ring via linear cluster-weighted models. *Computational Statistics & Data Analysis*,
450 **71**, 159–182.
- Ingrassia, Salvatore, Punzo, Antonio, Vittadini, Giorgio, & Minotti, Simona C. 2015.
The generalized linear mixed cluster-weighted model. *Journal of Classification*,
32(1), 85–113.
- Lachos, Victor H, Bandyopadhyay, Dipankar, & Garay, Aldo M. 2011. Heteroscedastic
455 nonlinear regression models based on scale mixtures of skew-normal distributions.
Statistics & Probability Letters, **81**(8), 1208–1217.

- Liang, Faming. 2008. Clustering gene expression profiles using mixture model ensemble averaging approach. *JP Journal of Biostatistics*, **2**, 57–80.
- Lin, Tsung-I. 2010. Robust mixture modeling using multivariate skew t distributions. *Statistics and Computing*, **20**(3), 343–356.
- Lin, Tsung-I, Lee, Jack C, & Yen, Shu Y. 2007. Finite mixture modelling using the skew normal distribution. *Statistica Sinica*, **17**(3), 909–927.
- Lindsay, Bruce G. 1995. Mixture models: theory, geometry and applications. *Pages i–163 of: NSF-CBMS Regional Conference Series in Probability and Statistics*, vol. 5. JSTOR.
- Liu, Min, & Lin, Tsung-I. 2014. A skew-normal mixture regression model. *Educational and Psychological Measurement*, **74**(1), 139–162.
- McLachlan, Geoffrey, & Krishnan, Thriyambakam. 2007. *The EM algorithm and extensions*. Vol. 382. John Wiley & Sons.
- McLachlan, Geoffrey, & Peel, David. 2000. *Finite mixture models*. John Wiley & Sons.
- Neykov, Neyko, Filzmoser, Peter, Dimova, R, & Neytchev, Plamen. 2007. Robust fitting of mixtures using the trimmed likelihood estimator. *Computational Statistics & Data Analysis*, **52**(1), 299–308.
- Plataniotis, Kostantinos N. 2000. Gaussian mixtures and their applications to signal processing. *Advanced Signal Processing Handbook: Theory and Implementation for Radar, Sonar, and Medical Imaging Real Time Systems*.
- Punzo, Antonio, & Ingrassia, Salvatore. 2016. Clustering bivariate mixed-type data via the cluster-weighted model. *Computational Statistics*, **31**(3), 989–1013.
- Punzo, Antonio, & McNicholas, Paul D. 2014. Robust clustering in regression analysis via the contaminated Gaussian cluster-weighted model. *arXiv preprint:1409.6019*.

- Rousseeuw, Peter J. 1985. Multivariate estimation with high breakdown point. *Mathematical Statistics and Applications*, **8**, 283–297.
- Rufibach, Kaspar. 2007. Computing maximum likelihood estimators of a log-concave density function. *Journal of Statistical Computation and Simulation*, **77**(7), 561–574.
- Song, Weixing, Yao, Weixin, & Xing, Yanru. 2014. Robust mixture regression model fitting by Laplace distribution. *Computational Statistics & Data Analysis*, **71**, 128–137.
- Stephens, Matthew. 2000. Dealing with label switching in mixture models. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, **62**(4), 795–809.
- Verkuilen, Jay, & Smithson, Michael. 2012. Mixed and mixture regression models for continuous bounded responses using the beta distribution. *Journal of Educational and Behavioral Statistics*, **37**(1), 82–113.
- Walther, Guenther. 2002. Detecting the presence of mixing with multiscale maximum likelihood. *Journal of the American Statistical Association*, **97**(458), 508–513.
- Wang, Shaoli, Yao, Weixin, & Hunter, David. 2012. Mixtures of linear regression models with unknown error density. *Unpublished manuscript*.
- Wu, Qiang, & Yao, Weixin. 2016. Mixtures of quantile regressions. *Computational Statistics & Data Analysis*, **93**, 162–176.
- Yao, Weixin. 2015. Label switching and its solutions for frequentist mixture models. *Journal of Statistical Computation and Simulation*, **85**(5), 1000–1012.
- Yao, Weixin, & Lindsay, Bruce G. 2009. Bayesian mixture labeling by highest posterior density. *Journal of the American Statistical Association*, **104**(486), 758–767.
- Yao, Weixin, Wei, Yan, & Yu, Chun. 2014. Robust mixture regression using the t-distribution. *Computational Statistics & Data Analysis*, **71**, 116–127.

Zeller, Camil B, Lachos, Vctor H, & Vilca-Labra, Filidor E. 2011. Local influence analysis for regression models with scale mixtures of skew-normal distributions. *Journal of Applied Statistics*, **38**(2), 343–368.