1 Fos Expression in the Olfactory Pathway of High- and Low-Sexually Performing
2 Rams Exposed to Urine from Estrous or Ovariectomized Ewes
3 AJ Mirto^a, KJ Austin^a, VA Uthlaut^a, CE Roselli^b, BM Alexander^a.

4

5 ^aUniversity of Wyoming, Laramie, WY
6 Department of Animal Science
7 Dept 3684
8 1000 E. University Ave.
9 Laramie, WY 82071 USA
10 amirto@uwyo.edu
11 kaustin@uwyo.edu
12 balex@uwyo.edu

13
14 ^bOregon Health and Science University, Portland, OR
15 Department of Physiology & Pharmacology
16 3181 SW Sam Jackson Park Rd.
17 Portland, OR 97239 USA
18 rosellc@ohsu.edu

19
20 **Corresponding Author:**

21 Brenda Alexander
22 Email: balex@uwyo.edu
23 Mobile: (307) 760-0331
24 Department of Animal Science
25 Dept 3684
26 1000 E. University Ave.
27 Laramie, WY 82071 USA

28
29

30

31

32

33

34

35

36

37 **ABSTRACT**

38 Exposure to estrous ewe urine stimulates investigation and mounting activity in
39 sexually active but not sexually inactive rams. It was hypothesized sexual
40 indifference may result from an inability to detect olfactory cues or an interruption
41 of the pathway from detection of the olfactory stimulus to the motor response.
42 Sexually active (n=4) and inactive (n=3) rams were exposed to urine from estrous
43 ewes. An additional group of sexually active rams (n=3) were exposed to urine from
44 ovariectomized ewes. Rams were exsanguinated following 1 h of exposure to
45 stimulus. Neural activity was determined in tissues of interest by the presence of
46 fos and fos-related proteins detected by immunohistochemistry procedures.
47 Sexually active rams exposed to urine from ovariectomized ewes had more ($P \leq$
48 0.05) fos-positive cells in the olfactory bulb, but fewer ($P = 0.03$) fos-positive cells in
49 the cortical amygdala compared to sexually active rams exposed to urine from
50 estrous ewes. Sexually inactive rams had similar ($P \geq 0.13$) numbers of fos positive
51 neurons in the olfactory bulb and medial amygdala but fewer ($P \leq 0.04$) in the
52 central amygdala, bed nucleus of the stria terminalis and the medial preoptic area
53 compared to sexually active rams exposed to urine from estrous ewes. Sexual
54 inactivity was not associated with decreased hypothalamic function since fos
55 activity was similar ($P \geq 0.14$) among groups in the suprachiasmatic and ventral
56 medial nucleus. Sexual inactivity is not likely due to an impaired ability to detect or
57 process olfactory stimuli by the main olfactory bulb and medial-cortical amygdala.
58 Sexually inactive rams may have reduced attentiveness to sexual stimuli and/or

decreased responsiveness of regions in the brain which regulate reproductive behaviors.

KEYWORDS

Rams, Sexual Behavior, olfactory pathway, c-fos

INTRODUCTION

Rams use olfaction to identify ewes in estrus, and display a preference for urine from ewes during the peri-estrous period (Blissitt et al., 1994a). Olfactory sexual cues facilitate sexual interest in rams and shorten the interval to mounting and ejaculation (Maina and Katz, 1999). However, regardless of the sexual stimuli, approximately 20% of rams are either slow to initiate sexual activity or fail to mount sexually-receptive females (Alexander et al., 1999).

Both the main and accessory olfactory organs detect chemosensory cues. Although the vomeronasal organ is not required for sexually experienced rams to discriminate among estrous and non-estrous females, a functional vomeronasal organ does enhance the expression of reproductive behaviors (Blissitt et al., 1990). As opposed to rodents (Coquelin et al., 1984; Meredith, 1986; Keverne, 1999), the main olfactory bulb identifies sexually relevant odors in rams and is required for the expression of sexual behavior (Blissitt et al., 1990). Olfactory sexual cues are likely processed directly through the amygdala (Martel and Baum, 2009) influencing the bed nucleus of the stria terminalis (BNST) and preoptic area of the hypothalamus (Petrulis, 2013).

Immediate early gene expression is used as a marker of neural activity as a first and transient indicator of gene expression following sensory stimulation in the

central nervous system (Halpern and Martinez-Marcos, 2003). Expression of fos and fos-related proteins is used to evaluate neural activity following exposure to a broad range of stimuli(Bialy and Kaczmarek, 1996). Basal levels of *c-fos* expression are low in the brain and increase precipitously in brain regions known to be important for male sexual behaviors following sexual stimulation(Bialy and Kaczmarek, 1996).

This study was designed to determine if sexual disinterest was a result of an inability to detect sexually relevant olfactory cues, or whether the processing of those cues is compromised in sexually inactive rams.

MATERIALS AND METHODS

All animal procedures were approved by the University of Wyoming animal care and use committee.

Rams.

Mature Western white-faced (Rambouillet x Columbia) commercial rams (*Ovis aries*, 2-3 yrs of age) from the University of Wyoming flock were sequentially evaluated for sexual interest and preference. Rams were maintained as a cohort in a 8.5 x 25.6 m pen fed a forage-based diet with water provided ad libitum.

Ovariectomized ewes used as stimulus animals were kept in a distant pen prior to behavior testing. To determine sexual interest, rams were subjected to a serving capacity test. For the serving capacity test, rams were penned with two or three ewes in estrus for 20 min in an enclosed pen (3.7 m x 4.0 m). Rams not exhibiting mounting behavior were fitted with a marking harness and exposed overnight in their home pen to their pen mates (n = 12 ram cohorts) or to ewes (n = 6) in estrus. To insure high performing rams had preferential interest in estrous females, and

sexually inactive rams did not have latent sexual interest in males, rams were subjected to sexual preference test. Sexual preference tests were conducted by exposing rams to ewes in estrus and cohort rams restrained in a four-way stanchion (Perkins et al., 1992) giving the subject ram free and equal access to all stimulus animals. For classification, rams were tested each breeding season as lambs between 6-10 months of age, yearlings between 18-22 months, and at 2+ years. For each breeding season, each ram was tested in 3 serving capacity tests on three consecutive days followed one week later by 3 sexual preference tests. For previous and unrelated experiments, two sexually inactive rams were castrated following puberty (8 mo) and initial behavior evaluation. Serum concentrations of testosterone were maintained by four Synovex H implants (200 mg testosterone and 20 mg estradiol/implant; Pfizer Animal Health, New York, NY, USA) into their ears (2/ear) one month prior to behavior testing during each breeding season.

Rams were classified based on expression of sexual behaviors during serving capacity and sexual preference tests (Alexander et al., 1999). Investigatory behaviors exhibited were recorded and include numbers of ano-genital investigatory sniffs (nosing of the perineum), flehmen (upper lip curl), fore-leg kicks (foreleg is extended and then flexed in short choppy motions), vocalizations (low-pitched, guttural), and rear nudges (ram positions behind the ewe and contacts the rear of the ewe with his shoulder—often followed by a fore-leg kick) (Banks, 1964). Consummatory behaviors recorded include mount attempts (brief, incomplete mounting of the ewe from behind), mounts (full mount of the ewe without intromission), and ejaculations (mounting with intromission).

andejaculatory thrusting)(Banks, 1964). Investigatory and consummatory behaviors were combined for analysis. Sexually active rams exclusively mounted females during all behavior evaluations, and achieved ≥ 6 ejaculation during serving capacity tests. Rams failing to exhibit sexual interest, having a long latency (>10 min) to mount or achieved ≤ 3 ejaculations in serving capacity tests, and exhibited no interest or preference for males during preference tests were considered sexually inactive.

Ewes.

Urine collected from non-treated ovariectomized ewes ($n = 2$) was used as the control stimulus. Ovariectomized ewes ($n = 2$) were induced into estrus by exposure to progesterone for 14 d with an Eazi-Breed CIDR® for sheep (Pfizer Animal Health, New York, NY, USA). Following CIDR removal, ewes were treated with estradiol ($50\mu\text{g}$) on 2 consecutive days prior to urine collection. Ewes were placed in metabolism crates on the day of CIDR removal for urine collection. Fresh urine was collected on the day of ram exposure and maintained at 37°C .

Exposure to Olfactory Cues.

The final behavior testing occurred 1 month prior to exsanguination during the late breeding season. All rams were 2-3 years of age at the time of odor exposure which occurred within 3 d of each other during the month of November. To decrease stress on the day of exposure, rams were isolated from ewes and placed in indoor treatment pens two days prior to exposure and provided with *ad libitum* access to grass hay and water. Rams were individually penned to prevent sexual contact but allowed visual and olfactory contact with cohort rams. Ten

milliliters of ewe urine from either estrous or ovariectomized ewes was placed on a stack of 3"x3" cotton pads inside of a facemask. Rams were exposed to the olfactory stimuli for one hour.

Blood and Tissue Collection.

Blood was collected from the jugular vein of each ram prior to and following olfactory exposure. Immediately following exposure, rams were anesthetized with intravenous injection of 1.5mL Ketamine/Xylazine mixture (1 part Xylazine to 10 parts Ketamine) then exsanguinated. Heads were removed and perfused via the carotid artery with 500mL of 0.9% saline with 75,000 units of heparin followed by 3L of 4% paraformaldehyde in 0.1M Sorensen's phosphate buffer. Olfactory bulbs and brains were removed from the cranium and the intact brain was postfixed overnight in a 4% paraformaldehyde solution.

Following the post-fix period, tissues were dissected using surface landmarks. The amygdala was dissected on the rostral/caudal plane at the level of the optic chiasm(Glass et al., 1984)(Alexander et al., 2001). The preoptic area of the hypothalamus was identified anterior of the optic chiasm, posterior to the anterior commissure, and lateral to the third ventricle(Glass et al., 1984)(Alexander et al., 2001). The ventromedial hypothalamus was dissected caudal to the preoptic area and lateral to the third ventricle with the fornix located in the anterior portion(Alexander et al., 2001). Fixed brain tissue was cryoprotected in a 20% sucrose fixative solution at 4°C and then rapidly frozen on dry ice and stored at -80°C until sectioned.

Testosterone Radioimmunoassay Procedure.

174 Blood samples were allowed to clot overnight at 4°C then centrifuged at
175 2,000 x g for 20 min. Serum was separated and stored at -20°C until analysis. Serum
176 concentrations of testosterone were analyzed by solid-phase commercially available
177 RIA (Diagnostic Products Corporation, Los Angeles, CA, USA). Standards were
178 prepared by serial dilutions of a stock solution in charcoal-stripped wether
179 serum. Standard or sample were dispensed (0.1 mL) in duplicate into antibody-
180 coated tubes and incubated with radiolabeled testosterone for 4h at room
181 temperature. Samples were analyzed in a single assay with an intra-assay
182 coefficient of variation of 10%.

183 ***Immunohistochemistry Staining Procedure.***

184 Fos and fos related proteins (FRP) were quantified in fixed frozen tissues cut
185 into 40µm coronal sections on a cryostat and stored in cryoprotectant (50% 0.05M
186 sodium phosphate buffer, 30% Ethylene Glycol, 20% Glycerol) at -20°C. Every third
187 section for a total of 24 sections was collected for each tissue. Tissue integrity of the
188 main olfactory bulb was problematic making mounting of consecutive sections
189 difficult; however, 24 total intact sections were stained. Immunohistochemistry was
190 carried out on free-floating sections that were washed overnight in 0.1 M phosphate
191 buffer with 0.9% saline and 0.4% TritonX-100 (PBSTx) to remove cryoprotectant.
192 After inactivation of endogenous peroxidases with 1% H₂O₂ and further PBS
193 washings, the sections were incubated for 2 h in PBS containing 4% normal goat
194 serum diluted in PBSTx. Sections were then incubated at 4°C in affinity-purified
195 rabbit polyclonal antibody (FRP, Ab-2, PC38, Calbiochem, San Diego, CA, USA)
196 diluted 1:20,000 in PBSTx. Following incubation, sections were washed and placed

in a solution of PBTX + 4% NDS with biotinylated goat anti-rabbit (dilution 1:200) for 1 h. The sections were subsequently washed and incubated for 1 h in avidin-biotin-horseradish peroxidase (HRP) complex (dilution 1:100 in PBS, Vector Laboratories, Burlingame, CA, USA). HRP was visualized using 3,3'-diaminobenzidine as the chromagen and reacted for 10 min. Sections were mounted on Superfrost Plus microslides (VWR International, LLC., Radnor, PA, USA) and dried on a drying rack overnight. Mounted sections were dehydrated in ascending ethanol solutions, cleared with xylene, and coverslipped with Permount (Fisher Scientific, Fair Lawn, NJ, USA). Nonspecific binding was determined in tissue incubated in absence of anti-fos primary antibody.

Analysis of Immunohistochemistry Staining.

Sections were viewed using an Olympus Vannox microscope equipped with a digital camera. Images were not altered with the exception of adjusting contrast as necessary. For each region of interest (Fig. 1), 9-12 sections were analyzed using the current version of ImageJ software (USA National Institute of Health) for fos immunoreactive positive cells. A grey scale threshold established the parameters of the fos-positive nuclei, and size exclusion parameters were adjusted by slide based on the size of the positive nuclei. The program subsequently quantified positive cells within a region of interest. Fos-positive cells on the border of the region of interest were excluded from quantification. Fos-positive cells were verified and artifacts removed by visual inspection without knowledge of ram group.

218 Density of FRP-positive neurons was defined as total number of Fos-IR
219 positive cells divided by the area of the region of interest. Cell counts were
220 transformed to average number/mm² by dividing the count by area.

221 ***Statistics.***

222 Cell counts from each region of interest were averaged for statistical analysis.
223 GLM procedure (one-way ANOVA for unbalanced sample size) in SAS (Ver 9.3, SAS
224 Institute, Cary, NC, USA) was used to analyze the data, means $\pm$ SEM are reported. To
225 determine effect of stimulus on serum concentration of testosterone and neural
226 expression of fos and fos related peptides, preplanned comparisons of stimulus
227 exposure (estrous vs ovariectomized ewe urine) in sexually active rams was
228 conducted. Effect of ram classification (sexually active vs inactive) was determined
229 by comparing sexually active to inactive rams with both groups exposed to urine
230 from ewes in estrus.

231 **RESULTS**

232 ***Behavior Testing.***

233 Total investigatory and consummatory behaviors exhibited during the final
234 two serving capacity tests and behaviors exhibited towards rams during the final
235 two sexual preference tests are reported in Table 1. As expected, rams classified as
236 sexually active exhibited the greatest numbers of investigatory (35.5 ± 9.7) and
237 consummatory behaviors (21.4 ± 18.4) towards ewes in estrous. Sexually inactive
238 rams exhibited few investigatory behaviors towards ewes in estrous (4 ± 5.2) and
239 only one sexually inactive achieved intromission.

240 ***Testosterone.***

Basal concentrations of serum testosterone (1.1 ± 0.2 ng/mL) did not differ ($F(2, 7) = 1.1, P = 0.4$) among sexually active and inactive rams prior to urine exposure. Serum concentrations of testosterone in sexually active rams did not change ($F(1, 5) = 0.8, P = 0.4$) following exposure to urine from estrous or ovariectomized ewes (-0.1 ± 0.1 vs 0.3 ± 0.5 ng/mL, respectively). Due to testosterone-replacement in two of three sexually inactive rams, change in serum concentrations of testosterone could not be evaluated in sexually inactive rams.

Fos Immunostaining

Absence of anti-fos antibody resulted in no staining (Fig. 2). Sexually active rams exposed to urine from ovariectomized ewes had greater numbers of FRP-positive neurons in all layers of the olfactory bulb including the granular ($F(1, 5) = 19.4, P = 0.007$) and mitral and tufted layer ($F(1, 5) = 6.61, P = 0.05$) compared to sexually active rams exposed to urine from estrous ewes (Fig. 3). Differences were not noted ($F(1, 5) = 0.2, P > 0.7$) among sexually active and inactive rams exposed to urine from ewes in estrus.

Olfactory stimulation of urine from estrous ewes evoked a greater response in the cortical ($F(1, 5) = 9.0, P = 0.03$) but not the medial ($F(1, 5) = 3.3, P = 0.13$) or central ($F(1, 5) = 5.3, P = 0.07$) nuclei of the amygdala when compared to sexually active rams exposed to urine from ovariectomized ewes (Fig. 4).

Sexually inactive rams had fewer ($F(1, 5) = 11.3, P = 0.02$) numbers of fos positive neurons in the central amygdala when exposed to urine from estrous ewes than sexually active rams exposed to similar stimuli, but differences in the medial amygdala were not noted ($F(1, 5) = 3.3, P \geq 0.13$). Sexually inactive rams had fewer

($F(1, 5) = 7.6, P = 0.04$) numbers of fos positive neurons in the BNST than sexually active rams following exposure to estrous ewe urine (Fig. 5). Hypothalamic responsiveness as measured by numbers of fos-positive neurons was similar among rams in the ventral medial ($F(1, 5) = 3.1, P \geq 0.14$) and suprachiasmatic ($F(1, 5) = 0.3, P \geq 0.60$) nuclei (Fig. 5). Numbers of fos-positive neurons in the medial preoptic area (mPOA), however, were fewer ($F(1, 5) = 34.7, P = 0.002$) for sexually inactive compared to sexually active rams exposed to urine from estrous ewes (Fig. 5). Differences in fos activity in the mPOA were not noted among sexually active rams exposed to urine from ovariectomized vs estrous ewes (Fig. 5).

DISCUSSION

Sexually active rams placed among randomly cycling females quickly identify ewes in estrus. Although ewes do exhibit limited proceptive behavior such as seeking and affiliation (Lindsay, 1974), rams primarily use olfactory cues to discriminate the estrus state of ewes (Blissitt *et al.*, 1994). Exposure of rams to ewes in estrus increases serum concentrations of LH and testosterone (Ungerfeld and Silva, 2004), but olfactory cues alone are not sufficient to elicit this increase (Gonzalez *et al.*, 1991a). Similarly, in the present study, sexually active rams exposed to urine from estrous ewes did not experience a rise in testosterone in response to olfactory stimuli. Testosterone may influence fos activity independent of stimulus (De Lorme and Sisk, 2013); therefore, it is important to note serum concentrations of testosterone were similar among groups prior to the presentation of olfactory stimulus and did not rise in response to stimulus.

287 Urine from ovariectomized ewes elicited a greater fos response in the main
288 olfactory bulb in sexually active rams than did urine from estrus ewes. It is possible
289 the chemical composition of ovariectomized ewe urine is sufficiently more
290 complex than urine from ewes in estrus to elicit this increased response. Regardless
291 of urine complexity, sexually experienced rams clearly differentiate among ewes.
292 Ewes will characteristically void urine when approached by a ram if they are not in
293 estrus (Bland and Jubilan, 1987). In response to main olfactory stimuli (Hart, 1987),
294 rams sniff and sample voided urine and typically follow with a flehmen response.
295 The flehmen response facilitates transport of the urine to the vomeronasal
296 organ (Ladewig and Hart, 1980). Although it is not clear how rams learn to associate
297 particular urine composition to a willing vs unwilling mate, classic conditioning may
298 be involved. Classic conditioning response was demonstrated when a foot shock
299 was paired with the presentation of an odor, rats had increased fos activity in the
300 main as well as accessory olfactory bulb when compared to presentation of odor
301 alone (Funk and Amir, 2000). Differences in fos activity, however, were not
302 apparent among high and low sexually performing rams exposed to urine from
303 estrous ewes. Thus, it appears rams with low libido have an equal ability to detect
304 potentially sexually evocative stimuli. It should be noted, however, the urine
305 stimulus in the present study primarily provided volatile cues, and non-volatile cues
306 from urine saturating the facial mask would have been limited. Because of the study
307 design, potential differences in the ability to detect non-volatile sexual cues cannot
308 be disregarded.

309 Initial recognition and discrimination of a ewe's reproductive status likely
310 occurs at the level of the olfactory bulb(Blissitt et al., 1994b). However, the motor
311 response to that stimuli requires further processing of the stimulus. Olfactory
312 information through the main and accessory bulb is transmitted to the amygdala
313 (Brennan and Keverne, 1997). In sheep, the cortical amygdala receives input from
314 both the main and accessory olfactory bulbs while the medial amygdala receives
315 input only from the accessory bulb (Meurisse et al., 2009). Although fos activity in
316 the medial amygdala was similar among sexually evocative and non-evocative
317 stimuli, fos activity was decreased in the cortical amygdala when rams were exposed
318 to urine from ovariectomized ewes compared to exposure to a putative sexual
319 stimuli. It is unexpected that fos activity in the medial amygdala was not different
320 among rams exposed to urine from ovariectomized or estrous ewes especially since
321 flehmen occurs most often toward non-estrous females and connections with the
322 accessory olfactory bulb are primarily to the medial amygdala (Meurisse et al.,
323 2009). In absence of a flehmen response, however, minimal fluid is transported into
324 the vomeronasal organ (Hart, 1987). In the present study the face mask likely
325 prohibited flehmen, and may account for the lack of differentiation in the medial
326 amygdala. Lesions to the cortical amygdala in Syrian hamsters did not influence
327 odor preference or discrimination, but did influence behavior patterns associated
328 with sexual satiety(Maras and Petrulis, 2008). Neural activity in the cortical
329 amygdala may facilitate the appropriate behavioral response to olfactory stimuli
330 (i.e. mounting vs feeding). A dense interconnection of these two amygdala nuclei
331 (Meurisse et al., 2009) may also refine olfactory signals. Similar to fos activity in the

olfactory bulb, rams characterized as sexually inactive had similar fos activity in the medial and cortical amygdala as sexually active rams when exposed to a putative sexual stimuli. Similar fos activity was also noted in rams allowed to interact with ewes in estrus (Borja and Fabre-Nys, 2012). Our study along with Borja and Fabre-Nys (2012) clearly indicate that medial and cortical amygdala function is not responsible for the lack of sexual interest in rams with low libido.

Fos activity in the central amygdala was lower in sexually inactive rams compared to sexually active rams following exposure to urine from ewes in estrus. The central nucleus initiates a state of arousal towards non-specific stimuli (Gallagher et al., 1990). Sexually active rams may be more aware and vigilant than sexually inactive rams and that vigilance facilitates increased sexual activity. In a production setting, sexually active rams seem to investigate their environment more and track to the feed source more often without consuming feed (Uthlaut et al., 2011). In the present study, similar fos activity in the central amygdala of sexually active rams exposed to either putative sexually evocative or non-evocative stimuli lends credence to this hypothesis.

In sheep, both the medial as well as the cortical amygdala have reciprocal connections to the BNST and the mPOA (Meurisse et al., 2009). Although differences in fos expression in the medial and cortical amygdala was not noted, sexually inactive rams had fewer numbers of fos positive neurons in the BNST and mPOA when compared to sexually active rams exposed to urine from ewes in estrus. This contrasts with (Borja and Fabre-Nys, 2012), who noted less fos expression in the POA but not BNST of low libido rams with full access to ewes in estrus. Differences

in stimuli exposure may explain the differences. The POA is known to be essential for the expression of male reproductive behaviors—especially mounting and intromission (Hart and Leedy, 1985). Lower fos expression in the mPOA may be explained in the Borja and Fabre-Nys (2012) experiment by the reduced mounting activity in their low libido rams. However, in conjunction with present results, it seems plausible that inherent low activity in the mPOA may be at least partially responsible for the lack of sexual activity in low sexually performing rams. Similar levels of fos expression among sexually active rams exposed to putative sexually evocative and non-evocative stimuli suggests sexually active rams require more sensory input to elicit a sexual response. This is supported by evidence that olfactory stimulus alone does not cause an increase in serum concentrations of testosterone (Alexander et al., 1999)(Gonzalez et al., 1991b), present study.

Aberrant hypothalamic function could be cause for decreased sexual interest in rams as has been shown in males of many species with large hypothalamic lesions (for review: (Hart and Leedy, 1985). However, it does not appear sexually inactive rams have a global impairment/reduction in hypothalamic function—clearly low sexually performing rams have normal serum concentrations of testosterone (Alexander et al., 1999), grow similarly and are selected as flock sires (Uthlaut et al., 2011), and also fos expression in the suprachiasmatic and ventral medial hypothalamus is similar among all rams.

CONCLUSION

Sexual inactivity is not likely due to an impaired ability to detect or process olfactory stimuli from by the main olfactory bulb or medial and cortical amygdala, but may be caused by a reduced vigilance to sexual stimuli and/or decreased responsiveness of the BNST and mPOA. This study advances our knowledge of mechanisms which regulate reproductive behavior. A thorough understanding of such basic mechanisms is a prerequisite to the development of reliable methods to regulate behavior.

ACKNOWLEDGEMENT

Funded in part by 3R01RR014270-10S1 NIH/DHHS (CER, PI) and USDA-NRI 2007-55618-18176 (BMA, PI).

Figure Legends

Figure 1. Nuclei and tissue regions analyzed: Panel A, Ventral medial hypothalamus (VMH); Panel B, Bed Nucleus of the Stria Terminalis (BNST), medial Preoptic Area of the Hypothalamus (POA), and suprachiasmatic nuclei (SCN); Panel C, Central Amygdala (CeA), Medial Amygdala (MeA), and Cortical Amygdala (CoA). Other identified regions: optic tract (OT), optic chiasm (OC), arcuate nucleus of the hypothalamus (ARC), supraoptic nucleus of the hypothalamus (SON), third ventricle (3V), anterior commissure (AC), fiber tract (FT), and stria terminalis (ST).

Figure 2. Preoptic area of the hypothalamus stained for c-fos and fos-related proteins from treatment groups: (clockwise) sexually active rams exposed to

estrous ewe urine, sexually active rams exposed to ovariectomized ewe urine and sexually inactive rams exposed to urine from estrous ewes.

Figure 3. Average fos and fos-related proteins positive neurons in the granular and mitral and tufted layer of the olfactory bulb per mm². Sexually active (A) or inactive (I) rams were exposed to urine from ewes in estrus (EU) or ovariectomized ewes (OU). *Denotes difference compared to sexually active rams exposed to urine from ewes in estrus (A-EU) ($P < 0.05$).

Figure 4. Average fos and fos-related proteins positive neurons in the medial, cortical, and central amygdala per mm². Sexually active (A) or inactive (I) rams were exposed to urine from ewes in estrus (EU) or ovariectomized ewes (OU). *Denotes difference compared to sexually active rams exposed to urine from ewes in estrus (A-EU) ($P < 0.05$).

Figure 5. Average fos and fos-related proteins positive neurons in the bed nucleus of the stria terminalis (BNST), preoptic area of the hypothalamus (POA), ventromedial hypothalamus (VMH), and the suprachiasmatic nucleus (SCN) per mm². Sexually active (A) or inactive (I) rams were exposed to urine from ewes in estrus (EU) or ovariectomized ewes (OU). *Denotes difference compared to sexually active rams exposed to urine from ewes in estrus (A-EU) ($P < 0.05$).

Literature Cited

423 Alexander, B.M., Rose, J.D., Stellflug, J.N., Fitzgerald, J.A., Moss, G.E., 2001. Fos-like
 424 immunoreactivity in brain regions of domestic rams following exposure to rams or
 425 ewes. *Phys. Behav.* 73, 75-80.
 426 Alexander, B.M., Stellflug, J.N., Rose, J.D., Fitzgerald, J.A., Moss, G.E., 1999. Behavior
 427 and endocrine changes in high-performing, low-performing, and male-oriented
 428 domestic rams following exposure to rams and ewes in estrus when copulation is
 429 precluded. *J. Anim. Sci.* 77, 1869-1874.
 430 Banks, E.M., 1964. Some aspects of sexual behavior in domestic sheep, *ovis aries*.
 431 *Behaviour* 23, 249-279.
 432 Bialy, M., Kaczmarek, L., 1996. c-Fos expression as a tool to search for the
 433 neurobiological base of the sexual behaviour of males. *Acta Neurobiol. Exp. (Wars)*
 434 56, 567-577.
 435 Bland, K.P., Jubilan, B.M., 1987. Correlation of Flehmen by Male Sheep with Female
 436 Behavior and Estrus. *Anim. Behav.* 35, 735-738.
 437 Blissitt, M.J., Bland, K.P., Cottrell, D.F., 1990. Olfactory and Vomeronasal
 438 Chemoreception and the Discrimination of Estrous and Non-Estrous Ewe Urine
 439 Odors by the Ram. *Appl. Anim. Behav. Sci.* 27, 325-335.
 440 Blissitt, M.J., Bland, K.P., Cottrell, D.F., 1994a. Detection of Estrous-Related Odor in
 441 Ewe Urine by Rams. *J. Reprod. Fertil.* 101, 189-191.
 442 Blissitt, M.J., Bland, K.P., Cottrell, D.F., 1994b. Detection of Estrous-Related Odor in
 443 Ewe Urine by Rams. *J. Reprod. Fertil.* 101, 189-191.
 444 Borja, F., Fabre-Nys, C., 2012. Brain structures involved in the sexual behaviour of Ile
 445 de France rams with different sexual preferences and levels of sexual activity.
 446 *Behav. Brain Res.* 226, 411-419.
 447 Brennan, P.A., Keverne, E.B., 1997. Neural mechanisms of mammalian olfactory
 448 learning. *Prog. Neurobiol.* 51, 457-481.
 449 Coquelin, A., Clancy, A.N., Macrides, F., Noble, E.P., Gorski, R.A., 1984. Pheromonally
 450 induced release of luteinizing hormone in male mice: involvement of the
 451 vomeronasal system. *J. Neurosci.* 4, 2230-2236.
 452 De Lorme, K.C., Sisk, C.L., 2013. Pubertal testosterone programs context-appropriate
 453 agonistic behavior and associated neural activation patterns in male Syrian
 454 hamsters. *Phys. Behav.* 112-113, 1-7.
 455 Funk, D., Amir, S., 2000. Enhanced fos expression within the primary olfactory and
 456 limbic pathways induced by an aversive conditioned odor stimulus. *Neuroscience*
 457 98, 403-406.
 458 Gallagher, M., Graham, P.W., Holland, P.C., 1990. The amygdala central nucleus and
 459 appetitive Pavlovian conditioning: lesions impair one class of conditioned behavior.
 460 *J. Neurosci.* 10, 1906-1911.
 461 Glass, J.D., Amann, R.P., Nett, T.M., 1984. Effects of season and sex on the distribution
 462 of cytosolic estrogen receptors within the brain and the anterior pituitary gland of
 463 sheep. *Biol. Reprod.* 30, 894-902.
 464 Gonzalez, R., Levy, F., Orgeur, P., Poindron, P., Signoret, J.P., 1991a. Female effect in
 465 sheep. II. Role of volatile substances from the sexually receptive female; implication
 466 of the sense of smell. *Reprod. Nutr. Dev.* 31, 103-109.

Gonzalez, R., Levy, F., Orgeur, P., Poindron, P., Signoret, J.P., 1991b. Female effect in sheep. II. Role of volatile substances from the sexually receptive female; implication of the sense of smell. *Reprod. Nutr. Dev.* 31, 103-109.

Halpern, M., Martinez-Marcos, A., 2003. Structure and function of the vomeronasal system: an update. *Prog. Neurobiol.* 70, 245-318.

Hart, B., Leedy, M., 1985. Neurological Bases of Male Sexual Behavior, in: Adler, N., Pfaff, D., Goy, R. (Eds.), *Reproduction*, Springer US, pp. 373-422.

Hart, B.L., 1987. Roles of the olfactory and vomeronasal systems in behavior. *Vet. Clin. N. Am.-Food A.3*, 463-475.

Keverne, E.B., 1999. The vomeronasal organ. *Science* 286, 716-720.

Ladewig, J., Hart, B.L., 1980. Flehmen and vomeronasal organ function in male goats. *Phys. Behav.* 24, 1067-1071.

Lindsay, D.R.a.F., I. C., 1974. Ram seeking activity associated with oestrous behavior in ewes. *Anim. Behav.* 20, 452-456.

Maina, D., Katz, L.S., 1999. Scent of a Ewe: transmission of a social cue by conspecifics affects sexual performance in male sheep. *Biol. Reprod.* 60, 1373-1377.

Maras, P.M., Petrulis, A., 2008. The posteromedial cortical amygdala regulates copulatory behavior, but not sexual odor preference, in the male Syrian hamster (*Mesocricetus auratus*). *Neuroscience* 156, 425-435.

Martel, K.L., Baum, M.J., 2009. A centrifugal pathway to the mouse accessory olfactory bulb from the medial amygdala conveys gender-specific volatile pheromonal signals. *European Journal of Neuroscience* 29, 368-376.

Meredith, M., 1986. Vomeronasal organ removal before sexual experience impairs male hamster mating behavior. *Physiol. Behav.* 36, 737-743.

Meurisse, M., Chaillou, E., Levy, F., 2009. Afferent and efferent connections of the cortical and medial nuclei of the amygdala in sheep. *J. Chem. Neuroanat.* 37, 87-97.

Perkins, A., Fitzgerald, J.A., Price, E.O., 1992. Luteinizing hormone and testosterone response of sexually active and inactive rams. *J. Anim. Sci.* 70, 2086-2093.

Petrulis, A., 2013. Chemosignals and hormones in the neural control of mammalian sexual behavior. *Front. Neuroendocrin.* 34, 255-267.

Ungerfeld, R., Silva, L., 2004. Ewe effect: endocrine and testicular changes in experienced adult and inexperienced young Corriedale rams used for the ram effect. *Anim. Reprod. Sci.* 80, 251-259.

Uthlaut, V.A., Moss, G.E., Stobart, R.H., Larson, B.A., Alexander, B.M., 2011. Sexual performance and production traits in white-faced yearling rams. *Small Ruminant Res.* 100, 63-66.

Figure 1

Table 1. Total number of investigatory and consummatory behaviors exhibited by sexually active (A) and sexually inactive (I) rams in the final two behavior tests.

Ram	Group	Serving Capacity Test		Behavior Toward Rams	
		Invest ^a	Consum ^b	Invest ^a	Consum ^b
8507	A	33	17	4	0
8011	A	25	11	3	0
8027	A	39	5	2	0
8045	A	48	31	1	0
8114	A	31	15	1	0
8576	A	46	63	0	0
8587	A	41	18	3	0
85136	A	21	11	2	0
7044	I	1	0	5	0
7111	I	1	0	0	0
8507	I	10	1	4	0

^aInvest = Investigatory behaviors expressed and include ano-genital sniffs, flehmen, foreleg kick and vocalizations

^bConsum = Consummatory behaviors expressed and include attempted mounts, mounts, and mounts with intromission or ejaculation