1

2 **Characterization of Extended-Spectrum β -Lactamase (ESBL) Producing**
3 **Non-Typhoidal *Salmonella* (NTS) from Imported Food Products**

4

5

6 Dongryeoul Bae¹, Chorng-Ming Cheng², and Ashraf A. Khan^{1*}

7

8 *Division of Microbiology, National Center for Toxicological Research, U.S. Food and Drug*
9 *Administration, Jefferson, Arkansas 72079¹ and Pacific Regional Laboratory-Southwest, U.S.*
10 *Food and Drug Administration, Irvine, California 92612²*

11

12

13

14 Running title: Resistance to β -lactam antibiotics in *S. enterica* isolates

15

16

17 Keywords: *Salmonella enterica*, Food, ESBL, Antimicrobial resistance, Conjugation

18

19

20

21 * Corresponding author.

22 Mailing address: Division of Microbiology, National Center for Toxicological Research, U.S.
23 Food and Drug Administration, Jefferson, AR 72079. Phone: (870) 543 7601. Fax: (870) 543
24 7307. E-mail: Ashraf.Khan@fda.hhs.gov.

25

Abstract

Food contaminated with extended-spectrum β -lactamase (ESBL)-producing *Salmonella enterica* has emerged as an important global issue due to the international food-products trade. Therefore, the purpose of this study was to investigate whether imported food products can serve as a reservoir for non-typhoidal *Salmonella* (NTS) that can transmit β -lactam-resistance to humans through ingestion of the contaminated food. NTS isolates ($n = 110$) were collected from various imported food products ($n = 3,480$) from 2011 to 2013. The NTS isolates were analyzed by serotyping, antimicrobial susceptibility tests, and plasmid profiling. *S. ser. Weltevreden*, *S. ser. Newport*, *S. ser. Senftenberg*, *S. ser. Virchow*, *S. ser. Enteritidis*, *S. ser. Typhimurium*, and *S. ser. Bareilly* were the most prevalent serovars. Nine NTS strains were resistant to ampicillin and/or one or more cephalosporins (MIC $> 32 \mu\text{g/ml}$). Polymerase chain reaction (PCR) detection revealed that all nine isolates carried the *bla*_{TEM-1} β -lactamase gene, with or without the *bla*_{CTX-M-9} or *bla*_{OXA-1} genes. Two isolates, PSS_913 and PSS_988, exhibited decreased susceptibility to extended-spectrum cephalosporins and ampicillin. Plasmids ranging in size from less than 8 to over 165 kbp, from all of the 9 resistant isolates, belonged to the IncHI1, IncI1, IncN, or IncX groups. Conjugation experiments and Southern hybridization, using *bla*_{TEM-1}, confirmed the plasmid-mediated transfer of ESBL genes, which resulted in increased MICs of β -lactams for *E. coli* transconjugants. The contamination of imported food products by NTS with conjugative plasmid-borne ESBL genes may contribute to the spread of ESBL-producing NTS and compromise the therapeutic activity of extended-spectrum β -lactam antibiotics.

1. Introduction

The contamination of food by non-typhoidal *Salmonella* (NTS) is becoming global concerns due to the higher contamination and antimicrobial resistance rates in imported food products than in domestic food products (Akiyama and Khan 2011, 2012; Akiyama et al., 2011, 2012, Mezal et al, 2013; Ponce et al., 2008; Skov et al., 2007; Zhao et al., 2003; Zhao et al., 2006). *Salmonella* is the most commonly isolated genus of foodborne pathogens in the United States (Crim et al., 2014). Even though more than 2,500 NTS serovars have been reported (Scallan et al., 2011), these *Salmonella* serovars, such as *S. ser. Enteritidis*, *S. ser. Typhimurium*, *S. ser. Heidelberg*, and *S. ser. Newport*, have been mostly linked to human *salmonella* outbreaks in the US (CDC, 2013a; Jackson et al., 2013). NTS serovars are estimated to cause over 1M cases of gastroenteritis, enteric fever, diarrhea, meningitis, and bacteremia, resulting in approximately 20,000 hospitalizations and 380 deaths annually in the US alone (CDC, 2013b; Fey et al., 2000; Hellerstein, 2003; Hohmann, 2001; Scallan et al., 2011; Soler et al., 2006). NTS infection in most cases is caused by ingestion of contaminated food (CDC, 2015; Tauxe, 1991; Zhao et al., 2008). NTS strains resistant to multiple antimicrobials have emerged worldwide and have become a serious public health problem (Chen et al., 2004; Dahshan et al., 2010; Gebreyes and Thakur, 2005). The reduced susceptibility of *Salmonella* to ampicillin, ceftriaxone, and ciprofloxacin, has been of globally concern. As a result, the World Health Organization (WHO) has supported a project, called the WHO Global Salm-Surv (WHO GSS) program, to conduct global surveillance and identify antimicrobial resistance of *Salmonella*, since 2000 (Hendriksen et al., 2009). In spite of the international cooperation program to improve public health through food safety enhancement, the incidence of *Salmonella* infection has not been conspicuously decreased (CDC, 2011; Chai et al., 2012).

The antimicrobial resistance of *Salmonella* isolated from humans, animals and foods has been monitored by the National Antimicrobial Resistance Monitoring System (NARMS) sponsored by the Centers for Disease Control and Prevention (CDC), the U.S. Food and Drug Administration's Center for Veterinary Medicine (FDA-CVM) and the U.S. Department of Agriculture's Agricultural Research Service (USDA-ARS) since 1996 (CDC, 2013b). Recently, Jackson and colleagues (2013) reported the most common NTS serovars that caused outbreaks during 1998 to 2008 by consuming food commodities, including eggs, chicken, pork, beef, fruit, turkey, and vegetables.

β -Lactam antibiotics, agents for the treatment of *Salmonella* infection, act to inhibit bacterial cell wall biosynthesis (Tomasz, 1979; Wise and Park, 1965). In response to the agents, bacteria have developed resistance to β -lactams by synthesizing a β -lactamase, which cleaves the β -lactam ring and consequently inactivates the antibiotic. Ceftriaxone, a third-generation of cephalosporin, has been widely used to treat invasive *Salmonella* infection and severe diarrhea (Al-Mashhadani et al., 2011). The resistance of NTS to extended-spectrum cephalosporins (ESCs) has been reported (CDC, 2013b; Dunne et al., 2000; Yates and Amyes, 2005). The prevalence of ceftriaxone resistant NTS was reported in 1996 (0.1%), in 1997 (0.4%), and in 1998 (0.5%) [CDC, 2013b]. Eighty percent of the ceftriaxone-resistant isolates had MIC > 16 μ g/mL. Four of the 2008 (0.2%) isolates in 2002 and 8 of 1865 (0.4%) isolates in 2003 were resistant to ceftriaxone (MIC 16 μ g/mL or higher). Since 2003, NARMS has reported the resistance rate of NTS to ceftriaxone as MIC $\geq$ 2 μ g/mL (CDC, 2013b). Thus, data for the ceftriaxone susceptibility of NTS in the NARMS annual reports show prevalence rates of 4.4%

90 (2002 and 2003), 3.3% (2004 and 2007), 2.9% (2005 and 2010), 3.7% (2006), 3.1% (2008),
91 3.4% (2009), 2.9% (2010), and 2.5% (2011) (CDC, 2013b).

92 Therefore, the aims of this study were to characterize *Salmonella* food isolates resistant to
93 extended-spectrum β -lactams (including ESCs) and examine transferability of antibiotic
94 resistance phenotypes or genotypes from NTS food isolates to other enteric bacteria for
95 determining the possible food sources and transport routes on ESBL or AmpC β -lactamases-
96 acquired NTS.

2. Materials and methods

2.1 Isolation, identification, and serotyping of *S. enterica* serovar isolates

S. enterica strains were recovered from imported food products, including fresh, frozen, and dried vegetables, fruits, meats, and seafood, during 2011 to 2013. The *Salmonella* strains were isolated, identified, and serotyped by the U.S. FDA's Pacific Regional Laboratory-Southwest (Irvine, California) using BAM procedures (USFDA/CFSAN, 2014). The stock cultures in LB broth (Difco Laboratories, Detroit, MI) were maintained at -80 °C until use.

2.2 Conjugation

The ampicillin or cephalosporin resistant *Salmonella* isolates (donors) and a sodium azide resistant *E. coli* J53 strain (recipient) were used for conjugation experiments to determine transferability of β -lactam antibiotic resistance genes. The broth mating conjugation method was performed by the method of Tran et al. (Tran et al., 2010). MacConkey (BD, Franklin Lakes, NJ) and LB agar plates containing NaN₃ (300 mg/mL, Sigma Chemical Co., St. Louis, MO) with ampicillin (100 μ g/mL, USB, Cleveland, OH) or ceftriaxone (64 μ g/mL, Sigma Chemical Co.) were used to select transconjugants harboring genes conferring β -lactam resistance.

2.3 Antimicrobial susceptibility assay

Antimicrobial susceptibility assays were conducted by the broth microdilution method according to the procedures specified by the Clinical and Laboratory Standards Institute (CLSI, 2014). In brief, one hundred microliter inocula of *S. enterica* isolates and *E. coli* J53 transconjugants (about 1.0×10^6 /ml) were added to 96-well microtiter plates containing 100 μ l of LB broth with ampicillin and cephalosporins (cephalothin - 1st, cefoxitin - 2nd, ceftriaxone - 3rd,

and cefepime - 4th generation, Sigma Chemical Co.). The concentrations of cephalosporins ranged from 0.06 to 1024 µg/mL. The inoculum cultures were incubated at 37 °C for 18 h. *Escherichia coli* ATCC 25922 was used as the assay control strain.

2.4 Plasmid and genomic DNA isolation and detection of genes involved in ESBLs

Nine ampicillin and cephalosporin (MIC >32 µg/mL) resistant NTS food isolates and 8 *E. coli* J53 transconjugants were cultured at 37 °C overnight in LB broth. Plasmids were isolated as previously described (Klaenhammer, 1984). Five milliliters of a bacterial culture was used for plasmid isolation and 1 mL of the culture was used for genomic DNA extraction, using the Qiagen DNeasy Blood and Tissue kit (Qiagen, Valencia, CA). The plasmid DNA was loaded on 0.8% agarose gels and electrophoresed using a CHEF Mapper XA (Bio-Rad Laboratories, Hercules, CA) at 6 V/cm for 7 h. A BAC-TrackerTM Supercoiled DNA Ladder (Epicentre Biotechnologies, Madison, WI) was used as a molecular weight marker. The agarose gel incubated with ethidium bromide was visualized by using Bio-Rad Gel DOC XR imaging system and stored at 4 °C until ready to perform the Southern blot. Primers were used for detection of genes involved in the resistance of β-lactam antibiotics and for determination of plasmid type (Table 1). For the further confirmation of plasmid-mediated β-lactam resistant isolates, PCR was performed followed by Southern hybridization using probes specific for *bla*_{TEM-1}, *bla*_{CTX-M-9}, and *bla*_{OXA-1}.

2.5 Southern blot hybridization

Southern hybridization analysis was performed according to the instructions of the DIG DNA Labeling and Detection Kit (Roche Diagnostics, Indianapolis, IN). The *bla*_{TEM-1}, *bla*_{CTX-M-9},

139 and *bla*_{OXA-1} PCR products were labelled with digoxigenin (DIG)-11-dUTP. Plasmid DNAs from
140 *Salmonella* isolates and transconjugants were transferred to a positively charged nylon
141 membrane (Millipore, Bedford, MA) by (Gebreyes and Thakur, 2005; Khan et al, 2009). The
142 membrane was then UV-cross linked and rinsed. After prehybridization for 30 min at 65 °C, the
143 membrane was hybridized in DIG Easy Hyb solution containing DIG-labelled DNA probe at
144 45 °C for 18 h. The membrane was then washed, blocked, and incubated in antibody solution
145 containing alkaline phosphate-conjugated anti-DIG antibody (Roche). DIG-labelled DNA probe
146 was detected with disodium 3-(4-methoxyspiro-{1,2-dioxetane-3,2'-(5'-chloro) tricyclo [3.3.1.1
147 3'7] decan}-4-yl) phenyl phosphate (CSPD, Roche) following the manufacturer's instructions.

3. Results

3.1 Distribution of NTS strains from imported food products

A total of 110 NTS strains from 3,840 food samples were assigned to 51 diverse serovars. The prevalence of NTS in the imported food products represents 2.86%. The most common serovars included *S. ser. Weltevreden* (19.1%), *S. ser. Newport* (8.2%), *S. ser. Senftenberg* (8.2%), *S. ser. Virchow* (6.4%), *S. ser. Enteritidis* (5.5%), *S. ser. Typhimurium* (4.5%), and *S. ser. Bareilly* (4.5%) [Table 2]. Table 2 also shows that the NTS strains were mostly recovered from seafood (44.5%), vegetables (28.2%), or freshwater fishes (17.3%). *S. ser. Weltevreden* (28.6%) and *S. ser. Senftenberg* (16.3%), *S. ser. Weltevreden* (12.9%) and *S. ser. Enteritidis* (9.7%), and *S. ser. Virchow* (21.0%) were predominant in seafood, vegetables, and freshwater fishes, respectively. The food products contaminated with NTS strains were mostly imported from India (17.3%), Taiwan (17.3%), Vietnam (15.5%), and Indonesia (11.8%) [Table 2].

3.2 Characterization of ESBL-producing NTS isolates

One hundred ten NTS isolates were characterized by antimicrobial susceptibility and PCR analyses. Nine (8.2%) NTS isolates (PSS_903, PSS_913, PSS_928, PSS_948, PSS_955, PSS_959, PSS_970, PSS_984, and PSS_988) were resistant to ampicillin and/or cephalosporins. The 9 NTS serovars included *S. ser. Enteritidis*, *S. ser. Indiana*, *S. ser. Weltevreden*, *S. ser. Newport*, *S. ser. Typhimurium*, and *S. ser. Give*, and were recovered from frozen seafood, amphibian, freshwater fish, pepper, or ginger products that had been imported from Indonesia, Vietnam, China, or Taiwan (Table 3). *S. ser. Enteritidis* was the most prevalent among the β -lactam resistant NTS isolates.

PCR detection revealed that all ampicillin and/or cephalosporin-resistant NTS strains carried the *bla*_{TEM-1} β -lactamase gene, with or without the *bla*_{CTX-M-9} or *bla*_{OXA-1} gene, involved in resistance to extended-spectrum β -lactams (Table 3). The 9 strains were resistant to cephalosporins and/or ampicillin (MIC > 32 μ g/mL). *bla*_{TEM-1} was detected in all 9 ampicillin-resistant isolates. Of ampicillin and 3rd and 4th cephalosporin-resistant strains, *S. ser. Indiana* PSS_913 harbored *bla*_{CTX-M-9} with *bla*_{TEM-1}, whereas *S. ser. Enteritidis* PSS_988 harbored only *bla*_{TEM-1}. The two isolates (PSS_913 and PSS_988) exhibited decreased susceptibility to ESCs and ampicillin. In addition, *S. ser. Weltevreden* PSS_928 and *S. ser. Typhimurium* PSS_955 contained *bla*_{CTX-M-9} and *bla*_{OXA-1} with *bla*_{TEM-1}, respectively (Table 3). Even though only two (1.82%) of the 110 NTS food isolates were resistant to ceftriaxone and other cephalosporins, the MICs of *S. ser. Indiana* PSS_913 and *S. ser. Enteritidis* PSS_988 to ceftriaxone were more than 512 μ g/mL. PSS_913 was resistant to all cephalosporins (MIC $\geq$ 32 mg/L) used in the study, whereas PSS_988 was susceptible to only cefoxitin (MIC of 8 mg/L) (Table 4).

3.3 Horizontal transfer of β -lactam-resistance phenotypes and genotypes

The resistance of *E. coli* J53 transconjugants to ampicillin and cephalosporins was determined. All transconjugants were resistant to ampicillin. Most of them were also resistant to cephalothin and cefoxitin, but susceptible to ceftriaxone and cefepime, while strains of the transconjugants *E. coli* J53_913 and J53_988 were resistant to ceftriaxone and cefepime, as in the *Salmonella* donor strains. Plasmid profiles of the 9 ESBL-producing strains were identified by DNA bands with different numbers and sizes, ranging from less than 8 kb to 150 kb (Fig. 1 A). Plasmids in all transconjugants except J53_984 were shown to be transferred (Table 4). The transfer of *bla*_{TEM-1} was found on various conjugative plasmids of NTS isolates and *E. coli* J53

191 transconjugants by Southern hybridization analysis (Fig. 1 B and C). In addition, *bla*_{CTX-M-9} was
192 transferred to the transconjugant *E. coli* J53_913, but not to J53_928. *bla*_{OXA-1} was not
193 transferred to *E. coli* J53. The resistance of *S. ser. Give* PSS_984 to ampicillin or cephalosporins
194 was not transferred to *E. coli* J53. When the plasmids were classified using the PCR-based
195 replicon typing assay, five isolates (PSS_948, PSS_955, PSS_970, PSS_984, and PSS_988) were
196 grouped with IncHI1, IncI1, IncN, and/or IncX (Table 1). The *bla*_{PSE-1}, *bla*_{SHV-1}, *bla*_{CTX-M},
197 *bla*_{CMY-2}, *bla*_{FOX}, *bla*_{IMP}, *bla*_{KPC}, or *bla*_{VIM} genes associated with ESBL or AmpC β -lactamase
198 (Grimm et al., 2004; Grobner et al., 2009; Paterson et al., 2003; Perez-Perez and Hanson, 2002;
199 Simarro et al., 2000) were not amplified from the plasmid or genomic DNAs of the isolates with
200 the primers used in the study (Table 1).

4. Discussion

The introduction of food safety management systems has played a greater role to reduce carriage of pathogens between trading nations. The amount and variety of imported food products consumed in the United States have increased; approximately 17 percent of total US food consumption is imported (USDA-ERS, 2015). Many studies on antimicrobial resistance of NTS isolated from humans, food animals, retail meat, etc. have been conducted, whereas studies on NTS isolated from imported vegetables, fruits, or seafood are less common (Chen et al., 2004; Egervarn et al., 2014; Fey et al., 2000; Gebreyes and Thakur, 2005; Perez-Perez and Hanson, 2002; Sjölund-Karlsson et al., 2013; Zhao et al., 2008). This study showed that ESBLs-producing NTS strains were highly resistant to ampicillin and ESCs. Even though the prevalence (1.8%) of NTS food isolates resistant to cephalosporins was relatively low when compared to previous NARMS data from 2002 (4.4%) to 2011 (2.5%), the high MICs of *S. Indiana* PSS_913 and *S. ser. Enteritidis* PSS_988 to ceftriaxone were noteworthy (CDC, 2013a; Dunne et al., 2000). The high MIC data may decrease the therapeutic activity of ESCs on both Gram-negative and positive bacteria.

Few studies have been conducted on the prevalence, serotyping, molecular characterization, and antimicrobial resistance of *Salmonella* food isolates imported into the US (Van Doren, et al., 2013a; Van Doren, et al., 2013b). This study on 110 *Salmonella* food isolates originated from over 20 countries provides the prevalence of ESBL-producing NTS strains. The food isolates become more concerned with increased antimicrobial-resistance. Interestingly, there was no antimicrobial-resistant NTS strain in food products imported from India, although the prevalence of *S. enterica* isolates was the highest among the exporting countries (Table 2).

However, we have isolated and reported earlier several multi-drug resistant NTS strains from

food samples originated from India (Akiyama et al., 2012). Further study to monitor and characterize foodborne pathogens from food products and food processing facilities in importing countries may be required to reduce foodborne pathogen transmission.

Several molecular-based studies of antibiotic resistance determinants have shown that ceftriaxone-resistant NTS strains mostly harbor *bla*_{CMY}, or *bla*_{SHV}-type genes with *bla*_{TEM-1} (Chen et al., 2004; Koh et al., 2008; Su et al., 2005). In this study, the resistance phenotypes of NTS strains and their transconjugants to ampicillin and ceftriaxone are most likely mediated by harboring and transferring *bla*_{TEM-1} with or without *bla*_{CTX-M-9} (Table 3). *bla*_{TEM-1} was also detected from all β -lactam resistant NTS strains and the transconjugants (data not shown). Combined, these data suggest that the resistance of the NTS strains to extended-spectrum β -lactams was mostly transferred to other enteric bacteria by plasmid-mediated *bla*_{TEM-1}. In this study, conjugation was performed at 37 °C to transfer plasmid harboring β -lactam resistant genes between NTS and another enteric bacterium. Consequently, a transconjugant was not obtained from PSS_984 harbored an IncHI1 plasmid. Previous studies (Sherburne et al., 2000; Phan et al., 2009; Taylor, 2009) have shown that the transfer of IncHI plasmids between 22 to 28 °C presented higher frequencies than 37 °C. Conjugation at 37 °C used in this study may be a primary reason for failure of horizontal gene transfer. Our data may support the suggestions that IncHI1 plasmid may disseminate antibiotic resistance to other bacteria in natural environments but not food-producing animal (Maher and Taylor, 1993; Taylor, 2009). In addition, PSS_984 was recovered from ginger powder. Four strains (PSS_903, PSS_913, PSS_928, and PSS_959) were not typed with the primer sets used. A further study for a new plasmid type on the unclassified plasmid(s) may be needed.

246 In conclusion, this study reports that NTS food isolates were contaminated with
247 conjugative plasmid-borne ESBLs, suggesting that ESBL-producing *Salmonella* in foods could
248 be a reservoir and a donor of ESBL genes (Egervarn et al., 2014; Sjölund-Karlsson et al., 2013).
249 The emergence and acquisition of ESBLs may attenuate the effectiveness of β -lactam antibiotics
250 used in the treatment of salmonellosis. Therefore, we suggest that our monitoring data for the
251 prevalence of ESBL-producing NTS from imported food products, combined with molecular
252 characterization of extended-spectrum β -lactam-resistance determinants in the NTS, provide
253 information that can contribute significantly to enhance food safety and promote public health.

Acknowledgements

We thank Drs. Carl E. Cerniglia and John B. Sutherland for critical review of the manuscript. The authors would like to thank Christine Summage-West for technical help and Gwendolyn Anderson and Stephanie Horton for *Salmonella* isolates.

The views presented in this article do not necessarily reflect those of the US Food and Drug Administration. This research was supported in part by an appointment to the Research Participation Program at the National Center for Toxicological Research administered by the Oak Ridge Institute for Science and Education through an interagency agreement between the U.S. Department of Energy and the U.S. Food and Drug Administration.

References

- Akiyama, T., and Khan, A. A. 2011. Molecular characterization of strains of fluoroquinolone-resistant *Salmonella enterica* serovar Schwarzengrund carrying multidrug resistance isolated from imported foods. *J. Antimicrob Chemother.* 67, 101-110.
- Akiyama, T., Khan, A. A., Cheng, C., and Stefanova, R. 2011. Molecular Characterization of *Salmonella enterica* serovar Saintpaul isolated from imported seafood, environmental and clinical samples. *Food Microbiol.* 28, 1124-1128.
- Akiyama, T., and Khan, A. A. 2012. Isolation and characterization of small qnrS1-carrying plasmids from imported seafood isolates of *Salmonella enterica* that are highly similar to plasmids of clinical isolates. *FEMS Immunol Med Microbiol.* 64, 429-432.
- Akiyama T., Khan, A. A., Nawaz, M. S., and Summage-West, C. 2012. Molecular characterization of high-level ciprofloxacin-resistant *Salmonella enterica* with multiple antibiotic resistance and class 1 integrons isolated from imported foods. *Int J Antimicrob Agents.* 39, 263-264.
- Al-Mashhadani, M., Hewson, R., Vivancos, R., Keenan, A., Beeching, N.J., Wain, J., Parry, C.M., 2011. Foreign travel and decreased ciprofloxacin susceptibility in *Salmonella enterica* infections. *Emerg Infect Dis.* 17, 123-125.
- Carattoli, A., Bertini, A., Villa, L., Falbo, V., Hopkins, K.L., Threlfall, E.J., 2005. Identification of plasmids by PCR-based replicon typing. *J Microbiol Methods* 63, 219-228.
- CDC, 2011. Vital signs: incidence and trends of infection with pathogens transmitted commonly through food--foodborne diseases active surveillance network, 10 U.S. sites, 1996-2010. *MMWR Morb Mortal Wkly Rep* 60, 749-755.
- CDC. 2013a. Foodborne Outbreak Online database (Food). Available at <http://wwwn.cdc.gov/foodborneoutbreaks/Default.aspx>.
- CDC, 2013b. National Antimicrobial Resistance Monitoring System for Enteric Bacteria (NARMS): Human Isolates Final Report, 2011. U.S. Department of Health and Human Services, CDC, Atlanta, Georgia.
- CDC, 2015. Foodborne Outbreak Online Database. Atlanta, GA: US Department of Health and Human Services, CDC; 2013. Available at <http://wwwn.cdc.gov/foodborneoutbreaks/default.aspx>.
- Chai, S.J., White, P.L., Lathrop, S.L., Solghan, S.M., Medus, C., McGlinchey, B.M., Tobin-D'Angelo, M., Marcus, R., Mahon, B.E., 2012. *Salmonella enterica* serotype Enteritidis: increasing incidence of domestically acquired infections. *Clin Infect Dis* 54 Suppl 5, S488-497.

- Chen, S., Zhao, S., White, D.G., Schroeder, C.M., Lu, R., Yang, H., McDermott, P.F., Ayers, S., Meng, J., 2004. Characterization of multiple-antimicrobial-resistant *Salmonella* serovars isolated from retail meats. *Appl Environ Microbiol* 70, 1-7.
- CLSI, 2014. Performance standards for antimicrobial susceptibility testing; twenty-fourth informational supplement (M100-S24). Clinical and Laboratory Standards Institute, Wayne, PA.
- Crim, S.M., Iwamoto, M., Huang, J.Y., Griffin, P.M., Gilliss, D., Cronquist, A.B., Cartter, M., Tobin-D'Angelo, M., Blythe, D., Smith, K., Lathrop, S., Zansky, S., Cieslak, P.R., Dunn, J., Holt, K.G., Lance, S., Tauxe, R., Henao, O.L., Centers for Disease, C., Prevention, 2014. Incidence and trends of infection with pathogens transmitted commonly through food--Foodborne Diseases Active Surveillance Network, 10 U.S. sites, 2006-2013. *MMWR Morb Mortal Wkly Rep* 63, 328-332.
- Dahshan, H., Shahada, F., Chuma, T., Moriki, H., Okamoto, K., 2010. Genetic analysis of multidrug-resistant *Salmonella enterica* serovars Stanley and Typhimurium from cattle. *Vet Microbiol* 145, 76-83.
- Dunne, E.F., Fey, P.D., Kludt, P., Reporter, R., Mostashari, F., Shillam, P., Wicklund, J., Miller, C., Holland, B., Stamey, K., Barrett, T.J., Rasheed, J.K., Tenover, F.C., Ribot, E.M., Angulo, F.J., 2000. Emergence of domestically acquired ceftriaxone-resistant *Salmonella* infections associated with AmpC beta-lactamase. *JAMA* 284, 3151-3156.
- Egervarn, M., Borjesson, S., Byfors, S., Finn, M., Kaibe, C., Englund, S., Lindblad, M., 2014. *Escherichia coli* with extended-spectrum beta-lactamases or transferable AmpC beta-lactamases and *Salmonella* on meat imported into Sweden. *Int J Food Microbiol* 171, 8-14.
- Fey, P.D., Safranek, T.J., Rupp, M.E., Dunne, E.F., Ribot, E., Iwen, P.C., Bradford, P.A., Angulo, F.J., Hinrichs, S.H., 2000. Ceftriaxone-resistant *Salmonella* infection acquired by a child from cattle. *N Engl J Med* 342, 1242-1249.
- Gebreyes, W.A., Thakur, S., 2005. Multidrug-resistant *Salmonella enterica* serovar Muenchen from pigs and humans and potential interserovar transfer of antimicrobial resistance. *Antimicrob Agents Chemother* 49, 503-511.
- Grimm, V., Ezaki, S., Susa, M., Knabbe, C., Schmid, R.D., Bachmann, T.T., 2004. Use of DNA microarrays for rapid genotyping of TEM beta-lactamases that confer resistance. *J Clin Microbiol* 42, 3766-3774.
- Grobner, S., Linke, D., Schutz, W., Fladerer, C., Madlung, J., Autenrieth, I.B., Witte, W., Pfeifer, Y., 2009. Emergence of carbapenem-non-susceptible extended-spectrum beta-lactamase-producing *Klebsiella pneumoniae* isolates at the university hospital of Tübingen, Germany. *J Med Microbiol* 58, 912-922.

- Hellerstein, S., 2003. Antibiotic treatment for urinary tract infections in pediatric patients. *Minerva Pediatr* 55, 395-406.
- Hendriksen, R.S., Mikoleit, M., Carlson, V.P., Karlslose, S., Vieira, A.R., Jensen, A.B., Seyfarth, A.M., DeLong, S.M., Weill, F.X., Lo Fo Wong, D.M., Angulo, F.J., Wegener, H.C., Aarestrup, F.M., 2009. WHO Global Salm-Surv external quality assurance system for serotyping of *Salmonella* isolates from 2000 to 2007. *J Clin Microbiol* 47, 2729-2736.
- Hohmann, E.L., 2001. Nontyphoidal salmonellosis. *Clin Infect Dis* 32, 263-269.
- Jackson, B.R., Griffin, P.M., Cole, D., Walsh, K.A., Chai, S.J., 2013. Outbreak-associated *Salmonella enterica* serotypes and food commodities, United States, 1998-2008. *Emerg Infect Dis* 19, 1239-1244.
- Khan, A. A., Ponce, E., Nawaz, M., Cheng, Chong-Ming, Khan, J. A., and Summace-West, C. 2009, Identification and Characterization of Class 1 Integron Resistance Gene Cassettes among *Salmonella* strains from imported seafood, *Appl Environ Microbiol* 75, 1192-1196.
- Klaenhammer, T.R., 1984. A general method for plasmid isolation in lactobacilli. *Current Microbiology* 10, 23-28.
- Koh, T.H., Koh, A.E., Hamdan, A., Khoo, B.C., Yu, V.Y., Raymond, R.T., Tee, N.W., 2008. Ceftriaxone-resistant *Salmonella* spp. in Singapore. *Ann Acad Med Singapore* 37, 900-901.
- Maher, D., Sherburne, R., Taylor, D.E., 1993. H-pilus assembly kinetics determined by electron microscopy. *J Bacteriol* 175, 2175-2183.
- Maynard, C., Fairbrother, J.M., Bekal, S., Sanschagrin, F., Levesque, R.C., Brousseau, R., Masson, L., Lariviere, S., Harel, J., 2003. Antimicrobial resistance genes in enterotoxigenic *Escherichia coli* O149:K91 isolates obtained over a 23-year period from pigs. *Antimicrob Agents Chemother* 47, 3214-3221.
- Mezal, E., Stefanova, R., and Khan, A.A., 2013. Isolation and Molecular Characterization of *Salmonella enterica* serovar Javiana from Food, Environmental and Clinical Samples. *Int J Food Microbiol.* 164, 113-118.
- Paterson, D.L., Hujer, K.M., Hujer, A.M., Yeiser, B., Bonomo, M.D., Rice, L.B., Bonomo, R.A., International Klebsiella Study, G., 2003. Extended-spectrum beta-lactamases in *Klebsiella pneumoniae* bloodstream isolates from seven countries: dominance and widespread prevalence of SHV- and CTX-M-type beta-lactamases. *Antimicrob Agents Chemother* 47, 3554-3560.

- Perez-Perez, F.J., Hanson, N.D., 2002. Detection of plasmid-mediated AmpC beta-lactamase genes in clinical isolates by using multiplex PCR. *J Clin Microbiol* 40, 2153-2162.
- Phan, M.D., Kidgell, C., Nair, S., Holt, K.E., Turner, A.K., Hinds, J., Butcher, P., Cooke, F.J., Thomson, N.R., Titball, R., Bhutta, Z.A., Hasan, R., Dougan, G., Wain, J., 2009. Variation in *Salmonella enterica* serovar typhi IncHI1 plasmids during the global spread of resistant typhoid fever. *Antimicrob Agents Chemother* 53, 716-727.
- Ponce, E., Khan, A. A., Cheng, Chong-Ming, Summage-West, C., and Cerniglia, C. E. 2008. Prevalence and characterization of *Salmonella enterica* serovar Weltevreden from imported seafood. *Food Microbiol.* 25, 29-35.
- Scallan, E., Hoekstra, R.M., Angulo, F.J., Tauxe, R.V., Widdowson, M.A., Roy, S.L., Jones, J.L., Griffin, P.M., 2011. Foodborne illness acquired in the United States--major pathogens. *Emerg Infect Dis* 17, 7-15.
- Sherburne, C.K., Lawley, T.D., Gilmour, M.W., Blattner, F.R., Burland, V., Grotbeck, E., Rose, D.J., Taylor, D.E., 2000. The complete DNA sequence and analysis of R27, a large IncHI plasmid from *Salmonella* typhi that is temperature sensitive for transfer. *Nucleic Acids Res* 28, 2177-2186.
- Simarro, E., Navarro, F., Ruiz, J., Miro, E., Gomez, J., Mirelis, B., 2000. *Salmonella enterica* serovar Virchow with CTX-M-like beta-lactamase in Spain. *J Clin Microbiol* 38, 4676-4678.
- Sjölund-Karlsson, M., Howie, R.L., Blickenstaff, K., Boerlin, P., Ball, T., Chalmers, G., Duval, B., Haro, J., Rickert, R., Zhao, S., Fedorka-Cray, P.J., Whichard, J.M., 2013. Occurrence of beta-lactamase genes among non-Typhi *Salmonella enterica* isolated from humans, food animals, and retail meats in the United States and Canada. *Microb Drug Resist* 19, 191-197.
- Skov, M.N., Andersen, J.S., Aabo, S., Ethelberg, S., Aarestrup, F.M., Sorensen, A.H., Sorensen, G., Pedersen, K., Nordentoft, S., Olsen, K.E., Gerner-Smidt, P., Baggesen, D.L., 2007. Antimicrobial drug resistance of *Salmonella* isolates from meat and humans, Denmark. *Emerg Infect Dis* 13, 638-641.
- Soler, P., Gonzalez-Sanz, R., Bleda, M.J., Hernandez, G., Echeita, A., Usera, M.A., 2006. Antimicrobial resistance in non-typhoidal *Salmonella* from human sources, Spain, 2001-2003. *J Antimicrob Chemother* 58, 310-314.
- Su, L.H., Wu, T.L., Chia, J.H., Chu, C., Kuo, A.J., Chiu, C.H., 2005. Increasing ceftriaxone resistance in *Salmonella* isolates from a university hospital in Taiwan. *J Antimicrob Chemother* 55, 846-852.
- Tauxe, R.V., 1991. *Salmonella*: A postmodern pathogen. *J food prot* 54, 563-568.

- Taylor, D.E., 2009. Thermosensitive nature of IncHI1 plasmid transfer. *Antimicrob Agents Chemother* 53, 2703.
- Tomasz, A., 1979. The mechanism of the irreversible antimicrobial effects of penicillins: how the beta-lactam antibiotics kill and lyse bacteria. *Annu Rev Microbiol* 33, 113-137.
- Tran, Q.T., Mahendran, K.R., Hajjar, E., Ceccarelli, M., Davin-Regli, A., Winterhalter, M., Weingart, H., Pages, J.M., 2010. Implication of porins in beta-lactam resistance of *Providencia stuartii*. *J Biol Chem* 285, 32273-32281.
- USDA-ERS, 2015. Import shares of consumption using the volumn method in U. S. Agricultural Trade database. Available at <http://www.ers.usda.gov/topics/international-markets-trade/us-agricultural-trade/import-share-of-consumption.aspx>.
- USFDA/CFSAN, 2014. Bacteriological Analytical Manual (BAM) Chapter 5. U.S. Food and Drug Administration, Silver Spring, MD.
- Van Doren, J.M., Blodgett, R.J., Pouillot, R., Westerman, A., Kleinmeier, D., Ziobro, G.C., Ma, Y., Hammack, T.S., Gill, V., Muckenfuss, M.F., Fabbri, L., 2013. Prevalence, level and distribution of *Salmonella* in shipments of imported capsicum and sesame seed spice offered for entry to the United States: observations and modeling results. *Food Microbiol* 36, 149-160.
- Van Doren, J.M., Kleinmeier, D., Hammack, T.S., Westerman, A., 2013. Prevalence, serotype diversity, and antimicrobial resistance of *Salmonella* in imported shipments of spice offered for entry to the United States, FY2007-FY2009. *Food Microbiol* 34, 239-251.
- Wise, E.M., Jr., Park, J.T., 1965. Penicillin: its basic site of action as an inhibitor of a peptide cross-linking reaction in cell wall mucopeptide synthesis. *Proc Natl Acad Sci U S A* 54, 75-81.
- Yates, C., Amyes, S., 2005. Extended-spectrum beta-lactamases in non-typhoidal *Salmonella* spp. isolated in the UK are now a reality: why the late arrival? *J Antimicrob Chemother* 56, 262-264.
- Zhao, S., Datta, A.R., Ayers, S., Friedman, S., Walker, R.D., White, D.G., 2003. Antimicrobial-resistant *Salmonella* serovars isolated from imported foods. *Int J Food Microbiol* 84, 87-92.
- Zhao, S., McDermott, P.F., Friedman, S., Qaiyumi, S., Abbott, J., Kiessling, C., Ayers, S., Singh, R., Hubert, S., Sofos, J., White, D.G., 2006. Characterization of antimicrobial-resistant *Salmonella* isolated from imported foods. *J Food Prot* 69, 500-507.
- Zhao, S., White, D.G., Friedman, S.L., Glenn, A., Blickenstaff, K., Ayers, S.L., Abbott, J.W., Hall-Robinson, E., McDermott, P.F., 2008. Antimicrobial resistance in *Salmonella*

483 *enterica* serovar Heidelberg isolates from retail meats, including poultry, from 2002 to
484 2006. Appl Environ Microbiol 74, 6656-6662.
485

Table 1. Primer sequences used in the study

Genes	Primer Sequence (5'-3')		Amplicon size (bp)	References
	Forward	Reverse		
For extended-spectrum β-lactamase resistance				
<i>bla</i> _{OXA-1}	GCAGCGCCAGTGCATCAAC	CCGCATCAAATGCCATAAGTG	888	(Maynard et al., 2003)
<i>bla</i> _{PSE-1}	AGTAGGGCAGGCAATCACAC	GCGATCCGCAATGTTCCATC	421	This study
<i>bla</i> _{SHV-1}	GCAAAACGCCGGGTATTTC	GGTTAGCGTTGCCAGTGCT	940	(Grobner et al., 2009)
<i>bla</i> _{TEM-1}	ATGAGTATTCAACATTTCCG	TTAATCAGTGAGGCACCTAT	851	(Grimm et al., 2004)
<i>bla</i> _{CTX-M}	CGCTTTGCGATGTGCAG	ACCGCGATATCGTTGGT	550	(Paterson et al., 2003)
<i>bla</i> _{CTX-M-9}	GTGACAAAGAGAGTGCAACGG	ATGATTCTCGCCGCTGAAGCC	856	(Simarro et al., 2000)
<i>bla</i> _{CMY-2}	TGGCCAGAACTGACAGGCAAA	TTTCTCCTGAACGTGGCTGGC	462	(Perez-Perez and Hanson, 2002)
<i>bla</i> _{FOX}	AACATGGGGTATCAGGGAGATG	CAAAGCGCGTAACCGGATTGG	190	(Perez-Perez and Hanson, 2002)
<i>bla</i> _{IMP}	CATGGTTTGGTGGTTCTTGT	ATAATTTGGCGGACTTTGGC	447	(Grobner et al., 2009)
<i>bla</i> _{KPC}	CAGCTCATTCAAGGGCTTTC	AGTCATTTGCCGTGCCATAC	533	(Grobner et al., 2009)
<i>bla</i> _{VIM}	AGTGGTGAGTATCCGACAG	ATGAAAGTGCGTGGAGAC	261	(Grobner et al., 2009)
For replicon typing				
A/C	GAGAACCAAAGACAAAGACCTGGA	ACGACAAACCTGAATTGCCTCCTT	465	(Carattoli et al., 2005)
B/O	GCGGTCCGGAAGCCAGAAAAC	TCTGCGTTCGCCAAGTTCGA	159	(Carattoli et al., 2005)
FIA	CCATGCTGGTTCTAGAGAAGGTG	GTATATCCTTACTGGCTTCCGCAG	462	(Carattoli et al., 2005)
FIC	GTGAACTGGCAGATGAGGAAGG	TTCTCCTCGTCGCCAAACTAGAT	262	(Carattoli et al., 2005)
FIIA /FIIS	CTGTGCTAAGCTGATGGC	CTCTGCCACAAACTTCAGC	270	(Carattoli et al., 2005)
FIB	GGAGTTCTGACACACGATTTTCTG	CTCCCGTCGCTTCAGGGCATT	702	(Carattoli et al., 2005)
HI1	GGAGCGATGGATTACTTCAGTAC	TGCCGTTTCACCTCGTGAGTA	471	(Carattoli et al., 2005)
HI2	TTTCTCCTGAGTCACCTGTTAACAC	GGCTCACTACCGTTGTCATCCT	644	(Carattoli et al., 2005)
I1	CGAAAGCCGGACGGCAGAA	TCGTCGTTCCGCCAAGTTCGT	139	(Carattoli et al., 2005)
K/B	GCGGTCCGGAAGCCAGAAAAC	TCTTTCACGAGCCCGCCAAA	160	(Carattoli et al., 2005)
L/M	GGATGAAAACATCAGCATCTGAAG	CTGCAGGGGCGATTCTTTAGG	785	(Carattoli et al., 2005)
N	GTCTAACGAGCTTACCGAAG	GTTTCAACTCTGCCAAGTTC	559	(Carattoli et al., 2005)
T	TTGGCCTGTTTGTGCCTAAACCAT	CGTTGATTACACTTAGCTTTGGAC	750	(Carattoli et al., 2005)
W	CCTAAGAACAACAAAGCCCCCG	GGTGCGGGCATAGAACCGT	242	(Carattoli et al., 2005)
X	AACCTTAGAGGCTATTTAAGTTGCTG AT	TGAGAGTCAATTTTTATCTCATGTTTT AGC	376	(Carattoli et al., 2005)

Table 2. *Salmonella enterica* isolated from imported food products during 2011 to 2013 by an US FDA's regional laboratory

by serovar	<i>S. Welteverden</i>		<i>S. Newport</i>	<i>S. Senftenberg</i>	<i>S. Virchow</i>	<i>S. Enteritidis</i>	<i>S. Bareilly</i>	<i>S. Typhimurium</i>	Others ¹			
	21 (19.1) ²		9 (8.2)	9 (8.2)	7 (6.4)	6 (5.5)	5 (4.5)	5 (4.5)	48 (43.6)			
by country	India	Taiwan	Vietnam	Indonesia	Mexico	Thailand	China	Philippines	Pakistan	Australia	Bangladesh	Others ³
	19 (17.3)	19 (17.3)	17 (15.5)	13 (11.8)	8 (7.3)	8 (7.3)	5 (4.5)	4 (3.6)	3 (2.7)	2 (1.8)	2 (1.8)	10 (9.1)
by food source	Seafood		Freshwater fishes			Vegetables		Fruits		Others ⁴		
	49 (44.5)		19 (17.3)			31 (28.2)		7 (6.4)		4 (3.6)		

¹ include *Salmonella enterica* serovar Litchfield, Agoveve, Nchanga, Oranienburg, Paratyphi B, Rubislaw, Stanley, Javiana, Anatum, Aschersleben, Braenderup, Carmel, Cerro, Chester, Cubana, Farmsen, Fischerstrasse, Give, Giza, Hvittingfoss, Indiana, Infantis, Lansing, Livingstone, London, Lucianna, Michigan, Orion, Oslo, Panama, Pomona, Putten, Richmond, Rissen, and Saintpaul. Four isolates were not serotypable.

² presents number and percentage (%) of total 110 food isolates.

³ include Denmark, France, Guatemala, Guyana, Ivory Coast, Lebanon, Malaysia, Turkey, and Panama. An isolate was unknown.

⁴ include animal and dairy foods.

Table 3

Table 3. Epidemiologic and molecular characterization of extended-spectrum β -lactamase producing *S. enterica* food isolates

ID	Serotype	Food Product	Year	Country	Extended-spectrum β -lactam resistance ¹	Extended-spectrum β -lactam resistance determinants	Plasmid size (Kbp)	Replicon Type
PSS_903	<i>S. ser. Enteritidis</i>	Frozen octopus	2011	Indonesia	AMP	<i>bla</i> _{TEM-1}	55, 35, <8	- ²
PSS_913	<i>S. ser. Indiana</i>	Frozen tilapia	2011	China	AMP, CET, FOX, CRO, FEP	<i>bla</i> _{TEM-1} , <i>bla</i> _{CTX-M-9}	150, 80, 25	-
PSS_928	<i>S. ser. Weltevreden</i>	Frozen whole blue anchovy	2012	Vietnam	AMP	<i>bla</i> _{TEM-1} , <i>bla</i> _{CTX-M-9}	115, 95, 25	-
PSS_948	<i>S. ser. Newport</i>	Frozen whole snakehead	2012	Vietnam	AMP	<i>bla</i> _{TEM-1}	>165, 50, 25	IncN
PSS_955	<i>S. ser. Typhimurium</i>	Frozen tilapia whole	2012	Taiwan	AMP	<i>bla</i> _{TEM-1} , <i>bla</i> _{OXA-1}	55, 35, 13, 11, 9, <8	IncI1
PSS_959	<i>S. ser. Enteritidis</i>	Szechuan pepper	2012	China	AMP	<i>bla</i> _{TEM-1}	50, 35	-
PSS_970	<i>S. ser. Litchfield</i>	Frozen farm raised perch	2012	Taiwan	AMP	<i>bla</i> _{TEM-1}	90	IncI1
PSS_984	<i>S. ser. Give</i>	Ginger powder	2012	Vietnam	AMP	<i>bla</i> _{TEM-1}	50, <8	IncHI1, X
PSS_988	<i>S. ser. Enteritidis</i>	White pepper powder	2013	Taiwan	AMP, CET, CRO, FEP	<i>bla</i> _{TEM-1}	80, 70, 38	IncI1, N

¹ Extended-spectrum β -lactam antibiotics: AMP, ampicillin; CET, cephalothin; FOX, cefoxitin; CRO, ceftriaxone; FEP, cefepime

² - represents that a replicon with primer sets used in a PCR-based replicon typing assay was not detected.

Table 4. MICs (mg/L) of *Salmonella* isolates and *E.coli* J53 transconjugants to ampicillin and cephalosporins and the conjugative plasmid size

ID	Ampicillin	Cephalothin	Cefoxitin	Ceftriaxone	Cefepime	Plasmid (Kbp)
PSS_903	> 64	16	16	<1	<1	55, 30, <8
¹ J53_903	> 64	16	8	<1	<1	30
PSS_913	> 64	>1024	32	>1024	128	150, 80, 25
J53_913	> 64	>1024	8	64	2	150, 80, 25
PSS_928	> 64	8	2	<1	<1	115, 95, 25
J53_928	> 64	64	64	<1	<1	115, 95, 25
PSS_948	> 64	8	8	<1	<1	>165, 50, 25
J53_948	> 64	32	32	<1	<1	50
PSS_955	> 64	4	4	<1	<1	55, 35, 13, 11, 9, <8
J53_955	> 64	64	64	<1	<1	13, 9
PSS_959	> 64	8	4	<1	<1	50, 38
J53_959	> 64	8	16	<1	<1	38
PSS_970	> 64	16	4	<1	<1	90
J53_970	> 64	16	8	<1	<1	90
PSS_984 ²	> 64	8	4	<1	<1	50, <8
PSS_988	> 64	>1024	8	512	32	80, 70, 38
J53_988	> 64	>1024	8	32	2	80, 70, 38

¹ represents *E.coli* J53 transconjugants.

² *S. Give* J53_984 transconjugant resistant to β -lactams was not obtained.

Figure 1

Figure Legends

Figure 1. Plasmid profiles and Southern blots of NTS isolates and transconjugants of *E. coli* J53 resistant to β -lactam antibiotics. (A) Plasmid profiles of the 9 NTS isolates; (B) Southern hybridization of the 9 NTS isolates; (C) *E. coli* J53 and its transconjugants using a *bla*_{TEM-1} probe. J53_984 transconjugants are not shown in Fig. 1 C, since the resistance of PSS_984 was not transferred to *E. coli* J53.