1 How do rubber (*Hevea brasiliensis*) plantations behave under seasonal water stress in
2 northeastern Thailand and central Cambodia?
3 Tomo'omi Kumagai^{a,*}tomoomikumagai@gmail.com, Ryan G. Mudd^b, Thomas W.
4 Giambelluca^b, Nakako Kobayashi^a, Yoshiyuki Miyazawa^c, Tiva Khan Lim^d, Wen Liu^b,
5 Maoyi Huang^e, Jefferson M. Fox^f, Alan D. Ziegler^g, Song Yin^d, Sophea Veasna Mak^d,
6 Poonpipope Kasemsap^h
7 ^aHydrospheric Atmospheric Research Center, Nagoya University, Chikusa-ku, Nagoya
8 464-8601, Japan
9 ^bDepartment of Geography, University of Hawai'i at Mānoa, Honolulu, HI 96822, USA
10 ^cResearch Institute for East Asia Environments, Kyushu University, Motooka, Fukuoka
11 811-0395, Japan
12 ^dCambodian Rubber Research Institute, Phnom Penh, Cambodia
13 ^eAtmospheric Sciences and Global Change Division, Pacific Northwest National Laboratory,
14 Richland, WA 99352, USA
15 ^fEast-West Center, Honolulu, HI 96848, USA
16 ^gDepartment of Geography, National University of Singapore, Singapore 117570
17 ^hDepartment of Horticulture, Faculty of Agriculture, Kasetsart University, Bangkok 10900,
18 Thailand
19 *Corresponding author at: Hydrospheric Atmospheric Research Center, Nagoya University,
20 Furo-cho, Chikusa-ku, Nagoya, Aichi 464-8601, Japan. Tel.: +81 52 789 3478; Fax: +81 52
21 788 6206.
22

28 **Abstract**

29 Delineating the characteristics of biosphere-atmosphere exchange in rubber (*Hevea*
30 *brasiliensis* Müll. Arg.) plantations, which are rapidly expanding throughout mainland
31 Southeast Asia, is important to understanding the impacts of the land-use change on
32 environmental processes. In attempt to shed new light on the impacts of conversion to rubber,
33 we have conducted eddy flux measurements over a 3-year period in two rubber plantation
34 sites: (1) Som Sanuk, located in northeastern Thailand; and (2) Cambodian Rubber Research
35 Institute (CRRRI), located in central Cambodia. Both sites have a distinct dry season. We used
36 a combination of actual evapotranspiration (E_T) flux measurements and an inverted version
37 of a simple 2-layer E_T model for estimating the mean canopy stomatal conductance (g_s). The
38 potential water balance (precipitation (P) – potential evaporation (E_{T_POT})) for each season
39 (i.e., December-February: DJF, March-May: MAM, June-August: JJA, and
40 September-November: SON) revealed when and how the water use is controlled. In the
41 seasons when actual water balance ($P - E_T$) was negative (DJF and MAM), the deficit was
42 compensated with soil water from the previous season at depths of 0-2 m (Thailand site) and
43 0-3 m (Cambodia site). At both sites, the reference value of g_s (g_{sref}) and the sensitivity of g_s
44 to atmospheric demand (m) appeared to be less in DJF and MAM than each in the other two
45 3-month periods (seasons). On average, in a whole year, m/g_{sref} was less in Thailand ($<< 0.6$)
46 than in Cambodia (near 0.6 for part of the year), suggesting that there was less sufficient
47 stomatal regulation at the Thailand site, where there might be little risk of water
48 stress-induced hydraulic failure because of its higher annual rainfall amount. In comparison,
49 at CRRRI where annual $P - E_{T_POT}$ was negative, there was stricter stomatal regulation,
50 preventing excessive xylem cavitation.

51

52 *Keywords:* Tropical deciduous trees; Stomatal control; Transpiration; Water vapor flux;

53 Eddy covariance; Soil moisture

54

54 1. Introduction

55 Rubber (*Hevea brasiliensis* Müll. Arg.) is among the major economic tree crops of tropical
56 areas throughout the world, especially Southeast Asia. The Association of Natural Rubber
57 Producing Countries (ANRPC, 2010) estimated that from 2003 to 2010 more than 1,500,000
58 ha of land were converted to rubber in southern China, Thailand, Vietnam, and Cambodia
59 (see Fig.1). Rubber is currently expanding rapidly in non-traditional areas of Myanmar, Laos,
60 and India. In the uplands of mainland Southeast Asia, about 9% of the current vegetation,
61 including evergreen broadleaf trees and swidden-related secondary vegetation, is projected
62 to be replaced by tree plantations and diversified farming systems by 2050 (Fox et al., 2012).
63 Almost half of this transition is expected to be due to the expansion of rubber cultivation, the
64 most rapidly expanding tree crop in the region, resulting in a projected fourfold increase in
65 land area devoted to rubber plantation (Fox et al., 2012, 2014).

66 *Hevea* rubber originates from the tropical rainforests of the Amazon in South America.
67 Much of the Southeast Asia growing areas are outside the native range, which is intrinsically
68 warm and moist throughout the year (Priyadarshan, 2003; Priyadarshan et al., 2005).
69 However, selective breeding of *Hevea* clones has facilitated the expansion of rubber
70 plantations into relatively cool and dry environments, such as those found in montane and
71 monsoonal regions of mainland Southeast Asia. Rubber is now cultivated profitably in many
72 “non-traditional” areas previously determined as unsuitable (Qiu, 2009).

73 Numerous potential negative ecological and environmental consequences of converting
74 primary and secondary forests into rubber plantations have been suggested: decrease in
75 biodiversity; reduction of total biomass carbon; alteration of the hydrological regime; and
76 acceleration of erosion (Wu et al., 2001; Mann, 2009; Qiu, 2009; Ziegler et al., 2009a, b;

77 Guardiola-Claramonte et al., 2010; Tan et al., 2011; Ziegler et al., 2012). On the other hand,
78 some studies reported that rubber plantations replacing non-forested lands could act as a
79 carbon sinks (Wauters et al., 2008). Further, while eddy covariance measurements in rubber
80 revealed high plantation water use and a large proportion of energy being used to drive
81 evapotranspiration for most of the year (Tan et al., 2011, Giambelluca et al., submitted for
82 publication, hereinafter referred to as submitted manuscript, 2015), the results of sapflow
83 measurements of individual rubber trees showed water use of rubber plantations may not be
84 as high as anticipated (Isarangkool Na Ayutthaya et al., 2011; Carr, 2012; Kobayashi et al.,
85 2014), but these findings might be related to differences in regional climate, geography,
86 physiological adaptations, and measurement method. Of concern to our research, large-scale
87 land use changes to rubber plantations may have important implications for local-to-regional
88 carbon and water balances, but they still remain unclear. Hindering the ability to predict and
89 understand these changes is lack of information on the characteristics of mass (i.e., CO₂ and
90 H₂O) exchange between rubber plantation canopy and atmosphere, which is the subject of
91 this investigation.

92 The region encompassing northeastern Thailand and central Cambodia has strong
93 seasonality in precipitation (see Fig. 3 in Kumagai et al. (2005)): well-defined wet and dry
94 seasons are clearly apparent. Although seasonality in atmospheric variables such as air
95 temperature and humidity affect the hydrologic fluxes above and within vegetation,
96 fluctuations in soil moisture and plant water status, influenced by precipitation variations,
97 are the most important controls on the hydrological, atmospheric and ecological processes
98 within the Soil–Plant–Atmosphere Continuum (SPAC) (see Kumagai et al., 2009b). Thus, it
99 is logical to expect that current rubber plantations in northeastern Thailand and central

100 Cambodia must cope with strong seasonal water stress via the SPAC context, such as
101 limiting tree water use by regulating stomatal behavior in response to soil moisture deficit,
102 and compensating for lack of water near the surface during the dry season by exploiting
103 deeper soil water. Further, it is important to note that a previous study (Sangsing et al., 2003)
104 found different rubber clones exhibit varying degrees of adaptability to water stress, thus it
105 would be expected that water use efficiency and risk of hydraulic fracture vary with clone
106 type in a particular climate.

107 Environmental control of canopy mass exchange and tree water use can be characterized
108 in terms of the relationship between canopy conductance (G_C) and environmental factors.
109 Canopy conductance is thought to result from the bulk leaf stomatal regulations within the
110 vegetation canopy; and it is generally calculated from the inverted Penman-Monteith
111 equation with measurements of canopy transpiration (e.g., Wullschleger et al., 2000; Delzon
112 et al., 2004; Kumagai et al., 2004, 2008; Herbst et al., 2008; Kumagai et al., 2009a; Tateishi
113 et al., 2010). Because both CO_2 and H_2O exchanges are regulated by stomata, G_C is a major
114 index of the exchange efficiency and environmental control. However, estimation from
115 above-canopy evapotranspiration (E_T), such as observed using eddy covariance system and
116 above-canopy environmental factors, leads to surface conductance (G_S), not G_C . Briefly, G_C
117 reflects the physiological behavior of the canopy vegetation, while G_S also incorporates the
118 exchange characteristics of the understory vegetation and/or soil surface (see Kelliher et al.,
119 1995).

120 The present study is built on a previous study that revealed that the rubber E_T , obtained by
121 eddy flux measurements in northeastern Thailand and central Cambodia, is much higher than
122 regional E_T in large-scale estimates in mainland Southeast Asia (Giambelluca et al.,

submitted for publication). Therefore, our goal is to understand environmental controls on G_C related to the ecosystem water resources in the SPAC context, and how the studied rubber plantations can maintain such high E_T despite existence of strong seasonal water stress. In the present study, we go further and use the eddy fluxes measurements combined with a simple two-layer vegetation evaporation model (e.g., Kelliher et al., 1995) to separate G_C from the G_S . We principally investigate the differences in G_C and the environmental controls between the rubber plantation sites, as well as between rubber and other vegetation. This analysis represents an important step in understanding the impacts of the rapid expansion of rubber plantations on water and carbon cycling in the region.

2. Materials and methods

2.1. Site description

We present an analysis of canopy stomatal behavior of rubber trees at two sites in monsoonal regions of mainland Southeast Asia with contrasting rainfall regimes (Fig. 1): (1) Som Sanuk, Bueng Kan Province, northeast Thailand (18°11'N, 103°25'E, 210 m a.s.l.); and (2) Cambodian Rubber Research Institute (CRRI) in Kampong Cham Province, central Cambodia (11°57'N, 105°34'E, 57 m a.s.l.). For each site, we present the results for two complete years of measurements: February 2009–June 2011 at Som Sanuk; and September 2009–February 2012 at CRRI.

2.1.1. Som Sanuk site

The study site at Som Sanuk is centered within a 2.4-ha smallholder property planted with clone RRIM-600 rubber trees. The average slope of the terrain surrounding the

146 meteorological tower is less than 1°. The A horizon of the soil is very shallow (< 10 cm),
147 light gray in color, and has little organic material (< 2%). The dull yellow B horizon is 30-70
148 cm thick, sandy, and is 20-40 cm thicker along tree rows than between rows as a result of the
149 practice of mounding soil along each row before planting. Below the B horizon is a BW
150 horizon, approximately 85-cm thick, with ferralitic concretions. Few roots penetrate the BW
151 horizon, below which is found a gray/brown clay layer. During the two project years (March
152 2009-February 2010 and March 2010-February 2011), annual precipitation was 2215 and
153 2020 mm, and mean annual temperature was 26.2 and 26.0 °C, respectively. About 90% of
154 the total annual rainfall occurs during May to September. November to January is relatively
155 cool. Air temperature is generally highest in April. Wind is predominantly from the northeast
156 during October to April, and from the southwest during June through August. The climatic
157 characteristics at Som Sanuk are presented in Fig. 2.

158 Of the 1360 trees originally planted on the property in 1991, 1260 trees had survived after
159 20 years. Trees were planted with a spacing of 7.0 m × 2.5 m. The mean canopy height was
160 approximately 19 m and the mean diameter at 1.7 m height was 18.9 cm (standard deviation
161 (SD) = 3.0 cm) in January 2010. During 2010-2012, mean stem diameter growth rate was 0.6
162 cm year⁻¹. Approximately 1.2 kg tree⁻¹ of fertilizer (50/50 mix of manure and chemical
163 fertilizers, typically N-P-K: 20-10-12) is applied twice per year. Latex tapping was initiated
164 in about 1997 or 1998. With the exception of a few small riparian areas with shrub vegetation,
165 mature rubber trees form a continuous canopy in all directions for at least 500 m.

166

167 2.1.2. *Cambodian Rubber Research Institute site*

168 A 6.25-ha plot of clone RRIC-100 in an experimental rubber plantation within the 994-ha

169 Cambodian Rubber Research Institute (CRRRI) was used for the observations in central
170 Cambodia (see Kumagai et al., 2013). The experimental rubber plantation is on a level plain
171 set in red basaltic latosols. The soil texture is clay with about 10 % fine sand, 10 % coarse silt,
172 20 % fine silt and 60 % clay. The climatic characteristics at CRRRI are presented in Fig. 3. The
173 climate is governed by the Asian Monsoon, which produces two distinct seasons: a wet
174 season (approximately May-October) and a dry season (approximately November-April).
175 Annual precipitation in 2010 and 2011 was 1332 and 1545 mm, respectively. Rainy seasons
176 extended from late-May to late-November in 2010, and late-April to mid-November in 2011.
177 The mean annual temperature was significantly higher in 2010 (28.0 °C) than in 2011
178 (27.0 °C). For both years, maximum monthly averages occurred in April-May; minimums
179 occurred in December-January.

180 All the trees at the site were planted in 2004 (i.e., 8 years old in 2012) using a regular
181 spacing of 6-m in north-south direction and 3-m in east-west direction, resulting in a
182 potential tree density of 555 trees ha⁻¹. Due to tree mortality, approximately 20% of trees in
183 mature stands were missing, producing a tree density of 458 trees ha⁻¹. Chemical fertilizer
184 (N-P-K: 15-15-15) was used only during the first 4 years after planting. Latex tapping was
185 initiated in November, 2010. Stem diameters were measured at 1.7 m height above the
186 ground so as not to overlap the tapping panel, which starts at 1.3 m. Mean stem diameter was
187 13.3 cm (SD = 2.3 cm) in February 2010. This increased to 15.3 cm (SD = 2.3 cm) in March
188 2012. Mean canopy height was 11.4 m in January 2010, 12.9 m in March 2011, and 14.3 m in
189 March 2012. Various species of herbaceous plants and shrubs occupied the understory; and
190 they fully cover the ground except for along the tappers path.

191

192 2.2. *Meteorological measurements*

193 Canopy towers 26.5-m and 30-m-tall were constructed at Som Sanuk and CRRI,
194 respectively, for micrometeorological and eddy flux measurements. Air temperature,
195 humidity (HMP45C, Campbell Scientific, Logan, UT), and four-component radiation (at
196 Som Sanuk; CNR1, Kipp & Zonen, Delft, Netherlands, and at CRRI; NR01, Hukseflux,
197 Delft, Netherlands) sensors were installed at the top of each tower. At Som Sanuk, a rain
198 gauge (TE-525, Texas Electronics, Inc., Dallas, TX) was mounted at a height of 20 m on the
199 southwest side of the tower. At CRRI, two rain gauges (TE-525) were mounted at a height of
200 15 m on the southwest and northeast sides of the tower. To minimize bias from tower
201 interference and wind-related measurement error, rain gauges were mounted close to the
202 canopy top. At CRRI, rainfall from the upwind gauge was used based on the mean wind
203 direction observed in each 30-min interval. Samples were taken at 10 Hz for radiation, air
204 temperature, and humidity and averaged over 30 min. Precipitation (P) was measured by
205 counting tipping bucket tips (CR3000 and CR23X, Campbell Scientific) at an interval of
206 30-min.

207 Photosynthetic photon flux density (PPFD) below the canopy was measured below the
208 canopy at a height of approximately 3.0 m at Som Sanuk with line quantum sensors (SQ-311,
209 Apogee Instruments, Logan, UT). At CRRI, intensive observations on the transmitted PPFD
210 through the canopy were conducted at approximately 1-m above the forest floor by walking
211 along 53.6-m and 60.3-m straight transects with a quantum sensor (MIJ-14PAR,
212 Environmental Measurement Japan Co. Ltd., Fukuoka, Japan) mounted at one end of
213 supporting arm (see Kumagai et al., 2013). The data on PPFD below the canopy at Som
214 Sanuk and CRRI were used for estimating radiative energy absorbed within the rubber tree

215 canopy and determining the relationship between the absorbed radiative energy and the
216 amount of foliage for each site.

217 Volumetric soil water content (θ) was measured through a depth profile at each site using
218 time domain reflectometers (TDR; CS616, Campbell Scientific). TDRs were placed at
219 depths (orientations) of 0.04 m (horizontal), 0.04-0.34 m (vertical), 0.48-0.75 m (vertical),
220 0.80 m (horizontal), 1.50-1.80 m (vertical), and 2.20-2.50 m (vertical) at Som Sanuk, and at
221 0.04 m (horizontal), 0.35-0.65 m (vertical), 1.05-1.35 m (vertical), 2.03-2.33 m (vertical),
222 and 3.08-3.38 m (vertical) at CRRI. At each site, soil heat flux (G) was estimated from
223 measurements taken at a depth of 8 cm with four soil heat flux plates (HFP01, Hukseflux)
224 and the calculated change in soil heat storage in the upper 8 cm layer based on soil
225 temperature and soil moisture measurements. The data were recorded with a data logger
226 (CR23X) at 30-min intervals.

227

228 2.3. Eddy covariance measurements

229 The latent heat flux (LE_T , where L is the latent heat of vaporization of water) and sensible
230 heat flux (H) were measured using the eddy covariance technique. At each site, a
231 three-dimensional sonic anemometer (CSAT3, Campbell Scientific) and an open-path
232 CO_2/H_2O analyzer (LI-7500, Li-Cor, Lincoln, NE) were installed (at 26.5 and 30.1 m height,
233 at Som Sanuk and CRRI, respectively). The CSAT3 and LI-7500 sensor orientations were
234 switched seasonally toward the southwest in wet season, and the north in dry season (the
235 seasonal predominant wind directions). Reorienting the sensors was necessary to minimize
236 effects of turbulence from the tower structure on the eddy covariance measurements. Wind
237 speeds and gas concentration time series were all sampled and stored at 10 Hz on a

238 datalogger (CR3000). All variances and covariances required for the eddy covariance flux
239 estimates were computed over a 30-min averaging interval. The details of screening, filtering,
240 and post-processing of 10 Hz data to compute 30-min fluxes are given by Giambelluca et al.
241 (submitted for publication).

242 The ratio of mean annual $LE_T + H$ to mean annual available energy (A), which was
243 estimated as net radiation minus G and heat storages in within-canopy air and above-ground
244 biomass, was 0.94 for Som Sanuk and 0.88 for CRRI. Here, we employed a “moving
245 window energy closure ratio $(LE_T + H)/A$ ” method for closing the canopy surface energy
246 balance for each 30-min interval. LE_T data gaps were filled with values estimated from the
247 relationship between actual LE_T and A . The details of energy closure adjustment and gap
248 filling are also given by Giambelluca et al. (submitted for publication).

249 In this study, we identified and removed 30-min averaged eddy fluxes during which
250 rainfall occurred. This was done because our main objective is to investigate environmental
251 control of G_C , derived from dry canopy E_T . Importantly, the open-path analyzer was
252 mounted in an inclined orientation and the values of AGC (Automatic Gain Control) were
253 used to eliminate abnormal flux values, so as to reduce problems of rain and dew droplets
254 sitting on the window. Despite high annual precipitation at these sites, rainfall during the wet
255 season frequently occurs in short bursts, thus wet sensor interruptions are relatively minor
256 compared to sites with similar rainfall totals that are more frequently wet.

257

258 2.4. Leaf area index measurements

259 The silhouette area index of leaves plus stems and branches (plant area index; PAI) was
260 measured using a pair of plant canopy analyzers (PCA) (LAI-2000 or LAI-2200, Li-Cor) at

261 irregular time intervals over the course of the study at an interval of 6 and 5 m at Som Sanuk
262 and CRRI, respectively, along transects oriented diagonally with respect to tree rows. The
263 PAI measurements were conducted frequently during leaf shedding and flushing periods in
264 2010 and 2012, and periodically throughout other years to approximate the annual cycle of
265 foliage accumulation and loss throughout the study period. While optical PCA
266 measurements are limited, estimation of the seasonal pattern of PAI is based on the best
267 available data at the sites, and the analysis to estimate the pattern of the annual cycle in each
268 year is independent of the flux data. Because the measurements in 2011 were scarce or
269 unavailable, seasonal variation was estimated based on the measured value near the time of
270 maximum PAI in August 2011 and on the annual cycles of 2010 and 2012, and 2013, years
271 which had more frequent PCA observations. Bi-weekly litterfall collections were also used
272 to verify the patterns of leaf loss during the leaf shedding period. Additionally, the timing of
273 leaf fall and leaf area recovery were estimated based on albedo measurements and visual
274 observations.

275 Kucharik et al. (1998) and Bréda (2003) suggested that PAI detected by PCA denotes leaf
276 area index (LAI) in the period when the foliage can mask the woody area, but stems cannot
277 be overlooked in the leaf-fall period. Also, Pokorný and Marek (2000) confirmed that a
278 common use of the LAI-2000 tends to underestimate LAI because the large zenith angle
279 readings include the effect of woody areas, especially, stems. Therefore, PCA measurements
280 were adjusted by masking the ring at the lowest angle (centered at 68° zenith angle) which
281 can only see stems. Masking consecutive lower angles in mature rubber plantations caused a
282 consistent increase in calculated LAI with PCA. The studied rubber canopies at both Som
283 Sanuk and CRRI were completely dormant (i.e., LAI = 0) on the day of the lowest PAI, but

the period for such dormant days is less than one month (see Figs. 2 and 3). Therefore, we could assume that for both Som Sanuk and CRRI, the time course of LAI was equal to that of PAI, except for January, when LAI was calculated by subtracting the minimum PAI during the studied period from the yearly PAI. At both sites, January is the leaf-fall period, and thus, was not used for the examination of the environmental stomatal control (described later).

2.5. Calculation of soil water balance

Because slopes at both Som Sanuk and CRRI are gentle, lateral soil water movement can be neglected. Vertically integrated soil water balance equation is then given by

$$\int_{t_1}^{t_2} [P(t) - E_T(t)] dt - \int_0^z [\theta(z, t_2) - \theta(z, t_1)] dz = 0 \quad (1)$$

where t is time, t_1 and t_2 are temporal points in the time course, and z is soil depth. It should be noted that leakage at z is neglected in Eq. (1). The first term in the right-hand side of Eq. (1) was referred to as actual water balance, $P - E_T$. The second term denotes the difference

between soil water content at t_2 and t_1 (Δw). $\int_0^z \theta dz$ in the second term was computed as

follows: (1) estimating θ_{10} , θ_{50} , θ_{100} , θ_{200} , and θ_{300} (where each subscript denotes the soil depth (in cm) for each θ) by weighted-averaging the measured θ profiles (see Figs. 2 and 3); and (2) integrating θ along z such as $300\theta_{10} + 450\theta_{50} + 750\theta_{100} + 500\theta_{200}$ and $300\theta_{10} + 450\theta_{50} + 750\theta_{100} + 1000\theta_{200} + 500\theta_{300}$ (in mm) for the 0-2 m (at Som Sanuk) and the 0-3 m (at CRRI) soil layers, respectively.

Potential evaporation (E_{T_POT}) was an expression represented by Priestley and Taylor (1972):

$$E_{T_POT} = \alpha_{PT} \frac{\Delta A}{L(\Delta + \gamma)} \quad (2)$$

where α_{PT} is the Priestley-Taylor coefficient (= 1.26), Δ is the rate of change of saturation water vapor pressure with temperature, A is the total available energy ($R_n - G$; where R_n is net radiation), and γ is the psychometric constant. $\int_{t_1}^{t_2} [P(t) - E_{T_POT}(t)] dt$ was referred to as potential water balance, $P - E_{T_POT}$.

310

2.6. Calculation of canopy stomatal conductance

The surface conductance (G_s), representing the efficiency of exchange between the top of the vegetation surface and the atmosphere, was calculated by applying the actual data of LE_T to the inverted Penman-Monteith equation as:

$$G_s = \frac{G_a \lambda E_T}{\Delta A + \rho C_p G_a D - (\Delta + \gamma) LE_T} \quad (3)$$

where ρ is the density of dry air, C_p is the specific heat of air at a constant pressure, and D is the vapor pressure deficit above the canopy. Aerodynamic conductance, G_a , was estimated as u_*^2 / u , where u_* is the friction velocity and u is wind speed above the canopy.

Canopy conductance (G_c), representing the efficiency of the canopy transpiration (LE_c), was calculated according to Kelliher et al. (1995) with some modifications:

$$G_c = \frac{G_a \lambda E_c}{\Delta A_c + \rho C_p G_a D - (\Delta + \gamma) LE_c}, \quad (4)$$

for which

$$A_c = A - A_s \quad (5)$$

$$A_s = A \exp(-aPAI^b) \quad (6)$$

In Eqs. (5) and (6), A_c and A_s are the total available energy absorbed by leaves within the canopy and at the vegetation floor, respectively; and a and b are the extinction coefficient and fitting parameter, respectively. The parameters a and b were estimated using nonlinear least squares regression between values of A_s calculated from Eq. (6) and obtained by the radiation observations above and below the canopy. On the basis of our regression analysis, we obtained values of $a = 0.34$ and $b = 1.47$ for both Som Sanuk and CRRI. LE_c can be expressed as a contribution from the canopy to the total vegetation latent heat flux:

$$LE_c = LE_T - LE_s, \quad (7)$$

for which

$$LE_s = \alpha_s \frac{\Delta A_s}{\Delta + \gamma}, \quad (8)$$

where LE_s is latent heat flux at the vegetation floor, and α_s is an empirical parameter representing evaporative efficiency. The parameter α_s was estimated to be 0.3 and 0.5 for Som Sanuk and CRRI, respectively, under the leaf-less (i.e., $LAI = 0$) conditions, where LE_T can be assumed to be equal to LE_s . Also, we assumed that the α_s determined in the dry season might not cause large error in estimating LE_s during the wet season because of relatively small values of A_s in the leafy and wet season.

We assumed that G_c is the parallel sum of the individual leaf stomatal conductances (e.g., Raupach and Finnigan, 1988), giving

$$g_s = \frac{G_c}{LAI} \quad (9)$$

where g_s is mean canopy stomatal conductance per unit leaf area ($m s^{-1}$, canopy stomatal

conductance: see Pataki and Oren, 2003). Note that for the quality control, we removed the g_s values obtained when $D \leq 0.6$ kPa (Ewers and Oren 2000).

2.7. Stomatal conductance model

To facilitate the interpretation of the influence of environmental variables on stomatal behavior, a multiplicative-type function is commonly used with determining parameters contained in each function by appropriate nonlinear regression analysis (see Jarvis et al., 1976). Oren et al. (1999) focused on the relationship between stomatal conductance and D as follows (see Kumagai 2011):

$$g_s = g_{sref} - m \ln D \quad (10)$$

where g_{sref} is the reference value of g_s (i.e., the value of g_s at $D = 1$ kPa) and m is the sensitivity of g_s to D (i.e., $-dg_s / d\ln D$). The model of Oren et al. (1999), i.e., Eq. (10), has been developed to analyze a leaf-level stomatal (g_s) response to atmospheric evaporative demand (D) employing the mean canopy stomatal conductance derived from individual tree canopy conductance, which is estimated by the sapflow measurement, divided by total leaf area for an individual tree (see Pataki and Oren, 2003). In other words, the stomatal characteristics such as the parameters g_{sref} and m are usually estimated for individual trees. In this study, we used ecosystem-level evapotranspiration (E_T) observed by eddy covariance measurements. Separating dry-canopy evaporation, i.e., canopy transpiration (E_c), from E_T and individual tree level stomatal behavior derived from canopy level one because of the even-sized trees within each study site enabled us to compare the Oren et al. model parameters obtained in the study sites and other studies.

A boundary line analysis on the relationships between D and g_s can be used for

determining the parameters g_{sref} and m (e.g., Schäfer et al., 2000; Ewers et al., 2001, 2005). The boundary line was derived by regressing Eq. (10) with data obtained as follows: (1) selecting g_s when solar radiation, R_s , $\geq 600 \text{ W m}^{-2}$; (2) partitioning the g_s response to D into 0.5 kPa D intervals; (3) removing outliers within each interval of D ($p < 0.05$ Smirnov-Grubbs test); and (4) selecting data above the mean plus one standard deviation of g_s and averaging the selected data for each D interval. Midday data (1000-1400 LT) were used for the analyses examining the environmental control of g_s . The maximum contributions to rainfall were in the mid-night and early morning at Som Sanuk and in the mid-night at CRRI. In addition, the data were selected on the condition of high R_s , under which the open-path sensor window is likely to dry out quickly after the morning rainfall. The qualified data at Som Sanuk was 76.8 % and 79.8 % at CRRI, a high ratio considering the high annual rainfall at these sites.

380

381 **3. Results**

382 *3.1. Environmental and eddy flux data*

For both sites, there was little seasonality in the mean daily R_s (Figs. 2 and 3): a decrease in R_s caused by cloud cover in the wet season was compensated by the higher solar elevation during the wet season (see Kumagai et al., 2009b). The seasonal variation in solar elevation along with the annual moisture cycle produces cool (November to mid-February) and hot dry (March to May) seasons. The additional effect of cloud cover in the wet season results in sinusoidal variations in T_a : annual maximum and minimum T_a appear in hot and cool dry seasons, respectively, for both sites (Figs. 2 and 3). Therefore, for both sites, D also shows a seasonal and sinusoidal variation, but unlike T_a , minimum D appears in the wet season; and

391 the maximum in hot dry season (Figs. 2 and 3).

392 Wang and LinHo (2002) developed a universal way to define the onset and withdrawal
393 dates of the wet season over the entire Asian summer monsoon region. According to their
394 result using the global climatological 5-day mean rainfall dataset for the 20-year period
395 (1979-1998), the onset and withdrawal dates (DOY) were 135 and 275, respectively, at Som
396 Sanuk, and 155 and 295, respectively, at CRRI. This suggested that on the normal conditions,
397 onset and withdrawal of the wet season are prone to be 20 days earlier at Som Sanuk than at
398 CRRI, although the wet season length for the two sites are the same (ca. 140 days).

399 Here, we applied Wang and LinHo's (2002) criteria with January mean precipitation for
400 2010-2011 at Som Sanuk and 2010-2012 at CRRI for each studied year (Figs. 2 and 3). As a
401 result, at Som Sanuk, the onset and withdrawal dates (DOY) were 131 and 256, respectively,
402 in 2009, when the onset came earlier and the period of wet season became shorter than for
403 normal years. There, in 2010, the onset and withdrawal dates were 126 and 266, respectively,
404 coming earlier than for normal years, but the length of wet season was the same as a normal
405 year. In comparison, at CRRI, the onset and withdrawal dates were 121 and 316, respectively,
406 in 2010, and 101 and 305, respectively, in 2011. There, for both studied years, the onset and
407 withdrawal were much earlier and later than for normal years, inducing much longer wet
408 season (ca. 200 days).

409 For both sites, the periods of full-leaf-fall were, at most, around one month (Figs. 2 and 3).
410 This also means that rubber trees are leafy even during most parts of dry season. Annual rate
411 of P was 2118 mm, compared with E_T estimate of 1128 mm at Som Sanuk. These values at
412 CRRI were 1439 and 1272 mm, respectively (see Figs. 2 and 3; Giambelluca et al., submitted
413 for publication). After adjustment for energy closure, the annual E_T became 1221 mm for

414 Som Sanuk and 1459 mm for CRRI (Giambelluca et al., submitted for publication). Thus,
415 two major questions arise: (1) what factors sustain such high E_T of rubber plantations,
416 especially at CRRI, where the annual P and E_T were similar; and (2) what factors sustain the
417 persistence of the E_T during the dry season, especially during the late (hot) part of the dry
418 season, when the ecosystem water resources must be limited.

419

420 3.2. Tree water use and soil water balance

421 E_{T_POT} , E_T , and P for each site are compared for each season (i.e., December-February:
422 DJF, March-May: MAM, June-August: JJA, and September-November: SON) in Fig. 4.
423 Negative values of potential water balance ($P - E_{T_POT}$) indicate seasons when the ecosystem
424 experienced water resource deficits, i.e., in DJF and MAM 2010 and DJF and MAM 2011 at
425 Som Sanuk and in MAM 2010, DJF and SON 2011, and DJF 2012 at CRRI (Fig. 4). Actual
426 water balance ($P - E_T$) was almost zero, indicating that tree water use regulation might be a
427 major controlling factor in MAM of 2011 at Som Sanuk and in SON of 2011 at CRRI.

428 We found that in the season when $P - E_T$ was negative (DJF and MAM 2010 and DJF
429 2011 at Som Sanuk, and MAM 2010, DJF 2011 and 2012 at CRRI), the $P - E_T$ was
430 compensated with residual soil water stored during the previous season (Δw) in the 0-2 m
431 soil layer at Som Sanuk and 0-3 m soil layer at CRRI (Fig. 4). Here, we should note that the
432 Δw might reflect the $P - E_T$ plus leakage from the bottom of soil layer and, therefore,
433 represents the maximum limit of the actual compensation for the $P - E_T$. Such compensation
434 for the tree water use deficit might have been achieved by a combination of utilization of the
435 carry-over soil water and the tree water use regulation.

436

437 3.3. *Tree water use regulation via stomatal behavior*

438 Seasonal patterns of midday (1000-1400 LT) averaged G_s , G_c and g_s , as major indices for
439 depicting the tree water regulation, are compared at both sites in Fig. 5. In the entire studied
440 period, the G_s reached as high as 25 mm s^{-1} at Som Sanuk and 30 mm s^{-1} at CRRI in the wet
441 seasons; and the minimum value was about 2 mm s^{-1} at both sites in the dry seasons (Fig. 5a,
442 c). On the other hand, maximum and minimum values of the G_c were approximately 25 mm
443 s^{-1} at Som Sanuk and 30 mm s^{-1} at CRRI in the wet seasons; and they were almost zero at
444 both sites in the dry season when trees were leaf-less (Fig. 5a, c). The differences between G_s
445 and G_c , due to the contribution of understory and soil (floor) evapotranspiration ($E_{T_{fl}}$) to the
446 total E_T , were somewhat constant (at most 2 mm s^{-1}), and thus, the $E_{T_{fl}}$ was not thought to
447 vary appreciably throughout the studied period. It is reasonable to assume $E_{T_{fl}}$ to be
448 invariant, given that some partial cancelations emerge – for example lower radiative energy
449 reaching the understory and soil in the leafy and wet season is likely to result in the lower
450 $E_{T_{fl}}$; however, low soil moisture in the leaf-less and dry season also induces lower $E_{T_{fl}}$.
451 Although g_s was estimated as G_c divided by LAI (see Eq. (9)), the seasonal variation in the g_s
452 was similar to the variation in G_c in that the g_s gradually increased with leaf-out and
453 leaf-expansion, and decreased with leaf-fall (Fig. 5b, d, see also Figs. 2 and 3). The gradual
454 increase in g_s after leaf expansion might be due to a maturation process in the photosynthetic
455 capacity of leaves (see Wilson et al., 2001). Interestingly, the g_s decreased after August at
456 Som Sanuk and after September-October at CRRI, corresponding to an earlier withdrawal of
457 wet season at Som Sanuk (in September) than at CRRI (in November). This steep decrease in
458 the g_s also might be due to change in the leaf-scale physiological status, i.e., leaf-senescence.
459

460 3.4. Analyses of stomatal conductance

461 The representative relationship between D and g_s in MAM at both sites (Fig. 6) shows a
462 general tendency of g_s to decline with increasing D . We obtained the reference conductance,
463 g_{sref} , and the rate of decline in g_s , m , by fitting Eq. (10) to the upper boundaries of the D - g_s
464 data. The MAM analysis period encompasses the end of the dry season and start of the wet
465 season, i.e., the onset of a wet season was in mid-May at Som Sanuk (see Fig. 6a) and early
466 May at CRRI (see Fig. 6b). Thus, although there was not a significant difference in
467 hydro-climate conditions between Som Sanuk and CRRI, g_{sref} and m were significantly
468 higher at CRRI than at Som Sanuk.

469 Fig. 7 compares seasonal variations in g_{sref} and m at Som Sanuk and CRRI. As ranges of D
470 in the D - g_s plots (see Fig. 6) are critical for the estimations of g_{sref} and m (e.g., Herbst et al.,
471 2008; Schmidt-Walter et al., 2014), we further estimated g_{sref} and m with a narrower range of
472 D (upper limit: 2 kPa) (Fig. 7). In the common study year 2010, even if taking into
473 consideration the lowered values by the narrower D range estimations, both g_{sref} and m were
474 larger at CRRI than at Som Sanuk. The g_{sref} values were robust regardless of the range of D
475 values, while m values were appreciably altered, in particular, much for CRRI, by the
476 narrower range estimations. At Som Sanuk, g_{sref} reached a minimum when $P - E_T < 0$; and
477 the variation in m is similar to that for g_{sref} (Fig. 7a). On the other hand, at CRRI, minimum
478 g_{sref} did not coincide with a condition of $P - E_T < 0$; minimum g_{sref} appeared in MAM, 2011,
479 which, despite being mostly within the dry season, had a positive water balance ($P - E_T > 0$,
480 Fig. 7b). At CRRI, g_{sref} and m was lower from SON and DJF in 2011-2012 as compared to
481 the same seasons in 2010-2011 (Fig. 7b), probably because the wet season withdrawal was
482 earlier in 2011 than in 2010 and other normal year (see Fig. 4), and it may also be explained

483 by differences in leaf emerging and senescence patterns between 2010 and 2011 as the stand
 484 matured, as lower leaves remained on the trees until the end of the dry season in 2010, and a
 485 secondary flushing of lower leaves emerged in mid-2010. The CRR g_s started decreasing in
 486 mid-October in 2011, however not until early December in 2010 (see Fig. 5). Note that it was
 487 confirmed that difference in LAI estimates between SON 2010 (about 4.0) and 2011 (about
 488 4.5) did not significantly alter the decreasing trend in the g_s in 2011.

489 The 0.6 proportionality (m/g_{sref}) results from the regulation of the minimum leaf water
 490 potential (ψ_L) to prevent excessive xylem cavitation, while a value less than 0.6 denotes less
 491 strict regulation of ψ_L (Oren et al., 1999; Ewers et al., 2005, 2007). We showed seasonal
 492 variations in the m/g_{sref} ratio estimated by both the conventional and the narrower D range
 493 methods with the 0.6 proportionality at Som Sanuk and CRR in Fig. 8. Rubber trees at Som
 494 Sanuk had a proportionality significantly lower than 0.6, with only one point greater than 0.5
 495 (for all points: $m/g_{sref} = 0.16-0.51$; cf., the proportionality 0.51 appeared when $P - E_T < 0$),
 496 implying these trees allow the minimum ψ_L to drop with increasing D . While for CRR, five
 497 of eight points were close to the 0.6 proportionality line (for all points: $m/g_{sref} = 0.19-0.64$),
 498 showing somewhat strict stomatal regulation of the minimum ψ_L . The CRR
 499 proportionalities that were much less than 0.6 (Fig. 8) were apparently due to much lower
 500 g_{sref} and m from mid-2011-2012 (see Fig. 7b). Here, we have to note that the theoretical
 501 proportionality decreases from 0.68 to 0.53 as the D range broadens from 1-3 to 1-5 kPa
 502 (Oren et al., 1999), implying that the 0.6 proportionality might be valid only for a D range of
 503 1-4 kPa. Thus, we recognize that although estimating the proportionality using the narrower
 504 D range would contain some additional uncertainty in the data interpretation, the evidence
 505 suggests a tendency for rubber trees at CRR to regulate their stomatal behavior more strictly

506 than those at Som Sanuk when $P - E_T < 0$ (Fig. 8).

507

508 **4. Discussion**

509 Guardiora-Claramonte et al. (2008) surmised that the introduction of rubber plantations
510 may reduce stream flow leading to drier conditions throughout a catchment area, mainly
511 because of high water use by rubber trees at the very peak of the dry season (cf. Fox et al.,
512 2014). As rubber trees originate from the Amazonian basin where it is rainy for most of the
513 year, they are intrinsically adapted to moist environments (Priyadarshan et al., 2005). In
514 mainland of Southeast Asia, however, a dry season extends from approximately November
515 to April, during which rubber trees shed their leaves over a short period of time (Figs. 2 and
516 3). They then initiate leaf-out by the beginning of February, well before the onset of wet
517 season. Rubber trees in this region appear to remain physiologically active for producing
518 new leaves and buds during the dry season by depleting soil water from deep within the soil
519 (see Fox et al., 2014).

520 Following on previous work, our main concern is to confirm that rubber plantations act as
521 “water pumps” (see Tan et al., 2011) and to determine how they consume such a large
522 volume of water despite the long dry season in this region. Using multi-year E_T flux
523 observations, our prior work revealed that rubber E_T is higher than that of other
524 tree-dominated land covers in mainland Southeast Asia (Giambelluca et al., submitted for
525 publication). Herein, we examined how the rubber trees used the ecosystem water resources,
526 especially in the dry season, and then, how environmental and biological factors affect their
527 canopy stomatal behavior under both water stress and non-stress conditions. We used a
528 combination of the E_T flux data and an inverted version of a simple 2-layer E_T model to

529 estimate stomatal regulation of tree water use, that is, g_s . This analysis enabled us to extract
530 the rubber canopy conductance from total surface conductance, including the canopy and the
531 understory and soil effects. As a result, we demonstrated how each studied rubber plantation
532 copes with strong seasonal drought in terms of tree water use strategy.

533 Potential water balance ($P - E_{T_POT}$) for each season can represent the need for tree water
534 use control, mainly by stomatal regulations, and needs for accessing previously stored soil
535 water (Fig. 4). Rubber trees in the season when actual water balance ($P - E_T$) was negative
536 rely on carried-over soil water from the previous season. Investigation of the temporal
537 variation in soil moisture profiles revealed that such negative $P - E_T$ was compensated with
538 soil water carried over from the previous season in the 0-2 m soil layer in Som Sanuk and the
539 0-3 m soil layer in CRRI. Many studies have reported that majority of tropical trees can
540 avoid severe water stress via deep soil water utilization during the dry periods (e.g., Nepstad
541 et al., 1994; Markewitz et al., 2010; Stahl et al., 2013). On the other hand, according to a
542 global analysis of main root distributions (Jackson et al., 1996), 97 % of main roots in
543 tropical deciduous forests are, on average, in the top 1 m. In reality, empirical knowledge for
544 rubber plantation management in CRRI suggested that the rubber tree main-rooting depth
545 should reach at most down to 1 m. Notwithstanding such insufficient rooting depth, soil
546 moisture profiles and visual observations of some deep fine roots showed the studied rubber
547 trees evidently extracted water from 2 and 3 m deep in the soil. Stahl et al. (2013) pointed out
548 that the vertical distribution of root biomass does not perfectly reflect the depth of tree water
549 uptake, and thus we need further investigation about the ability of the fine root system to
550 explore the deeper soil layers.

551 At both sites, g_s started to decrease earlier than the initiation of leaf-fall and withdrawal of

the wet season (see Figs. 2, 3 and 5). Interestingly, despite almost the same timing for leaf-fall at Som Sanuk and CRRI, the g_s started to decrease about one month earlier at Som Sanuk than at CRRI. This might be because the wet season withdrawal occurs more than one month earlier at Som Sanuk than at CRRI, as inferred from our observations and those of Wang and LinHo (2002). Also, at both sites, the intrinsic capacity of g_s appeared to start to recover immediately after the initiation of leaf-out, and the value of g_s gradually increased as the wet season proceeded (see Figs. 2, 3 and 5). These trends implied that the increases and decreases in the g_s were induced by not only environmental perturbation but also temporal variation in intrinsic leaf physiological traits such as photosynthetic capacity (see Kitajima et al., 2002).

There are promising indices for examination of the environmental and biological control of stomatal behavior, i.e., the reference value of g_s (g_{sref}) and the sensitivity of g_s to D (m) (Oren et al., 1999). Among these parameters, g_{sref} is thought to represent intrinsic and maximal capacity of g_s . Compared with the reported values of g_{sref} , collected from various biomes (Oren et al., 1999), maximum g_{sref} observed in this study (estimated to be $360 \text{ mmol m}^{-2} \text{ s}^{-1}$) was in their upper range; and even the minimum value ($\approx 150 \text{ mmol m}^{-2} \text{ s}^{-1}$), which appeared in the mid dry season, was somewhat common in Oren et al.'s (1999) study. Such relatively high g_{sref} throughout a year were maintained by the availability of deep soil water, which, therefore, is thought to be a major cause for the high rubber E_T . It was found that g_{sref} reached an annual peak in JJA at Som Sanuk and in JJA to SON at CRRI. Further, it started to decrease at the withdrawal of the wet season, i.e., in SON at Som Sanuk and in DJF at CRRI (Fig. 7). Also, at both sites, g_{sref} reached its lowest point each year when $P - E_T < 0$, except for DJF 2010-11 at CRRI when g_{sref} remained high. The process of full leaf-fall and

575 full leaf-expansion was completed within the period of DJF, when $P - E_{T_POT} < 0$.

576 Considering the above-mentioned behavior, rubber trees seem to avoid risk of water stress

577 during the dry season by reducing activity, then maximizing activity when soil water is

578 sufficient (see Kumagai and Porporato, 2012). The 0.6 proportionality (m/g_{sref}) was

579 predicted from the examination of results of the plant transpiration model coupled with the

580 hydraulic constraint model in which plants control transpiration rate via stomatal regulation

581 to protect the transport system from excessive loss of hydraulic function (Oren et al., 1999;

582 Cochard et al., 2002; Ewers et al., 2005; Katul et al., 2009). Thus, it was demonstrated that

583 rubber trees maintained strict stomatal regulation to keep the difference between leaf water

584 potential (ψ_L) and soil water potential constant in the period MAM 2010-JJA 2011 at CRRI.

585 There was less sufficient stomatal regulation in the period SON 2011-DJF 2012 at CRRI and

586 MAM 2009-SON 2010 at Som Sanuk (see Herbst et al., 2008; Kumagai et al., 2008;

587 Naithani et al., 2012) (Fig. 8). Observations at CRRI showed somewhat constant midday ψ_L

588 throughout the year, even in the dry season, partly confirming the strict stomatal regulation

589 characteristics (data not shown here). Kumagai et al. (2008) hypothesized that the

590 proportionality < 0.6 for Japanese cedar would confer a carbon uptake advantage to tree

591 species suitable for moist soil conditions, as there would be little risk of water stress-induced

592 hydraulic failure and g_s would not have to decrease with increasing atmospheric demand. In

593 this study, the proportionalities of ≈ 0.6 were realized at CRRI, where the ecosystem water

594 resources were less than at Som Sanuk and the rubber trees had to extract water from the 0-3

595 m soil layer to compensate the water deficit under severe drought conditions (see Fig. 4). The

596 values less than 0.6 were produced from somewhat moist conditions, such as the whole study

597 period at Som Sanuk where there is more abundant precipitation, and the period SON

2011-DJF 2012 at CRRI, when the annual precipitation was about 200 mm higher than in 2010. These findings are consistent with the hypothesis of Kumagai et al (2008).

In terms of regulation of water status, plants can be categorized as “isohydric” or “aniso-hydric” (see McDowell et al., 2008). Isohydric plants regulate transpiration by stomatal closure to maintain constant ψ_L , thus avoiding drought-induced hydraulic failure. In contrast, aniso-hydric plants have less stomatal sensitivity and allow ψ_L to decrease with decreasing soil moisture. On intra-annual time scales, the studied rubber trees had sensitive stomatal regulation under drought conditions, i.e., they exhibited isohydric behavior. However, aniso-hydric behavior was observed in the whole period at Som Sanuk and after SON 2011 at CRRI under moist environments where there was little risk of hydraulic failure (see Kumagai and Porporato, 2012). This alternating isohydric and aniso-hydric behavior makes these trees efficient in terms of water-use, as it allows the rubber trees to take advantage in their productivity under moist conditions and avoid risk of hydraulic failure under drought conditions.

5. Summary and implications

Results of this investigation suggest that for the rubber trees at Som Sanuk, where there is larger precipitation than at CRRI, water uptake from shallower soil depth and less stomatal regulation are sufficient under moist conditions, while at CRRI, the uptake from a deeper soil depth and strict stomatal regulation are necessary to cope with drier conditions. Because the studied sites are among the regions having the strongest seasonality in precipitation (see Kumagai et al., 2005), the strategy of these rubber trees for avoiding severe water stress in the dry season exhibits an exquisite balance between tree water use and the ecosystem water

621 resource, which has drastic seasonal variation. This also means that there might be a “tipping
622 point” (see Lenton et al., 2008) in the studied ecosystems, in that even a tiny perturbation in
623 the ecosystem water resources via changes in rainfall regimes could alter their state, balance
624 and development. For example, a close relationship between the interannual variations of the
625 monsoon onset and El Niño–Southern Oscillation (ENSO) on the Indochina Peninsula was
626 identified, and El Niño events appear to be associated with late monsoon onset leading to a
627 reduction in annual precipitation (Zhang et al., 2002). Global warming is likely to cause
628 changes in Pacific regional climate that might alter ENSO activity in the future (e.g.,
629 Timmermann et al., 1999). While it is still not clear how ENSO will be affected by global
630 warming, it is possible that the frequency or amplitude of ENSO events could increase
631 (Collins et al., 2010). Therefore, there is an urgent need to further examine the current and
632 future perturbations in the hydrological, atmospheric and ecological processes responding to
633 seasonal and interannual (e.g., ENSO-related) variation in precipitation in the region. One
634 approach would be to combine an investigation of the impact of land use change (i.e., from
635 pre-existing vegetation to rubber plantation; see Fig.1; Fox et al., 2012) with this study’s
636 result of rubber plantations-atmosphere exchange characteristics.

637

638

638 **Acknowledgements.**

639 The study in Cambodia was initiated as cooperative project between Cambodian Rubber
640 Plantation Department, Cambodian Rubber Research Institute (CRRI), University of
641 Hawai‘i, and Kyushu University, supported by the Global COE (Centers of Excellence)
642 Program (GCOE) of the Japan Society for the Promotion of Science (JSPS), National
643 Aeronautics and Space Administration (NASA) grants NNG04GH59G and NNX08AL90G,
644 and a grant from Kyushu University. We acknowledge help given by the staff of CRRI and
645 other contributors, including Tsuyoshi Kajisa, Khun Kakada, Nobuya Mizoue, Makiko
646 Tateishi, and Tetsukazu Yahara. TK, NK, and YM were supported by the project
647 “Estimation and simulation of carbon stock change of tropical forest in Asia (2011-2014)”
648 funded by the Ministry of Agriculture, Forestry and Fisheries, Japan. TK was supported by
649 in part by a Grant-in-Aid for Scientific Research (# 23405028) and the granted project
650 “Program for risk information on climate change” from the Ministry of Education, Science
651 and Culture, Japan. MH was supported by the Office of Science of the U.S. Department of
652 Energy (DOE) through the Earth System Modeling program. The Pacific Northwest
653 National Laboratory (PNNL) is operated for the US DOE by Battelle Memorial Institute
654 under contract DE-AC05-76RLO1830. Rubber mapping work was funded by NASA Earth
655 System Science grants NNG04GH59G and NX08AL90G and the Geo-Informatics and
656 Space Technology Development Agency of Thailand. ADZ was supported by National
657 University of Singapore (NUS) grant R-109-000-092-133 and Asia-Pacific Network for
658 Global Change Research (APN) grant #ARCP2008-01CMY.

659

660

660 **References**

- 661 Association of Natural Rubber Producing Countries (ANRPC). Natural Rubber Trends &
662 Statistics Archive. May 2010. <http://www.anrpc.org/html/archive.aspx> (29 January 2012,
663 date last accessed).
- 664 Bréda, N.J.J., 2003. Ground-based measurements of leaf area index: a review of methods,
665 instruments and current controversies. *J. Exp. Bot.* 54, 2403-2417.
- 666 Carr, M.K.V., 2012. The water relations of rubber (*Hevea brasiliensis*): a review. *Expl.*
667 *Agric.* 48, 176-193.
- 668 Cochard, H., Coll, L., Roux, X.L., Amélio, T., 2002. Unraveling the effects of plant
669 hydraulics on stomatal closure during water stress in walnut. *Plant Physiol.* 128, 282-290.
- 670 Collins, M., An, S.-I., Cai, W., Ganachaud, A., Guilyardi, E., Jin, F.-F., Jochum, M.,
671 Lengaigne, M., Power, S., Timmermann, A., Vecchi, G., Wittenberg, A., 2010. The
672 impact of global warming on the tropical Pacific Ocean and El Niño. *Nature Geosci.* 3,
673 391-397.
- 674 Delzon, S., Sartore, M., Burlett, R., Dewar, R., Loustau, D., 2004. Hydraulic responses to
675 height growth in maritime pine trees. *Plant Cell Environ.* 27, 1077-1087.
- 676 Ewers, B.E., Gower, S.T., Bond-Lamberty, B., Wang, C.K., 2005. Effects of stand age and
677 tree species on canopy transpiration and average stomatal conductance of boreal forests.
678 *Plant Cell Environ.* 28, 660-678.
- 679 Ewers, B.E., Mackay, D.S., Samanta, S., 2007. Interannual consistency in canopy stomatal
680 conductance control of leaf water potential across seven tree species. *Tree Physiol.* 27,
681 11-24.
- 682 Ewers, B.E., Oren, R., 2000. Analyses of assumptions and errors in the calculation of

683 stomatal conductance from sap flux measurements. *Tree Physiol.* 20, 579-589.

684 Ewers, B.E., Oren, R., Phillips, N., Strögren, M., Linder, S., 2001. Mean canopy stomatal
685 conductance responses to water and nutrient availabilities in *Picea abies* and *Pinus taeda*.
686 *Tree Physiol.* 21, 841-850.

687 Fox, J.M., Castella, J.-C., Ziegler, A.D., Westley, S.B., 2014. Rubber Plantations Expand in
688 Mountainous Southeast Asia: What Are the Consequences for the Environment?
689 AsiaPacific Issues, Analysis from the East-West Center No. 114, May 2014,,East-West
690 Center, Honolulu.

691 Fox, J., Vogler, J.B., Sen, O.L., Giambelluca, T.W., Ziegler, A.D., 2012. Simulating
692 land-cover change in montane mainland Southeast Asia. *Environ. Manage.* 49, 968-979.

693 Giambelluca, T.W., Mudd, R.G., Liu, W., Ziegler, A.D., Kobayashi, N., Kumagai, T.,
694 Miyazawa, Y., Lim, T.K., Huang, M., Fox, J., Yin, S., Mak, S.V., Kasemsap, P.,
695 Evapotranspiration of rubber (*Hevea brasiliensis*) cultivated at two sites in southeast
696 Asia. *Water Resour. Res.*, submitted for publication.

697 Guardiola-Claramonte, M., Troch, P.A., Ziegler, A.D., Giambelluca, T.W., Durcik, M.,
698 Vogler, J.B., Nullet, M.A., 2010. Hydrologic effects of the expansion of rubber (*Hevea*
699 *brasiliensis*) in a tropical catchment. *Ecohydrol.* 3, 306-314.

700 Guardiola-Claramonte, M., Troch, P.A., Ziegler, A.D., Giambelluca, T.W., Vogler, J.B.,
701 Nullet, M.A., 2008. Local hydrologic effects of non-native vegetation in a tropical
702 catchment. *Ecohydrol.* 1, 13-22.

703 Herbst, M., Rosier, P.T.W., Morecroft, M.D., Gowing, D.J., 2008. Comparative
704 measurements of transpiration and canopy conductance in two mixed deciduous
705 woodlands differing in structure and species composition. *Tree Physiol.* 28, 959-970.

706 Isarangkool Na Ayutthaya, S., Do, F.C., Pannangpetch, K., Junjittakarn, J., Maeght, J.-L.,
 707 Rochetau, A., Cochard, H., 2011. Water loss regulation in mature *Heavea brasiliensis*:
 708 effects of intermittent drought in the rainy season and hydraulic regulation. *Tree Physiol.*
 709 31, 751-762.

710 Jackson, R.B., Canadell, J., Ehleringer, J.R., Mooney, H.A., Sala, O.E., Schulze, E.D., 1996.
 711 A global analysis of root distributions for terrestrial biomes. *Oecologia* 108, 389-411.

712 Jarvis, P.G., 1976. The interpretation of the variations in leaf water potential and stomatal
 713 conductance found in canopies in the field. *Philos. Trans. R. Soc. Lond. Biol.* 273,
 714 593-610.

715 Katul, G.G., Palmroth, S., Oren, R., 2009. Leaf stomatal responses to vapour pressure deficit
 716 under current and CO₂-enriched atmosphere explained by the economics of gas exchange.
 717 *Plant Cell Environ.* 32, 968-979.

718 Kelliher, F.M., Luening, R., Raupach, M.R., Schulze, E.-D., 1995. Maximum conductances
 719 for evaporation from global vegetation types. *Agric. For. Meteorol.* 73, 1-16.

720 Kitajima, K., Mulkey, S.S., Samaniego, M., Wright, J.S., 2002. Decline of photosynthetic
 721 capacity with leaf age and position in two tropical pioneer tree species. *Am. J. Bot.* 89,
 722 1925-1932.

723 Kobayashi, N., Kumagai, T., Miyazawa, Y., Matsumoto, K., Tateishi, M., Lim, T.K., Mudd,
 724 R.G., Ziegler, A.D., Giambelluca, T.W., Yin, S., 2014. Transpiration characteristics of a
 725 rubber plantation in central Cambodia. *Tree Physiol.* 34, 285-301.

726 Kumagai, T. 2011. Transpiration in forest ecosystems. In: Levia, D.F., Carlyle-Moses, D.,
 727 Tanaka, T. (Eds.), *Forest Hydrology and Biogeochemistry: Synthesis of Past Research*
 728 *and Future Directions*, Ecological Studies 216. Springer, New York, pp. 389-406.

729 Kumagai, T., Aoki, S., Otsuki, K., Utsumi, Y., 2009a. Impact of stem water storage on
 730 diurnal estimates of whole-tree transpiration and canopy conductance from sap flow
 731 measurements in Japanese cedar and Japanese cypress. *Hydrol. Process.* 23, 2335-2344.
 732 Kumagai, T., Aoki, S., Shimizu, T., Otsuki, K., 2008. Transpiration and canopy conductance
 733 at two slope positions in a Japanese cedar forest watershed. *Agric. For. Meteorol.* 148,
 734 1444-1455.
 735 Kumagai, T., Mudd, R.G., Miyazawa, Y., Liu, W., Giambelluca, T.W., Kobayashi, N., Lim,
 736 T.K., Jomura, M., Matsumoto, K., Huang, M., Chen, Q., Ziegler, A., Yin, S., 2013.
 737 Simulation of canopy CO₂/H₂O fluxes for a rubber (*Hevea brasiliensis*) plantation in
 738 central Cambodia: the effect of the regular spacing of planted trees. *Ecol. Model.* 265,
 739 124-135.
 740 Kumagai, T., Porporato, A., 2012. Strategies of a Bornean tropical rainforest water use as a
 741 function of rainfall regime: isohydric or anisohydric? *Plant Cell Environ.* 35, 61-71.
 742 Kumagai, T., Saitoh, T.M., Sato, Y., Morooka, T., Manfroi, O.J., Kuraji, K., Suzuki, M.,
 743 2004. Transpiration, canopy conductance and the decoupling coefficient of a lowland
 744 mixed dipterocarp forest in Sarawak, Borneo: dry spell effects. *J. Hydrol.* 287, 237-251.
 745 Kumagai, T., Saitoh, T.M., Sato, Y., Takahashi, H., Manfroi, O.J., Morooka, T., Kuraji, K.,
 746 Suzuki, M., Yasunari, T., Komatsu, H., 2005. Annual water balance and seasonality of
 747 evapotranspiration in a Bornean tropical rainforest. *Agric. For. Meteorol.* 128, 81-92.
 748 Kumagai, T., Yoshifuji, N., Tanaka, N., Suzuki, M., Kume, T., 2009b. Comparison of soil
 749 moisture dynamics between a tropical rainforest and a tropical seasonal forest in
 750 Southeast Asia: impact of seasonal and year-to-year variations in rainfall. *Water Resour.*
 751 *Res.* 45, W04413, doi:10.1029/2008WR007307.

752 Kucharik, C.J., Norman, J.M., Gower, S.T., 1998. Measurements of branch area and
 753 adjusting leaf area index indirect measurements. *Agric. For. Meteorol.* 91, 69-88.
 754 Lenton, T.M., Held, H., Kriegler, E., Hall, J.W., Lucht, W., Rahmstorf, S., Schellnhuber,
 755 H.J., 2008. Tipping elements in the Earth's climate system. *Proc. Natl. Acad. Sci. U. S. A.*
 756 105, 1786-1793.
 757 Li, Z., Fox, J. 2011. Mapping rubber tree growth in mainland Southeast Asia using
 758 time-series MODIS 250 m NDVI and statistical data. *Appl. Geogr.* 32, 420-432.
 759 Mann, C.C., 2009. Addicted to rubber. *Science* 325, 564-566.
 760 Markewitz, D., Devine, S., Davidson, E.A., Brando, P., Nepstad, D.C., 2010. Soil moisture
 761 depletion under simulated drought in the Amazon: impacts on deep root uptake. *New*
 762 *Phytol.* 187, 592-607.
 763 McDowell, N., Pockman, W.T., Allen, C.D., Breshears, D.D., Cobb, N., Kolb, T., Plaut, J.,
 764 Sperry, J., West, A., Williams, D.G., Yepez, E.A., 2008. Mechanisms of plant survival
 765 and mortality during drought: why do some plants survive while others succumb to
 766 drought? *New Phytol.* 178, 719-739.
 767 Naithani, K.J., Ewers, B.E., Pendall, E., 2012. Sap flux-scaled transpiration and stomatal
 768 conductance response to soil and atmospheric drought in a semi-arid sagebrush
 769 ecosystem. *J. Hydrol.* 464-465, 176-185.
 770 Nepstad, D.C., de Carvalho, C.R., Davidson, E.A., Jipp, P.H., Lefebvre, P.A., Negreiros,
 771 G.H., da Silva, E.D., Stone, T.A., Trumbore, S.E., Vieira, S., 1994. The role of deep roots
 772 in the hydrological and carbon cycles of Amazonian forests and pastures. *Nature* 372,
 773 666-669.
 774 Oren, R., Sperry, J.S., Katul, G.G., Pataki, D.E., Ewers, B.E., Phillips, N., Schäfer, K.V.R.,

1999. Survey and synthesis of intra- and interspecific variation in stomatal sensitivity to
 vapour pressure deficit. *Plant Cell Environ.* 22, 1515-1526.

Pataki, D.E., Oren, R., 2003. Species differences in stomatal control of water loss at the
 canopy scale in a mature bottomland deciduous forest. *Advances in Water Resources* 26,
 1267-1278.

Pokorný, R., Marek, M.V., 2000. Test of accuracy of LAI estimation by LAI-2000 under
 artificially changed leaf to wood area proportions. *Biol. Plant.* 43, 537-544.

Priestley, C.H.B., Taylor, R.J., 1972. On the assessment of surface heat flux and evaporation
 using large-scale parameters. *Mon. Weather Rev.* 100, 81-92.

Priyadarshan, P.M., 2003. Breeding *Hevea brasiliensis* for environmental constraints. *Adv.*
Agron. 79, 351-400.

Priyadarshan, P.M., Hoa, T.T.T., Huasun, H., Goncalves, P.d.S. 2005. Yielding potential of
 rubber (*Hevea brasiliensis*) in sub-optimal environments. In: Kang, M.S. (Ed.), *Genetic*
and Production Innovations in Field Crop Technology: New Development in Theory and
Practice. The Haworth Press, Inc., Philadelphia, pp. 221-247.

Qiu, J., 2009. Where the rubber meets the garden. *Nature* 457, 246-247.

Raupach, M., Finnigan, J., 1988. 'Single-layer models of evaporation from plant canopies are
 incorrect but useful, whereas multilayer models are correct but useless': *Discuss. Aust. J.*
Plant Physiol. 15, 705-716.

Sangsing, K., Kasemsap, P., Thanisawanyangkura, S., Sangkhasila, K., Gohet, E., Thaler, P.,
 Cochard, H., 2004. Xylem embolism and stomatal regulation in two rubber clones
 (*Hevea brasiliensis* Muell. Arg.). *Trees* 18, 109-114.

Schäfer, K.V.R., Oren, R., Tenhunen, J.D., 2000. The effect of tree height on crown level

798 stomatal conductance. *Plant Cell Environ.* 23, 365-375.

799 Schmidt-Walter, P., Richter, F., Herbst, M., Schuldt, B., Lamersdorf, N. P. (2014)

800 Transpiration and water use strategies of a young and a full-grown short rotation coppice

801 differing in canopy cover and leaf area. *Agric. For. Meteorol.* 195, 165-178.

802 Stahl, C., Herault, B., Rossi, V., Burban, B., Brechet, C., Bonal, D., 2013. Depth of soil

803 water uptake by tropical rainforest trees during dry periods: does tree dimension matter?

804 *Oecologia* 173, 1191-1201.

805 Tan, Z.-H., Zhang, Y.-P., Song, Q.-H., Liu, W.-J., Deng, X.-B., Tang, J.-W., Deng, Y., Zhou,

806 W.-J., Yang, L.-Y., Yu, G.-R., Sun, X.-M., Liang, N.-S., 2011. Rubber plantations act as

807 water pumps in tropical China. *Geophys. Res. Lett.* 38, L24406

808 doi:10.1029/2011GL050006.

809 Tateishi, M., Kumagai, T., Suyama, Y., Hiura, T., 2010. Differences in transpiration

810 characteristics of Japanese beech trees, *Fagus crenata*, in Japan. *Tree Physiol.* 30,

811 748-760.

812 Timmermann, A., Oberhuber, J., Bacher, A., Esch, M., Latif, M., Roeckner, E., 1999.

813 Increased El Niño frequency in a climate model forced by future greenhouse warming.

814 *Nature* 398, 694-697.

815 Wang, B., LinHo, 2002. Rainy season of the Asian-Pacific summer monsoon. *J Climate* 15,

816 386-398.

817 Wauters, J.B., Coudert, S., Grallien, E., Jonard, M., Ponette, Q., 2008. Carbon stock in

818 rubber tree plantations in Western Ghana and Mato Grosso (Brazil). *For. Ecol. Manage.*

819 255, 2347-2361.

820 Wilson, K.B., Baldocchi, D.D., Hanson, P.J., 2001. Leaf age affects the seasonal pattern of

821 photosynthetic capacity and net ecosystem exchange of carbon in a deciduous forest.
822 *Plant Cell Environ.* 24, 571-583.

823 Wu, Z.-L., Liu, H.-M., Liu, L.-Y., 2001. Rubber cultivation and sustainable development in
824 Xishuangbanna, China. *Int. J. Sustain. Dev. World Ecol.* 8, 337-345.

825 Wullschleger, S.D., Wilson, K.B., Hanson, P.J., 2000. Environmental control of whole-plant
826 transpiration, canopy conductance and estimates of the decoupling coefficient for large
827 red maple trees. *Agric. For. Meteorol.* 104, 157-168.

828 Zhang, Y., Li, T., Wang, B., Wu, G., 2002. Onset of the summer monsoon over the
829 Indochina Peninsula: Climatology and interannual variations. *J. Climate* 15, 3206-3221.

830 Ziegler, A.D., Bruun, T.B., Guardiola-Claramonte, M., Giambelluca, T.W., Lawrence, D.,
831 Nguyen, T.L., 2009. Environmental consequences of the demise in swidden agriculture
832 in SE Asia: hydrology and geomorphology. *Hum. Ecol.* 37, 361-373.

833 Ziegler, A.D., Fox, J.M., Xu, J., 2009. The rubber Juggernaut. *Science* 324, 1024-1025.

834 Ziegler, A.D., Phelps, J., Yuen, J.Q., Webb, E.L., Lawrence, D., Fox, J.M., Bruun, T.B.,
835 Leisz, S.J., Ryan, C.M., Dressler, W., Mertz, O., Pascual, U., Padoch, C., Koh, L.P., 2012.
836 Carbon outcomes of major land-cover transitions in SE Asia: great uncertainties and
837 REDD+ policy implications. *Glob. Change Biol.* 18, 3087-3099.

838

839

839 **Figure Captions**

840 **Fig. 1.** Locations of rubber plantations in mainland Southeast Asia (red dots) shown along
841 with the study sites: Som Sanuk in northeastern Thailand and Cambodian Rubber Research
842 Institute (CRRI) in central Cambodia. Map is based on Li and Fox (2011).

843

844 **Fig. 2.** Climatic data at Som Sanuk in Thailand: (from the top): daily precipitation (P ; bars);
845 volumetric soil moisture content at each depth (θ ; lines); daily maximum (thick line) and
846 minimum air temperature (T_a); mean daytime vapor pressure deficit (D); daily total solar
847 radiation (R_s); plant area index (PAI); and daily total evapotranspiration from a whole
848 ecosystem (E_T). Shaded columns represent periods of wet season.

849

850 **Fig. 3.** Climatic data at Cambodian Rubber Research Institute (CRRI) in Cambodia: (from
851 the top): daily precipitation (P ; bars); volumetric soil moisture content at each depth (θ ;
852 lines); daily maximum (thick line) and minimum air temperature (T_a); mean daytime vapor
853 pressure deficit (D); daily total solar radiation (R_s); plant area index (PAI); and daily total
854 evapotranspiration from a whole ecosystem (E_T). Shaded columns represent periods of wet
855 season.

856

857 **Fig. 4.** Upper panel: precipitation (P ; asterisks), whole ecosystem evapotranspiration (E_T ;
858 closed circles), and potential evaporation (E_{T_POT} ; open circles) for each season (i.e.,
859 December-February: DJF; March-May: MAM; June-August: JJA; and
860 September-November: SON) at Som Sanuk and Cambodian Rubber Research Institute
861 (CRRI). Lower panel: actual water balance ($P - E_T$) and residual soil water stored during the

previous season (Δw : difference between mean soil water at a given season and the previous season) in 0-1, 0-2 and 0-3 m soil layers in the season of $P - E_T < 0$.

Fig. 5. Panels a and c: seasonal patterns of midday (1000-1400 LT) averaged surface conductance (G_s ; closed circles) and canopy conductance (G_c ; open circles) at Som Sanuk, Thailand and the Cambodian Rubber Research Institute (CRRI), Cambodia. Panels b and d: canopy stomatal conductance (g_s) at Som Sanuk (b) and CRRI (d).

Fig. 6. Representative relationships between vapor pressure deficit (D) and canopy stomatal conductance (g_s ; pluses) on a 30-min basis (1000-1400 LT) in March-May (MAM), 2009 at (a) Som Sanuk and in MAM 2010 at (b) Cambodian Rubber Research Institute (CRRI). Circles represent the data selected by the boundary line analysis (see text; section 2.7), to which Eq. (10) was fit (lines).

Fig. 7. Seasonal variations in canopy stomatal conductance (g_s) at vapor pressure deficit (D) = 1 kPa (g_{sref} ; closed circles) and sensitivity of g_s to D (m ($= -dg_s/d\ln D$); (open circles) at (a) Som Sanuk and (b) Cambodian Rubber Research Institute (CRRI). Values of g_{sref} (crosses) and m (pluses) estimated with a narrower range of D (upper limit: 2 kPa) are also shown. December-February: DJF, March-May: MAM, June-August: JJA, and September-November: SON. Note that the narrower D range estimations for g_{sref} and m were not made in JJA, 2009 at Som Sanuk and in SON, 2010, SON, 2011 and DJF, 2011-2012 at CRRI because of their original narrow range of D . Shaded columns represent periods when actual water balance ($P - E_T$) was negative (see Fig. 4).

885

886 **Fig. 8.** Seasonal variations in ratio of m (sensitivity of canopy stomatal conductance (g_s) to
887 vapor pressure deficit (D); $-dg_s/d\ln D$) to g_{sref} (g_s at $D = 1$ kPa) (m / g_{sref} : closed circles) at (a)
888 Som Sanuk and (b) Cambodian Rubber Research Institute (CRRI). Values of m / g_{sref}
889 estimated with a narrower range of D (upper limit: 2 kPa) are also shown (open circles). Note
890 that the narrower D range estimations for g_{sref} and m were not made in JJA, 2009 at Som
891 Sanuk and in SON, 2010, SON, 2011 and DJF, 2011-2012 at CRRI because of their original
892 narrow range of D . The theoretical ratio 0.60 (Oren et al., 1999) is shown by a solid line for
893 each site.

894

894 **Figures**

895

896

897

898 Figure 1

899

900

900

Som Sanuk

901

902

903 Figure 2

904

905

905

CRRI

906

907

908 Figure 3

909

910

910

911

912

913 Figure 4

914

915

915

916

917

918 Figure 5

919

920

920

921

922

923 Figure 6

924

925

925

926

927

928 Figure 7

929

930

930

931

932

933 Figure 8

934

935