Calnexin is a novel sero-diagnostic marker for lung cancer

Makoto Kobayashi^{1, 2}, Ryo Nagashio^{1, 3}, Shi-Xu Jiang⁴, Keita Saito¹, Benio Tsuchiya¹, Shinichiro Ryuge⁵, Ken Katono⁵, Hiroyasu Nakashima⁶, Eriko Fukuda⁷, Naoki Goshima⁷, Yukitoshi Satoh⁶, Noriyuki Masuda⁵, Makoto Saegusa⁴, Yuichi Sato^{*1, 3}

¹Department of Applied Tumor Pathology, Graduate School of Medical Sciences, Kitasato University, Kanagawa, Japan

²Research Fellow of the Japan Society for the Promotion of Science, Japan

³Department of Molecular Diagnostics, School of Allied Health Sciences, Kitasato University, Kanagawa, Japan

Departments of ⁴Pathology, ⁵Respiratory Medicine and ⁶Thoracic and Cardiovascular Surgery, School of Medicine, Kitasato University, Kanagawa, Japan

⁷Molecular Profiling Research Center for Drug Discovery, National Institute of Advanced Industrial Science and Technology, Tokyo, Japan

*Corresponding author: Yuichi Sato, Department of Molecular Diagnosis, School of Allied Health Sciences, Kitasato University, 1-15-1 Kitasato, Minami-Ku, Sagamihara, Kanagawa 252-0373, Japan. Tel.: +81 42 778 8013; Fax: +81 42 778 9854; E-mail: yuichi@med.kitasato-u.ac.jp

Keywords: Lung cancer, Calnexin, Sero-diagnostic marker, Antibody-based proteomics

Abstract

To develop sero-diagnostic markers for lung cancer, we generated monoclonal antibodies using lung adenocarcinoma (AC)-derived A549 cells as antigens by employing the random immunization method. Hybridoma supernatants were immunohistochemically screened for antibodies with AMeX-fixed and paraffin-embedded A549 cell preparations. Positive clones were monocloned twice through limiting dilutions. From the obtained monoclonal antibodies, one designated as KU-Lad-001 was recognized as calnexin (CANX) based on immunoprecipitation and MADLI TOF/TOF-MS analysis. To evaluate the utility of this antibody as a sero-diagnostic marker for lung cancer, we performed reverse-phase protein array analysis with samples of 195 lung cancer patients and 100 healthy controls. The CANX expression levels were significantly higher in lung cancer patients than in healthy controls ($P < 0.0001$), and the area under the curve of ROC was 0.980, with 96.9% specificity and 99.0% sensitivity. Furthermore, since CANX was also detected in stage I disease, the serum CANX levels should be applicable markers discriminating lung cancer patients from healthy controls and possibly used in the detection of early lung cancer. To our knowledge, the present results provide evidence that CANX may be a novel sero-diagnostic marker for lung cancer.

1. Introduction

According to the National Institute of Health, a biomarker is defined as “a characteristic used to measure and objectively evaluate normal biological processes, pathogenic processes, or pharmacological responses to therapeutic intervention”. Therefore, biomarkers provide a tremendous opportunity to promote personalized medicine, the early detection of disease, and generate diagnostic and predictive information [1]. Hanash et al. reported that to define biomarker signatures from the earliest stages of tumor development to advanced stages and metastasis, we need to perform integrative omics studies which include the genome, transcriptome, proteome, and metabolome [2]. Especially, blood-based biomarkers show potential in cancer screening and their utility could aid progression from general population risk assessment to treatment-response evaluation and recurrence monitoring.

To obtain useful sero-diagnostic markers, we place emphasis on an antibody-based proteomics approach through the random immunization method. We previously reported the application of monoclonal antibodies (MoAbs) obtained from random immunization as a useful method to obtain tissue and/ or sero-diagnostic markers for lung cancer [3, 4]. In this report, we focused on anti-calnexin (CANX) MoAb obtained from the random immunization method, which was found to be a useful sero-diagnostic

marker for lung cancer, and significantly higher CANX levels were noted in sera from patients with lung cancer than in those of healthy controls by reverse-phase protein array (RPPA).

2. Materials and Methods

2.1 Cell line and sera

A549 cells derived from lung adenocarcinoma (AC) were stored at -80 °C for proteomics analysis or fixed in AMeX and embedded in paraffin for immunohistochemistry (IHC) [5]. Sera from 195 patients with lung cancer and 100 healthy controls were used in this study (Table 1). All samples were collected in accordance with the ethical guidelines and written consent mandated, and this study was approved by the Ethics Committee of Kitasato University School of Medicine (B07-06).

2.2 Generation of monoclonal antibodies and antigen identification of Ku-Lad-001 antibody

The generation of monoclonal antibodies was previously described [3] [4]. The antigen of one antibody designated as Ku-Lad-001 was identified by immunoprecipitation (IP) and MALDI-TOF/TOF MS analysis and characterized by immunoblotting (IB) and immunohistochemistry (IHC) [6]. Recombinant CANX protein and Venus protein as negative controls with FLAG-tag were prepared using a wheat germ cell-free system [7].

2.3 Production of ascites by hybridoma producing Ku-Lad-001 antibody

Five-hundred μL of pristane was injected into mice intraperitoneally 1 week prior to inoculation with hybridomas. Hybridomas were harvested and washed with PBS-, centrifuged for 10 min at $200 \times g$, and then the supernatant was discarded. Hybridomas were resuspended in 10 mL of PBS-, and cell numbers were counted. After adjusting the cell concentration to 2.5×10^6 cells/0.5 mL with PBS-, hybridomas were intraperitoneally injected into the mice. After a couple of weeks, we harvested ascites including the highly-concentrated Ku-Lad-001 antibody. After centrifugation of the ascites, the supernatants were collected and stored at -80°C .

2.4 Reverse-phase protein array analysis

RPPA analysis was performed in almost the same way as described by Matsubara et al. but with some modifications [8]. Serum samples were diluted 1:100 with 0.01 % TrironX-100/ PBS- and spotted onto ProteoChip glass slides (Proteogen, Seoul, Korea) in duplicate in a 640-spot/slide format using Glass Slide Microarrayer (V&P Scientific Inc., San Diego, CA, USA). The spotted slides were immobilized in a moist chamber for 4 h at 37°C . After being blocked with 0.5 % Casein sodium (Wako Pure Chemical

Industries, Osaka, Japan) for 1 h at room temperature (RT), the slides were incubated with 500-times diluted Ku-Lad-001 ascites for 1 h at RT. Then, the slides were incubated with 100-times diluted biotinylated anti-mouse IgG (BA-2000, Vector Laboratories, Burlingame, CA, USA) for 1 h at RT and 1,000-times diluted streptavidin-horseradish peroxidase conjugate (GE Healthcare Bio-Sciences, Pittsburgh, PA, USA) for 30 min at RT, respectively. The peroxidase activity was detected using the 100-times diluted Tyramide Signal Amplification Cyanine 5 System (PerkinElmer Life Sciences, Massachusetts, MA, USA) for 20 min at RT. The slides were counterstained with 2,000-times diluted Alexa Fluor 546-labeled goat anti-human IgG (Life Technologies, Carlsbad, CA, USA) for 5 min at RT. Finally, the stained slides were scanned on a microarray scanner (Genepix 4000b, Molecular Devices, Sunnyvale, CA, USA). The fluorescence intensity, defined as the median net value of duplicate samples, was determined using the Genepix pro 6.0 software package (Molecular Devices). RPPA analysis was performed in triplicate on different days.

2.5 Statistical analysis

Serum CANX levels in patients with lung cancer and healthy controls were analyzed using the Mann-Whitney U-test. The area under the curve (AUC) and best cut-off point

were calculated employing receiver-operating characteristics (ROC) analysis with the StatFlex statistical software package (version 5.0, Artiteck, Osaka, Japan).

3. Results

3.1 Production of KU-Lad-001 MoAb and antigen identification

The antibody titer was tested by IHC with 100-times diluted sera of immunized mice as the first antibody on AMeX-fixed A549 cells. As a result, the sera from immunized mice contained antibodies that reacted with various components of A549 cells (Fig. 1A). Using AMeX-fixed A549 cell preparations for the IHC screening of hybridomas, we finally established 188 MoAbs including KU-Lad-001 MoAb. In order to identify the KU-Lu-6 antigen protein, we performed IP with lysate from A549 cells (Fig. 1B). The antigenic protein was observed at approximately 90 kDa. After analysis employing MALDI-TOF/TOF MS and a MASCOT search, the protein was determined as CANX (accession: P27824), which is composed of 592 amino acids with an observed M.W. of 90 Da (Fig. 1B). The immunoglobulin isotype of KU-Lad-001 antibody was determined as IgG1, κ . The result was confirmed by IB analysis with recombinant CANX protein using KU-Lad-001 ascites, which reacted with CANX protein but not Venus protein (Fig. 1C-a). The FLAG-tag reaction with recombinant CANX protein is shown in Fig. 1C-b.

3.2 Significantly elevated serum CANX levels in patients with lung cancer

The serum levels of CANX were significantly higher in lung cancer patients than in healthy controls ($P < 0.0001$, Figure 1D). Relative values of serum CANX levels ranged

from 0.00 to 4.31 (median: 3.74) in lung cancer patients, but from 0.00 to 3.29 (median: 2.13) in healthy controls (Table 1). The AUC between lung cancers and healthy controls was 0.980 (Fig. 1E). When an optimal cut-off value of 2.49 for CANX was applied, the diagnostic sensitivity and specificity for lung cancer were 99.0 and 96.9%, respectively, and the negative and positive predictive values were 96.4 and 90.5%, respectively. Clinicopathological characteristics of the patients are shown in Table 1. Serum CANX levels were not correlated with any clinicopathological factors. RPPA analysis was performed in triplicate on different days, and similar results were obtained from each experiment.

4. Discussion

CANX is a type I integral membrane phosphoprotein residing in the endoplasmic reticulum. Its intraluminal domain has been deduced to function as a lectin chaperone coordinating the timing of folding of newly synthesized N-linked glycoproteins of the secretory pathway [9]. There have been many reports that CANX plays important roles in protein synthesis and folding through the Calreticulin/ CANX cycle [10] [11] [12]. However to our knowledge, there have been no reports that CANX was secreted in serum and a useful sero-diagnostic marker. Okayama et al. reported that CANX was associated with a poor prognosis in lung AC patients, and they provided evidence that CANX played an important role in tumor invasion and metastasis by CANX SiRNA transfection of A549 cells, but they did not report detailed clinical parameters or the expression of CANX in lung AC tissues [13]. The present study revealed that serum CANX levels in lung cancer patients were significantly higher than those in healthy controls ($P < 0.001$), with 99.0% sensitivity and 96.9% specificity. Furthermore, CANX could be also used as a biomarker to differentiate lung cancer patients from healthy controls, with an AUC of 0.98 (Figure 1E). It is interesting that CANX levels were not associated with histological types of tumors or clinicopathological factors, including invasion and metastasis of lung cancer in the present study. However, CANX is

detected in sera from patients with stage I lung cancer (Table 1), indicating that CANX protein may be a useful early sero-diagnostic marker for lung cancer.

5. Conclusion

Using random immunization methods, we generated 188 MoAbs that specifically reacted with A549 cancer cells, including one designated as KU-Lad-001 which recognized CANX. The serum levels of CANX were significantly higher in lung cancer patients than in healthy controls. Furthermore, CANX was detected in the sera of patients with stage I disease. Our results suggest that serum CANX is a useful early sero-diagnostic marker for lung cancer.

7. Acknowledgements

This study was supported in part by JSPS KAKENHI Grant Number 23590414, the JST-SENTAN (Development of Systems and Technology for Advanced Measurement and Analysis: Life Innovation Area) program from the Japan Science and Technology Agency, the Japan Society for the Promotion of Science fellows (25-5519), the 2013-2014 Project Study from the Graduate School of Medical Sciences, and the Research Project (No. 2013-1004) from the School of Allied Health Sciences, Kitasato University.

8. References

- [1] Solier C, Langen H. Antibody-based proteomics and biomarker research - current status and limitations. *Proteomics* 2014; 14: 774-783.
- [2] Hanash SM, Taguchi A. Mouse to human blood-based cancer biomarker discovery strategies. *Cold Spring Harb Protoc* 2014; 2014: 144-149.
- [3] Nagashio R, Sato Y, Matsumoto T, Kageyama T, Satoh Y, Shinichiro R, et al. Expression of RACK1 is a novel biomarker in pulmonary adenocarcinomas. *Lung Cancer* 2010; 69: 54-59.
- [4] Kobayashi M, Matsumoto T, Ryuge S, Yanagita K, Nagashio R, Kawakami Y, et al. CAXII Is a sero-diagnostic marker for lung cancer. *PLOS ONE* 2012; 7: e33952.
- [5] Sato Y, Mukai K, Watanabe S, Goto M, Shimosato Y. The AMeX method. A simplified technique of tissue processing and paraffin embedding with improved preservation of antigens for immunostaining. *Am J Pathol* 1986; 125: 431-435.
- [6] Kobayashi M, Nagashio R, Ryuge S, Murakami Y, Yanagita K, et al. Acquisition of useful sero-diagnostic autoantibodies using the same patients' sera and tumor tissues. *Biomed Res* 2014; 35: 133-143.
- [7] Goshima N, Kawamura Y, Fukumoto A, Miura A, Honma R, Satoh R, et al. Human protein factory for converting the transcriptome into an in vitro-expressed proteome. *Nature Methods* 2008; 5: 1011-1017.
- [8] Matsubara J, Ono M, Honda K, Negishi A, Ueno H, Okusaka T, et al. Survival prediction for pancreatic cancer patients receiving gemcitabine treatment. *Mol Cell Proteomics* 2010; 9: 695-704.
- [9] Cameron PH, Chevet E, Pluquet O, Thomas DY, Bergeron JJ. Calnexin phosphorylation attenuates the release of partially misfolded alpha1-antitrypsin to the secretory pathway. *J Biol Chem* 2009; 284: 34570-34579.
- [10] Ellgaard L, Helenius A. Quality control in the endoplasmic reticulum. *Nat Rev Mol*

Cell Biol 2003; 4: 181-191.

[11] Deprez P, Gautschi M, Helenius A. More than one glycan is needed for ER glucosidase II to allow entry of glycoproteins into the calnexin/calreticulin cycle. Mol Cell 2005; 19: 183-195.

[12] Johnson S, Michalak M, Opas M, Eggleton P. The ins and outs of calreticulin: from the ER lumen to the extracellular space. Trends Cell Biol 2001; 11: 122-129.

[13] Okayama A, Miyagi Y, Oshita F, Nishi M, Nakamura Y, Nagashima Y, et al. Proteomic Analysis of Proteins Related to Prognosis of Lung Adenocarcinoma. J Proteome Res 2014; 13: 4686-4694.

9. Figure legends

Figure 1. Production of anti-CANX monoclonal antibody and its antigenic identification, and serum levels in patients with lung cancer and healthy controls.

(A) The antibody titer was tested immunohistochemically using 100-times diluted sera of immunized mice as the first antibody on AMeX-fixed A549 cells used as an immunogen. The sera of immunized mice contained antibodies that reacted with various cell components, such as the nucleus (↑), plasma membrane (↑↑), and cytoplasm (▲). (B) Immunoprecipitation with KU-Lad-001 antibody. Immunoblot analysis using KU-Lad-001 hybridoma supernatant as the first antibody, lane 1: A549 lysate; lane 2: A549 lysate combined with protein G; lane 3: KU-Lad-001 antibody combined with protein G; lane 4: A549 lysate combined with KU-Lad-001 antibody. Lanes 2 to 3 are negative controls, and immunoprecipitated product with KU-Lad-001 antibody was detected in lane 4. (C) Confirmation of identified antigenic protein. KU-Lad-001 antibody reacted with recombinant CANX protein, but not with recombinant Venus protein. The recombinant CANX protein also reacted with anti-FLAG-tag antibody. (D) The median CANX level in the sera from healthy controls was 2.13, but it was 3.14 in sera from lung cancer patients ($P < 0.0001$). (E) Receiver-operating characteristic curve analysis of CANX as a serum marker for lung cancer. The

corresponding area under the curve was 0.980 for CANX, with a 96.9% specificity and 99.0% sensitivity for lung cancer, while a cut-off value of 2.49 was employed.

Table 1. Serum calnexin levels in patients with lung cancer and healthy controls

Characteristics	AC_mean	AC (n)	SCC_mean	SCC (n)	SCLC_mean	SCLC (n)	LCNEC_mean	LCNEC (n)	Healthy_mean	Healthy (n)
Age										
<70	3.68	81	3.74	17	3.89	14	3.98	6	1.09	2
≥70	3.51	50	3.8	14	3.59	8	3.8	5	1.7	98
Gender										
Male	3.62	63	3.78	31	3.78	19	3.97	9	1.42	25
Female	3.61	68	0	0	3.79	3	3.56	2	1.78	75
Stage										
I	3.68	78	3.72	10	3.85	1	3.93	3		
II	3.78	12	3.79	8	3.58	1	3.73	3		
III	3.39	25	3.83	13	3.71	7	4.1	4		
IV	3.54	16	0	0	3.83	13	3.51	1		
Tumor differentiation										
Well	3.72	38	3.66	6						
Moderate	3.61	39	3.87	11						
Poor	3.45	28	3.7	7						
Unknown	3.67	26	3.81	7						
Tumor size										
<3 cm	3.63	66	3.74	9	3.52	4	3.87	4		
≥3 cm	3.63	51	3.82	16	3.9	9	3.98	6		
Unknown	3.85	14	3.72	6	3.78	9	3.52	1		

T-factor									
T1	3.62	68	3.74	9	3.59	4	3.99	3	
T2	3.62	36	3.7	10	3.76	3	3.86	6	
T3	3.73	16	3.88	8	3.82	4	3.52	1	
T4	3.45	11	3.87	4	3.84	11	4.26	1	
Nodal status									
N1-4	3.43	41	3.78	17	3.78	21	3.87	6	
N0	3.7	90	3.78	14	3.85	1	3.93	5	
Distant metastasis									
M1	3.54	16	0	0	3.81	14	3.94	10	
M0	3.63	115	3.78	31	3.73	8	3.52	1	

Figure 1