1
2
3
4 A complex origin for the Kelso Dunes, Mojave National
5 Preserve, California, USA: A case study using a simple
6 geochemical method with global applications
7
8
9

10 Daniel R. Muhs

11 *U.S. Geological Survey, MS 980, Box 25046, Federal Center, Denver, Colorado 80225*
12 *USA*

13 Nicholas Lancaster

14 *Desert Research Institute, Division of Earth & Ecosystem Sciences, 2215 Raggio*
15 *Parkway, Reno, NV 89512 USA*

16 Gary L. Skipp

17 *U.S. Geological Survey, MS 980, Box 25046, Federal Center, Denver, Colorado 80225*
18 *USA*

19
20
21
22
23 KEY WORDS: Kelso Dunes, Mojave Desert, mineralogy, geochemistry, provenance
24
25
26
27

28 **Abstract**

29 The Kelso dune field in southern California is intriguing because although it is of limited
30 areal extent ($\sim 100 \text{ km}^2$), it has a wide variety of dune forms and contains many active
31 dunes ($\sim 40 \text{ km}^2$), which is unusual in the Mojave Desert. Studies over the past eight
32 decades have concluded that the dunes are derived primarily from a single source,
33 Mojave River alluvium, under a dominant, westerly-to-northwesterly wind regime. The
34 majority of these studies did not, however, present data to support the Mojave River as
35 the only source. We conducted mineralogical and geochemical studies of most of the 14
36 geomorphically defined dune groups of the Kelso dune field as well as potential sand
37 sources, alluvial sediments from the surrounding mountain ranges. Results indicate that
38 sands in the nine western dune groups have K/Rb and K/Ba (primarily from K-feldspar)
39 compositions that are indistinguishable from Mojave River alluvium
40 (westerly/northwesterly winds) and Budweiser Wash alluvium (southwesterly winds),
41 permitting an interpretation of two sources. In contrast, sands from the five eastern dune
42 groups have K/Rb and K/Ba values that indicate significant inputs from alluvial fan
43 deposits of the Providence Mountains. This requires either rare winds from the east or
44 southeast or, more likely, aeolian reworking of distal Providence Mountain fan sediments
45 by winds from the west, at a rate greater than input from the Mojave River or other
46 western sources. The results indicate that even a small dune field can have a complex
47 origin, either from seasonally varying winds or complex alluvial-fan-dune interaction.
48 Application of K/Rb and K/Ba in K-feldspar as a provenance indicator could be used in
49 many of the world's ergs or sand seas, where dune origins are still not well understood or
50 are controversial. Four examples are given from Africa and the Middle East where such
51 an approach could yield useful new information about dune sand provenance.

1. Introduction

Dune fields occupy large areas of the world's arid and semiarid regions, mostly (though not entirely) in the subtropical deserts and in rain-shadowed mid-latitude zones (Wilson, 1973; Cooke et al., 1993; Lancaster 1995; Livingstone and Warren, 1996; Goudie, 2002; Pye and Tsoar, 2009; Warren, 2013; Lorenz and Zimbelman, 2014). Although there have been many studies of desert dune fields, much of the focus has been on genesis of dune forms, sedimentary structures found in aeolian sand, and chronology of dune deposition as an indicator of paleoclimate. In general, there has been much less work done on understanding dune sediment provenance, with many studies simply assuming an underlying or nearby rock or sediment as the primary source or ignoring the issue of provenance altogether. Nevertheless, dune field evolution cannot really be fully understood without identifying the source sediment or sediments. Further, 'aeolian system sediment state' (as defined by Kocurek and Lancaster, 1999) cannot be assessed without provenance information. Aeolian system sediment state includes evaluation of whether a dune field is transport-limited, sediment-supply-limited or sediment-availability-limited. The latter two conditions can only be assessed if the source of sand supply is known.

It is fair to ask why an understanding of dune sand provenance is important. Assessment of whether a dune field could grow larger in the future requires understanding whether there are supplies of sand available for growth. Dune field growth in turn has important implications for ecosystems that are hosted by the dune field itself, if it is stabilized, or for impacts on ecosystems that are downwind of a dune field, if it is active now and could expand in that direction with enhanced supplies of sediment.

75 Paleoclimatic interpretations of dune fields are dependent on assessment of sand
76 provenance. Very commonly, geomorphologists interpret evidence of past activity of a
77 dune field as an indicator of an arid paleoclimate and evidence of past stability of a dune
78 field as an indicator of a more humid paleoclimate. This need not be the case, however,
79 if past periods of activity are in fact linked to enhanced sediment supply and past periods
80 of stability are linked to diminished sediment supply. Aeolian sediments are unusual
81 among geologic records of climate change in that they are direct indicators of
82 atmospheric circulation. Aeolian sediments that lack geomorphic indicators of paleowind
83 can still be used for inferring paleoclimate if sand provenance can be determined.
84 Further, if a stratigraphic section composed of alternating beds of aeolian sand exists,
85 changes in paleowind direction can be determined if provenance of each aeolian sand be
86 can be ascertained.

87 The Kelso Dunes are situated in the Mojave Desert region of southern California, one
88 of a number of modest-sized dune fields in the southwestern USA and northwestern
89 Mexico (Fig. 1). Unlike many of the other Mojave Desert dunes, however, parts of the
90 Kelso Dunes are unvegetated and active. Because of their unusual nature, both from a
91 geomorphological viewpoint and as the home for a number of desert plants and animals,
92 the Kelso Dunes and surrounding areas are now protected as part of Mojave National
93 Preserve (Fig. 2). The dunes host a number of rare and sensitive desert plants (Thorne et
94 al., 1981; Pavlik, 1989; André, 2014), as well seven species of endemic insects and a
95 lizard that, while not strictly endemic, is rare outside of Mojave National Preserve
96 (Schoenherr, 1992, p. 470).

97 Speculation about the origin of the Kelso Dunes is found in a surprisingly large
98 number of studies that span more than eight decades. In his pioneering exploratory study
99 of the Mojave Desert, Thompson (1929) thought that the Kelso Dunes probably
100 originated from sediments of the Mojave River (Fig. 3), although he presented no direct
101 evidence to support this hypothesis. More than three decades passed before there were
102 any detailed studies of the dunes, but in a now-classic paper Sharp (1966) provided
103 mineralogical and particle size data for Kelso Dunes and proposed that, in agreement
104 with Thompson (1929), the Mojave River was likely the main source for the dunes. In a
105 follow-up study some years later, Sharp (1978) reiterated the importance of the Mojave
106 River as a source. Nevertheless, in his earlier study Sharp (1966) also noted, on the basis
107 of particle size data, that 'local' sources could be important. Norris and Webb (1976)
108 agreed with the early studies, citing the Mojave River as the main source, but also
109 suggested that Soda Lake could be an important source. Yeend et al. (1984) reported
110 mineralogical data for the Kelso Dunes, but departed from earlier studies in proposing
111 that in addition to the Mojave River, Kelso Wash could have been an important source
112 for the dunes, as well as sediments derived from the Granite Mountains and Providence
113 Mountains. Paisley et al. (1991) returned to the concept of the Mojave River as the
114 primary source. Lancaster (1993, 1994) conducted detailed mapping of the different
115 geomorphic units within the Kelso Dunes and, in agreement with Yeend et al. (1984)
116 hypothesized that the Mojave River, Soda Lake, Kelso Wash, the Granite Mountains, and
117 the Providence Mountains could all have been contributors to the dunes. Three papers on
118 the luminescence geochronology of the dunes proposed the Mojave River and Soda and
119 Silver Lakes as sources (Edwards, 1993; Clarke, 1994), or a combination of these three

sources and the Granite Mountains (Wintle et al., 1994), although no mineralogical or geomorphic data were presented to test these hypotheses. Ramsey et al. (1999) presented both traditional mineralogical data and thermal infrared remote sensing data to propose that the dunes could have had a complex origin, with contributions from the Mojave River, Kelso Wash, the Providence Mountains, and the Granite Mountains. Lancaster and Tchakerian (2003) agreed with the multiple-source concept articulated by Ramsey et al. (1999). Further, Sweeney et al. (2013) reported evidence that fans and dry washes of the Providence Mountains are an important source of aeolian sand that is a part of soil Av horizons in the region. Nevertheless, Warren (2013) and Lorenz and Zimbelman (2014) returned to the simpler model of the Mojave River as the primary source of the Kelso Dunes.

Dune fields of course can be derived from either a single source or multiple sources, and sources can also change over time. A complex origin for a desert dune field, with multiple or changing sources, can be challenging to document, but recent studies have shown that multiple sources have contributed to dune fields in a number of arid regions. For example, in addition to Ramsey et al. (1999), cited above, for the Kelso Dunes, Scheidt et al. (2011) demonstrated that the Gran Desierto dune field in northwestern Mexico is derived in part from ancestral Colorado River sediments, but also has had contributions from local sources. Although the Negev and Sinai dunes of Israel and Egypt are derived primarily from Nile River sands, Muhs et al. (2013a) showed that sand derived from local carbonate rocks have also made contributions to the dunes.

Herein, we use simple geochemical methods for assessing the origin of Kelso Dunes using the composition (K, Ba, Rb) of one of the most common minerals found in dune

fields, K-feldspar. Both feldspars and micas host K as a major element, but the major K-bearing mineral phase in most dune fields is K-feldspar, either microcline or orthoclase. Micas, the second most common K-bearing group of minerals, are rarely reported in dune fields, and it is not difficult to understand why this is the case. K-feldspar is approximately seven times as abundant as biotite in granite and it is about five times as abundant as biotite in granodiorite (Wedepohl, 1969, p. 248). These two rock types are estimated to comprise ~78% of the upper continental crust of the Earth. Thus, on the surface of the Earth, there is simply less mica available, compared to K-feldspar, for delivery to dune fields from average upper crustal rocks. Thus, bulk analyses of sediment for concentrations of K, as well as Rb and Ba (trace elements that follow K; see Heier and Adams, 1964; Lange et al., 1966), typically will reflect K-feldspar compositions. Muhs et al. (2008, 2013b) showed that K/Rb and Ba/Rb have the ability to distinguish loess deposits of different provenance in both Alaska and the North American midcontinent. Comparison of K-feldspar composition of Kelso Dunes with compositions of candidate source sediments can potentially identify the most important source or sources of aeolian sand. To our knowledge, K/Rb and K/Ba values have not been applied to deciphering provenance of sand in dune fields. Here, we apply this to the Kelso Dunes as a case study, and then suggest how the methods could be applied to other dune fields in west Africa, southwest Africa, north Africa, and the Middle East, where the sources of sand are unknown or controversial.

2. Study area

2.1. Geology of the Kelso Dunes area

The Kelso Dunes are situated in the arid Mojave Desert of southern California, a region that is part of the Basin and Range physiographic province. The Kelso Dunes occupy a structural basin surrounded by small, dominantly granitic mountain ranges (with maximum elevations ranging from ~1200 m to ~2200 m), in turn flanked by alluvial fans (Figs. 2, 3). Sediments from these fan deposits and nearby dry-wash drainages are the major candidates for sources of aeolian sand. Miller et al. (1991) provide a good summary of the bedrock geology of the mountain ranges surrounding the Kelso Dunes. To the north, the Kelso Mountains are composed of early Proterozoic granitoids and gneiss, as well as Cretaceous monzogranite. East of the dune field, the Providence Mountains are comprised of early Proterozoic gneiss, Jurassic and Cretaceous granite, monzogranite, and quartz monzonite, as well as carbonate and siliciclastic sedimentary rocks. To the south, the Granite Mountains are composed of Jurassic and Cretaceous granite, granitoids, granodiorite, monzogranite, diorite, and quartz diorite gneiss. Budweiser Wash, situated largely to the west of the Kelso Dunes, carries sediment from both the Granite Mountains, described above, as well as the Old Dad Mountains (composed of Jurassic granitoids and Tertiary rhyolite) and the Bristol Mountains (composed largely of Jurassic granitoids and metavolcanic rocks). The source of Kelso dune sand cited most frequently, as mentioned above, is the Mojave River, which heads in the San Bernardino Mountains. In the upper drainage basin of this river, the major bedrock types are Cretaceous or Jurassic granodiorite and quartz monzonite, along with some metamorphic rocks of uncertain age (Bortugno and Spittler, 1986).

The age of aeolian sand at Kelso Dunes has been estimated primarily on the basis of luminescence geochronology. From infrared stimulated luminescence (IRSL) methods,

some parts of the Kelso Dunes appear to have formed by the late Pleistocene (~16 ka), based on data in Clarke (1994). She also reports that some aeolian sands, particularly in the eastern parts of the dune field (units I-V of Lancaster [1993, 1994]; see below), are interbedded with alluvial fan deposits from the Providence Mountains, observations confirmed by detailed mapping and stratigraphy conducted by McDonald et al. (2003). More recent studies by Sweeney et al. (2015) show that there may have been a long and complex history of alternating aeolian and fluvial sedimentation for dunes in this area, as well as in sand ramps along the margins of Kelso Dunes. Nevertheless, IRSL geochronology shows that the major landforms in Kelso Dunes formed or were reformed during the Holocene and particularly in the past ~4,000 years (Edwards, 1993; Clarke, 1994; Wintle et al., 1994). All aeolian sands sampled in the present study were taken either from active dune sand or, where stabilized, from dunes considered to be of late Holocene age.

2.2. Climatology of the Kelso Dunes area

The climate of the Kelso Dunes area is warm and arid. All data presented here are for the 1981-2010 period of record from the two weather stations closest to the Kelso Dunes, Daggett, California (N34°52'; W116°47'), ~94 km to the west and Baker, California (N35°16'; W116°04'), ~46 km to the northwest. Mean July temperatures at Daggett and Baker are 31.6°C and 30.9°C, respectively, with mean January temperatures of 9.3°C (Daggett) and 9.2°C (Baker). Mean monthly low temperatures are above freezing in all months. Aridity of the Mojave Desert is due to its position in the lee of southern California's mountains. The San Bernardino Mountains (to the south), San Gabriel Mountains (to the southwest), Tehachapi Mountains (to the northwest), and the coastal

mountain ranges (to the northwest) are high enough that much, though not all, of the moisture from frontal systems derived from the north Pacific Ocean is blocked from transport to the east, leaving the Mojave Desert in a rain shadow. Thus, mean annual precipitation at Daggett is just 105.4 mm and at Baker it is 104.9 mm. Measurable precipitation, on average, is recorded in all months, but is typically highest in winter (December, January, and February) from Pacific-derived frontal systems. Rainfall in the winter months accounts for approximately 58% (Daggett) to 49% (Baker) of the mean annual precipitation. There is a smaller, secondary rainfall peak, primarily in the form of convective storms, in late summer/early fall (July, August, and September). Late summer/early fall precipitation is due to a moderate monsoonal flow of tropical moisture derived from the Pacific Ocean and Gulf of California off the coasts of Baja California and mainland Mexico. This warm-season precipitation accounts for ~22% (Daggett) to 28% (Baker) of the mean annual rainfall. During El Niño years, when warm, tropical waters of the equatorial western Pacific Ocean migrate eastward, winter precipitation in the Mojave Desert is significantly higher, whereas during La Niña years, drier winters occur (Hereford et al., 2004).

Winds in the Mojave Desert are important in understanding both dune geomorphology and potential source sediments. Of particular importance are those winds above the threshold velocity for transport of sand-sized particles. For loose, non-vegetated, medium-sized (500-250 μm) sand particles, the threshold velocity is ~5-6 m/s, when recorded with an anemometer that has a standard height of ~10 m. Resultant annual drift potentials (RDP), using the method of calculation in Fryberger and Dean (1979), show a wide range of values in the areas of dunes in the southwestern USA and northwestern

Mexico, although all show dominant sand-moving winds from the northwest, west, or southwest (Fig. 1). RDP for Daggett, California is ~600 vector units, putting it in the 'high-energy wind environment' class of localities of Fryberger and Dean (1979).

Although the resultant drift direction for the area around Daggett is northwest-to-southeast (Fig. 1), there is a significant seasonal variability in wind directions in the Mojave National Preserve area. We generated wind roses for four representative months of the year (Fig. 4) for the Mojave 'Sink' area (location shown on Fig. 3). The plots show the frequency of winds above a threshold velocity of 5 m/s and therefore display only winds strong enough to entrain sand-sized particles. Common to all four seasons are frequent winds from the west and southwest. In addition, however, fall (represented by October) and winter (represented by February) frequently have winds from the northwest and north. Summer (represented by July) frequently has winds from the southeast, likely due to the monsoonal flow of air from the south.

3. Methods

In the Kelso Dunes, aeolian sands from most of the major geomorphic units (Fig. 5) defined by Lancaster (1993, 1994) were collected for mineralogical and geochemical analyses. For source sediments, we collected samples from the following drainages: (1) Mojave River wash, from Afton to ~1 km east of Crucero (n=36); (2) Budweiser Wash, along an ~8 km reach between the Granite Mountains and the Bristol Mountains (n=10); (3) fan deposits of the Granite Mountains, collected along a ~16 km transect to the north of the mountains, from just west of Cottonwood Wash to Budweiser Wash (n=27); (4) Providence Mountain fan deposits, collected to the west of the mountain front, on a 4.4

km transect along the Vulcan Mine Road and a 5.4 km transect along the pipeline access road to the south of the Vulcan Mine Road (n=25); and (5) Kelso Wash, sampled from eight separate small washes, along a 1.8 km transect northwest of the town of Kelso (n=15). As discussed above, a number of investigators have hypothesized that sediments in the basins of Silver Lake and Soda Lake playas could have been sources of sand for the Kelso Dunes. Our examination of these areas, however, indicates very little sand-sized material is available for aeolian transport, except along the margins of the playas. Moreover, as pointed out by Wells et al. (2003) and Enzel et al. (2003), the vast majority of sediment reaching these basins, including the minor sand-sized component, is ultimately derived from the Mojave River, which we sampled separately.

Aeolian sands from each of the major geomorphic units were analyzed as bulk samples, with no pretreatments other than pulverization. Alluvial samples from the potential source sediments were pretreated to yield a particle size distribution similar to that of aeolian sand. After disaggregation and removal of coarse gravel, the samples were placed in de-ionized water, a Na-pyrophosphate dispersant was added, particles in suspension were stirred, and then allowed to sit overnight. After this treatment, the samples were stirred again by ultrasonication and then wet-sieved to remove fine gravel and coarse sand ($>500\text{ }\mu\text{m}$). Silt and clay were removed by sieving with either a $53\text{ }\mu\text{m}$ or $63\text{ }\mu\text{m}$ sieve; no materials finer than $53\text{ }\mu\text{m}$ are present in any of the samples. Thus, the final sediment separates contained grains ranging from medium sand to very fine sand, similar to the size range of well-sorted dune sands at Kelso Dunes (see data in Lancaster, 1993, 1994).

Mineralogy was determined by X-ray diffractometry (XRD) on pulverized, randomly packed powder mounts. Percentages of quartz, K-feldspar, plagioclase, and calcite in Kelso dune sands and samples from other dune fields were estimated using a method developed by Dr. Amir Sandler of the Geological Survey of Israel (written communication, 9 August 2012), wherein the following factors are applied to XRD peak heights at two-theta, summed, and calculated as percentages: quartz, 26.6° , x 1.0; K-feldspar, 27.4° , x 3.0; plagioclase, 27.8° , x 3.5; and calcite, 29.4° , x 1.1. Crouvi et al. (2009), Enzel et al. (2010) and Muhs et al. (2013a) have used this method previously and results show good agreement with major element geochemistry. Abundances of K, Ca, Ti, Rb, Sr, Y, Zr, Nb, Ba, La, and Ce were determined by energy-dispersive X-ray fluorescence (ED XRF), following the methods in Muhs et al. (1995). USGS rock standard GSP-1 was included with each sample batch and resulting concentrations for this standard agree well with published values for most elements (see Muhs et al., 1995, their Table 2). For this rock standard, ED XRF element concentrations agree with published values within 5% (K), 0.4% (Rb), and 0.8% (Ba). Precision for these elements (± 1 sigma) is 0.9% (K), 1.2% (Rb), and 0.8% (Ba). We note that there could be minor differences in element abundances determined by ED XRF compared to wavelength-dispersive X-ray fluorescence (WD XRF). In particular, concentrations of K by ED XRF may be biased high by ~5%, based on routine runs of USGS rock standards (e.g., a sample with a K_2O concentration of 3.5% by ED XRF might yield a concentration of 3.3% by WD XRF). Thus, we caution readers that element concentrations determined by WD XRF in the literature may be only approximately comparable to data presented here. Mineralogical interpretations of all geochemical data presented here are based on first

principles of geochemistry and element substitution in minerals (Mason and Moore, 1982).

4. Results

4.1. Geomorphology of Kelso Dunes

Lancaster (1993, 1994) showed that despite the small area occupied by the Kelso Dunes, there is a diverse geomorphology, and he identified 14 distinct landform units. These landform units can be distinguished on Landsat ETM+ imagery (Fig. 5), but we also examined larger-scale stereoscopic aerial photography (black-and-white and color aerial photographs from the U.S. Geological Survey's NAPP (National Aerial Photography Program; images acquired in 1989, 1994, and 1995) and oblique aerial photography to assess the geomorphology of the dune field (Figs. 6, 7). Three dune units (VI, VIII, and X) are dominantly vegetation-free (Figs. 6-8), and a fourth unit (III) is only partially vegetated (Fig. 6). The remaining dune units are all stabilized to one degree or another by desert psammophytic scrub vegetation, including big galleta grass (*Hilaria rigida*), dune panic grass (*Panicum urvilleaneum*), and creosote bush (*Larrea tridentata*) (Thorne et al., 1981) and experience little or no active sand transport at present (Figs. 5-8).

The three dominantly active units, VI, VIII, and X, exhibit the most distinctive suites of aeolian landforms and all are complex dunes. At the broadest scale, all three of these active dune units consist of southwest-to-northeast trending ridges that could be former linear dunes. If so, these orientations would imply winds dominantly from the southwest (or, less likely, from the northeast). Superimposed on these landforms are much smaller

barchanoid ridges, parabolic dunes, and occasional star dunes; the latter are rare in North American dune fields. Although the large linear ridges in these units could imply southwesterly winds during their formation, the smaller (and younger) barchanoid ridges and parabolic dunes suggest winds from the west to west-northwest. The star dunes are also superimposed on (and are therefore younger than) the large linear dunes and imply variable paleowinds during their time of formation (Table 1).

Most stabilized dune units in the eastern part of the area (those east of Cottonwood Wash; Figs. 5-7) indicate that recent dune-forming paleowinds were from the west or west-northwest, based on geomorphology (Table 1). Unit I consists largely of sheet sands, but it and Unit V also have some degraded parabolic dunes that indicate westerly (unit I) or northwesterly (unit V) paleowinds. Landforms are better preserved in units II and III and consist of linear dunes with superimposed parabolic dunes, and the latter imply northwesterly or westerly paleowinds. Unit IV consists of degraded linear dunes and sand sheets, but unlike the other eastern dune units, has linear dune orientations that could indicate the possibility of paleowinds from the southwest.

Stabilized dune units in the western part of the area (west of Cottonwood Wash) for the most part imply past winds that were generally westerly. Units VII, XII, and XIV all consist of parabolic dunes that indicate winds from the northwest (VII) or west (XII and XIV). Units IX and XIII consist either of linear dunes with superimposed parabolic dunes (IX) or linear dunes that grade southeastward into parabolic dunes (XIII) that indicate dune-forming winds from the west (IX) or northwest (XIII). The major exception to the overall pattern of westerly or northwesterly paleowinds in this part of the Kelso Dunes is unit XI. This unit consists of linear dunes that grade eastward into

parabolic dunes, but the orientations of both the linear dunes and the parabolic dunes indicate paleowinds from the west-southwest (Fig. 5).

4.2. Mineralogy of Kelso Dunes

There have been highly divergent estimates of the abundances of common minerals in the Kelso Dunes. Sharp (1966) identified quartz, K-feldspar, plagioclase, and a variety of heavy minerals in the Kelso Dunes, but did not make estimates of abundances of mineral species. However, Yeend et al. (1984), Paisley et al. (1991) and Ramsey et al. (1999) reported average quartz contents of 70-80%, ~90%, and ~42%, respectively. The wide range in estimates of mineral species abundances led us to re-examine the composition of the Kelso Dunes, and we conducted XRD analyses on 74 samples from 12 of the 14 dune units.

Based on our measurements of the Kelso Dunes, quartz averages 33.5% ($\pm 7.8\%$, one sigma), K-feldspar content averages 24.8% (± 6.1), and plagioclase averages 41% ($\pm 7.6\%$). Calcite was also detected in small amounts in a few samples, primarily in dune unit I. Based on the relative abundance of quartz, plagioclase, and K-feldspar, the Kelso Dunes have a bulk composition similar to that of granite (slightly more than half the samples) or granodiorite (slightly less than half the samples), based on the classification system of igneous rocks by Streckeisen (1976). Given the granitic composition of the most of the rocks in all the possible mountain ranges that could deliver sediments to the Kelso Dunes, these results are not surprising. In this regard, the Kelso Dunes have a mineralogical composition that is similar to other Mojave Desert dune fields, such as the Cadiz and Danby dunes, as well as the Sonoyta dunes, in Sonora, Mexico (Fig. 9). The

Cadiz and Danby dunes are likely derived from nearby granitic mountain ranges (Muhs et al., 2003) and the Sonoyta dunes are likely derived in part from sediments of the Sonoyta River (Scheidt et al., 2011), which drains rock bodies composed of granite and gneiss in northern Sonora. In contrast, the Algodones dunes in southern California and the Parker dunes in western Arizona are both thought to be derived primarily from quartz-rich Colorado River sediments (Muhs et al., 1995, 2003) and show much lower concentrations of K-feldspar and plagioclase compared to the Kelso Dunes (Fig. 9). In good agreement with the mineralogy, concentrations of K_2O and CaO in Kelso Dunes (Fig. 10a) also fall within the ranges for these elements in granites and granodiorites (Nockolds, 1954). Concentrations of K_2O and CaO in Kelso Dunes are similar to other dune fields in the Mojave Desert, such as the Cadiz-Danby dunes and the Dale Lake sand sheet, but differ from the quartz-rich Algodones and Parker dunes (Figs. 10b, c, d).

A number of previous workers have noted the presence of lag concentrations of heavy minerals on the crests of the active portions of the Kelso Dunes (Sharp, 1966; Paisley et al., 1991; Lancaster, 1994). These concentrations of dark minerals are visually striking and can be observed from considerable distances in the field (Fig. 8c). Sharp (1966) and Yeend et al. (1984) identified a number of heavy minerals in the Kelso Dunes, including amphibole, pyroxene, zircon, biotite, epidote, apatite, sphene (titanite), garnet, rutile, tourmaline, monazite, ilmenite, magnetite and cassiterite. Yeend et al. (1984) also reported that the surface concentrations of heavy minerals consist largely of amphibole and magnetite, with magnetite concentrations of up to 10%. Geochemical data support previous interpretations of the presence of these minerals (Fig. 11). Concentrations of Ti (and Nb, correlated positively with Ti; note, however, that some samples approach the

lower detection limit for Nb) could be due to the presence of sphene, rutile, ilmenite and magnetite (if present as titanomagnetite). Pyroxene, hornblende and biotite can also host trace amounts of Ti. The presence of zircon is confirmed with concentrations of Zr that are typically less than ~500 ppm, but range as high as ~2000 ppm in one sample. We also measured two of the light rare earth elements (REE), La and Ce, and Y - an element with properties similar to the REE. La, Ce and Y are often present in minerals such as sphene and zircon, and all three of these elements are positively correlated with concentrations of Ti and Zr. Abundances of these elements are extremely high in magnetite-rich surface lags (Fig. 11).

4.3. Geochemistry of Kelso Dunes

The relatively high concentrations of K-feldspar in the Kelso Dunes allow the use of K/Rb and K/Ba to distinguish geochemical signatures of possible source sediments. As a first step in testing some of the hypothesized source sediments for the Kelso Dunes, we examined the composition of the small dunes at Devils Playground, to the northwest of the main dune field. If Mojave River sediments were the main source of the Kelso Dunes, as proposed by many previous investigators, then the Devils Playground dunes should have a composition that is similar to the river sands, because this dune field is situated between the Mojave River sink and Kelso Dunes (Fig. 3). K/Ba and K/Rb values from these dunes are not significantly different from those in Mojave River sediments, which supports this hypothesis (Fig. 12a). Devils Playground dunes also show only partial overlap with the K/Rb and K/Ba fields defined by Providence Mountains fan alluvium and Budweiser Wash alluvium and no overlap with Kelso Wash alluvium (Figs.

12b, c). Nevertheless, Devils Playground dunes fall completely within the field defined by Granite Mountains fan alluvium (Fig. 12d).

To the southeast of the dunes in Devils Playground lie the western group of dune units in Kelso Dunes (units VI-XIV), situated west of Cottonwood Wash (Figs. 3, 5, 7). Aeolian sands from the western dune units have K/Ba and K/Rb compositions similar to Devils Playground dunes, Mojave River sands (Fig. 13a), Granite Mountains fan alluvium (Fig. 13b) and Budweiser Wash alluvium (Fig. 13d). As with the Devils Playground dunes, the western dune sands show only partial overlap with Providence Mountains fan alluvium (Fig. 13c), and no overlap at all with Kelso Wash alluvium (Fig. 13d).

The eastern dune units have compositions that differ significantly from both the Devils Playground dunes and western dunes. In contrast to the latter, the eastern dune units show somewhat less variable K/Rb, but much more variable K/Ba (Fig. 14). The eastern dune units show only a partial overlap with the composition of Mojave River sands (Fig. 14a) and no overlap with either Budweiser Wash or Kelso Wash (Fig. 14d). The dunes in this area do show, however, considerable overlap with the K/Ba and K/Rb fields defined by both Granite Mountains fan alluvium and Providence Mountains fan alluvium (Figs. 14b, c).

5. Discussion

5.1. Origin of the Kelso Dunes

One of the most fundamental differences between the results reported here and those of previous researchers is mineralogical composition. Our mineralogical results differ

435 significantly from those of Yeend et al. (1984) and Paisley et al. (1991) but are close to
436 those of Ramsey et al. (1999). We found that quartz averages ~33%, plagioclase
437 averages ~41%, and K-feldspar content averages ~25%. If Kelso Dunes sands had quartz
438 contents as high as ~90% (Paisley et al., 1991) or even ~80% (Yeend et al., 1984), they
439 would be considered to be mineralogically mature. Many of the world's larger dune
440 fields are in fact mineralogically mature (see review in Muhs, 2004), but based on
441 processes whereby dune fields can achieve such a composition, we would not expect
442 Kelso Dunes to be quartz-rich. Depletion of less-resistant minerals such as calcite,
443 plagioclase, and even K-feldspar can take place by chemical weathering, while dunes are
444 stabilized by vegetation under a humid climate. This mechanism is unlikely for many
445 dune fields found in semiarid or arid regions, although it cannot be dismissed entirely,
446 because in the past some deserts (including the Mojave Desert: see Spaulding, 1995)
447 were more humid than they are now. However, a more likely mechanism of calcite,
448 plagioclase, or K-feldspar depletion in dune fields is by mechanical breakdown, either
449 from abrasion or ballistic impacts during aeolian transport. Silt-sized particles of calcite,
450 plagioclase, or K-feldspar that result from this physical disintegration can then be
451 removed from a dune field altogether by aeolian transport in suspension, leaving behind a
452 residual, quartz-rich, sand grain population. This process can be important where there is
453 either long-distance aeolian transport of sand particles far from the source or there is
454 repeated reactivation and reworking of sand grains over long periods within a desert
455 basin. Based on the setting of Kelso Dunes, long-distance transport does not apply and
456 based on the IRSL chronology, reactivation and reworking of grains over many periods
457 of dune activity does not seem likely. Inheritance of a quartz-rich sand population can

also explain mineralogical maturity (see Muhs et al., 1995, 2003). However, both the bulk mineralogical composition of Kelso Dunes sands (Fig. 9) and their major element composition (Fig. 10) show that the dunes differ little from the composition of average granite or granodiorite. This, along with the granitic composition of the surrounding mountains, implies only short distances of aeolian transport and little or no reworking.

The *lack* of mineralogical maturity and abundance of K-feldspar in Kelso Dunes actually provides the key to assessing likely source sediments, through the use of K/Rb and K/Ba values. Geochemical data presented here do not provide a unique solution to the origin of Kelso Dunes. Nevertheless, although several scenarios of sediment transport pathways to the dunes can be envisioned, all of them require a revision of the simple, single-source, Mojave River wash model reported by numerous previous investigators.

Devils Playground dunes have K/Rb and K/Ba compositions that fall within the range of both Mojave River wash sediments and sediments from the Granite Mountains. The Providence Mountains and Budweiser Wash are less likely as sources for Devils Playground dunes and Kelso Wash does not appear to be an important source at all (Fig. 12). Granite Mountains fan alluvium as a source for Devils Playground dunes would require a dominance of winds from the southeast, which occur only during mid-to-late summer months (and with less frequency, in winter), whereas Mojave River wash sediments as a source would require westerly to northwesterly winds, which occur year-round. We infer that Devils Playground dunes are dominantly derived from Mojave River wash sediments, in agreement with hypotheses of many previous researchers.

The western dune units (VI-XIV) do not differ significantly in K/Rb and K/Ba composition from Devils Playground dunes or Mojave River sediments (compare Figs. 12 and 13). As with the Devils Playground dunes, Providence Mountains fan sediments and Kelso Wash sands can be eliminated as likely sources for the western dune units. The most straightforward interpretation is that the western dune fields are simply the downwind recipients in a sediment cascade from (1) the Mojave River sink to (2) the Devils Playground dunes to (3) the western dune units. The orientations of stable dunes in western dune units VII, IX, XIII, and XIV all imply winds out of the west or northwest, consistent with this interpretation. Furthermore, the orientations of the active, secondary barchanoid-ridge dunes, star dunes, and parabolic dunes of western dune units VI, VIII, and X also imply winds from the west or northwest. Winds from the west occur year-round in the Kelso Dunes (Fig. 4).

Although the concept of derivation of the western dunes exclusively from the Mojave River has appeal because of its simplicity, this explanation overlooks the fact that the western dune fields also have K/Rb and K/Ba compositions that fall within the ranges of both Granite Mountains fan alluvium and Budweiser Wash alluvium (Fig. 13). Because of where the western dunes are situated, these sources would require winds from the southwest, but winds from this direction do occur year-round (Fig. 4). Further, the southwest-to-northeast orientations of the large linear ridges of active dune units VI, VIII, and X (on which the barchanoid-ridge, star, and parabolic dunes are situated) and the stable linear dunes of unit XI are consistent with dune-forming winds dominantly from the southwest (Table 1). Thus, modern winds, dune orientations, and sand geochemistry would permit a model of aeolian delivery of sand from Budweiser Wash or fans of the

Granite Mountains to the Kelso Dunes. It is worth noting that the alternative interpretations presented here for the origin of the western dunes are not mutually exclusive. Southwesterly winds transporting sand from Budweiser Wash or fans of the Granite Mountains could have alternated with westerly winds transporting sand from the Mojave River. The results reported here permit a departure from the simple, single-source Mojave River model advocated by most previous researchers.

The greater variability in K/Ba values in sands from the eastern dune units (I-V), east of Cottonwood Wash, contrasts with the composition of the western dunes and Devils Playground dunes (Fig. 14). There is only a partial compositional overlap of these dune fields with Mojave River sands and no overlap with Budweiser Wash or Kelso Wash sands. However, the eastern dunes have compositions that fall well within the range of alluvium from both the Granite Mountains and the Providence Mountains. From these observations, it can be inferred that the Mojave River certainly cannot be the sole source of the eastern dune units. A complex origin could be inferred if we interpret the eastern dune units to be derived from a mix of Budweiser Wash sands and Kelso Wash fans, a possibility permitted by the compositional array shown in Fig. 14d. We question this interpretation, however, because it is difficult to envision how sands from Budweiser Wash could be transported to the eastern dunes without being mixed with sands from the Mojave River. A simpler explanation is that the eastern dunes are derived from the easternmost fans of the Granite Mountains, an interpretation also permitted by the K/Rb and K/Ba data.

The other alternative permitted by the data is derivation of the eastern dunes dominantly from fan sediments of the Providence Mountains. This would explain the

distinctive composition of the eastern dunes as compared with the western dunes, but is a bit more difficult to reconcile with contemporary wind regimes. Fan sediments from the Providence Mountains are situated mostly to the east of the eastern dune fields (Figs. 3, 5, 7). Winds from the southeast occur only in late summer and, with less frequency, in late winter (Fig. 4). Although derivation of the eastern dune fields from Providence Mountains fan sediments under easterly winds is possible, stratigraphic and geomorphic studies by McDonald et al. (2003) suggest another explanation. Work by these investigators on the geomorphology and stratigraphy of Providence Mountains fan deposits shows that older fan sediments are found beneath some of the aeolian sands of the eastern dune fields. Further, aeolian sands and fan sediments are interbedded in parts of the eastern dune fields and in places fan sediments are found west of some of the eastern dunes (see Fig. 8d for an example). Thus, distal fan sediments from the Providence Mountains, found to the west of the eastern dune fields, could have been the source of these dunes and would require only winds from the west, which occur year-round (Fig. 4). We note that Sweeney et al. (2013) found that fans and dry washes from the Providence Mountains are an important source, via aeolian transport, of the sand fraction in eastern Mojave Desert soil Av horizons, which supports our interpretations.

Based on thermal infrared multispectral scanner (TIMS) data, Ramsey et al. (1999) also inferred that fan sediments from the Providence Mountains were an important source of sand for the Kelso Dunes. Of the eastern dune fields, Ramsey et al. (1999) inferred that the greatest influence of Providence Mountains fan sediments, with respect to K-feldspar inputs, was in dune unit V, with the least influence on dune unit II. With the

data presented here, we agree with Ramsey et al. (1999) that Providence Mountains fan sediments were likely an important influence on the origin of the eastern dune fields.

We summarize our interpretations of the origin of the youngest aeolian sands in the Kelso Dunes region in two different scenarios shown on Landsat imagery (Figs. 15a,b). In the interpretation shown in Fig. 15a, western dunes are derived from Mojave River Wash sands from northerly (fall and winter) and westerly (year-round) winds, combined with possible inputs from Budweiser Wash from southwesterly winds (year-round). Eastern dunes are derived from Providence Mountain fan sediments from southeasterly winds (summer) and/or Granite Mountains fan sediments and Budweiser Wash sediments from southwesterly winds (year-round). In the interpretation shown in Fig. 15b, western dunes are derived from various sources as described in the first scenario. However, in this alternative sequence of events, the eastern dune units are derived from distal Providence Mountain fan sediments that were deposited close to Kelso Wash and then entrained by winds from the west (year-round) and transported back up onto proximal Providence Mountains fan sediments. This latter scenario would also imply that Cottonwood Wash (Figs. 5-7) has been a significant barrier to sand transport from sources to the west and northwest (i.e., the Mojave River). Of the two scenarios, we favor the second, because the dominance of westerly winds in the region supports it and it is consistent with stratigraphic evidence of interbedded aeolian sand and Providence Mountains fan deposits, documented by McDonald et al. (2003). With either interpretation, however, it is clear that the origin of the Kelso Dunes is more complex than the single-source, Mojave River model that has been envisaged by most investigators for the better part of a century.

Information on the composition of Kelso Dunes provides constraints on the age of the dune field. Although local sources of sand are clearly important, the volume of sediment generated from them is likely insufficient compared to the volume of sand in the dune field, which is estimated to be $\sim 1 \text{ km}^3$ (Lancaster, 1993). The lower reaches of the Mojave River have a complex history of drainage integration (Enzel et al., 2003), and recent studies indicate that the Mojave River reached the Soda Lake basin and present day Mojave River Sink via Afton Canyon sometime after 25 ka (Reheis and Redwine, 2008; Reheis et al., 2012). Based on sediment composition, it is likely that the dune field has accumulated episodically over the past ~ 25 ka, although small dune areas fed by local sources may have existed prior to that time.

5.2. Use of K/Rb and K/Ba to understand aeolian sand provenance in large sand seas on other continents

One conclusion from our studies is that even though the Kelso Dunes are a relatively small dune field, more than one source of sand is apparent from the data we present here. Thus, it seems likely that larger dune fields, and certainly ergs or large sand seas, also likely have complex origins. The use of K/Rb and K/Ba in K-feldspar as a provenance indicator leads us to propose that this method could be applied to investigations of dune origins in many of the world's large sand seas. Sand seas cover a significant amount of land area in the low-latitude deserts of Africa, Asia, and Australia and at higher latitudes of Asia where structural basins are rimmed by mountains that provide rain-shadowed topographic lows (Fig. 16). Here we present some examples of dune fields on other continents whose origins are not understood or are controversial and where we believe the methodology applied to the Kelso Dunes could be used for determination of sand

provenance. In some cases, there are also good reasons why the approach would *not* be useful and we give examples of these as well.

5.2.1. Saharan and Sahel dunes of Mauritania and Senegal

Spectacular linear dunes are found over a large area of western Mauritania in the southern Sahara Desert (Fig. 16). Long axes of these dunes are aligned northeast-to-southwest, in good agreement with resultant drift directions (RDDs) computed by Fryberger and Ahlbrandt (1979) and Breed et al. (1979). Lancaster et al. (2002) documented at least three ages of dunes with luminescence dating in this region and showed that although all were generated by dominantly northeasterly winds, each phase of dune-building was associated with a slightly different wind regime. Many of these dunes are still active today (Figs. 17a, b). To the south in Senegal, dunes are also found but are stabilized by vegetation (Figs. 17a, c). In a now-classic paper, Grove and Warren (1968) hypothesized that the stabilized linear dunes in Senegal are continuations of what were once more extensive linear dunes originating farther north, in Mauritania. Further, they speculated that movement of aeolian sand would have blocked the Senegal River in its lower reaches, presumably filling the valley downstream and allowing aeolian sand migration to continue to the southwest. Mapping from early Landsat imagery by Breed et al. (1979) confirms the possibility of such a sequence of events. To our knowledge, this intriguing hypothesis has never been tested with any rigorous field and laboratory studies. An alternative explanation is that the stabilized dunes to the south derived their sand independently from the Senegal River at some hypothetical time or times when there was at least seasonally low discharge. Determination of K/Rb and K/Ba values in sands from the dunes in Mauritania, the Senegal River (upstream of the dunes, to avoid problems of

fluvial reworking of dune sand from the north), and dunes in Senegal (the Ferlo-Cayor dune field) could provide a test of this hypothesis. Dunes in Mauritania contain 0.2-0.7% K₂O (Lancaster et al., 2002), indicating that they are likely mineralogically mature, or rich in quartz. Nevertheless, even this amount of K₂O indicates the likely presence of K-feldspar in sufficient amounts to allow a provenance study using K/Rb and K/Ba.

5.2.2. Namib Sand Sea, southwestern Africa

The Namib Sand Sea is a large (~34,000 km²), active dune field along the coast of Namibia in southwestern Africa (Figs. 16, 18). A wide variety of aeolian landforms can be found in the Namib sand sea, including barchanoid ridges (particularly on the coast), linear dunes (inland, in the southern part), linear dunes with superimposed star dunes (inland in the southern part), and sheet sands (northern, eastern and southern margins) based on studies by Breed et al. (1979) and Lancaster (1989). Dune orientations indicate that major dune-forming winds came mainly from the south, which is consistent with modern resultant drift directions (Fig. 18). Lancaster (1989) has shown that physical properties of the dune sediments vary spatially: dunes are redder, and particles are more rounded and better sorted in a west-to-east direction (Fig. 18), suggesting that the youngest sediments are near the coast and older sediments are found inland. Based on data from cosmogenic isotopes, the Namib Sand Sea may have been in existence as a dune field for at least the past million years (Vermeesch et al., 2010).

The origin of dune sand in the Namib Sand Sea has been discussed for the better part of a century and debate continues to the present decade. Gevers (1936), Logan (1960), Hallam (1964) and Rogers (1977, 1979) all thought that sand in this dune field was ultimately derived from sediments of the Orange River, which is upwind of the Namib

640 Sand Sea (Fig. 18). Besler and Marker (1979, p. 159) and Besler (1980, 1984), however,
641 challenged this hypothesis and proposed that the Tsondab Sandstone (referred to as the
642 Namib Sandstone by these investigators), which crops out to the east of the sand sea,
643 below the Great Escarpment (Fig. 18), was the more likely source. These workers
644 envisioned fluvial erosion of the sandstone with delivery to the west, followed by aeolian
645 reworking to the north. Lancaster and Ollier (1983) conducted heavy mineral analyses of
646 both dunes in the Namib Sand Sea and possible sources, including the Tsondab
647 Sandstone, but not sands from the Orange River. They concluded that an Orange River
648 source from the south was likely, but pointed out that their data would permit an
649 interpretation that local additions from the Tsondab Sandstone could have been important
650 in the eastern part of the dune field. These investigators note that the light mineral
651 fraction of the Tsondab Sandstone is similar to that of the dune sands. Besler (1996), in a
652 follow-up study, maintained her assertion that the Tsondab Sandstone is a significant
653 source of Namib dune sand. White et al. (2007) took a different approach to studying the
654 Namib dunes and used remote sensing of Fe-oxide content in the dunes to infer at least
655 two sources of sand, one eroded from rocks of the Great Escarpment and the other
656 derived from southern, coastal sources. Vermeesch et al. (2010) determined U-Pb ages
657 of detrital zircons from both dunes of the Namib Sand Sea and one sample from
658 sediments of the Orange River. Although they interpret their age spectra from the
659 westernmost dunes to indicate an Orange River source, they also point out that dunes in
660 the eastern part of the sand sea have age spectra that differ significantly from those in the
661 western part of the dune field. These investigators did not provide data on the U-Pb ages
662 of zircons from the Tsondab Sandstone, so it is not possible to know at present if

contributions from this formation could be responsible for the difference in zircon age spectra. Garzanti et al. (2012) reiterated the U-Pb zircon age data of Vermeesch et al. (2010) and in addition provided detailed mineralogical data for four samples of Orange River sands and 12 Namib dune sands, as well as about a dozen sediment samples from rivers draining the Great Escarpment. These authors state that their mineralogical comparison of these samples ‘proves’ that most of Namib aeolian sand is derived from the Orange River. In making this rather categorical statement, the authors chose to ignore the larger heavy mineral data set presented by Lancaster and Ollier (1983), and even disregard the implications of the U-Pb zircon age spectra that they themselves presented. Finally, Stone (2013) presented a comprehensive review of all the previous studies and concluded that the majority of evidence pointed to the Orange River as the most important source, with local contributions from fluvial systems draining the Great Escarpment being secondary sources.

Lancaster and Ollier (1983) reported that the light mineral fractions of Namib Sand Sea sands consist of quartz, K-feldspar, and plagioclase. Garzanti et al. (2012) reported quartz contents of 66-79% and feldspar contents (presumably both plagioclase and K-feldspar) of 16-26%, with the remainder being rock fragments. The latter data agree with mineralogical and whole-sediment, major-element geochemical data reported by Muhs (2004). In that study, Muhs (2004) reported that quartz is the most important mineral in Namib dune sands, but all samples examined have abundant plagioclase, K-feldspar, and rock fragments. In addition, a few samples appear to have trace amounts of calcite and dolomite. Geochemical analyses of Muhs (2004) support the mineralogical data of Garzanti et al. (2012), as SiO₂ contents are 81-87%, Al₂O₃ contents are 5.0-6.8%, Na₂O

contents are 1.0-1.4% and K₂O contents are 1.0-2.0%. Thus, K/Rb and K/Ba analyses could easily be conducted on sediments of the Namib Sand Sea as there is sufficient K-feldspar, based on both mineralogical and geochemical data. Such analyses, if conducted over a major part of the dune field, along with analyses of Orange River sands, fluvial sands draining the Great Escarpment, and sands from the Tsondab Sandstone, could potentially provide information about whether there is more than one source for the Namib Sand Sea.

5.2.3. *Grand Erg Oriental, Algeria and Tunisia*

The Grand Erg Oriental is one of the largest sand seas in North Africa and may cover as much as 192,000 km² (Wilson, 1973). Despite its large size and mostly active condition (Fig. 19), there have been remarkably few studies of its origins, likely because of the inaccessibility of much of the dune field. Nevertheless, there have been a handful of studies that have offered hypotheses for its origin. Bellair (1953) is one of the few investigators who did mineralogical analyses of the Grand Erg Oriental. He reported that the dunes are dominated by quartz, but also locally contain gypsum and anhydrite. Using heavy mineral separates from dune sands collected along a ~400 km transect between El Oued and Ghudamis (Fig. 19), he found that the composition changed significantly ~300 km southeast of El Oued, near Sif-Fatima. From this, Bellair (1953) concluded that the Grand Erg Oriental was derived from diverse sources. Later researchers have based their hypotheses about dune origins in this large erg mainly on interpretations of maps, aerial photographs, and satellite imagery. Breed et al. (1979) interpreted Wilson (1973) to say that the Grand Erg Oriental was derived from alluvium in basins upwind of the dune field, but Wilson (1973) was actually not this specific. Although the Grand Erg Oriental

was a major focus of his study, with regard to sources Wilson (1973) stated simply that ergs in most deserts are derived from alluvium. Fryberger and Ahlbrandt (1979), interpreting Landsat imagery, hypothesized that large wadi systems draining bedrock uplands to the northwest were likely contributors to sand in the Grand Erg Oriental. Mainguet (1978, p. 20), however, rejected the idea of local alluvium being a major source of sand for the Grand Erg Oriental (and for other Saharan ergs as well) and thought that sand was derived from distant sources, hundreds or perhaps thousands of kilometers away. In a later paper (Mainguet, 1984), it was emphasized that the Grand Erg Oriental was a *source* of sand for other ergs, namely the Erg Chech (to the southwest) and the Erg Issaousane (to the south).

All of these hypotheses can be tested with a systematic program of sampling alluvial sands found upwind of the Grand Erg Oriental, transects of dunes within the erg itself, local sources of alluvium to the east and west of the erg, and sand within the Erg Chech and Erg Issaouane to the southwest and south. Analyses of K/Rb and K/Ba in K-feldspar would help to test the hypotheses of (1) diverse local contributions to the Grand Erg Oriental (Bellair, 1953); (2) wadi sources upwind of this sand sea (Fryberger and Ahlbrandt, 1979); and (3) erg-to-erg sand transport, with the Grand Erg Oriental being a source as well as a sink for wind-blown sand (Mainguet, 1984). Such an approach requires, of course, that K-feldspar is present in sands of the Grand Erg Oriental. This is not a trivial issue, as some dune sands in the Sahara are mineralogically mature, and absolutely dominated by quartz. To test for the feasibility of doing further studies, we sampled nine dune sand samples in the northeastern margin of the Grand Erg Oriental, in southern Tunisia (three samples from a 10-meter high dune at N33°31'42.4";

E08°47'11.5" and six samples from 4-6-meter high dunes within a 200-meter radius of N33°26'58.2"; E08°51'57.0"). We analyzed these sediments for mineralogy (by XRD) and major-element geochemistry (by WD XRF). Results indicate, in broad agreement with Bellair (1953), that quartz is dominant in these dune sands, with subordinate amounts of calcite and gypsum. In addition, however, all Grand Erg Oriental samples we studied contain measurable amounts of K-feldspar and some even contain plagioclase. Concentrations of K₂O range from 0.7-0.9% and Na₂O contents are 0.13-0.19%, confirming the presence of both species of feldspar inferred from XRD. Thus, the potential for learning more of the provenance of sand in the Grand Erg Oriental is high and could benefit from K/Rb and K/Ba analyses of K-feldspar in these dunes and potential sources.

5.2.4. *Sinai-Negev sand sea, Egypt and Israel*

The Sinai-Negev sand sea (Fig. 19) is a single geomorphic entity, but both names are applied here because the dune body is found in both Egypt (Sinai) and Israel (Negev). As has been the case with the history of studies of the Kelso Dunes, essentially one sediment source alone has been proposed for the Sinai-Negev dune field, namely sand from the Nile Delta (see review in Muhs et al., 2013a). Interestingly, identification of the Nile as a source for the Sinai-Negev erg sands seems to have been both a working assumption as well as an untested hypothesis (Neev et al., 1987; Goring-Morris and Goldberg, 1990; Pye and Tsoar, 2009; Amit et al., 2011; Roskin et al., 2011, 2012). Indeed, Davis et al. (2012) even proposed that the Nile has been a major source for aeolian sediments in Israel for the past ~2.5 million years, based on cosmogenic isotope evidence from quartz. And, in point of fact, the best argument for a Nile Delta source is the simple lack of

755 evidence for other likely sources (Tsoar et al., 2008), although this does not in and of
 756 itself constitute proof.

757 Other possible quartz-rich sources for the Sinai-Negev erg are, however, very limited.
 758 Areas in central Sinai within the Wadi El Arish drainage basin (Fig. 19) have bedrock
 759 dominated by Cretaceous or Eocene rocks. Although these rocks are composed mostly of
 760 carbonate facies, Bartov (1990) reports that sandstone facies are also present in two of the
 761 Cretaceous units. These rocks are situated in the upper drainage basin area of Wadi El
 762 Arish and thus provide a source that is upstream and upwind of much of the Sinai-Negev
 763 dune field. Farther north, in north-central Sinai, Lower Cretaceous sandstones, now part
 764 of what is called the Kurnub Group (Bartov, 1990), were formerly referred to as ‘Nubian
 765 Sandstone.’ These rocks are downwind of many of the dunes in the erg, however, so at
 766 most they constitute a potential source for only part of the erg.

767 Muhs et al. (2013a) conducted mineralogical and geochemical studies of the Sinai-
 768 Negev dune field, sand-sized particles from the Nile Delta, and sand-sized alluvium from
 769 Wadi El Arish. They report that ternary diagrams of Fe_2O_3 - MgO - TiO_2 , a geochemical
 770 proxy for heavy mineral assemblages, show that Nile Delta sands and Wadi El Arish
 771 alluvial sands in Sinai have distinctive compositions, with no overlap. Sinai-Negev dune
 772 sands fall mostly within or close to the field defined by Nile Delta sands, although some
 773 samples fall within the range of Wadi El Arish alluvial sands. Furthermore, most Sinai-
 774 Negev dune sands contain calcite, which is largely absent from Nile Delta sands, but is
 775 abundant in sands from Wadi El Arish. Thus, although the evidence points to the Nile
 776 Delta as the major source of heavy minerals in the Sinai-Negev dune field, Wadi El Arish
 777 has likely contributed most of the calcite to the dunes. An interesting question that arises,

therefore, what the relative contributions of Nile Delta sands and Wadi El Arish sands are to the light, silicate mineral fraction (quartz, plagioclase, K-feldspar) of Sinai-Negev dunes. Geochemical plots of mineralogical maturity ($\text{Log} [\text{Na}_2\text{O}/\text{K}_2\text{O}]$ vs. $\text{Log} [\text{SiO}_2/\text{Al}_2\text{O}_3]$) that measure feldspar depletion relative to quartz show that many Sinai-Negev dunes have compositions that fall within the range of Nile Delta sands, but some fall within the range of Wadi El Arish sands, and some could have had contributions from both sources. Investigations by Muhs et al. (2013a) show that K-feldspar is present in measurable amounts in Nile Delta sands, Wadi El Arish sands, and Sinai-Negev dune sands, and concentrations of K_2O in these sediments, with a couple of exceptions, range from 0.4-1.4%. A worthwhile effort, therefore, would be to analyze both Nile Delta sands and Wadi El Arish sands for K/Rb and K/Ba. If these two sand sources show distinctive K/Rb and K/Ba compositions, then it should be possible to determine the relative contributions of each source to the light, silicate mineral fraction of Sinai-Negev dunes.

5.2.5. Problems in applying the method to certain other dune fields

Although we propose here that K/Rb and K/Ba data from K-feldspars might yield much new significant information about the sources of dunes in many sand seas, we recognize that there are a number of places where such studies would not be feasible. The limiting factor in many dune fields worldwide is simply lack of K-feldspar. Dunes in many of the world's great sand seas are mineralogically mature, dominated by quartz, and of course at least some minimum amount of K-feldspar is required for K/Rb and K/Ba studies. As an example, consider the arid interior of Australia, where much of the landscape is covered with large fields of predominantly linear dunes (Fig. 16).

Mineralogical analyses by both microscopic examination of grains and X-ray diffraction show that only one sample from the Great Sandy Desert had even the slightest trace of feldspar, with all others showing only quartz (Muhs, 2004), confirming earlier findings of Goudie et al. (1993). Geochemical analyses support these findings, as all samples have SiO_2 contents of 95-98% and K_2O contents of only 0.03-0.08%. It is likely, given that quartz was essentially the only sand-sized grain observed, that the small amounts of Al_2O_3 (0.5-1.7%) and K_2O do not represent feldspar but are reflecting the presence of clay mineral coatings that were not completely removed in the pre-treatment process. Thus, we propose that use of U-Pb age spectra of detrital zircons, such as those conducted by Pell et al. (1999, 2000) for the Great Victoria, Simpson, Strzelecki, and Tirari Deserts (Fig. 16) is a more appropriate method for provenance studies of mineralogically mature Australian dunes.

Although we presented data to show that the Grand Erg Oriental has sufficient K-feldspar to make K/Rb and K/Ba studies feasible, this is not true for all dune fields in the Sahara. The Zallaf sand sea, the easternmost arm of the larger Ubari dune field of Libya (McKee and Tibbitts, 1964; Walker, 1979; Breed et al., 1979), is such an example (Fig. 16). Muhs (2004) showed that in this dune field, quartz is the only mineral in some samples and K-feldspar is present only in very small amounts in other samples. The dominance by quartz with only minor amounts of (or no) K-feldspar in sands from the Zallaf sand sea is supported by major element geochemistry, with SiO_2 contents of 97.9-98.5% and K_2O contents of only 0.12-0.17%.

6. Summary and conclusions

823 The Kelso Dunes are unusual among dune fields of the western USA in that there is a
824 wide variety of dune forms occupying a small area. In addition, they are also partially
825 active, which is rare among dune fields in the deserts of the western USA. Over eight
826 decades, there has been much speculation about the origin of the dunes, but the majority
827 of investigators have favored a simple, Mojave-River-dominated source. Nevertheless,
828 with a couple notable exceptions, these studies have not presented mineralogical or
829 geochemical data to ascertain possible dune sources and thus are of a speculative nature.

830 Analysis of the geomorphology of Kelso Dunes suggests the possibility of derivation
831 from more than one source. Many of the dunes have orientations indicating dominantly
832 westerly or northwesterly winds during their formation, consistent with a Mojave River
833 source. On the other hand, other dunes show an orientation indicating southwesterly
834 winds or have a complex nature, with the earlier phase of dune development indicating
835 formation under southwesterly winds.

836 In previous studies that have presented mineralogical data, two report that the Kelso
837 Dunes are quartz-rich. Given the likelihood of short transport distances and the granitic
838 composition of all the surrounding mountain ranges, a dominance of quartz in Kelso
839 Dunes, indicating mineralogical maturity, would be surprising. Mineralogical and
840 geochemical data presented here indicate that the dunes are far less quartz-rich than
841 previous studies have indicated, with relatively high abundances of plagioclase and K-
842 feldspar. The presence of K-feldspar can be used to identify the source or sources of
843 aeolian sand in Kelso Dunes, through K/Rb and K/Ba values in this mineral. These
844 values reflect compositions of K-feldspars from different alluvial sources and ultimately
845 K-feldspars from different source rocks. Although there is some overlap among different

alluvial sources, several of these can be distinguished from one another, allowing some inferences about what fed the Kelso Dunes. Comparison of K/Rb and K/Ba in aeolian sands with alluvial sources indicates that sands in the nine western dune groups have compositions that are indistinguishable from Mojave River alluvium and Budweiser Wash alluvium. Mojave River alluvium would require northwesterly or westerly winds whereas Budweiser Wash alluvium would require southwesterly winds to build dunes. Thus, whereas many previous interpretations have inferred a single Mojave River source, the new results indicate that two sources could have supplied sand to the western dune fields. In contrast, sands from the five eastern dune groups have K/Rb and K/Ba values that indicate significant inputs from alluvial fan deposits of the Providence Mountains. This requires either rare winds from the southeast or, more likely, aeolian reworking of distal Providence Mountain fan sediments by winds from the west, at a rate greater than that from Mojave River or other western source inputs. Overall, the results indicate that even a small dune field can have a complex origin, either from seasonally varying winds or complex fan-dune interaction. Our case study of the Kelso dunes demonstrates the potential of using K/Rb and K/Ba in K-feldspar as an indicator of sand provenance that can be applied to understand sediment sources and transport pathways in many sand seas worldwide.

7. Acknowledgments

This study was supported by the Climate and Land Use Change Research and Development Program of the U.S. Geological Survey. Although field sampling of the dunes predated establishment of Mojave National Preserve, we wish to acknowledge the interest and support of National Park Service personnel in the progress of this study.

Debra Hughson of Mojave National Preserve provided encouragement and valuable feedback on our results. George Kennedy and Jossh Beann assisted with field work and Brian Marshall, Shannon Mahan, and DeAnna Laurel assisted in the laboratory. Tracy Rowland kindly provided the oblique aerial photograph shown in Figure 7. Dr. Nabil Gasmi (Département de Géographie, Faculté des Lettres et des Sciences Humaines de Sousse, Tunisia) took Muhs to dune sample sites in the Grand Erg Oriental (merci beaucoup, Nabil). We appreciate helpful discussions with Mark Sweeney (University of South Dakota) and Eric McDonald (Desert Research Institute). Finally, we thank Jeff Pigati and Elly Brouwers (both USGS), Mark Sweeney, and editor Andy Plater (University of Liverpool) for helpful comments on an earlier version of this paper. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. government.

References

- Amit, R., Enzel, Y., Crouvi, O., Simhai, O., Matmon, A., Porat, N., McDonald, E., Gillespie, A.R., 2011. The role of the Nile in initiating a massive dust influx to the Negev late in the middle Pleistocene. *Geological Society of America Bulletin* 123, 873-889.
- André, J.M., 2014. Floristic diversity and discovery in the California desert. *Fremontia* 42, 3-8.
- Bartov, Y., 1990. Geological photomap of Israel & adjacent areas. Jerusalem, Geological Survey of Israel, scale 1:750,000.
- Bellair, P., 1953. Diagramme minéralogique du Grand Erg Oriental d'El Oued à Ghadamès. *Compte Rendu Sommaire des Séances de la Société Géologique de France* 5-6, 99-101.
- Besler, H., 1980. Die Dünen-Namib: Entstehung und Dynamik eines Ergs. *Stuttgarter Geographische Studien* 96.

- 892 Besler, H., 1984. The development of the Namib dune field according to sedimentological and
 893 geomorphological evidence. In: Vogel, J.C., ed., Late Cainozoic Palaeoclimates of the Southern
 894 Hemisphere. Rotterdam: Balkema.
- 895 Besler, H., 1996. The Tsondab Sandstone in Namibia and its significance for the Namib Erg. South
 896 African Journal of Geology 99, 77–87.
- 897 Besler, H., Marker, M.E., 1979. Namib sandstone: A distinct lithological unit. Transactions of the
 898 Geological Society of South Africa 82, 155-160.
- 899 Bortugno, E.J., Spittler, T.E., 1986. Geologic map of the San Bernardino quadrangle, California. State of
 900 California, Division of Mines and Geology, scale, 1:250,000.
- 901 Bowler, J.M., Wyrwoll, K.-H., Lu, Y., 2001. Variations of the northwest Australian summer monsoon over
 902 the last 300,000 years: the paleohydrological record of the Gregory (Mulan) Lakes System.
 903 Quaternary International 83-85, 63-80.
- 904 Breed, C.S., Fryberger, S.C., Andrews, S., McCauley, C., Lennartz, F., Gebel, D., Horstman, K., 1979.
 905 Regional studies of sand seas using Landsat (ERTS) imagery. In: McKee, E.D. (Ed.), A Study of
 906 Global Sand Seas. U.S. Geological Survey Professional Paper 1052, pp. 305-397.
- 907 Clarke, M.L., 1994. Infra-red stimulated luminescence ages from aeolian sand and alluvial fan deposits
 908 from the eastern Mojave Desert, California. Quaternary Science Reviews (Quaternary
 909 Geochronology) 13, 533-538.
- 910 Cooke, R., Warren, A., and Goudie, A., 1993. Desert Geomorphology. UCL Press Limited, London, 526
 911 pp.
- 912 Crouvi, O., Amit, R., Porat, N., Gillespie, A.R., McDonald, E.V., Enzel, Y., 2009. Significance of primary
 913 hilltop loess in reconstructing dust chronology, accretion rates, and sources: An example from the
 914 Negev Desert, Israel. Journal of Geophysical Research 114, F02017, doi: 10.1029/2008JF001083.
- 915 Davis, M., Matmon, A., Rood, D.H., Avnaim-Katav, S., 2012. Constant cosmogenic nuclide
 916 concentrations in sand supplied from the Nile River over the past 2.5 m.y. Geology 40, 359-362.

- 917 Edwards, S.R., 1993. Luminescence dating of sand from the Kelso Dunes, California. In: Pye, K. (ed.),
 918 The Dynamics and Environmental Context of Aeolian Sedimentary Systems. Geological Society
 919 Special Publication No. 72, 59-68.
- 920 Enzel, Y., Wells, S.G., Lancaster, N., 2003. Late Pleistocene lakes along the Mojave River, southeast
 921 California. Geological Society of America Special Paper 368, 61-77.
- 922 Enzel, Y., Amit, R., Crouvi, O., Porat, N., 2010. Abrasion-derived sediments under intensified winds at the
 923 latest Pleistocene leading edge of the advancing Sinai-Negev erg. Quaternary Research 74, 121-131.
- 924 Fryberger, S.G., Ahlbrandt, T.S., 1979. Mechanisms for the formation of eolian sand seas. Zeitschrift für
 925 Geomorphologie 23, 440-460.
- 926 Fryberger, S.G., Dean, G., 1979. Dune forms and wind regime. In: McKee, E.D. (Ed.), A Study of Global
 927 Sand Seas. U.S. Geological Survey Professional Paper 1052, pp. 137-169.
- 928 Garzanti, E., Andò, S., Vezzoli, G., Lustrino, M., Boni, M., Vermeesch, P., 2012. Petrology of the Namib
 929 Sand Sea: Long-distance transport and compositional variability in the wind-displaced Orange
 930 Delta. Earth-Science Reviews 112, 173-189.
- 931 Gevers, T.W., 1936. The morphology of western Damaraland and the adjoining Namib Desert. South
 932 African Geographical Journal 19, 61-79.
- 933 Goring-Morris, A.N., Goldberg, P., 1990. Late Quaternary dune incursions in the southern Levant:
 934 Archaeology, chronology and palaeoenvironments. Quaternary International 5, 115-137.
- 935 Goudie, A.S., 2002. Great warm deserts of the world: Landscapes and evolution. Oxford, Oxford
 936 University Press, 480 pp.
- 937 Goudie, A.S., Stokes, S., Livingstone, I., Bailiff, I.K., Allison, R.J., 1993. Post-depositional modification
 938 of the linear sand ridges of the West Kimberley area of north-west Australia. The Geographical
 939 Journal 159: 306-317.

- 940 Grove, A.T., Warren, A., 1968. Quaternary landforms and climate on the south side of the Sahara.
941 Geographical Journal 134, 194-208.
- 942 Hallam, C.D., 1964. The geology of the coastal diamond deposits of southern Africa. In: Houghton, S.E.,
943 ed., The Geology of Some Ore Deposits in South Africa. Johannesburg: Geological Society of
944 South Africa.
- 945 Heier, K.S., and Adams, J.A.S., 1964. The geochemistry of the alkali metals. Physics and Chemistry of
946 the Earth 5, 253-381.
- 947 Helm, P.J., Breed, C.S., 1999. Instrumented field studies of sediment transport by wind. U.S. Geological
948 Survey Professional Paper 1598-B, 29-51.
- 949 Hereford, R., Webb, R.H., Longpre, C.I., 2004. Precipitation history of the Mojave Desert region, 1893-
950 2001. U.S. Geological Survey Fact Sheet FS 117-03. Available at:
951 <http://geopubs.wr.usgs.gov/fact-sheet/fs117-03/>.
- 952 Kocurek, G., and Lancaster, N., 1999. Aeolian system sediment state: theory and Mojave Desert Kelso
953 dune field example. Sedimentology 46, 505-515.
- 954 Lange, I.M., Reynolds, R.C., Lyons, J.B., 1966. K/Rb ratios in coexisting K-feldspars and biotites from
955 some New England granites and metasediments. Chemical Geology 1, 317-322.
- 956 Lancaster, N., 1988. Development of linear dunes in the southwestern Kalahari, southern Africa. Journal
957 of Arid Environments 14, 233-244.
- 958 Lancaster, N., 1989. The Namib Sand Sea: Dune forms, processes and sediments. Rotterdam, A.A.
959 Balkema, 180 pp.
- 960 Lancaster, N., 1993. Kelso Dunes. National Geographic Research & Exploration 9, 444-459.
- 961 Lancaster, N., 1994. Controls on aeolian activity: some new perspectives from the Kelso Dunes, Mojave
962 Desert, California. Journal of Arid Environments 27, 113-125.

- 963 Lancaster, N., 1995. *Geomorphology of Desert Dunes*. Routledge, London, 290 pp.
- 964 Lancaster, N., Ollier, C.D., 1983. Sources of sand for the Namib sand sea. *Zeitschrift für Geomorphologie*
 965 Suppl. 45, 71-83.
- 966 Lancaster, N., Tchakerian, V.P., 2003. Late Quaternary eolian dynamics, Mojave Desert, California.
 967 Geological Society of America Special Paper 368, 231-249.
- 968 Lancaster, N., Kocurek, G., Singhvi, A. Pandey, V., Deynoux, M., Ghienne, J.-F., Lo, K., 2002. Late
 969 Pleistocene and Holocene dune activity and wind regimes in the western Sahara Desert of
 970 Mauritania. *Geology* 30, 991-994.
- 971 Livingstone, I., Warren, A., 1996. *Aeolian geomorphology: An introduction*. Longman, London, 211 pp.
- 972 Logan, R.F., 1960. *The Namib Desert, South West Africa*. Washington, D.C., National Academy of
 973 Sciences.
- 974 Lorenz, R.D., Zimbelman, J.R., 2014. *Dune worlds: How windblown sand shapes planetary landscapes*.
 975 Springer, 308 pp.
- 976 Maman, S., Blumberg, D.G., Tsoar, H., Mamedov, B., Porat, N., 2011. The Central Asian ergs: A study by
 977 remote sensing and geographic information systems. *Aeolian Research* 3, 353-366.
- 978 Mainguet, M., 1978. The influence of trade winds, local air-masses and topographic obstacles on the
 979 aeolian movement of sand particles and the origin and distribution of dunes and ergs in the Sahara
 980 and Australia. *Geoforum* 9, 17-28.
- 981 Mainguet, M., 1984. Space observation of Saharan aeolian dynamics. In: El-Baz, F., ed., *Deserts and arid*
 982 *lands*. The Hague, Martinus Nijhoff Publishers, pp. 59-77.
- 983 Mason, B., Moore, C.B., 1982. *Principles of Geochemistry*. Wiley, New York, 344 pp.
- 984 McDonald, E.V., McFadden, L.D., Wells, S.G., 2003. Regional response of alluvial fans to the
 985 Pleistocene-Holocene climatic transition, Mojave Desert, California. Geological Society of
 986 America Special Paper 368, 189-205.

- 987 McKee, E.D., Tibbitts, G.C., Jr., 1964. Primary structures of a seif dune and associated deposits in Libya.
988 *Journal of Sedimentary Petrology* 34, 5-17.
- 989 Miller, D.M., Miller, R.J., Nielson, J.E., Wilshire, H.G., Howard, K.A., Stone, P., 1991. Preliminary
990 geologic map of the East Mojave National Scenic Area, California. U.S. Geological Survey Open-
991 File Report 91-435, 8 pp, scale 1:100,000.
- 992 Muhs, D.R., 2004. Mineralogical maturity in dune fields of North America, Africa and Australia.
993 *Geomorphology* 59, 247-269.
- 994 Muhs, D.R., Zárate, M., 2001. Late Quaternary eolian records of the Americas and their paleoclimatic
995 significance. In: Markgraf, V. (Ed.), *Interhemispheric climate linkages*. Academic Press, San
996 Diego, pp. 183-216.
- 997 Muhs, D.R., Bush, C.A., Cowherd, S.D., Mahan, S., 1995. Geomorphic and geochemical evidence for the
998 source of sand in the Algodones dunes, Colorado Desert, southeastern California. In: Tchakerian,
999 V.P. (Ed.), *Desert Aeolian Processes*. Chapman and Hall, London, pp. 37-74.
- 1000 Muhs, D.R., Reynolds, R., Been, J., Skipp, G., 2003. Eolian sand transport pathways in the southwestern
1001 United States: Importance of the Colorado River and local sources. *Quaternary International* 104,
1002 3-18.
- 1003 Muhs, D.R., Bettis, E.A. III, Aleinikoff, J., McGeehin, J.P., Beann, J., Skipp, G., Marshall, B.D., Roberts,
1004 H.M., Johnson, W.C., Benton, R., 2008. Origin and paleoclimatic significance of late Quaternary
1005 loess in Nebraska: Evidence from stratigraphy, chronology, sedimentology, and geochemistry.
1006 *Geological Society of America Bulletin* 120, 1378-1407.
- 1007 Muhs, D.R., Roskin, J., Tsoar, H., Skipp, G., Budahn, J.R., Sneh, A., Porat, N., Stanley, J.-D., Katra, I.,
1008 Blumberg, D.G., 2013a. Origin of the Sinai-Negev erg, Egypt and Israel: mineralogical and
1009 geochemical evidence for the importance of the Nile and sea level history. *Quaternary Science*
1010 *Reviews* 69, 28-48.
- 1011 Muhs, D.R., Bettis, E.A. III, Roberts, H.M., Harlan, S., Paces, J.B., Reynolds, R., 2013b. Chronology and
1012 provenance of last-glacial (Peoria) loess in western Iowa and paleoclimatic implications.
1013 *Quaternary Research* 80, 468-481.

- 1014 Neev, D., Bakler, N., Emery, K.O., 1987. Mediterranean coasts of Israel and Sinai. New York, Taylor &
1015 Francis, 130 p.
- 1016 Nockholds, S.R., 1954. Average chemical compositions of some igneous rocks. Geological Society of
1017 America Bulletin 65, 1007-1032.
- 1018 Norris, R.M., Webb, R.W., 1976. Geology of California. John Wiley & Sons, New York, 379 pp.
- 1019 Paisley, E.C.I., Lancaster, N., Gaddis, L.R., Greeley, R., 1991. Discrimination of active and inactive sand
1020 from remote sensing: Kelso Dunes, Mojave Desert, California. Remote Sensing of the
1021 Environment 37, 153-166.
- 1022 Pavlik, B.M. 1989. Phytogeography of sand dunes in the Great Basin and Mojave deserts. Journal of
1023 Biogeography 16, 227-238.
- 1024 Pell, S.D., Chivas, A.R., Williams, I.S., 1999. Great Victoria Desert: development and sand provenance.
1025 Australian Journal of Earth Sciences 46, 289-299.
- 1026 Pell, S.D., Chivas, A.R., Williams, I.S., 2000. The Simpson, Strzelecki and Tirari Deserts: development
1027 and sand provenance. Sedimentary Geology 130, 107-130.
- 1028 Pye, K., Tsoar, H., 2009. Aeolian sand and sand dunes. Berlin: Springer-Verlag, 458 pp.
- 1029 Ramsey, M.S., Christensen, P.R., Lancaster, N., Howard, D.A., 1999. Identification of sand sources and
1030 transport pathways at the Kelso Dunes, California, using thermal infrared remote sensing.
1031 Geological Society of America Bulletin 111, 646-662.
- 1032 Reheis, M.C., Redwine, J.L., 2008. Lake Manix shorelines and Afton Canyon terraces: Implications for
1033 incision of Afton Canyon. Geological Society of America Special Paper 439, 227-259.
- 1034 Reheis, M.C., Bright, J., Lund, S.P., Miller, D.M., Skipp, G., Fleck, R.J., 2012. A half-million-year record
1035 of paleoclimate from the Lake Manix Core, Mojave Desert, California. Palaeogeography,
1036 Palaeoclimatology, Palaeoecology 365-366, 11-37.
- 1037 Rogers, J., 1977. Sedimentation on the continental margins off the Orange River and the Namib Desert.
1038 Joint Geological Survey/University of Cape Town Marine Geoscience Group Bulletin 7.
- 1039 Rogers, J., 1979. Dispersal of sediment from the Orange River along the Namib desert coast. South
1040 African Journal of Science 75, 567.

- 1041 Roskin, J., Porat, N., Tsoar, H., Blumberg, D.G., Zander, A.M., 2011. Age, origin and climatic controls on
1042 vegetated linear dunes in the northwestern Negev Desert (Israel). *Quaternary Science Reviews* 30,
1043 1649-1674.
- 1044 Roskin, J., Blumberg, D.G., Porat, N., Tsoar, H., Rozenstein, O., 2012. Do dune sands redden with age?
1045 The case of the northwestern Negev dunefield, Israel. *Aeolian Research* 5, 63-75.
- 1046 Scheidt, S., Lancaster, N., Ramsey, M., 2011. Eolian dynamics and sediment mixing in the Gran Desierto,
1047 Mexico, determined from thermal infrared spectroscopy and remote-sensing data. *Geological*
1048 *Society of America Bulletin* 123, 1628-1644.
- 1049 Schoenherr, A.A., 1992. A natural history of California. Berkeley, University of California Press, 772 pp.
- 1050 Sharp, R.P., 1966. Kelso Dunes, Mojave Desert, California. *Geological Society of America Bulletin* 77,
1051 1045-1074.
- 1052 Sharp, R.P., 1978. The Kelso dune complex. In: Greeley, R., Womar, M.B., Papson, R.P., Spudis, P.D.
1053 (eds.), *Aeolian features of southern California: A comparative planetary geology guidebook*.
1054 NASA Office of Planetary Geology, Washington, D.C., pp. 54-63.
- 1055 Soller, D.R., Reheis, M.C., 2004. Surficial materials in the conterminous United States. U.S. Geological
1056 Survey Open-File Report OFR-03-275.
- 1057 Spaulding, W.G., 1995. Environmental change, ecosystem responses, and the Late Quaternary
1058 development of the Mojave Desert. In: *Late Quaternary environments and deep history: A tribute*
1059 *to Paul S. Martin (D.W. Steadman and J.L. Mead, eds.)*, pp. 139-164. The Mammoth Site of Hot
1060 Springs, South Dakota.
- 1061 Stone, A.E.C., 2013. Age and dynamics of the Namib Sand Sea: A review of chronological evidence and
1062 possible landscape development models. *Journal of African Earth Sciences* 82, 70-87.
- 1063 Streckeisen, A.L., 1976. To each plutonic rock its proper name. *Earth-Science Reviews* 12, 1-34.

- 1064 Sun, J., Muhs, D.R., 2007. Dune fields: Mid-latitudes. In: Elias, S. (ed.), The Encyclopedia of Quaternary
1065 Sciences. Amsterdam, Elsevier, p. 607-626.
- 1066 Sweeney, M.R. McDonald, E.V., Markley, C.E., 2013. Alluvial sediment or playas: What is the dominant
1067 source of sand and silt in desert soil vesicular A horizons, southwest USA. Journal of
1068 Geophysical Research: Earth Surface 118, 257-275.
- 1069 Sweeney, M.R., McDonald, E.V., Hanson, P., Chabela, L.P., 2015. Sand dunes exert control on alluvial
1070 base level and soil formation, eastern Mojave Desert, USA. Geological Society of America
1071 Abstracts with Programs 47 (7), 737.
- 1072 Thomas, D.S.G., Shaw, P.A., 1991. The Kalahari environment. Cambridge, Cambridge University Press.
- 1073 Thompson, D.G., 1929. The Mohave Desert region California: A geographic, geologic, and hydrologic
1074 reconnaissance. U.S. Geological Survey Water-Supply Paper 578, 759 pp.
- 1075 Thorne, R.F., Prigge, B.A., Henrickson, J., 1981. A flora of the higher ranges and the Kelso dunes of the
1076 Eastern Mojave Desert in California. *Aliso* 10, 71–186.
- 1077 Tsoar, H., Blumberg, D.G., Wenkart, R., 2008. Formation and geomorphology of the north-western Negev
1078 sand dunes. In: Breckle, S.-W., Yair, A., and Veste, M., eds., *Arid Dune Ecosystems: The*
1079 *Nizzana Sands in the Negev Desert*. Berlin, Springer, 25-48.
- 1080 Vermeesch, P., Fenton, C.R., Kober, F., Wiggs, G.F.S., Bristow, C.S., Xu, S., 2010. Sand residence times
1081 of one million years in the Namib Sand Sea from cosmogenic isotopes. *Nature Geoscience*,
1082 doi:10.1038/NGEO985.
- 1083 Walker, T.R., 1979. Red color in dune sand. U.S. Geological Survey Professional Paper 1052D, 61-81.
- 1084 Warren, A., 2013. *Dunes*. John Wiley & Sons, Ltd., Chichester, 219 pp.
- 1085 Wedepohl, K.H., 1969. *The handbook of geochemistry, volume 1*. Berlin, Springer-Verlag, 442 pp.

- 1086 Wells, S.G., Brown, W.J., Enzel, Y., Anderson, R.Y., McFadden, L.D., 2003. Late Quaternary geology
1087 and paleohydrology of pluvial Lake Mojave, southern California. Geological Society of America
1088 Special Paper 368, 79-114.
- 1089 White, K., Walden, J., Gurney, S.D., 2007. Spectral properties, iron oxide content and provenance of
1090 Namib dune sands. Geomorphology 86, 219-229.
- 1091 Wilson, I.G., 1973. Ergs. Sedimentary Geology 10, 77-106.
- 1092 Wintle, A.G., Lancaster, N., and Edwards, S.R., 1994. Infrared stimulated luminescence dating of late-
1093 Holocene aeolian sands in the Mojave Desert, California, USA. The Holocene 4, 74-78.
- 1094 Yeend, W., Dohrenwend, J.C., Smith, R.S.U., Goldfarb, R., Simpson, R.W., Jr., Munts, S.R., 1984.
1095 Mineral resources and mineral resource potential of the Kelso Dunes Wilderness Study Area
1096 (CDCA-250), San Bernardino County, California. U.S. Geological Survey Open-File Report 84-
1097 647, 19 pp.

1098 **Figure captions**

1099 **FIGURE 1:** Map showing the distribution of aeolian sand (brown shades – online
1100 version) in arid and semiarid regions of the southwestern United States and
1101 northern Mexico and modern sand transport directions (resultant drift directions
1102 and resultant drift potential, calculated using methods of Fryberger and Dean
1103 [1979]). Aeolian sand distribution is from compilation in Muhs and Zárate (2001)
1104 and references therein; see also Soller and Reheis (2004). Abbreviations for wind
1105 direction localities, which approximate the station locations: D, Daggett; Cl,
1106 China Lake; TP, Twenty-nine Palms; EC, El Centro; I, Indio; B, Blythe; YF,
1107 Yucca Flat; LV, Las Vegas; GS, Gold Spring; F, Farmington; G, Gallup; H,
1108 Holloman Air Force Base; EP, El Paso. Resultant annual drift potentials

1109 calculated by the authors except for Gold Spring, which is from Helm and Breed
1110 (1999) and Holloman Air Force Base, which is from Fryberger and Dean (1979).

1111 FIGURE 2. Map showing a part of eastern California and adjacent Nevada, the location
1112 of Mojave National Preserve (brown shading – online version), major towns and
1113 highways, and location of the Devils Playground dune field and Kelso Dunes.

1114 FIGURE 3. Landsat ETM+ image (Path 39; Row 36), 20 May 2006, band 1, showing the
1115 Kelso Dunes and surrounding area. Numbered features indicate hypothesized
1116 source sediments; PM, Providence Mountains; GM, Granite Mountains.

1117 FIGURE 4: Wind roses for four representative months of the year from the Mojave
1118 ‘Sink’ area (see Fig. 3 for location); period of record, 1988-2009. Only shown are
1119 frequencies of winds above the threshold velocity for medium sand (~5 m/s) when
1120 considered at an instrument measurement height of 10 m. Note scale change for
1121 the month of May. Graphs compiled from data from the Western Regional
1122 Climate Center of the Desert Research Institute, Reno, Nevada USA
1123 (<http://www.wrcc.dri.edu>).

1124 FIGURE 5. Enlargement of a portion of the Landsat ETM+ image shown in Fig. 3,
1125 showing aeolian landform groups (Roman numerals) in the Kelso Dunes
1126 identified by Lancaster (1993, 1994) on the basis of geomorphology.

1127 FIGURE 6: U.S. Geological Survey aerial photograph (National Aerial Photography
1128 Program or NAPP, acquired 24 July 1989) of a portion of the Kelso Dunes area;
1129 Roman numerals designate aeolian landform groups identified by Lancaster
1130 (1993, 1994; see also Lancaster and Tchakerian, 2003). Cottonwood Wash, in the

1131 center of the photograph, divides what is designated herein as ‘western dune
1132 groups’ and ‘eastern dune groups.’

1133 FIGURE 7: Oblique aerial photograph (acquired at an altitude of ~10,000 m), looking
1134 south, of the Kelso dunes and adjacent mountain ranges and drainages. Roman
1135 numerals designate aeolian landform groups identified by Lancaster (1993, 1994;
1136 see also Lancaster and Tchakerian, 2003). Photo courtesy of Tracy Rowland.

1137 FIGURE 8: Ground photographs of the Kelso Dunes, showing some of the major dune
1138 groups shown in Figs. 5-7 and surrounding mountain ranges. In photograph (c),
1139 red arrows point to heavy mineral concentrations on dune crests. Photos by D.
1140 Muhs.

1141 FIGURE 9: Ternary plots of percentages of quartz, K-feldspar, and plagioclase
1142 (calculated as a sum of 100% for these three mineral species) for the Kelso
1143 Dunes, shown in all four diagrams as red squares. See text for methods of
1144 estimating mineral content percentages. Shown for comparison are ranges of
1145 compositions for these three minerals for (a) the Algodones Dunes, California
1146 (mineralogical data from this study; samples from Muhs et al., 1995); (b) San
1147 Nicolas Island, California dunes and the Parker Dunes, Arizona; (c) the
1148 Cadiz/Danby Dunes, California (mineralogical data from this study; samples from
1149 Muhs et al., 2003); and (d) the Soynota dunes, Sonora, Mexico (mineralogical
1150 data and samples from this study, but see Scheidt et al., 2011 for additional data).

1151 FIGURE 10: Plots showing weight-percent concentrations of K₂O (representing mostly
1152 K-feldspar) and CaO (representing mostly plagioclase possibly with minor

1153 amphibole and calcite) from the Kelso Dunes, shown as red squares in all graphs.
 1154 (a) Concentrations of K_2O and CaO in Kelso Dunes compared to ranges in
 1155 granites and granodiorites (data from Nockolds, 1954); and (b), (c), (d)
 1156 Concentrations of K_2O and CaO in Kelso Dunes compared to similar data for the
 1157 Algodones Dunes, California (data from Muhs et al., 1995), the Parker Dunes,
 1158 Arizona, the Cadiz/Danby Dunes, California, and the Dale Lake sand sheet,
 1159 California (data from Muhs et al., 2003).

1160 FIGURE 11: Plots of concentrations of Ti vs. Ce, Y, and Nb, and Zr vs. Ce and Y in the
 1161 Kelso dunes, all geochemical indicators of the heavy mineral suite.

1162 FIGURE 12: Plots of K/Ba vs. K/Rb, representing mostly K-feldspar compositions, for
 1163 aeolian sands from Devils Playground (Fig. 3), shown as solid green circles,
 1164 compared to range of compositions for these element ratios for hypothesized
 1165 source sediments, shown in Fig. 3.

1166 FIGURE 13: Plots of K/Ba vs. K/Rb, representing mostly K-feldspar compositions, for
 1167 dune sands from the western groups of Kelso Dunes (units VI through XIV on
 1168 Figs. 5-7), shown as open black circles, compared to range of compositions for
 1169 these element ratios for hypothesized source sediments, shown in Fig. 3.

1170 FIGURE 14: Plots of K/Ba vs. K/Rb, representing mostly K-feldspar compositions, for
 1171 dune sands from the eastern groups of Kelso Dunes (units I through V shown on
 1172 Figs. 5-7), shown as open blue circles, compared to range of compositions for
 1173 these element ratios for hypothesized source sediments, shown in Fig. 3.

1174 FIGURE 15: Alternative interpretations of the origin of the eastern dunes in the Kelso
 1175 Dunes region based on geochemical data presented in Figs. 12-14. Blue arrows in
 1176 both (a) and (b) (see online version) represent sand-transporting winds, as shown
 1177 in Fig. 4. In interpretation (a), western dunes are derived from the Mojave River
 1178 by western winds (year-round) and Granite Mountain fans and Budweiser Wash
 1179 sediments by southwesterly winds (year-round). Eastern dunes are derived from
 1180 Providence Mountain fan sediments from southeasterly winds (summer only). In
 1181 interpretation (b), western dunes are derived as in (a), but eastern dunes are
 1182 derived from distal Providence Mountain fan sediments deposited close to Kelso
 1183 Wash and then entrained by winds from the west and blown back up onto
 1184 proximal Providence Mountains fan sediments. Images in both (a) and (b) are
 1185 from Landsat ETM+ (Path 39; Row 36), 20 May 2006, band 1, as in Fig. 3.

1186 FIGURE 16: Map showing the distribution of aeolian sand, whether active or stabilized,
 1187 in Africa, Asia, and Australia. Sources of information: northern Africa: Wilson
 1188 (1973); southern Africa: Lancaster (1988, 1989) and Thomas and Shaw (1991);
 1189 Sinai-Negev: Muhs et al. (2013a); Arabian Peninsula and India: Wilson (1973);
 1190 Central Asia: Maman et al. (2011); China: Sun and Muhs (2007) and sources
 1191 therein; Australia: Bowler et al. (2001).

1192 FIGURE 17: (a) Moderate Resolution Imaging Spectrometer (MODIS) natural-color
 1193 image from NASA's Terra satellite, acquired 14 January 2015, showing mostly
 1194 active dune fields (Azefal, Akchar, and Aoukar) in Mauritania (north of the
 1195 Senegal River) and the stabilized Ferlo-Cayor dune field in Senegal (south of the
 1196 Senegal River). Blue arrows show resultant drift directions (RDDs) for aeolian

1197 sand, from Fryberger and Ahlbrandt (1979) and Breed et al. (1979). Image
 1198 courtesy of Jeff Schmaltz, LANCE MODIS Rapid Response Team at NASA
 1199 Goddard Space Flight Center. (b) Ground photo (by N. Lancaster) of active linear
 1200 dune in the Azefal sand sea of Mauritania. (c) Ground photo (by D. Muhs) of
 1201 stable dune in the Ferlo-Cayor dune field, just north of Dakar, Senegal.

1202 FIGURE 18: (a) MODIS natural-color image from NASA's Terra satellite, acquired on
 1203 26 June 2011, showing the Namib Sand Sea of Namibia, southwestern Africa, and
 1204 rivers to the north and south. Light blue arrows show RDDs, from Breed et al.
 1205 (1979) and Lancaster (1989). Dark blue arrows mark drainages from possible
 1206 bedrock sources of sand adjacent to the Great Escarpment. Image courtesy of
 1207 MODIS Rapid Response Team at NASA Goddard Space Flight Center (b)
 1208 Ground photo of Namib dunes at their southernmost occurrence, near possible
 1209 source areas, Elizabeth Bay area, just south of Luderitz (c) Ground photo of dunes
 1210 in the Namib Sand Sea farther north in the main body of the dune field, where
 1211 sand is redder (both ground photos by N. Lancaster).

1212 FIGURE 19: (a) False-color Landsat ETM+ mosaic of parts of Algeria, Libya, and
 1213 Tunisia showing the Grand Erg Oriental. Mosaic constructed from individual
 1214 Landsat ETM+ images acquired from 1999-11-11 to 2001-06-97; image is N-32-
 1215 30, courtesy of the University of Maryland. Blue arrows (see online version)
 1216 show RDDs, from Fryberger and Ahlbrandt (1979) and Breed et al. (1979). (b)
 1217 Ground photograph (by D. Muhs) of active dunes on the northeastern margin of
 1218 the Grand Erg Oriental (N33°26'58.2"; E08°51'57.0"), southeast of Chott El
 1219 Jerid.

1220 FIGURE 20: (a) MODIS image of the Sinai-Negev dune field of Egypt and Israel,
1221 acquired from the NASA Aqua satellite on 6 February 2003. Also shown are
1222 possible source sediment areas for this dune field, the Nile Delta and the Wadi El
1223 Arish drainage basin. MODIS image courtesy of Jeff Schmaltz, MODIS Rapid
1224 Response Team, NASA Goddard Space Flight Center. Blue arrows indicate
1225 RDDs (from Muhs et al., 2013a). (b) Ground photograph (by D. Muhs) showing
1226 dunes of the Sinai-Negev dune field along the Egypt-Israel border, west of
1227 Halamish, Israel.
1228

Table 1. Geomorphology of the Kelso Dunes

Unit from Lancaster (1993, 1994)	Geography	Condition	Geomorphology	Most recent paleowind	Samples for XRF (n)	Samples for XRD (n)
I	Eastern	Stabilized	Degraded parabolic dunes and sand sheets	W?	9	5
II	Eastern	Stabilized	Linear dunes with superimposed parabolic dunes	WNW	9	6
III	Eastern	Active; stabilized	Linear dunes with superimposed parabolic dunes	W	5	4
IV	Eastern	Stabilized	Degraded linear dunes and sand sheets	SW?	10	8
V	Eastern	Stabilized	Degraded parabolic dunes	NW, W	5	1
VI	Western	Active	Linear dune with superimposed parabolic dunes, barchanoid ridges, and star dunes	SW;W or WNW	25	19
VII	Western	Stabilized	Degraded parabolic dunes	NW?	7	6
VIII	Western	Active	Linear dune with superimposed barchanoid ridges, star dunes and parabolic dunes	SW; W, WNW	0	0
IX	Western	Stabilized	Degraded linear dunes with superimposed, degraded parabolic dunes	W	9	4
X	Western	Active	Linear dune with superimposed barchanoid ridge, star, and parabolic dunes	SW; W, WNW	0	0
XI	Western	Stabilized	Linear dunes grading eastward into parabolic dunes	WSW	9	9
XII	Western	Stabilized	Parabolic dunes	W	3	1
XIII	Western	Stabilized	Linear dunes grading southeastward into parabolic dunes	NW	9	8
XIV	Western	Stabilized	Parabolic dunes	W	4	3
Devils Playground	Western	Stabilized	Sand sheets, streaks	W	9	0

Figure 1

Figure 2

Figure 4

FEBRUARY

MAY

JULY

OCTOBER

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9

Figure 10

(a)

(b)

(c)

(d)

Figure 11

Figure 12

(a)

(b)

(c)

(d)

Figure 13

(a)

(b)

(c)

(d)

Figure 14

Figure 15

Figure 16

Figure 18

Figure 19

Figure 20

NORTH
→