Effect of bacteria isolates in Powdery mildew control in flowering dogwoods (*Cornus florida* L.)

M.T. Mmbaga^{ac}, F.A. Mrema^b, L. Mackasmie^{ac} and E. Rotich^a

^aTennessee State University, College of Agriculture, Human and Natural Sciences, Department of Agricultural and Environmental Sciences, Nashville TN, ^bOtis L. Floyd Nursery Research Center, McMinnville, TN 37110; ^bAlcorn State University, Research and Applied Sciences, MS 39096-7500

Abstract

Five bacterial isolates collected from dogwood leaves were evaluated for powdery mildew control in shadehouse and greenhouse environments by using foliar sprays and/or root drenching. Two isolates displayed superior bioactivity and suppressed powdery mildew similar to conventional fungicide thiophanate methyl (Cleary's 3336F®). The two bacteria disrupted powdery mildew spore germination and ruptured spore membranes causing spore lyses. Bacterial filtrates without bacterial cells were also effective in suppressing powdery mildew and disrupting spore germination and suggested the involvement of secondary metabolites. The two biocontrol agents (BCAs) colonized roots endophytically and promoted plant growth.

Keywords: Ornamentals, fungicides, *Oidium* spp.; *Erysiphe* (Sect. *Microsphaera*) *pulchra*; biological control.

1. Introduction

Powdery mildew caused by *Erysiphe* (Sect. *Microsphaera*) *pulchra* (Cook & Peck, Braun & Takamatsu) is one of the most economically important diseases affecting dogwood production in the southeastern USA (Leigh et al., 1998; McRitchie, 1994. Mmbaga, 1998; Mmbaga, 2000; Windham, 1994). The disease causes stunted growth, premature defoliation, reduced esthetic value and overall decline of infected trees (Chartfield and Rose, 1996; Mmbaga, 1998; Mmbaga, 2000; Daughtrey and Hagan, 2001; Smith, 1999). Current management of this disease is primarily with fungicides (Windham, 1994; Mmbaga, 2000; Mmbaga and Sheng, 1999), but this has increased production costs (Li et al., 2009) and caused environmental concerns (Mmbaga and Sheng, 1999). Many small nurseries have ceased or terminated dogwood production because profit margins were not sufficient to

cover fungicide and labor costs associated with routine fungicide applications every two weeks from May to October (**Li et al 2010**). In addition to cost, chemical fungicides also harm non-target organisms that play a role in protecting plants against pathogens (**Kiss, 2003; Elad, 2000; Elad et al., 1996**) and pose health hazards to humans, other animals including wildlife as a result of accidental exposures to the toxic chemicals. Biological agents for controlling powdery mildew will offer alternative disease management options that are presumably safer, sustainable, and less damaging to the environment than traditional chemical fungicides (**Elad, 2000; Elad et al., 1996; Kiss, 2003; Mmbaga et al., 2008; Mmbaga and Sauvé, 2009**).

Bacteria belonging to the genera *Bacillus*, *Agrobacterium*, *Pseudomonas* and *Streptomyces* have previously been used to control plant diseases (**Gardner and Fravel, 2002**). A combination of biological control and other user-friendly methods such as resistant cultivars (**Hartman et al., 2003**) and cultural practices that reduce pathogen proliferation may substantially reduce the amount of fungicide used in powdery mildew control in nursery production of dogwood. Filamentous fungi, bacteria and yeast isolated from dogwood foliage in the wild have shown potential in controlling dogwood powdery mildew (**Mmbaga et al., 2008; Mmbaga and Sauvé, 2009**). Although some biological agents are often isolated from plant leaves as epiphytes, some have been reported to have an endophytic phase and interact with their host in symbiotic and other types of relationships (**Nejad and Johnson, 2000; Mmbaga and Sauvé, 2009; Pal et al 2006**). The objective of this study was to confirm the efficacy of selected bacterial isolates in powdery mildew disease control, assess their ability to colonize plants through the roots and their potential mechanism of action.

2. Materials and methods.

2.1 Efficacy of bacterial isolates in powdery mildew disease control

54 Four sets of experiments were conducted to (1) evaluate five isolates previously isolated from the
55 wild (**Mmbaga et al., 2008**) and stored for two years at -80°C, (2) evaluate isolates (B17A and B17B)
56 that showed superior bioactivity, (3) evaluate method of application as foliar sprays and root drenching,
57 and (4) assess the potential mechanism of action on powdery mildew spore germination. All
58 experiments were repeated ones.

59 ***2.1.1. Bacterial isolates inoculum preparation, plant material for biological control of powdery***
60 ***mildew.***

61 All bacterial isolates used in this study were previously isolated from dogwood leaves collected
62 from two natural and uncultivated forest habitats (**Mmbaga et al., 2008**). These isolates had been stored
63 at -80°C in 15% aqueous glycerol and were revived and grown in nutrient agar (NA) and visually
64 checked for purity.

65 Pure cultures were grown in nutrient broth containing 1.0 g meat extract; 1.0 g yeast extract; 5.0
66 g peptone; and 5.0 g sodium chloride per L. After 24 hours growth in nutrient broth (NB), cells were
67 pelleted by centrifugation, washed twice in sterile water and then re-suspended in sterile water
68 containing 0.05% Tween 20. Bacterial suspensions of each isolate were adjusted to a concentration of
69 5.0×10^9 cells ml⁻¹ and used as inoculum. Experimental plants consisted of 5-month old seedlings
70 grown in 3.75 L (1 gallon) containers using Morton's Nursery Mix consisting of 1:1:1 sand:bark:loam
71 (Morton's Horticultural Supplies Inc., McMinnville. TN). Plants were fertilized in early May using
72 water-soluble Miracle-Gro™, consisting of 18-24-16 (Nitrogen:Phosphorus:Potassium) at the rate of 18 g
73 per 3780 mL (w/v) and with controlled-release fertilizer (Nutricote Total™ 18-6-8), applied at the rate
74 of 12 g per container. Fungicide Cleary's 3336™ F (41.25% thiophanate-methyl, Cleary Chemical
75 Corp., Dayton, NJ) was used as a positive control at the recommended rate of 1.56 ml/L and water was
76 used as the non-treated control. Greenhouse environment was maintained at $26 \pm 3^\circ\text{C}$ and shadehouses

were covered with 65% shade cloth. Powdery mildew treatments were applied at 7-10 days intervals and their efficacy in suppressing powdery mildew severity was evaluated using a disease rating scale of 0-5 where, 0 = no disease symptoms, 1 = 1-10%, 2 = 11-25%, 3 = 26-50%, 4 = 51-75% and 5 = 75-100% of foliage covered with powdery mildew symptoms.

2.1.2. Efficacy of five bacterial BCA isolates after two years in cold storage

This study was conducted in greenhouse environment to compare five bacterial isolates with a fungicide positive control and non-treated, water control. Foliar spray method was used to apply treatments and plants were allowed to air dry for approximately 1 h and then placed in an incubation chamber maintained at 100% RH for 24 h to allow establishment of the biological control agents on plant surfaces prior to powdery mildew infection. Inoculation with powdery mildew pathogen was by using spore settling tower technique whereby spores from previously infected plants were blown into the air above test plants and allowed to float down onto test plants as described in **Mmbaga et al., 2008**. A continuous supply of air-borne powdery mildew inoculum was from previously infected plants scattered in the experimental area; powdery mildew control treatment was repeated every 7-10 days.

2.1.3. Evaluation of two isolates B17A and B17B as BCA for powdery mildew control.

Isolate 17A that showed superior efficacy as a powdery mildew BCA after two years in cold storage was designated B17A and compared to isolate B17B that was previously selected for superior efficacy against powdery mildew. Efficacies of these two isolates were compared on susceptible (S) and moderately resistant (MR) plants in shade-house environment under 65% shade-cloth and in greenhouse environment. Susceptible plants used in this study were dogwood selections 295, 327, 400 and moderately resistant plants were selections R11 and R12. Each plant selection was replicated by six individual plants per treatment and plants were arranged in a randomized complete block design. A

101 continuous supply of air-borne powdery mildew inoculum was from previously infected plants scattered
102 in the experimental area; treatments with BCA was initiated when test plants showed first disease
103 symptoms. In addition, BCA applications were done in the evening (approx. 6:00 pm) and natural dew
104 was the only source of free moisture for BCA plant colonization. Plants were sprayed to runoff using a
105 hand-held atomizer for even distribution of BCA suspension.

106

107 ***2.1.4. Efficacy of biocontrol agents applied by foliage spray and by root drenching.***

108 Treatments for disease control were initiated when the first disease symptoms were observed and
109 biocontrol isolate B17A and B17B suspensions of 5.0×10^9 bacterial cells ml^{-1} were applied using hand-
110 held atomizer and the foliage sprays or root drenching methods. Foliage spray treatments were done
111 using an atomizer to deliver the inoculum uniformly and plants were sprayed to run off; root drenching
112 was done using 20 ml inoculum per plant. Treatments were applied in late evening and dew was the
113 only source of free moisture for BCA plant colonization as described above; treatments were repeated
114 every 7-10 d until end of August or early September following grower practice for fungicide
115 applications.

116 In order to examine roots for endophytic presence of BCA when root drenching with BCA was
117 used, root clearing was done in 10% potassium hydroxide (KOH), acidified with 20% HCl, and stained
118 in 0.1% toluidine blue as described by **Phillip and Hayman, (1970)** with slight modifications. The
119 cleared roots were examined under a compound microscope; the presence of bacterial cells inside root
120 tissues was assessed at 400X-1000X magnifications. Root colonization with biological control agents
121 was also evaluated on plants generated from BCA inoculated seed in which seed coating with BCA was
122 done on emerged radicles soon after germinating. In this study, surface sterilized seed were first
123 vernalized to break seed dormancy at 37°C as described in **Evans and Blazich (1999)**. When vernalized

seed started germinating, they were drenched with BCAs (isolates B17A and B17B) using a bacterial suspensions of 5.0×10^9 bacterial cells ml^{-1} and planted in sterile soil in 15 cm x 20 cm containers; the non-treated control was drenched with sterile water. A replication of four containers with three plants in each container was used for each treatment and arranged in a randomized complete block experimental design. At the end of each study, roots were harvested and cleared for microscopic observations as described above.

2.2. *Effect of B17A and B17B bacterial suspensions and bacterial filtrates on spore germination*

Bacterial isolates B17A and B17B grown in nutrient agar for 24 hours were used to make suspensions of 5.0×10^9 bacterial cells ml^{-1} in sterile water and the bacterial suspension was sprayed onto the nutrient agar plates to form a thin film; sterile water was used as a negative control. The plates were incubated for approx. 12 hours then powdery mildew conidiospores were dusted onto the treated media, with spores being in direct contact with the bacteria. Germination of the conidiospores was monitored over time using a compound microscope (Leitz Diavert inverted microscope, Germany), at 200X and 320X magnification; germination was determined by any bulge or protrusions beyond the spore walls that either formed elongated or short germ tubes (**Pal and Gardener, 2006**). Spores that germinated were counted as a percentage all spores observed per field view at 200X magnification.

In addition, detached leaves were surface sterilized with 10% Clorox bleach (0.6% sodium hypochlorite) and placed on two layers of wet paper towels in an incubation chamber, two leaves per container. The leaves were brush-inoculated with powdery mildew spores and allowed to develop colonies of 2.5 - 3.5 cm diam. with fluffy mycelia visible without magnification. The leaves were then sprayed to runoff using bacterial suspensions or filtrates of the B17A and B17B with a replication of four containers and two leaves per container for each treatment. Experimental design used a randomized

complete block design and sterile water was used as non-treated control. Bacterial filtrates were prepared by growing isolates B17A and B17B in nutrient broth incubated on a shaker for 48 hours at 37 °C. The cultures were then centrifuged and filtered through 0.20 µm filters to obtain cell-free supernatant filtrates. The filtrate was observed under a compound microscope at 400X-1000X magnifications to confirm that they did not contain any bacterial cells before the filtrate was used to spray leaves to run off.

2.3. *Data analysis*

Disease severity and spore germination data analyses were performed using general linear models procedure of SAS (SAS/STAT 1990). Multiple comparisons between pairs of means were performed using a series of t-tests according to SAS procedures in PROC. ANOVA (**Gomez and Gomez 1984, SAS/STAT 1990**). The least significant differences (LSD) were calculated according to Fisher's protected LSD test at $P \leq 0.05$.

3. RESULTS

3.1 *Efficacy of bacterial isolates in powdery mildew disease control*

All five bacterial isolates that were retrieved from storage suppressed powdery mildew severity compared to non-treated control. Isolate B017, had superior efficacy and was statistically similar to fungicide control (**Fig. 1**). In addition to controlling powdery mildew, plants treated with isolate B017 grew larger than those treated with other bacteria or with conventional fungicide thiophanate methyl (data not shown).

Fig. 1(A-D): Efficacy of five bacterial isolates on powdery mildew disease severity on dogwood seedlings as compared to conventional fungicide (-■-) and non-treated control (-○-); (A) All five bacterial isolates tested, and B-D individual isolates compared to conventional fungicide thiophenate methyl, and water control; (B) B017- (-△-); (C) B08- (-▲-); and (D) B103- (-○-).

3.1.2. Evaluation of two isolates B17A and B17B as BCA for powdery mildew control.

A comparison of B017 (B17A) and another isolate B17B showed similar efficacy in suppressing powdery mildew disease (**Fig. 2**). At the end of the study, both susceptible and moderately resistant plants treated with BCA isolates had similar disease severity and had significantly lower disease than the non-treated control at $p < 0.0001$ (**Fig. 2**). The efficacy of the two BCA isolates in controlling powdery

178 mildew was statistically similar to conventional fungicide on both susceptible and moderately resistant
 179 selections (**Fig. 2**).

180

181 **3.1.3. Efficacy of biocontrol agents applied by foliage spray and by root drenching.**

182 All plants inoculated with BCAs by either foliage sprays or root drenching developed
 183 significantly less disease than the non-treated controls in both shadehouse and greenhouse experiments
 184 (**Fig. 3**). However, BCA application by foliage sprays was slightly more effective than root drenching
 185 (**Fig. 3**).

186

Fig. 2. Effect of two biological control agents (BCA B17A and B17B) on powdery mildew disease severity on susceptible (S) and moderately resistant (MR) plants as shown in S-(295) and R (R12 compared to conventional fungicide thiophenate methyl and water-treated (control) by foliar sprays in greenhouse environment; disease severity was rated on a scale of 1-5 in which 1 = 1-10%, 2 = 11-25%, 3 = 26-50%, 4 = 51-75% and 5 = 75-100% of foliage covered with powdery mildew symptoms. Similar results were obtained from shadehouse environment.

187
 188

Fig. 3. Effect of two biological control agents (B17A and B17B) on powdery mildew disease severity on susceptible dogwood seedlings maintained in greenhouse (GH) and shadehouse (SH) environments where foliar sprays and root drenching method of application were compared with conventional fungicide thiophenate methyl and water-treated (control); disease severity was rated on a scale of 1-5 in which 1 = 1-10%, 2 = 11-25%, 3 = 26-50%, 4 = 51-75% and 5 = 75-100% of foliage covered with powdery mildew symptoms. Similar results were obtained in repeated experiments

Roots that had been treated with BCA bacterial suspensions showed the presence of B17A and B17B bacterial cells inside root parenchyma cells and in some xylem tissues, indicating endophytic colonization of the roots (**Fig. 4**). Colonization of stem or leaves internal tissues was not examined.

Fig. 4. Microscopic observation of Cleared dogwood roots that had been treated with two bacterial biocontrol agents (BCA), isolates B17A and B17B by root drenching showing bacterial cells inside parenchyma cells (arrows)indicating endophytic colonization of the roots by the BCA. The roots were cleared in 10% potassium hydroxide (KOH), 1% hydrochloric acid (HCl) and counterstained in 1% toluidine blue-O.

3.2. Effect of B17A and B17B bacterial suspensions and bacterial filtrates on spore germination

Evaluation of conidiospore germination on growth media treated with B17A and B17B showed fewer spores germinated compared to the non-treated control; the BCA treated plates had significantly lower germination percentage than the water-treated controls at $p < 0.0001$ when the spores had a direct

203 contact with the BCA and when the BCA and powdery mildew spores were in separate sections of
 204 divided plates with no physical contact between them (**Fig. 5**).

205
 206 **Fig. 5.** Effect of biological control agents (BCAs) isolates B17A and B17B on powdery mildew
 207 conidiospores germination on nutrient agar (NA): (a) when powdery mildew spores were added on the
 208 NA media 12 hours after treatment with BCA and water treated control were in separate plate and (b)
 209 when BCA treatment and powdery mildew spores were in separate sections of divided plates with no
 210 physical contact between BCA and powdery mildew spores; water was used as (control). The
 211 experiment was repeated ones.
 212

213 Some spores exposed to BCA formed germ tubes that did not continue to develop after the initial
 214 bulge stage (**Fig. 6**). At 36 hours post-inoculation, bacterial cells were observed inside powdery mildew
 215 conidiospores and some spores appeared to be lysed and bacterial cells were flowing out of the
 216 conidiospores (**Fig. 6**). Powdery mildew colonies that were sprayed with BCA suspensions and BCA
 217 filtrates with no bacterial cells stopped expanding and collapsed; spores collected from treated colonies
 218 showed lower germination percentage and had fewer germ tubes than the non-treated controls. Some
 219 spores formed germ tube initials, but the germ tubes did not elongate compared to the non-treated
 220 control; some spore lyses were also observed on BCA filtrate treatments, but not in the non-treated
 221 water control

Fig. 6: Colonization of powdery mildew spores by bacterial biological control agents (A) The conidia germination showing a gem tube (Ca. 8 h after bacteria inoculation). (B) Gem tube after 15 hours and (C-D) after 36 hours showing ruptured/lysed spores.

4. DISCUSSIONS

Numerous microorganisms are naturally present on leaves of dogwoods and the majority of these are saprophytic (Mmbaga et al., 2008; Mmbaga and Sauvé, 2009). All five isolates tested for biocontrol activity reduced powdery mildew disease severity compared to the non-treated control (Figs. 1-2), but two isolates (B17A and B17B) displayed efficacy similar to that of conventional fungicides and demonstrated high potential as biological control agents (BCA). In the initial experiments, post-inoculation incubation under 100% humidity chambers was used to facilitate BCA-plant colonizations; this process facilitated the identification of isolates that had best potential as biocontrol agents. However, after this initial selection, moisture from natural dew formation was used to enhance BCA-plant colonization. It is reasonable to presume that these natural conditions were favorable to plant colonization with BCAs because the treatments were effective in suppressing powdery mildew. Results showed that the two BCAs, B17A and B17B, may provide effective alternatives to conventional fungicides in controlling powdery mildew in container-grown dogwoods in shadehouse environment (Figs. 2-3).

Weekly applications of isolates B17A and B17B were as effective as a conventional fungicide thiophanate methyl, but neither the fungicide nor BCAs eliminated or stopped the disease completely. The two isolates B17A and B17B were highly effective when they were allowed to colonize plants before the pathogen was introduced and when BCA application was initiated soon after initial disease

245 symptoms were observed and natural colonization was used without additional moisture. These results
246 show that the two bacterial isolates B17A and B17B offer potential microbial alternatives to
247 conventional fungicides in powdery mildew management. Weekly applications of the BCAs were used
248 to maintain high bacteria populations on leaf surfaces following reports from other studies (**Lindow et.**
249 **al., 2002**).

250 Previous studies have shown that seedling populations are more susceptible to powdery mildew
251 than adult plants (**Li et al., 2010**); thus seedling populations make excellent test plants for studies on
252 powdery mildew control. The two isolates were highly effective in controlling powdery mildew in
253 seedling populations and it is reasonable to presume that they would be highly effective on adult plants.
254 Seedling plants used in this study are also expected to have high genetic variability including variation
255 in powdery mildew susceptibility because dogwoods are self-sterile and obligately out-crossing (open
256 pollinated) (**Reed 1999**), thus the BCA isolates are likely to be effective on diverse populations.
257 Although host resistance is the best method for controlling powdery mildew, none of the commercial
258 cultivars have high resistance. Consequently, when disease pressure is high, growers often use fungicide
259 applications. Results from this study show that biological-based integrated disease management to
260 control powdery mildew in nursery production of flowering dogwoods can reduce the need for
261 conventional fungicides. Our results showed that moderately resistant plants sustained less disease
262 severity than susceptible plants when powdery mildew was not controlled (**Fig. 2**), but both B17A and
263 B17B BCAs were as effective on susceptible and moderately resistant plants (**Figs. 2-3**). The two BCA
264 isolates were statistically similar to conventional fungicide in controlling powdery mildew on both
265 susceptible and resistant selections (**Fig. 3**).

266 Foliage and root inoculation methods for BCA application suppressed powdery mildew and
267 reduced disease severity significantly compared to the non-treated control (**Fig. 3**), and the difference

268 between foliar sprays and root drenching methods of BCA application was not statistically significant.
269 It is possible that root inoculation with the two BCAs suppressed powdery mildew by Induced systemic
270 resistance (ISR), which may also include resistance to other fungal pathogens including pathogenic soil
271 microbes (**Haas and Defago, 2005**). A slight advantage of foliage sprays over root inoculations was
272 observed and this may be due to the direct effect of the bacterial cells on spores and on spore
273 germination as indicated by the spore lysis, reduced spore germination, and disruption of germ tube
274 walls (**Figs. 5-6**). Perhaps the combination of the two methods of application may be advantageous and
275 possibly improve efficacy by providing early application in the form of seed treatment followed by
276 foliar sprays; further study is needed to confirm the concept.

277 The effect of the two bacterial BCAs filtrates in disrupting spore germination suggests that the
278 BCA may produce deleterious compounds or secondary metabolites as reported in other fungal disease
279 control mechanisms from bacterial BCAs (**Zhou et al., 2007**). The effect of BCA isolates B17A and
280 B17B on spore germination including destruction of germ tube formations suggests that antibiosis may
281 be one of the mechanisms of action. Chemical compounds produced by the two BCA isolates have not
282 been reported in this article, but compounds that caused lyses of spore membranes and germ tube walls
283 may include various catabolic enzymes and some volatile compounds (**Schulz, et al., 2010**). In
284 addition, the BCA isolates displayed plant growth promoting properties that may be associated with
285 mechanisms reported for other growth promoting BCAs (**Haas and Defago, 2005; Singh et al., 2003**).
286 The mechanism of action for plant-growth promoting bacteria is not understood, but theories include
287 induced systemic resistance in the host plant (**Haas and Defago, 2005**).

288 Although the two BCAs isolates were isolated from leaf epiphytes, they colonized roots and
289 displayed endophytic colonization of treated plants. Root drenching with the BCA suppressed powdery
290 mildew severity and promoted plant growth. The colonization process of most bacterial endophytes

remains largely unknown. It is reasonable to presume that soil may be their natural habitat and the starting point for plant colonization and endophytic migration to the foliage, but studies on their presence in the rhizosphere are needed to further explore potential for commercial application of the selected BCAs (McSpadden and Fravel, 2002). The presence of bacterial cells inside parenchyma cells and some vascular tissues of cleared roots suggested a potential avenue for introducing the BCAs into dogwood plants. Kilic and Yuen (2000), Hoffland, et al. (1995), Leeman, et al. (1995; 1996) and Van Loon, et al. (1998) proposed that bacterial endophytes trigger induced systemic resistance (ISR) , it is likely that the two bacterial isolates are associated with ISR, but studies are required to confirm that and determine other mechanisms of action.

300

301 **Acknowledgements:**

302 The authors would like to thank Terry Kirby, Terri Simmons, Debbie Eskandarnia, and Candace
303 Stubblefield for their technical support during this study. This study was partly funded by a USDA grant
304 no. 203-38814-13921 and by USDA NIFA Grant no. TENX-2010-38821-21477.

305

306 **Literature Cited:**

307 Chartfield, J.A., Rose, M.A., 1996. Ornamental Plants Annual Report and Research Summaries.

308 Ohio State University Special Circular No. 152.

309 Chin-A-Wong T.F, Bloomberg G.V, Mulders I.H, Dekkers L.C, Lugtenberg B.J., 2000. Root

310 colonization by phenazine-1-carboxamide-producing bacterium *Pseudomonas chlororaphis*

311 PCL1391 is essential for biocontrol of tomato foot and root rot. *Mol Plant Microbe Interact* **13**

312 (12): 1340–1345.

313 Daughtrey, M.L., and Hagan, A.K., 2001. Dogwood diseases in Diseases

314 of Woody Ornamentals and Trees in Nurseries. Jones R. K., Benson, D.M. (eds). American
 315 Phytopathological Society, St. Paul, MN., Pages 124-132.

316 Elad, Y. 2000. Biological control of foliar pathogens by means of *Trichoderma harziunum* and potential
 317 modes of action. Crop protection 19: 709-714.

318 Elad, Y., Malathrakis, E., Dik, A.J., 1996. Biological control of *Botrytis*-incited diseases and powdery
 319 mildews in greenhouse crops. Crop protection. Vol. 15(3): 229-240.

320 Evans, E., Blazich, F.A., 1999. Overcoming Seed Dormancy: Trees and Shrubs. North Carolina
 321 Cooperative Extension Service, 1/99 HIL-8704.

322 Glick, B., 1995. The enhancement of plant growth by free-living bacteria. Can. J. Microbiol. 41:109-
 323 117.

324 Gaffney, T.D., Lam, S.T., Ligon, J., Gates, K., Frazelle, A., Di Maio, J., Hill, S., Goodwin, S.,
 325 Torkewitz, N., Allshouse A.M., 1994. Global regulation of expression of antifungal factors by a
 326 *Pseudomonas fluorescens* biological control strain. Mol. Plant-Microbe Interact. 7:455-463.

327 Gomez, K.A., Gomez, A.A., 1984. Statistical procedures for agricultural research. John Wiley and
 328 Sons, NY, 679 p.

329 Haas, D., Blumer, C., Keel. C., 2000. Biocontrol ability of fluorescent pseudomonads genetically
 330 dissected: importance of positive feedback regulation. Curr. Opin. Biotechnol. 11:290-297

331 Haas, D., Defago, G., 2005. Biological control of soil-borne pathogens by fluorescent pseudomonads.
 332 *Nature Reviews in Microbiology* 3 (4): 307–319.

333 Hartman, J., Dixon, E., Mmbaga, M.T., 2003. Evaluation of Tennessee dogwood selections for
 334 powdery mildew. Pest Management-Diseases.

335 Hoffland, E., Pieterse, C.M.L., Bik, L., van Pelt, J.A., 1995. Induced systemic resistance in radish is not
 336 associated with accumulation of pathogenesis-related proteins. *Physiol. Mol. Plant Pathol.*
 337 46:309-320.

338 Kai, M.M., Haustein, F., Molina, A., Petri, B., Scholz, B. P., 2009. Bacterial volatiles and their action
 339 potential. *Appl. Microbiol. Biotechnol.* 81: 1001-1012.

340 Kilic-Ekici, O. and Yuen, G.Y., 2004. Comparison of strains of *Lysobacter enzymogenes* and PGPR for
 341 induction of resistance against *Bipolaris sorokiniana* in tall fescue. *Biol. Control* 30:446-455.

342 Kiss, L. 2003. A review of fungal antagonists of powdery mildews and their potential as biocontrol
 343 agents. *Pest Manag. Sci.* Vol. 59: 475-483.

344 Leeman, M., van Pelt, J.A., Denouden, F. M., Heinsbroek, M., Bakker, P., and Schippers, B., 1995.
 345 Induction of systemic resistance against *Fusarium* wilt of radish by lipopolysaccharides of
 346 *Pseudomonas fluorescens*. *Phytopathology* 85:1021-1027.

347 Leeman, M., E., Denouden, M., van Pelt, J.A., Dirkx, F., Steijl, H., Bakker, P., Schippers, B., 1996. Iron
 348 availability affects induction of systemic resistance to *Fusarium* wilt of radish by *Pseudomonas*
 349 *fluorescens*. *Phytopathology* 86:149-155.

350 Leigh, A.K., Windham, M.T., Trigiano, R. N., 1998. Natural occurrence of *Microsphaera pulchra*
 351 and *Phyllactina guttata* on two *Cornus* species. *Plant Dis.* 82, 383-385.

352 Li, Y., Mmbaga, M.T., Windham, A.S., Windham, M.T., Trigiano, R.N., 2009. Powdery mildew of
 353 dogwoods: Current Status and Future Prospects. *Plant Dis.* 93-11-1084.

354 Lindow, S.E., Hecht-Poinar, E. I, and Elliott, V. J. ,(Eds) 2002. *Phyllosphere Microbiology*. The
 355 American Phytopathological Society. St. Paul, Minnesota. 398p.

356 McSpadden Gardener, B. B., Fravel, D. R., 2002. Biological Control of Plant Pathogens: Research,
 357 Commercialization, and Application in the USA.
 358 <http://www.apsnet.org/online/feature/biocontrol/top.html>
 359 McRitchie, J.J. 1994. Powdery mildew of flowering dogwood. Florida Department of Agriculture
 360 Consumer Service Plant Pathology Circular No. 368.
 361 Mmbaga, M.T. 1998. Effect of powdery mildew on growth of dogwood. Tenn. Nurserymen's Assoc.
 362 Newsl. 8:16–17.
 363 Mmbaga, M.T. 2000. Winter survival and source of primary inoculum of powdery mildew of dogwood
 364 in Tennessee. Plant Dis. Vol. 84:574-579.
 365 ———. 2002. Ascocarp formation and survival and primary inoculum production in *Erysiphe* (sect.
 366 *Microsphaera*) *pulchra* in dogwood powdery mildew. Ann. Appl. Biol. 141:153–161.
 367 Mmbaga, M.T., Sheng, H., 2002. Evaluation of biorational products for powdery mildew management
 368 in *Cornus florida*. J. Environ. Hort. 20:113–117.
 369 Mmbaga, M.T., Sheng, H., 2002. Evaluation of biorational products for powdery mildew management in
 370 *Cornus florida*. J. Environ. Hort. 20(2): 113-117.
 371 Mmbaga, M.T., Sauve, R.J., Mrema, F.A. 2008. Identification of microorganisms for biological control
 372 of powdery mildew in *Cornus florida*. Biological Control 44: 67–72.
 373 Mmbaga M.T., Sauv, R.J., 2009. Epiphytic microbial communities on foliage of fungicide treated and
 374 non-treated flowering dogwoods. Biological Control 49: 97-104.
 375 Nejad, P., Johnson, P.A., 2000. Endophytic bacteria induced growth promotion and wilt disease
 376 suppression in oilseed rape and tomato. Biological Control. Vol. 18:208-215.
 377 Pal, K., Gardener, K., McSpadden B., 2006. Gardener, Biological Control of Plant Pathogens. The Plant
 378 Health Instructor, 2006. doi: 10.1094/PHI-A-2006-1117-02.

379 Phillips J.M., Hayman D.S., 1970. Improved procedures for clearing roots and staining parasitic and
 380 vesicular-arbuscular mycorrhizal fungi for rapid assessment of infection. Trans. Brit. Mycol.
 381 Soc. 55, 158-161.

382 Reed, S.M., 1999. Development of a labor-efficient hand pollination procedure for flowering dogwood.
 383 J. Environ. Hort. 17:92-94.

384 Singh, U., Birinchi, P., Sarma, K., Singh D.P., 2003. Effect of Plant Growth-Promoting Rhizobacteria
 385 and Culture Filtrate of *Sclerotium rolfsii* on Phenolic and Salicylic Acid Contents in Chickpea
 386 (*Cicer arietinum*). Current Microbiology vol. 46: 0131 - 0140

387 SAS Institute. 1990. SAS/STAT Users Guide, Version 6. Statistical Analysis Systems Institute,

388 Schulz, S., Dickschat, J.S., Kunze, B., Wagner-Dobler, I., Diestel, R., Sasse, F., 2010. Biological activity of
 389 volatiles from marine and terrestrial bacteria. Mar. Drugs. 8:2976-2987.

390 Smith, V.L., 1999. First report of powdery mildew on *Cornus florida* in Connecticut caused by
 391 *Microshaera pulchra*. Plant Disease 83, 782.

392 Windham, A.S., 1994. Disease management of woody ornamentals in nurseries and
 393 commercial landscapes. Univ. Tenn. Agric. Ext. Serv. PB 1234.

394 Van Loon, L.C., Bakker, P.A., Pieterse, C. M. J., 1998. Systemic resistance induced by rhizosphere
 395 bacteria. Annu. Rev. Phytopathol. 36:453-483.

396 Zhou C.S., Mo, M.H. Q., Gu, Y., Zhou, J.P., Zhang, K.Q., 2007. Possible contributions of volatile-
 397 producing bacteria to soil fungistasis. *Soil Biol. and Biochem.* 39: 2371-2379. Inc., Cary, NC.

398

399 **Fig. 1(A-D):** Efficacy of five bacterial isolates on powdery mildew disease severity on dogwood
 400 seedlings as compared to conventional fungicide (-■-) and non-treated control (-□-); (A). All five

401 bacterial isolates tested, and B-D individual isolates compared to conventional fungicide thiophenate
402 methyl, and water control; (B) B017- (-Δ-); (C) B08- (-▲-); and (D) B103- (-O-).

403 **Fig. 2.** Effect of two biological control agents (BCA B17A and B17B) on powdery mildew disease
404 severity on susceptible (S) and moderately resistant (MR) plants as shown in S-(295) and R (R12
405 compared to conventional fungicide thiophenate methyl and water-treated (control) by foliar sprays in
406 greenhouse environment; disease severity was rated on a scale of 1-5 in which 1 = 1-10%, 2 = 11-25%,
407 3 = 26-50%, 4 = 51-75% and 5 = 75-100% of foliage covered with powdery mildew symptoms. Similar
408 results were obtained from shadehouse environment.

409 **Fig. 3.** Effect of two biological control agents (B17A and B17B) on powdery mildew disease severity on
410 susceptible dogwood seedlings maintained in greenhouse (GH) and shadehouse (SH) environments
411 where foliar sprays and root drenching method of application were compared with conventional
412 fungicide thiophenate methyl and water-treated (control); disease severity was rated on a scale of 1-5 in
413 which 1 = 1-10%, 2 = 11-25%, 3 = 26-50%, 4 = 51-75% and 5 = 75-100% of foliage covered with
414 powdery mildew symptoms. Similar results were obtained in repeated experiments

415 **Fig. 4.** Microscopic observation of Cleared dogwood roots that had been treated with two bacterial
416 biocontrol agents (BCA), isolates B17A and B17B by root drenching showing bacterial cells inside
417 parenchyma cells (arrows) indicating endophytic colonization of the roots by the BCA. The roots were
418 cleared in 10% potassium hydroxide (KOH), 1% hydrochloric acid (HCl) and counterstained in 1%
419 toluidine blue-O.

420

421 **Fig. 5.** Effect of biological control agents (BCAs) isolates B17A and B17B on powdery mildew
422 conidiospores germination on nutrient agar (NA): (a) when powdery mildew spores were added on the
423 NA media 12 hours after treatment with BCA and water treated control were in separate plate and (b)

424 when BCA treatment and powdery mildew spores were in separate sections of divided plates with no
425 physical contact between BCA and powdery mildew spores; water was used as (control). The
426 experiment was repeated ones.

427 **Fig. 6:** Colonization of powdery mildew spores by bacterial biological control agents (A) The conidia
428 germination showing a gem tube (Ca. 8 h after bacteria inoculation). (B) Gem tube after 15 hours and
429 (C-D) after 36 hours showing ruptured/lysed spores.

Effect of bacteria isolates in Powdery mildew control in flowering dogwoods (*Cornus florida* L.)

M.T. Mmbaga^{ac}, F.A. Mrema^b, L. Mackasmiel^{ac} and E. Rotich^a

^aTennessee State University, College of Agriculture, Human and Natural Sciences, Department of Agricultural and Environmental Sciences, Nashville TN, ^bOtis L. Floyd Nursery Research Center, McMinnville, TN 37110; ^bAlcorn State University, Research and Applied Sciences, MS 39096-7500

Graphic Abstract