1 **Acebes-Doria et al.: *H. halys* nymphal injury on tree fruit**

2 **Crop Protection**

3 Insect Damage Assessment

Corresponding author:

Angelita L. Acebes-Doria

Virginia Tech AHSAREC

595 Laurel Grove Road

Winchester, VA 22602

aacebes@vt.edu

11 **Injury to apples and peaches at harvest from feeding by *Halyomorpha halys* (Stål)**

12 **(Hemiptera: Pentatomidae) nymphs early and late in the season**

14 ANGELITA L. ACEBES-DORIA¹, TRACY C. LESKEY² and J. CHRISTOPHER BERGH¹

15 ¹Alson H. Smith Jr. Agricultural Research and Extension Center, Virginia Tech, Winchester, VA

16 22602

17 ²USDA-ARS, Appalachian Fruit Research Station, 2217 Wiltshire Rd, Kearneysville, WV,

18 25430

Abstract

Halyomorpha halys (Stål) (Hemiptera: Pentatomidae) is an invasive species that has become an important orchard pest in the Mid-Atlantic region of the US. Adults and nymphs feed on tree fruit. Feeding injury from adults has been characterized but the injury from nymphs has not been examined systematically. Since the four plant-feeding instars of *H. halys* (second through fifth) differ substantially in size, it is plausible that the effects of their feeding on fruit injury and injury expression may differ among them. We compared feeding injury at harvest from young nymphs (second plus third instars), older nymphs (fourth plus fifth instars), and adults that were caged on ‘Smoothie Golden’ apples and ‘Redhaven’ peaches in early June (peach and apple), late July (peach), and late August (apple). Individual apples and peaches were caged at fruit set and assigned to the following treatments (n = 28/treatment): 1) control (no *H. halys*), 2) young nymphs or 3) adults early in the season, and 4) young nymphs, 5) older nymphs or 6) adults later in the season. Fruit in each treatment were exposed to 3-4 young nymphs, two older nymphs or 1-2 adults placed in the cages for 96 h and evaluated for external and internal feeding injury within 36 h after harvest. No injury was recorded from unexposed peaches or apples. The percentage of injured fruit and number of injuries per fruit varied significantly among the exposed treatments. Early season feeding by young nymphs yielded the least injury to peaches and apples. In apples, the highest percentage of injured fruit and number of injuries per fruit were caused by late season feeding by adults. In peaches, early season adult feeding produced the highest percentage of injured fruit and injuries per fruit. More internal than external injury was recorded on peach and no such difference was observed on apple. The implications of these findings on *H. halys* management in fruit orchards are discussed.

Keywords: tree fruit, brown marmorated stink bug, *Malus domestica*, *Prunus persica*

Introduction

Halyomorpha halys (Stål) (Hemiptera: Pentatomidae) is an invasive and polyphagous pest from Asia and has become a serious threat to Mid-Atlantic tree fruit orchards. Its initial and widespread outbreak in this region in 2010 resulted in feeding injury that caused losses to the Mid-Atlantic apple crop at harvest valued at \$37 million and a complete loss of peaches in some Maryland orchards (Leskey et al., 2012a). Aside from the fact that *H. halys* populations have not yet shown indications of being adequately suppressed by native parasitoids and predators, several basic aspects of its biology and behavior contribute to its status as a key orchard pest. First, it is highly polyphagous (Rice et al., 2014) and present throughout most of fruiting period of orchard crops (Leskey et al., 2015). Many of its host plants are common at the forest edge adjacent to orchards in the Mid-Atlantic region (Acebes, 2016) and can support the development of large, localized populations that can pose a season-long risk to nearby orchards. The ability of *H. halys* to invade orchards season-long is facilitated by the strong flight dispersal capacity of adults (Wiman et al., 2014; Lee and Leskey, 2015) and the ability of nymphs to walk considerable distances in search of food plants (Lee et al., 2014). Finally, unlike the native stink bug pests of tree fruit, both nymphs and adults of *H. halys* are known to feed on fruit.

While the injury symptoms from feeding by adult *H. halys* on apples, *Malus domestica* Borkh., and peaches, *Prunus persica* (L.), have been well characterized (Nielsen and Hamilton, 2009), those associated with feeding by nymphs have not. Similar to other phytophagous stink bug species, *H. halys* has five nymphal instars that vary considerably in size (Hoebeke and Carter, 2003), of which the second through fifth feed on plants (Panizzi and Slansky Jr, 1985). Studies of other stink bug species indicate that the different nymphal stages have different feeding habits (Panizzi and Machado-Neto, 1992; Simmons and Yeargan, 1988) and may

therefore differ in the extent to which they cause economic injury to tree fruit and/or the nature of injury expression resulting from their feeding. In the Mid-Atlantic region where this study was conducted, *H. halys* has two generations which overlap spatially and temporally from mid-season onward (Leskey et al., 2012b; Bakken et al., 2015). During a portion of each growing season, *H. halys* captures in pheromone-baited traps consist predominantly of nymphs (Leskey et al., 2015); thus, it is important to understand the relative risk of injury from feeding by the different nymphal instars to apples and peaches at points during the season and the extent to which nymphal feeding contributes to economic injury at harvest.

Here, we characterized, quantified, and compared the injury to apple and peach fruit at harvest from feeding early and later in the season by **early instar** nymphs, **late instar** nymphs, and adults of *H. halys*.

Material and Methods

Insects and Study Site. Adult male and female *H. halys* collected from an overwintering population inside a dwelling in Stephens City, Virginia, USA in April 2015 were held in mixed-sex groups of 50-60 in 30.5 cm³ screened cages (BioQuip Products, Inc., Rancho Dominguez, CA, USA) in a laboratory room at ~25°C, ~70% RH, and a 16:8 (L:D) photo regimen. Cages were provisioned regularly with a diet of popcorn kernels, barley, buckwheat, soybeans, dried figs, dry roasted, unsalted peanuts and sundried tomatoes in a Petri dish and a fresh apple in a separate dish. Water was provided in a small plastic container with a cotton wick inserted through the lid. Folded paper towels on the cage floor and three to four freshly excised, compound leaves of tree of heaven, *Ailanthus altissima* (Mill.) Swingle, in a water-filled vase were provided as oviposition substrates. Starting in mid-May, egg masses in the cages were

collected at 2-d intervals and held in Petri dishes in the rearing room described above. After the eggs had hatched, a piece of wet cotton placed in each dish provided water for the nymphs. Upon molting to the second instar, nymphs were transferred to 30.48cm³ screened cages provisioned with the same diet described above. Second and third instars from these laboratory colonies were used in the early season field experiment. Additionally, subsets of the field-collected, overwintered adults reared in the laboratory since April were used in the early season field experiments. Due to fewer laboratory-reared insects available for the late season experiments, a combination of laboratory-reared and field-collected summer generation nymphs and adults were used. These were randomly assigned to the respective treatments. Insects from the field were collected by hand and sweep netting from tree of heaven and catalpa, *Catalpa speciosa* Warder, and held under the laboratory conditions described above for ≥ 24 h prior to use in the experiments.

Trees in a 1.1-ha experimental block of 7-yr-old ‘Smoothee Golden’ apples, and a 0.2-ha experimental block of 14-yr-old ‘Redhaven’ peaches, at Virginia Tech’s Alson H. Smith Jr. Agricultural Research and Extension Center (AHS AREC), Winchester, Virginia, USA were used. The orchards were treated with a routine disease management program but no insecticides were used during the study.

Inclusion Cage Experiments. The cylindrical, open-ended cages (17.8 cm diam, 17.8 cm long) used by Joseph et al. (2015) were employed (Fig. 1). These were constructed from semi-rigid, black plastic mesh (13 × 3 mm openings) covered by a 5-gal paint strainer (Ace Hardware Corp, Oak Brook, IL, USA) with one open end. A wooden dowel (0.7 cm diam, 18 cm long) placed perpendicularly in the center of cage was secured at both ends to the cage wall. Soon after fruit set (in early May), one or two peach and apple fruit growing near branch tips

115

116 **Figure 1. Cage used to enclose (A) apple and (B) peach fruit from early May until harvest. Insets without the netting highlight**
117 **the distance between the fruit and the cage wall.**

were enclosed within the cages. In mid-May the number of fruit per cage was reduced to one and only fruit that showed no evidence of injury from any insect pest were selected. Plastic twist ties were used to attach the dowel to the branch, to secure one edge of the plastic cage to the branch so the fruit was not touching the cage wall, and to close the opening of the mesh sleeve around the branch. To preserve the natural conditions as well as possible, four to seven peach leaves and three to five apple leaves were allowed to remain in each cage. Nymphal survivorship on a diet of peach or apple leaves alone is extremely low (Acebes-Doria et al. 2016), so feeding predominantly on fruit was assumed.

In total, 168 apples on 28 trees (6 fruit/tree) and 168 peaches on 14 trees (12 fruit/tree) were caged. Predetermined numbers of insects from the laboratory culture were transferred to small lidded plastic cups and released in each cage. To assess the effects of *H. halys* feeding either early or later in the fruiting period of each crop, there were two separate 96-hr exposure periods. For both apples and peaches, the early season exposure occurred from 6 to 10 June, 2015. Since fourth and fifth instars would not be expected in the field at the time of the early season experiment (Acebes, 2016), the following treatments were used; 1) control (not exposed to *H. halys*), 2) four second and third instars, and 3) two adults, with 28 replicates each. The instars were identified based on their size and the morphological features described by Hoebeke and Carter (2003). The late season exposure dates differed between the crops, based on differences in projected harvest dates and fruit maturity; peaches and apples were exposed from 20 to 24 July and from 21 to 25 August 2015, respectively. Since all nymphal stages would be expected in field populations of *H. halys* at the time of the late season exposures (Acebes, 2016), the treatments included, 1) control (not exposed to *H. halys*), 2) three second and third instars, 3) two fourth and fifth instars, and 4) one to two adults, each with 28 replicates. At the end of each

96-hr exposure period, the number of live and dead **insects** in each cage was recorded and the insects were removed from the cages. Of all cages to which **insects** were introduced, only 2.5% contained no live individuals at the end of the exposure period. However, the **insects** in those cages were assumed to have fed on the fruit for some portion of the 96-hr exposure and the fruit from them were included in the evaluation. The caged peaches and apples were harvested on 31 July and 11 September, respectively, coinciding with the approximate harvest dates for both crops in this region. Those that had dropped from the branch prior to harvest were excluded from the evaluation, which represented only 4.8 % of each crop. Fruit were stored at 4°C and evaluated for injury within 36 hr of harvest. The average daily temperature and rainfall during the exposure periods were recorded using a WatchDog weather station (Model 900ET, Spectrum Technologies, Inc., Plainfield, IL, USA) at the AHS AREC.

Injury Evaluation. Criteria used to evaluate injury to the apples and peaches followed Joseph et al. (2015). For apples, external injury was recorded as the number of ‘punctures’ and/or ‘dimples’ (Figure 2A) and discrete areas of surface discoloration with or without distinct depressions (Figs. 3A and 4A, respectively). Internal injury to apples was recorded as the number of areas of necrotic tissue after peeling the skin and slicing each fruit to the core in all quadrants (Figs. 2C, 3B and 4B). For peaches, external injury was recorded as the number of areas of gummosis (gummy, whitish exudate associated with tissue damage on stone fruit) (Fig. 5) and the number of surface depressions or deformations (Figs. 6A and 7A). Internal injury was recorded as the number of discrete areas of necrotic tissue after fruit were peeled and sliced to the core in all quadrants (Figs. 5B, 5C, 6B and 7B). To adjust the injury recorded based on differences in the number of insects per cage, the total number of injury sites on each fruit was divided by the number of bugs to which that fruit had been initially exposed.

Figure 2. Early season feeding injury from *H. halys* adults on 'Smoother Golden' apples; (A) External injury: punctures (inset) and 'dimpling' (B) Internal injury (less apparent) below the surface, (C) Evidence of deeper internal necrosis.

Figure 3. Late season feeding injury from third and fourth *H. halys* instars on 'Smoother Golden' apples; (A) External injury: surface discoloration with distinct depressions, (B) Internal injury (brown necrosis) below the surface, (C) No evidence of deeper internal necrosis.

Figure 4. Late season feeding injury from *H. halys* adults on 'Smoother Golden' apples; (A) External injury: surface discoloration without distinct depressions, (B) Internal necrosis just below the surface, (C) No evidence of deeper internal necrosis.

189

190

191 **Figure 5. Early season feeding injury from *H. halys* adults on 'Redhaven' peaches; (A) External injury: gummosis and surface**
192 **depressions (B) Internal injury (brown necrosis) below the surface, (C) Deeper internal necrosis.**

193

194

195

196

197

198

199

Figure 6. Late season feeding injury from third and fourth *H. halys* instars on 'Redhaven' peaches; (A) External injury: shallow depressions, (B) Internal injury (white necrosis) below the surface, (C) No evidence of deeper internal necrosis.

208

209 **Figure 7. Late season feeding injury from *H. halys* adults on 'Redhaven' peaches; (A) External injury: shallow depressions, (B)**

210 **Internal injury (white necrosis) below the surface, (C) No evidence of deeper internal necrosis.**

Data Analyses. Analyses were conducted using JMP[®] Pro version 11 (SAS Institute Inc., Cary, NC, 2007) and outcomes were considered significant at $P < 0.05$. The number of injured apples and peaches were compared among treatments using nominal logistic regression. When the likelihood ratio chi-square test indicated a significant overall treatment effect, the difference between two treatments was examined using the odds ratio. The number of each injury type per fruit and mean daily rainfall data were transformed using $\log(x+1)$ to satisfy the assumptions of parametric tests. Differences in the number of external injuries and the number of internal injuries per fruit among treatments in each experiment were analyzed using one-way analysis of variance (ANOVA). For external injuries, each injury type was analyzed separately. One-way ANOVA was also used to compare mean daily temperatures and rainfall among the exposure periods. When a significant treatment effect was found from ANOVA, means were compared using Tukey's HSD test. Differences between the number of surface depressions and internal injuries per fruit were compared using the pairwise t-test.

Results

Of all nymphs and adults released into cages, 86% were recovered alive after the 96-hr exposure period and 97.5% of cages contained ≥ 1 live individual. For both the early and late exposure periods, control fruit showed no evidence of injury (Tables 1 and 3), indicating that the cages effectively excluded *H. halys* and other pests, as was reported by Joseph et al. (2015). Mean ($\pm$ SE) daily temperatures recorded during the exposure periods on 6–10 June, 20–24 July and 21–25 August 2015 were $21.1 \pm 0.6^{\circ}\text{C}$, $24.0 \pm 0.9^{\circ}\text{C}$, and $22.3 \pm 0.5^{\circ}\text{C}$, respectively, which differed significantly ($F_{2,12} = 4.60$, $P = 0.03$). The mean daily temperature on 20–24 July 2015 was significantly higher than on 6–10 June and 21–25 August 2015, which did not differ. Mean ($\pm$ SE)

234 **Table 1. Percentage of caged ‘Smoother Golden’ apples injured at harvest following exposure to *H. halys* nymphs or adults for**
 235 **96 h early and late in the season.**

Treatment	No. of fruit evaluated	Percentage of apples with external injury		Percentage of apples with internal injury
		Punctures/ Dimpling ¹	Discolored depressions ²	
Control	23	0.00 b	0.00 d	0.00 d
<u>Early season</u>				
Young nymphs (2 nd /3 rd instars)	26	0.00 b	15.38 c	23.08 c
Adults	28	85.71 a	25.00 c	71.43 b
<u>Late season</u>				
Young nymphs (2 nd /3 rd instars)	28	3.57 b	64.29 b	60.71 b
Older nymphs (4 th /5 th instars)	28	0.00 b	92.86 a	75.00 ab
Adults	27	3.70 b	88.89 a	92.59 a
Statistics		$\chi^2 = 101.86$	$\chi^2 = 98.22$	$\chi^2 = 74.91$
		$P < 0.0001$	$P < 0.0001$	$P < 0.0001$

236 Means within columns followed by the same letter are not significantly different based on the χ^2 value from the odds ratio comparison
237 at $P < 0.05$.

238 ¹ Punctures and/or dimpling without discoloration (Figure 2A)

239 ² Surface discoloration with or without distinct depressions (Figs 3A and 4A, respectively)

daily rainfall during the respective exposure periods was 0.05 cm, 0.06 cm, and 0.0 cm, and did not differ significantly ($F_{2,12} = 0.56$, $P = 0.58$).

Apples. The percentage of apples showing external and internal injury at harvest differed significantly among the treatments (Table 1). The percentage of apples with punctures/dimpling was significantly highest on fruit exposed to adults early in the season. Exposure to older nymphs and adults later in the season resulted in the highest percentages of apples with discolored areas with or without distinct depressions (Table 1). Feeding by young nymphs later in the season also resulted in substantial surface injury that was significantly greater than earlier feeding by young nymphs and adults. The highest percentages of apples that incurred internal injury were recorded from those exposed to adults and older nymphs late in the season, followed by those exposed to adults early in the season and young nymphs later in the season.

Table 2 shows the severity of injury (i.e. number of injuries per fruit) to apples at harvest. Exposure to adults early in the season resulted in significantly more punctures/dimpling than all other treatments, which did not differ. Fruit exposed to adults late in the season showed the greatest numbers of discolored depressions (Table 2), followed by those exposed to older nymphs during the same period. Exposure to young nymphs and adults early in the season caused the fewest discolored depressions, and were comparable to those from exposure to young nymphs later in the season. Similarly, the numbers of internal injury sites were significantly highest from exposure to adults late in the season.

259 **Table 2. Mean ($\pm$ SE) number of injuries per fruit on caged ‘Smoother Golden’ apples at harvest following exposure to *H.***
260 ***halys* nymphs or adults for 96 h early and late in the season.**

Treatment	No. of fruit evaluated	Mean $\pm$ SE no. external injury*		Mean $\pm$ SE no. internal injury
		Punctures/ Dimpling ¹	Discolored depressions ²	
Control	23	0.00 b	0.00 d	0.00 d
<u>Early season</u>				
Young nymphs (2 nd /3 rd instars)	26	0.00 b	0.04 $\pm$ 0.02 d	0.07 $\pm$ 0.03 d
Adults	28	2.80 $\pm$ 0.42 a	0.18 $\pm$ 0.06 cd	0.95 $\pm$ 0.17 bc
<u>Late season</u>				
Young nymphs (2 nd /3 rd instars)	28	0.01 $\pm$ 0.01 b	0.60 $\pm$ 0.12 c	0.45 $\pm$ 0.11 cd
Older nymphs (4 th /5 th instars)	28	0.00 b	2.30 $\pm$ 0.51 b	2.07 $\pm$ 0.57 b
Adults	27	0.02 $\pm$ 0.02 b	5.56 $\pm$ 0.68 a	5.72 $\pm$ 0.67 a
Statistics		$F_{5,154} = 75.04$	$F_{5,154} = 57.06$	$F_{5,154} = 44.72$
		$P < 0.0001$	$P < 0.0001$	$P < 0.0001$

261 *No. of injuries per fruit divided by the corresponding no. of stink bugs released in each cage.

262 Means with columns followed by the same letter are not significantly different at $\alpha = 0.05$ using Tukey's HSD.

263 ¹ Punctures and/or dimpling without discoloration (Figure 2A)

264 ² Surface discoloration with or without distinct depressions (Figs 3A and 4A, respectively).

265

266

267

268

269

270

271

272

273

274

275

276

277

278

Peaches. The percentage of peaches that incurred external and internal injury varied significantly among treatments (Table 3). A high percentage of fruit exposed to adults early in the season expressed gummosis, and this injury was not otherwise detected. Significantly more peaches with surface depressions were recorded from fruit exposed to adults early in the season, followed by adults late in the season. Exposure to both young and older nymphs late in the season resulted in intermediate percentages of peaches with surface depressions that were not statistically different. The lowest percentage of fruit with this injury was recorded from fruit exposed to young nymphs early in the season. The percentages of fruit showing internal necrosis at harvest were significantly highest from adult exposures early and late in the season, followed by those exposed to older nymphs in late season. Young nymphs caused significantly more expressions of this injury from exposures during the late season than the early season, but both exposure periods resulted in a significantly lower percentage of injured fruit compared with the other treatments.

The number of internal and external injuries per peach at harvest differed significantly among the treatments (Table 4). Feeding by adults early in the season resulted in the greatest numbers of surface depressions ($P < 0.05$), followed by adults late in the season with all other treatments having significantly less and statistically equal numbers of depressions. Internal injuries were significantly more abundant on fruit exposed to adults early and late in the season, followed by those exposed to older nymphs late in the season. Feeding by young nymphs early and late in the season resulted in equivalent low levels of injury.

Comparisons between the number of surface depressions (external injury) and internal injury pooled across all treatments showed that peaches expressed more internal than external

301 **Table 3. Percentage of caged ‘Redhaven’ peaches injured at harvest following exposure to *H. halys* nymphs or adults for 96 h**
302 **early and late in the season.**

Treatment	No. of fruit evaluated	Percentage of peaches with external injury		Percentage of peaches with internal injury
		Gummosis	Surface depressions	
Control	26	0.00 b	0.00 d	0.00 d
<u>Early season</u>				
Young nymphs (2 nd /3 rd instars)	25	0.00 b	4.00 d	8.00 d
Adults	25	72.00 a	100.00 a	100.00 a
<u>Late season</u>				
Young nymphs (2 nd /3 rd instars)	28	0.00 b	46.43 c	39.29 c
Older nymphs (4 th /5 th instars)	28	0.00 b	39.29 c	82.14 b
Adults	28	0.00 b	75.00 b	96.43 ab
Statistics		$\chi^2 = 82.90$	$\chi^2 = 103.70$	$\chi^2 = 74.91$
		$P < 0.0001$	$P < 0.0001$	$P < 0.0001$

303 Means within columns followed by with the same letter are not significantly different based on the χ^2 value from the odds ratio
304 comparison at $P < 0.05$.

305 **Table 4. Mean ($\pm$ SE) no. of injuries per fruit on caged ‘Redhaven’ peaches at harvest following exposure to *H. halys* nymphs**
 306 **and adults for 96 h early and late in the season.**

Treatment	No. of fruit evaluated	Mean $\pm$ SE no. external injury*		Mean $\pm$ SE no. internal injury
		Gummosis	Surface depressions	
Control	26	0.00 b	0.00 c	0.00 c
<u>Early season</u>				
Young nymphs (2 nd /3 rd instars)	25	0.00 b	0.02 $\pm$ 0.02 c	0.03 $\pm$ 0.02 c
Adults	25	1.5 $\pm$ 0.32 a	4.00 $\pm$ 0.41 a	6.88 $\pm$ 0.51 a
<u>Late season</u>				
Young nymphs (2 nd /3 rd instars)	28	0.00 b	0.39 $\pm$ 0.15 c	0.54 $\pm$ 0.15 c
Older nymphs (4 th /5 th instars)	28	0.00 b	0.54 $\pm$ 0.14 c	2.18 $\pm$ 0.34 b
Adults	28	0.00 b	3.11 $\pm$ 0.66 b	7.93 $\pm$ 0.99 a
Statistics		$F_{5,154} = 41.51$	$F_{5,154} = 45.77$	$F_{5,154} = 102.32$
		$P < 0.0001$	$P < 0.0001$	$P < 0.0001$

307 *No. of injuries per fruit divided by the corresponding no. of stink bugs released in each cage.

308 Means within columns followed by the same letter are not significantly different at $\alpha = 0.05$ using Tukey’s HSD.

injury per fruit ($t = 6.98$, $df = 159$, $P < 0.0001$), while there was no difference between these two measures of injury in apples ($t = 1.41$, $df = 159$, $P = 0.16$).

Discussion

This study provided strong evidence that injury to apples and peaches from feeding by *H. halys* differed according to the life stage to which fruit were exposed and to the time at which feeding occurred. The types of injury recorded were similar to those described previously (Nielsen and Hamilton, 2009; Leskey et al., 2012c; Joseph et al., 2015). However, data from field studies by Leskey et al. (2012c) and Joseph et al. (2015) reasonably assumed that most of the injury recorded was from feeding by *H. halys*, based on recent grower experiences and on much higher populations of *H. halys* than native stink bugs species in orchards, but did not differentiate between injury caused by adults and nymphs. In a similar cage study by Nielsen and Hamilton (2009), apples and peaches were exposed only to adult *H. halys*. Here, we have systematically characterized the differences between adult and nymphal feeding injury and showed that injury to apples and peaches was quantitatively and qualitatively different. Exposure to second and third instars early in the season consistently resulted in lowest levels of external and internal injury to apples and peaches. However, with the exception of gummosis in peaches, young nymphs did cause both external and internal injury to both crops. Later in the season, when the fruit were at or near full maturation, second and third instars caused significantly or numerically more external and internal injury to both crops than they did early in the season. Most injury assessments following exposures late in the season revealed that older nymphs caused significantly or numerically more injury than younger nymphs. With the exception of punctures/dimpling from adult feeding on apples early in the season, more apples exposed to

adults late in the season were injured and showed more injury than those exposed to adults early in the season. Also, punctures alone would not likely be recorded as economic injury at harvest if not associated with other surface injury. Feeding by adults and older nymphs late in the season caused similar percentages of apples to show external and internal injury at harvest, although the severity of injury was higher for adults than older nymphs. In peaches, high percentages of fruit showed surface depressions and internal necrosis following exposure to adults early or late in the season, with similar levels of injury severity between the exposure periods. During the late season exposure, adults caused significantly or numerically more injured fruit than older nymphs and significantly greater numbers of injuries per fruit.

Early season injuries on apples were expressed mainly as punctures/dimpling and as less apparent internal injury, although some fruit showed necrotic areas deeper in the flesh (Fig. 2). The less pronounced expression of injury to apples exposed to *H. halys* early in the season may possibly be attributed to the rapid rate of cell division at that time. It may be helpful for growers to monitor stink bug injury by not only peeling the skin of fruit that have evident external injury but also to slice the fruit to check for deeper internal damage. Late season injury on apple from older nymphs and adults was expressed as discolored surface depressions and as internal tissue necrosis which was mostly close to the surface (Figs. 3 and 4). These findings suggest that apples are more susceptible to economic injury at harvest from *H. halys* feeding late in the season than earlier, concurring with previous findings by Nielsen and Hamilton (2009). Similarly, the most susceptible developmental stages of soybean for *H. halys* injury were also in the late season, which coincided with the largest annual *H. halys* populations that consisted of older nymphs and adults (Nielsen et al., 2011). Although some level of external injury and minor internal injury from *H. halys* may be tolerated on Mid-Atlantic processing apples, the degree of tolerance each

year may vary inversely with crop yield and quality. Heavy internal injury can adversely affect fruit acceptability for some processed products, such as frozen apple slices, and both external and internal fruit defects from *H. halys* feeding are problematic for growers who produce apples for fresh market.

On peaches, gummosis was an indicator of external feeding injury from *H. halys* early in the season and was only recorded from fruit exposed to adults during that period (Fig. 5), concurring with previous findings (Joseph et al., 2015) that gummosis was associated with early season stink bug feeding injury and was often accompanied by surface discoloration or depressions and internal necrosis. Early season injury on peach was more pronounced than late season injury, with obvious surface discoloration and deformations and deeper brownish internal necrosis (Fig. 5), whereas injuries from nymphs and adults later in the season were expressed mainly as shallow surface depressions and superficial whitish necrosis (Figs. 6 and 7). During our study, we never observed the classic ‘cat-facing’ injury on peaches, which is manifest as shallow to deep irregular-shaped deformations and depressions on the fruit surface (Howitt 1993) and are usually attributed to feeding by native stink bugs or plant bugs (Rings 1957). Since Mid-Atlantic peaches are a high-value, fresh market crop, these results imply that attack by *H. halys* either early or late in the season can negatively affect peach marketability.

The lack of apparent peach injury at harvest from feeding by young nymphs early in the season suggests that early season injury is mainly caused by overwintered adults (Leskey et al., 2012c), conforming with observations that overwintered adults start moving into orchards in mid-April (Leskey et al., 2015). Leskey et al. (2014) found that overwintered adults were more susceptible to insecticide applications than F₁ summer generation adults, suggesting the utility of early season insecticide applications targeting *H. halys* on peach. On the other hand, the use of

stronger, broad spectrum insecticides may be more useful later in the season, when populations are mostly composed of older nymphs and adults than young nymphs. Additionally, since less injury to apples was recorded from feeding by both nymphs and adults early than later in season, managing *H. halys* in apple orchards in the early season may not be as critical as later, when populations consist predominantly of older nymphs and adults (Acebes 2016).

Our findings that the numbers of external and internal injuries on apples did not differ significantly but that peaches showed significantly more internal than external injury confirm those of Joseph et al. (2015) and affirm that this can be an especially problematic issue for grading and marketing peaches if some fruit are not cut and inspected prior to sale.

With respect to ongoing efforts to develop and refine *H. halys* management decision thresholds based on cumulative captures in pheromone-baited traps in orchards, our results suggest that the inclusion of fourth and fifth instars may be important to consider. To date, provisional thresholds have relied only on adult captures (Short et al., unpublished), but nymphs are equally attracted to pheromone traps (Weber et al., 2014; Leskey et al., 2015) and feeding injury from the later instars is qualitatively and quantitatively comparable to that from adults. Moreover, *H. halys* is generally considered a perimeter-driven threat; both nymphs and adults are capable of dispersing from wild hosts into orchards and have been captured in orchards in substantial numbers, particularly in the border areas adjacent to woodlands (Bergh and Leskey, unpublished results) where highest injury from *H. halys* has been recorded (Leskey et al., 2012b; Joseph et al., 2014). Acebes-Doria et al. (2016) showed that *H. halys* nymphs often require a diet of ≥ 2 host plants for optimal performance and survivorship and Acebes (2016) documented the walking dispersal of nymphs onto and off of fruit and wild tree host along the orchard-woodland interface. In combination, these studies suggest that combining trap-based thresholds

with management tactics at the orchard perimeter (e.g. border sprays, Blaauw et al., 2015 or “attract and kill”, Morrison et al., 2015) may be promising tactics for this pest.

Temperature is known to affect the feeding activity of *H. halys* adults. Wiman et al. (2014) reported that optimal feeding activity was observed between 16°C and 17°C and that no feeding occurred above 26.5°C - 29.5°C. As well, more injury to blueberries from late instars and adults was observed in Oregon than in New Jersey and attributed to optimal temperatures in Oregon (Wiman et al., 2015). Whereas those studies addressed the feeding activity and amount of injury to blueberries from *H. halys*, the present study focused primarily on the expression of injury at harvest, with the amount of injury being a secondary measurement. Mean daily temperatures during the different exposure periods ranged from 19°C to 25°C, implying that *H. halys* would feed during at least portions of each 96-hr period, which was confirmed by the various injuries recorded on peaches and apples from both early and late season exposures and from all life stages. There is no information on the effects of abiotic factors on the expression of injury from *H. halys* feeding on apples or peaches. Aside from extreme abiotic conditions, which were not encountered during this study, annual conditions of temperature and rainfall during the period between when fruit were exposed and harvest would be expected to fluctuate around some yearly average. Since most of the types of injuries recorded were the result of a physiological response of fruit that are not expressed immediately after feeding but rather develop over some period of time, we contend that normal variations in abiotic conditions are not likely to result in differences in injury expression at harvest. Consequently, despite data from a single season, we feel that the injury recorded at harvest was representative of that caused by nymphs and adults at the different exposure timings and phenological stages of fruit development and would not likely change appreciably among years.

In conclusion, *H. halys* nymphs caused injury to tree fruit and the extent to which the injuries were expressed varied significantly depending on the developmental stage of the nymphs and the period during which feeding occurred. Our results confirmed previous findings regarding the stage of fruit development during which apple and peach fruit were more vulnerable to feeding injury from *H. halys* adults and provide useful information that has implications for developing temporally and spatially precise management programs against *H. halys* in apple and peach orchards.

Acknowledgements

We sincerely grateful to Jean Engelman, Eliezer Doria and the 2015 summer interns, Sophia Dutterer, Rachel Wilk, Jason Chou and Brendan Yarbrough for their excellent technical assistance. This material is based upon work that is supported by the National Institute of Food and Agriculture, U.S. Department of Agriculture, under award number #2011-51181-30937.

References

- Acebes, A.L., 2016. Host plant effects on the biology, behavior and ecology of brown marmorated stink bug, *Halyomorpha halys* (Stål) (Hemiptera: Pentatomidae). Doctoral Dissertation. Virginia Polytechnic Institute and State University
<http://hdl.handle.net/10919/64910>.
- Acebes-Doria, A.L., Leskey, T.C., Bergh, J.C., 2016. Host plant effects on *Halyomorpha halys* (Hemiptera: Pentatomidae) nymphal development and survivorship. Environ. Entomol. 45, 663-670.

446 Bakken, A.J., Schoof, S.C., Bickerton, M., Kamminga, K.L., Jenrette, J.C., Malone, S., Abney,
 447 M.A., Herbert Jr, D.A., Reisig, D., Kuhar, T.P., Walgenbach, J.F., 2015. Occurrence of
 448 brown marmorated stink bug (Hemiptera: Pentatomidae) on wild hosts in nonmanaged
 449 woodlands and soybean fields in North Carolina and Virginia. Environ. Entomol. 44,
 450 1011-1021.

451 Blaauw, B.R., Polk, D., Nielsen, A.L., 2015. IPM-CPR for peaches: Incorporating behaviorally-
 452 based methods to manage *Halyomorpha halys* and key pests in peach. Pest. Manag. Sci.
 453 71, 1513-1522.

454 Hoebeke, E.R., Carter, M.E., 2003. *Halyomorpha halys* (Stål) (Heteroptera: Pentatomidae): a
 455 polyphagous plant pest from Asia newly detected in North America. Proc. Entomol. Soc.
 456 Washington 105, 225-237.

457 Howitt, A.H., 1993. Common tree fruit pests. Michigan State University Extension. North
 458 Central Region Extension Publication #63. 252 p.

459 Joseph, S.V., Stallings, J.W., Leskey, T.C., Krawczyk, G., Polk, D., Butler, B., Bergh, J.C.,
 460 2014. Spatial distribution of brown marmorated stink bug (Hemiptera: Pentatomidae)
 461 injury at harvest in mid-Atlantic apple orchards. J. Econ. Entomol. 107, 1839-1848.

462 Joseph, S.V., Nita, M., Leskey, T.C., Bergh, J.C., 2015. Temporal effects on the incidence and
 463 severity of brown marmorated stink bug (Hemiptera: Pentatomidae) feeding injury to
 464 peaches and apples during the fruiting period in Virginia. J. Econ. Entomol. 108, 592-
 465 599.

466 Lee, D.-H., Nielsen, A.L., Leskey, T.C., 2014. Dispersal capacity and behavior of nymphal
 467 stages of *Halyomorpha halys* (Hemiptera: Pentatomidae) evaluated under laboratory and
 468 field conditions. J. Insect Behav. 27, 639-651.

469 Lee, D.H., Leskey, T.C., 2015. Flight behavior of foraging and overwintering brown marmorated
 470 stink bug, *Halyomorpha halys* (Hemiptera: Pentatomidae). Bull. Entomol. Res. 105, 566-
 471 573.

472 Leskey, T.C., Hamilton, G.C., Nielsen, A.L., Polk, D.F., Rodriguez-Saona, C., Bergh, J.C.,
 473 Herbert, D.A., Kuhar, T.P., Pfeiffer, D., Dively, G.P., Hooks, C.R.R., Raupp, M.J.,
 474 Shrewsbury, P.M., Krawczyk, G., Shearer, P.W., Whalen, J., Koplinka-Loehr, C., Myers,
 475 E., Inkley, D.B., Hoelmer, K., Lee, D.H., Wright, S.E., 2012a. Pest status of the brown
 476 marmorated stink bug, *Halyomorpha halys* in the USA. Outlooks Pest Manag. 23, 218-
 477 226.

478 Leskey, T.C., Wright, S.E., Short, B.D., Khrimian, A., 2012b. Development of behaviorally based
 479 monitoring tools for the brown marmorated stink bug, *Halyomorpha halys* (Stål)
 480 (Heteroptera: Pentatomidae) in commercial tree fruit orchards. J. Entomol. Sci. 47, 76-85.

481 Leskey, T.C., Short, B.D., Butler, B.R., Wright, S.E., 2012c. Impact of the invasive brown
 482 marmorated stink bug, *Halyomorpha halys* (Stål), in mid-Atlantic tree fruit orchards in
 483 the United States: case studies of commercial management. Psyche: J. Entomol. 2012,
 484 535062. DOI: 10.1155/2012/535062

485 Leskey, T.C., Short, B.D., Lee, D.H., 2014. Efficacy of insecticide residues on adult *Halyomorpha*
 486 *halys* (Stål) (Hemiptera: Pentatomidae) mortality and injury in apple and peach orchards.
 487 Pest. Manag. Sci. 70, 1097-1104.

488 Leskey, T.C., Agnello, A., Bergh, J.C., Dively, G.P., Hamilton, G.C., Jentsch, P., Khrimian, A.,
 489 Krawczyk, G., Kuhar, T.P., Lee, D.-H., Morrison, W.R., Polk, D.F., Rodriguez-Saona,
 490 C., Shearer, P.W., Short, B.D., Shrewsbury, P.M., Walgenbach, J., Weber, D.C., Welty,
 491 C., Whalen, J., Wiman, N.G., Zaman, F., 2015. Attraction of the invasive *Halyomorpha*

492 *halys* (Hemiptera: Pentatomidae) to traps baited with semiochemical stimuli across the
 493 United States. Environ. Entomol. 44, 746-756.

494 Morrison, W.R.I., Lee, D.-H., Short, B.D., Khrimian, A., Leskey, T.C., 2015. Establishing the
 495 behavioral basis for an attract-and-kill strategy to manage the invasive *Halyomorpha*
 496 *halys* in apple orchards. J. Pest Sci. 89, 81-96.

497 Nielsen, A.L., Hamilton, G.C., 2009. Seasonal occurrence and impact of *Halyomorpha halys*
 498 (Hemiptera: Pentatomidae) in tree fruit. J. Econ. Entomol. 102, 1133-1140.

499 Nielsen, A.L., Hamilton, G.C., Shear, P.W., 2011. Seasonal phenology and monitoring of the non-
 500 native *Halyomorpha halys* (Hemiptera: Pentatomidae) in soybeans. Environ. Entomol.
 501 40: 231-238.

502 Panizzi, A.R., Machado-Neto, E., 1992. Development of nymphs and feeding habits of nymphal
 503 and adult *Edessa meditabunda* (Heteroptera: Pentatomidae) on soybean and sunflower.
 504 Ann. Entomol. Soc. Am. 85, 477-481.

505 Panizzi, A., Slansky Jr, F., 1985. Review of phytophagous pentatomids (Hemiptera: Pentatomidae)
 506 associated with soybean in the Americas. Fla. Entomol., 184-214.

507 Rice, K.B., Bergh, C.J., Bergmann, E.J., Biddinger, D.J., Dieckhoff, C., Dively, G., Fraser, H.,
 508 Garipey, T., Hamilton, G., Haye, T., Herbert Jr, D.A., Hoelmer, K., Hooks, C.R.R.,
 509 Jones, A.L., Krawczyk, G., Kuhar, T.P., Martinson, H.M., Mitchell, W., Nielsen, A.L.,
 510 Pfeiffer, D., Raupp, M.J., Rodriguez-Saona, C., Shearer, P.W., Shrewsbury, P.M.,
 511 Venugopal, P.D., Whalen, J., Wiman, N.G., Leskey, T.C., Tooker, J., 2014. Biology,
 512 ecology, and management of brown marmorated stink bug (Hemiptera: Pentatomidae). J.
 513 Integr. Pest Manag. 5, A1-A13.

514 Rings, R.W., 1957. Types and seasonal incidence of stink bug injury to peaches. J. Econ. Entomol.
515 50, 599-604.

516 Short, B.D., Khrimian, A., Leskey, T.C. Development of pheromone-based decision support tools
517 for management of *Halyomorpha halys* in apple orchards. J. Pest Sci, (submitted).

518 Simmons, A.M., Yeargan, K.V., 1988. Feeding frequency and feeding duration of the green stink
519 bug (Hemiptera: Pentatomidae) on soybean. J. Econ. Entomol. 81, 812-815.

520 Weber, D.C., Leskey, T.C., Walsh, G.C., Khrimian, A., 2014. Synergy of aggregation pheromone
521 with methyl (E, E, Z)-2, 4, 6-decatrienoate in attraction of *Halyomorpha halys*
522 (Hemiptera: Pentatomidae). J. Econ. Entomol. 107, 1061-1068.

523 Wiman, N.G., Walton, V.M., Shearer, P.W., Rondon, S.I., 2014. Electronically monitored labial
524 dabbing and stylet ‘probing’ behaviors of brown marmorated stink bug, *Halyomorpha*
525 *halys*, in simulated environments. PLoS One 9, e113514.

526 Wiman, N.G., Walton, V.M., Shearer, P.W., Rondon, S.I., Lee, J.C., 2014. Factors affecting flight
527 capacity of brown marmorated stink bug, *Halyomorpha halys* (Hemiptera:
528 Pentatomidae). J. Pest Sci. 88, 37-47.

529 Wiman, N.G., Parker, J.E., Rodriguez-Saona, C., Walton, V.M., 2015. Characterizing damage of
530 brown marmorated stink bug (Hemiptera: Pentatomidae) in blueberries. J. Econ.
531 Entomol. 108, 1156-1163.