**Differing Foraging Strategies Influence Mercury (Hg) Exposure in an
Antarctic Penguin Community**

Michael J. Polito^{1*}, Rebecca L. Brasso², Wayne Z. Trivelpiece³, Nina Karnovsky⁴, William P.
Patterson⁵, Steven D. Emslie⁶

¹ Louisiana State University, Department of Oceanography and Coastal Sciences, Baton Rouge,
LA, 70802, USA *mpolito@lsu.edu

² Southeast Missouri State University, Department of Biology, One University Plaza, MS 6200,
Cape Girardeau. MO 63701

³ National Marine Fisheries Service, Antarctic Ecosystem Research Division, Southwest
Fisheries Science Center, La Jolla, CA 92037-1508, USA

⁴ Department of Biology, Pomona College, Claremont, CA 91711, USA

⁵ University of Saskatchewan, Department of Geological Sciences, Saskatoon, Saskatchewan,
Canada S7N 5E2

⁶ University of North Carolina Wilmington, Department of Biology and Marine Biology,
Wilmington, NC, 28403, USA

Abstract: Seabirds are ideal model organisms to track mercury (Hg) through marine food webs as they are long-lived, broadly distributed, and are susceptible to biomagnification due to foraging at relatively high trophic levels. However, using these species as biomonitors requires a solid understanding of the degree of species, sexual and age-specific variation in foraging behaviors which act to mediate their dietary exposure to Hg. We combined stomach content analysis along with Hg and stable isotope analyses of blood, feathers and common prey items to help explain inter and intra-specific patterns of dietary Hg exposure across three sympatric *Pygoscelis* penguin species commonly used as biomonitors of Hg availability in the Antarctic marine ecosystem. We found that penguin tissue Hg concentrations differed across species, between adults and juveniles, but not between sexes. While all three penguins species diets were dominated by Antarctic krill (*Euphausia superba*) and to a lesser extent fish, stable isotope based proxies of relative trophic level and krill consumption could not by itself sufficiently explain the observed patterns of inter and intra-specific variation in Hg. However, integrating isotopic approaches with stomach content analysis allowed us to identify the relatively higher risk of Hg exposure for penguins foraging on mesopelagic prey relative to congeners targeting epipelagic or benthic prey species. When possible, future seabird biomonitoring studies should seek to combine isotopic approaches with other, independent measures of foraging behavior to better account for the confounding effects of inter and intra-specific variation on dietary Hg exposure.

Capsule Abstract: Differences in penguin foraging ecology identified from stable isotope and stomach content analyses influence mercury exposure with species foraging on mesopelagic prey at higher risk.

Keywords: Mercury; Stable isotope analysis; *Pygoscelis* penguins; Antarctica

1. Introduction

Mercury (Hg) is a potent neurotoxin leading to physiological and reproductive impairments in wild birds (Brasso and Cristol 2008, Evers et al. 2008, Wada et al. 2009). In marine systems, seabirds are often used as biomonitors to assess the distribution, bioavailability and toxicity of Hg across the world's oceans (Kojadinovic et al. 2007, Day et al. 2012; Brasso et al. 2015). Seabirds are ideal model organisms to track Hg through marine food webs as they are long-lived, have wide-spread foraging ranges, and are susceptible to biomagnification due to feeding at relatively high trophic levels (Burger and Gochfeld 2000, Bond and Diamond 2009).

However, using seabirds as biomonitors for Hg bioavailability requires a solid understanding of species-specific foraging ecologies and other aspects of their life-histories. This is because species living in the same breeding location could be exposed to different concentrations of dietary Hg due to differences in prey preferences, small-scale foraging habitat use, and larger-scale migration and dispersal strategies (Anderson et al. 2009, Carravieri et al. 2014). Therefore, when comparing Hg exposure across and within seabird communities interspecific variation in foraging ecology must be explicitly taken into account. Even in single species studies, factors such as age, sex and individual-based differences in foraging ecology and thus Hg exposure necessitate an understanding of the degree of intra-specific variation in these factors within populations (Becker et al. 2000, Pedro et al. 2015).

The use of stable isotope analysis in applied studies of the diets and foraging ecology of seabirds has developed rapidly in recent years (Bond and Jones 2009). Nitrogen isotopic values ($\delta^{15}\text{N}$) in seabird tissues are commonly used to infer trophic level and diets (Polito et al. 2011a, Brasso and Polito 2013), while carbon isotopic values ($\delta^{13}\text{C}$) help trace trends in marine habitat use (inshore/benthic vs. offshore/epipelagic; Cherel and Hobson 2007). These relationships have

led stable isotope analysis to be often incorporated into modern ecotoxicological investigations (Jardine et al. 2006). Integration of stable isotope and Hg analyses have allowed researchers to track biomagnification of Hg within marine food webs (Atwell et al. 1998) and identify trophic differences among sympatric species that explain patterns of Hg exposure (Blévin et al. 2013, Pouilly et al. 2013). In addition, stable isotope-based dietary studies, including those using dietary mixing models, avoid many of the digestive and temporal biases inherent to more traditional methods such as stomach content analysis (Karnovsky et al. 2012). Even so, stable isotope-based dietary mixing models are limited by their inability to estimate the dietary contribution of specific prey species with overlapping isotopic values (Phillips et al. 2005). Often these limitations can be overcome in seabirds by combining isotopic approaches with the analysis of otoliths and other identifiable prey remains recovered from stomach contents (Polito et al. 2011a).

The objective of this study was to identify how differences in the foraging strategies of the three sympatric *Pygoscelis* penguin species influenced their exposure to Hg in the Antarctic Peninsula. *Pygoscelis* penguins are commonly used as biomonitors for Hg availability in Antarctic marine ecosystem as they forage primarily on Antarctic krill (*Euphausia superba*), nest in large, readily accessible breeding colonies, and tissues such as blood, feathers, guano and eggshells can be used to track Hg exposure across much of their annual cycle (Brasso et al. 2012, 2014, Celis et al. 2015). However, studies integrating stable isotope and Hg analyses in *Pygoscelis* penguins have often reported species-specific differences in tissue Hg concentrations which do not correlate with stable isotope based estimates of species relative trophic position (e.g. Brasso et al. 2012, 2014, 2015). This unpredicted uncoupling of trophic position and Hg concentration has hampered the interpretation and use of these species as biomonitors of Hg

availability across and within studies. As such a greater understanding of how the inter and intra-specific variation in the foraging habitats and diets of these species influences their exposure to Hg in the Antarctic marine food web is required.

To address this issue we examined variation in the foraging ecology and Hg exposure of Adélie (*Pygoscelis adeliae*), chinstrap (*P. antarctica*), and gentoo (*P. papua*) penguins breeding on King George Island in the Antarctic Peninsula region. Specifically, we use a combination of stable isotope and stomach content analysis to identify species, age and sex-specific dietary and foraging habitat trends and relate them tissue Hg concentrations reflecting both short-term dietary exposure (blood) as well as over longer annual, or inter-molt, time periods (feathers). Blood is used here as measure of short-term Hg exposure as concentrations in blood represent an integrated signal of recent dietary uptake and internal tissue redistribution of Hg (Evers et al. 2005, Ackerman et al. 2008). For example, lab dosing studies of non-molting seabirds indicate blood Hg concentrations reflect dietary Hg exposure within days of dosing with half-life of incorporation averaging between 30 and 60 days (Bearhop et al. 2000a). Due to the reasonable overlap in the timeframe of integration blood Hg and stable isotope values are often compared in ecotoxicological studies of seabird (e.g. Bond and Diamond 2009, Hipfner et al. 2011). For example, the half-life of stable isotope values in whole blood of a 3-5 kg penguin is predicted to be approximately 10-30 days based on allometric and empirical studies (Carleton and Martinez del Rio 2005, Barquete et al. 2013). In contrast, Hg concentrations in feathers provide a temporally integrative signal of annual or inter-molt body Hg burden in birds (Evers et al. 2008, Bond and Diamond 2009). During molt feather Hg concentrations are highly correlated with blood Hg concentrations as both are reflective of remobilized Hg from body tissues (Bearhop et al. 2000a). Stable isotope analyses feathers provides a similarly integrated signal of diets during

the inter-molt period owing to the unique pattern of catastrophic molt and fasting in adult penguins (Stonehouse 1967) when the bulk of feathers are synthesized via endogenous reserves (Cherel et al. 2005a, Brasso et al. 2013) . Finally, we examine Hg concentrations and stable isotope values in the tissues of four representative penguin prey species to provide additional predictive power and explain individual and group patterns of dietary Hg exposure in these penguins.

2. Methods

2.1 Study site and sample collection

We conducted fieldwork within the Antarctic Specially Protected Area (ASPA) no. 128 along the western shores of Admiralty Bay, King George Island, South Shetland Islands, Antarctica (62°10'S, 58°27'W). All three species of *Pygoscelis* penguins breed sympatrically at this location (Trivelpiece et al. 1987). We sampled penguins during December 2010, coinciding with the late incubation period for chinstrap penguins and the early brood period for Adélie and gentoo penguins. We captured 20-22 actively breeding adults per species and 20 non-breeding juvenile gentoo penguins (approximately one year old) identified by white head patches that did not reach the eye, incomplete white eye-rings and the lack of a brood patch (Trivelpiece et al. 1985). Individuals were sexed using a combination of molecular and morphometric methods (see Polito et al. 2012 for details). From each individual, we collected plucked breast feathers and whole blood samples (< 0.5 ml) by venipuncture of the interdigital vein. Blood samples were stored frozen (-20°C) prior to analysis.

In January 2009 we collected penguin prey samples during trawls conducted by the U.S. Antarctic Marine Living Resources (US AMLR program) around the South Shetland Islands and

Northern Antarctic Peninsula. Prey species were representative of the major components of penguin's diet at Admiralty Bay: Antarctic krill (*Euphausia superba*), epipelagic fish (*Pleuragramma antarcticum*), benthic fish (*Lepidonotothen squamifrons*) and mesopelagic myctophid fish (*Electrona antarctica*; Volkman et al. 1980, Karnovsky 1997). We collected a total of ten, 10 g samples of krill (approximately 20 individuals per sample) and ten whole fish per species and stored them frozen (-80°C) prior to analysis. While we could not assess to what effect differences in penguin and prey collection years may have had on our study, previous work suggest little inter-annual variability in the Hg and stable isotope values of *Pygoscelis* penguins and their prey (Polito et al. 2013, Brasso et al. 2014).

2.2 Mercury analysis

Prior to analysis, we cleaned adult feathers in a series of acetone and deionized water baths and allowed them to air dry under a fume hood for ~24 h. Whole prey samples were homogenized and freeze-dried to constant weight. We analyzed individual adult feathers and aliquots of whole blood and whole prey (~0.02 g) for total Hg via atomic absorption spectrometry on a Direct Mercury Analyzer DMA-80. Only one feather was analyzed from each individual as Hg concentrations do not vary significantly among body feathers in penguins (Brasso et al. 2013). Because nearly all Hg in blood, feathers and prey tissue is present in the form of methylmercury (except possibly in krill where percent methylmercury is unknown), a measurement of total Hg concentration was used as a proxy for this highly bioavailable form (Bloom 1992, Bond and Diamond 2009, Payne and Taylor 2010). Each set of 20 samples analyzed was preceded and followed by two method blanks, a sample blank, and two samples each of standard reference material (DORM-3, DOLT-4; fish protein, and dogfish liver certified

reference materials, respectively, provided by National Research Council Canada). All Hg concentrations are reported as parts per million (ppm); blood and prey values are presented as wet weight (ww), and feathers are reported as fresh weight (fw). As blood samples were analyzed as wet weight we did not assess the effect of hematocrit values on reported Hg concentrations. Mean percent recoveries for standard reference materials were $100.5 \pm 2.1\%$ (DORM-3) and $103.5 \pm 2.2\%$ (DOLT-4). The relative percent difference among duplicate samples was 2.1%. Detection limit of the assay was 0.005 ng Hg.

2.3 Stable isotope analysis

Prior to isotopic analysis, the tip (oldest part) of each feather was cut and discarded, as this section is grown at sea and may not reflect endogenous reserves (Cherel et al. 2005a). We cleaned feathers using a 2:1 chloroform : methanol rinse, air-dried under a fume hood for ~24 h and cut them into small fragments with stainless steel scissors. Prior to isotopic analysis whole blood samples and prey homogenates were freeze-dried to constant weight. We extracted lipids from prey homogenates using a Soxhlet apparatus with a 1:1 Petroleum-Ether: Ethyl-Ether solvent mixture for 8 h (Seminoff et al. 2007). Lipid-extracted krill were not acidified prior to isotopic analysis. Whole blood was not lipid-extracted prior to isotopic analysis as the lipid component of avian blood is less than 1% of the total wet mass and unlikely to be of sufficient magnitude to influence $\delta^{13}\text{C}$ values (see Bearhop et al. 2000b). Furthermore, the C:N ratios of penguin blood in our study ranged from 3.2-3.5 and was not correlated with $\delta^{13}\text{C}$ value ($R^2 = 0.01$, $P = 0.521$). We flash-combusted (Thermo-Finnigan & Costech ECS4010 elemental analyzers) approximately 0.5 mg of each whole blood, feather and prey sample loaded into tin cups and analyzed for carbon and nitrogen isotopes ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$) through interfaced Thermo

Finnigan Delta Plus XL and Delta V Plus continuous-flow stable isotope ratio mass spectrometers (CFIRMS). Raw δ values were normalized on a two-point scale using glutamic acid reference materials with low and high values (i.e. USGS-40 ($\delta^{13}\text{C} = -26.4\text{‰}$, $\delta^{15}\text{N} = -4.5\text{‰}$) and USGS-41 ($\delta^{13}\text{C} = 37.6\text{‰}$, $\delta^{15}\text{N} = 47.6\text{‰}$)). Sample precision based on repeated sample and reference material was 0.1‰ and 0.2‰, for $\delta^{13}\text{C}$, and $\delta^{15}\text{N}$, respectively.

Stable isotope ratios are expressed in δ notation in per mil units (‰), according to the following equation:

$$\delta X = [(R_{\text{sample}} / R_{\text{standard}}) - 1] \cdot 1000 \quad (1)$$

Where X is ^{13}C or ^{15}N and R is the corresponding ratio $^{13}\text{C} / ^{12}\text{C}$ or $^{15}\text{N} / ^{14}\text{N}$. The R_{standard} values were based on the Vienna PeeDee Belemnite (VPDB) for $\delta^{13}\text{C}$ and atmospheric N_2 for $\delta^{15}\text{N}$.

2.4 Penguin dietary analysis

We used the SIAR Bayesian multi-source isotopic mixing model in the R environment (R Development Core Team 2007) to quantify the diet composition of penguins in our study. The SIAR model estimates probability distributions of multiple source contributions to a mixture while accounting for the observed variability in source and mixture isotopic signatures, dietary isotopic fractionation, and elemental concentration (Parnell et al. 2010). We used the SIAR model with $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values from penguin and prey tissues to predict the relative contribution of krill, epipelagic fish, benthic fish and mesopelagic myctophid fish to the diets of penguins in our study. We incorporated diet to whole blood discrimination factors derived from captive studies of four piscivorous birds (see review by Cherel et al. 2005b; $\delta^{15}\text{N}$: $+2.7 \pm 0.4$; $\delta^{13}\text{C}$: $+0.0 \pm 0.7$) and diet to feather discrimination factor from a captive feeding study of *Pygoscelis* penguins ($\delta^{15}\text{N}$: $+3.5 \pm 0.4$; $\delta^{13}\text{C}$: $+1.3 \pm 0.5$; Polito et al. 2011b) and ran 1 million

iterations, thinned by 15, with an initial discard of the first 40,000 resulting in 64,000 posterior draws. Finally, we conducted a sensitivity analyses to examine the effect of variation in discrimination factors on mixing model results (e.g. Bond and Diamond 2011) by incorporating discrimination factors ranging 1‰ higher and lower than those estimated from the above captive feeding studies.

While isotopic mixing models provide a robust estimate of the relative amounts of fish and krill in penguin diets, they can have difficulty estimating the relative contribution of different fish species to penguin diets due to overlap in stable isotopes values (Polito et al. 2011a). Therefore we used fish otoliths recovered from penguin stomach contents to assess the fish composition of penguin diets at Admiralty Bay and complement our isotopic analyses. Direct dietary information was not available for the year or individuals sampled in our study, therefore we assess long-term differences in the fish composition of penguin diets at our study site using a 21-year dataset of penguin diets during the chick-rearing period at Admiralty Bay from 1988-89 to 2008-09. Detailed methods are described in Hinke et al. (2007), but briefly we collected stomach samples using gastric lavage from four to five breeding adults per species, replicated on 5-8 day intervals, resulting in approximately 20–40 samples per species per year and 1699 samples over the 21 year period (Adélie penguin = 619, chinstrap penguin = 549, gentoo penguin = 479). We recovered fish otoliths by swirling diet samples in a dark-bottomed pan and identified otoliths to the lowest possible taxonomic level and three habitat groupings (benthic, epipelagic and mesopelagic myctophids) using an internal reference collection and a published guide (Williams and McEldowney 1990, Karnovsky 1997). We calculated the frequency occurrence and the minimum number of individuals (MNI) of each fish taxa in each penguins species diet over this 21 year period following standard methods (Polito et al. 2002).

This method allowed us to assess the relative importance of epipelagic, benthic and mesopelagic fish to the diets of the adult Adélie, chinstrap and gentoo penguin populations at Admiralty Bay. Unfortunately, no data were available to help assess the fish component of juvenile gentoo penguin diets at Admiralty Bay.

2.5 Statistical Analysis

Statistical analysis was conducted using SAS (Version 9.1, SAS Institute 1999). Penguin and prey tissue Hg concentrations were log-transformed prior to analyses to generate data sets that did not differ from a normal distribution (Shapiro-Wilk, $P > 0.05$ in all cases). We used separate ANOVAs to test for differences among groups (adult Adélie penguins, adult chinstrap penguins, adult gentoo penguins, and juvenile gentoo penguins) and between sexes for penguin blood and feather Hg, $\delta^{13}\text{C}$, and $\delta^{15}\text{N}$ values. We used similar ANOVAs approach to examine differences in Hg, $\delta^{13}\text{C}$, and $\delta^{15}\text{N}$ values among penguin prey species. Tukey's HSD was used for all post-hoc comparisons. Relationships between blood and feather Hg, $\delta^{13}\text{C}$, and $\delta^{15}\text{N}$ values, and Hg concentrations and stable isotope values within tissue were tested using Pearson correlation.

We then used a generalized linear modeling approach with a normal distribution and an identity-link function to test multiple alternative hypotheses regarding the influence of species, age, sex, trophic position ($\delta^{15}\text{N}$), and foraging habitat ($\delta^{13}\text{C}$) on Hg concentrations in penguin blood and feathers. We conducted separate analyses by tissue for both adults of all three species as well as for gentoo penguins only to test for differences between age classes. We parameterized a global model for all adults using log transformed Hg as the dependent variable, species and sex as grouped factors, $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ as covariates, and all two-way interaction terms. We

parameterized a global model for gentoo penguins only using log transformed Hg as the dependent variable, age and sex as grouped factors, $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ as continuous variables, and all two-way interaction terms. Continuous variables ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$) that were significantly correlated were not included in the same models. All possible model subsets of our global model were compared using Akaike Information Criteria adjusted for small sample sizes (AIC_c ; Akaike 1973). The model with the lowest AIC_c score was selected as the most parsimonious model. Models with ΔAIC_c scores (difference in AIC_c between a given model and the model with the lowest AIC_c) ≤ 2.0 were interpreted to be equally competitive with the most parsimonious model and any model with a $\Delta\text{AIC}_c \leq 10.0$ was considered well supported. Model fits were further assessed by AIC weight (w_i) which is a measure of the relative likelihood that a given model is the best among a set of models fitted (Burnham and Anderson 2002). All means are presented $\pm$ SD and statistical significance was defined at $P < 0.05$.

3. Results

3.1 Hg concentrations in penguins and their prey

Penguin blood Hg concentration differed among penguin groups ($F_{3,83} = 3.79$, $P < 0.001$) but not between sexes ($F_{1,83} = 3.08$, $P = 0.083$) and did not exhibit a significant interaction between group and sex ($F_{3,83} = 2.73$, $P = 0.051$; Fig. 1). Similarly, penguin feather Hg concentrations differed among penguin groups ($F_{3,83} = 18.25$, $P < 0.00$) but not between sexes ($F_{1,83} = 0.64$, $P = 0.425$; Fig. 1), though there was a significant interaction between group and sex ($F_{3,83} = 3.62$, $P = 0.0169$). Inter-specific patterns were similar in both blood and feathers, with adult chinstrap penguins having the highest mean Hg concentration, followed by adult Adélie and gentoo penguins which were both significantly higher than juvenile gentoo penguins (Table

1). In addition, gentoo penguins tended to have the most variable blood and feather Hg concentrations. This was especially true of adult gentoo penguins which had ~3-5 times higher coefficient of variation (CV) relative to other penguins species (Table 1).

Penguin Hg concentrations were significantly and positively correlated between blood and feathers from the same individual ($r = 0.66$, $P < 0.001$). However, this trend appears to be driven primarily by adult gentoo penguins ($r = 0.65$, $P = 0.002$), as blood and feathers Hg concentrations were not significantly correlated within other penguin groups (All $P > 0.8$).

Hg concentrations also differed among representative penguin prey species ($F_{3,83} = 9.36$, $P < 0.001$). Antarctic krill (*E. superba*) Hg concentrations were significantly lower than all three fish species (Table 2). In addition, epipelagic fish (*P. antarcticum*) had lower Hg concentrations relative to benthic fish (*L. squamifrons*) and myctophid (*E. antarctica*) fish species (Table 2).

3.2 Stable isotope values in penguins and their prey

Penguin blood $\delta^{13}\text{C}$ values differed among penguin groups ($F_{3,83} = 3.79$, $P = 0.014$), but not between sexes ($F_{1,83} = 3.45$, $P = 0.067$) and did not exhibit a significant interaction between group and sex ($F_{3,83} = 0.66$, $P = 0.582$; Fig. 2). Significant differences between penguin groups were driven by higher blood $\delta^{13}\text{C}$ values found adult Adélie penguins relative to chinstrap penguins (Table 1). Similarly, penguin feather $\delta^{13}\text{C}$ values differed among penguin groups ($F_{3,83} = 4.36$, $P = 0.007$), but not between sexes ($F_{1,83} = 0.63$, $P = 0.429$) and did not exhibit a significant interaction between group and sex ($F_{3,83} = 1.02$, $P = 0.387$; Fig. 2). Significant differences between penguin groups were driven by higher feather $\delta^{13}\text{C}$ values found in adult chinstrap penguins and juvenile gentoo penguin, relative to adult Adélie penguins (Table 1).

gentoo penguins tended to have the most variable blood and feather $\delta^{13}\text{C}$ values in comparison to other species (Table 1).

Penguin blood $\delta^{15}\text{N}$ values differed among penguin groups ($F_{3,83} = 14.07$, $P < 0.001$), between sexes ($F_{1,83} = 16.99$, $P < 0.001$) and exhibited a significant interaction between group and sex ($F_{3,83} = 4.40$, $P = 0.007$; Fig. 2). Significant differences between penguin groups were driven by higher blood $\delta^{15}\text{N}$ values found in adult Adélie and chinstrap penguins relative to juvenile gentoo penguins (Table 1). While statically significant, the difference between males and females in all species combined ($+0.2\text{‰}$) was equal to our level of analytical precision for $\delta^{15}\text{N}$. In addition, significant differences between sexes were found only in adult Adélie and gentoo penguins, with males having higher blood $\delta^{15}\text{N}$ values in both cases (Fig. 2). Penguin feather $\delta^{15}\text{N}$ values differed among penguin groups ($F_{3,83} = 5.14$, $P = 0.003$), but not between sexes ($F_{1,83} = 1.92$, $P = 0.170$) and did not exhibit a significant interaction between group and sex ($F_{3,83} = 0.92$, $P = 0.434$; Fig. 2). Significant differences between penguin groups were driven by higher blood $\delta^{15}\text{N}$ values found in adult gentoo penguins relative to juvenile gentoo penguins (Table 1). Similar to Hg concentration and $\delta^{13}\text{C}$ values, gentoo penguins tended to have the most variable blood and feather $\delta^{15}\text{N}$ values in comparison to other species (Table 1).

Penguin stable isotope values were significantly and positively correlated between blood and feathers from the same individual for $\delta^{15}\text{N}$ ($r = 0.51$, $P < 0.001$), but not $\delta^{13}\text{C}$ ($P = 0.720$). Within penguin groups the only significant correlations found between tissue stable isotope values were $\delta^{13}\text{C}$ in adult gentoo penguins ($r = 0.63$, $P = 0.002$) and $\delta^{15}\text{N}$ for adult ($r = 0.82$, $P < 0.001$) and juvenile ($r = 0.49$, $P = 0.025$) gentoo penguins (all other comparisons $P > 0.121$). Penguin tissue $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values were also significantly and positively correlated with each other for both blood ($r = 0.49$, $P < 0.001$) and feathers ($r = 0.32$, $P = 0.003$). Within penguin

groups the only significant correlations between tissue $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values were adult (Blood: $r = 0.94$, $P < 0.001$; Feather: $r = 0.55$, $P = 0.009$) and juvenile (Blood: $r = 0.70$, $P < 0.001$; Feather: $r = 0.82$, $P < 0.001$) gentoo penguins (all other comparisons $P > 0.152$).

Prey stable isotope values differed across taxa for both $\delta^{13}\text{C}$ ($F_{3,83} = 22.09$, $P < 0.001$) and $\delta^{15}\text{N}$ ($F_{3,83} = 239.31$, $P < 0.001$). Krill had lower $\delta^{13}\text{C}$ values than both epipelagic fish and benthic fish and $\delta^{15}\text{N}$ values that were lower than all three fish species (Table 2). Within fish species $\delta^{13}\text{C}$ values differed between all species, with epipelagic fish having the lowest values and benthic fish having the highest values (Table 2). Epipelagic and mesopelagic myctophid fish did not differ in $\delta^{15}\text{N}$ values, though both species had lower $\delta^{15}\text{N}$ values relative to benthic fish (Table 2).

3.3 Penguin diets at Admiralty Bay

Isotopic mixing models using discrimination factors estimated from captive feeding studies predicted that krill was more important than fish prey in the diet of penguins at Admiralty Bay with mean estimates ranging from 54.2 to 65.8% of the diet depending on species, age and the type of tissue examined (Fig. 3). However, isotopic mixing models had difficulty separating the fish component of penguin diets among epipelagic, mesopelagic and benthic fishes with a higher degree of overlap in 95% credibility intervals (CI) in isotopic models using both blood and feather values (Fig. 3). One exception to this generalization is the model using blood isotopic values that predicted a higher mean contribution of benthic fishes (30.0%; CI: 12.5-60.7%) relative to mesopelagic fishes (4.7%; CI: 0.0-11.6%) in the diets of adult gentoo penguins at Admiralty Bay. Predicted contributions of the proportion of krill vs. other prey (i.e. fish) were sensitive to variation in discrimination factors with changes of ~15% per 1‰ shift in

discrimination factors (Table S1 of the Supplementary Material). However, differences in the relative importance of krill were consistent among penguin groups across all discrimination factors employed and all models had similar difficulty separating the fish component of penguin diets. Therefore, we restrict our further discussion to mixing models that employed the discrimination factors estimated from captive feeding studies.

From 1988-89 to 2008-09 a total of 1,996 otoliths were recovered from 1,699 stomach samples with 93.6% of otoliths identifiable to taxa in three habitat groups (epipelagic, mesopelagic, benthic fishes). Using this dataset we found strong preferences in the types of fish consumed by each penguin species during the chick rearing period. Epipelagic fish were the most common fish found in adult Adélie penguin diets by frequency occurrence and percentage of individuals (Fig. 4). In contrast, mesopelagic fish were most common in adult chinstrap penguin diets and benthic fish were the most common in adult gentoo penguin diets. Otolith data also validated our use of representative epipelagic, mesopelagic, benthic fish taxa for Hg and stable isotope analysis as *P. antarcticum* (47.4%), *E. antarctica* (60.0%), and *Lepidonotothen* sp. (28.26%) were the single most common fish prey found in adult Adélie, chinstrap and gentoo penguin stomach samples respectively.

3.4 Influence of species, age, sex, and foraging ecology on penguin Hg concentrations

When examined across all species, sexes and ages, feathers and blood Hg concentrations were significantly and positively correlated with $\delta^{15}\text{N}$ values, but not $\delta^{13}\text{C}$ values (Fig. 5). When examining only adults, based on AIC_c analyses the most parsimonious model found to explain variation in both feathers and blood Hg concentrations included species, $\delta^{15}\text{N}$ and their interaction (species* $\delta^{15}\text{N}$) as covariates (Table S2 of the Supplementary Material). For both

feathers and blood, Hg concentrations differed across species and while they were not significantly correlated with $\delta^{15}\text{N}$ values in adults only, a significant species* $\delta^{15}\text{N}$ interaction was found (Table 3). This interaction was a result of significant positive correlations between Hg concentrations and $\delta^{15}\text{N}$ values in adult gentoo penguin feathers ($r = 0.75$, $P < 0.001$) and blood ($r = 0.63$, $P = 0.002$) samples and the lack of relationship between these parameters in adult Adélie and chinstrap penguins (all $P > 0.450$). While no other competitive models (i.e., $\Delta\text{AIC}_c \leq 2.0$) were found to predict adult feather Hg concentration, a model that included an effect of species, $\delta^{13}\text{C}$ and their interaction (species* $\delta^{13}\text{C}$) was competitive in explaining variation in adult blood Hg concentration ($\Delta\text{AIC}_c = 1.7$; Table S2). This alternate model for adult blood provided similar results, with a significant effect of species ($F_{2,63} = 4.61$, $P = 0.014$), a non-significant effect of $\delta^{13}\text{C}$ values ($F_{2,63} = 2.61$, $P = 0.1120$), and a significant species* $\delta^{13}\text{C}$ ($F_{2,63} = 4.75$, $P = 0.012$) due to positive correlations between blood Hg concentrations and $\delta^{13}\text{C}$ values in adult gentoo penguin feathers ($r = 0.60$, $P = 0.004$) but not other species (all $P > 0.319$). Even so AIC_c weights illustrated that the model using blood $\delta^{15}\text{N}$ values had twice as much support as the model using blood $\delta^{13}\text{C}$ values as a covariate (Table S2).

When examining feathers Hg concentrations in adult and juvenile gentoo penguins the model containing only $\delta^{15}\text{N}$ as a covariate was selected as the most parsimonious with no other competitive models found (Table S3 of the Supplementary Material). Gentoo penguin feathers Hg concentrations were positively correlated with $\delta^{15}\text{N}$ values both across (Table 3) and within age classes (Adults: $r = 0.75$, $P < 0.001$; Juveniles: $r = 0.81$, $P < 0.001$). The most parsimonious model explaining variation in gentoo penguin blood Hg concentrations included age, $\delta^{15}\text{N}$ and their interaction (age* $\delta^{15}\text{N}$) as covariates (Table S3). The only significant term in this model was the age* $\delta^{15}\text{N}$ interaction, because adult blood Hg concentrations were positively correlated

with $\delta^{15}\text{N}$ values ($r = 0.63$, $P = 0.002$), but not juvenile gentoo penguin ($r = -0.18$, $P = 0.431$). A model that included the above three covariates as well two others (sex, sex* $\delta^{15}\text{N}$) was also competitive ($\Delta\text{AIC}_c = 1.6$; Table S3), but resulted in all model covariates being non-significant ($P > 0.279$).

4. Discussion

We found that tissue Hg concentrations differed across penguin species, between adults and juveniles, but not between sexes. Adult chinstrap penguins had the highest mean tissue Hg concentrations followed in descending order by adult Adélie penguins, adult gentoo penguins and juvenile gentoo penguins. Interestingly, while tissue Hg concentrations across all species and age classes were significantly and positively correlated with $\delta^{15}\text{N}$ values, these relationships were driven exclusively by adult gentoo penguins which also had the most variable tissue Hg concentrations and stable isotope values. Furthermore, similar to previous studies (Brasso et al. 2012, Brasso et al. 2014) the higher tissue Hg concentrations found in adult chinstrap penguins could not be explained using $\delta^{15}\text{N}$ values, a proxy for trophic position, or by the stable isotope-based dietary mixing models used in this study. However, integrating data from penguin stomach content analysis with prey species Hg concentrations and stable isotope values revealed that the differences in epipelagic, benthic and mesopelagic prey selection by penguins are stronger drivers of Hg exposure, relative to trophic position alone.

4.1 Interspecific trends in penguin foraging ecology and Hg exposure

While all three species of *Pygoscelis* penguin diets are dominated by Antarctic krill in the Antarctic Peninsula region (Volkmann et al. 1980), previous studies have found a general pattern

of foraging niche partitioning of these species during the breeding season (Trivelpiece et al 1987, Lynnes et al. 2002). This occurs via slight differences in diet, foraging area, and depth utilization (Kokubun et al. 2010, Miller et al. 2010, Wilson 2010). For example, past stable isotope and conventional studies at King George Island and sites in the Antarctic Peninsula indicate higher krill consumption and greater use of offshore foraging habitats by Adélie and chinstrap penguin relative to gentoo penguins (Trivelpiece et al 1987, Miller and Trivelpiece 2008, Miller et al. 2010, Polito et al. 2015). The results of our otolith analysis agree with these previous studies and help to identify the differential use of epipelagic, mesopelagic, and benthic foraging habitats across the three species of *Pygoscelis* penguins.

Interestingly, the foraging habitat differences identified via otolith analysis were not as clearly delineated in the stable isotope values of penguin tissues. Isotopic mixing models predictions of the fish component of penguins diets overlapped broadly and only were able to predict a higher mean contribution of benthic fishes relative to mesopelagic fishes in the diets of adult gentoo penguins. Stable isotope-based dietary mixing models in this and other studies provided a robust estimate of the relative amounts of fish and krill in penguin diets due to their large difference in $\delta^{15}\text{N}$ values (e.g. Polito et al. 2011a, Polito et al. 2015). However, these models have difficulty estimating the relative dietary contribution of different fish species to penguin diets due to a high degree of overlap in the stable isotope values of epipelagic, mesopelagic, and benthic fishes around Antarctica (Cherel et al. 2010, Cherel et al. 2011).

Differences in the diets and foraging ecology of *Pygoscelis* penguin species appear to have significant implication for their relative exposure to Hg in the Antarctic marine food web. For example, adult gentoo penguins often have a more mixed diet of krill and fish relative to their congeners (Trivelpiece et al. 1987) and this has the potential to lead to greater Hg exposure

due their higher trophic level of diet (Carravieri et al. 2013, Brasso et al. 2015). However, while differences in relative krill vs. fish consumption has been observed in previous studies (Miller et al. 2010, Polito et al. 2015), we found no evidence of differences in trophic level (i.e. $\delta^{15}\text{N}$ values in blood or feathers) or predicted krill consumption (from isotopic mixing models) among adult Adélie, chinstrap and gentoo penguins at Admiralty Bay. Therefore the differences in Hg exposure among adult *Pygoscelis* penguins in our study are not likely related to difference in level of krill consumption or trophic levels *per se*. Instead, the observed differences in penguin species relative selection of epipelagic, mesopelagic, and benthic prey are likely stronger drivers of the observed differences in tissue Hg concentrations across *Pygoscelis* penguin species.

For example, we found that while adult Adélie, gentoo and chinstrap penguins shared a similar trophic level of diet (based on $\delta^{15}\text{N}$), chinstrap penguins have relatively higher feather and blood tissue Hg concentrations. In addition, past studies suggest that this pattern of chinstrap penguins having higher Hg concentrations than Adélie and gentoo penguins is inter-annually and seasonally consistent across much the Antarctic Peninsula region (Brasso et al. 2012, Brasso et al. 2014). While not tested until now, these past studies hypothesized that elevated Hg concentrations in chinstrap penguin tissues may result from exploiting mesopelagic prey with higher Hg concentrations rather than foraging at a higher trophic level than their congeners (Brasso et al. 2012, Brasso et al. 2014, Brasso et al. 2015). Marine predators that forage in the mesopelagic realm or on mesopelagic prey have been documented to have elevated exposure to Hg compared to their trophically similar, epipelagic-feeding counterparts due to enhanced bioaccumulation of Hg at depth (Montiero et al. 1996, Montiero and Furness 1997, Choy et al. 2009). We found strong evidence to support this hypothesis when comparing tissue Hg concentrations and stable isotope values across the three most common epipelagic, mesopelagic,

and benthic fish species found in the diets of *Pygoscelis* penguins. Specifically, while epipelagic and mesopelagic myctophid fish shared similar $\delta^{15}\text{N}$ values, mesopelagic fish had higher Hg concentrations relative to epipelagic fishes. In addition, while mesopelagic fish occupied a slightly lower $\delta^{15}\text{N}$ values than benthic fish, these two prey species have similar Hg concentrations. These results substantiate the relatively higher risk of Hg exposure for chinstrap penguins foraging on mesopelagic fish relative to congeners foraging at a similar trophic level but targeting epipelagic or benthic prey species.

4.2 *Intraspecific trends in penguin foraging ecology and Hg exposure*

Relationships between feather and blood tissue Hg concentration and stable isotope values provide additional insights into seasonal and individual variation in penguin foraging ecology and Hg exposure. Whole blood reflects the short-term Hg exposure incorporated from food during blood formation as well as some component of residual Hg residing in other tissues (Bearhop et al. 2000a,c). Similarly, the stable isotope values of whole blood in a penguin 3-5kg penguin integrate short-term dietary information over a period 20-60 days prior to collection (Carleton and Martinez del Rio 2005, Barquete et al. 2013). Therefore the adult and juvenile whole blood samples examined in this study provide insight into species foraging ecology and Hg exposure during a short time period during the late incubation and the early chick-rearing periods. In contrast, the stable isotope values and Hg concentrations in adult penguin feathers reflect an integrated signal of endogenous tissue Hg and diets over the inter-molt period prior to the annual catastrophic molt in which all body feathers are lost and re-grown over a period of 2-3 weeks while fasting (Stonehouse 1967, Brasso et al. 2013). Furthermore, as non-breeding juvenile gentoo penguins (one year olds) have not yet completed their first adult molt, their

feather stable isotope values reflect an integrated dietary history and Hg exposure from the food parents provide them during feather synthesis prior to fledgling (Hobson and Clark 1992, Tierney et al. 2008, Polito et al. 2011a).

When examined across all species and age classes, tissue Hg concentrations and stable isotope values were significantly and positively correlated between blood (short-term signal) and feathers (long-term signal) from the same individual. This result suggests some degree of seasonal consistency in the diets and foraging ecology and Hg exposure at the individual level in *Pygoscelis* penguins. However, this global trend was driven primarily by adult and, to a lesser extent, juvenile gentoo penguins. Gentoo penguins also tended to have the most variable blood and feather Hg concentration and stable isotope values in comparison to other species. This high degree of individual variation in Hg exposure reflects similar individual variation in trophic position ($\delta^{15}\text{N}$) as individuals foraging at higher trophic positions also had higher Hg concentrations. Two previous studies, on Antarctic (Brasso et al. 2014) and sub-Antarctic (Carravieri et al. 2013) islands also have found that individual variation in trophic position led to increased variation in Hg concentrations in gentoo penguins at the population level.

Previous studies indicate the potential for sexual differences in diets in gentoo penguins, with larger males diving deeper and consuming a higher proportion of fish than females (Volkman et al. 1980, Bearhop et al. 2006, Miller et al. 2010). Such sexual differences in diets, when they occur, might be expected to lead to differential exposure to Hg between males and females (Becker et al. 2002). For example, Pedro et al. (2015) found higher Hg exposure in males relative to females as well as higher Hg exposure in individuals feeding on a higher proportion of fish. In contrast, we found little evidence for differences in stable isotope values between sexes within species and no evidence of differences in tissue Hg concentration.

However, differing results across studies might be expected due to temporal and/or spatial variation in prey and foraging habitat availability that mediate the degree and consistency of sexual variation in *Pygoscelis* penguin diets as observed in other seabird species (Miller et al. 2010, Phillips et al. 2011).

While no sexual differences were apparent, we did observe differences in tissue Hg concentrations between age classes of gentoo penguins in our study. Both blood and feather Hg concentrations were significantly higher in adults relative to juveniles. However, these differences do not seem to be entirely age-related *per se* as differences in Hg exposure between age classes was influenced primarily by the generally higher and more variable trophic position ($\delta^{15}\text{N}$) occupied by adults relative to juveniles. As such, feather $\delta^{15}\text{N}$ values were the strongest predictor of feather Hg concentration regardless of age class, and age class differences in blood Hg concentrations were no longer significant once $\delta^{15}\text{N}$ values were taken into account. This indicates that individual gentoo penguins feeding on higher trophic levels may be at greater risk of Hg exposure regardless of their age.

4.3 Conclusions and implications for biomonitoring

The tissue Hg concentrations observed in *Pygoscelis* penguins in this study were comparatively low relative to those found in other studies of seabirds in the Southern Hemisphere (Anderson et al. 2009, Blévin et al. 2013, Carravieri et al. 2013, Brasso et al. 2015). Hg concentrations in feathers averaged <0.50 ppm and those in blood samples averaged <0.10 ppm. In comparison, adverse effects in terms of reproductive impairment in other aquatic and marine birds have been reported at feather Hg concentrations between 5.0-40.0 ppm and blood Hg concentrations > 3.0 ppm (Wolfe et al. 1998, Burger and Gochfeld 2000, Evers et al. 2008). The highest feather and blood Hg concentrations observed in this study from an individual

gentoo penguin and were 1.98 ppm and 0.34 ppm, respectively. Thus, it is unlikely that *Pygoscelis* penguins in our study or other areas of the Antarctic (e.g. Brasso et al. 2015) are experiencing negative impacts from Hg exposure.

While the potential for adverse effects due to Hg exposure is currently low in *Pygoscelis* penguins, this study provides important baseline data that will aid in the design and interpretation of future studies seeking to use *Pygoscelis* penguins as biomonitors for Hg availability in the Antarctic marine ecosystem. Specifically this study highlights how inter- and, to a lesser extent, intraspecific variation in foraging habitats and diet influences a species relative exposure to Hg. In addition, our findings indicate that stable isotope analysis alone was not sufficiently robust to identify the differences in penguin foraging ecology that most influenced Hg exposure. It was only by combining isotopic analyses with data from stomach content analysis that we were able to identify the key differences in foraging habitat and prey selection among *Pygoscelis* penguins acting as strong drivers of Hg exposure. Furthermore, this approach allowed us to substantiate for the first time, the relatively higher risk of Hg exposure for chinstrap penguins foraging in mesopelagic habitats relative to congeners targeting epipelagic or benthic prey species. Therefore, in future studies of Hg exposure in penguins and other seabirds we recommend combining isotopic approaches with other, independent measures of diets and foraging habitat to help account for the potentially confounding effects of inter and intraspecific variation in foraging ecologies.

5. Acknowledgements

Many thanks to the US-AMLR program especially S. Trivelpiece, G. Watters, C. Jones, C. Reiss, A. Will, P. Chilton and K. Boysen and the many other researchers who have worked at

553 the Copacabana field camp who helped gather data used in this study. We also thank Raytheon
554 Polar Services, the Laurence M. Gould, the M/V Ushuaia, John Evans, Melissa Rider and Ethan
555 Norris for transportation and logistical support. D. Besic, B. Drummond, K. Durenberger, T.
556 Prokopiuk, J. Seminoff and E. Unger provided assistance with sample preparation and stable
557 isotope analysis. Animal use in this study was conducted under approved animal use protocols
558 from the University of North Carolina Wilmington Animal Care and Use Committee (A0910-20)
559 and in accordance to Antarctic Conservation Act permits provided by the U.S. National Science
560 Foundation (NSF) to S. Emslie (2006-001) and G. Watters (2011-005). Funding was provided in
561 part by a NSF Office of Polar Programs grant to S. Emslie (ANT-0739575).

6. References

- Ackerman JT, Eagles-Smith CA, Takekawa JY, Bluso JD, Adelsbach TL. 2008. Mercury concentrations in blood and feathers of prebreeding forster's tern in relation to space use of San Francisco Bay, California, USA, habitats. *Environmental Toxicology and Chemistry* 27, 897-908.
- Akaike, H., 1973. Information theory and an extension of the maximum likelihood principle. In: Petrov BN, Csaki F (eds.) *Proceedings of the Second International Symposium on Information Theory*. Akademiai Kiado, Budapest, pg 267-281.
- Anderson, O.R.J., Phillips, R.A., McDonald, R.A., Shore, R.F., McGill, R.A.R., Bearhop, S. 2009. Influence of trophic position and foraging range on mercury levels within a seabird community. *Marine Ecology Progress Series* 375, 277–288.
- Atwell, L., Hobson, K.A., Welch, H.E. 1998. Biomagnification and bioaccumulation of mercury in an arctic marine food web: insights from stable nitrogen isotope analysis. *Canadian Journal of Fisheries and Aquatic Sciences* 55(5), 1114-1121.
- Barquete, V., Strauss, V., Ryan, P.G., 2013. Stable isotope turnover in blood and claws: a case study in captive African penguins. *Journal of Experimental Marine Biology and Ecology* 448, 121-127.
- Bearhop, S., Ruxton, G. D., Furness, R.W. 2000a. Dynamics of mercury in blood and feathers of great skuas. *Environmental Toxicology and Chemistry* 19(6), 1638-1643.
- Bearhop, S., Teece, M.A., Waldron, S., Furness, R.W. 2000b. Influence of lipid and uric acid on $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of avian blood: implications for trophic studies. *The Auk* 117(2), 504-507.

584 Bearhop, S., Waldron, S., Thompson, D., Furness, R., 2000c. Bioamplification of mercury in
 585 great skua *Catharacta skua* chicks: the influence of trophic status as determined by stable
 586 isotope signatures of blood and feathers. *Marine Pollution Bulletin* 40(2), 181-185.

587 Bearhop, S., Phillips, R.A., McGill, R., Cherel, Y., Dawson, D.A., Croxall, J.P. 2006. Stable
 588 isotopes indicate sex-specific and long-term individual foraging specialisation in diving
 589 seabirds. *Marine Ecology Progress Series* 311, 157-164.

590 Becker, P.H., González-solís, J., Behrends, B., Croxall, J., 2002. Feather mercury levels in
 591 seabirds at South Georgia: influence of trophic position, sex and age. *Marine Ecology*
 592 *Progress Series* 243:261-269.

593 Blevin, P., Carravieri, A., Jaeger, A., Chastel, O., Bustamante, P., Cherel, Y. 2013. Wide range
 594 of mercury contamination in chicks of Southern Ocean seabirds. *PLoS One* 8(1), e54508.

595 Bloom, N.S., 1992. On the chemical form of mercury in edible fish and marine invertebrate
 596 tissue. *Canadian Journal of Fisheries and Aquatic Science* 49, 1010-1017.

597 Bond, A.L., Diamond, A.W. 2009. Total and methyl mercury concentrations in seabird feathers
 598 and eggs. *Archives of Environmental Toxicology Chemistry* 56, 286-291.

599 Bond, A.L., Diamond, A.W. 2011. Recent Bayesian stable-isotope mixing models are highly
 600 sensitive to variation in discrimination factors. *Ecological Applications* 21(4), 1017-1023.

601 Bond, A.L., Jones, I.L., 2009. A practical introduction to stable-isotope analysis for seabird
 602 biologists: approaches, cautions and caveats. *Marine Ornithology* 37(183), e188.

603 Brasso, R.L., Cristol, D.A., 2008. Effects of mercury exposure on the reproductive success of
 604 tree swallows (*Tachycineta bicolor*). *Ecotoxicology* 17, 133-141.

605 Brasso, R.L., Polito, M.J., 2013. Trophic calculations reveal the mechanism of population-level
606 variation in mercury concentrations between marine ecosystems: Case studies of two
607 polar seabirds. *Marine Pollution Bulletin* 75, 244-249.

608 Brasso, R.L., Polito, M.J., Lynch, H.J., Naveen, R., Emslie, S., 2012. Penguin eggshell
609 membranes reflect homogeneity of mercury in the marine food web surrounding the
610 Antarctic Peninsula. *Science of the Total Environment* 439, 165-171.

611 Brasso, R.L., Drummond, B.E., Borrett, S.R., Chiaradia, A., Polito, M.J., Raya Rey, A., 2013.
612 Unique pattern of molt leads to low intraindividual variation in feather mercury
613 concentrations in penguins. *Environmental Toxicology and Chemistry* 32, 2331-2334.

614 Brasso, R.L., Polito, M.J., Emslie, S.D., 2014. Multi-tissue analyses reveal limited inter-annual
615 and seasonal variation in mercury exposure in an Antarctic penguin community.
616 *Ecotoxicology* 23(8), 1494-1504.

617 Brasso, R. L., Chiaradia, A., Polito, M. J., Rey, A. R., Emslie, S. D. 2015. A comprehensive
618 assessment of mercury exposure in penguin populations throughout the Southern
619 Hemisphere: Using trophic calculations to identify sources of population-level variation.
620 *Marine Pollution Bulletin* 97:408-418

621 Burger, J., Gochfeld, M. 2000 Metal levels in feathers of 12 species of seabirds from Midway
622 Atoll in the northern Pacific Ocean. *Science of the Total Environment* 257, 37-52.

623 Burnham, K.P., Anderson, D.R., 2002. Model selection and multimodel inference: a practical
624 information-theoretic approach. Springer Science & Business Media.

625 Carleton, S.A., del Rio, C.M. 2005. The effect of cold-induced increased metabolic rate on the
626 rate of ^{13}C and ^{15}N incorporation in house sparrows (*Passer domesticus*). *Oecologia*
627 144(2), 226-232.

628 Carravieri, A., Bustamante, P., Churlaud, C., Cherel, Y., 2013. Penguins as bioindicators of
629 mercury contamination in the Southern Ocean: birds from the Kerguelen Islands as a case
630 study. *Science of the Total Environment* 454, 141-148.

631 Carravieri, A., Cherel, Y., Blévin, P., Brault-Favrou, M., Chastel, O., Bustamante, P., 2014.
632 Mercury exposure in a large subantarctic avian community. *Environmental Pollution* 190,
633 51-57.

634 Celis, J.E., Espejo, W., Barra, R., Gonzalez-Acuña, D., Gonzalez, F., Jara, S., 2015. Assessment
635 of trace metals in droppings of Adélie penguins (*Pygoscelis adeliae*) from different
636 locations of the Antarctic Peninsula area. *Advances in Polar Science* 26, 1-7.

637 Cherel, Y., Hobson, K.A., 2007. Geographical variation in carbon stable isotope signatures of
638 marine predators: a tool to investigate their foraging areas in the Southern Ocean. *Marine*
639 *Ecology Progress Series* 329, 281–287.

640 Cherel, Y., Hobson, K. A., Bailleul, F., Groscolas, R., 2005a. Nutrition, physiology, and stable
641 isotopes: new information from fasting and molting penguins. *Ecology* 86(11), 2881-
642 2888.

643 Cherel, Y., Hobson, K.A., Hassani, S. 2005b. Isotopic discrimination between food and blood
644 and feathers of captive penguins: implications for dietary studies in the wild.
645 *Physiological and Biochemical Zoology* 78(1), 106-115.

646 Cherel, Y., Fontaine, C., Richard, P., Labatc, J.P., 2010. Isotopic niches and trophic levels of
647 myctophid fishes and their predators in the Southern Ocean. *Limnology and*
648 *Oceanography* 55(1), 324-332.

649 Cherel, Y., Koubbi, P., Giraldo, C., Penot, F, Tavernier, E., Moteki, M., Ozouf-Costaz, C.,
650 Causse, R., Chartier, A., Hosie, G., 2011. Isotopic niches of fishes in coastal, neritic and
651 oceanic waters off Adélie land, Antarctica. *Polar Science* 5, 286-297.

652 Choy, C.A., Popp, B.N., Kaneko, J., Drazen, J.C. 2009. The influence of depth of mercury levels
653 in pelagic fishes and their prey. *Proceedings of the National Academy of Sciences* 106,
654 13865-13869.

655 Day, R.D., Roseneau, D.G., Vander Pol, S.S., Hobson, K.A., Donard, O.F., Pugh, R.S., ...
656 Becker, P.R., 2012. Regional, temporal, and species patterns of mercury in Alaskan
657 seabird eggs: Mercury sources and cycling or food web effects? *Environmental Pollution*
658 166, 226-232.

659 Evers, D.C., Burgess, N.M., Champoux, L., Hoskins, B., Major, A., Goodale, W.M., Taylor,
660 R.J., Poppenga, R., Daigle, T. 2005. Patterns and interpretation of mercury exposure in
661 freshwater avian communities in northeastern North America. *Ecotoxicology* 14: 193-221

662 Evers, D.C., Savoy, L.J., DeSorbo, C.R., Yates, D.E., Hanson, W., Taylor, K.M., Siegel, L.S.,
663 Cooley, J.H., Bank, M.S., Major, A., Munney, K., Mower, B.F., Vogel, H.S., Schoch, N.,
664 Pokras, M., Goodale, M.W., Fair, J., 2008. Adverse effects from environmental mercury
665 loads on breeding common loons. *Ecotoxicology* 17, 69-81

666 Hipfner, J.M., Hobson, K.A., Elliott, J.E. 2011. Ecological factors differentially affect mercury
667 levels in two species of sympatric marine birds of the North Pacific. *Science of the Total*
668 *Environment* 409(7), 1328-1335.

669 Hinke, J.T., Salwicka, K., Trivelpiece, S.G., Watters, G.M., Trivelpiece, W.Z., 2007. Divergent
670 responses of *Pygoscelis* penguins reveal a common environmental driver. *Oecologia*
671 153(4), 845-855.

672 Hobson K.A., Clark, R.G., 1992. Assessing avian diets using stable isotopes I: turnover of ^{13}C in
673 tissues. *Condor* 94, 181-188.

674 Jardine, T.D., Kidd, K.A., Fisk, A.T., 2006. Applications, considerations, and sources of
675 uncertainty when using stable isotope analysis in ecotoxicology. *Environmental Science*
676 *and Technology* 40, 7501-7511.

677 Karnovsky, N.J., 1997. The fish component of *Pygoscelis* penguin diets. Doctoral dissertation,
678 Montana State University, Bozeman.

679 Karnovsky, N.J., Hobson, K.A., Iverson, S.J., 2012. From lavage to lipids: estimating diets of
680 seabirds. *Marine Ecology Progress Series* 451, 263-284.

681 Kojadinovic, J., Bustamante, P., Churlaud, C., Cosson, R. P., Le Corre, M., 2007. Mercury in
682 seabird feathers: Insight on dietary habits and evidence for exposure levels in the western
683 Indian Ocean. *Science of the Total Environment* 384(1), 194-204.

684 Kokubun, N., Takahashi, A., Mori, Y., Watanabe, S., Shin, H.C., 2010. Comparison of diving
685 behavior and foraging habitat use between chinstrap and gentoo penguins breeding in the
686 South Shetland Islands, Antarctica. *Marine Biology* 157(4), 811-825.

687 Lynnes, A., Reid, K., Croxall, J., Trathan, P., 2002. Conflict or co-existence? Foraging
688 distribution and competition for prey between Adélie and Chinstrap Penguins. *Marine*
689 *Biology* 141(6), 1165-1174.

690 Miller, A.K., Trivelpiece, W.Z., 2008. Chinstrap penguins alter foraging and diving behavior in
691 response to the size of their principle prey, Antarctic krill. *Marine Biology* 154(2), 201-
692 208.

693 Miller, A.K., Kappes, M.A., Trivelpiece, S.G., Trivelpiece, W.Z., 2010. Foraging-niche
694 separation of breeding gentoo and chinstrap penguins, South Shetland Islands, Antarctica.
695 Condor 112(4), 683-695.

696 Montiero, L.R., Costa, V., Furness, R.W., Santos, R.S., 1996. Mercury concentrations in prey
697 fish indicate enhanced bioaccumulation in mesopelagic environments. Marine Ecology
698 Progress Series 141, 21-25.

699 Montiero, L.R., Furness, R.W., 1997. Accelerated increase in mercury contamination in north
700 Atlantic mesopelagic food chains as indicated by time series of seabird feathers.
701 Environmental Toxicology and Chemistry 16, 2489-2493.

702 Payne, E.J., Taylor, D.L., 2010. Effects of diet composition and trophic structure on mercury
703 bioaccumulation in temperate flatfishes. Archives of Environmental Contaminants and
704 Toxicology 58, 431-443.

705 Parnell, A.C., Inger, R., Bearhop, S., Jackson, A.L., 2010. Source partitioning using stable
706 isotopes: coping with too much variation. PloS One, 5(3), e9672.

707 Pedro, S., Xavier, J.C., Tavares, S., Trathan, P.N., Ratcliffe, N., Paiva, V.H., Medeiros, R.,
708 Vieira, R.P., Ceia, F.R., Pereira E., Pardal, M.A., 2015. Mercury accumulation in gentoo
709 penguins *Pygoscelis papua*: spatial, temporal and sexual intraspecific variations. Polar
710 Biology 38, 1335-1343.

711 Phillips, R.A., McGill, R.A., Dawson, D.A., Bearhop, S., 2011. Sexual segregation in
712 distribution, diet and trophic level of seabirds: insights from stable isotope analysis.
713 Marine Biology 158(10), 2199–2208.

714 Phillips, D.L., Newsome, S.D., Gregg, J.W., 2005. Combining sources in stable isotope mixing
715 models: alternative methods. Oecologia 144(4), 520-527.

716 Polito, M., Emslie, S.D., Walker, W., 2002. A 1,000-year record of Adélie penguin diets in the
717 southern Ross Sea. *Antarctic Science* 14, 327-332.

718 Polito, M.J., Trivelpiece, W.Z., Karnovsky, N.J., Ng, E., Patterson, W.P., Emslie, S.D. 2011a.
719 Integrating stomach content and stable isotope analyses to quantify the diets of
720 Pygoscelid penguins. *PLoS ONE* 6(10)::e26642.

721 Polito, M.J., Abel, S., Tobias, C.R., Emslie, S.D., 2011b. Dietary isotopic discrimination in
722 gentoo penguin (*Pygoscelis papua*) feathers. *Polar Biology* 34(7), 1057-1063.

723 Polito, M.J., Clucas, G.V., Hart, T., Trivelpiece, W.Z., 2012. A simplified method of
724 determining the sex of *Pygoscelis* penguins using bill measurements. *Marine Ornithology*
725 40, 89-94.

726 Polito, M.J., Reiss, C.S., Trivelpiece, W.Z., Patterson, W.P., Emslie, S.D., 2013. Stable isotopes
727 identify an ontogenetic niche expansion in Antarctic krill (*Euphausia superba*) from the
728 South Shetland Islands, Antarctica. *Marine Biology* 160(6), 1311-1323.

729 Polito, M.J., Trivelpiece, W.Z., Patterson, W.P., Karnovsky, N.J., Reiss, C.S., Emslie, S.D.,
730 2015. Contrasting specialist and generalist patterns facilitate foraging niche partitioning
731 in sympatric populations of *Pygoscelis* penguins. *Marine Ecology Progress Series* 519,
732 221-237.

733 Pouilly, M., Rejas, D., Pérez, T., Duprey, J., Molina, C.I., Hubas, C., Guimarães, J.D., 2013.
734 Trophic structure and mercury biomagnification in tropical fish assemblages, Iténez
735 River, Bolivia. *PLoS ONE* 8, e65054.

736 Seminoff, J.A., Bjorndal, K.A., Bolten, A.B., 2007. Stable carbon and nitrogen isotope
737 discrimination and turnover in pond sliders *Trachemys scripta*: insights for trophic study
738 of freshwater turtles. *Copeia* 2007(3), 534-542.

739 Stonehouse, B., 1967. The general biology and thermal balances of penguins. *Advances in*
740 *Ecological Research* 4, 131-196.

741 Tierney, M., Southwell, C., Emmerson, L.M., Hindell, M.A., 2008. Evaluating and using stable-
742 isotope analysis to infer diet composition and foraging ecology of Adélie penguins
743 *Pygoscelis adeliae*. *Marine Ecology Progress Series* 355:297-307.

744 Trivelpiece, S.G., Trivelpiece, W.Z., Volkman, N.J., 1985. Plumage characteristics of juvenile
745 pygoscelid penguins. *Ibis*, 127(3), 378-380.

746 Trivelpiece, W.Z., Trivelpiece, S.G., Volkman, N.J., 1987. Ecological segregation of Adelie,
747 gentoo, and chinstrap penguins at King George Island, Antarctica. *Ecology*, 351-361.

748 Volkman, N.J., Presler, P., Trivelpiece, W., 1980. Diets of pygoscelid penguins at King George
749 Island, Antarctica. *Condor*, 373-378.

750 Wada, H., Cristol, D.A., McNabb, F.M.A., Hopkins, W.A., 2009. Suppressed adrenocortical
751 responses and thyroid hormone levels in birds near a mercury-contaminated river.
752 *Environmental Science and Technology*, 43, 6031-6038.

753 Williams, R., McEldowney, A., 1990. A guide to the fish otoliths from waters off the Australian
754 Antarctic Territory, Heard and Macquarie Islands. ANARE Res. Notes (Vol. 75),
755 Australian Antarctic Division, Australia

756 Wilson, R.P., 2010. Resource partitioning and niche hyper-volume overlap in free-living
757 Pygoscelid penguins. *Functional Ecology* 24(3), 646-657.

758 Wolfe, M.F., Schwarzbach, S., Sulaiman, R.A., 1998. Effects of mercury on wildlife: a
759 comprehensive review. *Environmental Toxicology and Chemistry* 17, 146-160.

Figure Captions

Figure 1. Total Hg concentration in blood ($\mu\text{g g}^{-1}$ ww) and feathers ($\mu\text{g g}^{-1}$ fw) from three penguin species (Adélie: ADPE; Chinstrap: CHPE; Gentoo: GEPE) at King George Island, Antarctica. Sample sizes provided below each bar; significant differences between sexes are indicated with asterisks (Tukey's HSD, $p < 0.05$).

Figure 2. Stable carbon ($\delta^{13}\text{C}$) and nitrogen ($\delta^{15}\text{N}$) isotope values (‰) in blood and feathers from three penguin species (Adélie: ADPE; Chinstrap: CHPE; Gentoo: GEPE) at King George Island, Antarctica. Sample sizes provided below each bar; significant differences between sexes are indicated with asterisks (Tukey's HSD, $p < 0.05$).

Figure 3. Predicted dietary composition of three penguin species (Adélie: ADPE; Chinstrap: CHPE; Gentoo: GEPE) at King George Island, Antarctica based on blood and feather stable isotope values and a multi-source Bayesian isotopic mixing model. Proportions are presented as mean $\pm$ Bayesian 95% credibility intervals.

Figure 4. The proportion of minimum number of individuals and frequency occurrence of epipelagic, mesopelagic and benthic fish species based on a 21 year dataset of fish otoliths recovered from adult stomach contents of three penguin species (Adélie: ADPE; Chinstrap: CHPE; Gentoo: GEPE) at King George Island, Antarctica.

Figure 5. Relationships between stable carbon ($\delta^{13}\text{C}$) and nitrogen ($\delta^{15}\text{N}$) isotope values (‰) and total Hg concentration in blood ($\mu\text{g g}^{-1}$ ww) and feathers ($\mu\text{g g}^{-1}$ fw) from three penguin species

783 (Adélie: ADPE; Chinstrap: CHPE; Gentoo: GEPE) at King George Island, Antarctica. Test
784 statistics reflect Pearson correlation analyses across all species and age groups.

Table 1. Mean±SD, range, and coefficient of variation (CV; %) of total Hg concentration ($\mu\text{g g}^{-1}$ ww or fw) and stable carbon ($\delta^{13}\text{C}$) and nitrogen ($\delta^{15}\text{N}$) isotope values (‰) in blood and feathers from three penguin species at King George Island, Antarctica. Within each tissue, means not sharing a letter were found to differ significantly (Tukey's HSD, $p<0.05$)

Tissue	Penguin species (age)	<i>n</i>	Hg		$\delta^{13}\text{C}$		$\delta^{15}\text{N}$	
			Concentration ($\mu\text{g g}^{-1}$ ww or fw)	CV	Ratio (‰)	CV (%)	Ratio (‰)	CV (%)
Blood	Adélie (adults)	22	0.040±0.013 ^a (0.022 to 0.084)	32.76	-24.6±0.3 ^a (-25.1 to -24.1)	-1.23	8.5±0.3 ^a (7.8 to 9.1)	3.98
	Chinstrap (adults)	20	0.197±0.048 ^b (0.070 to 0.294)	24.43	-24.9±0.2 ^b (-25.4 to -24.7)	-0.79	8.7±0.2 ^a (8.3 to 9.0)	2.15
	Gentoo (adults)	21	0.082±0.104 ^a (0.019 to 0.342)	126.89	-24.7±0.4 ^{ab} (-25.0 to -23.0)	-1.79	8.5±0.6 ^{ab} (8.0 to 10.6)	7.39
	Gentoo (juvenile)	20	0.022±0.008 ^c (0.007 to 0.039)	36.85	-24.8±0.6 ^{ab} (-25.9 to -23.4)	-2.22	8.0±0.2 ^b (7.7 to 8.5)	2.97
Feather	Adélie (adults)	22	0.346±0.140 ^a (0.182 to 0.693)	40.58	-24.1±1.1 ^a (-27.4 to -22.4)	-4.60	9.1±0.4 ^{ab} (8.0 to 10.1)	4.57
	Chinstrap (adults)	20	0.742±0.243 ^b (0.398 to 1.290)	32.79	-23.3±1.0 ^b (-24.5 to -21.5)	-4.39	9.0±0.2 ^{ab} (8.6 to 9.3)	2.31
	Gentoo (adults)	21	0.514±0.513 ^a (0.180 to 1.976)	99.86	-23.6±0.7 ^{ab} (-24.5 to -21.8)	-2.87	9.4±0.6 ^a (8.6 to 11.3)	6.30
	Gentoo (juvenile)	20	0.297±0.341 ^c (0.079 to 1.523)	114.85	-23.0±1.0 ^b (-24.3 to -20.6)	-4.37	8.7±1.0 ^b (7.7 to 11.3)	10.97

Table 2. Mean±SD, range, and coefficient of variation (CV; %) of total Hg concentration ($\mu\text{g g}^{-1}$ ww) and stable carbon ($\delta^{13}\text{C}$) and nitrogen ($\delta^{15}\text{N}$) isotope values (‰) in four representative penguin prey taxa. Means not sharing a letter were found to differ significantly (Tukey's HSD, $p < 0.05$).

Prey group (species)	n	Hg		$\delta^{13}\text{C}$		$\delta^{15}\text{N}$	
		Concentration ($\mu\text{g g}^{-1}$ ww)	CV	Ratio (‰)	CV (%)	Ratio (‰)	CV (%)
Krill (<i>E. superba</i>)	6	0.002±0.001 ^a (0.001 to 0.002)	34.74	-26.4±0.8 ^a (-27.3 to -25.4)	-2.91	3.1±0.3 ^a (2.5 to 3.5)	11.05
Epipelagic fish (<i>P. antarcticum</i>)	10	0.008±0.002 ^b (0.005 to 0.011)	22.85	-25.1±0.2 ^b (-25.3 to -24.8)	-0.65	9.0±0.3 ^b (8.5 to 9.4)	3.53
Mesopelagic fish (<i>E. antarctica</i>)	10	0.031±0.017 ^c (0.017 to 0.069)	53.31	-25.8±0.6 ^a (-26.9 to -24.7)	-2.48	9.0±0.4 ^b (8.2 to 9.3)	3.94
Benthic fish (<i>L. squamifrons</i>)	10	0.041±0.021 ^c (0.012 to 0.083)	50.37	-24.2±0.7 ^c (-25.3 to -23.4)	-2.79	9.6±0.8 ^c (7.9 to 10.5)	8.39

Table 3. Model components, including: selected parameters, degrees of freedom (df), F value (F), P-value (P), parameter estimates (β) and standard errors (SE from the AIC_c selected models that best explain variation in total Hg for penguin feather and blood in adults as well as for only gentoo penguins. Na is shown for estimates of multi-level parameters.

Comparison	Tissue	Parameter	df	F	P	β	SE
All adults	Feather	Species	2	7.67	0.0011	Na	Na
		$\delta^{15}\text{N}$	1	1.44	0.2357	0.3756	0.0648
		Species* $\delta^{15}\text{N}$	1	7.31	0.0015	Na	Na
	Blood	Species	2	4.59	0.0124	Na	Na
		$\delta^{15}\text{N}$	1	1.16	0.2854	0.4401	0.0813
		Species* $\delta^{15}\text{N}$	1	4.33	0.0178	Na	Na
Gentoo only	Feather	$\delta^{15}\text{N}$	1	78.21	< 0.0001	0.3228	0.0365
	Blood	Age	1	3.83	0.058	Na	Na
		$\delta^{15}\text{N}$	1	1.03	0.3177	0.1471	0.2718
		Age* $\delta^{15}\text{N}$	1	4.12	0.0496	Na	Na