**The partitioning behavior of persistent toxicant organic contaminants in
eutrophic sediments: Coefficients and effects of fluorescent organic matter and
particle size**

**Wei He, Chen Yang, Wenxiu Liu, Qishuang He, Qingmei Wang, Yilong Li, Xiangzhen Kong,
Xinyu Lan, Fuliu Xu***

**MOE Laboratory for Earth Surface Processes, College of Urban & Environmental Sciences, Peking
University, Beijing 100871, China**

***Correspondence author: Tel./Fax: +86–10–62756122, Email: xufl@urban.pku.edu.cn**

Capsule abstract

The partition mechanism of PAEs in the sediment – pore water system in a eutrophic Chinese lake
was found to be different from that of PAHs and OCPs

Abstract

In the shallow lakes, the partitioning of organic contaminants into the water phase from the solid phase might pose a potential hazard to both benthic and planktonic organisms, which would further damage aquatic ecosystems. This study determined the concentrations of polycyclic aromatic hydrocarbons (PAHs), organochlorine pesticides (OCPs), and phthalate esters (PAEs) in both the sediment and the pore water from Lake Chaohu and calculated the sediment – pore water partition coefficient (K_D) and the organic carbon normalized sediment – pore water partition coefficient (K_{OC}), and explored the effects of particle size, organic matter content, and parallel factor fluorescent organic matter (PARAFAC-FOM) on K_D . The results showed that log K_D values of PAHs (2.61 – 3.94) and OCPs (1.75 – 3.05) were significantly lower than that of PAEs (4.13 – 5.05) ($p < 0.05$). The chemicals were ranked by log K_{OC} as follows: PAEs (6.05 – 6.94) > PAHs (4.61 – 5.86) > OCPs (3.62 – 4.97). A modified MCI model can predict K_{OC} values in a range of log 1.5 at a higher frequency, especially for PAEs. The significantly positive correlation between K_{OC} and the octanol – water partition coefficient (K_{OW}) were observed for PAHs and OCPs. However, significant correlation was found for PAEs only when excluding PAEs with lower K_{OW} . Sediments with smaller particle sizes (clay and silt) and their organic matter would affect distributions of PAHs and OCPs between the sediment and the pore water. Protein-like fluorescent organic matter (C2) was associated with the K_D of PAEs. Furthermore, the partitioning of PARAFAC-FOM between the sediment and the pore water could potentially affect the distribution of organic pollutants. The partitioning mechanism of PAEs between the sediment and the pore water might be different from that of PAHs and OCPs, as indicated by their associations with influencing factors and K_{OW} .

Keywords: persistent toxicant organic contaminants; partition coefficient (K_D); organic carbon

normalized partition coefficient (K_{OC}); organic matter; particle size; fluorescent organic matter (FOM)

1. Introduction

In an aquatic ecosystem, sediment traps and releases persistent toxic organic contaminants (PTOCs) such as polycyclic aromatic hydrocarbons (PAHs), organochlorine pesticides (OCPs), and phthalate esters (PAEs) via the settling of suspended particulate matter and the resuspension of sedimentary solids (He et al., 2013a; Liu et al., 2013; Qin et al., 2014). To visualize the environmental behavior and the bioaccumulation of PTOCs in sediment, an equilibrium-partitioning model (EqP) is typically employed (Kraaij et al., 2003; Ololade et al., 2012). In brief, the EqP model assumes that the partitioning of PTOCs between three compartments, the lipids in organisms, the pore water, and the sediment organic matter, is in equilibrium (Kraaij et al., 2003). The pore water links the sediment solids and organisms in that model (Ololade et al., 2012) and exchanges with the overlying water (Booij et al., 2003). In addition, the mobile pore water were much earlier to be ingested by benthos than sediment. Furthermore, most toxicity data and bioaccumulation calculation for aquatic organisms were based on exposure in water. Those are reasons why pore water are more appropriate for risk assessment than the sediment (Hawthorne et al., 2007a). Thus, ecological risk and environmental quality standards for PTOCs in sediment is generally assessed by directly determining the PTOCs in the pore water or indirectly calculated based on the PTOCs in the sediment and the partitioning behavior (Neff et al., 2005; Dueri et al., 2008; Smedes et al., 2013; Charrasse et al., 2014). Although direct measurement of the PTOCs in the pore water is more accurate than drawing inferences from sediment concentration (Arp et al., 2009; Yates et al., 2013),

we still need alternative indirect methods when the PTOC concentration in the pore water are not available (Neff et al., 2005; Persson et al., 2005). Furthermore, the partitioning behavior of PTOCs in sediment is of great concern because it can predict the PTOC content of the pore water and their potential risk (Kraaij et al., 2003; Shi et al., 2012; Smedes et al., 2013; Li et al., 2015). Recently employed equilibrium passive sampling methods for the direct risk assessment of PTOCs in sediment are also based on the PTOC partitioning behavior (Haftka et al., 2010; Smedes et al., 2013; Yates et al., 2013; Charrasse et al., 2014).

To date, most studies have focused on the sediment – pore water partitioning of PAHs and their derivatives in harbors (Arp et al., 2011), coastal seas (Ololade et al., 2012; Smedes et al., 2013; Yates et al., 2013), wetlands (Cao et al., 2015), rivers (Hawthorne et al., 2006; Yu et al., 2009; Shi et al., 2012), and lakes (Qiao et al., 2008; Haftka et al., 2010). In both anthropogenically impacted harbor sediment and a natural mangrove wetland sediment, PAH partitioning was highly variable and typically correlated with organic matter in sediment solids (Gao et al., 1998; Arp et al., 2011). In detail, clay and sand particles, containing more organic matter, trap more PAHs compared with other sediment solids (Gao et al., 1998). Haftka et al. (2010) found that the distribution between PAHs in the pore water and PAHs in the sediment exhibited a significant dependence on the phases containing more organic matter. Temperature might decrease the partitioning of PAHs into the particulate phase of the sediment via thermal adsorption (Haftka et al., 2010). Chemical compositions and structures of both PAH and organic matter essentially associate the PAH's partitioning between pore water and sediment solids (Qiao et al., 2008; Shi et al., 2012; Qin et al., 2014). For example, PAHs with a high number of aromatic rings (e.g., indeno(1,2,3-cd)pyrene, dibenzo(a,l)pyrene) are seldom detected in water phases because their aromatic rings help to highly associate with sediment particles through

π - π electron donor-acceptor interactions (Qiao et al., 2008; Arp et al., 2009; Tao et al., 2014). The organic matter fractions most affected include black carbon (Hawthorne et al., 2007b), soot carbon (Qiao et al., 2008), and aromatic carbon (Haftka et al., 2010). Moreover, coal combustion contributes more highly to the PAH partitioning than vehicular emissions and emissions of petrogenic origin (Shi et al., 2012). Conversely, the sediment – pore water partitioning of OCPs and PAEs is of less concern than their sediment – overlying water partitioning and PAH sediment – pore water partitioning (Williams et al., 1995; tenHulscher et al., 1997; Dueri et al., 2008; Dai et al., 2013).

Due to PTOC association with organic matter (shown by organic carbon, OC), the OC-normalized partitioning coefficient (K_{OC}) is typically calculated to indicate the diverse partitioning behaviors of various PTOCs into/onto organic matter in the field (*in situ*) or the lab (*ex situ*) (Williams et al., 1995; Arp et al., 2009; Shi et al., 2012; Cao et al., 2015). In addition to the experimental measurement of K_{OC} , two other popular categories for the estimation of K_{OC} based on organic matter content of the sediment exist ((e.g., a carbon model and an amorphous organic carbon (AOC) and black carbon (BC) model) and a quantitative-structure activity relationship (QSAR) approach (e.g., a K_{OW} related model, Raoult's law, a poly-parameter linear free energy relationship (PP-LFERs)) (Arp et al., 2009; Hawthorne et al., 2011; Martinez et al., 2013; Charrasse et al., 2014). Multiple models have been typically employed to estimate the K_{OC} of PTOCs in sediment in recent studies (Martinez et al., 2013; Charrasse et al., 2014).

To our knowledge, few studies have investigated the sediment – pore water partitioning of PTOCs, including PAHs, OCPs, and PAEs in eutrophic lakes based on field investigations. Although ratio of the organic matter content (as indicated by the OC) in the sediment to that in the pore water might affect PTOC partitioning behavior, the parallel factor fluorescent organic matter's

(PARAFAC-FOM's) effects have not been sufficiently investigated. Furthermore, PARAFAC-FOM has been shown to associate with PTOCs in previous studies (Hur et al., 2014). Thus, It is worth investigating whether the distribution behavior of sediment organic matter revealed by FOM also affects PTOC partitioning behavior. In summary, this study focuses on 1) discriminating the partitioning behavior of PAHs, OCPs, and PAEs in a eutrophic lake, 2) the applicability of multiple models to estimate the K_{OC} of various PTOCs, and 3) the effects of particle size, organic matter content and PARAFAC-FOM on PTOC partitioning.

2. Materials and methods

2.1 Sampling and pretreatment

Nineteen surface sediment samples were collected from locations and with methods used in previous studies (Jiang et al., 2014; Kang et al., 2015; He et al., 2016a). Briefly, in May 2011, ten lake sites (L01~L10) and nine river estuarine sites (R01, R03~R10) were determined in a large, shallow, and eutrophic fresh water lake, Lake Chaohu (Fig. 1). This lake is located in one of the most developed areas of China, the Yangtze River Delta Economic Zone, and has experienced serious nutrient, heavy metal, and persistent organic pollutant contamination due to a significant increase in human activity during the last 35 years (Jiang et al., 2014; Qin et al., 2014; He et al., 2016a). After collection using a stainless steel Ponar grab sampler (Omwoma et al., 2015; He et al., 2016a), homogenized surface sediment (<15 cm based on the angle of that system), samples were centrifuged at 5000 rpm for 30 min in the laboratory (Yu et al., 2009; Ololade et al., 2012). Generally, to expel the particulate organic matter or mineral solids and obtain pore water, 0.1~0.7 μm pore size membrane were employed (Wang et al., 2011). The pore water was filtered using 0.7 μm glass fiber

membrane and stored at 4 °C until analysis in this study. Then, the sediment was freeze-dried, ground with an agate mortar, and sifted through a 200-mesh sieve. Before extraction and cleanup, a composite sample, mixed thoroughly with a sub-sample, was prepared to minimize the sampling error caused by the sampler.

Fig. 1 Location of Lake Chaohu and the sampling sites.

2.2 Reagents and materials

A standard mixed stock of 25 PAHs, 25 OCPs, and 19 PAEs as well as the internal standards (ISs) and surrogate standards (SSs) were purchased from AccuStandard Inc., NY, USA and are listed in Table S1. The ISs for PAHs were acenaphthene-D10, anthracene-D10, chrysene-D12, naphthalene-D8, and perylene-D12, and PAH SSs were 2-Fluorobiphenyl and p-Terphenyl-D14. The IS and SS for OCPs were pentachloronitrobenzen and 1-bromo-2-nitrobenzene, respectively. The PAE ISs included 3,4,5,6,D4- diethyl phthalate and 3,4,5,6,D4-di-n-octyl phthalate, and their SS was 3,4,5,6,D4- di-n-butyl phthalate. HPLC-grade methanol (MET), n-hexane (HEX), acetone (ACE), and dichloromethane (DCM) were supplied by J & K Chemical Ltd. in China. A solid phase extract (SPE) device and C18 SPE cartridges (ENVI-18, 6 ml, 500 mg) were purchased from Supelco Co., Bellefonte, Pennsylvania, USA. Anhydrous sodium sulfate (analytical grade, baked at 650 °C for 6 h before use), alumina (200-300 mesh), silica gel (200-300 mesh), sodium hydroxide (NaOH, analytical grade), concentrated sulfuric acid (H₂SO₄, 98%, analytical grade), and copper granules were purchased from Sinopharm Chemical Reagent Co., Ltd. (Shanghai, China). Both the silica gel and the alumina were baked at 450 °C for 6 h and activated for 16 h at 130 °C before use. Ultrapure water (3 g) was added to 100 g of silica gel or alumina to form neutral silica gel and alumina. All glassware, including the Soxhlet extractor, columns, flasks, bottles, vial insert tubes, and vials, were

ultrasonically cleaned with distilled water, washed with ultrapure water, and baked at 450 °C for 6 h. The microwave-extract ion cylinders, plugs, and lids were ultrasonically cleaned with distilled water, washed with ultrapure water, rinsed with ACE twice, and placed on filter paper to dry.

2.3 Extraction and cleanup

The same extraction and cleanup procedures of PAHs and OCPs for both the sediment and the pore water as described in previous studies were used (Liu et al., 2013; Qin et al., 2014). In brief, the pore water was extracted by an SPE system after the SPE cartridges were prewashed with DCM and conditioned with MET and ultrapure water. The SPE cartridges were N₂-dried and eluted with DCM. The sediment was extracted with a HEX/ACE (1:1, v/v) mixture using the microwave-assisted extraction (MAE) system MARS 5 (CEM Corp., Matthews, NC, USA). The crude extract was concentrated to 1 mL, transferred onto an anhydrous sodium sulfate/ silica gel/alumina (1 cm/12 cm/12 cm) column (10-mm i.d.), and eluted with a HEX/DCM (1:1, v/v) mixture for cleanup. Both the pore water and sediment refined extracts were concentrated to a volume of 1 mL with a vacuum rotary evaporator. After the IS was added, the sample vial was stored at -20 °C until analysis by gas chromatography-mass spectrometry (GC-MS).

The PAE extraction and cleanup procedures were also from previous studies (He et al., 2013b; Kang et al., 2015). The pore water was acidified to pH ~ 2.0 before following the same SPE procedures as for PAHs and OCPs. The prewashing and eluting solvents were replaced by HEX and HEX/DCM (1:1, v/v). The sediment was extracted with DCM for 48 h in a Soxhlet extractor. Activated copper granules were added to remove elemental sulfur. Concentrated extracts were transferred onto an anhydrous sodium sulfate/alumina/silica gel (1 cm/12 cm/12 cm) column (10-mm i.d.), cleaned with DCM/HEX (3:7, v/v), and eluted with ACE/HEX (2:8, v/v). The refined elutes

were treated the same as the PAHs and OCPs described above.

The sediment organic matter, also known as the alkaline-extractable organic matter (AEOM), was extracted from the sediment following procedures employed in previous studies (Hur et al., 2014; He et al., 2016c). Briefly, the sediment sample was soaked with 0.1 mol L⁻¹ NaOH and shaken for 24 h with nitrogen purging before centrifugation and filtration using a pre-washed 0.45- μ m pore-sized membrane. The supernatant was diluted, and the pH was adjusted to ~7. Interfering cations were removed from the alkaline extracts with a cation exchange resin. The purified AEOM solution was stored at 4 °C until spectroscopic analysis.

2.4 Instrument analysis

The instrument conditions for the GC-MS analysis of PAHs, OCPs, and PAEs are listed in Text S1. Quantitation was performed using an internal calibration method based on a six-point calibration curve for individual PTOCs. The method detection limits (MDL) are listed in Table S1. Total organic carbon (TOC) in the sediment was analyzed by a TOC 5000A coupled with an SSM-5000A sampler (Shimadzu Corp., Japan). Potassium acid phthalate was used to calibrate the TOC analyzer. The AEOM and pore water were characterized using a PerkinElmer Lambda 35 spectrometer with a 1-cm cuvette to scan the absorption spectra from 200 to 800 nm, and the fluorescent excitation emission matrix (EEM) was determined with a fluorescence spectrometer (Hitachi F4500) at the excitation/emission wavelengths of 200 – 400/220 – 600 nm with excitation at a 5-nm step and emission at a 2-nm step. According to an inter-laboratory standard method proposed by Murphy et al. (2010), the EEM data were corrected with a water blank and both primary and secondary inner filter effects using the automatic Matlab code FDOMcorrect. The particle size was measured by a Fritsch A22 laser particle analyzer. Clay, silt, fine sand, and coarse sand fractions were identified based on

the particle size distribution.

2.5 QA/QC

The extracted sediment and ultrapure water were spiked with PAHs, OCPs, and PAEs in triplicate to test the recovery and precision of the analytical procedures. The average recovery percentages for most target PTOCs were within a range of 70% – 130% for sediment, but some PTOCs were not within a reasonable range for the pore water (Table S1). Thus, the results of sediment – pore water partitioning behavior of those chemicals should be suspect when used for modeling or employed for other deductions. The SS was added to all samples before extraction to evaluate the recovery, and the recovery was in a range of 70% – 130% for both the sediment and the pore water. The MDLs of individual PAHs, OCPs, and PAEs were 0.021 – 0.083, 0.021– 0.21, and 0.013 – 0.13 ng g⁻¹ dry weight (dw) for 8 g of sediment and 0.05 – 0.1, 0.005 – 0.1, 0.1 – 1 ng L⁻¹ for 1 L of pore water (Table S1). Three blank samples were analyzed for solvent and glassware contamination to correct the sample data.

2.6 Data handling

2.6.1 EEM parallel factor fluorescent organic matter (EEM-PARAFAC-FOM)

We extracted the PARAFAC based on a total of 38 EEMs in the pore water and AEOM using the Matlab toolbox (DOMFluor) as shown in Fig. S1 (Stedmon and Bro, 2008). The core consistency diagnostic combined with the explained variances was used to identify the number of PARAFAC components (Yang et al., 2015). As shown in Fig. S1 and Table S2, a 6-component model was finally chosen to describe the EEM features for all of the samples the core consistency was good (67.4%) and most of the variance was explained (99.6%) (Fig. S1). According to the suggestion of Parr et al. (2014), a self-edited Matlab code (comPARAFAC.m) was employed to compare the identified

components with a library containing 38 PARAFAC models (Table S3) (He and Hur, 2015). Based on the excellent coincidence of $mTCC > 0.95$ and results from the relevant literature, the following 6 components were identified: the protein-like C1 and C2 components were similar to tryptophan and tyrosine, respectively, and indicated lability of the DOM and bacterial production (Fellman et al., 2011). However, a protein-like C5 component was not observed in the library, so we assumed that it was a tryptophan-like FOM that could absorb high energy light when compared with C1. It might be refractory protein-like organic matter. A C3, similar to a marine humic-like FOM, was strongly associated with recent biological activity (Yamashita et al., 2010). Both C4 and C6 were humic-like FOM. C4 was close to the “unknown” component 1 identified by Cory and McKnight (2005). C6 had spectral features of an oxidized quinone-like DOM moiety, which was also attributed to biologically labile organic matter primarily rich in aliphatic carbon (Singh et al., 2013). The normalized concentration in Raman units (RU) of each PARAFAC component was estimated based on the F_{max} output from DOMFluor (He et al., 2016c).

2.6.2 Partitioning models

After measuring the PTOCs in both the pore water and the sediment, the partitioning coefficient (K_D) and the organic carbon normalized partitioning coefficient (K_{OC}) were calculated:

$$K_D = C_s / C_{pw}, \quad (1)$$

$$K_{OC} = K_D / f_{OC}, \quad (2)$$

where C_s is the sediment concentration, C_{pw} is the pore water concentration, and f_{OC} is the sediment fraction of organic carbon.

In this study, the K_{OC} was estimated by QSAR models, including the linear relationship between K_{OC} and the octanol – water partition coefficient (K_{OW}) and between the K_{OC} and the molecular

connectivity indices (MCI) of the chemicals, using EPI Suite™ v.4.11. The formulas for those relationships were described in the help documents of EPI Suite. We also attempted to use the Raoult's law (Arp et al., 2009):

$$\log K_{OC} = -\log(S_L^* \cdot MW_{OC}). \quad (3)$$

where S_L^* is the sub-cooled saturated molar water solubility for the chemicals, and MW_{OC} is the molar weight of the total organic phase of the sediment (223 g mol⁻¹). The relevant parameters for each chemical that was estimated or cited are summarized in Table S4.

2.6.3 Statistical analysis

Statistical analyses, including an independent t-test, Spearman's correlation, and a principal component analysis (PCA), were conducted with SPSS v20 software. A significant value was assigned with $p < 0.05$. The PCA was used to show the association between the PTOCs and influencing factors.

3. Results and discussion

3.1 Occurrence and spatial variations of PAHs, OCPs, and PAEs in pore water and sediment

In the eutrophic Lake Chaohu, PAEs were detected at the highest concentration, followed by PAHs and OCPs in the surface water (He et al., 2013b; Liu et al., 2013; Qin et al., 2014). This rank pattern was also observed in both the sediment and the pore water. Furthermore, total PAEs were 100-fold greater than PAHs and 1000-fold greater than OCPs (Table 1). We also found that individual PTOCs had different detection frequencies (DFs) between the sediment and the pore water. Overall, PTOCs in the sediment had higher DFs than the pore water (Table S5). High-ring PAHs were seldom detected in pore water but often detected in sediment, indicating that high-ring PAHs

preferentially partition in the solid phase of the sediment. Although low-ring PAHs were detected at high concentrations in the pore water, their concentrations were still much lower than in the sediment, indicating that the solid phase of the sediment trapped more PAHs than the dissolved phases. The large order of magnitudes differences between the PAHs concentrations in the pore water and the sediment were similar to those in Ondo coastal area, Nigeria and Teufelsweiher Pond in Germany (Gao et al., 1998; Ololade et al., 2012), but lower than in the Lanzhou Reach of the Yellow River in China (Yu et al., 2009).

Table 1 PAHs, OCPs, PAEs in the pore water and the sediment collected from limnic and riverine areas of Lake Chaohu, the particle sizes and total organic carbon (TOC) of the sediment, and the PARARAC-FOMs in both the AEOM and the pore water.

The DFs of the OCPs were diverse with the chemical species. More OCPs were detected in the solid phases. Only hexachlorbenzene (HCB) was detected in both sediment and pore water. The four hexachlorocyclohexane isomers (HCHs) had higher DFs in the pore water, whereas dichlorodiphenyltrichloroethane (DDT) and its degradation products (dichlorodiphenyldichloroethylene (DDE) and dichlorodiphenyldichloroethane (DDD)) had higher DFs in the sediment. More Aldrin (Ald) was detectable in the pore water than isodrin (ID), whereas ID was only detected in the sediment. Generally, HCHs in sediment were 1~100-fold greater than in the pore water (Dai et al., 2013). However, three orders of magnitude separated the sediment and pore water concentrations in Lake Chaohu, indicating that the eutrophic sediment might trap more HCHs. Similar to PAHs, more PAEs were detected in the sediment than the pore water. Nevertheless, PAEs in sediment were at a 1000-fold greater concentration than in the pore water and the PAEs in the pore water were actually much greater than in the surface water (He et al., 2013b), indicating a

potential surface water – pore water – sediment distribution pattern for PAEs.

In Lake Chaohu, the limnic and riverine locations can be significantly discriminated by slit, coarse and, C2-FOM, C3-FOM, and C5-FOM in the sediment and C1-FOM, C2-FOM, C5-FOM, and C6-FOM in the pore water ($p < 0.05$) (Table 1). Typically, limnic sediment contained more slit and FOMs but less coarse sand than riverine sediment. It seemed that PAHs and PAEs in lake ecosystem were carried by river nets as indicated by the larger concentrations in riverine sediment and pore water. Furthermore, the seriously polluted rivers were caused by anthropogenic effects such as coal combustion and plastic discarding without reasonable treatment (He et al., 2013b; Qin et al., 2014). Conversely, OCPs were detected at higher concentrations in the limnic sediments and pore water. It disclosed that discharge of OCPs into rivers became weaker and historic residuals in their sediment and pore water were gradually released into overlying water by the strong perturbation in rivers (Arp et al., 2011).

3.2 Sediment – pore water partition coefficients of PAHs, OCPs, and PAEs

The apparent sediment – pore water partition coefficients (K_D) of individual PTOCs in Lake Chaohu were calculated in Table S5, summarized in Table 2, and selectively shown in Fig. 2. K_D could be obtained for most PAHs and PAEs (> 60%) but for no more than 50% of OCPs because of inconsistent detection in both the sediment and the pore water. Significant differences in the partitioning behavior of PAHs (0.24 – 0.63) and OCPs (0.05 – 0.31) was indicated by their DF values ($p < 0.05$). Compared with PAHs and OCPs, the DF range for PAEs was moderate. In detail, the partitioning behaviors of 10 PAHs (naphthalene, acenaphthylene, acenaphthene, fluorine, phenanthrene, anthracene, fluoranthene, pyrene, retene, benz(a)anthracene,), 2 OCPs (α -HCH and

HCB), and 5 PAEs (diethyl phthalate (DEP), diisobutyl phthalate (DIBP), di-n-butyl phthalate (DnBP), benzyl butyl phthalate (BBP), di-(2-ethylhexyl) phthalate (DEHP)) were clearly observed at most sampling locations ($DF > 0.8$). Those PTOCs' high occurrences in both sediment and pore water reminded us that those chemicals typically exchanged among multiple media such as sediment, pore water, and overlying water (Booij et al., 2003; Kraaij et al., 2003). Those partitioning behaviors would cause their ubiquity in aquatic ecosystem, which pose a potential risk to both the aquatic ecosystem and human water safety (Neff et al., 2005; Dueri et al., 2008; Arp et al., 2011; He et al., 2013b). The variation of K_D values was not significantly different between river and lake locations, as indicated by an independent t test ($p > 0.05$, Fig. 2), indicating that similar partitioning behavior occurred in both the limnic and riverine areas of Lake Chaohu. The K_D values of PAHs (2.61 – 3.94) and OCPs (1.75 – 3.05) was significantly lower than for PAEs (4.13 – 5.05) ($p < 0.05$, Table 2), indicating that more PAEs were adsorbed by the sediment solids.

Table 2 The variance of the ratio, detection frequency, partition coefficient ($\log K_D$), and organic carbon-water partition coefficient ($\log K_{OC}$) of PAHs, OCPs, and PAEs.

Fig. 2 The measured sediment-pore water partition coefficients (K_D) of PAHs, OCPs, and PAEs ($DF > 0.15$) from limnic and riverine locations in Lake Chaohu. The error bar indicates the standard deviation.

The apparent K_{OC} values of the individual PTOCs are shown in Table 3 and summarized in Table 2. According to the $\log K_{OC}$ values, the PTOCs can be significantly ranked: PAEs (6.05 – 6.94) > PAHs (4.61 – 5.86) > OCPs (3.62 – 4.97). As shown in Fig. 3a, most average K_{OC} values of PAHs were scattered below 1:1 line, indicating they were larger than the corresponding K_{OC} in the EPI Suite database. The differences between the observed and the recorded K_{OC} values of the PAHs

showed that the sediment in Lake Chaohu had larger adsorption capacity for PAHs and can be linked by a factor within 1 to 30 (Hawthorne et al., 2011). Most OCPs were within the 1:30 line except Ald, which was 2 orders of magnitude lower than the values in the literature. No significant differences were present for the 4 HCH isomers, which had a range of 3.72 to 4.04. The experimental K_{OC} values of DDE, DDT, HCB, and methoxychlor (MOC) were close to the literature values. Interestingly, Fig. 3a shows that the PAEs were scattered outside the 1:30 line, indicating very large differences between the experimental values and the literature values. We deduced that different affinities of PAEs for the sediment might exist in Lake Chaohu. That is, organic matter did not solely adsorb PAEs and minerals might also adsorb the predominant PAEs, especially PAEs with shorter carbon chains (e.g., dimethyl phthalate (DMP), DEP, DIBP, DnBP) (Williams et al., 1995; Gao et al., 1998; Cao et al., 2015; He et al., 2016b).

Fig. 3 Comparison of experimental $\log K_{OC}$ values for individual PTOCs in Lake Chaohu sediments with the (a) experimental data from the EPI Suite database ($\log K_{OC,exp}$) and estimates of (b) the octanol – water partition coefficient ($\log K_{OC,ow}$), (c) molecular connectivity indices ($\log K_{OC,MCI}$), and (d) Raoult's law ($\log K_{OC,s}$). Figures containing more information refers to Fig. S2 in supplementary materials.

3.3 Applicability of various K_{OC} evaluation models

Compared with $\log K_{OC,ow}$, $\log K_{OC,MCI}$, and $\log K_{OC,s}$, the applicability of the K_{OC} evaluation models are shown in Fig. 3. Overall, the estimates using MCI showed the best performance as indicated by that more positions were scattered along the 1:1 line or within the 1:30 lines. This model predicted $K_{OC,MCI}$ values of PTOCs, PAHs, and OCPs in a range of $\log 1.5$ with frequencies of 69%,

94%, and 91%, respectively. Raoult's law did not show good performance, as in previous studies (Hawthorne et al., 2011; Charrasse et al., 2014). This model predicted the $K_{OC,S}$ values of PTOCs, PAHs, and OCPs in a range of log 1.5 with frequencies of 44%, 44%, and 73%, respectively. PAEs were not be predicted by all three models and 83% of the species were outside the 1:30 line. Raoult's law gave negative values for DMP and DEP, but both the observed and the literature values were positive (Williams et al., 1995), indicating the inapplicability of Raoult's law for the estimation of the K_{OC} of PTOCs in Lake Chaohu. As suggested by Hawthorne et al. (2011), a weathering factor (WF) was employed to correct the MCI model. After investigating frequency variations of the PTOCs within a ranges of log 1.5 (also within the 1:30 line) with increasing WF, we proposed WF = 1.0 as a correction factor. The modified MCI model can predict $K_{OC,MCI}$ values of PTOCs, PAHs, OCPs, PAEs in a range of log 1.5 with frequencies of 82.1%, 88%, 91%, and 69%, respectively. Although several PAHs could not be reasonably predicted, the modified MCI model can estimate significantly more PAEs than the original model (frequency = 17%). Although the estimates for the K_{OW} model were similar to the MCI model, the modified K_{OW} model was not as greatly improved as the MCI model (Fig. S3).

Generally, K_{OC} is positively correlated with K_{OW} (Hawthorne et al., 2006; Qiao et al., 2008; Arp et al., 2009; Charrasse et al., 2014; Cao et al., 2015). Significant positive correlations between K_{OC} and K_{OW} were observed for PAHs and OCPs after excluding outliers and their linear relationships are shown in Fig. 4. The ratio of log K_{OC} to log K_{OW} ($\log K_{OC} / \log K_{OW}$) was > 1 for PAHs based on the linear relationships, indicating that sediment carbon in Lake Chaohu was had greater affinity for PAHs than for n-octanol (Qiao et al., 2008). According to the OCPs $\log K_{OC} - \log K_{OW}$ formula, $\log K_{OC} / \log K_{OW}$ was > 1 when $\log K_{OW}$ was < 4.05 . That is to say, OCPs with lower K_{OW} values ($<$

4.05) might also have greater affinity for the sediment organic carbon than n-octanol. On the contrary, no credible linear relationship could be obtained for PAEs unless that we excluded PAEs with low K_{OW} (<4) such as DMP, DEP, di-(2-ethoxyethyl) phthalate (DEEP). A better linear relationships could be conducted after excluding more individual PAEs such as DIBP and dinonyl phthalate (DNP). It disclosed that PAEs' distribution between sediment and water could not predicted by their distribution between n-octanol and water, especially for those PAEs with low K_{OW} . PAEs with low K_{OW} preferred to distribute in water in n-octanol – water system, whereas they had more affinity with sediment particles in sediment – pore water system. Thus, we concluded that partitioning mechanism of PAEs between the sediment and the pore water might differ from PAHs and OCPs based on their relationships with K_{OW} .

To our knowledge, sediment consist of both organic matter and minerals. Organic matter interacted with minerals through both hydrophilic way (electrostatic forces and/or ligand exchange reactions) and hydrophobic interactions (He et al., 2016b). PTOCs had anthropogenic origins, anyhow but also a group of organic chemicals. Their interaction with sediment particles was similar to organic matter and largely depended on the their properties and sediment composition. For example, PAHs were generally adsorbed through π - π electron donor-acceptor interactions between PAHs and aromatic rings of sediment organic matter (SOM) or/and between PAHs and aluminium hydroxides in surface of minerals (Arp et al., 2009; Tao et al., 2014). OH-groups and coordinated exchangeable cations were probably active positions for the OCPs' sorption on the sediment mineral particles (Sprynskyy et al., 2008). Furthermore, OCPs with larger $\log K_{OW} > 4.5$ generally had larger molecular size, which prohibited them entering micro-pores of mineral particles or microscopic structure of SOM (Pignatello and Xing, 1996). PAEs are amphiphilic chemicals, consisting of both

hydrophobic fractions (e.g., aromatic rings, aliphatics) and hydrophilic fraction (ester linkage) (Wen et al., 2013). The hydrophobic interaction between PAEs and sediment particles is similar to PAHs and OCPs. Furthermore, oxygen (O) atom in ester linkage of PAEs can form hydrogen bond with sediment particles, which might strengthen their affinity with sediment particles (Zheng et al., 2007).

Fig. 4 Linear relationships between $\log K_{OC}$ and $\log K_{OW}$ for PAHs (a), OCPs (b), and PAEs (c).

Retene, cyclopenta(c,d)pyrene, methoxychlor, DMP, DEP, DIBP, DEEP, and DNP were excluded to obtain the slope and intercept of linear regression.

In this study, simple LFER models were employed to describe the PTOC distribution in the sediment although those models have been criticized as having a large uncertainty and limited applications to describe a large number of chemicals (Cao et al., 2015). Goss and Schwarzenbach (2001) proposed the PP-LFER model because it considers all interactions involved in partitioning using separate parameters. They also emphasized that more effort should be made to adapt and improve the more comprehensive PP-LFER model rather than to modify existing one-parameter LFER (Goss and Schwarzenbach, 2001). Nguyen et al. (2005) have tested the applicability of PP-LFER to unify various groups of chemicals, including monoaromatic hydrocarbons, halogenated hydrocarbons, PAHs, and polar chemicals. Nevertheless, the parameters required by PP-LFER are difficult to obtain. However, we still propose that PP-LFERs be used to test K_{OC} estimates in a future study.

3.4 Effects of fluorescent organic matters and particle sizes on K_D

Sediment is mainly consists of organic matter and minerals. Both compartments have been demonstrated to bind or adsorb organic contaminants (Gao et al., 1998; Haftka et al., 2010).

Furthermore, the pore water, which contains organic matter, also has been shown to effect the distribution of organic contaminants (Haftka et al., 2010). To test whether minerals, total organic matter, and the distribution of fluorescent organic matter between the sediment and the pore water affects the PTOCs' sediment – pore water partitioning behavior, a correlation analysis was conducted and the results are shown Table S6. We expected to find that both PAHs and OCPs were positively correlated with the total organic carbon (TOC) of the sediment, but some PAEs were negatively correlated with the TOC, especially those that contained tyrosine-like organic matter (C2-FOM). Most PAHs and OCPs seemed to be positively associated with both sediment minerals (clay, slit, and coarse sand) and the TOC, whereas PAE associations with sediment properties were rather inconsistent and depended on the PAE species. PARAFAC-FOM partitioning between the sediment and the pore water (SPC) potentially affected the distribution of organic contaminants as indicated by a negative association between most PTOCs and SPC. This phenomenon demonstrated that FOM exchange from the pore water to the sediment might inhibit the PTOC exchange from the pore water to the sediment but promote PTOC exchange from the sediment to the pore water. That is to say, FOM might compete with PTOC in binding to sites in the sediment solids. Furthermore, tyrosine-like organic matter (C2) was the signature FOM that demonstrated the association between FOM and PTOC partitioning behavior.

PCA also visually revealed individual PTOC associations with influencing factors (Fig. 5). Component 1 (PC1) and component 2 (PC2) explained 53.7% and 27.6% of the variance, respectively. And PC1 and PC2 were associated with organic matter and minerals, respectively. The positive side of PC1 denoted the sediment TOC and mainly protein-like FOM's partitioning between the sediment and the pore water (C1, C2, and C5), whereas the positive side of PC2 indicated

sediment minerals with lower particle sizes (silt and clay). Although TOC was located between clay and sand fractions and in the lower right quadrant, it did not mean that TOC was more associated with sand. Actually, TOC, coarse sand, and clay were located in the positive direction of PC1, indicating their association with organic matter. Previous investigation also showed that sediment organic matter distributed more in those fractions (Gao et al., 1998). Most PAHs were in the upper right quadrant and therefore associated with clay, C1, and C4, indicating that those PAHs' partition between pore water and sediment solids were affected by particles' specific surficial areas, aromatic ring-containing FOM, and humic-like substances. Their interactions mechanisms included both hydrophilic way (electrostatic forces and/or ligand exchange reactions) and hydrophobic interactions (Arp et al., 2009; Tao et al., 2014). However, naphthalene (Nap) and acenaphthylene (Acy) were in the lower right quadrant and associated with sand, C2 and C5. It seemed that the sediment – pore water partitioning behaviors of PAHs with lower aromatic rings were manipulated by particles coated with protein-like substances (C2 and C5 FOM). The regulation mechanisms was still unknown and proposed to be investigated in the future study. Both C3 and C6 were labile FOM derived from recent biological activity (Yamashita et al., 2010; Singh et al., 2013). Those recently produced FOMs possibly associated with α -HCH's and HCB's distribution. On the contrary, α -HCH's other isomers were associated with C1 and C4. Their different partitioning behaviors might be caused by their structures or/and bioavailability. Nevertheless, the present study still could not offer direct evidences for those speculations. Still, some credible results showed that some PTOCs' partitioning behaviors were associated with TOC, silt, clay and protein-like FOMs after combining with correlation analysis above.

Fig. 5 Principal component analysis' loading plot of PTOC partition coefficients and influencing

factors. The circles, squares, diamonds, and pentangles denote PAHs, OCPs, PAEs, and influencing factors.

3.5 Implication to of PTOCs' partition in sediment – pore water system

PTOC partitioning in a sediment – pore water system were of great concern because the PTOC risk might be transported from the water phase, trapped in the sediment, and potentially released into the water phase by resuspension (Liu et al., 2013; Qin et al., 2014). The K_{OC} values in Lake Chaohu were relatively higher than those calculated under laboratory conditions, suggesting that the measured pore water PTOC concentrations were lower than the predicted values. That is to say, not all of the sediment PTOCs are available to distribute rapidly into the pore water (Qiao et al., 2008). It also indicates caution when predicting the pore water concentration based on literature K_{OC} values (Arp et al., 2009; Yates et al., 2013). Otherwise, the ecological risks of PTOCs' might be overestimated. Furthermore, the relatively lower K_D values of OCPs results in more release from the sediment. Considering OCPs' high-toxicity, they might pose a potential risk as previously emphasized by the priority screening of PTOCs in Lake Chaohu (He et al., 2014). This study demonstrated that sediment minerals and fluorescent organic matter in the sediment/pore water might affect the PTOC partitioning behavior and that different partitioning mechanism might exist for PAHs, OCPs, and PAEs (Fig. 6); i.e., PAHs and OCPs had similar association with influencing factors and K_{OW} , whereas PAE associations with influencing factors and K_{OW} were different from PAHs and OCPs.

Fig. 6 The different partitioning mechanism of PAHs, OCPs, and PAEs and the potential influencing factors. The bold outline denotes the association degree between the partitioning

behavior and influencing factors.

4. Conclusions

In the sediment – pore water system of Lake Chaohu, sediment PTOCs had higher DFs than in the pore water. The spatial variation revealed that currently anthropogenic activity affected the PAHs and PAEs and historic contamination of OCPs. PAH log K_D values (2.61 – 3.94) and those of OCPs (1.75 – 3.05) was significantly lower than for PAEs (4.13 – 5.05) ($p < 0.05$). The chemicals were ranked by log K_{OC} as follows: PAEs (6.05 – 6.94) > PAHs (4.61 – 5.86) > OCPs (3.62 – 4.97). Both the MCI model and the K_{OW} models predicted PTOC K_{OC} values in range of log 1.5 with similar frequencies. However, the modified MCI model could predict $K_{OC,MCI}$ values in a range of log 1.5 with a higher frequency, especially for PAEs. Significant positive correlations between K_{OC} and the octanol – water partition coefficient (K_{OW}) were observed for PAHs and OCPs and linear relationships were also obtained. However, significant correlation was found for PAEs only when excluding PAEs with lower K_{OW} values. Sediments of smaller particle sizes (clay and silt) and organic matter would have different distributions of PAHs and OCPs between the sediment and the pore water. Protein-like organic matter (C2) was associated with the K_D of PAEs. Furthermore, PARAFAC-FOM's partitioning between the sediment and the pore water could potentially affect the distribution of organic pollutants. The partitioning mechanism of PAEs between the sediment and the pore water might differ from that of PAHs and OCPs, as indicated by their associations with influencing factors and K_{OW} .

Acknowledgments

This work was supported by the National Project for Water Pollution Control in China

(2012ZX07103-002) and the National Science Foundation of China (41503083, 41271462, 41030529).

References

- Arp, H.P.H., Breedveld, G.D., Cornelissen, G., 2009. Estimating the in situ sediment-porewater distribution of PAHs and chlorinated aromatic hydrocarbons in anthropogenic impacted sediments. *Environ. Sci. Technol.* 43, 5576-5585.
- Arp, H.P.H., Villers, F., Lepland, A., Kalaitzidis, S., Christanis, K., Oen, A.M.P., Breedveld, G.D., Cornelissen, G., 2011. Influence of historical industrial epochs on pore water and partitioning profiles of polycyclic aromatic hydrocarbons and polychlorinated biphenyls in Oslo Harbor, Norway, Sediment Cores. *Environ. Toxicol. Chem.* 30, 843-851.
- Booij, K., Hoedemaker, J.R., Bakker, J.F., 2003. Dissolved PCBs, PAHs, and HCB in pore waters and overlying waters of contaminated harbor sediments. *Environ. Sci. Technol.* 37, 4213-4220.
- Cao, Q.M., Wang, H., Qin, J.Q., Chen, G.Z., Zhang, Y.B., 2015. Partitioning of PAHs in pore water from mangrove wetlands in Shantou, China. *Ecotox. Environ. Safe.* 111, 42-47.
- Charrasse, B., Tixier, C., Hennebert, P., Doumenq, P., 2014. Polyethylene passive samplers to determine sediment-pore water distribution coefficients of persistent organic pollutants in five heavily contaminated dredged sediments. *Sci. Total. Environ.* 472, 1172-1178.
- Cory, R.M., McKnight, D.M., 2005. Fluorescence spectroscopy reveals ubiquitous presence of oxidized and reduced quinones in dissolved organic matter. *Environ. Sci. Technol.* 39, 8142-8149.
- Dai, G.H., Liu, X.H., Liang, G., Gong, W.W., Tao, L., Cheng, D.M., 2013. Evaluating the sediment-water exchange of hexachlorocyclohexanes (HCHs) in a major lake in North China. *Environ. Sci.-Process Impacts* 15, 423-432.
- Dueri, S., Castro-Jimenez, J., Comenges, J.M.Z., 2008. On the use of the partitioning approach to derive Environmental Quality Standards (EQS) for persistent organic pollutants (POPs) in sediments: A review of existing data. *Sci. Total. Environ.* 403, 23-33.
- Fellman, J.B., Petrone, K.C., Grierson, P.F., 2011. Source, biogeochemical cycling, and fluorescence characteristics of dissolved organic matter in an agro-urban estuary. *Limnol. Oceanogr.* 56, 243-256.
- Gao, J.P., Maguhn, J., Spitzauer, P., Kettrup, A., 1998. Distribution of polycyclic aromatic hydrocarbons (PAHs) in pore water and sediment of a small aquatic ecosystem. *Int. J. Environ. Anal. Chem.* 69, 227-242.
- Goss, K.U., Schwarzenbach, R.P., 2001. Linear free energy relationships used to evaluate equilibrium partitioning of organic compounds. *Environ. Sci. Technol.* 35, 1-9.
- Haftka, J.J.H., Govers, H.A.J., Parsons, J.R., 2010. Influence of temperature and origin of dissolved organic matter on the partitioning behavior of polycyclic aromatic hydrocarbons. *Environ. Sci. Pollut. Res.* 17, 1070-1079.
- Hawthorne, S.B., Azzolina, N.A., Neuhauser, E.F., Kreitinger, J.P., 2007a. Predicting bioavailability of sediment polycyclic aromatic hydrocarbons to *Hyaella azteca* using equilibrium partitioning,

supercritical fluid extraction, and pore water concentrations. *Environ. Sci. Technol.* 41, 6297-6304.

- Hawthorne, S.B., Grabanski, C.B., Miller, D.J., 2006. Measured partitioning coefficients for parent and alkyl polycyclic aromatic hydrocarbons in 114 historically contaminated sediments: Part 1. KOC values. *Environ. Toxicol. Chem.* 25, 2901-2911.
- Hawthorne, S.B., Grabanski, C.B., Miller, D.J., 2007b. Measured partition coefficients for parent and alkyl polycyclic aromatic hydrocarbons in 114 historically contaminated sediments: Part 2. Testing the KOCKBC two carbon-type model. *Environ. Toxicol. Chem.* 26, 2505-2516.
- Hawthorne, S.B., Grabanski, C.B., Miller, D.J., Arp, H.P.H., 2011. Improving Predictability of Sediment-Porewater Partitioning Models using Trends Observed with PCB-Contaminated Field Sediments. *Environ. Sci. Technol.* 45, 7365-7371.
- He, W., Bai, Z.-L., Liu, W.-X., Kong, X.-Z., Yang, B., Yang, C., Jørgensen, S.E., Xu, F.-L., 2016a. Occurrence, spatial distribution, sources, and risks of polychlorinated biphenyls and heavy metals in surface sediments from a large eutrophic Chinese lake (Lake Chaohu). *Environ. Sci. Pollut. Res.*, 1-14, doi:10.1007/s11356-11015-16001-11356.
- He, W., Chen, M., Schlautman, M.A., Hur, J., 2016b. Dynamic exchanges between DOM and POM pools in coastal and inland aquatic ecosystems: A review. *Sci. Total. Environ.* 551-552, 415-428.
- He, W., Hur, J., 2015. Conservative behavior of fluorescence EEM-PARAFAC components in resin fractionation processes and its applicability for characterizing dissolved organic matter. *Water Res.* 83, 217-226.
- He, W., Jung, H., Lee, J.-H., Hur, J., 2016c. Differences in spectroscopic characteristics between dissolved and particulate organic matters in sediments: Insight into distribution behavior of sediment organic matter. *Sci. Total. Environ.* 547, 1-8.
- He, W., Qin, N., Kong, X.-Z., Liu, W.-X., He, Q.-S., Wang, Q.-M., Yang, C., Jiang, Y.-J., Yang, B., Wu, W.-J., Xu, F.-L., 2014. Water quality benchmarking (WQB) and priority control screening (PCS) of persistent toxic substances (PTSs) in China: Necessity, method and a case study. *Sci. Total. Environ.* 472, 1108-1120.
- He, W., Qin, N., Kong, X., Liu, W., He, Q., Ouyang, H., Wang, Q., Yang, B., Yang, C., Jiang, Y., Xu, F., 2013a. Polybrominated diphenyl ethers (PBDEs) in the surface sediments and suspended particulate matter (SPM) from Lake Chaohu, a large shallow Chinese lake. *Sci. Total. Environ.* 463-464, 1163-1173.
- He, W., Qin, N., Kong, X., Liu, W., He, Q., Ouyang, H., Yang, C., Jiang, Y., Wang, Q., Yang, B., Xu, F., 2013b. Spatio-temporal distributions and the ecological and health risks of phthalate esters (PAEs) in the surface water of a large, shallow Chinese lake. *Sci. Total. Environ.* 461-462, 672-680.
- Hur, J., Lee, B.-M., Shin, K.-H., 2014. Spectroscopic characterization of dissolved organic matter isolates from sediments and the association with phenanthrene binding affinity. *Chemosphere* 111, 450-457.
- Jiang, Y.-J., He, W., Liu, W.-X., Qin, N., Ouyang, H.-L., Wang, Q.-M., Kong, X.-Z., He, Q.-S., Yang, C., Yang, B., Xu, F.-L., 2014. The seasonal and spatial variations of phytoplankton community and their correlation with environmental factors in a large eutrophic Chinese lake (Lake Chaohu). *Ecol. Indic.* 40, 58-67.
- Kang, L., Wang, Q.-M., He, Q.-S., He, W., Liu, W.-X., Kong, X.-Z., Yang, B., Yang, C., Jiang, Y.-J.,

- Xu, F.-L., 2015. Current status and historical variations of phthalate ester (PAE) contamination in the sediments from a large Chinese lake (Lake Chaohu). *Environ. Sci. Pollut. Res.*, 1-13.
- Kraaij, R., Mayer, P., Busser, F.J.M., Bolscher, M.V., Seinen, W., Tolls, J., 2003. Measured pore-water concentrations make equilibrium partitioning work - A data analysis. *Environ. Sci. Technol.* 37, 268-274.
- Li, Y.-L., He, W., Liu, W.-X., Kong, X.-Z., Yang, B., Yang, C., Xu, F.-L., 2015. Influences of binding to dissolved organic matter on hydrophobic organic compounds in a multi-contaminant system: Coefficients, mechanisms and ecological risks. *Environ. Pollut.* 206, 461-468.
- Liu, W.-X., He, W., Qin, N., Kong, X.-Z., He, Q.-S., Ouyang, H.-L., Xu, F.-L., 2013. The residues, distribution, and partition of organochlorine pesticides in the water, suspended solids, and sediments from a large Chinese lake (Lake Chaohu) during the high water level period. *Environ. Sci. Pollut. Res.* 20, 2033-2045.
- Martinez, A., O'Sullivan, C., Reible, D., Hornbuckle, K.C., 2013. Sediment pore water distribution coefficients of PCB congeners in enriched black carbon sediment. *Environ. Pollut.* 182, 357-363.
- Murphy, K.R., Butler, K.D., Spencer, R.G.M., Stedmon, C.A., Boehme, J.R., Aiken, G.R., 2010. Measurement of Dissolved Organic Matter Fluorescence in Aquatic Environments: An Interlaboratory Comparison. *Environ. Sci. Technol.* 44, 9405-9412.
- Neff, J.M., Stout, S.A., Gunster, D.G., 2005. Ecological risk assessment of polycyclic aromatic hydrocarbons in sediments: identifying sources and ecological hazard. *Integr. Environ. Assess. Manag.* 1, 22-33.
- Nguyen, T.H., Goss, K.U., Ball, W.P., 2005. Polyparameter linear free energy relationships for estimating the equilibrium partition of organic compounds between water and the natural organic matter in soils and sediments. *Environ. Sci. Technol.* 39, 913-924.
- Ololade, I.A., Oladoja, N.A., Lajide, L., Ololade, O.O., Ejelolu, B.C., Akinnifesi, T.A., Alaremu, A.G., 2012. Partitioning of polycyclic aromatic hydrocarbons in sediment and porewater from Ondo coastal area, Nigeria. *Environmentalist* 32, 363-370.
- Omwoma, S., Lalah, J.O., Virani, M., Schramm, K.W., Henkelmann, B., 2015. Dioxin-like PCBs and PCDD/Fs in surface sediments near the shore of Winam Gulf, Lake Victoria. *Chemosphere* 118, 143-147.
- Parr, T.B., Ohno, T., Cronan, C.S., Simon, K.S., 2014. comPARAFAC: a library and tools for rapid and quantitative comparison of dissolved organic matter components resolved by Parallel Factor Analysis. *Limnol. Oceanogr.-Meth.* 12, 114-125.
- Persson, N.J., Bucheli, T.D., Gustafsson, O., Broman, D., Naes, K., Ishaq, R., Zebuhr, Y., 2005. Testing common sediment-porewater distribution models for their ability to predict dissolved concentrations of POPs in The Grenlandsfjords, Norway. *Chemosphere* 59, 1475-1485.
- Pignatello, J.J., Xing, B.S., 1996. Mechanisms of slow sorption of organic chemicals to natural particles. *Environ. Sci. Technol.* 30, 1-11.
- Qiao, M., Huang, S., Wang, Z., 2008. Partitioning characteristics of PAHs between sediment and water in a shallow lake. *J. Soil. Sediment.* 8, 69-73.
- Qin, N., He, W., Kong, X.-Z., Liu, W.-X., He, Q.-S., Yang, B., Wang, Q.-M., Yang, C., Jiang, Y.-J., Jorgensen, S.E., Xu, F.-L., Zhao, X.-L., 2014. Distribution, partitioning and sources of polycyclic aromatic hydrocarbons in the water-SPM-sediment system of Lake Chaohu, China. *Sci. Total. Environ.* 496, 414-423.

- Shi, G.L., Tian, Y.Z., Guo, C.S., Feng, Y.C., Xu, J., Zhang, Y., 2012. Sediment-pore water partition of PAH source contributions to the Yellow River using two receptor models. *J. Soil. Sediment.* 12, 1154-1163.
- Singh, S., Inamdar, S., Scott, D., 2013. Comparison of two PARAFAC models of dissolved organic matter fluorescence for a mid-Atlantic forested watershed in the USA. *J. Ecosyst.* 2013, 532424.
- Smedes, F., van Vliet, L.A., Booij, K., 2013. Multi-Ratio Equilibrium Passive Sampling Method to Estimate Accessible and Pore Water Concentrations of Polycyclic Aromatic Hydrocarbons and Polychlorinated Biphenyls in Sediment. *Environ. Sci. Technol.* 47, 510-517.
- Sprynskyy, M., Ligor, T., Buszewski, B., 2008. Clinoptilolite in study of lindane and aldrin sorption processes from water solution. *J. Hazard. Mater.* 151, 570-577.
- Stedmon, C.A., Bro, R., 2008. Characterizing dissolved organic matter fluorescence with parallel factor analysis: a tutorial. *Limnol. Oceanogr.-Meth.* 6, 572-579.
- Tao, W., Xin, J., Congying, W., Fang, W., Yongrong, B., Guifen, Y., 2014. Adsorption of phenanthrene on Al (oxy)hydroxides formed under the influence of tannic acid. *Environ. Earth Sci.* 71, 773-782.
- tenHulscher, T.E.M., vanNoort, P.C.M., vanderVelde, L.E., 1997. Equilibrium partitioning theory overestimates chlorobenzene concentrations in sediment porewater from lake Ketelmeer, the Netherlands. *Chemosphere* 35, 2331-2344.
- Wang, D.-Z., Dong, H.-P., Xie, Z.-X., Dai, M.-H., Hong, H.-S., 2011. Metaproteomic characterization of dissolved organic matter in the water column of the South China Sea. *Limnol. Oceanogr.* 56, 1641-1652.
- Wen, Z.-D., Gao, D.-W., Li, Z., Ren, N.-Q., 2013. Effects of humic acid on phthalate adsorption to vermiculite. *Chem. Eng. J.* 223, 298-303.
- Williams, M.D., Adams, W.J., Parkerton, T.F., 1995. Sediment sorption coefficient measurements for 4 phthalate esters: Experimental results and model theory. *Environ. Toxicol. Chem.* 14, 1477-1486.
- Yamashita, Y., Cory, R.M., Nishioka, J., Kuma, K., Tanoue, E., Jaffe, R., 2010. Fluorescence characteristics of dissolved organic matter in the deep waters of the Okhotsk Sea and the northwestern North Pacific Ocean. *Deep-Sea Res. Pt. II* 57, 1478-1485.
- Yang, L., Han, D.H., Lee, B.M., Hur, J., 2015. Characterizing treated wastewaters of different industries using clustered fluorescence EEM-PARAFAC and FT-IR spectroscopy: Implications for downstream impact and source identification. *Chemosphere* 127, 222-228.
- Yates, K., Pollard, P., Davies, I., Webster, L., Moffat, C., 2013. Silicone rubber passive samplers for measuring pore water and exchangeable concentrations of polycyclic aromatic hydrocarbons concentrations in sediments. *Sci. Total. Environ.* 463, 988-996.
- Yu, Y., Xu, J., Wang, P., Sun, H.W., Dai, S.G., 2009. Sediment-porewater partition of polycyclic aromatic hydrocarbons (PAHs) from Lanzhou Reach of Yellow River, China. *J. Hazard. Mater.* 165, 494-500.
- Zheng, Z., He, P.J., Shao, L.M., Lee, D.J., 2007. Phthalic acid esters in dissolved fractions of landfill leachates. *Water Res.* 41, 4696-4702.

Figure captions

Fig. 1 Location of Lake Chaohu and the sampling sites.

Fig. 2 The measured sediment – pore water partition coefficients (K_D) of PAHs, OCPs, and PAEs ($DF > 0.15$) from limnic and riverine locations in Lake Chaohu. The error bar indicates the standard deviation.

Fig. 3 Comparison of experimental $\log K_{OC}$ values for individual PTOCs in Lake Chaohu sediments with the (a) experimental data from the EPI Suite database ($\log K_{OC,exp}$) and estimates of (b) the octanol-water partition coefficient ($\log K_{OC,OW}$), (c) molecular connectivity indices ($\log K_{OC,MCI}$), and (d) Raoult's law ($\log K_{OC,S}$). Figures containing more information refers to Fig. S2 in supplementary materials.

Fig. 4 Linear relationships between $\log K_{OC}$ and $\log K_{OW}$ for PAHs (a), OCPs (b), and PAEs (c). Retene, cyclopenta(c,d)pyrene, methoxychlor, DMP, DEP, DIBP, DEEP, and DNP were excluded to obtain the slope and intercept of linear regression.

Fig. 5 Principal component analysis' loading plot of PTOC partition coefficients and influencing factors. The circles, squares, diamonds, and pentangles denote PAHs, OCPs, PAEs, and influencing factors.

Fig. 6 The different partitioning mechanism of PAHs, OCPs, and PAEs and the potential influencing factors. The bold outline denotes the association degree between the partitioning behavior and influencing factors.

Fig. 1 Location of Lake Chaohu and the sampling sites.

Fig.2 The measured sediment – pore water partition coefficients (K_D) of PAHs, OCPs, and PAEs ($DF > 0.15$) from limnic and riverine locations in Lake Chaohu. The error bar indicates the standard deviation.

Fig. 3 Comparison of experimental $\log K_{OC}$ values for individual PTOCs in Lake Chaohu sediments with the (a) experimental data from the EPI Suite database ($\log K_{OC, exp}$) and estimates of (b) the octanol-water partition coefficient ($\log K_{OC, OW}$), (c) molecular connectivity indices ($\log K_{OC, MCI}$), and (d) Raoult's law ($\log K_{OC, S}$). Figures containing more information refers to Fig. S2 in supplementary materials.

Fig. 4 Linear relationships between $\log K_{OC}$ and $\log K_{OW}$ for PAHs (a), OCPs (b), and PAEs (c). Retene, cyclopenta(c,d)pyrene, methoxychlor, DMP, DEP, DIBP, DEEP, and DNP were excluded to obtain the slope and intercept of linear regression.

Fig. 5 Principal component analysis' loading plot of PTOC partition coefficients and influencing factors. The circles, squares, diamonds, and pentangles denote PAHs, OCPs, PAEs, and influencing factors.

Fig. 6 The different partitioning mechanism of PAHs, OCPs, and PAEs and the potential influencing factors. The bold outline denotes the association degree between the partitioning behavior and influencing factors.

Table 1 PAHs, OCPs, PAEs in the pore water and the sediment collected from limnic and riverine areas of Lake Chaohu, the particle sizes and total organic carbon (TOC) of the sediment, and the PARARAC-FOMs in both the AEOM and the pore water.

	Limnic locations	Riverine locations	All locations
Sediment			
PAHs (ng g ⁻¹)	1,467.76 (539.69 – 2,395.84) ^a	2,549.24 (197.66 – 4,900.83) ^a	1,980.04 (869.48 – 3,090.60)
OCPs (ng g ⁻¹)	123.27 (13.18 – 233.35) ^a	62.25 (7.60 – 116.90) ^a	94.36 (35.20 – 153.53)
PAEs (μg g ⁻¹)	322.56 (256.35 – 388.77) ^a	497.87 (378.56 – 617.18) ^b	405.60 (332.38 – 478.82)
Clay (%)	19.2 (15.8 – 22.6) ^a	16.0 (11.8 – 20.2) ^a	17.7 (15.2 – 20.2)
Slit (%)	42.8 (38.0 – 47.6) ^a	32.3 (25.0 – 39.5) ^b	37.8 (33.2 – 42.4)
Fine sand (%)	37.0 (29.6 – 44.5) ^a	48.6 (37.3 – 59.9) ^a	42.5 (35.9 – 49.0)
Coarse sand (%)	1.0 (0.3 – 1.7) ^a	3.2 (1.2 – 5.2) ^b	2.0 (1.0 – 3.1)
TOC (%)	1.19 (0.91 – 1.47) ^a	1.86 (0.46 – 3.27) ^a	1.51 (0.88 – 2.13)
C1 (R.U.)	0.016 (0.013 – 0.018) ^a	0.010 (0.004 – 0.016) ^a	0.013 (0.010 – 0.016)
C2 (R.U.)	0.010 (0.009 – 0.011) ^a	0.007 (0.006 – 0.008) ^b	0.009 (0.008 – 0.010)
C3 (R.U.)	0.060 (0.055 – 0.065) ^a	0.047 (0.036 – 0.058) ^b	0.054 (0.048 – 0.060)
C4 (R.U.)	0.021 (0.020 – 0.022) ^a	0.024 (0.020 – 0.027) ^a	0.022 (0.020 – 0.024)
C5 (R.U.)	0.002 (0.001 – 0.002) ^a	0.004 (0.002 – 0.006) ^b	0.003 (0.002 – 0.004)
C6 (R.U.)	0.023 (0.019 – 0.027) ^a	0.024 (0.016 – 0.032) ^a	0.023 (0.019 – 0.027)
Pore water			
PAHs (ng L ⁻¹)	0.51 (0.15 – 0.87) ^a	0.65 (0.33 – 0.97) ^a	0.58 (0.36 – 0.80)
OCPs (ng L ⁻¹)	0.069 (n.d. – 0.19) ^a	0.031 (0.014 – 0.048) ^a	0.051 (n.d. – 0.11)
PAEs (μg L ⁻¹)	0.076 (n.d. – 0.18) ^a	0.043 (0.003 – 0.082) ^a	0.060 (0.006 – 0.114)
C1 (R.U.)	0.417 (0.327 – 0.507) ^a	0.051 (0.025 – 0.077) ^b	0.244 (0.143 – 0.344)
C2 (R.U.)	0.137 (0.090 – 0.185) ^a	0.048 (0.028 – 0.068) ^b	0.095 (0.063 – 0.128)
C3 (R.U.)	0.022 (0.018 – 0.025) ^a	0.026 (0.023 – 0.029) ^a	0.024 (0.021 – 0.026)
C4 (R.U.)	0.011 (0.010 – 0.012) ^a	0.012 (0.010 – 0.013) ^a	0.011 (0.011 – 0.012)
C5 (R.U.)	0.082 (0.066 – 0.098) ^a	0.023 (0.016 – 0.030) ^b	0.054 (0.037 – 0.071)
C6 (R.U.)	0.033 (0.031 – 0.034) ^a	0.028 (0.025 – 0.031) ^b	0.031 (0.029 – 0.032)
Ratio of sedimentary FOM to pore water FOM			
C1 _{sed} /C1 _{pw}	0.041 (0.032 – 0.050) ^a	0.293 (0.070 – 0.517) ^b	0.161 (0.048 – 0.273)
C2 _{sed} /C2 _{pw}	0.086 (0.061 – 0.111) ^a	0.181 (0.108 – 0.254) ^b	0.131 (0.091 – 0.171)
C3 _{sed} /C3 _{pw}	2.901 (2.336 – 3.466) ^a	1.820 (1.408 – 2.232) ^b	2.389 (1.972 – 2.806)
C4 _{sed} /C4 _{pw}	1.870 (1.689 – 2.051) ^a	2.039 (1.762 – 2.316) ^a	1.950 (1.800 – 2.100)
C5 _{sed} /C5 _{pw}	0.025 (0.011 – 0.039) ^a	0.210 (0.107 – 0.313) ^b	0.113 (0.050 – 0.176)
C6 _{sed} /C6 _{pw}	0.697 (0.581 – 0.813) ^a	0.853 (0.571 – 1.135) ^a	0.771 (0.635 – 0.907)

Note: The differences between limnic and riverine locations were tested by independent t-test. The range values in the bracket were 95% confidence interval of the mean values. When the minimum data was < 0, it was replaced by “n.d.”, The superscripts after the data indicate the similarity and differences. The same alphabet indicates that the compared groups are similar. Different superscripts represent significant difference between the compared groups ($p < 0.05$).

Table 2 The variance of the ratio, detection frequency, partition coefficient ($\log K_D$), and organic carbon – water partition coefficient ($\log K_{OC}$) of PAHs, OCPs, and PAEs.

Chemical groups	Measured ratio (%)	Detection frequency	$\log K_D$	$\log K_{OC}$
PAHs	64.0	0.44 (0.24 – 0.63) ^a	3.28 (2.61 – 3.94) ^a	5.23 (4.61 – 5.86) ^a
OCPs	44.0	0.18 (0.05 – 0.31) ^b	2.40 (1.75 – 3.05) ^a	4.30 (3.62 – 4.97) ^b
PAEs	63.2	0.33 (0.13 – 0.52) ^{ab}	4.59 (4.13 – 5.05) ^b	6.50 (6.05 – 6.94) ^c

Note: measured ratio indicated the fraction (%) of the chemicals, whose $\log K_D$ or $\log K_{OC}$ could be measured in this study; the range values in the bracket were 95% confidence interval of the mean values. The differences among PAHs, OCPs, and PAEs were tested by independent t-test. The superscripts after the data indicate the similarity and differences. The same alphabet indicates that the compared groups are similar. Different superscripts represent significant difference between the compared groups ($p < 0.05$).

Table 3 The detection frequency (DF > 0), arithmetic mean, minimum (min), and maximum (max) of apparent $\log K_{OC}$ for PAHs, OCPs, and PAEs in sediment-pore water system of Lake Chaohu as well as chemicals' estimated $\log K_{OC}$ by octanol-water partition coefficient ($\log K_{OC, OW}$), molecular connectivity indices ($\log K_{OC, MCI}$), Raoult's law ($\log K_{OC, S}$), and measured $\log K_{OC}$ summarized in EPI Suite database ($\log K_{OC, exp}$).

Chemicals (Abbr.)	$\log K_{OC}$		$\log K_{OC, OW}$	$\log K_{OC, MCI}$	$\log K_{OC, S}$	$\log K_{OC, exp}$
	DF	Mean (min-max)				
Nap	1.00	4.10 (3.44-4.66)	2.86	3.19	1.27	2.96
Acy	1.00	4.60 (3.64-6.59)	3.42	3.70	1.63	3.75
Ace	1.00	4.38 (3.57-5.33)	3.40	3.70	2.25	3.59
Fle	1.00	4.55 (3.98-5.46)	3.63	3.96	2.64	3.70
Phe	1.00	4.62 (3.81-5.56)	3.87	4.22	2.84	4.35
Ant	0.95	4.62 (3.90-5.58)	3.86	4.21	4.27	4.31
Flu	1.00	5.56 (4.40-6.40)	4.48	4.74	3.54	4.80
Pyr	1.00	5.36 (4.14-6.18)	4.24	4.74	3.83	4.90
Ret	0.89	4.39 (3.51-5.34)	5.51	5.12	4.82	N.A.
CcdP	0.05	2.74 (2.74-2.74)	4.95	5.25	5.14	N.A.
BaA	0.89	5.69 (3.83-6.81)	5.00	5.25	5.04	5.30
Chr	0.68	6.25 (5.56-7.03)	5.04	5.26	5.71	N.A.
BbF	0.16	6.69 (5.84-7.33)	5.02	5.78	5.88	N.A.
BkF	0.11	6.64 (5.90-7.38)	5.30	5.77	6.15	5.60
BeP	0.11	6.99 (6.75-7.22)	5.59	5.78	5.25	N.A.
BaP	0.05	6.56 (6.56-6.56)	5.32	5.77	5.84	5.95
α -HCH	0.89	3.72 (2.19-4.90)	3.59	3.45	2.81	3.04
β -HCH	0.63	3.88 (2.10-5.50)	3.59	3.45	3.74	3.04
γ -HCH	0.42	3.80 (2.69-4.87)	3.59	3.45	2.25	3.04
δ -HCH	0.68	4.04 (2.55-5.33)	3.59	3.45	1.62	3.04
p,p'-DDE	0.16	5.31 (4.85-5.79)	5.65	5.07	4.55	4.82
o,p'-DDD	0.05	4.73 (4.73-4.73)	5.09	5.08	4.16	N.A.
o,p'-DDT	0.05	5.37 (5.37-5.37)	5.89	5.24	4.27	N.A.
p,p'-DDT	0.53	4.99 (4.13-6.17)	6.00	5.23	5.46	5.31
HCB	1.00	4.01 (3.00-5.06)	4.24	3.79	5.31	3.74
Ald	0.05	2.02 (2.02-2.02)	5.64	4.91	4.98	4.69
MOC	0.05	5.38 (5.38-5.38)	3.85	4.43	4.19	4.90
DMP	0.79	6.50 (5.66-7.34)	1.68	1.50	-0.66	1.60
DEP	0.89	6.63 (5.49-7.50)	2.13	2.02	-0.03	1.84
DIBP	0.89	6.61 (5.58-7.37)	3.07	2.91	2.30	3.14
DnBP	0.89	5.54 (4.17-6.52)	3.28	3.06	2.05	3.14
BMPP	0.05	6.17 (6.17-6.17)	4.35	3.96	3.99	N.A.
DEEP	0.32	7.89 (7.25-8.40)	1.77	1.84	-0.26	N.A.
BBP	0.89	6.35 (4.97-7.18)	3.41	3.86	2.72	3.72
HEHP	0.11	6.14 (5.80-6.49)	4.93	4.72	5.29	N.A.
DBEP	0.05	5.31 (5.31-5.31)	2.86	2.88	1.89	N.A.
DEHP	0.89	6.95 (6.27-7.63)	5.00	5.08	3.81	4.94

DnOP	0.37	7.28 (6.27-8.02)	5.27	5.15	4.90	4.38
DNP	0.05	6.57 (6.57-6.57)	6.06	5.67	7.66	N.A.