Title: Genetic and biogeochemical investigation of sedimentary nitrogen cycling communities responding to tidal and seasonal dynamics in Cape Fear River Estuary

Authors: Jessica A. Lisa^{1,2*}, Bongkeun Song^{1,2}, Craig R. Tobias³, David E. Hines¹

Affiliations:

¹Department of Biology and Marine Biology, University of North Carolina Wilmington, NC

²Department of Biological Sciences, Virginia Institute of Marine Science, College of William & Mary, Gloucester Point, VA

³Department of Marine Sciences, University of Connecticut, CT

*Current address of corresponding authors:

Jessica A. Lisa

Virginia Institute of Marine Science, College of William & Mary, 1375 Greate Road, Gloucester Point, VA 23062

Tel: 804.684.7382

E-mail: jalisa@vims.edu

Running Title: Temporal Dynamics of nitrogen cycling

Subject Categories: Nitrogen cycle; Denitrification; Nitrification; DNRA; Anammox;

Polymerase chain reaction;

ABSTRACT

Tidal and seasonal fluctuations in the oligohaline reaches of estuaries may alter geochemical features that influence structure and function of microbial communities involved in sedimentary nitrogen (N) cycling. In order to evaluate sediment community responses to short-term (tidal) and long-term (seasonal) changes in different tidal regimes, nitrogen cycling rates and genes were quantified in three sites that span a range of tidal influence in the upper portion of the Cape Fear River Estuary. Environmental parameters were monitored during low and high tides in winter and spring. ^{15}N tracer incubation experiments were conducted to measure nitrification, denitrification, anaerobic ammonium oxidation (anammox), and dissimilatory nitrate reduction to ammonia (DNRA). Abundances of functional genes including bacterial and archaeal ammonia monooxygenases (*amoA*), nitrite reductases (*nirS* and *nrfA*), nitrous oxide reductase (*nosZ*), and hydrazine oxidoreductase (*hzo*) were measured using quantitative PCR assays. Denitrification rates were highest among the measured N cycling processes while bacteria carrying *nrfA* genes were most abundant. A discernable pattern in the short-term variation of N cycling rates and gene abundance was not apparent under the different tidal regimes. Significant seasonal variation in nitrification, denitrification, and anammox rates as well as bacterial *amoA*, *nirS* and *nosZ* gene abundance was observed, largely explained by increases in substrate availability during winter, with sediment ammonium playing a central role. These results suggest that the coupling of nitrification to N removal pathways is primarily driven by organic carbon mineralization and independent of tidal or salinity changes. Finally, changes in denitrification and nitrification activities were strongly reflected by the abundance of the respective functional genes, supporting a linkage between the structure and function of microbial communities.

1.0 INTRODUCTION

Estuaries are highly dynamic systems that form the transition between freshwater and marine environments. The interface between these two environments supports a tight coupling of biogeochemical activities that are important in regulating the amount of fixed nitrogen (N) in coastal systems. Microbially mediated N recycling processes such as nitrification and dissimilatory nitrate reduction to ammonium (DNRA) occur simultaneously with the removal processes of denitrification and anaerobic ammonium oxidation (anammox) in estuarine sediments (Capone et al., 2008). Each pathway can be influenced by a wide range of interacting factors including: temperature, salinity, dissolved oxygen (DO), substrate availability (nitrate + nitrite; $\text{NO}_3^- + \text{NO}_2^-$), ammonium (NH_4^+), dissolved organic carbon (DOC), and hydrogen sulfide (H_2S ; Cornwell et al., 1999; Dalsgaard et al., 2005; Seitzinger et al., 2006). These environmental factors change over short and long temporal scales in estuarine ecosystems and thus, differentially influence the activities and structure of N cycling microbial communities on these time scales.

Tidal fluctuations in the oligohaline reaches of aquatic systems result in short-term changes in sediment and porewater chemistry, primarily as a result of movement of the water masses during the ebb and flood of the tide (Mortimer et al., 1999; Rocha and Cabral, 1998; Usi et al., 1998). Such changes in the supply of oxygen and salinity influence the availability of organic material as well as the species and distribution of N, creating a bottom up control on biogeochemical cycling. Expansion of the oxic zone as a result of tidal forcing may support higher rates of remineralization and nitrification in intertidal estuarine sediments (Mortimer et al., 1999; Rocha and Cabral, 1998). Enhanced nitrification can provide substrate for denitrification when NO_3^- diffuses to anoxic zones; however, inhibitory effects on denitrification

1
2
3
4 in intertidal sediments can also be sustained through prolonged oxygen exposure during ebb tide
5
6 or flushing of nutrients during flood tide (Thompson et al., 1995; Usi et al., 1998). In addition,
7
8 distinct differences in microbial community structure have been observed between high and low
9
10 tides, linked to variability in environmental conditions (Chauhan et al., 2009; Kara and Shade,
11
12 2009; Olapade, 2012). Temperature, light and nutrient availability contributed to greater bacterial
13
14 diversity and abundance observed during high and outgoing tide (Olapade, 2012) while
15
16 significant shifts in phylotype abundance occurred in response to elevated supply of dissolved
17
18 organics during low tide (Chauhan et al., 2009). These studies primarily examine the effects of
19
20 tidal fluctuation on planktonic microbial communities. However, little is known about the effects
21
22 of tidal exchange on benthic communities in the oligohaline reaches of estuaries, where a major
23
24 portion of N cycling takes place. Variations in N cycling activities and microbial community
25
26 structure are likely to occur as a result of changes in sediment chemistry and the movement of
27
28 different water masses with changing tides, greatly influencing estuarine N cycling.
29
30
31
32
33
34
35

36 Longerscale temporal shifts corresponding to seasonality influence sedimentary N
37
38 cycling processes in various ecosystems (Berounsky and Nixon, 1990; Dunn et al., 2013;
39
40 Eriksson et al., 2003; Jorgensen and Sorensen, 1985; Lisa et al., 2014; Rysgaard et al., 1995).
41
42 Literature reports of peak N cycling activity vary both for season and for the reaction described;
43
44 peaks often coincide with temperature as well as oxygen dynamics (Dunn et al., 2013; Rysgaard
45
46 et al., 1995) or weather events that provide additional sources of substrate to support elevated
47
48 processes (Eriksson et al., 2003; Lisa et al., 2014). Significant shifts in microbial community
49
50 composition and metabolism in have been observed seasonally (Desnues et al., 2007; Kristensen,
51
52 1993). This trend is also evident in autotrophic microbial communities where particular ammonia
53
54 oxidizing archetypes are more prevalent during specific seasons and correlated with temperature,
55
56
57
58
59
60
61
62
63
64
65

DON and NH_4^+ concentrations (Bouskill et al., 2011). The seasonal reoccurrence of ammonia oxidizing organisms suggests the return of indigenous communities following large perturbations in more stable systems.

The greatest changes in geochemical conditions occur in oligohaline reaches of estuaries where fluctuations in salinity and nutrients occur tidally and seasonally, yet little is known about the response of sedimentary N cycling activities and sediment microbial communities to short or long-term changes under these environmental conditions. The extent to which temporal variability in the environment influences both activities and structure of microbial communities together has largely escaped attention and can have implications for coupling of sedimentary N cycling processes and the overall ability to remove N from the ecosystem. It is therefore important to consider the effects of dynamic environmental conditions on microbial community structure and function when examining the temporal effects on estuarine biogeochemical cycling over short and long-term scales. The objective of this study was to investigate the linkage between the abundance and activities of microbial communities responsible for N cycling processes under short and long term changes in a tidal estuarine ecosystem. We examined the effects of temporal changes on nitrification, denitrification, anammox, and DNRA activities and respective functional gene abundance in subtidal estuarine sediments at three tidal sites in the oligohaline reaches of the Cape Fear River Estuary, NC, USA.

2.0 MATERIALS AND METHODS

2.1 Study System

The Cape Fear River Estuary (CFRE), located in southeastern North Carolina (Figure 1), makes up a 72 km long portion of the lower Cape Fear River proper and empties directly into

1
2
3
4 Long Bay (Mallin et al. 1999, Dafner et al. 2007). The Cape Fear River proper is a sixth-order
5
6 brown water stream originating in the Piedmont (Mallin et al., 1999). Two fifth-order blackwater
7
8 tributaries originating in the coastal plains, The Black and Northeast Cape Fear Rivers, also
9
10 empty into the lower portion of the Cape Fear River proper.
11
12

13
14 The CFRE watershed encompasses 23,310 km², the largest in North Carolina, and
15
16 supports one fifth of the population of the state. One half of the land within the basin is forested
17
18 while the remaining fifty percent is dedicated to cropland and pastureland or is urbanized (Lin et
19
20 al., 2006). Within the Cape Fear River basin, the most industrialized river basin in the state, are
21
22 641 licensed point discharges, a harbor, and state port in Wilmington (Mallin et al., 2000, 1999).
23
24 High levels of inorganic nutrients enter the system through these point discharges and additional
25
26 nonpoint sources such as runoff from urban, suburban, and livestock facilities.
27
28
29
30
31
32

33 *2.2 Seasonal Sampling of the CFRE*

34

35
36 Sampling was conducted during two seasons (winter and spring) over the course of a tidal
37
38 cycle (high and low tide) in the upper portion of the estuary in February and May of 2012.
39
40 Sampling sites were selected based upon tidal salinity variation. Salinity, a conservative
41
42 environmental parameter often used as a proxy for changes in other geochemical conditions, was
43
44 used to monitor the magnitude of tidal influence on our study sites. Sites included, Indian Creek
45
46 (IC; 34.2842N, 77.9981W), where salinity is <0.1 and invariant with tidal stage. Two other sites,
47
48 Navassa (NAV; 34.2589N, 77.9846W) and Horseshoe Bend (HB; 34.2422N, 77.9681W), had
49
50 larger changes in salinity throughout the tidal cycle, ranging from 0 to 15. At each site,
51
52
53 sampling was conducted in the channel and at the west bank (subtidal) during low and high tide.
54
55
56 This sampling scheme allowed for seasonal comparison between winter and spring at each site
57
58
59
60
61
62
63
64
65

1
2
3
4 under the two extreme tidal conditions. The design enabled us to deconvolute the effects of short
5
6 and long- temporal variation in geochemical conditions on N cycling activities and community
7
8 structure, seasonally and tidally.
9

10 11 12 13 14 *2.3 Environmental Parameter Measurements*

15
16 Environmental parameters including water column depth, temperature, salinity and
17
18 dissolved oxygen (DO), were measured within 1 meter of the bottom at the time of sampling
19
20 using a 6820 multi parameter YSI data sonde (YSI Incorporated, Yellow Springs, OH). Water
21
22 samples were 0.7 μ m (GF/F) filtered and stored on ice prior to nutrient analysis. Ammonium and
23
24 NO₃⁻ concentrations were measured spectrophotometrically on a Bran Luebbe segmented flow
25
26 nutrient auto analyzer using phenol hypochlorite and Cd-reduction/azo dye methods, respectively
27
28 following modified standard EPA methods (Long and Martin, 1997). Samples designated for
29
30 DOC and TDN analysis were stored in glass vials, preserved with H₃PO₄ and refrigerated until
31
32 analysis. The samples were analyzed within one week on a Shimadzu 5050A analyzer following
33
34 standard operating procedures.
35
36
37
38
39
40

41 Sediment samples were collected using a petite ponar grab (Wildco, Buffalo NY). The top
42
43 2 cm of sediment were collected and divided into aliquots to be characterized for solid phase
44
45 sediment properties, N cycling rate determinations, and microbial molecular analyses. Sediment
46
47 percent organic content was determined by loss on ignition (LOI) of dried sediments (500°C for
48
49 4 hours). Sediment NH₄⁺ (free plus extractable) was measured by phenol hypochlorite following
50
51 KCl extraction using a 1:1 ratio of 2M KCl to sediments (Mackin and Aller, 1984). Sediments
52
53 from each site were stored at -80°C for molecular analysis.
54
55
56
57
58
59
60
61
62
63
64
65

2.4 N cycling rates

Nitrification rates were determined under aerobic conditions using an isotope pool dilution method modified from Wessel & Tietema (1992). Incubations were conducted using 20g of sediment incubated in 80mL filtered site water amended with 70 μ M NO₃⁻ at 0.7 15N atom% (Carini et al. 2010). Initial and final samples were taken at time (t) t_i and t_f from the same incubation container and conducted for 24 hours in triplicate. ¹⁵NO₃⁻ was measured on an Elemental Analyzer Isotope Ratio Mass Spectrometer (EA-IRMS; Delta V, Thermo Fisher Scientific) following ¹⁵NO₃⁻ isolation using the Devarda's reduction and alkaline acid trap diffusion technique (Sigman et al. 1997, Holmes et al. 1998). Dilution of the isotope pool from t_i to t_f was used to calculate the rates of nitrification (Tobias et al. 2003).

Sediment slurry incubation experiments with ¹⁵N tracer were conducted to measure potential rates of denitrification and anammox following the method of Lisa et al. (2014). Production of ²⁹N₂, and ³⁰N₂ was measured on a Gas Bench - Isotopic Ratio Mass Spectrometer (GB-IRMS; Delta V Plus, Thermo Fisher Scientific, Waltham, MA) and used to calculate the rates of anammox and denitrification following the method of Thamdrup & Dalsgaard (2002) as modified by Song and Tobias (2011). Percent anammox (%anammox) was estimated based on the rates of anammox and total N₂ production in each sample.

The accumulation of ¹⁵NH₄⁺ in the same incubations was measured to calculate potential rates of DNRA (Tobias et al. 2001) according to the modifications of Song et al. (2014). Briefly, NH₄⁺ was isolated from the slurry by alkaline acid trap diffusion following the addition of 7 mL of 40 ppt NaCl solution, 0.15 g MgO, and 3 μ moles of unlabeled NH₄⁺ carrier. DNRA rate calculations were based on the concentration and ¹⁵N mole fraction excess of extractable NH₄⁺ as well as the ¹⁵NO₃⁻ mole fraction (Tobias et al. 2001). Percent DNRA (%DNRA) was estimated

1
2
3
4 based on the contribution of DNRA to total NO_3^- reduction (denitrification + anammox +
5
6 DNRA).
7
8
9

10 11 *2.5 Quantitative PCR Assays of functional genes* 12 13

14 Sediment DNA was extracted following the cetyltrimethylammonium bromide (CTAB)
15
16 method (DeAngelis et al., 2010; Griffiths et al., 2000) with modifications. Modifications
17
18 included the increase of wet sediment from 0.5 to 0.75 g and the use of a Thermo Savant Fast
19
20 Prep FP 120 Cell Disrupter (Qbiogene Inc. Carlsbad, CA) for cell disruption. DNA extractions
21
22 were carried out on IC and HB sites from winter and spring sampling events only.
23
24
25

26 Real time PCR assays were used to quantify abundance of functional genes in order to
27
28 obtain a quantitative measurement of the genetic potential a site has to carry out a particular
29
30 biogeochemical reaction. Primers targeting genes encoding the catalytic subunits of relevant
31
32 enzymes were used to quantify the abundances of microorganisms capable of N cycling
33
34 processes. Thermal cycling, fluorescent data collection, and data analysis were carried out using
35
36 the ABI 7500 Fast Real Time PCR System (Version 1.4). Assays were carried out in a volume of
37
38 20 μL containing 0.5 to 1.0 ng of template DNA and SYBR green using Go-Taq qPCR Master
39
40 Mix (Promega Corporation, Madison, WI). PCR specificity was confirmed using gel
41
42 electrophoresis to ensure amplification of the desired target and monitored by analysis of
43
44 dissociation curves. Standards were derived through a serial dilution of plasmids carrying
45
46 respective gene targets obtained from environmental samples. Standards were quantified using
47
48 Agilent 2200 TapeStation System (Agilent Technologies, Santa Clara, CA) following a digestion
49
50 with ECoR1. Information on PCR efficiency, R^2 values are reported below for each functional
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

gene. Efficiency of each qPCR reaction was calculated from the slope of the standard curve using the equation $\text{Efficiency} = -1 + 10^{(-1/\text{slope})}$.

Abundances of ammonia oxidizing bacteria and archaea were quantified by targeting ammonia monooxygenase (*amoA*) genes under the following PCR conditions: 95°C for 10 m; 50 cycles consisting of 95°C for 15 s, 53°C for 45 s, 72°C for 30 s, and 80°C for 30s (data acquisition); followed by dissociation step consisting of 95°C for 15 s, 60°C for 1 m, 95°C for 15 s, 60°C for 15 s. Bacterial *amoA* gene fragments were amplified using the PCR primers amoA-1F and amoA-2R (Gao et al., 2014; Rotthauwe et al., 1997) and archaeal *amoA* gene fragments were detected with Arch-amoAF and Arch-amoAR (Francis et al., 2005; He et al., 2007). Efficiencies for AOB and AOA were 85.09% and 98.03%, respectively. R^2 values were 0.99 and 0.98 for AOB and AOA, respectively.

Denitrifying bacterial abundance was assessed by quantifying nitrite reductase (*nirS*) and nitrous oxide reductase (*nosZ*) genes. Primers used to quantify *nirS* genes were nirS1F (Braker et al., 1998) and nirS-q-R (Mosier and Francis, 2010). The thermal cycling conditions were modified as follows: 95°C for 15 m; 38 cycles of 95°C for 15 s, 62.5°C for 30 s, 72°C for 30 s, and 84°C for 35 s (data acquisition), with a dissociation step of 95°C for 15 s, 60°C for 1 m, 95°C for 15 s, 60°C for 15 s. qPCR of *nosZ* genes were carried out using the primers nosZ2F and nosZ2R (Henry et al., 2006) under the following modified conditions: 95°C for 10 m; 50 cycles of 95°C for 15 s, 55°C for 45 s, 72°C for 35 s, and 75°C for 35 s (data acquisition), and a dissociation step of 95°C for 15 s, 60°C for 1 m, 95°C for 15 s, 60°C for 15 s. Efficiencies for *nirS* and *nosZ* were 83.30% and 74.17%, respectively. R^2 values were 0.99 for both *nirS* and *nosZ*.

Abundance of anammox bacteria was quantified using qPCR primers HZOQPCR1F and HZOQPCR1R following the methods of Long et al. (2013). Efficiency and R^2 values for *hzo* qPCR were 72.57% and 0.99, respectively. DNRA bacterial abundance was determined by quantifying *nrfA* gene abundance following the method of Song et al. (2014). Efficiency and R^2 values for the *nrfA* qPCR reaction were 60.29% and 0.99.

2.6 Statistical Analysis

Data were examined for normality using the Shapiro-Wilk Test and log transformed to meet the assumptions of statistical inference for parametric tests as well as reduce the range of the data. Paired-T Tests were used to identify differences among activities and gene abundance between tides and seasonally between paired samples. Pearson's product moment correlation analysis was used to investigate correlations between environmental conditions, potential N cycling activities and functional gene abundance. Linear regression analyses between N transformation rates and respective bacterial functional gene abundances were conducted to determine if relationships between the variables exist. Due to the robust nature of the analyses, $\alpha \leq 0.05$ was retained to delineate significant relationships between all response and explanatory variables. All statistical analyses were conducted in R (version 2.15.3 Copyright 2013 The R Foundation for Statistical Computing).

3.0 RESULTS

3.1 Environmental Parameters

Changes in salinity were negligible at the IC site ranging from 0.01 to 0.10 during both seasons (Supplementary Table 1). Changes in salinity during the winter sampling event were

greater at the two downstream sites, with salinity varying from 0.1 to 6.4 and up to 8.0 within the 6-hour tidal exchange at NAV and HB, respectively. The overall salinity at the downstream sites was higher during the spring low tide sampling, accompanied by lower amplitude of change between tidal stages. Bottom water temperature was constant throughout the sites and reflected typical seasonal conditions. DO in bottom water remained $>6 \text{ mg L}^{-1}$ during all sampling events.

Water column dissolved nutrients remained spatially uniform at the three sites within a tide and season, but were quite variable between tidal stage and season (Supplementary Table 2). Bottom water NO_3^- levels were elevated in the winter but remained relatively constant between tidal stages. Seasonal and tidal trends for bottom water NH_4^+ were not observed. DOC and TDN levels were both highest in winter. DOC was highly variable with the incoming tide while TDN consistently decreased with the incoming tide. The ratio of bottom water Carbon:Nitrogen ($\text{DOC}:\text{NO}_x$) was also higher during the winter and elevated at high tide at all sites.

Sediment %organics were variable across tidal stage and season and increased with downstream movement towards NAV and HB (Supplementary Table 3). Sediment extractable NH_4^+ was consistently low for both seasons. The greatest increase in sediment extractable NH_4^+ was also observed downstream at NAV and HB, with the incoming tide where the largest changes in salinity occurred.

3.2 N Cycling Rates & Their Correlation to Environmental Parameters

Tidal variations in all process rates - nitrification, denitrification, anammox, and DNRA - between low and high tide were not significantly different from one another ($p>0.05$; Figure 2) nor where the differences between bank and channel (data not shown). These data were pooled for subsequent analysis. Seasonal variations were specific to particular N cycling processes.

Several environmental parameters, including temperature, DO, and dissolved nutrients, showed strong seasonal variation and correlated with particular N cycling pathways, with sediment extractable NH_4^+ taking on a strong central role in connecting the N cycling activities.

Nitrification rates ranged from 0.02 to 5.72 nmols N g^{-1} wet sediment hour^{-1} and were significantly higher in the winter ($p < 0.001$; Supplementary Table 4a). Nitrification was negatively correlated with temperature, and positively correlated with DO levels, DOC, TDN, bottom water NH_4^+ , as well as sediment extractable NH_4^+ (Table 1).

Denitrification was the highest of all N cycling processes, ranging from 2.50 to 44.21 nmols N g^{-1} wet sediment hour^{-1} , with significantly higher rates also occurring in the winter ($p = 0.008$; Supplementary Table 4b). Temperature negatively correlated with denitrification rates (Table 1). Interestingly, increases in DO and sediment extractable NH_4^+ were accompanied by increases in denitrification activities.

Anammox activities were on the same order of magnitude as nitrification, ranging from 0.17 to 4.77 nmols N g^{-1} wet sediment hour^{-1} (Supplementary Table 4c). Highest anammox rates were also observed during winter ($p < 0.05$). Anammox rates were inversely correlated with temperature, but positively responded to increasing DO levels, DOC, TDN and bottom NH_4^+ (Table 1). Once again, sediment extractable NH_4^+ correlated with anammox N_2 production.

DNRA was the lowest of all N cycling processes measured throughout the study, ranging from 0.00 to 1.89 nmols N g^{-1} wet sediment hour^{-1} (Supplementary Table 4d). Differences in DNRA both tidally and seasonally were not significant for DNRA ($p > 0.05$). DNRA was positively correlated with salinity and negatively correlated with bottom water nitrate (Table 1). Sediment extractable NH_4^+ was also positively correlated with DNRA activities.

Mean %anammox was 11.58 and overall %anammox spanned the broad range of 1.9 to 44.6% contribution to N₂ production. %DNRA also varied, ranging from 0 to 24.7% with a mean of 2.56% contribution to NO₃⁻ reduction. Seasonal, spatial, and tidal variations and correlations with environmental parameters were not significant for either %anammox or %DNRA (data not shown).

3.3 Functional Gene Abundance & Its Correlation to Environmental Parameters

Variations in functional gene copy numbers showed no discernable pattern with respect to the different tidal stages and were not significantly different from one another between low and high tides (Figure 3). However, seasonal trends in gene abundance were evident and correlated with particular environmental parameters.

Bacterial and archaeal *amoA* genes showed very similar distribution patterns in abundance. Bacterial *amoA* gene abundance ranged from 1.60×10^3 to 3.78×10^5 , while archaeal *amoA* ranged from 2.31×10^3 to 2.53×10^5 gene copies g⁻¹ wet sediment (Supplementary Table 5a and 5b). Bacterial *amoA* gene copy number was significantly higher in the winter ($p=0.040$), while archaeal *amoA* gene abundance was not seasonally influenced ($p>0.05$). As a result, bacterial *amoA* genes comprised a higher proportion of total *amoA* gene abundance during the winter, but this trend did not hold during the spring. Abundance of AOB and AOA also correlated with differing environmental parameters. Bacterial *amoA* gene abundance was negatively correlated with temperature and positively correlated with DO (Table 1). Alternatively, archaeal *amoA* gene abundance was correlated with DOC (Table 1). Positive correlations with sediment extractable NH₄⁺ were common in both AOB and AOA communities.

Abundance of *nirS* genes ranged from 5.91×10^5 to 7.53×10^7 and *nosZ* genes ranged from 1.82×10^6 to 7.10×10^7 copies g^{-1} wet sediment (Supplementary Table 5c and 5d). Both *nirS* and *nosZ* genes were also most abundant during winter ($p=0.031$ and 0.045 respectively), but did not differ tidally. Several environmental parameters were commonly correlated with both *nirS* and *nosZ* gene abundance; these parameters included temperature, DO, and sediment extractable NH_4^+ (Table 1). TDN was the only parameter that correlated with *nirS* gene abundance.

Abundance of *hzo* genes ranged from 1.38×10^4 to 5.80×10^5 gene copies g^{-1} wet sediment (Supplementary Table 5e). Seasonal variations in *hzo* gene abundance were not present ($p>0.05$). Environmental parameters examined in this study showed no correlations with *hzo* gene abundance (Table 1). Bacteria possessing *nrfA* were the most abundant out of all groups in the examined communities, ranging from 5.79×10^6 to 1.34×10^9 gene copies g^{-1} wet sediment (Supplementary Table 5f). Abundance of *nrfA* was not significantly different between two seasons ($p>0.05$). Sediment organics and sediment extractable NH_4^+ positively correlated with *nrfA* gene copy number (Table 1).

3.4 Relationships Among Rates and Functional Gene Abundance

Changes in N cycling activities strongly reflected changes in their respective functional genes for denitrification and nitrification but not for anammox and DNRA (Figure 4). Nitrification rates were positively and significantly correlated with both bacterial and archaeal *amoA* gene abundances. Similarly, increases in denitrification rates significantly reflected increases in both *nirS* and *nosZ* gene abundances. However, anammox and DNRA activities were not correlated with abundance of their respective genes.

4.0 DISCUSSION

Overall, sedimentary N cycling processes in term of the rates and gene copy numbers measured in this study were comparable to previous studies conducted in similar environments. Denitrification rates and gene copy numbers were consistent with that of other estuaries of similar trophic status (Dalsgaard et al., 2005; Dong et al., 2009; Henderson et al., 2010; Henry et al., 2006; Seitzinger et al., 2006; Smith et al., 2015, 2007). Nitrification rates were within the lower quartile of published values (Caffrey et al., 2007) but gene abundance (AOA and AOB) were not remarkably different from others (AOB (Bernhard et al., 2007; Stehr et al., 1995) and AOA (Moin et al., 2009; Mosier and Francis, 2008)). Anammox, %anammox and *hzo* gene copy number were also consistent with the range of activities (Dalsgaard et al., 2005; Nicholls and Trimmer, 2009) and gene abundance (Dang et al., 2010; Lisa et al., 2014) reported in other estuarine systems. DNRA activities, however, were substantially lower than those measured in other estuaries (Tobias et al. 2001, Gardner et al. 2006, McCarthy et al. 2008) despite the abundance of *nrfA* genes similar to those found in other estuaries (Dong et al., 2009; Smith et al., 2015; Song et al., 2014). The low rates of DNRA in this study are consistent with findings in oligohaline systems (Giblin et al., 2013) and suggest that the conditions necessary to support DNRA were not met. Instead, denitrification is favored as the primary dissimilatory NO_x^- reduction process in the tidal reaches of the CFRE. Denitrification tends to dominate under higher NO_3^- conditions while DNRA is favored under low NO_3^- and high labile organic carbon environments (Koop-Jakobsen and Giblin, 2010; Smith et al., 2015; Tobias et al., 2001).

Short-term temporal variation in N cycling processes and functional gene abundance was observed in the upper CFRE during tidal exchange, however a significant, clear pattern was not

1
2
3
4 found. These results are contrary to several studies that observed significant changes in activities
5
6 and structure of microbial communities in intertidal sediments exposed to the atmosphere during
7
8 low tide. The difference in findings could be a result of a reduced capacity for solute exchange
9
10 due to slower diffusion in continually inundated sediments of CFRE. Under the circumstances
11
12 of continual inundation observed in this study, advection or seepage is less likely to accelerate
13
14 solute exchange as it would in intertidal systems thereby not affecting N cycling communities or
15
16 resulting activities simply with the changing tide. The difference in findings of this study could
17
18 also have to do with the slow growth rate of some of the microbes that mediate the various N
19
20 cycling processes. We might expect rapid changes in the structure and activities of denitrifying
21
22 and DNRA communities relative to nitrifying and anammox communities due to the versatile
23
24 lifestyles of the organisms. Additionally, we would also expect to see the most changes reflected
25
26 in the expression of the functional genes, not necessarily at the DNA level.
27
28
29
30
31
32

33 Nitrification, denitrification, and anammox rates and respective gene abundance (with the
34
35 exception of *hzo*) did show significant changes over the long-term, with elevated values observed
36
37 during winter. Temperature had the strongest correlation with nitrification, denitrification, and
38
39 anammox rates, as well as functional gene abundance in ammonia oxidizing and denitrifying
40
41 communities in the CFRE; however, co-variation between temperature and nutrients supply is
42
43 the likely factor contributing to higher winter activities.
44
45
46
47

48 Generally, metabolic processes and functional gene abundance positively respond to
49
50 increases in temperature, as is sometimes the case with nitrification rates and AOB abundance
51
52 (Berounsky and Nixon, 1990; Cébron et al., 2003), denitrification and denitrifier abundance
53
54 (Nowicki, 1994; Szukics et al., 2010). However, the inverse relationships between nitrification,
55
56 denitrification and anammox with temperature in this study, as well as the studies conducted by
57
58
59
60
61
62
63
64
65

Kemp et al. (1990) and Bernhard et al. (2007), suggest seasonally variable environmental factors other than temperature have a stronger control on activities and structure of microbial communities in the oligohaline reaches of CFRE.

The inverse relationships between N cycling processes and gene abundance with temperature in this study can be explained by a higher supply of nutrients, particularly DOC, TDN, and sediment extractable NH_4^+ , to the upper reaches of the CFRE during the winter. Taking the relationship of the three variables into consideration, it is likely that elevated N supply during the winter and lack of competition for DIN is driving the availability of dissolved N and sediment extractable NH_4^+ ($r=0.82$, $p<0.001$; $r=0.47$, $p=0.024$, respectively). This elevated supply of NH_4^+ during the winter may be responsible for higher winter nitrification activities and AOB abundance that can in turn support elevated denitrification and anammox activities in the CFRE.

Although not directly measured in this study, our data suggest the abundance of ammonia oxidizers and nitrification activities are, at least in part, coupled to denitrification and anammox rates and respective communities. This conclusion is corroborated by an earlier study in the upper CFRE, which showed 43% of denitrification activities were coupled to nitrification (Hines et al., 2012) as well as a significant positive correlation between nitrification and denitrification rates in this study ($p<0.001$, $r=0.733$). These findings are also supported by the occurrence of substrate induced stimulation of coupled nitrification-denitrification and nitrification-anammox in other marine and estuarine systems (Caffrey et al., 2007; Cornwell et al., 1999; Crowe et al., 2012; Lam et al., 2007; Rysgaard et al., 1995; Seitzinger, 1994). Caffrey et al (2007) also noted that sediment NH_4^+ concentrations, as a result of organic matter remineralization, were highly central to nutrient cycling; results from this study suggest this is also the case in the CFRE.

This genetic approach combined with the use of stable isotope tracer techniques to examine multiple N transformation processes offers a unique and holistic view of microbial community dynamics in the CFRE sediments under varying environmental conditions. A strong positive relationship between denitrifier community abundance and potential denitrification rates was observed in this study. Similar relationships between denitrification rates and *nirS* gene abundance were observed in other estuarine sediments, with an increase in rates and gene abundance supported by higher substrate concentrations (Dong et al., 2009; Smith et al., 2015). Despite low nitrifying activities, the reflection of changes in nitrification rates with changes in AOB and AOA gene abundance, indicate that a large portion of the nitrifying community is active. Significant correlations between nitrification rates with *amoA* gene abundance have also been observed in other coastal systems, linked to changes in salinity and substrate concentrations (Bernhard et al., 2007; Caffrey et al., 2007; Petersen et al., 2012). In the CFRE sediments, sediment NH_4^+ was the most important environmental factors correlated with denitrification and nitrification rates as well as functional gene abundance, highlighting the importance of organic carbon remineralization and the tight coupling of these metabolic processes with the abundance of their respective microbial communities.

A significant correlation between gene abundance and rates of anammox communities was not observed in this study, as in other estuarine sediments (Dong et al., 2009). Similarly, increases in the numbers of *nrfA* genes did correlate with increases in DNRA rates. While positive relationships between *nrfA* copies and DNRA rates have been observed (Dong et al. 2009, Song et al. 2014), decoupling between the genetic potential and rates for the process has been observed within the same system (Smith et al., 2015).

1
2
3
4 When we consider the what is known about the physiological nature of the organisms
5
6 involved in N metabolisms and the environmental conditions of the system, we can begin to
7
8 explain why genetic potential for a particular metabolic process may or may not predict the
9
10 potential for that process. In the case of nitrification and denitrification, gene abundance was
11
12 strongly correlated with activities, and largely a result of the favorable conditions of the system
13
14 as well as the ability of the organisms to respond to these conditions. The supply of NH_4^+
15
16 supported populations of both bacterial and archaeal nitrifiers that were active despite the rapidly
17
18 changing environment. These environmental conditions also supported a high abundance of
19
20 denitrifiers, directly and possibly through the coupling of nitrification. The strong correlations
21
22 between genetic potential and rates of nitrification and denitrification suggest that the
23
24 microorganisms present in these dynamic environments are well adapted and poised to respond
25
26 to the continually changing conditions in the tidal oligohaline reaches of the CFRE.
27
28
29
30
31
32

33 Alternatively, a decoupling between the genetic potential and activities of anammox and
34
35 DNRA is in part due to the physiology of the microorganisms and/or unfavorable environmental
36
37 conditions. Given the diversity of organisms capable of DNRA pathway (Kartal et al., 2007;
38
39 Mohan et al., 2004; Rutting et al., 2011; Simon, 2002), it is not particularly surprising that a
40
41 correlation between activities and *nrfA* gene abundance was not observed in the tidal reaches of
42
43 the CFRE. On the other hand, anammox bacteria prefer stable environmental conditions with a
44
45 continual and simultaneous low supply of substrate and oxygen. Thus, anammox bacteria are
46
47 typically out-competed for substrate by heterotrophic processes such as denitrification and
48
49 DNRA dynamic environments such as estuarine sediments (Dalsgaard et al., 2005; Dong et al.,
50
51 2011). The slow growing nature of anammox, low genetic potential, in a rapidly changing
52
53
54
55
56
57
58
59
60
61
62
63
64
65

environment put anammox at a disadvantage and resulted in the decoupling of gene abundance and observed anammox activities.

5.0 CONCLUSIONS

A unique experimental approach that combined the use of molecular and stable isotope techniques was used to examine multiple N transformation processes under varying environmental conditions. Denitrification activities were highest among the measured N cycling processes while bacteria capable of DNRA were most abundant. The strong centrality of sediment NH_4^+ levels and the potential for coupled nitrification-denitrification and nitrification-anammox supports the role of nutrients and substrate availability as drivers of these processes rather than temperature and salinity. Alternatively, salinity was found to positively influence DNRA activities. We observed significant correlations between nitrification activities and *amoA* gene abundances in both AOB and AOA, suggesting that both bacteria and archaea are equally important to NH_4^+ oxidation in the CFRE. Similar trends occurred with denitrification activities and abundance of denitrifying bacterial communities possessing both *nirS* and *nosZ* genes. The strong correlation between nitrifier and denitrifier microbial communities and rate measurements implies that abundances of microbial members is important in determining the magnitude of nitrification and denitrification activities.

6.0 ACKNOWLEDGMENTS

Funding for this research was provided by the US National Science Foundation, grant numbers (DEB1329273, DEB1020944, OCE0851435, EAR1024662). The authors thank L. B. Cahoon, S. Carini, M. A. Mallin, R. Whitehead and S. Borrett for their contributions to this

1
2
3
4 study. The authors also acknowledge K. Duernberger and M. McIever at UNCW's Center for
5
6 Marine Science, as well as The Laboratories of B. Song and C. R. Tobias, in particular A.

7
8 Arfken, K. Fridey, J. Taylor for assistance in seasonal sampling and processing. Finally, the
9
10 authors thank M. Hanke for providing help with manuscript figures as well as I. C. Anderson, D.
11
12 A. Bronk, M. J. Brush and B. B. Ward for providing comments and feedback on this manuscript.
13
14

15
16 This article is contribution number XXXX of the Virginia Institute of Marine Science, College of
17
18 William & Mary.
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

7.0 LITERATURE CITED

- Bernhard, A.E., Tucker, J., Giblin, A.E., Stahl, D.A., 2007. Functionally distinct communities of ammonia-oxidizing bacteria along an estuarine salinity gradient. *Environ. Microbiol.* 9, 1439–47.
- Berounsky, V.M., Nixon, S.W., 1990. Temperature and annual cycle of nitrification in waters of Narragansett Bay. *Limnol. Oceanogr.*
- Bouskill, N.J., Eveillard, D., O'Mullan, G., Jackson, G.A., Ward, B.B., 2011. Seasonal and annual reoccurrence in betaproteobacterial ammonia-oxidizing bacterial population structure. *Environ. Microbiol.* 13, 872–86.
- Braker, G., Fesefeldt, A., Witzel, K.P., 1998. Development of PCR primer systems for amplification of nitrite reductase genes (*nirK* and *nirS*) to detect denitrifying bacteria in environmental samples. *Appl. Environ. Microbiol.* 64, 3769–75.
- Caffrey, J.M., Bano, N., Kalanetra, K., Hollibaugh, J.T., 2007. Ammonia oxidation and ammonia-oxidizing bacteria and archaea from estuaries with differing histories of hypoxia. *ISME J.* 1, 660–2.
- Capone, D.G., Bronk, D.A., Mulholland, M.R., Carpenter, E.J., 2008. Nitrogen in the Marine Environment, Estuaries.
- Carini, S.A., McCarthy, M.J., Gardner, W.S., 2010. An isotope dilution method to measure nitrification rates in the northern Gulf of Mexico and other eutrophic waters. *Cont. Shelf Res.* 30, 1795–1801.
- Cébron, A., Berthe, T., Garnier, J., 2003. Nitrification and Nitrifying Bacteria in the Lower Seine River and Estuary (France). *Appl. Environ. Microbiol.* 69, 7091–7100.

- Chauhan, A., Cherrier, J., Williams, H.N., 2009. Impact of sideways and bottom-up control factors on bacterial community succession over a tidal cycle. *Proc. Natl. Acad. Sci. U. S. A.* 106, 4301–4306.
- Cornwell, J.C., Kemp, W.M., Kana, T.M., 1999. Denitrification in coastal ecosystems : methods , environmental controls , and ecosystem level controls , a review. *Aquat. Ecol.* 33, 41–54.
- Crowe, S.A., Canfield, D.E., Mucci, A., Sundby, B., Maranger, R., 2012. Anammox, denitrification and fixed-nitrogen removal in sediments from the Lower St. Lawrence Estuary. *Biogeosciences* 9, 4309–4321.
- Dafner, E. V., Mallin, M.A., Souza, J.J., Wells, H.A., Parsons, D.C., 2007. Nitrogen and phosphorus species in the coastal and shelf waters of Southeastern North Carolina, Mid-Atlantic U.S. coast. *Mar. Chem.* 103, 289–303.
- Dalsgaard, T., Thamdrup, B., Canfield, D.E., 2005. Anaerobic ammonium oxidation (anammox) in the marine environment. *Res. Microbiol.* 156, 457–64.
- Dang, H., Chen, R., Wang, L., Guo, L., Chen, P., Tang, Z., Tian, F., Li, S., Klotz, M.G., 2010. Environmental factors shape sediment anammox bacterial communities in hypernutrified Jiaozhou Bay, China. *Appl. Environ. Microbiol.* 76, 7036–47.
- DeAngelis, K.M., Silver, W.L., Thompson, A.W., Firestone, M.K., 2010. Microbial communities acclimate to recurring changes in soil redox potential status. *Environ. Microbiol.* 12, 3137–49.
- Desnues, C., Michotey, V.D., Wieland, A., Zhizang, C., Fourçans, A., Duran, R., Bonin, P.C., 2007. Seasonal and diel distributions of denitrifying and bacterial communities in a hypersaline microbial mat (Camargue, France). *Water Res.* 41, 3407–3419.

- 1
2
3
4 Dong, L.F., Smith, C.J., Papaspyrou, S., Stott, A., Osborn, A.M., Nedwell, D.B., 2009. Changes
5
6 in benthic denitrification, nitrate ammonification, and anammox process rates and nitrate
7
8 and nitrite reductase gene abundances along an estuarine nutrient gradient (the Colne
9
10 estuary, United Kingdom). *Appl. Environ. Microbiol.* 75, 3171–9.
11
12
13
14 Dong, L.F., Sobey, N., Smith, C.J., Rusmana, I., Phillips, W., Stott, A., Osborn, A.M., Nedwell,
15
16 D.B., 2011. Dissimilatory reduction of nitrate to ammonium , not denitrification or
17
18 anammox , dominates benthic nitrate reduction in tropical estuaries. *Limnology Oceanogr.*
19
20 56, 279–291.
21
22
23
24 Dunn, R.J.K., Robertson, D., Teasdale, P.R., Waltham, N.J., Welsh, D.T., 2013. Benthic
25
26 metabolism and nitrogen dynamics in an urbanised tidal creek: Domination of DNRA over
27
28 denitrification as a nitrate reduction pathway. *Estuar. Coast. Shelf Sci.* 131, 271–281.
29
30
31 Eriksson, P.G., Svensson, J.M., Carrer, G.M., 2003. Temporal changes and spatial variation of
32
33 soil oxygen consumption, nitrification and denitrification rates in a tidal salt marsh of the
34
35 Lagoon of Venice, Italy. *Estuar. Coast. Shelf Sci.* 58, 861–871.
36
37
38 Francis, C. a, Roberts, K.J., Beman, J.M., Santoro, A.E., Oakley, B.B., 2005. Ubiquity and
39
40 diversity of ammonia-oxidizing archaea in water columns and sediments of the ocean. *Proc.*
41
42 *Natl. Acad. Sci. U. S. A.* 102, 14683–8.
43
44
45 Gao, J., Luo, X., Wu, G., Li, T., Peng, Y., 2014. Abundance and diversity based on amoA genes
46
47 of ammonia-oxidizing archaea and bacteria in ten wastewater treatment systems. *Appl.*
48
49 *Microbiol. Biotechnol.* 98, 3339–54.
50
51
52
53 Gardner, W., McCarthy, M., An, S., Sobolev, D., Sell, K.S., Brock, D., 2006. Nitrogen fixation
54
55 and dissimilatory nitrate reduction to ammonium (DNRA) support nitrogen dynamics in
56
57 Texas estuaries. *Limnol. Oceanogr.* 51, 558–568.
58
59
60
61
62
63
64
65

- Giblin, A., Tobias, C.R., Song, B., Weston, N., Banta, G.T., Rivera-Monroy, V.H., 2013. The Importance of Dissimilatory Nitrate Reduction to Ammonium (DNRA) in the Nitrogen Cycle of Coastal Ecosystems. *Oceanography* 26, 124–131.
- Griffiths, R.I., Whiteley, A.S., O'Donnell, A.G., Bailey, M.J., 2000. Rapid Method for Coextraction of DNA and RNA from Natural Environments for Analysis of Ribosomal DNA- and rRNA-Based Microbial Community Composition. *Appl. Environ. Microbiol.* 66, 5488–5491.
- He, J.-Z., Shen, J.-P., Zhang, L.-M., Zhu, Y.-G., Zheng, Y.-M., Xu, M.-G., Di, H., 2007. Quantitative analyses of the abundance and composition of ammonia-oxidizing bacteria and ammonia-oxidizing archaea of a Chinese upland red soil under long-term fertilization practices. *Environ. Microbiol.* 9, 2364–74.
- Henderson, S.L., Dandie, C.E., Patten, C.L., Zebarth, B.J., Burton, D.L., Trevors, J.T., Goyer, C., 2010. Changes in denitrifier abundance, denitrification gene mRNA levels, nitrous oxide emissions, and denitrification in anoxic soil microcosms amended with glucose and plant residues. *Appl. Environ. Microbiol.* 76, 2155–64.
- Henry, S., Bru, D., Stres, B., Hallet, S., Philippot, L., 2006. Quantitative detection of the *nosZ* gene, encoding nitrous oxide reductase, and comparison of the abundances of 16S rRNA, *narG*, *nirK*, and *nosZ* genes in soils. *Appl. Environ. Microbiol.* 72, 5181–9.
- Hines, D.E., Lisa, J.A., Song, B., Tobias, C.R., Borrett, S.R., 2012. A network model shows the importance of coupled processes in the microbial N cycle in the Cape Fear River Estuary. *Estuar. Coast. Shelf Sci.* 106, 45–57.

- Holmes, R.M., McClelland, J.W., Sigman, D.M., Fry, B., Peterson, B.J., 1998. Measuring ^{15}N - NH_4 in marine, estuarine and fresh waters: An adaptation of the ammonia diffusion method for samples with low ammonium concentrations. *Mar. Chem.* 60, 235–243.
- Jorgensen, B.B., Sorensen, J., 1985. Seasonal cycles of O_2 , NO_3^- , and SO_4^{2-} reduction in estuarine sediments: the significance of an NO_3^- reduction maximum in spring. *Mar. Ecol. Prog. Ser.* 24, 65–74.
- Kara, E., Shade, A., 2009. Temporal dynamics of South End tidal creek (Sapelo Island, Georgia) bacterial communities. *Appl. Environ. Microbiol.* 75, 1058–1064.
- Kartal, B., Kuypers, M.M.M., Lavik, G., Schalk, J., Op den Camp, H.J.M., Jetten, M.S.M., Strous, M., 2007. Anammox bacteria disguised as denitrifiers: nitrate reduction to dinitrogen gas via nitrite and ammonium. *Environ. Microbiol.* 9, 635–42.
- Kemp, W., Sampou, P., Caffrey, J., Mayer, M., Henrikson, K., Boynton, W., 1990. Ammonium recycling versus denitrification Chesapeake Bay sediments. *Limnol. Oceanogr.* 35, 1545–1563.
- Koop-Jakobsen, K., Giblin, A.E., 2010. The effect of increased nitrate loading on nitrate reduction via denitrification and DNRA in salt marsh sediments. *Limnol. Oceanogr.* 55, 789–802.
- Kristensen, E., 1993. Seasonal Variations in Benthic Community Metabolism and Nitrogen Dynamics in a Shallow, Organic-Poor Danish Lagoon. *Estuar. Coast. Shelf Sci.* 36, 565–586.
- Lam, P., Jensen, M.M., Lavik, G., McGinnis, D.F., Müller, B., Schubert, C.J., Amann, R., Thamdrup, B., Kuypers, M.M.M., 2007. Linking crenarchaeal and bacterial nitrification to anammox in the Black Sea. *Proc. Natl. Acad. Sci. U. S. A.* 104, 7104–7109.

- Lin, J., Xie, L., Pietrafesa, L.J., Shen, J., Mallin, M. a., Durako, M.J., 2006. Dissolved oxygen stratification in two micro-tidal partially-mixed estuaries. *Estuar. Coast. Shelf Sci.* 70, 423–437.
- Lisa, J., Song, B., Tobias, C., Duernberger, K., 2014. Impacts of freshwater flushing on anammox community structure and activities in the New River Estuary, USA. *Aquat. Microb. Ecol.* 72, 17–31.
- Long, A., Heitman, J., Tobias, C., Philips, R., Song, B., 2013. Co-occurring anammox, denitrification, and codenitrification in agricultural soils. *Appl. Environ. Microbiol.* 79, 168–76.
- Long, S.E., Martin, T., 1997. Method for the determination of chemical substances in marine and estuarine environmental matrices - 2nd edition U.S. Environmental Protection Agency. National Service for Environmental Publications: Washington DC.
- Mackin, J.E., Aller, R.C., 1984. Ammonium adsorption in marine sediments. *Limnol. Oceanogr.* 29, 250–257.
- Mallin, M.A., Burkholder, J.M., Cahoon, L.B., Posey, M.H., 2000. North and South Carolina Coasts. *Mar. Pollut. Bull.* 41, 56–75.
- Mallin, M.A., Cahoon, L.B., McIver, M.R., Parsons, D.C., Shank, G.C., 1999. Alternation of Factors Limiting Phytoplankton Production in the Cape Fear River Estuary. *Estuaries* 22, 825.
- McCarthy, M.J., McNeal, K.S., Morse, J.W., Gardner, W.S., 2008. Bottom-water Hypoxia Effects on Sediment–Water Interface Nitrogen Transformations in a Seasonally Hypoxic, Shallow Bay (Corpus Christi Bay, TX, USA). *Estuaries and Coasts* 31, 521–531.

- 1
2
3
4 Mohan, S.B., Schmid, M., Jetten, M., Cole, J., 2004. Detection and widespread distribution of
5
6 the *nrfA* gene encoding nitrite reduction to ammonia, a short circuit in the biological
7
8 nitrogen cycle that competes with denitrification. *FEMS Microbiol. Ecol.* 49, 433–443.
9
10
11 Moin, N.S., Nelson, K.A., Bush, A., Bernhard, A.E., 2009. Distribution and diversity of archaeal
12
13 and bacterial ammonia oxidizers in salt marsh sediments. *Appl. Environ. Microbiol.* 75,
14
15 7461–8.
16
17
18 Mortimer, R.J.G., Krom, M.D., Watson, P.G., Frickers, P.E., Davey, J.T., Clifton, R.J., 1999.
19
20 Sediment-Water Exchange of Nutrients in the Intertidal Zone of the Humber Estuary, UK.
21
22 *Mar. Pollut. Bull.* 37, 261–279.
23
24
25 Mosier, A.C., Francis, C.A., 2008. Relative abundance and diversity of ammonia-oxidizing
26
27 archaea and bacteria in the San Francisco Bay estuary. *Environ. Microbiol.* 10, 3002–16.
28
29
30 Mosier, A.C., Francis, C.A., 2010. Denitrifier abundance and activity across the San Francisco
31
32 Bay estuary. *Environ. Microbiol. Rep.* 2, 667–76.
33
34
35 Nicholls, J.C., Trimmer, M., 2009. Widespread occurrence of the anammox reaction in estuarine
36
37 sediments. *Aquat. Microb. Ecol.* 55, 105–113.
38
39
40 Nowicki, B.L., 1994. The effect of temperature, oxygen, salinity, and nutrient enrichment on
41
42 estuarine denitrification rates measured with a modified nitrogen gas flux technique. *Estuar.*
43
44 *Coast. Shelf Sci.* 38, 137–156.
45
46
47 Olapade, O.A., 2012. Diel Fluctuations in the Abundance and Community Diversity of Coastal
48
49 Bacterioplankton Assemblages over a Tidal Cycle. *Microb. Ecol.* 63, 96–102.
50
51
52 Petersen, D.G., Blazewicz, S.J., Firestone, M., Herman, D.J., Turetsky, M., Waldrop, M., 2012.
53
54 Abundance of microbial genes associated with nitrogen cycling as indices of
55
56
57
58
59
60
61
62
63
64
65

- biogeochemical process rates across a vegetation gradient in Alaska. *Environ. Microbiol.* 14, 993–1008.
- Rocha, C., Cabral, A.P., 1998. The Influence of Tidal Action on Porewater Nitrate Concentration and Dynamics in Intertidal Sediments of the Sado Estuary. *Estuaries* 21, 635–645.
- Rotthauwe, J.H., Witzel, K.P., Liesack, W., 1997. The ammonia monooxygenase structural gene *amoA* as a functional marker: molecular fine-scale analysis of natural ammonia-oxidizing populations. *Appl. Environ. Microbiol.* 63, 4704–12.
- Rutting, T., Boeckx, P., Muller, C., Klemetsson, L., 2011. Assessment of the importance of dissimilatory nitrate reduction to ammonium for the terrestrial nitrogen cycle. *Biogeosciences* 8, 1779–1791.
- Rysgaard, S., Christensen, P.B., Nielsen, L.P., 1995. Seasonal variation in nitrification and denitrification in estuarine sediment colonized by benthic microalgae and bioturbating infauna. *Mar. Ecol. Prog. Ser.*
- Seitzinger, S., 1994. Linkages between organic matter mineralization and denitrification in eight riparian wetlands. *Biogeochemistry*.
- Seitzinger, S., Harrison, J. a, Böhlke, J.K., Bouwman, a F., Lowrance, R., Peterson, B., Tobias, C., Van Drecht, G., 2006. Denitrification across landscapes and waterscapes: a synthesis. *Ecol. Appl.* 16, 2064–90.
- Sigman, D.M., Altabet, M. a., Michener, R., McCorkle, D.C., Fry, B., Holmes, R.M., 1997. Natural abundance-level measurement of the nitrogen isotopic composition of oceanic nitrate: an adaptation of the ammonia diffusion method. *Mar. Chem.* 57, 227–242.
- Simon, J., 2002. Enzymology and bioenergetics of respiratory nitrite ammonification. *FEMS Microbiol. Rev.* 26, 285–309.

- 1
2
3
4 Smith, C.J., Dong, L.F., Wilson, J., Stott, A., Osborn, A.M., Nedwell, D.B., 2015. Seasonal
5
6 variation in denitrification and dissimilatory nitrate reduction to ammonia process rates and
7
8 corresponding key functional genes along an estuarine nitrate gradient. *Front. Microbiol.* 6,
9
10 1–11.
11
12
13
14 Smith, C.J., Nedwell, D.B., Dong, L.F., Osborn, a M., 2007. Diversity and abundance of nitrate
15
16 reductase genes (*narG* and *napA*), nitrite reductase genes (*nirS* and *nrfA*), and their
17
18 transcripts in estuarine sediments. *Appl. Environ. Microbiol.* 73, 3612–22.
19
20
21 Song, B., Lisa, J.A., Tobias, C.R., 2014. Linking DNRA community structure and activity in a
22
23 shallow lagoonal estuarine system. *Front. Microbiol.* 5, 1–10.
24
25
26 Song, B., Tobias, C.R., 2011. Molecular and stable isotope methods to detect and measure
27
28 anaerobic ammonium oxidation (anammox) in aquatic ecosystems. In: *Methods in*
29
30 *Enzymology*. Elsevier Inc., pp. 63–89.
31
32
33
34 Stehr, G., Bottcher, B., Dittberner, P., Rath, G., Koops, H.-P., 1995. The ammonia-oxidizing
35
36 nitrifying population of the River Elbe estuary. *FEMS Microbiol. Ecol.* ... 17, 177–186.
37
38
39 Szukics, U., Abell, G.C.J., Hödl, V., Mitter, B., Sessitsch, A., Hackl, E., Zechmeister-
40
41 Boltensern, S., 2010. Nitrifiers and denitrifiers respond rapidly to changed moisture and
42
43 increasing temperature in a pristine forest soil. *FEMS Microbiol. Ecol.* 72, 395–406.
44
45
46 Thamdrup, B., Dalsgaard, T., 2002. Production of N₂ through Anaerobic Ammonium Oxidation
47
48 Coupled to Nitrate Reduction in Marine Sediments. *Appl. Environ. Microbiol.* 68, 1312–
49
50 1318.
51
52
53 Thompson, S.P., Paerl, H.W., Go, M.C., 1995. Seasonal Patterns of Nitrification and
54
55 Denitrification in a Natural and a Restored Salt Marsh. *Estuaries*.
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4 Tobias, C., Anderson, I., Canuel, E., Macko, S., 2001. Nitrogen cycling through a fringing
5
6 marsh-aquifer ecotone. *Mar. Ecol. Prog. Ser.* 210, 25–39.
7
8
9 Tobias, C., Giblin, A., McClelland, J., Tucker, J., Peterson, B., 2003. Sediment DIN fluxes and
10
11 preferential recycling of benthic microalgal nitrogen in a shallow macrotidal estuary. *Mar.*
12
13 *Ecol. Prog. Ser.* 257, 25–36.
14
15
16 Usi, T., Koike, I., Ogura, N., 1998. Tidal Effect on Dynamics of Pore Water Nitrate in Intertidal
17
18 Sediment of a Eutrophic Estuary. *J. Oceanogr.* 54, 205–216.
19
20
21 Wessel, W., Tietema, A., 1992. Calculating gross N transformation rates of ^{15}N pool dilution
22
23 experiments with acid forest litter: Analytical and numerical approaches. *Soil Biol.*
24
25 *Biochem.* 24, 931–942.
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

8.0 Table and Figure Legends

Table 1. Correlation analysis of N cycling activities and gene abundance with environmental parameters. Bold values indicate significant correlations.

Supplementary Table 1. Water column physical parameters in the Cape Fear River Estuary.

Supplementary Table 2. Water column dissolved nutrients in the Cape Fear River Estuary.

Supplementary Table 3. Sediment characteristics of the Cape Fear River Estuary.

Supplementary Table 4. Tidal and seasonal fluctuations in sedimentary N cycling rates in the Cape Fear River Estuary. NA indicates Not Available. N cycling values are presented in $\text{nmol N g}^{-1} \text{ sediment hr}^{-1}$. "SD" indicates standard deviation.

Supplementary Table 5. Abundance of relevant N cycling genes in the Cape Fear River Estuary during low and high tides. NA indicates Not Available.

Figure 1. Sampling sites in the Cape Fear River Estuary, NC, USA. Sampling sites in the upper, oligohaline portion of the CFRE include Indian Creek (IC), Navassa (NAV) and Horseshoe bend (HB).

Figure 2. Fluctuation of sedimentary nitrogen cycling rates at high and low tides. N cycling values are presented in $\text{nmol N g}^{-1} \text{ sediment hr}^{-1}$ for two of six total sites, the banks of Indian Creek (IC) and Horseshoe bend (HB) during winter. Rate measurements are presented as follows: A) nitrification, B) denitrification, C) anammox, and D) dissimilatory nitrate reduction to ammonium (DNRA). Error bars represent standard deviation.

Figure 3. Sedimentary nitrogen cycling functional gene abundance during low and high tides. Gene copy numbers are presented for two of six total sites, the banks of Indian Creek (IC) and Horseshoe bend (HB) during winter only. Genes include: A) bacterial ammonia monooxygenase (Bacterial *amoA*), B) archaeal ammonia monooxygenase (Archaeal *amoA*), C) nitrite reductase (*nirS*), D) nitrous oxide reductase (*nosZ*), E) hydrazine oxidoreductase (*hzo*), and F) cytochrome C nitrite reductase (*nrfA*). Error bars represent standard deviation.

Figure 4. Correlation analyses of nitrogen cycling rates and respective functional gene abundance in the Cape Fear River Estuary. Correlations are presented for: A) nitrification and bacterial ammonia monooxygenase (Bacterial *amoA*), B) nitrification and archaeal ammonia monooxygenase (Archaeal *amoA*), C) denitrification and nitrite reductase (*nirS*), D) denitrification and nitrous oxide reductase (*nosZ*), E) anammox and hydrazine oxidoreductase (*hzo*), and F) dissimilatory nitrate reductase to ammonium (DNRA) and cytochrome C nitrite reductase (*nrfA*). Error bars represent standard deviation.

Table 1

	Bottom Water												Sediment					
	Salinity		Temperature		DO		DOC		TDN		Nitrate		Ammonium		Organics		Ammonium	
	<i>r</i>	<i>p</i> -value	<i>r</i>	<i>p</i> -value	<i>r</i>	<i>p</i> -value	<i>r</i>	<i>p</i> -value	<i>r</i>	<i>p</i> -value	<i>r</i>	<i>p</i> -value	<i>r</i>	<i>p</i> -value	<i>r</i>	<i>p</i> -value	<i>r</i>	<i>p</i> -value
Nitrification	-0.07	0.747	-0.82	<0.001	0.80	<0.001	0.78	<0.001	0.68	<0.001	-0.27	0.205	0.52	0.010	0.14	0.526	0.75	0.000
Denitrification	0.19	0.384	-0.45	0.027	0.43	0.037	0.38	0.066	0.30	0.149	-0.40	0.052	0.32	0.122	0.27	0.215	0.67	0.000
Anammox	-0.08	0.722	-0.57	0.003	0.55	0.005	0.49	0.016	0.49	0.015	-0.14	0.525	0.32	0.128	0.01	0.950	0.61	0.002
DNRA	0.48	0.034	0.00	0.996	-0.03	0.898	-0.03	0.912	-0.14	0.560	-0.74	<0.001	-0.05	0.831	0.41	0.079	0.49	0.034
Bacterial <i>amo</i> A	0.17	0.569	-0.68	0.008	0.66	0.011	0.44	0.111	0.51	0.060	-0.30	0.305	0.37	0.199	0.18	0.529	0.62	0.023
Archaeal <i>amo</i> A	0.12	0.713	-0.52	0.081	0.53	0.073	0.40	0.194	0.42	0.169	-0.35	0.271	0.16	0.615	0.28	0.380	0.64	0.035
<i>nir</i> S	-0.27	0.332	-0.65	0.009	0.67	0.006	0.37	0.180	0.59	0.021	-0.13	0.641	0.29	0.298	-0.21	0.462	0.69	0.006
<i>nos</i> Z	0.25	0.368	-0.59	0.021	0.55	0.033	0.39	0.153	0.44	0.105	-0.39	0.154	0.36	0.194	0.30	0.284	0.66	0.011
<i>hzo</i>	-0.13	0.642	-0.47	0.080	0.49	0.061	0.41	0.125	0.45	0.094	-0.16	0.577	0.04	0.892	0.20	0.469	0.47	0.092
<i>nrf</i> A	0.39	0.156	-0.06	0.840	0.03	0.913	0.18	0.533	-0.09	0.756	-0.46	0.084	0.06	0.845	0.82	<0.001	0.55	0.041

Figure 1

Figure 2

Figure 4