The Mexican Drought Atlas:

Tree-Ring Reconstructions of the Soil Moisture Balance During the Late Pre-Hispanic, Colonial, and Modern Eras

David W. Stahle^a, Edward R. Cook^b, Dorian J. Burnette^c, Jose Villanueva^d,
Julian Cerano^d, Jordan N. Burns^e, Daniel Griffin^f, Benjamin I. Cook^g, Rodolfo Acuña^h,
Max C.A. Torbenson^a, Paul Sjeznerⁱ, and Ian M. Howard^a

Quaternary Science Reviews (re-submission June 27, 2016)

^aDepartment of Geosciences

University of Arkansas

Fayetteville, AR 72701

^bLamont-Doherty Earth Observatory

Columbia University

Palisades, NY 10964

^cDepartment of Earth Sciences

University of Memphis

Memphis, TN 38152

23 ^dINIFAP: Centro Nacional de Investigacion Disciplinaria Relacion Agua-Suelo-Planta-
24 Atmosfera, Lerdo, Mexico

25

26 ^eThe Nature Conservancy

27 Denver, CO 80203

28

29 ^fDepartment of Geography, Environment, and Society

30 University of Minnesota

31 Minneapolis, MN 55455

32

33 ^gNASA Goddard Institute for Space Studies

34 New York, NY 10025

35

36 ^hUniversidad Nacional Autonoma de Mexico

37 Mexico City, Mexico

38

39 ⁱLaboratory of Tree-Ring Research

40 University of Arizona

41 Tucson, AZ 85721

42

Abstract

Mexico has suffered a long history and prehistory of severe sustained drought. Drought over Mexico is modulated by ocean-atmospheric variability in the Atlantic and Pacific, raising the possibility for long-range seasonal climate forecasting, which could help mediate the economic and social impacts of future dry spells. The instrumental record of Mexican climate is very limited before 1920, but tree-ring chronologies developed from old-growth forests in Mexico can provide an excellent proxy representation of the spatial pattern and intensity of past moisture regimes useful for the analysis of climate dynamics and climate impacts. The Mexican Drought Atlas (MXDA) has been developed from an extensive network of 252 climate sensitive tree-ring chronologies in and near Mexico. The MXDA reconstruction extends from AD 1400 to 2012 and were calibrated with the instrumental summer (JJA) self-calibrating Palmer Drought Severity Index (scPDSI) on a 0.5° latitude/longitude grid extending over land areas from 14-34°N and 75-120°W using Ensemble Point-by-Point Regression (EPPR) for the 1944-1984 period. The grid point reconstructions were validated for the period 1920-1943 against instrumental gridded scPDSI values based on the fewer weather station observations available during that interval. The MXDA provides a new spatial perspective on the historical impacts of moisture extremes over Mexico during the past 600-years, including the Aztec Drought of One Rabbit in 1454, the drought of El Año de Hambre in 1785-1786, and the drought that preceded the Mexican Revolution of 1909-1910.

The El Niño/Southern Oscillation (ENSO) is the most important ocean-atmospheric forcing of moisture variability detected with the MXDA. In fact, the

reconstructions suggest that the strongest central equatorial Pacific sea surface temperature (SST) teleconnection to the soil moisture balance over North America may reside in northern Mexico. This ENSO signal has stronger and more time-stable correlations than computed for either the Atlantic Multidecadal Oscillation or Pacific Decadal Oscillation. The extended Multivariate ENSO Index is most highly correlated with reconstructed scPDSI over northern Mexico, where warm events favor moist conditions during the winter, spring, and early summer. This ENSO teleconnection to northern Mexico has been strong over the past 150 years, but it has been comparatively weak and non-stationary in the MXDA over central and southern Mexico where eastern tropical Pacific and Caribbean/tropical Atlantic SSTs seem to be more important. The ENSO teleconnection to northern Mexico is weaker in the available instrumental PDSI, but analyses based on the millennium climate simulations with the Community Earth System Model suggest that the moisture balance during the winter, spring, and early summer over northern Mexico may indeed be particularly sensitive to ENSO forcing. Nationwide drought is predicted to become more common with anthropogenic climate change, but the MXDA reconstructions indicate that intense “All Mexico” droughts have been rare over the past 600 years and their frequency does not appear to have increased substantially in recent decades.

1. Introduction

Aridity prevails over most of Mexico, including the Valley of Mexico, the seat of powerful pre-Hispanic city-states and the location of modern Mexico City, one of the largest urban centers in the world. The impressive pre-Hispanic archaeological record of cultural development and decline in Mexico appears to have been influenced in part by severe decadal drought (Coe 1984, Hodell et al. 1995, Brenner et al. 2001, Haug et al. 2003). The post-Conquest colonial and modern record is rich in detail on prolonged drought, famines, epidemic disease, international migration, and warfare. However, Mexico does not have a high quality nationwide instrumental climate record before 1920, and the available weather recording stations are in fact sparsely distributed prior to 1940 (Juaregui 1979, Douglas 2007). There are also very few centuries-long, high-resolution paleoclimate records in Mexico with which to test hypotheses concerning climate change, the internal and external forcing of climate variability, or the possible role of climate in socioeconomic extremes during the late pre-Hispanic, colonial, or early modern eras. In this article we describe the Mexican Drought Atlas (MXDA) that provides annually resolved and spatially detailed tree-ring reconstructions of the summer moisture balance from AD 1400 to 2012 and now covers the entire Republic of Mexico. These reconstructions are most reliable for the last 400 years during the colonial and modern periods when the network of predictor tree-ring chronologies is most spatially complete and internally well replicated.

Severe sustained drought has caused serious socioeconomic impacts in Mexican history (Liverman 1990; O'Hara and Metcalfe 1997) and has been implicated in political

unrest and social change among colonial and modern societies (Florescano 1980, 1986; Dell 2012). Ironically, human influences may now be aggravating drought severity and persistence over 21st century Mexico, as may have occurred during the pre-Hispanic era with widespread anthropogenic deforestation in the Mayan Lowlands (Cook et al. 2012). Extensive regional land cover changes can impact moisture exchange between the land surface and atmosphere, and may be partially responsible for the dramatic escalation of maximum afternoon temperatures recorded during the last 40 years over central Mexico (Englehart and Douglas 2005, Pascual et al. 2015). Simultaneously, the accumulation of anthropogenic greenhouse gases may be causing both rising global temperatures and the intensification of the Hadley circulation, with potential regional consequences over Mexico (Seager et al. 2007, Feng and Fu 2013, Cook et al. 2015a). Modern industrialized Mexico is not immune to drought and this vulnerability crosses many socioeconomic sectors, from subsistence farming to industrial agriculture, and from municipal and industrial water supply to hydroelectric power generation.

The recurrence and societal impacts of drought may be partially predictable because Mexican hydroclimatic variability is often modulated by large-scale ocean-atmospheric interactions, including the El Niño/Southern Oscillation (ENSO). However, the impact of these remote influences on Mexican climate varies by latitude and season, and from one event to another. There are also many unanswered questions regarding the stability of these large-scale teleconnections, and the magnitude of climate variance they might actually explain, which have bearing on the potential predictability of seasonal drought and wetness over Mexico. Tree-ring chronologies can provide an

exactly-dated, annually-resolved extension of the climate record into the centuries prior to instrumental observations to better understand the magnitude and duration of drought and wetness extremes, their frequency of recurrence, and the climate dynamics involved.

The North American Drought Atlas (NADA) was developed by Cook et al. (1999, 2004, 2007, 2010) and provided tree-ring reconstructions of the Palmer Drought Severity Index on a regularized grid over most of North America, including much of Mexico. However, very few tree-ring chronologies were available for inclusion in the NADA from central and southern Mexico. The NADA was also restricted to chronologies dating back to AD 1700 or earlier. The network of climate sensitive tree-ring chronologies in and near Mexico has been greatly expanded since the development of the NADA, including many new chronologies at least 150-years in length that now cover most of the Republic of Mexico, largely exclusive of the Baja and Yucatan peninsulas. The Mexican Drought Atlas has been developed to utilize this greatly expanded network of tree-ring chronologies to reconstruct the summer (JJA) self-calibrating Palmer Drought Severity Index (scPDSI) on a 0.5° latitude/longitude grid. These gridded reconstructions are linked continuously with the instrumental PDSI data and extend from CE 1400-2012. The MXDA now provides a modern and pre-modern perspective on hydroclimatic variability over Mexico that will be useful in the study of social and environmental change. The MXDA will also be useful for investigating the influence of ENSO and other ocean-atmospheric interactions on Mexican climate and will help document the natural climate background to anthropogenic climate changes now apparently taking place over Mexico.

This article begins with a review of the large-scale influences on climate variability and the socioeconomic impact of drought over Mexico. The network of tree-ring chronologies and the calculation of the MXDA are described, along with the derived calibration and validation statistics. The MXDA is then used to document the spatial extent and severity of extreme droughts and pluvials in Mexico over the past 600-years, including during drought events identified with pre-Hispanic and historical records. The history of inter-annual to decadal variability in soil moisture is estimated for four large sub-regions of Mexico where the tree-ring network is best replicated. The influence of ENSO, the Atlantic Multidecadal Oscillation (AMO), and the Pacific Decadal Oscillation (PDO) on instrumental and reconstructed PDSI are examined beginning in the 1870s when direct measurements of ENSO and the AMO become available. The ENSO teleconnection appears to influence development of a north-south anti-phasing or “dipole” in moisture variability over Mexico, but climate modeling suggests that anthropogenic global warming may overwhelm this gradient and lead to an increase in nationwide “All Mexico” dryness. The frequency of these two spatial patterns of moisture variability is examined over the past 600-years with the new tree-ring reconstructions in the MXDA.

2. Natural climate variability, tree growth, and anthropogenic climate change in Mexico

The precipitation climatology of Mexico is subject to a substantial seasonal cycle (Mosino and Garcia 1974, Cavazos and Hastenrath 1990, Metcalfe et al. 2015).

177 Modest cool season precipitation amounts over Mexico are often associated with
178 “*nortes*” (the penetration of midlatitude frontal systems). The major warm season
179 (JJAS) precipitation maxima is part of the North American Monsoon System (NAMS)
180 and represents more than 60% of the annual precipitation total for much of northwestern
181 Mexico (Adams and Comrie 1997, Higgins et al. 1999). In southern Mexico warm
182 season precipitation is influenced by the “*temporales*,” episodic surges of the Inter-
183 Tropical Convergence Zone (ITCZ) and increased precipitation into Central America
184 (Portig 1965, Metcalfe et al. 2015). Surges from the ITCZ into Central America may
185 cover up to 800 km per day and are influenced in part by troughs in the geopotential
186 height field over central North America (Portig 1965).

187 The interannual variability of climate in Mexico and Central America appears to
188 be mediated by at least three large-scale modes of ocean-atmospheric variability, most
189 notably by ENSO (Cavazos and Hastenrath 1990, Magaña et al. 2003). The AMO and
190 the PDO have also been linked empirically and experimentally with the severity and
191 persistence of drought over Mesoamerica (Pavia et al. 2006, Mendez and Magaña
192 2010, Stahle et al. 2011b, Feng et al. 2011). The impact of these remote influences on
193 Mesoamerican climate, particularly the ENSO teleconnection, varies by latitude and
194 season such that El Niño events tend to be associated with wetness in winter over
195 northern Mexico and dryness in summer over southern Mexico and Central America
196 (Magaña et al. 2003, Seager et al. 2009). Large-scale forcing of the NAMS appears to
197 be weaker than the teleconnections linked with cool season precipitation (Nicholas and
198 Battisti 2008), but ENSO influences (Castro et al. 2001), SSTs in the Atlantic and
199 Pacific, and anomalies in the warm pools of the Caribbean and tropical eastern North

Pacific have been implicated in the inter-annual variability of the NAMS (Carleton et al. 1990, Higgins et al. 1999, Castro et al. 2001, Sheppard et al. 2002, Vera et al. 2006, Douglas 2007, Metcalfe et al. 2015).

The main moisture signal recorded by long-lived trees in and near Mexico occurs during the winter, spring, and early summer (e.g., Cleaveland et al. 2003, Villanueva et al. 2007, Stahle et al. 2011a). In Mexico itself, the radial growth of Douglas-fir (*Pseudotsugamenziesii*) and Montezuma baldcypress (*Taxodiummucronatum*), two of the principal species used in the MXDA, appears to frequently end during the *canicula* or the mid-summer drought that typically occurs in July and August (Magaña et al. 1999). However, heavy precipitation late in the summer season, including rainfall associated with landfalling tropical systems, can greatly enhance warm season totals and is capable of reversing long-term hydrological drought in some drainage basins (e.g., Nicholas and Battisti 2008; Woodhouse et al. 2007). Unfortunately, these very late season rainfall events are not well represented in the Mexican tree-ring proxies. Nevertheless, the available tree-ring chronologies are highly correlated with the moisture balance during the early summer season, and with the summer-long average (JJA) of the scPDSI due to the real world persistence of soil moisture regimes and the built in month-to-month persistence structure of the Palmer drought index (see section IIIb, below). The JJA scPDSI has therefore been selected as the best variable for reconstruction in the MXDA using the available tree-ring data. Heavy precipitation in the late summer and autumn season may influence soil moisture and tree growth in the following year and can be captured to some extent in the MXDA. But dramatic reversals in the moisture balance from the early to late warm season may not be well

represented by the MXDA, and this may explain a few differences noted between the high quality PDSI estimates in the MXDA and selected historical evidence for late summer moisture (e.g., 1521-1524 and 1808-1811, discussed below).

Major fluctuations in precipitation and temperature have occurred over Mexico during the instrumental period and larger changes are projected by the end of the 21st century (Seager et al. 2009). The two most extreme and prolonged precipitation deficits occurred over Mexico during the 1950s and from the 1990s to early 2000s (Stahle et al. 2009). The severity and persistence of the recent 'Turn-of-the-century' drought may have been a partial consequence of the significant increase in surface air temperatures over Mexico (Cortez Vazquez 2006, Pascual et al. 2015). The impacts of early 21st century droughts have been severe for the agricultural and water resources sectors (Federman et al. 2014, Carter et al. 2015), and have induced widespread emigration from drought-affected sectors of Mexico into the United States (Feng et al. 2010, Hunter et al. 2013).

Ensemble analyses of 27 climate models driven by business-as-usual accumulations of greenhouse gases and land cover changes suggest that Mexico, the southwestern United States, and Central America will experience progressive drying over the 21st century (Feng and Fu 2013). The sharply rising surface air temperatures over Mexico (Pascual et al. 2015) may be an indication that the projected anthropogenic climate change is already underway. In fact, the intense warming is dominated by a rise in maximum daily temperature that may be a partial consequence of land cover change in Mexico (Englehart and Douglas 2005). Overgrazing, deforestation, and the extensive land conversion for human use and settlement are

believed to have reduced evaporative cooling and sharply increased the sensible to latent heat flux, favoring higher daily temperature maxima and overwhelming the potential cooling effect of increased surface albedo (e.g., Balling 1988, Englehart and Douglas 2005). The influence of anthropogenic land cover change on Mexican climate may have some precedent in the archaeological record. Model simulations of the climate consequences of deforestation by the ancient Maya in the lowland Yucatan indicate heightened drought conditions (Cook et al. 2012), which in turn may have contributed to the decline of certain Mayan city-states near the close of the Classic Period. The potential parallels with the land surface changes taking place across modern Mexico are provocative, and the “anthropogenic drought” that threatens modern society may have precedent in the long history of cultural and environmental change in Mexico.

3. The socioeconomic impacts of drought in Mexico

The modern economy of Mexico varies from subsistence agriculture in some of the most poverty-stricken regions in the Western Hemisphere to the globally integrated society of Mexico City. Mexico remains vulnerable to drought across this full spectrum of economic development, but especially among the poor practicing subsistence agriculture on *Ejidos* (i.e., communal lands; Dell 2012). The impact of drought on small stakeholders may have been heightened by economic changes arising from the North American Free Trade Agreement (NAFTA) and other reforms, which have tended to increase production costs, crop prices, and price volatility with serious negative impacts

on small producers (Liverman et al 1999, Wilder and Lankao 2006). Drought sensitivity among the poor may have also been increased by market-style changes in Mexican water law that have led to higher water prices in some cases (Seager et al. 2009). This political and economic squeezing of the poor is not new in Mexico (Simpson 1966). Large landowners and speculators controlled some of the best agricultural property in the 18th century and were able to manipulate crop prices during shortages, which aggravated the effects of drought among the rural poor (Florescano 1980, 1986; Liverman 1999).

Drought impacts are usually most acute among the poor (Eakin 2000), but the impacts of recent drought in Mexico have not been restricted to the underprivileged. The drought of 2011-2013 necessitated rural food relief in northern Mexico, but also resulted in an estimated 40% reduction in national agricultural production (Allen et al 2012). Central Mexico may be the most vulnerable region of the country to anthropogenic climate change because of the high population density, the enormous water and hydroelectric power consumption, and its position between the semi-arid north and more humid south (Seager et al. 2009). The socioeconomic vulnerability of urbanized central Mexico arises in part from the energy sector. Mexico City receives 70% of its electrical supply from hydroelectric facilities in southern Mexico, especially Chiapas (Lanza et al. 2002). Annual precipitation totals have declined by some measures over Chiapas from 1950 to 2000 (Golicher 2006, Douglas 2007, Seager et al. 2009) simultaneously with the sharp rise in maximum temperatures over central Mexico (Pascual et al. 2015), which has increased evapotranspiration, energy demand, and water consumption. This escalating consumption has led to declining surface and sub-

surface water supplies, and has placed increased pressure on water supply and distribution systems (Liverman 1999, Wilder and Lankao 2006).

Climate model simulations forecast increasing aridity over Mexico and Central America (Seager et al. 2009). Drier conditions are expected to appear by mid-century and increase in severity by 2100 (Feng and Fu 2013). The simulations also suggest that the aridity gradient will intensify southward across southern Mexico and into Central America, roughly in parallel with a gradient of increasing poverty. In fact, the spatial signature of future anthropogenic drought in Mexico may be nationwide (Seager et al. 2009), an “All Mexico” drought anomaly pattern that has not occurred frequently in the instrumental record and would pose a major challenge to Mexican society.

Consequently, some of the most severe future drought impacts may be felt among the most vulnerable populations in Latin America. The Mexican Drought Atlas has been developed to advance understanding of past drought and wetness extremes, the natural climate dynamics involved in drought recurrence, and the anthropogenic factors that may be leading to increased frequency and severity of drought over Mexico.

4. Development of the Mexican Drought Atlas

4.1. Tree-ring data

The forests of Mexico are among the most diverse on Earth and include the centers of biodiversity for the genera *Quercus* and *Pinus* (oaks and pines; Standley 1967). However, most tree species native to tropical Mexico do not form reliable, anatomically distinctive annual growth rings, including some native oak and pine

species. Fortunately, two of the most important and long-lived species for dendroclimatology in North America, Douglas-fir and Montezuma baldcypress, reach the southern limits of their distribution in Mexico and Guatemala (Figure 1). Douglas-fir is extensively present in the Sierra Madre Occidental of northwest Mexico, but is only found in small isolated populations in the Sierra Madre Oriental of northeast Mexico, the Trans-Mexican Volcanic Belt of central Mexico, and in the Sierra Madre del Sur of southern Mexico (Martinez 1963, Little 1991; Figure 1a). Tree-ring data from 49 Douglas-fir sites are currently available for Mexico, a few of which are over 500 years long. Montezuma baldcypress, the National Tree of Mexico, is widely distributed across Mexico, excluding the Baja and Yucatan peninsulas (Martinez 1963; Figure 1b). Eighteen Montezuma baldcypress ring-width collections were used for the MXDA, all obtained from riparian habitats above 300m in elevation where individuals can achieve great age, but where stands can also be heavily disturbed by human activity. Montezuma baldcypress is closely related to swamp baldcypress (*T. distichum*; Figure 1b) of the southeastern United States, and both species are reliable proxies of precipitation and the PDSI in spite of their riverine habitat (Stahle et al. 1988, 1998, Stahle and Cleaveland 1992, Davidson et al. 2006, Villanueva et al. 2007). A few tree-ring chronologies have also been developed in Mexico from *Abies* sp. (Meko et al. 2013) and several species of *Pinus* (Diaz et al. 2001, Fule et al. 2011, Cerano-Paredes et al. 2013), but they do not tend to be as long or widely distributed as Douglas-fir or baldcypress.

The network of Mexican chronologies has been augmented for the MXDA with chronologies available from the southern United States and western Guatemala (Figure

2). Some of these chronologies were included in the North American Drought Atlas (Cook et al. 2004, 2007, 2010), but many have been developed since the NADA was created and others did not meet the minimum length requirements of the NADA. Only a few tree-ring chronologies from Mexico are more than 600-years long and many are less than 300-years in length (Figure 2). For this reason, the MXDA has been limited to the period after AD 1400 when several tree-ring chronologies are available from northern, central, and southern Mexico.

A total of 252 ring width collections from the southern United States, Mexico, and western Guatemala were used to develop the MXDA, including 83 from Mexico. The signal free method of ring width standardization (Melvin and Briffa 2008, Cook et al. 2014) was used to detrend each measured ring width time series and to compute all tree-ring chronologies used for climate reconstruction with the MXDA. Signal freedetrending is designed to preserve high-to-medium frequency variance (e.g., less than the mean length of the component series) that is common to the ring-width samples being processed. It also helps avoid backward propagating trend distortion effects caused by the fitted growth curves being biased by a large common growth signal towards the end of the tree-ring time series caused by, for example, global warming signal found in many boreal conifers (Melvin and Briffa 2008) or from severe decadal drought that has afflicted much of Mexico and the southwestern United States for the past 20 years. The ring width data were detrended with an age-dependent cubic smoothing spline (Cook and Peters 1981, Melvin et al. 2007) where the degree of spline flexibility and ring width data tracking are a function of the length (age) of the individual time series. This detrending approach ensures that the low frequency variability in the

derived chronology will be based primarily on the longest ring width time series from the oldest trees.

The ring width indices were computed as residuals from the spline curve fit to the raw measurements after the ring widths were first power transformed to reduce bias in the resulting detrended tree-ring indices (Cook and Peters 1997). The derived ring width indices were then averaged into the mean index chronology using the biweight robust mean (Hoaglin et al. 2000) to reduce the influence of outliers on the calculation of the mean. After all chronologies were computed with the signal free method some excessive low frequency variance remained in a few individual series, especially among some long baldcypress chronologies from the southern United States. To ensure a more homogenous expression of centennial timescale variance across the MXDA tree-ring chronology network, a 500-year smoothing spline was applied to all mean index chronologies prior to their use in reconstruction. This modest level of high-pass filtering, uniformly applied to all chronologies, preserves annual, decadal, to centennial-scale variability across the network but guards against spurious multi-centennial timescale tree growth variance that might arise from human or other non-climatic disturbances.

4.2. Instrumental PDSI and reconstruction methods

The station-based observations of monthly temperature and precipitation compiled by the Climatic Research Unit at the University of East Anglia were used in gridded form (Harris et al. 2014) to develop the gridded self-calibrating Palmer drought severity index (scPDSI) by van der Schrier et al. (2013). These gridded scPDSI values include 1501 grid points on a 0.5° latitude/longitude field from 14° to 34°N and 75° to

120°W over Mexico and adjacent land areas (Figure 3), and were used as the target instrumental climate field for reconstruction from tree rings. Most Mexican tree-ring chronologies respond to precipitation primarily in the late winter, spring, and early summer (Cleaveland et al. 2003, Villanueva et al. 2007, Stahle et al. 2011a). Nonetheless, the high month-to-month Markovian persistence term of 0.897 prescribed in the calculation of the Palmer index (Palmer 1965), intended to model true persistence in the soil moisture balance, means that June PDSI is very highly correlated with PDSI subsequently recorded in July and August (Figure 4). The area of weakest correlation between June and subsequent July-August average scPDSI over subtropical North America is actually computed for portions of western Mexico influenced by the North American Monsoon (Figure 4), but even here the correlation during the instrumental period 1901-2012 does not drop below $r = +0.60$. The tree-ring chronologies used to develop the MXDA are well correlated with the JJA-average scPDSI and with Palmer drought indices throughout the year. Consequently, the summer (JJA) average scPDSI, when evapotranspiration demand is often highest, is a reasonable though not perfect compromise for development of the moisture balance reconstructions in the MXDA. It is true that tree-ring chronologies used for the MXDA are not well correlated with late summer or autumn rainfall extremes, but the statistical and historical evidence presented below indicates that the MXDA does provide important new insights into the socioeconomic and environmental impacts of droughts and pluvials over Mexico, and the internal and external climate factors that drive those moisture changes.

The Ensemble Point-by-Point Regression (EPPR) method of Cook et al (2010, 2015b) was used for reconstruction of the scPDSI at each grid point. The instrumental

scPDSI extend from 1901 to 2012, but the instrumental precipitation and temperature measurements used to compute the scPDSI are subject to large uncertainties in the early 20th century over Mexico (Jauregui 1979, Douglas 2007). Therefore, the reconstructions were calibrated using the best quality period of data from 1944 to 1984, and statistically validated by comparison with the lesser quality withheld instrumental scPDSI available from 1920 to 1943 prior to the calibration interval.

Three levels of EPPRensembles were used to compute the MXDA:

1. An 8-member ensemble average was computed where the candidate chronologies were selected from a 500-km search radius and were then weighted by the power of their correlation with summer scPDSI at the target grid point. See Cook et al (2015b) for the exact method of power weighting also used in the MXDA.

2. Same as ensemble 1, but using a 1000km search radius. The reason for employing this larger search radius was to bracket the likely correlation decay e-folding distance of 500-1000 km for scPDSI over the MXDA domain (cf. Cook et al 2015b). Secondly, this larger search radius includes additional tree-ring predictors of scPDSI for reconstruction.

3. The joint average of ensembles 1 and 2, a 16-member ensemble, yielded a MXDA with a common interval of reconstruction for all 1501 grid points of 1400-2012. This 16-member joint ensemble is used for the MXDA.

5. Results

5.1. Calibration

The results of the calibration and validation of the tree-ring reconstructions of scPDSI at each grid point over the MXDA domain are mapped in Figure 5. These statistics indicate that it is possible to reconstruct the summer scPDSI over most of Mexico, including areas with few or no local tree-ring chronologies (Figure 5a). The fraction of variance explained by regression in the calibration period (the coefficient of multiple determination or CRSQ, 1944-1984) is above 40% over most of Mexico, and is above 50% for the regions best sampled by the proxy tree-ring chronologies (Figures 2 and 5a). The cross validation reduction of error calculated for the calibration period (CVRE) uses a leave-one-out procedure to estimate the reduction of error statistic and provides a more rigorous estimate of the variance explained by the tree-ring data than the CRSQ (Cook et al 1999, 2010). The spatial pattern of explained variance represented by the CVRE conforms with the CRSQ and again indicates that over 40% of the variance is represented by the tree-ring data over most of Mexico (Figure 5b). Considering the large uncertainties in the instrumental scPDSI for portions of Mexico during the verification period (1920-1943, see below), the CVRE may provide the most reasonable estimate of out-of-sample validation skill currently possible for the MXDA.

5.2. Validation

The verification period statistics indicate that the validation of the tree-ring estimates against withheld instrumental scPDSI during the interval from 1920 to 1943 is generally restricted to those regions of Mexico with the most numerous tree-ring chronology predictors. There are exceptions, however, such as Chihuahua where there

are many tree-ring records but poor statistical validation during the period 1920-1943 (cf. Figures 2 and 5cde). The square of the Pearson correlation coefficient during the verification period (VRSQ) is highest over the state of Durango, over Central Mexico, and along the US/Mexico borderlands (Figure 5c) where some 30 to 40% of the variance is shared by the instrumental and reconstructed data and where several high quality tree-ring chronologies were available for reconstruction. The spatial pattern for the verification period reduction of error statistic (VRE)(Figure 5d) is very similar to the VRSQ (Figure 5c), but the verification period coefficient of efficiency (VCE)(Figure 5e) indicates some reconstruction skill for only about half of Mexico. The VCE is the most rigorous of these verification statistics because it is tested against the mean of the instrumental scPDSI during the validation period (Cook et al 1999) and the areas of Mexico without local tree-ring data generally do not pass this test. Some of this weakness may simply be attributed to the lack of local tree-ring data, but not all. High quality chronologies are available for Chihuahua, for example, but the VRE and especially the VCE statistics do not indicate reconstruction skill in those areas. Much of this apparent lack of skill is likely due to few and/or lower-quality instrumental climate observations over portions of Mexico during the 1920-1943 validation period [see Jauregui (1979) for information on the sparse and discontinuous precipitation recording stations in Mexico during the late 19 and early 20th centuries]. This interpretation is supported by the sharp increase in validation along the U.S. side of the northern border of Mexico (Figure 5de) because the communities and weather stations there did not suffer the turmoil witnessed in Mexico during the early 20th century.

The RBAR map (Figure 5f) shows the average correlation between the 16 individual ensemble reconstructions at each grid point over their full length (1400-1984). It is intended to show how consistent the reconstruction ensembles are as a function of correlation weighting and search radius. The highest RBAR values are located in the southwestern and southeastern United States and for central and southern Mexico.

6. Analysis and Discussion

6.1. The spatial pattern and intensity of significant droughts in Mexican history

Serious social impacts have resulted from severe droughts during the historical era in Mexico, and some reconstructed with proxy records may have contributed to the decline of certain pre-Hispanic city-states. The tree-ring data currently available for Mexico are unfortunately still too sparsely distributed prior to 1400 for a robust estimation of the spatial pattern and intensity of drought or wetness during the rise and fall of ancient Mesoamerican societies. However, drought seems to be a far too frequent occurrence in Mexico to always result in profound social change. Rural subsistence agriculture in Mexico has evolved in the presence of repeated dryness and many drought coping strategies have been developed. These strategies include the development of drought selected crop varieties, a flexible ratio of drought-sensitive to drought-tolerant crop varieties, social exchange networks for the redistribution of food in times of scarcity, the last resort use of famine foods such as agave, cactus, and mesquite fruits, and of course irrigation (Liverman 2001). But even these drought coping systems can be overwhelmed during times of exceptional drought, and they may be

further stressed if drought continues to deepen in response to anthropogenic climate change over Mexico.

The long historical record in pre-Hispanic, colonial, and modern Mexico is rich in detail on the socioeconomic and environmental impacts of severe drought, beginning with the “Drought of One Rabbit” in 1454 recorded in Aztec iconography and oral history. The Aztec counted time with inter-related religious and solar calendars that together recycled every 52 years (Caso 1971, QuiñonesKeber 1995). The sequential order of the 52-year calendar cycles is not obvious, but was organized around major political events by Aztec scribes. The sequencing of the 52-year cycles was then correlated with the Julian calendar by Jesuits and surviving Aztec scribes during the 16th century. These calendar correlations have been improved with ingenious historical, astronomical, and archaeological research, especially using solar eclipse dates recorded by both Aztec and Western systems (Aveni and Calnek 1999, Milbrath 2013). It is therefore possible to relate some events recorded in traditional Aztec history to the Western calendar and to exactly dated tree-ring reconstructed moisture anomalies during the late Aztec era (Therrell et al. 2004, Stahle and Dean 2011).

Perhaps the most infamous pre-Hispanic Aztec famine occurred in the year 1454, a year of “One Rabbit,” the first year of the 52-year Aztec calendar (QuiñonesKeber 1995, Chimalpahin 1998). The few surviving Aztec codices (pictorial histories of pre-Hispanic city-states), and oral histories communicated to the Spanish in the 16th century, mention this famine in 1454 and allude to crop failure due to drought and frost (QuiñonesKeber 1995, Therrell et al. 2004). The Nahuatl-speaking Aztec appear to have been superstitious about the year One Rabbit, and informants reported

to Spanish writers that One Rabbit years were often linked with some sort of calamity for the Aztec. Statistical analysis suggests that poor tree growth and inferred drought tended to occur in the year immediately preceding most One Rabbit years during and preceding the Aztec era, which might have reduced crop supplies and caused famine in the following year (Therrell et al. 2004). The best-documented pre-Hispanic example of this apparent “Curse of One Rabbit” occurred in the famine of 1454. The gridded PDSI reconstructions now available in the MXDA are based on more tree-ring chronologies and indicate that the most intense dryness lasted for three years from 1452-1454 and was concentrated over Central Mexico (Figure 6a).

Rain fed agriculture in the highlands of rural Mexico can be a precarious enterprise. Maize yields in particular can be limited by early growing season drought that reduces germination, height growth, and flowering prior to the *canicula*, and by freezing temperatures in autumn before the maize has fully matured (Eakin 2000, Therrell et al. 2006, Skoufias and Vinha 2012). The co-occurrence of early summer drought and fall frost can greatly reduce maize yields and lead to famine conditions that might not be fully alleviated until the next adequate harvest. The Famine of One Rabbit in 1454 may have been a case in point because in addition to the three-years of drought reconstructed in the MXDA, a fall frost may have also occurred in central Mexico in 1453, based on limited historical and proxy data (Therrell et al. 2004).

The conquest of the Aztec empire by a handful of Spanish conquistadores and their indigenous allies is one of the most incredible military campaigns in world history. The story has been recounted by Hernan Cortez himself (1986 translation), and perhaps most faithfully by Bernal Diaz (1963 translation), one of Cortez’s lieutenants in

the Conquest. The fall of the Aztec empire was a clear-cut military victory by the Spanish and the alliance of native groups shrewdly organized by Cortez. But the subsequent collapse of native population appears to have been the complicated result of imported and indigenous disease, drought, and harsh treatment following conquest (Acuña Soto et al. 2002). The estimates of the native population of Mexico in the early 16th century prior to Conquest are uncertain, but as many as 20 million native people may have perished in Mexico during the 16th century (Cook and Simpson 1948, Gerhard 1993). Most of this population collapse occurred in just three relatively brief episodes of epidemic disease in 1519-1520 (smallpox), the 1540s, and the 1570s (Acuña Soto et al. 2002). Extreme climatic conditions may have played a role in the ecology and infections associated with these diseases, especially the epidemics of hemorrhagic fever known as *cocoliztli* in the 1540s and 1570s (Acuña Soto et al. 2002 and 2005).

The Aztec apocalypse in 1521 came not only with military conquest and smallpox disease, it also appears to have been attended by severe sustained drought. Dry conditions over central Mexico may have begun as early as 1514 (Stahle et al. 2011a), but according to the MXDA were most extreme from 1521 to 1524 (Figure 6b). What role, if any, this drought may have had in the tumultuous aftermath of conquest and disease is not presently known. However, the spatial pattern of the most severe four-years of the Conquest-era drought is interesting because it was nearly nationwide (Figure 6b), a pattern predicted to become more common in the event of anthropogenic climate change.

The worst famine of the colonial era in Mexico occurred in 1786, and is referred to as *El Año de Hambre*, the year of hunger (Florescano and Swan 1995, Therrell 2005).

Two to three years of drought and an early fall frost in 1785 again appear to have led to crop failure and famine in 1786 (Therrell 2005, Therrell et al. 2006). An estimated 300,000 people died during *El Año de Hambre* due to both famine and an outbreak of epidemic typhus in 1785-1787 (Cooper 1965, Burns et al. 2014). The MXDA indicates that drought conditions were most serious during the two-year period from 1785 to 1786 when drought extended over most of Mexico, most severely over central and northeastern Mexico (Figure 6c).

The Mexican Revolution began in November 1910 and was stimulated in part by grievances related to agrarian reform, but drought conditions have long been suspected as a contributing factor (e.g., Tutino 1986). The MXDA indicates at least two years of drought over most of Mexico in 1909 and 1910 (Figure 6d). A recent analysis based on the available instrumental observations of drought indicate that the highest rates of insurgent activity occurred in those Mexican municipalities that suffered the most severe drought conditions just prior to the Revolution (Dell 2012). Similar analyses indicate that some of the highest rates of illegal immigration to the United States have tended to come from the most drought-afflicted areas of Mexico (Hunter et al. 2013).

In fact, Dell (2012) argues that relatively modest events, such as the droughts possibly involved in insurgency during the Revolution, can have persistent and highly non-linear socioeconomic effects. The efforts of the Mexican state to promote stability in insurgent regions using the agrarian reforms of the *Ejido* system are cited as an example. The unintended consequences of the *Ejido* system of small communally held farming systems may have included reduced borrowing against land collateral, thus limiting agricultural business development and *Ejido* incomes. Consequently, Dell (2012)

argues that the Mexican GDP might have been 124% higher in 1995 without the land reforms following the Revolution. Due in part to these persistent economic inefficiencies, many of the municipalities that were most heavily involved in insurgency during the Mexican Revolution remain poverty stricken today (Dell 2012). Drought conditions during the Revolution were certainly not the only cause of these hypothesized political and economic outcomes, but the argument that drought (or prolonged wetness) can lead to long term social and environmental change has been abundantly illustrated in the aftermath of other severe droughts and pluvials, including the Dust Bowl drought in the U.S. Great Plains during the 1930s (Lockeretz 1978, McLeman et al. 2014, Lee and Gill 2015).

The preceding examples might wrongly suggest that droughts and pluvials reconstructed in the MXDA are all well supported by available historical or proxy evidence on climate. Documentary information regarding the Conquest Drought of 1521-1524 and climate conditions during the Mexican War of Independence seem to contradict the MXDA. Historical descriptions of lake levels in the Basin of Mexico and Michoacan suggest wet conditions beginning in 1517 and reaching a maximum in 1522 (O'Hara and Metcalfe 1997). This documentary information strongly disagrees with the tree-ring reconstructions for the period 1521-1524 (Figure 6b) which are based on high quality Douglas-fir and Montezuma baldcypress chronologies from all four regions of Mexico, especially central and northeast Mexico (e.g., Figure 9, below). Assuming the historical information is correct, there must be an explanation for these dramatic differences in the hydroclimatology of central Mexico during the 1520s. One possibility

could have been dry conditions during the winter, spring and early summer, which were then reversed by heavy rains late in the summer season.

Drought from 1808-1811, identified with historical evidence, is believed to have caused “acute agricultural hardship” and contributed to the revolt against colonial rule that initiated the Mexican War of Independence in September of 1810 (O’Hara and Metcalfe 1997). However, the tree-ring reconstructions in the MXDA do not indicate uniform dryness during this time period. Drought is estimated in 1808 but normal to above average conditions over central Mexico are reconstructed for 1809, 1810, and 1811 (Figure 7). These apparent differences between documentary evidence and the exactly dated PDSI estimates in the MXDA need to be rectified and doing so could lead to a deeper understanding of the climate, social, and environmental history of Mexico.

6.2. Regional drought and wetness reconstructions for Mexico

There are important geographical gradients in the seasonal variability of climate in Mexico (Metcalfe et al. 2015), including an out-of-phase dipole-like pattern of warm season precipitation totals that sometimes develops between southern and northern Mexico (Higgins et al. 1999, Therrell et al. 2002). These spatial differences arise in part from large-scale ocean-atmospheric forcing associated with ENSO (Magaña et al. 2003, Ogelsby et al. 2012), and possibly from land surface interactions associated with the northward progression of the Mexican monsoon (Higgins et al. 1999). At the same time, the tree-ring chronologies used to estimate the scPDSI in the MXDA are not evenly distributed across the country but are most numerous in certain regions of Mexico (Figure 2). We have therefore computed four large regional averages of instrumental

and reconstructed JJA scPDSI from the MXDA to describe and analyze the history of moisture variability over Mexico. These four regions are northwest (NW; 22-31°N, 102.5-111°W), northeast (NE; 22-27°N, 97-102.5°W), central (C; 15-22°N, 95-103°W), and southeast Mexico (SE; 14-19°N, 90-94°W; Figure 3). These regions include the best coverage of tree-ring chronologies and highest quality estimates of PDSI in the MXDA, and they roughly conform with other regional studies of moisture variability in Mexico (e.g., Englehart and Douglas 2002, Giddings et al. 2005, Bravo et al. 2012).

The instrumental JJA scPDSI are plotted from 1920 to 2012 for the four regions in Figure 8 (the gridded scPDSI for Mexico begin in 1901 but the earliest years are based on very sparse observations). Inter-annual and decadal variability appears to be greatest in NW and NE Mexico, where the most severe and sustained droughts occurred in the 1950-1960s and 1990-2000s (Figure 8ab). Central and SE Mexico recorded near normal to above average soil moisture during these two 20th century northern Mexico drought regimes (Figure 8cd). Instrumental PDSI is positively correlated among the three central and northern regions, but not with southern Mexico (Table 1). The quality of the instrumental PDSI data declines prior to 1944 simply due to the fewer available weather stations (Englehart and Douglas 2002, Giddings et al. 2005), and this may be reflected by the generally lower correlations computed among the regional averages for 1920-1943 compared with the 1944-2012 time period (Table 1). There is some weak negative correlation between SE and NW or NE Mexico in the instrumental scPDSI (Table 1) possibly reflecting in part the dipole-like structure in moisture levels over Mexico in certain years (e.g., Figure 20, below).

The reconstructed JJA scPDSI are plotted from 1400 to 2012 for the same four regions of Mexico in Figure 9 (includes the instrumental data from 1985 to 2012). The regional reconstructions also exhibit higher inter-annual variability in the more arid NE and NW regions of Mexico, and only a few decadal droughts or pluvials are estimated to have occurred over all four regions with equal intensity (Figure 9). The sample size of predictor chronologies remains strong throughout this 600-year period in NW Mexico and to a lesser extent in NE Mexico, but in central Mexico the sharp decline of predictor tree-ring chronologies before ca 1600 (Figure 2) is evident in the temporal stability of the derived reconstructions (Figure 9). There is considerable overlap in the tree-ring predictors between the four regional reconstructions due to the variable search radii employed with EPPR (500- to 1000-km), and this may be reflected by the higher correlations among the three central and northern regions than computed for the instrumental data, especially when comparing the best replicated subperiods (i.e., 1944-2012 and 1700-1984 for the instrumental and reconstructed data, respectively, Table 1). There is also some weak negative correlation between SE and NW Mexico over the past 600 years that is marginally significant given the large sample size (Table 1). Most of this anti-phasing appears to arise from the occasional development of a dipole in reconstructed moisture extremes over SE and NW Mexico (see below) but it is not a strong feature of the inter-annual variability present in the sub-regions of the MXDA.

The decadal versions of the regional reconstructions reveal a few persistent droughts that extended over most of Mexico (Figure 9). The most intense and prolonged of these droughts are mapped in Figure 10 for the period after 1550 when the reconstructions are based on tree-ring chronologies that cover most of the country.

These five selected droughts lasted from four to eight years and were most severe over northern Mexico (Figure 10). The drought from 1566-1573 (Figure 10a) was part of the 16th century megadrought that lasted with little interruption for at least 20-years over northern Mexico (Stahle et al. 2000, 2007). The drought from 1666-1669 was one of the two most extreme sub-decadal to decadal droughts of the past 600-years over NE Mexico, and afflicted most of the country during this four-year drought regime (Figure 10b). This 1660s drought had well documented impacts among the Spanish and Native Americans of the southwestern United States (Stahle and Dean 2011), and was the most severe sustained drought indicated in the 470-year long tree-ring reconstruction of the standardized precipitation index during the summer monsoon season over Arizona, New Mexico, and northwest Mexico by Griffin et al. (2013). Severe sub-decadal droughts of five to six years duration are estimated to have occurred in the mid-18th and early 19th centuries (Figure 10cd).

The best-documented regime of decadal drought in Mexican history occurred in the mid-20th century. The 1950s drought is evident in both the reconstructed and instrumental PDSI (Figure 10ef) and it may have been the most extreme decadal drought in the 600-year reconstructions for NE and NW Mexico (Figure 9). The 1950s drought was most intense from 1951-1957 in both the instrumental and reconstructed data, but it began in the late 1940s and persisted into the 1960s over northern Mexico (Figure 9). In contrast, only select years were dry over central Mexico during the 1950s, and SE Mexico experienced above average moisture conditions for most this period (Figures 8, 9, and 10ef). The anti-phasing between southern and northern Mexico is driven in part by differences in ENSO forcing (Therrell et al. 2002, Magaña et al. 2003,

Seager et al. 2009) and the La Niña conditions that recurred during portions of the 1950s favored wetness in the south but extreme drought in the north.

The reconstructions for central and southeast Mexico are reasonably well replicated after 1550 and we use the MXDA to identify tree-ring reconstructed droughts that were concentrated in those regions. The most intense decadal droughts over central Mexico in the well-replicated period occurred from 1625-1634, 1696-1705, and 1894-1899 (Figures 9 and 11). The 1890s drought actually affected most of Mexico except for the Southeast and began as early as 1890 over central Mexico and lasted over some sectors until 1902 (not shown). The onset of the 1890s drought coincided with a six-year period of persistent La Niña conditions [based on the extended Multivariate ENSO Index (eMEI) of Wolter and Timlin 2011; Figure 20, see below]. However, the most intense six years of drought in the 1890s were concentrated over central Mexico (Figure 11c), and the severe socioeconomic impacts were reported in the international press (Seager et al. 2009).

The five most extreme sub-decadal to decadal droughts that were concentrated in Southeast Mexico are mapped in Figure 12. Two of these droughts occurred during the 20th century and the instrumental PDSI data are mapped along with the reconstructions for these particular events (Figure 12d-g). In some cases, when drought occurs over the southeast there is a tendency for wetter conditions to prevail over central and northern Mexico (Figure 12, Table 1). The occasional anti-phasing of tree-ring reconstructed moisture extremes between southern and northern Mexico was reported by Therrell et al. (2002), and has been detected in the new MXDA. The statistical significance of regional reconstructions recording extreme events of opposite

sign during the same year was tested using Monte Carlo simulation. For each regional reconstruction, the number of values below -1.5 and above +1.5 for the period 1700-2012 was counted and synthetic time series of corresponding length and event frequency were produced by random distribution. The synthetic time series were compared and the number of matches between all pairs of series was recorded. This procedure was repeated 10,000 times to calculate one-tailed critical values for the 95% and 99% probability levels. If the number of matches between opposite extremes in two reconstructed regional time series exceeds these levels, then the count of coinciding extreme values is higher than expected by chance.

The count of concurrent extremes of the opposite sign is significant when comparing reconstructed scPDSI in SE and NW Mexico ($p < 0.01$; note that extremes of the same sign are not significant). Most of the extremes in SE Mexico are actually positive (> 1.5) during the period from 1500-2012 (62 of 97 years). In fact, reconstructed PDSI over SE Mexico was above average (moist) for most of the 16th, 17th, and early 18th centuries (Figure 9d). ENSO and Atlantic forcing (Magaña et al. 2003, Seager et al. 2009), and land surface feedbacks involved in the onset of the Mexican monsoon (Higgins et al. 1999) may be involved in the dipole of moisture extremes between southern and northern Mexico, and the MXDA reconstructions suggest that this anti-phasing of regional moisture extremes has been a minor but statistically significant component of climate variability over Mexico for the past 500+ years.

The MXDA provides evidence for several “All Mexico” pluvials, some more intense and sustained than any witnessed during the modern instrumental era (Figure 13). One of the most extreme All Mexico wet episodes occurred from 1814-1817 (Figure

13e), a period that included global climate forcing from the April 1815 cataclysmic volcanic eruption of Tambora, Indonesia, the largest eruption in recorded history (Stothers 1984). However, the exceptional wet anomalies over Mexico in 1814 and 1815 largely predated the climatic forcing from Tambora and contributed to this exceptional four-year period of wetness (note the simultaneous dryness reconstructed for SE Mexico and the southeastern United States, Figure 13f). The most extreme All Mexico pluvial estimated with the MXDA occurred from 1483-1487 in the late-Aztec era (Figures 9 and 13a), but fewer tree-ring chronology predictors are available from the main areas of wetness in the late 15th century so the magnitude, duration, and spatial coverage of this exceptional pluvial will need confirmation with additional long tree-ring data from Mexico.

6.3. The ENSO, AMO, and PDO teleconnection to instrumental and reconstructed soil moisture over Mexico, 1871-2005

One of the strongest ENSO teleconnections worldwide has been detected in instrumental climate data and tree-ring chronologies from the southwestern United States and Mexico (Ropelewski and Halpert 1987, Cavazos and Hastenrath 1990, Allan et al. 1996, Cole and Cook 1998, Stahle et al. 1998, Magaña et al. 2003, Villanueva et al. 2007, Stahle et al. 2011b, Pompa-Garcia and Nemiga 2015). However, even in portions of this so-called TexMex sector there can be substantial inter-decadal variability in the strength of ENSO forcing (Cole and Cook 1998, Diaz et al. 2001). Because the quality and quantity of meteorological observations in Mexico decline in the early 20th century, and because the instrumental record of ENSO now extends back into

the 19th century, the new scPDSI reconstructions in the MXDA can contribute to the analysis and understanding of ENSO forcing to Mexican climate.

The correlation between the extended Multivariate ENSO Index (eMEI; available from 1871-2005) averaged for the February-May (FMAM) season and the scPDSI at 1501 grid points across Mexico is illustrated in Figure 14 using both tree-ring reconstructed and instrumental data. The correlations are restricted to the period from 1901-1984 in common to all series, but the results are similar for the full period of overlap between the eMEI and instrumental or reconstructed scPDSI (i.e., 1901-2005 and 1871-1984, respectively). The correlations between the eMEI and reconstructed scPDSI exceed +0.6 over portions of northern Mexico (Figure 14a), but only reach +0.5 over the same region for the instrumental data (Figure 14b). Slightly negative correlations with the eMEI are observed over southern Mexico for both the reconstructed and instrumental scPDSI. The negative correlations with the eMEI over southern Mexico might be higher using only summer precipitation observations (Magaña et al. 2003) instead of the scPDSI which models the moisture balance over several seasons or approximately 12 months.

Other large-scale ocean-atmospheric teleconnections to Mexican climate have been detected with analyses of instrumental data (e.g., Pavia et al. 2006, Seager et al. 2009, Mendez and Magaña 2010, Stahle et al. 2011b, Feng et al. 2011), and the AMO and PDO are also both correlated with instrumental and reconstructed JJA scPDSI in the MXDA (illustrated for the period 1901-1984 in Figure 15). However, these correlations are lower and less widespread than computed with the ENSO index (eMEI; Figure 14 and summarized in Table 2 for the four regions of Mexico), similar to the

results computed for ring width chronologies from Mexico by St. George (2014). The correlation between the AMO index and the reconstructed PDSI in Mexico for the period 1901-1984 is representative of the correlation for the full common period 1874-1984 (or even 1874-2012; not shown).

The ENSO teleconnection to reconstructed scPDSI over northern Mexico is not only stronger than either the AMO or PDO, it has also been reasonably stable since the 19th century. The correlation between the FMAM eMEI and the gridded reconstructions is computed for non-overlapping 30-year subperiods in Figure 16 (only 24-years for the final period 1961-1984). The strength of the correlations vary and even change sign over southern Mexico, but remain strong and positive over the north, even more so than over the southern and southwestern United States (Figure 16a-d). This ENSO signal in tree-ring reconstructed climate for northern Mexico is one of the strongest and most temporally stable in the world (Stahle et al. 1998, Christie et al 2009, Sano et al. 2012).

The Mexican soil moisture response to warm and cold ENSO events is also investigated using composite analysis in Figure 17. The average moisture anomalies in the reconstructed and instrumental PDSI data for Mexico during the most extreme El Niño and La Niña events recorded in the eMEI are mapped in Figure 17abcd (15 extremes each from 1871-1984 for the reconstructed PDSI and 10 each from 1901-1984 for instrumental PDSI). The symmetry of the reconstructed soil moisture response to warm and cold ENSO forcing is striking over northern Mexico (Figure 17ab), and is much stronger than the shift from wet to dry conditions evident in the instrumental observations (Figure 17cd). The drought conditions reconstructed over northern Mexico during La Niña conditions (Figure 17b) compare favorably with the instrumental moisture

anomalies for cold events (Figure 17d), but the wet conditions during El Niño events are more variable in the instrumental average than in the reconstructed data for warm regimes (Figure 17). Some of the differences may be due to the fact that the El Niño years being composited are not identical between the instrumental and reconstructed data, but the quality and quantity of instrumental climate data from Mexico during the early 20th century may also be involved.

The reconstructed symmetry of the wet response to warm events and the dry response to cold events (Figure 17ab) is reflected reasonably well in the maps of reconstructed scPDSI during each ENSO extreme from 1871-1984 (Figures 18 and 19). Reconstructed scPDSI over northern Mexico was moist during 11 of 15 El Niño extremes (Figure 18) and was dry during 13 of 15 La Niña extremes (Figure 19; based on the FMAM eMEI), although the magnitude of the moisture anomalies varied among events. The ENSO teleconnection to central and southern Mexico is much less consistent in the MXDA. The occurrence of drought over southern Mexico during El Niño conditions is strongly reconstructed for only four events in the MXDA (1878, 1897, 1973, 1983; Figure 18). Wetness is reconstructed over southern Mexico only during three La Niña events (1894, 1956, 1974; Figure 19). Again, the MXDA does not robustly estimate late summer precipitation, but these results suggest that the ENSO teleconnection to warm season moisture over central and southern Mexico witnessed in the relatively short instrumental records may not be entirely stable (e.g., Magaña et al. 2003). In both northern and southern Mexico the ENSO teleconnection may vary in response to the location of the SST anomaly in the eastern or central Pacific (e.g., Infanti and Kirtman 2016).

The stability of the large-scale ocean-atmospheric signal in the MXDA can be further evaluated by correlating the regional average scPDSI from northwest, central, and southern Mexico with Atlantic and Pacific SSTs over the past 142-years (1871-2012). Reconstructed JJA scPDSI in northwest Mexico is positively correlated with JFM-average SSTs in the tropical Pacific for their full common period 1871-1984 (Figure 20a; these correlations exceed +0.6 in the central equatorial Pacific). This positive correlation with tropical Pacific SSTs is also detected in the reconstructions for northeast Mexico and is generally stable during various subperiods of the late 19th and 20th century (not shown). Instrumental JJA scPDSI for northwest Mexico is correlated with SSTs over the same region of the tropical Pacific after 1901, but the positive correlations are not as strong or as centered along the equator as noted in the reconstructed data (Figure 20b).

Reconstructed JJA scPDSI in central Mexico are not correlated with winter SSTs, but they are negatively correlated with SSTs in the eastern tropical Pacific from 1951-1984 when the SST data are averaged for the summer season (MJJA; Figure 20c). The instrumental JJA Palmer drought indices are also negatively correlated with tropical Pacific SSTs in summer for the period 1951-1984 (and 1951-2012), but not as strongly as computed for the reconstructed data (Figure 20d). This negative SST relationship with summer moisture conditions in central Mexico has been previously reported (e.g., Seager et al 2009, Englehart and Douglas 2010, Stahle et al 2011), but analyses with the MXDA indicate that it has not been stable during the 20th century. The correlation between instrumental and reconstructed PDSI in central Mexico with tropical Pacific SSTs actually becomes weakly positive during the period 1901-1950 (not shown).

In southern Mexico, JFM-average SSTs in the Caribbean and tropical Atlantic are most highly and consistently correlated with reconstructed and instrumental PDSI (Figure 20e,f). These correlations are positive as has been reported for summer precipitation over southern Mexico and Central America (e.g. Wang et al. 2006), but they are observed only after 1950. Prior to 1951, the reconstructed or instrumental PDSI for southern Mexico are not correlated at all with SSTs in the Atlantic or Caribbean, and all correlations in the tropical Pacific are weak and inconsistent (not shown). Some of the instability in these SST correlations with regional PDSI in central and southern Mexico from the early to late 20th century may be due in part to data quality issues, but the strong and reasonably stable SST signal seen in the tree-ring reconstructions for northwest Mexico suggests that real non-stationarity may exist in the SST signals detected over central and southern Mexico.

Aside from this apparent temporal instability, the SST correlations with regional PDSI indicate that the ocean influences on summer PDSI change sharply from north to south across Mexico (Figure 20). These changes parallel the shift from the mid-latitude westerlies that prevail over northwest Mexico to the easterly trade winds over southern Mexico. The reconstructions also register the change in the seasonality and sign of ENSO-related SST forcing from northwest to central Mexico as reported for instrumental precipitation data (e.g., Magaña et al 2003, Englehart and Douglas 2010). However, these north-south changes in the SST signals detected with the MXDA are not as strong or as temporally stable as the ENSO and/or tropical Pacific SST signal embedded in the JJA scPDSI reconstructions for northwest (and northeast) Mexico, which appears to be

one of the strongest ENSO signals yet detected with proxy tree-ring data (e.g., Christie et al. 2009, Sano et al. 2012).

The tree-ring reconstructed JJA scPDSI over northwestern Mexico is highly correlated with cool season indices of ENSO, especially the eMEI averaged for the FMAM season (e.g., $r = 0.55$ for 1871-1984). However, the strongest reconstructed PDSI response to ENSO forcing we have detected in the MXDA is actually in an index of the north-south dipole-like structure of moisture regimes over Mexico, in spite of the apparent instability in the moisture response over southern Mexico to ENSO. To index this dipole-like structure over Mexico, the average moisture anomaly during the worst seven-years of the 1950s drought (1951-1957), which was characterized by severe drought in northern Mexico simultaneously with wetness over southern Mexico (Figure 10ef), was used in a map pattern correlation analysis (Cook et al. 1999). Reconstructed PDSI values at each grid point were correlated with the average for 1951-1957 at the same points for each year from 1400-2012. The MXDA-wide annual average of the map pattern correlation coefficients provides an estimate of the strength and direction of the dipole pattern for the past 600-years (Figure 21a; the correlations were inverted so that dry conditions during the 1950s are negative). Note the very strong dipole estimates during the severe sustained droughts of the late 16th century, the 1860s, and of course 1950s (Figure 21a). The spatial pattern of the 15 most positive and 15 most negative correlation coefficients is illustrated in Figure 21bc. This Mexican dipole index is correlated with the Nov-Feb eMEI at $r = 0.58$ ($p < 0.001$) for the common period 1871-2005, and ranged from 0.40 to 0.75 for discrete 30-year subperiods (Figure 21d; correlations using the Feb-May eMEI are only slightly lower).

909 These tree-ring analyses indicate that the main JJA PDSI response to ENSO
910 forcing over subtropical North America occurs in northern Mexico (Figures 14 and 16-
911 20). However, some analyses based on instrumental observations, including those of
912 JJA PDSI in Figures 14b and 17c,d, indicate that ENSO forcing over this sector is
913 stronger to the north or east. Part of the relatively weak instrumental climate response to
914 ENSO in northern Mexico may be due to the short and discontinuous record of
915 meteorological observations for that region. The tree-ring reconstructions are certainly
916 suggesting that the strength and temporal stability of the ENSO signal in North America
917 is strongest in northern Mexico.

918 To test the strength and stability of the ENSO signal over northern Mexico we
919 computed a central equatorial Pacific sea surface temperature (SST) average for the
920 Niño 4 region (i.e., 5N-5S, 160E-150W) using the millennium simulations developed
921 with the Community Earth System Model (CESM 1.2.z; Hurrell et al. 2013) for the Jan-
922 May season. This equatorial Pacific SST index was correlated with Jan-May
923 precipitation totals and average temperature in the 1156-year simulations over the North
924 American sector (850-2005; Figure 22). The area of positive precipitation response
925 includes northern Mexico, but is strongest in the model over southern California,
926 Arizona, and the Gulf of Mexico. The Jan-May temperature response, however, is
927 strongest over northern Mexico in approximately the same region of maximum ENSO
928 correlation computed with the tree-ring reconstructions (Figures 14 and 22). These
929 results suggest that the integration of both precipitation and temperature in the soil
930 moisture balance over northern Mexico during the winter-spring-early summer season
931 may indeed be particularly sensitive to ENSO forcing. The Niño 4 SST correlations with

precipitation and temperature in the CESM simulations also reproduce the change in sign of the coefficients from northern to southern Mexico (Figure 22).

6.4. Anthropogenic forcing of All Mexico drought?

Climate model simulations of precipitation over Mexico under business-as-usual anthropogenic radiative forcing suggest that a deepening pattern of nationwide dryness will emerge over Mexico and extend into Central America in the coming century (Seager et al. 2009, Feng and Fu 2013). This anthropogenic drying trend over Mexico and Central America is expected to arise from a northward shift in the jet stream and mean storm track in winter, and from a weakening and southward displacement of the convection zone over southern Mexico and Central America in summer (Seager et al. 2009). However, the pattern of drought and wetness over Mexico evident in the instrumental record for the past 30-years is not consistent with this projected “All Mexico” drying trend (Seager et al. 2009).

The tree-ring reconstructions in the MXDA can be used to estimate the frequency of nationwide drought to place the anticipated anthropogenic changes over Mexico in the context of the past 600 years. The gridded reconstructions have been used to compute the Drought Area Index (DAI) for mainland Mexico between 14-30°N and 90-110°W, exclusive of the Baja and Yucatan Peninsulas. The DAI is calculated as the percentage of grid points where the reconstructed scPDSI is ≤ -1.0 , and provides an estimate of the area of drought over mainland Mexico from 1400-2012. This MexicanDAI is plotted in Figure 23a and composite maps of the 15 lowest and highest DAI values depict the nationwide patterns of wetness and dryness measured by

extremes of this index (Figure 23bc). The reconstructed DAI indicates that 25.3% of mainland Mexico experiences drought indices of ≤ -1.0 in any given year (median DAI = 14.6%). The most widespread single year of dryness is reconstructed for 1819 when 93.6% of mainland Mexico is estimated to have suffered drought (Figure 23a). Regimes of widespread dryness over Mexico are reconstructed during the 1520s, 1570s, 1660s, 1860s, 1950s, and early 2000s, but the MexicanDAI indicates that intense All Mexico drought has been rare over the past 600 years.

The area of mainland Mexico under drought has increased since 1980 (Figure 23a). However, this drying trend has not been unprecedented in the reconstructed Mexican DAI since 1400. Instead, the tree-ring reconstructions indicate that truly nationwide dryness has not been common and nor persistent beyond the decadal timescale, which provides one measure for how unusual the model projected moisture levels may be in the late 21st century with business-as-usual anthropogenic climate forcing.

Another remarkable feature of the Mexican DAI is the 30-year absence of extensive dryness over mainland Mexico from 1824-1854(Figure 23a). A 20-year episode is also reconstructed to have been largely free of widespread droughtfrom 1930-1949. The tree-ring reconstructions are well replicated during both of these intervals and provide an interesting proxy perspective on the range of decadal moisture variability over Mexico that is not apparent in the short and spatially discontinuous instrumental observations.

7. Conclusions

978

979 The Mexican Drought Atlas provides an interesting new perspective on the
980 spatial variability of soil moisture over Mexico from 1400 to 2012. The best replicated
981 and most reliable subperiod extends from 1550-2012, and the first 150-years of the
982 Atlas need to be improved with the development of additional long tree-ring
983 chronologies, especially over central and southern Mexico. The reconstructed patterns
984 of drought and wetness are a rich source of independent climate information for
985 historical and environmental research, as illustrated for 1452-1454 and 1785-1786 when
986 drought contributed to severe famine in the Aztec and Colonial eras. The MXDA
987 also provides useful information on the dynamics responsible for climate variability over
988 Mexico, perhaps none more important than the potential predictability associated with
989 the El Niño/Southern Oscillation.

990 Analyses of the MXDA and of climate model simulations suggest that the
991 influence of ENSO on the moisture balance over North America has been strongest
992 in northern Mexico since 1871. The ENSO influence on reconstructed PDSI over
993 northern Mexico has been reasonably stable during the past 150 years, and it appears to
994 occur largely before the mid-summer drought. This seasonality is important for
995 socioeconomic reasons because moisture levels before and during the early wet
996 season can have a major effect on crop yields (Skoufias and Vinha 2012) and total
997 reservoir inflow (Nicholas and Battisti 2008) in Mexico. The strong correlation between
998 ENSO and instrumental precipitation (Caso et al. 2007) and reconstructed scPDSI in
999 the MXDA over northern Mexico during the winter and early growing season has
1000 allowed development of empirical and probability-based forecast models useful for

agriculture and water resource planning in Mexico (e.g., Magaña et al. 2003, Nicholas and Battisti 2008, Vazquez 2009).

The frequency and severity of decadal drought both appear to have been higher over northern Mexico than in more humid central and southern Mexico, especially during the last 450-years covered by the MXDA with the best-replicated tree-ring data. But extended droughts are reconstructed for central and southern Mexico, including the decade-long events from 1696-1705 over central Mexico and from 1811-1820 over southeastern Mexico. Droughts of this magnitude and duration would place heavy pressure on water and hydroelectric power supplies in urban Mexico and on subsistence agriculture in the southern highlands where the most extreme poverty in Mexico is concentrated (Landa et al. 2008). The CMIP5 climate model predictions indicate that these naturally occurring sub-decadal to decadal droughts will be made more frequent and widespread over all of Mexico by anthropogenic climate change (Seager et al. 2009). The MXDA reconstructions indicate that widespread “All Mexico” droughts have been uncommon over the past 600 years and their frequency does not appear to have increased significantly in recent decades. But if these nationwide anthropogenic droughts do become more frequent, they will certainly represent a major socioeconomic and humanitarian challenge for Mexico.

Acknowledgements

NSF grant numbers GEO/ATM-0753399, AGS-1266014, AGS-1266015, and AGS-1301587. Support was also provided by INIFAP, IAI, and WWF. Lamont-Doherty Earth Observatory contribution number xxxx. We thank Mark Brenner, Rosalinda Cervantes Martinez, Malcolm Cleaveland, Art Douglas, Vicenta Constante Garcia, Eladio Cornejo Olviedo, Juan Estrada Avalos, Song Feng, Ian Howard, Sarah Metcalfe, Susan Milbrath, Gene Paull, Daniel Stahle, Lorenzo Vazquez Salem, and Matthew for advice and assistance.

Appendix

The Mexican Drought Atlas can be accessed and analyzed at:

<http://www.djburnette.com/research/MXDA/Default.aspx>

where the maps of individual years or any combination of years can be composed and downloaded for the instrumental or reconstructed JJA scPDSI. The time series data for any single grid point or any contiguous regional selection of grid points can also be plotted and downloaded. Other analytical capabilities at the MXDA website include correlation analysis using uploaded time series, and map pattern correlation or congruence analyses using specified map targets.

Table 1. Correlation coefficients computed between the four sub-regions of Mexico using both instrumental and reconstructed PDSI. Three time periods are compared for both the instrumental (Inst) and reconstructed (Recon) data. The sub-regions are northwest (NW), northeast (NE), central (C), and southeast Mexico (SE).

Inst PDSI 1901-2012					Inst PDSI 1901-1943					Inst PDSI 1944-2012				
	NW	NE	C	SE		NW	NE	C	SE		NW	NE	C	SE
NW					NW					NW				
NE	.64				NE	.54				NE	.75			
C	.26	.55			C	.18	.63			C	.37	.47		
SE	-.11	-.08	.20		SE	-.16	-.05	.03		SE	-.08	-.12	.33	

Recon PDSI 1400-1984					Recon PDSI 1400-1699					Recon PDSI 1700-1984				
	NW	NE	C	SE		NW	NE	C	SE		NW	NE	C	SE
NW					NW					NW				
NE	.87				NE	.85				NE	.90			
C	.33	.51			C	.13	.34			C	.57	.70		
SE	-.12	-.06	.22		SE	-.16	-.06	.40		SE	-.08	-.07	-.03	

Table 2. Correlation coefficients computed between instrumental and reconstructed PDSI for the four sub-regions of Mexico and indices of the ENSO (eMEI), AMO, and PDO (*<0.05; **<0.01; ***<0.0001). Note the different seasons and time periods of comparison (dictated by the length of the available data). The sub-regions are northwest (NW), northeast (NE), central (C), and southeast Mexico (SE). The correlations with the AMO increase slightly if a FMAM season is used (e.g., $r = 0.30$ for reconstructed PDSI in NW Mexico).

	FMAM eMEI		Jun-Jul AMO		DJF PDO
Instrumental PDSI	1901-1984		1901-1984		1901-1984
NW Mexico	0.27 **		-0.20		0.26 **
NE Mexico	0.24 *		-0.20		0.16
C Mexico	-0.01		-0.06		0.15
S Mexico	0.11		-0.04		0.05
Reconstructed PDSI	1871-1984		1874-1984		1901-1984
NW Mexico	0.55 ***		-0.26 **		0.30 **
NE Mexico	0.47 ***		-0.24 *		0.28 **
C Mexico	0.13		-0.19 *		0.13
S Mexico	-0.05		-0.06		0.11

References Cited

- Adams, D.K., and A.C. Comrie, 1997. The North American monsoon. *Bulletin of the American Meteorological Society* 78:2197-2213.
- Acuña-Soto, R., D.W. Stahle, M.K. Cleaveland, and M.D. Therrell, 2002. Megadrought and megadeath in 16th century Mexico. *Emerging Infectious Diseases* 8:360-362.
- Acuña-Soto, R., D.W. Stahle, R.D. Griffin, and M.D. Therrell, 2005. When half the population died: the epidemic of hemorrhagic fevers of 1576 in Mexico. *FEMS Microbiology Letters* 240:1-5.
- Allen J, A. Assaf, J. Small, H. Riebeek, 2012. Drought causes Mexico food shortages. NASA earth observatory images, January 21, 2012 [cited 2012 Feb 1].
<http://earthobservatory.nasa.gov/IOTD/view.php?id=76983>
- Aveni, A.F., and E.E. Calnek, 1999. Astronomical considerations in the Aztec expression of history. *Ancient Mesoamerica* 10:87-98.
- Balling, R.C. Jr. 1988. The Climatic Impact of a Sonoran Vegetation Discontinuity. *Climatic Change*. 13:99-109.
- Bravo, J.L., E. Azpra, V. Zarraluqui, C. Gay, and F. Estrada, 2012. Cluster analysis for validated climatology stations using precipitation in Mexico. *Atmosfera* 25:339-354.
- Brenner, M., D.A. Hodell, J.H. Curtis, M.F. Rosenmeier, M.W. Binford, and M.B. Abbott, 2001.

- Abrupt climate change and Pre-Columbian cultural collapse. In *Interhemispheric Climate Linkages*, edited by V. Markgraf. Academic Press, New York. Pp. 87-103.
- Burns, J.N., R. Acuña-Soto, and D.W. Stahle, 2014. Drought and epidemic typhus, central Mexico, 1655-1918. *Emerging Infectious Diseases* 20:442-447.
- Carter, N.T., C.R. Seelko, and D.T. Shedd, 2015. *US-Mexico Water Sharing: background and recent developments*. Congressional Research Service 7-5700, p. 1-18.
- Caso, A., 1971. Calendrical systems of Central Mexico. *Handbook of Middle American Indians*, Vol. 10, R. Wauchope, Ed., University of Texas Press, 333–348.
- Caso, M., C. Gonzalez-Abraham, and E. Ezcurra, 2007. Divergent ecological effects of oceanographic anomalies on terrestrial ecosystems of the Mexican Pacific coast. *Proceedings of the National Academy of Sciences* 104:10530-10535.
- Cavazos, T. and S. Hastenrath, 1990. Convection and rainfall over Mexico and their modulation by the Southern Oscillation. *International Journal of Climatology* 10:377-386.
- Cerano-Paredes, J., J. Méndez-González, A. Amaro-Sánchez, J. Villanueva-Díaz, R. Cervantes-Martínez, E.A. Rubio-Camacho, 2013. Reconstrucción de precipitación invierno-primavera con anillos anuales de *Pinus douglasiana* en la reserva de la biósfera Sierra de Manantlán, Jalisco. *Revista Chapingo, Series ciencias forestales y del ambiente* 19:413-423.

1112 Chimalpahin, D.F., 1998. *Las Ocho Relaciones y el Memorial de Colhuacan (The Eight*
1113 *Relations and the Memorial of Colhuacan)*. Vol. 1. Consejo Nacional para la Cultura
1114 y las Artes, 433 pp.

1115 Christie, D.A., A. Lara, J. Barichivich, R. Villalba, M.S. Morales, and E. Cuq, 2009. El Niño-
1116 Southern Oscillation signal in the world's highest-elevation tree-ring chronologies from
1117 the Altiplano, Central Andes. *Palaeogeography, Palaeoclimatology,*
1118 *Palaeoecology* 281:309-319.

1119 Cleaveland, M.K., D.W. Stahle, M.D. Therrell, J. Villanueva-Díaz, and B.T. Burns, 2003.
1120 Tree-ring reconstructed winter precipitation in Durango, Mexico. *Climatic Change*
1121 59:369-388.

1122

1123 Coe, M.D., 1984. *Mexico*. Thames and Hudson Ltd., London. Third edition, 180pp.

1124

1125 Cole, J.E. and E.R. Cook, 1998. The changing relationship between ENSO variability and moisture
1126 balance in the continental United States. *Geophysical Research Letters* 25:4529-4532.

1127

1128 Cook, B.I., K.J. Anchukaitis, J.O. Kaplan, M.J. Puma, M. Kelley, and D. Gueyffier, 2012. Pre-
1129 Columbian deforestation as an amplifier of drought in Mesoamerica. *Geophysical Research*
1130 *Letters* 39, L16706, 1-5.

1131

1132 Cook, B.I., T.R. Ault, and J.E. Smerdon, 2015a. Unprecedented 21st century drought risk in the
1133 American Southwest and Central Plains drought risk in western North America. *Science*
1134 *Advances* 1, no. 1, e1400082, doi:10.1126/sciadv.1400082.

1135

1136 Cook, E.R., and K. Peters, 1981. The smoothing spline: a new approach to standardizing forest
1137 interior ring-width series for dendroclimatic studies. *Tree-Ring Bulletin* 41:45-53.

1138

1139 Cook, E.R., and K. Peters, 1997. Calculating unbiased tree-ring indices for the study of climatic and
 1140 environmental change. *The Holocene* 7:359-368.

1141

1142 Cook, E. R., and P. J. Krusic, 2005. Program ARSTAN: A Tree-Ring Standardization Program
 1143 Based on Detrending and Autoregressive Time Series Modeling, with Interactive
 1144 Graphics. Report on file, Tree-Ring Lab, Lamont Doherty Earth Observatory of Columbia
 1145 University, Palisades, NY.

1146

1147 Cook, E.R., D.M. Meko, D.W. Stahle, and M.K. Cleaveland, 1999. Drought reconstructions for
 1148 the continental United States. *Journal of Climate* 12:1145-1162.

1149

1150 Cook, E.R., C. Woodhouse, C.M. Eakin, D.M. Meko, and D.W. Stahle, 2004. Long-term aridity
 1151 changes in the western United States. *Science* 306:1015-1018.

1152

1153 Cook, E.R., R. Seager, M.A. Cane, and D.W. Stahle, 2007. North American Drought:
 1154 reconstructions, causes, and consequences. *Earth Science Review* 81:93-134.

1155

1156 Cook, E. R., R. Seager, R. R. Heim Jr, R. S. Vose, C. Herweijer, and C. Woodhouse, 2010.
 1157 Megadroughts in North America: placing IPCC projections of hydroclimatic change in a
 1158 long-term palaeoclimate context. *Journal of Quaternary Science* 25:48–61.

1159

1160 Cook ER, Anchukaitis KJ, Buckley BM, D'Arrigo R, Jacoby GC, Wright WE, 2010. Asian
 1161 monsoon failure and megadrought during the last millennium. *Science* 328:486-489

1162

1163 Cook, E. R., P. J. Krusic, and T. Melvin, 2014. Program RCSigFree. Tree-Ring Lab, Lamont

1164 Doherty Earth Observatory of Columbia University, Palisades, NY.

1165

1166 Cook, E.R., R. Seager, J. Kushnir, et al., 2015b. Old world megadroughts and pluvials during

1167 the Common Era. *Science Advances*, vol. 1, no. 10, e1500561, DOI:

1168 10.1126/sciadv.1500561.

1169

1170 Cook, S.F., and L.B. Simpson LB, 1948. *The Population of Central Mexico in the Sixteenth*

1171 *Century*. Ibero Americana Vol. 31, Berkeley: University of California Press; 1948.

1172

1173 Cooper, D.B., 1965. *Epidemic disease in Mexico City 1761-1813*. University of Texas Press,

1174 Austin.

1175

1176 Cortés, Hernán, 1986. *Letters* – available as *Letters from Mexico* translated by Anthony

1177 Pagden. Yale University Press, New Haven.

1178

1179 Cortez Vazquez, M., 2006. Mexico. In: State of the Climate in 2005, K.A. Shein, editor. Special

1180 Supplement to the *Bulletin of the American Meteorological Society* 87: S1-S102.

1181

1182 Davidson, G.R., B.C. Laine, S.J. Galicki, and S.T. Threlkeld, 2006. Root zone hydrology: why

1183 bald cypress in flooded wetlands grow more when it rains. *Tree-Ring Research* 62:3-12.

1184

1185 Dell, M., 2012. Path dependence in development: Evidence from the Mexican Revolution.

1186 Working paper, in preparation: <http://scholar.harvard.edu/dell/publications>

1187

1188 Díaz del Castillo, B., 1963. *The Conquest of New Spain*. Penguin Classics. J. M. Cohen (trans.),

1189 Harmondsworth, England

1190

1191 Diaz, S.C., R. Touchan, T.W. Swetnam, 2001. A tree-ring reconstruction of past precipitation
 1192 for Baja California Sur, Mexico. *International Journal of Climatology* 21:1007-1019.

1193

1194 Douglas, A.V., 2007. Summer precipitation variability over Mexico. AGU Joint Assembly,
 1195 Acapulco, Mexico, May 2007.

1196

1197 Eakin, H., 2000, Smallholder maize production and climatic risk: A case study from Mexico.
 1198 *Climatic Change* 45:19-36.

1199

1200 Englehart, P., and A. Douglas, 2002. Mexico's summer rainfall patterns: an analysis of regional
 1201 modes and changes in their teleconnectivity. *Atmosfera* 15:147-164.

1202

1203 Englehart, P., and A. Douglas, 2005. Changing behavior in the diurnal range of surface air
 1204 temperatures over Mexico. *Geophysical Research Letters* 32, L01701.

1205

1206 Federman, D.K., F.I. Arregun Cortes, M. Lopez Perez, 2014. Constructing a framework for
 1207 National Drought Policy: The way forward in Mexico. *Weather and Climate Extremes*
 1208 3:90-94.

1209

1210 Feng, Sh., A.B. Krueger, and M. Oppenheimer, 2010. Linkages among climate change, crop
 1211 yields and Mexico-US cross-border migration. *Proceedings of the National Academy of*
 1212 *Science* 107:14257-14262.

1213

1214 Feng S, Q Hu, and RJ Oglesby, 2011. Influence of Atlantic sea surface temperature on
 1215 persistent drought in the North America. *Climate Dynamics*, 37:569-586.

1216

1217 Feng, S, and Q Fu, 2013. Expansion of global drylands under a warming climate. *Atmospheric*

1218 *Chemistry and Physics*, 13:10081-10094.

1219

1220 Florescano, E., Ed., 1980. Analisis Historico de las Sequias en Mexico (Historical Analysis of

1221 Drought in Mexico). Comision del Plan Nacional Hidraulico, 158 pp.

1222

1223 ,Florescano, E., 1986. Precios del Maiz y Crisis Agricolas en Mexico: 1708–1810 (Prices of

1224 Maize and Agricultural Crises in Mexico: 1708-1810). Ediciones Era, 236 pp.

1225

1226 Florescano, E. and S. Swan. 1995. *Breve Historia de la Sequía en México* (A Brief History of

1227 Drought in Mexico). Universidad Veracruzana, Xalapa, Mexico. (Spanish).

1228

1229 Fulé, P.Z., M. Ramos-Gómez, C. Cortés-Montaña, and A.M. Miller, 2011. Fire regime in a

1230 Mexican forest under indigenous resource management. *Ecological Applications*

1231 21:764–775.

1232

1233 Gerhard, P., 1993. *A Guide to the Historical Geography of New Spain*. University of Oklahoma

1234 Press, Norman.

1235

1236 Giddings, L., M. Soto, B.M. Rutherford, and A. Maarouf, 2005. Standardized precipitation index

1237 zones for Mexico. *Atmosfera* 18:33-56.

1238

1239 Golicher JD, 2006. Correlations between precipitation patterns in southern Mexico and the El

1240 Niño sea surface temperature index. *Interciencia* 31, No. 2.

1241

- Griffin, D., C.A. Woodhouse, D.M. Meko, D.W. Stahle, H.L. Faulstich, C.L. Carillo, R. Touchan, C.L. Castro, S.W. Leavitt, 2013. North American Monsoon precipitation reconstructed from tree-ring latewood. *Geophysical Research Letters* doi: 10.1002/grl.50184.
- Harris, I., P.D. Jones, T.J. Osborn, and D.H. Lister, 2014. Updated high-resolution grids of monthly climatic observations—The CRU TS3.10 dataset. *International Journal of Climatology* 34:623–642.
- Haug, G.H., D. Gunther, L.C. Peterson, D.M. Sigman, K.A. Hughen, and B. Aeschlimann, 2003. Climate and the collapse of Maya civilization. *Science* 299:1731-1735.
- Higgins, R.W., Y. Chen, and A.V. Douglas, 1999. Interannual variability of the North American warm season precipitation regime. *Journal of Climate* 12:653-680.
- Hoaglin, D.C., F. Mosteller, J.W. Tukey, 2000. *Understanding Robust and Exploratory Data Analysis*. Wiley, 472 pp.
- Hodell, D.A., J.H. Curtis, and M. Brenner, 1995. Possible role of climate in the collapse of Classic Maya civilization. *Nature* 375:391-394.
- Hunter, L.M., S. Murray, and F. Riosmena, 2013. Rainfall patterns and US migration from rural Mexico. *International Migration Review* 47:874-909.
- Hurrell, J.W., M.M. Holland, P.R. Gent, 2013. The Community Earth System Model: A Framework for Collaborative Research. *Bulletin of the American Meteorological Society* 94:1339–1360.

1270

1271 Infanti, J.M., and B.P. Kirtman, 2016. North American rainfall and temperature prediction
 1272 response to the diversity of ENSO. *Climate Dynamics* 46:3007-3023.

1273 Jauregui, E., 1979. Algunos aspectos de las fluctuaciones pluviométricas en México en los
 1274 últimos cien años. *Boletín del Instituto de Geografía, UNAM* 9:39-63.

1275

1276 Landa, R., V. Magaña, and C. Neri, 2008. *Agua y Clima: Elementos para la Adaptación al*
 1277 *Cambio Climático*. Secretarí de Medio Ambiente y Recursos Naturales (SEMARNAT),
 1278 De. Tlalpan, México DF. 133 p.

1279

1280 Lanza-Espino, G. y J.L. GarcíaCalderón. 2002. Historical summary of the geology,
 1281 climate, hydrology, culture and natural resource utilization in the Basin of Mexico. In:
 1282 Fenn E., L.I. de Bauer and T. Hernández-Tejeda (Eds). *Urban Air Pollution and Forests;*
 1283 *Resources at Risk in the Mexico City Air Basin*. Springer, New York: 3-23.

1284

1285 Little, E.L., 1999. Digital representation of "Atlas of United States Trees" by Elbert L. Little, Jr.
 1286 U.S. Geological Survey. <http://esp.cr.usgs.gov/info/veg-clim/>

1287

1288 Liverman, D.M., 1990. Drought and Agriculture in Mexico: The case of Sonora and Puebla in
 1289 1970. *Annals of the Association of American Geographers* 80:49-72.

1290

1291 Liverman, D.M., 1999. Vulnerability and Adaptation to Drought in Mexico. *Natural Resources Journal*
 1292 39: 99-115.

1293

1294 Liverman, D.M., R.G. Varady, O. Chavez, R. Sanchez, 1999. Environmental issues along the
 1295 United States-Mexico Border: Drivers of change and responses of citizens and

1296 institutions. *Annual Review Energy and Environment* 24:607-643.

1297

1298 Liverman, D.M. 2001. Adaptation to drought in Mexico. In Wilhite, D., *Drought: A Global*

1299 *Assessment*, pp. 35-45. Routledge: New York.

1300 Lockeretz, W., 1978. The lessons of the Dust Bowl. *American Scientist* 66:560-569.

1301

1302 Magaña, V., J.A. Amador, and S. Medina, 1999. The midsummer drought over México and

1303 Central America. *Journal of Climate* 12:1577-1588.

1304

1305 Magaña, V., J.L. Vazquez, J.L. Perez, and J.B. Perez, 2003. Impact of El Niño on precipitation

1306 in Mexico. *Geofísica Internacional* 42:313-330.

1307

1308 Mantua, N.J., S.R. Hare, Y. Zhang, J.M. Wallace, and R.C. Francis, 1997. A Pacific

1309 interdecadal climate oscillation with impacts on salmon productin. *Bulletin of the*

1310 *American Meteorological Society* 78:1069-1079.

1311

1312 Martinez, M., 1963, *Las Pinaceas Mexicanas*, 400 pp., Universidad NacionalAutonoma de

1313 Mexio, Ciudad Universitaria, Mexico 20, D.F.

1314

1315 McLeman, R.A., J. Dupre, L.B. Ford, J. Ford, K. Gajewski, and G. Marchildon, 2014. What we

1316 learned from the Dust Bowl: lessons in science, policy, and adaptation. *Population and*

1317 *Environment* 35:417-440.

1318

1319 Meko, D.M., R. Touchan, J. Villanueva Diaz, D. Griffin, C.A. Woodhouse, C.L. Castro, C.

1320 Carrillo, and S.W. Leavitt, 2013. Sierra San Pedro Martir, Baja California, cool-season

- precipitation reconstructed from earlywood width of *Abiesconcolor* tree rings. *Journal of Geophyscial Research Biogeosciences*doi: 10.1002/2013JG002408.
- Melvin, T.M., K.R. Briffa, K. Nicolussi, andM. Grabner, 2007.Time-varying-response smoothing.*Dendrochronologia* 25:65–69.
- Melvin, T.M., and K.R. Briffa, 2008.A “signal-free” approach to dendroclimatic standardization. *Dendrochronologia*26:71-86.
- Mendez, M., Magaña, V., 2010. Regional aspects of prolonged meteorological droughts over Mexico and Central America.*Journal of Climate* 23:1175–1188.
- Metcalfe, S.E., J.A. Barron, and S.J. Davies, 2015. The Holocene history of the North American monsoon: ‘known knowns’ and ‘known unknowns’ in understanding its spatial and temporal complexity. *Quaternary Science Reviews* 120:1-27.
- Milbrath, S, 2013.*Heaven and Earth in Ancient Mexico*. University of Texas Press, Austin 224p.
- Mosiño, P.A., and E. García, 1974, The climate of Mexico. In *Climates of North America*, edited by R.A. Bryson and F.K. Hare, pp., 345-404, Elsvier, New York.
- Nicholas, R.E., and D.S. Battisti, 2008. Drought recurrence and seasonal rainfall prediction in the Rio Yaqui Basin, Mexico. *Journal of Applied Meteorology and Climatology* 47:991-1005.
- Oglesby, R., S. Feng, Q. Hu, and C. Rowe, 2012. The role of the Atlantic Multidecadal Oscillation on medieval drought in the North America: synthesizing results

1347 from proxy data and climate models. *Global and Planetary Change* 84:85, 56-65.

1348

1349 O'Hara, S.L., and S.E. Metcalfe, 1997. The climate of Mexico since the Aztec Period.

1350 *Quaternary International* 43/44:25-31.

1351

1352 Palmer, W.C., 1965. *Meteorological Drought*. Research Paper No. 45, US Dept Commerce,

1353 Weather Bureau.

1354

1355 Pascual, R., A. Albanil, J.L. Vazquez, 2015. Mexico (in "State of the Climate in 2014").

1356 *Bulletin of the American Meteorological Society* 96(7):S172-173.

1357

1358 Pavia EG, F Graef, and J Reyes, 2006. PDO-ENSO effects in the climate of Mexico. *Journal of*

1359 *Climate* 19:6433–6438

1360

1361 Portig, W.H., 1965. Central American rainfall. *Geographical Review* 55:68-90.

1362

1363 Quiñones-Keber, E., 1995. *Codex Telleriano-Remensis*. University of Texas Press, 365 pp.

1364 Ropelewski, C.F., and M.S. Halpert, 1987. Global and regional scale precipitation patterns a

1365 ssociated with the El Niño/Southern Oscillation. *Monthly Weather Review*, 115:1606-1

1366 626.

1367 Ropelewski, C.F., and M.S. Halpert, 1989. Precipitation patterns associated with the high index

1368 phase of the Southern Oscillation. *Journal of Climate*, 2:268-284.

1369

1370 Sano, M., C. Xu, and T. Nakatsuka, 2012. A 300-year Vietnam hydroclimate and ENSO

1371 variability record reconstructed from tree ring $\delta^{18}O$. *Journal of Geophysical Research*

1372 117, D12115.

1373

1374 Seager, R., M. Ting, I. Held, Y. Kushnir, J. Lu, G. Vecchi, H. Huang, N. Harnik, A. Leetma, N.

1375 Lau, C. Li, J. Velez, and N. Naik, 2007. Model projections of an imminent transition to a

1376 more arid climate in southwestern North America. *Science* 316:1181-1184.

1377

1378 Seager, R., M. Ting, M. Davis, M. Cane, N. Naik, J. Nakamura, C. Li, E. Cook, and D.W. Stahle,

1379 2009. Mexican drought: An observational, modeling and tree ring study of variability and

1380 climate change. *Atmosfera* 22:1-31.

1381

1382 Sheppard, P.R., A.C. Comrie, G.D. Packin, K. Augersbach, M.K. Hughes, 2002. The climate of

1383 the U.S. Southwest. *Climate Research* 21:219-238.

1384

1385 Simpson, L.B., 1966. *Many Mexicos*. University of California Press, Berkeley. 408 p.

1386

1387 Skoufias, E., and K. Vinha, 2013. The impacts of climate variability on household welfare in

1388 rural Mexico. *Population and Environment* 34:370-399.

1389

1390 St. George, S., 2014. An overview of tree-ring width records across the Northern Hemisphere.

1391 *Quaternary Science Reviews* 95:132-150.

1392

1393 Stahle, D.W., and M.K. Cleaveland, 1992. Reconstruction and analysis of spring rainfall over

1394 the Southeastern U.S. for the past 1000 years. *Bulletin of the American Meteorological*

1395 *Society*, 73:1947–1961.

1396

1397 Stahle, D.W., M.K. Cleaveland and J.G. Hehr, 1988. North Carolina climate changes

1398 reconstructed from tree– rings: A.D. 372 to 1985. *Science* 240:1517– 1519.

1399

1400 Stahle, D.W., R.D. D'Arrigo, P.J. Krusic, M.K. Cleaveland, E.R. Cook, R.J. Allan, J.E. Cole,

1401 R.B. Dunbar, M.D. Therrell, D.A. Gay, M.D. Moore, M.A. Stokes, B.T. Burns, J.

1402 Villanueva-Diaz, L.G. Thompson, 1998b. Experimental dendroclimatic reconstruction of

1403 the Southern Oscillation. *Bulletin of the American Meteorological Society* 79:2137-

1404 2152.

1405

1406 Stahle, D.W., E.R. Cook, M.K. Cleaveland, M.D. Therrell, D.M. Meko, H.D. Grissino-Mayer, E.

1407 Watson, and B.H., Luckman, 2000b. Tree-ring data document 16th century

1408 megadrought over North America, *Eos, Transactions of the American Geophysical*

1409 *Union* 81:212, 125.

1410

1411 Stahle, D.W., F.K. Fye, and E.R. Cook, 2007. Tree-ring reconstructed megadroughts over

1412 North America since AD 1300. *Climatic Change* 83:133-149.

1413

1414 Stahle, D.W., E.R. Cook, J. Villanueva Diaz, F.K. Fye, D.J. Burnette, R.D. Griffin, R. Acuña-

1415 Soto, R. Seager, and R.R. Heim Jr., 2009. Early 21st century drought in Mexico. *Eos*

1416 90:89-90.

1417

1418 Stahle, D.W., and J.S. Dean, 2011. North American tree rings, climatic extremes,

1419 and social disasters. *Dendroclimatology: Progress and Prospects*, M.K. Hughes, T.W.

1420 Swetnam, H.F. Diaz, editors. *Developments in Paleoenvironmental Research*, Vol. 11,

1421 Springer, pp. 297-327.

1422

1423 Stahle, D. W., J. Villanueva Diaz, D. J. Burnette, J. CeranoParedes, R. R. Heim, Jr., F. K. Fye,

1424 R. Acuña-Soto, M. D. Therrell, M. K. Cleaveland, and D. K. Stahle, 2011a. Major
 1425 Mesoamerican droughts of the past millennium, *Geophysical Research Letters* , 38,
 1426 L05703, doi:10.1029/2010GL046472.
 1427
 1428 Stahle, D. W., D.J. Burnette, J. Villanueva Diaz, R.R. Heim, Jr., F.K. Fye, J. CeranoParedes, R.
 1429 R. Acuña-Soto, M.K. Cleaveland, 2011b. Atlantic and Pacific influences on
 1430 Mesoamerican climate over the past millennium. *Climate Dynamics* DOI
 1431 10.1007/s00382-011-1205-z.
 1432
 1433 Stahle, D. W., D.J. Burnette, J. Villanueva Diaz, F.K. Fye, R.D. Griffin, M.K. Cleaveland, D.K.
 1434 Stahle, J.R. Edmondson, K. Wolff, 2012. Tree-ring analysis of ancient baldcypress trees
 1435 and subfossil wood. *Quaternary Science Reviews* 34:1-15.
 1436
 1437 Standley, P.C., 1967. *Trees and Shrubs of Mexico*. Smithsonian Institution, Washington DC,
 1438 1721 p.
 1439
 1440 Stothers, R.B., 1984. The Great Tambora Eruption in 1815 and its aftermath. *Science*
 1441 224:1191-1198.
 1442
 1443 Therrell, MD, 2005. Tree rings and 'El Año de Hambre' in Mexico. *Dendrochronologia* 22:203-
 1444 207.
 1445
 1446 Therrell, M.D., D.W. Stahle, M.K. Cleaveland, and J. Villanueva-Diaz, 2002. Warm season tree
 1447 growth and precipitation over Mexico. *Journal of Geophysical Research* 107(D14):ACL
 1448 6-1 to 6-8.
 1449

1450 Therrell, M.D., D.W. Stahle, R. Acuña Soto, 2004. Aztec drought and the curse of One Rabbit.
 1451 *Bulletin of the American Meteorological Society* 85:1263-1272.
 1452
 1453 Therrell, M.D., D.W. Stahle, M.K. Cleaveland, J. Villanueva-Diaz, E. Cornejo Oviedo, 2006.
 1454 Tree-ring reconstructed maize yield in Central Mexico: 1474-2001. *Climatic Change*
 1455 74(4)493-504.
 1456
 1457 Trouet, V., and G.J. van Oldenborgh, 2013. KNMI Climate Explorer: A web-based research tool
 1458 For high-resolution paleoclimatology. *Tree-Ring Research* 69:3–13.
 1459
 1460 Tutino, J., 1986. From Insurrection to Revolution in Mexico. Princeton University Press,
 1461 Princeton, New Jersey. 425 p.
 1462
 1463 van der Schrier, G., J. Barichivich, K.R. Briffa, and P.D. Jones, 2013. A scPDSI-based global
 1464 data set of dry and wet spells for 1901–2009. *Journal of Geophysical Research*
 1465 118:4025–4048.
 1466
 1467 Vazquez, J.L., 2009. Climate variability and changes in the climate extremes of Mexico. PhD
 1468 Dissertation, University of East Anglia, Norwich, UK..
 1469
 1470 Vera, C., W. Higgins, J. Amador, T. Ambrizzi, R. Gerreaud, D. Gochis, D. Gutzler, D.
 1471 Lettenmaier, J. Marengo, C.R. Mechoso, J. Nogue-Paegle, P.L. Silva Dias, and C.
 1472 Zhang, 2006. Toward a Unified View of the American Monsoon Systems. *Journal of*
 1473 *Climate* 19:4977-5000.
 1474
 1475 Villanueva, J., D. Stahle, B. Luckman, J. Cerano-Paredes, M. Therrell, M.K. Cleaveland,

- and E. Cornejo-Oviedo, 2007. Winter-spring precipitation reconstruction from tree rings for northeast Mexico. *Climatic Change* 83:117-131.
- Villanueva, J., D.W., Stahle, B.H. Luckman, J.Ceranno-Paredes, M.D. Therrell, R. Moran Martinez, and M.K. Cleaveland, 2007. Potencial dendrocronológico de *Taxodium mucronatum* Ten. y acciones para su conservación en México. *Ciencia Forestal en México* 32 (101), 9e38.
- Wang, C., D.B. Enfield, S. Lee, and C.W. Landsea, 2006. Influences of the Atlantic warm pool on western hemisphere summer rainfall and Atlantic hurricanes. *Journal of Climate* 19:3011-3028
- Wilder M. and P. R. Lankao, 2006. Paradoxes of decentralization: Water reform and social implications in México. *World Development* 34:1977-1995.
- Wolter, K., and M.S. Timlin, 2011. El Niño/Southern Oscillation behaviour since 1871 as diagnosed in an extended multivariate ENSO index (MEI.ext). *International Journal of Climatology* 31:1074-1087.
- Zhang, T., M.P. Hoerling, J. Perlwitz, D. Sun, and D. Murray, 2011. Physics of U.S. surface temperature response to ENSO. *Journal of Climate* 24:4874-4887.

1499

1500

1501

1502

1503

1504

1505

1506

1507

1508

1509

1510

Figure 1. The native distribution of Douglas-fir (*Pseudotsugamenziesii*; a) and baldcypress (*Taxodiumdistichum* and *T. mucronatum*; b) are mapped (from Little 1999). Small local populations are indicated by 'x' and additional populations may exist to the south of the known range for both species (indicated by '?'). A few local populations of *Taxodiummucronatum* have been added to the mapping by Little (1999) in Mexico, based on more recent field research by the University of Arkansas and INIFAP.

1511

1512

1513

1514 Figure 2. The spatial distribution of the 252 tree-ring chronologies used for
1515 reconstruction of the soil moisture balance in the MXDA. The start year of each
1516 chronology is classified into 200-year groups.

1517

1518

1519

Mexican Drought Atlas

Grid Points

1520

1521

1522 Figure 3. The 0.5° latitude/longitude grid of instrumental scPDSI used to calibrate the
1523 soil moisture reconstructions in the MXDA (1501 total grid points). The four sub-regions
1524 of Mexico analyzed in the text are also indicated.

1525

1526

1527

1528

1529

1530

1531

1532

Figure 4. The correlation between June scPDSI and the average of July and August scPDSI is computed and plotted for all 1501 grid points of instrumental data used for the MXDA. All available data were used from 1901-2012, but the pattern is very similar when restricted to the more data rich period from 1951-2012.

1533

1534

1535

1536

1537

1538

1539

1540

1541

Figure 5. Calibration and verification statistics calculated for the gridded scPDSI reconstructions are mapped for the 1944-1984 (a,b) and 1920-1943 (c,d,e) time periods, respectively. All abbreviations are defined in the text. The average correlation (RBAR) between the 16 individual reconstructions at each grid point (in both 8-member ensembles) is computed and mapped for the full period of reconstruction, 1400-1984 (f).

1542

Significant Droughts in Mexican History

1543

1544

1545

1546

1547

1548

Figure 6. The spatial pattern and intensity of tree-ring reconstructed scPDSI during four significant droughts in Mexican history are mapped using the online analytical tools now available with the MXDA (Appendix 1).

1549

1550

1551

1552

1553

1554

1555

1556

Figure 7. The tree-ring reconstructed scPDSI is mapped for four years near the onset of the Mexican War of Independence, which began with the “Grito de Dolores” on September 16, 1810.

1557

1558

1559

1560

1561

1562

1563

1564

1565

Figure 8. The instrumental scPDSI data were averaged and plotted for the four sub-regions of Mexico identified in Figure 3 for the period 1920-2012. Decadal variability is highlighted with a cubic smoothing spline (Cook and Peters 1981). Positive values of PDSI indicate wetness, negative values indicate drought.

1566

1567

1568

1569

1570

1571

1572

1573

Figure 9. The tree-ring reconstructed scPDSI data were averaged and plotted for the four sub-regions of Mexico (red). Decadal variability is highlighted with a cubic smoothing spline (black). The reconstructions for northwest (a) and northeast Mexico are reasonably well replicated back to 1400, but the reconstructions for central (c) and southeast Mexico (d) are based on few long chronologies before 1550.

1574

1575

1576

1577

1578

1579

1580

1581

1582

1583

Figure 10. The gridded tree-ring reconstructions for five of the most severe and sustained droughts over Mexico in the last 600-years are averaged and mapped. The most intense event in the well-replicated period after 1550 appears to have occurred during the 1950s and was concentrated over northern Mexico in both the reconstructed (e) and instrumental data (f; see also Figures 9 and 20). Note the spatial inconsistencies in instrumental scPDSI during this seven-year drought (f), especially in northern Durango.

Decadal Droughts of Central Mexico

Figure 11. The most severe and sustained reconstructed droughts that were concentrated in central Mexico appear to have occurred from 1625-1634 (a), 1696-1705 (b) and 1894-1899 (c), excluding the period from 1400-1550 when the reconstructions are based on few chronologies from this region (see also Figure 9).

Severe Droughts of Southeastern Mexico

1593

1594

1595

1596

1597

1598

1599

Figure 12. The most severe and sustained reconstructed droughts that were concentrated in southeastern Mexico are estimated to have occurred from 1585-1589 (a), 1767-1769 (b), 1811-1820 (c), 1936-1946 (d), and 1971-1975 (f) excluding the period before 1550 when the reconstructions for SE Mexico are based on few local chronologies (see also Figure 9). The instrumental scPDSI are averaged and mapped for the two 20th century droughts (e, g).

Major Mexican Pluvials

Figure 13. Several exceptionally wet episodes that covered most of Mexico and lasted from four to seven years are reconstructed with the MXDA, including one of the wettest periods from 1814-1817 (f) that included the aftermath of the Tambora volcanic eruption in April 1815.

Figure 14. (a) The ENSO teleconnection to reconstructed scPDSI over Mexico is illustrated with this map of correlation coefficients computed between each grid point and the Feb-May extended MEI (Wolter and Timlin 2011) during the 1901-1984 period. (b) Same as (a) using the instrumental scPDSI.

1613

1614

1615

1616

1617

1618

1619

1620

1621

Figure 15. The correlations between indices of the AMO and PDO are plotted for reconstructed (a,c) and instrumental JJA scPDSI (b,d) for the common time period of 1901-1984. The highest single grid point correlations are <0.5 for the AMO and the PDO in both the reconstructed and instrumental comparisons. The AMO indices were averaged annually for Jul-Jun (monthly anomalies from UK Met Office HadSST 3.1.1.0 for 25-60N, 7-75W, minus regression on global mean temperature). The PDO indices were averaged for the winter season (DJF; after Mantua et al. 1997).

Stability of the ENSO Teleconnection to Mexico

Figure 16. Same as Figure 14a for 30-year subperiods (only 24-years available in d).

1626

1627

1628

1629

1630

1631

1632

1633

1634

1635

1636

Figure 17. The average of reconstructed scPDSI is mapped for the 15 most extreme warm events (a) and 15 most extreme cold events (b) identified in the Jan-Feb extended MEI for the period 1871-1984 in common with the tree-ring estimates (extreme warm and cold years identified in Figures 18 and 19, respectively). The instrumental scPDSI is averaged and mapped for the 10 most extreme warm (c) and 10 most extreme cold events (d) for the period 1901-1984 to compare with the registration of the ENSO signal in the tree-ring reconstructions (warm and cold extremes also identified in Figures 18 and 19 for the 1901-1984 interval).

1637

Reconstructed PDSI in Mexico During El Nino Years

1638

1639 Figure 18. Reconstructed scPDSI is mapped for each of the 15 most extreme warm
1640 events recorded in the Feb-May extended MEI from 1871-1984.

1641

1642

1643

1644

1645 Figure 19. Reconstructed scPDSI is mapped for each of the 15 most extreme cold

1646 events recorded in the Feb-May extended MEI from 1871-1984.

1647

1649

1650

1651

1652

1653

1654

1655

1656

1657

1658

Figure 20. The significant ($p < 0.1$) correlation coefficients computed between reconstructed (left panels) and instrumental (right panels) JJA scPDSI in three subregions of Mexico and gridded SSTs from the HadISST data set are mapped (Rayner et al. 2003; using the KNMI *Climate Explorer*, Trouet and van Oldenborgh 2013). The length of the available data and the strength of signal dictated the seasons and time periods used. Note the shift in seasonality from northwest to central Mexico (a,b to c,d), and the change in signal strength from the Pacific to the Atlantic from northwest to southern Mexico (a,b to e,f).

Figure 21. (a) An index of the Mexican moisture dipole is plotted from 1400 to 2012 (y-axis inverted), based on a map pattern correlation with the north-south anomaly reconstructed for 1951-1957 (see Figure 10e). The 15 most extreme positive (b) and negative values (c) were averaged and mapped to illustrate the north-south spatial pattern over Mexico. (d) The Nov-Feb extended MEI (red) is plotted with the dipole index (black; both normalized) to highlight variations in the ENSO influence on the moisture gradient ($r = 0.58$, 1871-2005; correlations for 30-year sub-periods also noted).

1668

1669

1670

1671

1672

1673

1674

1675

Figure 22. The correlation between the central Pacific SST average (Niño 4 region) for the JFMAM season with JFMAM total precipitation (a) and JFMAM average temperature (b) using the millennium simulations computed with the Community Earth System Model. Note the generally positive correlations with precipitation and negative correlations with temperature over northern Mexico.

1676

1677

1678

1679

1680

1681

1682

1683

1684

1685

Figure 23. (a) The Drought Area Index (DAI) for the mainland of Mexico was computed as the percentage of grid points between 14-30°N and 90-110°W that were reconstructed as scPDSI ≤ -1.0 (i.e., exclusive of the Baja and Yucatan Peninsulas). Widespread droughts spanning most of mainland Mexico are indicated by the highest DAI values (red) and multi-decadal variability is indicated by the 30-year smooth curve (black). The most extreme “All Mexico” drought is reconstructed for 1819. The 15 years with the lowest (wet) and highest (dry) Drought Area Indices were

1686 averaged and mapped to document the nationwide impact of these extremes(b and c,
1687 respectively).

Significant Droughts in Mexican History

Graphical abstract: The Mexican Drought Atlas uses 252 climate sensitive tree-ring chronologies to reconstruct the summer Palmer Drought Severity Index at 1,501 0.5° latitude/longitude grid points over Mexico and adjacent land areas. Four severe droughts are depicted here, including three previously documented events (a,c,d) and the reconstructed drought that afflicted much of Mexico soon after Conquest (b).