Gradual and abrupt changes during the Mid-Pleistocene Transition

Heather L. Ford¹, Sindia Sosdian², Yair Rosenthal³, Maureen E. Raymo¹

¹Biology and Paleo Environment, Lamont-Doherty Earth Observatory, Palisades, New York, USA

²Earth and Ocean Sciences, Cardiff University, Cardiff, UK

³Department of Marine and Coastal Sciences and Department of Earth and Atmospheric Sciences, Rutgers University, New Brunswick, New Jersey, USA

*Corresponding author: hford@ldeo.columbia.edu

Abstract

During the Mid-Pleistocene Transition (MPT), the dominant glacial-interglacial cyclicity as inferred from the marine $\delta^{18}\text{O}$ records of benthic foraminifera ($\delta^{18}\text{O}_{\text{benthic}}$) changed from 41 kyr to 100 kyr years in the absence of a comparable change in orbital forcing. Currently, only two Mg/Ca-derived, high-resolution bottom water temperature (BWT) records exist that can be used with $\delta^{18}\text{O}_{\text{benthic}}$ records to separate temperature and ice volume signals over the Pleistocene. However, these two BWT records suggest a different pattern of climate change occurred over the MPT—a record from North Atlantic DSDP Site 607 suggests BWT decreased with no long-term trend in ice volume over the MPT, while South Pacific ODP Site 1123 suggests that BWT has been relatively stable over the last 1.5 Myr but that there was an abrupt increase in ice volume at ~900 kyr. In this paper we attempt to reconcile these two views of climate change across the MPT. Specifically, we investigated the suggestion that the secular BWT trend obtained from Mg/Ca measurements on *Cibicides wuellerstorfi* and *Oridorsalis umbonatus* species from North Atlantic Site 607 is biased by the possible influence of $\Delta[\text{CO}_3^{2-}]$ on Mg/Ca values in these species by generating a low-resolution BWT record using *Uvigerina* spp., a genus whose Mg/Ca

values are not thought to be influenced by $\Delta[\text{CO}_3^{2-}]$. We find a long-term BWT cooling of ~ 2 -
 3°C occurred from 1500 to ~ 500 kyr in the N. Atlantic, consistent with the previously generated
C. wuellerstorfi and *O. umbonatus* BWT record. We also find that changes in ocean circulation
likely influenced $\delta^{18}\text{O}_{\text{benthic}}$, BWT, and $\delta^{18}\text{O}_{\text{seawater}}$ records across the MPT. N. Atlantic BWT
cooling starting at ~ 1.2 Ma, presumably driven by high-latitude cooling, may have been a
necessary precursor to a threshold response in climate-ice sheet behavior at ~ 900 ka. At that
point, a modest increase in ice volume and thermohaline reorganization may have caused
enhanced sensitivity to the 100 kyr orbital cycle.

1. Introduction

Variations in the Earth's orbit, and consequently incoming solar radiation, act as a
pacemaker for glacial-interglacial cycles [Hays et al., 1976]. During the mid-Pleistocene
transition (MPT, ~ 1.2 to 0.65 Ma) the glacial-interglacial periodicity as recorded by marine $\delta^{18}\text{O}$
records of benthic foraminifera ($\delta^{18}\text{O}_{\text{benthic}}$) changed from primarily 41 to 100 kyr year cycles
without any obvious change in external orbital forcing [Shackleton and Opdyke, 1976; Pisias and
Moore, 1981; Imbrie et al., 1992; Raymo and Nisancioglu, 2003]. As the 100 kyr eccentricity
cycle has a relatively weak effect on incoming solar radiation [Imbrie et al., 1993], feedbacks
within the climate system must have amplified the glacial-interglacial response to the 100 kyr
cycle. However, after nearly 40 years of research, the processes and mechanisms that caused the
MPT remain enigmatic [Clark et al., 2006].

Explanations as to what caused the MPT are diverse (summarized in [McClymont et al.,
2013]), and they generally invoke changes in climate boundary conditions such as: (1) gradual
global cooling, possibly related to $p\text{CO}_2$ decrease [Berger and Jansen, 1994; Raymo et al., 1997],

(2) changes in ice sheet dynamics that allow for the build up of large ice sheets[*Clark et al.*, 2006; *Elderfield et al.*, 2012], and/or (3) thermohaline circulation reorganization [*Schmieder et al.*, 2000; *Sexton and Barker*, 2012; *Pena and Goldstein*, 2014]. Much of our understanding of the MPT comes from $\delta^{18}\text{O}_{\text{benthic}}$ records, which reflect a combination of bottom water temperature (BWT) and seawater oxygen isotope composition ($\delta^{18}\text{O}_{\text{seawater}}$)[*Shackleton*, 1967; *Pisias and Moore*, 1981; *Labeyrie et al.*, 1987; *Maasch*, 1988; *Ruddiman et al.*, 1989; *Saltzman and Maasch*, 1991; *Mudelsee and Schulz*, 1997; *Rutherford and D'Hondt*, 2000; *Lisiecki and Raymo*, 2005; 2007]. Despite years of study and statistical analysis, disagreement still exists as to whether the MPT reflects a gradual transition that started ~ 1.2 Ma and finished ~ 0.65 Ma[*Pisias and Moore*, 1981; *Ruddiman et al.*, 1989; *Rutherford and D'Hondt*, 2000; *Clark et al.*, 2006; *Sosdian and Rosenthal*, 2009; *McClymont et al.*, 2013] or represents an abrupt event centered around 0.9 Ma [*Maasch*, 1988; *Mudelsee and Schulz*, 1997; *Elderfield et al.*, 2012]. The timing of the MPT is particularly relevant to evaluating the mechanisms of climate change (e.g., a gradual response of ice to a gradual forcing, an abrupt response of ice to an abrupt forcing event, or a threshold (abrupt) response of ice to long-term gradual climate forcing). Because $\delta^{18}\text{O}_{\text{benthic}}$ records a number of environmental parameters (global ice volume, BWT, and local hydrography), part of the disagreement over the timing and character of the MPT may be related to the use of different $\delta^{18}\text{O}_{\text{benthic}}$ records, with local hydrographic effects obscuring global trends in temperature and ice volume in some records [*Clark et al.*, 2006; *Elderfield et al.*, 2012; *Bates et al.*, 2014].

Because $\delta^{18}\text{O}_{\text{benthic}}$ reflects BWT and $\delta^{18}\text{O}_{\text{seawater}}$ (e.g. ice volume and local hydrography), several environmental signals are imprinted on any given $\delta^{18}\text{O}_{\text{benthic}}$ record. To isolate BWT and $\delta^{18}\text{O}_{\text{seawater}}$ components of $\delta^{18}\text{O}_{\text{benthic}}$ records, the Mg/Ca ratio of benthic foraminiferal calcite is

widely used as an independent proxy of BWT on orbital and tectonic timescales [*Lear et al.*, 2000; *Billups and Schrag*, 2002; *Sosdian and Rosenthal*, 2009; *Elderfield et al.*, 2010; 2012].

Currently, only two high-resolution coupled benthic foraminifera Mg/Ca and $\delta^{18}\text{O}_{\text{benthic}}$ records exist across the MPT and their implications for how global ice volume changed across the MPT disagree. The first record is the N. Atlantic (the *Sosdian and Rosenthal* (2009) record from Deep Sea Drilling Project (DSDP) Site 607, 41.0° N, 33.0° W, 3427 m water depth), and the second is from the S. Pacific (the *Elderfield et al.* (2012) record from Ocean Drilling Program (ODP) Site 1123 located on the Chatham Rise, 41.8° S, 171.5° W, 3290 water depth) (Figure 1). The interpretation of Site 607 record has been challenged based on the observation that, in addition to temperature, changes in carbonate ion saturation ($\Delta[\text{CO}_3^{2-}]$) could influence Mg/Ca in benthic species thus biasing BWT estimates in some species (e.g. *Cibicides wuellerstorfi*; [*Yu and Broecker*, 2010]; but also see [*Sosdian and Rosenthal*, 2010]). The $\Delta[\text{CO}_3^{2-}]$ is defined as $[\text{CO}_3^{2-}]_{\text{in situ}} - [\text{CO}_3^{2-}]_{\text{saturation}}$.

In addition, the distinct differences in implied BWT histories at Sites 607 and 1123 (and thus climate interpretations) also merit further study. For instance, The N. Atlantic *Sosdian and Rosenthal* (2009) record at Site 607 suggests that significant BWT cooling ($\sim 2^\circ\text{C}$) occurred over an interval of a million years, from ~ 1.5 to 0.5 Ma. On the other hand, The S. Pacific *Elderfield et al.* (2012) record from Site 1123 shows relatively constant, near-freezing temperatures over the entire 1.5 Myr record. Although a significant volume of global water flows over Chatham Rise, this area is also sensitive to the relative mixture of northern and southern component water masses (Figure 1).

Given that these sites are in different oceans and at different depths, it is possible that they could be characterized by different BWT trends; however, when the Mg/Ca-derived BWTs are

subtracted from the $\delta^{18}\text{O}_{\text{benthic}}$ records at each site, two very different patterns of $\delta^{18}\text{O}_{\text{seawater}}$ are observed across the MPT implying seemingly contradictory global ice volume histories. Specifically, Site 607 shows no discernible long-term trend in $\delta^{18}\text{O}_{\text{seawater}}$ across the MPT [Sosdian and Rosenthal, 2009], whereas Site 1123 record shows an abrupt increase occurring at ~ 900 ka, which Elderfield et al. [2012] attributes to a large expansion of Antarctic ice volume. Instead of global cooling of the deep ocean across the early Pleistocene, the site 1123 record was interpreted as suggesting that an abrupt change in ice sheet dynamics and/or ice volume increase may have been the primary cause of the MPT.

Here we revisit DSDP Site 607 and generate a Mg/Ca-derived BWT record from 500 to 1500 ka using *mixed Uvigerina* spp.. We compare this record to the previously published *C. wuellerstorfi* and *Oridorsalis umbonatus* record [Sosdian and Rosenthal, 2009] in order to evaluate long-term changes in N. Atlantic BWT over the MPT. We also evaluate the possible influence of $\Delta[\text{CO}_3^{2-}]$ on Mg/Ca BWT reconstructions based on *Uvigerina* spp., *C. wuellerstorfi* and *Oridorsalis umbonatus*. We compare $\delta^{18}\text{O}_{\text{benthic}}$, BWT and $\delta^{18}\text{O}_{\text{seawater}}$ records from Site 607 with those from Site 1123 in the S. Pacific. Lastly, we consider how changes in ocean circulation could influence the interpretation of $\delta^{18}\text{O}_{\text{benthic}}$, BWT and $\delta^{18}\text{O}_{\text{seawater}}$ records from these two sites.

3. Materials and methods

Mg/Ca were measured on *mixed Uvigerina* spp. from DSDP 607 to estimate bottom water temperature. Previous calibration studies suggest that the Mg/Ca ratios of infaunal species such as *Uvigerina* spp. primarily respond to temperature [Elderfield et al., 2010] and show very little $\Delta[\text{CO}_3^{2-}]$ effect [Yu and Elderfield, 2007]. $\Delta[\text{CO}_3^{2-}]$ approaches zero at shallow depths within the sediment because pore waters come to rapid equilibration with calcium carbonate,

regardless as to whether the bottom water $\Delta[\text{CO}_3^{2-}]$ is oversaturated or undersaturated at particular location [Elderfield et al., 2006]. By contrast, calibration studies suggest that the Mg/Ca ratios of epifaunal species such as *Cibicidoideswuellerstorfi* respond to both temperature and changes in $\Delta[\text{CO}_3^{2-}]$ [Elderfield et al., 2006; Yu and Elderfield, 2007; Elderfield et al., 2010], possibly resulting in biased temperature reconstructions in areas with markedly large changes in bottom water $\Delta[\text{CO}_3^{2-}]$ [Yu and Broecker, 2010]. Thus, it is hypothesized that Mg/Ca measurements derived from *Uvigerina* spp. would more accurately reflect BWT variations through time [Elderfield et al., 2010; 2012].

Uvigerina spp. specimens were picked from the $>150\text{ }\mu\text{m}$ size fraction with approximately 5-15 specimens analyzed per sample, with a median of 7 specimens per sample. The average sampling resolution is one sample every $\sim 9,000$ years. The *Uvigerina* spp. measurements are relatively uniformly distributed between glacial and interglacial periods (Supplemental Figure 1). The vast majority ($>88\%$) of the samples are from the same core samples that were used to generate the Ruddiman et al., [1989] $\delta^{18}\text{O}_{\text{benthic}}$ record. Samples were cleaned using standard analytical procedures [Rosenthal et al., 1999; Martin and Lea, 2002] - samples were repeatedly rinsed in water and methanol to remove clays and cleaned with reductive and oxidative reagents. Owing to small sample weights ($\sim 100\text{ }\mu\text{g}$), not all samples were subjected to a weak acid leaching. Elemental measurements were made either at Lamont-Doherty Earth Observatory (LDEO) on a Thermo Scientific iCAP-Q inductively coupled plasma mass spectrometer (ICP-MS) or at Rutgers University on a Finnigan Element-XR ICP-MS. Al/Ca, Fe/Ca, Mn/Ca and Ti/Ca (Supplemental Figure 2) were used to monitor samples for contamination due to diagenetic overprinting or to residual detrital material ($n=6$ samples removed) [Rosenthal et al., 1999]. We find no obvious downcore trends in Mg/Ca and Al/Ca,

Fe/Ca, and Ti/Ca. Mn/Ca values are generally high ($\mu = 126.5 \mu\text{mol/mol}$, $\sigma = 53.56 \mu\text{mol/mol}$) and correlate with Mg/Ca ($R^2 = 0.33$, <0.001). These high Mn/Ca values do not appear to be due to Mn-Fe oxide and MnCO_3 contamination because all samples were reductively cleaned and we found that after repetitive weak acid leaches, the Mn/Ca values remained relatively high within a subset of samples. There is a weak correlation downcore between Mn/Ca and Mg/Ca but we believe this is a coincidence not caused by the same process. Elderfield et al. [2012] see a similar weak correlation downcore at Site 1123 between Mn/Ca and Mg/Ca that they attribute to diagenetic reorganization. A similar process may have occurred at Site 607.

Instrumental precision at LDEO and Rutgers was 0.5-1%. Long-term precision of a liquid consistency standard at LDEO and Rutgers University was between 1-2%. At LDEO, reproducibility of three consistency standards with Mg/Ca of 1.4840, 4.1325, 8.1035 mmol/mol were $\pm 1.3\%$, $\pm 1.8\%$, and $\pm 1.7\%$, respectively. At Rutgers University, reproducibility of consistency standards with Mg/Ca of 1.25, 3.32, and 7.51 mmol/mol were $\pm 1.2\%$, $\pm 1.2\%$, and $\pm 0.6\%$, respectively. To ensure inter-laboratory accuracy, standards from Rutgers University were run at LDEO. Inter-laboratory accuracy was found to be 3.0%, within the range of previously reported inter-laboratory comparisons [Rosenthal et al., 2004].

There are several species-specific calibrations to choose from to account for known differences in temperature sensitivity and offsets for various benthic foraminifera species [Lear et al., 2002; Healey et al., 2008; Yu and Elderfield, 2008; Sosdian and Rosenthal, 2009; Elderfield et al., 2012; Cappelli et al., 2015]. For ease of comparison with the site 1123 *Uvigerina* spp. record, we use a modified version of the temperature equation, and the same $\delta^{18}\text{O}$ equation, as used by Elderfield et al. (2010, 2012). To correct for known Mg loss during the reductive step [Barker et al., 2003; Rosenthal et al., 2004] used in our cleaning procedure, we adjusted

the Elderfield *Uvigerina* spp. paleotemperature equation by 10%: $Mg/Ca = 0.9 + 0.1 * BWT$. Previously published *Cibicides wuellerstorfi*, and *Oridorsalis umbonatus* paleotemperature equations were used for further BWT comparisons between species and to account for species offsets [Sodrian and Rosenthal, 2009]. Finally, to estimate $\delta^{18}O_{seawater}$, we used the published $\delta^{18}O_{benthic}$ record [Ruddiman et al., 1989], the Mg/Ca-derived BWT, and the $\delta^{18}O_{seawater}$ equation: $\delta^{18}O_{seawater} = \delta^{18}O_{benthic} + 0.27 - 0.25 * (16.9 - BWT)$ [O'Neil et al., 1969; Shackleton, 1974; Kim and O'Neil, 1997; Elderfield et al., 2010; 2012].

To investigate changes in ocean circulation across the MPT we use previously published $\delta^{13}C$ values of benthic foraminifera ($\delta^{13}C_{benthic}$) (Table 1) and calculated the percent Northern Component Water (%NCW) [Oppo and Fairbanks, 1987; Raymo et al., 1990; Flower et al., 2000; Venz and Hodell, 2002]. The %NCW is used to estimate the relative contribution of deep water formed in the N. Atlantic (i.e. North Atlantic Deep Water, NADW) at a particular site [Oppo and Fairbanks, 1987] and is calculated using the following equation:

$$\%NCW = \frac{\delta^{13}C_{SITE} - \delta^{13}C_{SCW}}{\delta^{13}C_{NCW} - \delta^{13}C_{SCW}} * 100$$

where the $\delta^{13}C_{SITE}$ is the $\delta^{13}C_{benthic}$ value at a given location of interest and $\delta^{13}C_{NCW}$ and $\delta^{13}C_{SCW}$ are the $\delta^{13}C_{benthic}$ values at the designated northern and southern (i.e. Antarctic Bottom Water) water mass end-member locations. Here, we use ODP Site 982 and ODP 1090 as a North Atlantic and South Atlantic end-member, respectively [Venz and Hodell, 2002].

4. Results and Discussion

4.1 Comparison of *Uvigerina* spp., *Cibicidoides wuellerstorfi*, and *Oridorsalis umbonatus* records at DSDP Site 607

The new Mg/Ca record derived from *Uvigerina* spp. at Site 607 shows a decreasing trend from 1500 to 500 kyrs (Figure 2). The *Uvigerina* spp. Mg/Ca record has a similar long-term trend and variability to the previously published *Cibicidoides wuellerstorfi*, and *Oridorsalis umbonatus* Mg/Ca records [Sosdian and Rosenthal, 2009]. Uncertainty in Mg/Ca values is based on the 95% confidence of Monte Carlo error propagation (1000 resampling, $\sigma = 0.128$ mmol/mol for *Uvigerina* spp., this study and $\sigma = \sim 0.1$ for *Cibicidoides wuellerstorfi*, and *Oridorsalis umbonatus* [Sosdian and Rosenthal, 2009]. Offsets in Mg/Ca values are likely related to vital effects.

Large changes in deep ocean chemistry occurred over the MPT [Raymo et al., 1997; Lisiecki, 2014] as evidenced from the CaCO_3 [Ruddiman et al., 1989], $\delta^{13}\text{C}_{\text{benthic}}$ [Ruddiman et al., 1989] and %NCW (Figure 2) at Site 607, changes which may have influenced *C. wuellerstorfi* and *O. umbonatus* Mg/Ca values [e.g. Yu and Broecker, 2010]. Specifically, after ~900 kyra step change in deep ocean chemistry is suggested by an increased amplitude of variability in CaCO_3 , $\delta^{13}\text{C}_{\text{benthic}}$, and %NCW, as well as a greater contribution of low $[\text{CO}_3^{2-}]$ SCW water at Site 607, particularly during glacial periods. However, given the similarity in the overall long-term Mg/Ca trends in *Uvigerina* spp., *C. wuellerstorfi* and *O. umbonatus*, if *Uvigerina* spp. Mg/Ca values primarily respond to changes in temperature [Elderfield et al., 2010; 2012], then much of the long-term trend in *C. wuellerstorfi* and *O. umbonatus* Mg/Ca values (Figure 2) may also be explained by temperature and not by long-term changes in deep

208 sea carbon chemistry (i.e. changes in $[\text{CO}_3^{2-}]$) during the 1500 to 500 kyr time interval at Site
 209 607.

210 Additionally, we performed an F test to determine whether the large amplitude changes in
 211 deep ocean chemistry after ~900 kyr also influenced the variability in Mg/Ca values of
 212 *O. umbonatus*, *C. wuellerstorfi* and *Uvigerina* spp.. We compared the Mg/Ca values between the
 213 600 – 800 ka and 1000 – 1250 ka, intervals of large and small amplitude changes in deep ocean
 214 chemistry, respectively, for each species and found no statistical difference Mg/Ca variability
 215 prior to or after ~900 kyr (Table 1). The lack of coherence and synchronicity in the long-term
 216 trend and variability between Mg/Ca and the deep ocean chemistry parameters suggests changes
 217 in $[\text{CO}_3^{2-}]$ do not dominate benthic foraminifera Mg/Ca values during the MPT at Site
 218 607 [Sosdian and Rosenthal, 2009; 2010].

219 The new BWT record derived from *Uvigerina* Mg/Ca values at Site 607 shows a
 220 decreasing trend in mean temperature of ~2-3°C from 1500 to 500 kyr (Figure 3), regardless of
 221 calibration choice, suggesting a long-term cooling of bottom waters occurred. In Figure 3,
 222 *O. umbonatus*, *C. wuellerstorfi* and *Uvigerina* spp. Mg/Ca values were converted to temperature
 223 using available low-temperature species-specific temperature calibrations [Lear et al., 2002;
 224 Healey et al., 2008; Yu and Elderfield, 2008; Sosdian and Rosenthal, 2009; Elderfield et al.,
 225 2012; Cappelli et al., 2015]. The magnitude of long-term temperature cooling between 1500 to
 226 500 kyr is dependent on calibration choice with *C. wuellerstorfi* broadly showing less cooling than
 227 *O. umbonatus* and *Uvigerina* spp.. These differences in temperature reconstructions are not related

to changes in the presence or absence of different species during interglacial and glacial periods as the species are generally found throughout the 1500 to 500 kyr interval(see Supplemental Figure 1 for *Uvigerina* spp. distribution). The previously published Mg/Ca records of *O. umbonatus* and *C. wuellerstorfi* [Sosdian and Rosenthal, 2009] suggest a $\sim 2^{\circ}\text{C}$ and $\sim 1\text{-}2^{\circ}\text{C}$ cooling trend from 1500 to 500 kyr respectively, depending on calibration choice, which is in broad agreement with the *Uvigerina* spp. record (Figure 3). Additionally, all species show a similar magnitude of cooling at the warm and cold range of temperatures which suggests there was a similar magnitude of cooling in the interglacial and glacial intervals over the MPT [Sosdian and Rosenthal, 2009]. Likewise, for the late Pleistocene, the previously published *O. umbonatus* and *C. wuellerstorfi* records are in good agreement with an ostracode Mg/Ca-BWT record [Dwyer et al., 1995], though there is also disagreement as to whether ostracode Mg/Ca values also respond to changes $\Delta[\text{CO}_3^{2-}]$ [Elmore et al., 2012; Farmer et al., 2012]. Regardless of species-specific calibration choice, the *Uvigerina* spp., *O. umbonatus* and *C. wuellerstorfi* records show a similarly decreasing BWT trend over the MPT.

Based on the above observations, we combine the *Uvigerina* spp., *C. wuellerstorfi* and *O. umbonatus* data to create a composite BWT record at Site 607 (Figure 4A). Mg/Ca values for each species were converted to temperature using the species-specific paleotemperature equations [Sosdian and Rosenthal, 2009; Elderfield et al., 2010; 2012] to account for species offsets and Mg/Ca-temperature sensitivities. Using the *Uvigerina* spp. BWT generated in this study and the previously published $\delta^{18}\text{O}_{\text{benthic}}$ record [Ruddiman et al., 1989], paired $\delta^{18}\text{O}_{\text{seawater}}$ values were calculated (blue dots, Figure 4B). We then combine our *Uvigerina* spp.-derived $\delta^{18}\text{O}_{\text{seawater}}$ values with the previously published *C. wuellerstorfi* and *O. umbonatus*-derived

250 $\delta^{18}\text{O}_{\text{seawater}}$ values [Sosdian and Rosenthal, 2009] to create a composite $\delta^{18}\text{O}_{\text{seawater}}$ record (Figure
 251 4). We note that during the early Pleistocene some warm BWT estimates result in reconstructed
 252 $\delta^{18}\text{O}_{\text{seawater}}$ values approaching +1.5‰, which seems unlikely considering the early Pleistocene is
 253 thought to have had less ice than the recent glacial period. Part of the issue may be that several
 254 species of benthic foraminifera are used in these reconstructions (e.g. the $\delta^{18}\text{O}_{\text{benthic}}$ is based on
 255 *C. wuellerstorfi* and the BWT is based on *Uvigerina* spp.) or that BWT and $\delta^{18}\text{O}_{\text{benthic}}$ records are
 256 not in phase (i.e. temperature leads ice volume, Sosdian and Rosenthal, [2009]). To account for
 257 the fact that these records are based on different species with different temperature and $\delta^{18}\text{O}$
 258 sensitivities, the composite BWT and $\delta^{18}\text{O}_{\text{seawater}}$ records were smoothed using a three point
 259 moving to reduce the error on the estimates (see Supplemental Text). By smoothing the records,
 260 we can focus on our primary goal of understanding the long-term trends in BWT and $\delta^{18}\text{O}_{\text{seawater}}$.
 261 The advantage of combining *Uvigerina* spp., *C. wuellerstorfi* and *O. umbonatus* analyses is that we
 262 are able to achieve a more continuous, high-resolution record because these species often alter in
 263 abundance with *Uvigerina* being generally more abundant during glacial periods.
 264 Interestingly, over the same period, the $\delta^{18}\text{O}_{\text{benthic}}$ record at Site 607 has a decreasing trend
 265 of 0.317‰ (Figure 5), which is equivalent to a $\sim 1.3^\circ\text{C}$ cooling using a sensitivity of 0.25‰ per
 266 $^\circ\text{C}$. A combination of possibilities could explain this trend: 1) the $\delta^{18}\text{O}_{\text{benthic}}$ record is dominated
 267 by BWT cooling and there was little ice volume increase over the MPT, 2) there was a change in
 268 ice sheet end member values and thus mean ocean $\delta^{18}\text{O}_{\text{seawater}}$ (e.g. [Winnick and Caves, 2015]),
 269 and/or 3) changes in ocean circulation may influence the $\delta^{18}\text{O}_{\text{seawater}}$ values at Site 607. Due to the
 270 variable constraints on $\delta^{18}\text{O}_{\text{seawater}}$ and the large error on the $\delta^{18}\text{O}_{\text{seawater}}$ estimate, we do not

intend to reconstruct sea level and instead focus on the long-term trends in BWT and $\delta^{18}\text{O}_{\text{seawater}}$ at Site 607.

4.2. Comparison of Site 607 and Site 1123 BWT records

The BWT, $\delta^{18}\text{O}_{\text{benthic}}$ and $\delta^{18}\text{O}_{\text{seawater}}$ records at Sites 607 and 1123 have dissimilar long-term trends from 1500 to 500 kyr (Figure 5). While Site 607 shows a subtle yet distinct long-term BWT cooling trend from 1500 to ~500 kyr (taking into account uncertainties discussed in previous section; Figure 4, 5), Site 1123 data show colder, near freezing temperatures characterizing the entire 1500 kyr record (Figure 5) [Elderfield et al., 2012]. The $\delta^{18}\text{O}_{\text{benthic}}$ records at Sites 607 and 1123 correlate well with the globally averaged benthic oxygen isotope LR04 stack [Lisiecki and Raymo, 2005] (Figure 5) except during the glacial periods between MIS 22 and MIS 18 when Site 1123 shows a pronounced ~0.3 to 0.5‰ increase during glacial periods relative to LR04. At MIS 16 the $\delta^{18}\text{O}_{\text{benthic}}$ records converge again.

Site 607 $\delta^{18}\text{O}_{\text{seawater}}$ values show no change in the long-term mean or amplitude from 1500 to 500 kyr (Figure 4, 5). On the other hand, at Site 1123, $\delta^{18}\text{O}_{\text{seawater}}$ values show an increase in the mean and the amplitude after MIS 22 (Figure 5). The site 1123 $\delta^{18}\text{O}_{\text{seawater}}$ record has been interpreted as showing a large build up of ice volume at MIS 22 [Elderfield et al., 2012], relative to the earlier interval, whereas the Site 607 $\delta^{18}\text{O}_{\text{seawater}}$ record shows no obvious, abrupt increase in ice volume occurred at MIS 22. If there was a large increase in glacial ice volume at MIS 22, relative to earlier intervals, as Elderfield et al. (2012) suggests, then it is difficult to reconcile the different $\delta^{18}\text{O}_{\text{seawater}}$ records at Site 1123 and 607 because a large increase in ice volume should be common to both records. This suggests that hydrographic changes may have occurred over

the MPT at one or both sites, changes that could have influenced the $\delta^{18}\text{O}_{\text{seawater}}$. In the next sections we explore possible hydrographic changes at Sites 1123 and 607.

4.3. Teasing apart local changes in ocean circulation and global climate

$\delta^{18}\text{O}_{\text{seawater}}$ values depend *globally* on the build-up/loss of ^{16}O on the continents on glacial-interglacial time scales and *locally* on water that flows over any abyssal location, integrating both global climate and regional bottom water properties imprinted by high latitude deep-water formation and ocean circulation. One implication of the Elderfield et al., [2012] record is that if the Site 1123 $\delta^{18}\text{O}_{\text{seawater}}$ is assumed to be the global $\delta^{18}\text{O}_{\text{seawater}}$ signal (as suggested by those authors), then the residual $\delta^{18}\text{O}_{\text{benthic}}$ temperature component in the LR04 stack as well as in the Sites 607, 1020, 1146, 1143, 846, 849 $\delta^{18}\text{O}_{\text{benthic}}$ records would suggest a warming of the deep Atlantic and Pacific water of nearly 2°C (e.g., $\sim 0.5\text{‰}$, assuming $0.25\text{‰}/^{\circ}\text{C}$) occurred during glacial MIS 22 (Figure 5). Indeed, if Elderfield et al.'s [2012] conclusions about global ice volume are correct, then these sites would seemingly reflect bottom water warming during glacial periods across the MPT, until MIS 16 when the $\delta^{18}\text{O}_{\text{benthic}}$ converge again (Figure 5). It is difficult to explain why large expanses of the deep Atlantic and Pacific oceans would be warming during glacial excursions across the MPT. Here we suggest that changes in ocean circulation may influence the magnitude of the abrupt change observed at Site 1123 and the record reflects both global and local changes. Numerous studies have documented changes in deep ocean circulation over the MPT using $\delta^{13}\text{C}$ of benthic foraminifera [Raymo et al., 1997; Venz et al., 1999; Venz and Hodell, 2002; Hodell et al., 2003; Ferretti et al., 2010; Poirier and Billups, 2014] and carbonate preservation [Schmieder et al., 2000; Sexton and Barker, 2012].

Here we examine how these circulation changes may have influence the $\delta^{18}\text{O}_{\text{benthic}}$ and Mg/Ca-derived $\delta^{18}\text{O}_{\text{seawater}}$ records from Site 1123 and 607.

4.3.1. Changes in ocean circulation from MIS 24 to 22

The $\delta^{18}\text{O}_{\text{benthic}}$ record at Site 1123 (and thus also the calculated $\delta^{18}\text{O}_{\text{seawater}}$ record) stands out as anomalous in comparison to the LR04 stack (Figure 5). Elderfield et al. [2012] argue that Site 1123 is representative of global conditions because a large volume of water flows over the Chatham Rise and into the Pacific (Figure 1) and LR04 is not representative of global conditions because it is biased to the Atlantic Ocean; however, Site 1123 $\delta^{18}\text{O}_{\text{benthic}}$ values are different from many of the records from the rest of the Pacific basin. In comparison to the previously published high-resolution mid-to-deep (i.e. >2000 m water depth, Table 2) $\delta^{18}\text{O}_{\text{benthic}}$ records from the South China Sea, California Margin, and equatorial Pacific (Figure 6), Site 1123 is ~0.4‰ heavier than most of the other Pacific records. Although Elderfield et al. [2012] point out Site 1123 shows good agreement with Site 677, Site 677 is located in the Panama Basin (sill depth 2300 to 2920 m [Lonsdale, 1977]) and is bathymetrically isolated from the rest of the deep eastern equatorial Pacific. The other Pacific records, including those from the deep eastern equatorial Pacific (e.g. Sites 846 and 849, water depth 3307 and 3849 m, respectively), show good agreement with LR04. The choice of benthic species used in generating the $\delta^{18}\text{O}_{\text{benthic}}$ records also does not appear to contribute to these discrepancies. Although many of these $\delta^{18}\text{O}_{\text{benthic}}$ records are based on *C. wuellerstorfi* or a mixture of *C. wuellerstorfi* and *Uvigerina* spp., as Elderfield et al. [2012] note, at Site 1123 during MIS 22, the abrupt shift in $\delta^{18}\text{O}_{\text{benthic}}$ is observed in both *C. wuellerstorfi* and *Uvigerina* spp. samples. The similarity between *C. wuellerstorfi* and *Uvigerina* spp. $\delta^{18}\text{O}_{\text{benthic}}$ samples appears to hold for the rest of the Pacific.

Consequently, we infer that the very positive $\delta^{18}\text{O}_{\text{benthic}}$ excursions at MIS 22 at Sites 677 and 1123 may reflect a local hydrographic effects, possibly due to the time-varying mixing of different water masses, that are not recorded in the rest of the Pacific basin (Figure 1).

In the Atlantic, the relative importance of NCW and SCW changed over the MPT [Raymo *et al.*, 1997; Venz and Hodell, 2002; Hodell *et al.*, 2003; Pena *et al.*, 2008; Lisiecki, 2014]. Significant changes in ocean circulation over the MPT may have created hydrographic conditions unique to Site 1123 that are not representative of deep Pacific Ocean or global conditions. This change could have had downstream consequences for the Pacific because a mixture of SCW and NCW flows into the deep Pacific [Hall *et al.*, 2001; Elderfield *et al.*, 2012]. We suggest that Site 1123 $\delta^{18}\text{O}_{\text{seawater}}$ record is likely not representative of mean ocean $\delta^{18}\text{O}_{\text{seawater}}$ because its location on the Chatham Rise appears to be sensitive to the mixing ratio of SCW and NCW (Figure 1).

In Figure 7, $\delta^{13}\text{C}_{\text{benthic}}$ and %NCW records show marked changes in glacial deep ocean circulation in the Atlantic across the MPT (for completeness, the Pacific basin can be found in Supplemental Figure 3). Prior to MIS 24, NCW filled the Atlantic above ~3500m during glacial periods [Raymo *et al.*, 1997; Venz and Hodell, 2002; Hodell *et al.*, 2003; Pena *et al.*, 2008; Lisiecki, 2014]. Beginning at MIS 24 and intensifying at MIS 22, Nd isotopes and $\delta^{13}\text{C}_{\text{benthic}}$ from intermediate and deep water horizons suggest bottom waters were predominantly ventilated by SCW [Raymo *et al.*, 1997; Venz and Hodell, 2002; Hodell *et al.*, 2003; Pena *et al.*, 2008; Lisiecki, 2014; Poirier and Billups, 2014]. These changes in Atlantic circulation may be related to an observed decrease in the mean flow speed over the Chatham Rise starting at MIS 22 (e.g. Site 1123; [Hall *et al.*, 2001; Venuti *et al.*, 2007]), a hydrographic change which may have contributed to the unique $\delta^{18}\text{O}_{\text{benthic}}$ signal recorded at Site 1123.

This change in Atlantic deep ocean circulation likely influenced the Site 607 $\delta^{18}\text{O}_{\text{benthic}}$ record as well as the records throughout the Atlantic. Elderfield et al. [2012] point out Site U1308 also shows an abrupt increase in $\delta^{18}\text{O}_{\text{benthic}}$ at MIS 22 which is similar to Site 1123 (Figure 8). However, this abrupt increase in $\delta^{18}\text{O}_{\text{benthic}}$ is absent in the Site 607 and LR04 records. Although Sites 607 and U1308 are both in the N. Atlantic, they experience different water masses because they are on different sides of the basin separated by the mid-Atlantic ridge. Site 607 is in the western Atlantic basin and ventilated by either NCW and SCW whereas U1308 is in the eastern Atlantic basin which is ventilated from the south by a mixture of NCW and SCW that flows over the Romanche Fracture Zone at the equator [Raymo et al., 1997]. The value of using a stacked record like LR04 is that it changes in hydrography are mostly averaged out. The LR04 stack, though biased toward the Atlantic, has at least 23 records that on average do not show an abrupt change in $\delta^{18}\text{O}_{\text{benthic}}$ at MIS 22. The LR04 stack combined with the observations in the Pacific this suggests that an abrupt, large magnitude change at MIS 22 is likely related to changes in ocean circulation ventilating that particular area rather than whole ocean changes in $\delta^{18}\text{O}$ related to ice volume.

4.3.2 Long-term cooling at Site 607

The cause of the long-term bottom water cooling found in at Site 607 and its relationship to the emergence of the 100 kyr cycle remains elusive. Long-term cooling that begins or intensifies at ~1.2 Ma is found in multiple mid- to high latitude sea surface temperature records [Lawrence et al., 2010; McClymont et al., 2013], suggesting that this long-term secular cooling trend is not limited to the deep ocean. Although this trend is observed in a number of surface locations in the N. Atlantic, the BWT cooling trend observed at Site 607 could also be due to cold,

southern component waters gradually penetrating further into the N. Atlantic over the MPT. $\delta^{13}\text{C}_{\text{benthic}}$ records from deep North Atlantic locations show gradually decreasing values starting at ~1.1 Ma [Raymo *et al.*, 1990; 1997; 2004; Ferretti *et al.*, 2005; Lisiecki, 2014; Poirier and Billups, 2014] consistent with an increasing contribution of an isotopically light, southern-sourced water mass. It is likely that the BWT cooling observed at Site 607 reflects both cooling in the deep ocean and **changing deep ocean circulation, such as** increasing influence of southern component waters [Sosdian and Rosenthal, 2009] **as shown in Figure 7.**

4.3.3 Long-term and Threshold Climate Change across the Mid-Pleistocene Transition

It is clear from the gradual BWT cooling at Site 607 as well as the abrupt changes at Site 1123 that the MPT exhibited both gradual and threshold-like responses to climate forcing. Gradually changing environmental conditions (**summarized by McClymont *et al.*, [2013]**), including N. Atlantic BWT and mid- to high-latitude SST cooling [this study, Sosdian and Rosenthal, 2009; Lawrence *et al.*, 2010; McClymont *et al.*, 2013], expanded sea-ice coverage [Kemp *et al.*, 2010], equatorward polar front migration [McClymont *et al.*, 2008; Lawrence *et al.*, 2010; Martinez-Garcia *et al.*, 2010], decreased deep ocean ventilation [Hall *et al.*, 2001; Venuti *et al.*, 2007], intensified ocean/atmospheric circulation [de Garidel-Thoron *et al.*, 2005; McClymont and Rosell-Mele, 2005], **and increased ice-sheet size** [Clark *et al.*, 2006] may have altered climate boundary conditions in a way that shifted the cryosphere across a threshold that permanently altered climate–ice sheet **response to orbital forcing**. Centered on this long-term trend, **the LR04 $\delta^{18}\text{O}_{\text{benthic}}$ stack shows a ~0.15‰ increase at MIS 22 (~900 ka**, Figure 5), likely related to ice volume growth, and a statistically significant emergence of the 100 kyr cycle [Maasch, 1988; Mudelsee and Stattegger, 1997; Clark *et al.*, 2006; Raymo *et al.*, 2006;

Lisiecki and Raymo, 2007; McClymont et al., 2013]. As southern Hemisphere summer insolation was anomalously low during the MIS 24 to 22 interval, this ice volume growth may have occurred in Antarctica (as suggested by [*Raymo et al., 2006*] and Elderfield et al. [2102]). However, we suggest the ice volume growth is not to the magnitude implied by record at Site 1123 ($\sim 0.3\text{‰}$ $\delta^{18}\text{O}_{\text{seawater}}$) due to possible local hydrographic effects as discussed above. Additional globally distributed, high-resolution $\delta^{18}\text{O}_{\text{seawater}}$ records are needed to fully deconvolve the influence of ice volume and circulation changes during the MIS 24 to MIS 22 transition.

As with the glacial cycles that characterize the late Pleistocene [*Shakun et al., 2015*], ice sheets may be largely responding to changes in global temperatures and atmospheric CO_2 on glacial-interglacial time scales and these ice sheets may have had a threshold response to global cooling during the MPT. For example, ice volume may have switched from a 41 kyr to a 100 kyr orbital signal because global SSTs underwent a similar transition during the MPT: SSTs are dominated by 41 kyr cycles during the early Pleistocene [*Herbert et al., 2010; McClymont et al., 2013*] and the 100 kyr cycle slowly emerges at ~ 1.2 Ma [*McClymont et al., 2013*] and fully dominates at ~ 900 ka [*Medina-Elizalde and Lea, 2005; McClymont et al., 2013*]. However, invoking a 41 kyr to 100 kyr transition in SST change as the mechanism for forcing the 41 kyr to 100 kyr transition in $\delta^{18}\text{O}_{\text{benthic}}$ records just moves the question of “why” to another proxy.

It is possible that thermohaline circulation reorganization [*Pena and Goldstein, 2014*] and the increased influence of the southern ocean in deep water circulation [*Raymo et al., 1997; 2004; Ferretti et al., 2005; Lisiecki, 2014; Poirier and Billups, 2014*] may have altered heat and salt transport and carbon cycling in the deep ocean. At ~ 900 ka, the BWT and $\delta^{18}\text{O}_{\text{seawater}}$ records at Site 607 and 1123 converge (Figure 5) suggesting enhanced connectivity between the Atlantic

and Pacific ocean basins. High-resolution atmospheric CO₂ estimates and records of deep ocean chemistry are necessary to fully understand the links between temperature, ocean circulation, CO₂, ice volume during the MPT.

5. Summary

We have reconstructed aMg/Ca-derived BWT record using *Uvigerina* spp. from DSDP Site 607 that shows an ~2-3°C cooling from ~1500 to 500 kyrs across the MPT. This gradual cooling trend is consistent with a previously published BWT record based on *C. wuellerstorfi* and *O. umbonatus* [Sosdian and Rosenthal, 2009] implying that Mg/Ca records based on epifaunal benthic species at Site 607 during the MPT are robust. We hypothesize that thermohaline reorganization starting at MIS 24 and intensifying at MIS 22 likely influenced the local hydrography at ODP Site 1123, consequently explaining the unusually large decrease in $\delta^{18}\text{O}_{\text{seawater}}$ at that site. Gradual N. Atlantic BWT and mid- to high-latitude sea surface temperature cooling likely pushed the climate system across a threshold starting at MIS 24. A modest increase in ice volume at ~900 ka, possibly in Antarctica, combined with thermohaline reorganization, created a new climate regime with enhanced sensitivity to the 100 kyr orbital cycle.

Acknowledgements

We would like to thank two anonymous reviewers that greatly improved the quality of the manuscript. Additionally, we would like to thank Stella Woodard and Blake Dyer for helpful discussions, Katherine Esswein for analytical support and Bilal Nadeem, Andrés Salazar and Jason Swann for laboratory assistance. Marine sediment samples were provided as part of the

International Ocean Discovery Program (IODP). Funding was provided by NSF grant OCE-1436014 (HLF and MER).

Table 1. F-test statistics comparing Mg/Ca values of *C. wuellerstorfi* and *O. umbonatus* and *Uvigerina* spp. between the 600 – 800 ka and 1000 – 1250 ka. No statistical difference in Mg/Ca variability is observed prior to or after MIS 22.

	F-test	Degrees of Freedom Numerator	Degrees of Freedom Denominator	P value
<i>C. wuellerstorfi</i>	1.5128	47	31	0.2251
<i>O. umbonatus</i>	0.7160	16	24	0.4948
<i>Uvigerina</i> spp.	1.0393	30	37	0.9027

461 **Table 2. Core Locations of Previously Published Benthic Stable Isotope Values.**

Site	Latitude	Longitude	Water Depth (m)	$\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ Reference	$\delta^{13}\text{C}$ Reference	$\delta^{18}\text{O}$ Reference
Atlantic						
GIK13519	5.7 N	19.9 W	2862	[Sarnthein <i>et al.</i> , 1994]		
GeoB1034	21.7 S	5.4 E	3731	Bickert and Wefer [1996]		
GeoB1211	24.5 S	7.5 E	4089	Bickert and Wefer [1996]		
DSDP 552	56 N	23.2 W	2301	[Shackleton and Hall, 1984]		
DSDP 607	41 N	33 W	3427	[Ruddiman <i>et al.</i> , 1989]		
ODP 664	0.1 N	23.2 W	3806	[Raymo <i>et al.</i> , 1997]		
ODP 925	4.2 N	43.5 W	3040	[Bickert <i>et al.</i> , 1997]		
ODP 926	3.7 N	42.9 W	3598	[Lisiecki <i>et al.</i> , 2008]		
ODP 927	5.5 N	44.5 W	3326	[Bickert <i>et al.</i> , 1997]		
ODP 928	5.5 N	43.7 W	4012	[Lisiecki <i>et al.</i> , 2008]		
ODP 929	6 N	43.7 W	4369	[Bickert <i>et al.</i> , 1997]		
ODP 980	55.5 N	14.7 W	2179	[Flower <i>et al.</i> , 2000]		
ODP 981	55.5 N	14.7 W	2173	[Raymo <i>et al.</i> , 2004]		
ODP 982	57.5 N	15.9 W	1146	[Venz <i>et al.</i> , 1999; Venz and Hodell, 2002]		
ODP 983	60.4 N	23.6 W	1983	[Raymo <i>et al.</i> , 2004]		
ODP 984	61.4 N	24.1 W	1650	[Raymo <i>et al.</i> , 2004]		
ODP 1063	33.7 N	57.6 W	4584	[Ferretti <i>et al.</i> , 2005; Poirier and Billups, 2014]		
ODP 1088	41.1 S	13.6 E	2081	[Hodell <i>et al.</i> , 2003]		
ODP 1089	40.9 S	9.9 E	4621	[Hodell <i>et al.</i> , 2001]		
ODP 1090	42.9 S	8.9 E	3699	[Venz and Hodell, 2002]		
ODP 1264	28.5 S	2.8 E	2505	[Bell <i>et al.</i> , 2014]		
ODP 1267	28.1 S	1.7 E	4355	[Bell <i>et al.</i> , 2014]		
IODP U1308	49.9 N	24.2 W	3871	[Hodell <i>et al.</i> , 2008]		
IODP U1313	41 N	33 W	3426	[Ferretti <i>et al.</i> , 2010]		
IODP U1314	56.4 N	27.9 W	2820	[Alonso-Garcia <i>et al.</i> , 2011]		
Pacific						
GeoB15016	27.5 S	71.1 W	956		[Martínez-Méndez <i>et al.</i> , 2013]	
PC72	0.1 N	139.4 W	4298			[Murray <i>et al.</i> , 2000]
RC13110	0.1 S	95.7 W	3231	[Mix <i>et al.</i> , 1991]		
ODP 677	1.2 N	83.7 W	3461	[Shackleton <i>et al.</i> , 1990]		
ODP 806	0.3 N	159.4 E	2520		[Bickert <i>et al.</i> , 1993]	
ODP 846	3.1 S	90.8 W	3307	[Mix <i>et al.</i> , 1995a]		
ODP 849	0.2 N	110.5 W	3849	[Mix <i>et al.</i> , 1995b]		
ODP 1020	41 N	126.4 W	3038			Liu, personal communication to L. Lisiecki [2002]
ODP 1123	41.8 S	171.5 W	3290	[Hall <i>et al.</i> , 2001; Harris, 2002; Elderfield <i>et al.</i> , 2012]		
ODP 1143	9.4 N	113.3 E	2772		[Cheng <i>et al.</i> , 2004]	[Tian <i>et al.</i> , 2002]
ODP 1146	19.5 N	116.3 E	2091	[Prell, 2003]		
ODP 1241	5.8 N	86.4 W	2027	[Lalicata and Lea, 2011]		

Figure Captions

Figure 1. Locations of sites: $\delta^{13}\text{C}_{\text{benthic}}$ from the Atlantic (squares, Table 1), $\delta^{18}\text{O}_{\text{benthic}}$ from the Pacific (circles, Table 2), and DSDP Site 607 and ODP Site 1123 with Mg/Ca-derived BWT records (stars) (A). Hydrographic transects for Site 607 (B) and Site 1123 (C). Atlantic transect of carbonate ion concentration $[\text{CO}_3^{2-}]$ [Key *et al.*, 2004]. Site 607 is in area sensitive to changes in $[\text{CO}_3^{2-}]$ on glacial-interglacial time scales. Site 1123 is located in an area sensitive to water mass mixing (panel modified from [Carter *et al.*, 1999]). Figure generated using Ocean Data View [Schlitzer] and [Troupin *et al.*, 2012].

Figure 2. DSDP Site 607 Mg/Ca values of *Uvigerina* spp. (blue, this study) and previously published *O. umbonatus* (pink) and *C. wuellerstorfi* (light green) [Sosdian and Rosenthal, 2009] (A). Deep ocean chemistry indicators %CaCO₃ (B), $\delta^{13}\text{C}_{\text{benthic}}$ (C) and %NCW (D).

Figure 3. Mg/Ca values converted to BWT using previously published species-specific temperature calibrations of L2002 [Lear *et al.*, 2002], SR2009 [Sosdian and Rosenthal, 2009], YE2008 [Yu and Elderfield, 2008], C2015 [Cappelli *et al.*, 2015], H2008 [Healey *et al.*, 2008] and E2012 [Elderfield *et al.*, 2012].

Figure 4. All species included for a composite BWT record at Site 607 using the species-specific temperature calibrations of [Sosdian and Rosenthal, 2009] and [Elderfield *et al.*, 2012]. (A). Previously published (grey) and new *Uvigerina* spp. (blue, this study) $\delta^{18}\text{O}_{\text{seawater}}$ values (B). Error bars on the left hand side indicate the 3 point smoothed combined error. Error for BWT

measurements is $\pm 1.4^{\circ}\text{C}$ and the propagated error for $\delta^{18}\text{O}_{\text{seawater}}$ estimates is $\pm 0.33\%$ (see Supplemental Text).

Figure 5. Comparison of LR04 (black), ODP Site 1123 (A, orange), DSDP Site 607 (B, blue), for $\delta^{18}\text{O}_{\text{benthic}}$, BWT (C), and $\delta^{18}\text{O}_{\text{seawater}}$ (D). Site 607 shows a long-term cooling trend over the MPT while Site 1123 has near freezing temperatures over length of the 1.5 Myr record.

Figure 6. Previously published, high-resolution, mid to deep ocean depth $\delta^{18}\text{O}_{\text{benthic}}$ records from the Pacific plotted in comparison to LR04. ODP Sites 1123 and 677 stand out as anomalous in comparison to most of the records from the Pacific Ocean basin and LR04.

Figure 7. Cross-section view of mean glacial $\delta^{13}\text{C}_{\text{benthic}}$ values and %NCW for MIS 38 and MIS 26 to MIS 16 for sites in the Atlantic Ocean basin. Figure generated using Ocean Data View [Schlitzer]. For completeness, the Pacific Basin is included in Supplemental Figure 2.

Figure 8. Previously published $\delta^{18}\text{O}_{\text{benthic}}$ records from Sites 607, U1308 and 1123 plotted in comparison to LR04. ODP Sites 1123 and U1308 stand out as anomalous in comparison to Site 607 and LR04.

509 **References**

- 510 Alonso-Garcia, M., F. J. Sierro, and J. A. Flores (2011), Arctic front shifts in the subpolar North
 511 Atlantic during the Mid-Pleistocene (800–400 ka) and their implications for ocean
 512 circulation, *Palaeogeogr Palaeoclimatol*, 311(3-4), 268–280, doi:10.1016/j.palaeo.2011.09.004.
- 513 Barker, S., M. Greaves, and H. Elderfield (2003), A study of cleaning procedures used for
 514 foraminiferal Mg/Ca paleothermometry, *Geochem. Geophys. Geosyst*, 4, 8407,
 515 doi:10.1029/2003GC000559.
- 516 Bates, S. L., M. Siddall, and C. Waelbroeck (2014), Hydrographic variations in deep ocean
 517 temperature over the mid-Pleistocene transition, *Quaternary Science Reviews*, 88(C), 147–
 518 158, doi:10.1016/j.quascirev.2014.01.020.
- 519 Bell, D. B., S. J. A. Jung, D. Kroon, L. J. Lourens, and D. A. Hodell (2014), Local and regional
 520 trends in Plio-Pleistocene $\delta^{18}\text{O}$ records from benthic foraminifera, *Geochem. Geophys.*
 521 *Geosyst*, 15(8), 3304–3321, doi:10.1002/2014GC005297.
- 522 Berger, W. H., and E. Jansen (1994), Mid-Pleistocene Climate Shift-The Nansen Connection,
 523 *The polar oceans and their role in shaping the global environment*, 295–311.
- 524 Bickert, T., W. B. Curry, and G. Wefer (1997), 16. Late Pliocene to Holocene (2.6-0 Ma)
 525 western equatorial Atlantic deep-water circulation: Inferences from benthic stable isotopes,
 526 *Proceedings of the Ocean Drilling Program, Scientific Results*, 154, 239–253.
- 527 Bickert, T., W. H. Berger, S. Burke, H. Schmidt, and G. Wefer (1993), 24. Late Quaternary
 528 Stable Isotope Record of Benthic Foraminifers: Sites 805 and 806, Ontong Java Plateau,
 529 *Proceedings of the Ocean Drilling Program, Scientific Results, Vol. 130*, 130, 411–420.
- 530 Billups, K., and D. P. Schrag (2002), Paleotemperatures and ice volume of the past 27 Myr
 531 revisited with paired Mg/Ca and $^{18}\text{O}/^{16}\text{O}$ measurements on benthic foraminifera,
 532 *Paleoceanography*, 17(1), 3–13–11, doi:10.1029/2000PA000567.
- 533 Cappelli, E. L. G., M. Regenberg, A. Holbourn, W. Kuhnt, D. Garbe-Schönberg, and N.
 534 Andersen (2015), Refining C. wuellerstorfi and H. elegans Mg/Ca temperature calibrations,
 535 *Marine Micropaleontology*, 121(C), 70–84, doi:10.1016/j.marmicro.2015.10.001.
- 536 Carter, R. M., I. N. McCave, C. Richter, and L. Carter (1999), Leg 181 Summary: Southwest
 537 Pacific Paleoceanography, *Proceedings of the Ocean Drilling Program, Initial Reports, Vol.*
 538 *130, 181*, 1–80.
- 539 Cheng, X., J. Tian, and P. Wang (2004), Data Report: Stable Isotopes from Site 1143, edited by
 540 W. Prell, P. Wang, D. K. Rea, and S. Clemens, *Proceedings of the Ocean Drilling Program,*
 541 *Scientific Results, Volume*, 1–8, doi:doi:10.2973/odp.proc.sr.184.221.2004.
- 542 Clark, P. U., D. Archer, D. Pollard, J. D. Blum, J. A. Rial, V. Brovkin, A. C. Mix, N. G. Pisias,
 543 and M. Roy (2006), The middle Pleistocene transition: characteristics, mechanisms, and
 544 implications for long-term changes in atmospheric pCO_2 , *Quaternary Science Reviews*,

- 545 25(23-24), 3150–3184, doi:10.1016/j.quascirev.2006.07.008.
- 546 de Garidel-Thoron, T., Y. Rosenthal, F. Bassinot, and L. Beaufort (2005), Stable sea surface
547 temperatures in the western Pacific warm pool over the past 1.75 million years, *Nature*,
548 433(7023), 294–298.
- 549 Dwyer, G. S., T. M. Cronin, P. A. Baker, M. E. Raymo, J. Buzas, and T. Corrège (1995), Late
550 Pliocene and Quaternary bottom water temperature change in the deep North Atlantic,
551 *Science*, 270, 1347–1351.
- 552 Dyez, K. A., and A. C. Ravelo (2014), Dynamical changes in the tropical Pacific warm pool and
553 zonal SST gradient during the Pleistocene, *Geophysical Research Letters*, 41(21), 7626–
554 7633, doi:10.1002/2014GL061639.
- 555 Elderfield, H., J. Yu, P. Anand, T. Kiefer, and B. Nyland (2006), Calibrations for benthic
556 foraminiferal Mg/Ca paleothermometry and the carbonate ion hypothesis, *Earth and*
557 *Planetary Science Letters*, 250(3), 633–649.
- 558 Elderfield, H., M. Greaves, S. Barker, I. R. Hall, A. Tripathi, P. Ferretti, S. Crowhurst, L. Booth,
559 and C. Daunt (2010), A record of bottom water temperature and seawater $\delta^{18}\text{O}$ for the
560 Southern Ocean over the past 440kyr based on Mg/Ca of benthic foraminiferal *Uvigerina*
561 spp, *Quaternary Science Reviews*, 29(1), 160–169.
- 562 Elderfield, H., P. Ferretti, M. Greaves, S. Crowhurst, I. N. McCave, D. Hodell, and A. M.
563 Piotrowski (2012), Evolution of Ocean Temperature and Ice Volume Through the Mid-
564 Pleistocene Climate Transition, *Science*, 337(6095), 704–709, doi:10.1126/science.1221294.
- 565 Elmore, A. C., S. Sosdian, Y. Rosenthal, and J. D. Wright (2012), A global evaluation of
566 temperature and carbonate ion control on Mg/Ca ratios of ostracoda genus *Krithe*, *Geochem.*
567 *Geophys. Geosyst.*, 13(9), n/a–n/a, doi:10.1029/2012GC004073.
- 568 Farmer, J. R., T. M. Cronin, and G. S. Dwyer (2012), Ostracode Mg/Ca paleothermometry in the
569 North Atlantic and Arctic oceans: Evaluation of a carbonate ion effect, *Paleoceanography*,
570 27(2), PA2212–14, doi:10.1029/2012PA002305.
- 571 Ferretti, P., N. J. Shackleton, D. Rio, and M. A. Hall (2005), Early-middle Pleistocene deep
572 circulation in the western subtropical Atlantic: southern hemisphere modulation of the North
573 Atlantic Ocean, *Geological Society, London, Special Publications*, 247(1), 131–145.
- 574 Ferretti, P., S. J. Crowhurst, M. A. Hall, and I. Cacho (2010), North Atlantic millennial-scale
575 climate variability 910 to 790 ka and the role of the equatorial insolation forcing, *Earth and*
576 *Planetary Science Letters*, 293(1-2), 28–41, doi:10.1016/j.epsl.2010.02.016.
- 577 Flower, B. P., D. W. Oppo, J. F. McManus, K. A. Venz, D. A. Hodell, and J. L. Cullen (2000),
578 North Atlantic intermediate to deep water circulation and chemical stratification during the
579 past 1 Myr, *Paleoceanography*, 15(4), 388–403.
- 580 Hall, I. R., I. N. McCave, N. J. Shackleton, G. P. Weedon, and S. E. Harris (2001), Intensified

- 581 deep Pacific inflow and ventilation in Pleistocene glacial times, *Nature*, 412(6849), 809–812.
- 582 Harris, S. (2002), Late Pliocene-Pleistocene Carbon and Oxygen Stable Isotopes from Benthic
583 Foraminifers at Ocean Drilling Program Site 1123 in the Southwest Pacific, in *Proceedings*
584 *of the Ocean Drilling Program, Scientific Results Volume 181*, vol. 181, edited by C. Richter,
585 pp. 1–20, Proceedings of the Ocean Drilling Program, Scientific Results Volume 181.
- 586 Hays, J. D., J. Imbrie, and N. J. Shackleton (1976), Variations in the Earth's orbit: Pacemaker of
587 the ice ages,
- 588 Healey, S. L., R. C. Thunell, and B. H. Corliss (2008), The Mg/Ca-temperature relationship of
589 benthic foraminiferal calcite: New core-top calibrations in the <4 °C temperature range,
590 *Earth and Planetary Science Letters*, 272(3–4), 523–530, doi:10.1016/j.epsl.2008.05.023.
- 591 Herbert, T. D., L. C. Peterson, K. T. Lawrence, and Z. Liu (2010), Tropical Ocean Temperatures
592 Over the Past 3.5 Million Years, *Science*, 328(5985), 1530–1534,
593 doi:10.1126/science.1185435.
- 594 Hodell, D. A., C. D. Charles, and F. J. Sierro (2001), Late Pleistocene evolution of the ocean's
595 carbonate system, *Earth and Planetary Science Letters*, 192(2), 109–124.
- 596 Hodell, D. A., J. E. T. Channell, J. H. Curtis, O. E. Romero, and U. Rohl (2008), Onset of
597 “Hudson Strait” Heinrich events in the eastern North Atlantic at the end of the middle
598 Pleistocene transition (>640 ka)? *Paleoceanography*, 23(4), n/a–n/a,
599 doi:10.1029/2008PA001591.
- 600 Hodell, D. A., K. A. Venz, C. D. Charles, and U. S. Ninnemann (2003), Pleistocene vertical
601 carbon isotope and carbonate gradients in the South Atlantic sector of the Southern Ocean,
602 *Geochem. Geophys. Geosyst.*, 4(1), 1–19, doi:10.1029/2002GC000367.
- 603 Imbrie, J., A. Berger, E. A. Boyle, S. C. Clemens, A. Duffy, W. R. Howard, G. Kukla, J.
604 Kutzbach, D. G. Martinson, and A. McIntyre (1993), On the structure and origin of major
605 glaciation cycles 2. The 100,000-year cycle, *Paleoceanography*, 8(6), 699–735.
- 606 Imbrie, J., E. A. Boyle, S. C. Clemens, A. Duffy, W. R. Howard, G. Kukla, J. Kutzbach, D. G.
607 Martinson, A. McIntyre, and A. C. Mix (1992), On the structure and origin of major
608 glaciation cycles 1. Linear responses to Milankovitch forcing, *Paleoceanography*, 7(6), 701–
609 738.
- 610 Kemp, A. E. S., I. Grigorov, R. B. Pearce, and A. C. N. Garabato (2010), Migration of the
611 Antarctic Polar Front through the mid-Pleistocene transition: evidence and climatic
612 implications, *Quaternary Science Reviews*, 29(17–18), 1993–2009,
613 doi:10.1016/j.quascirev.2010.04.027.
- 614 Key, R. M., A. Kozyr, C. L. Sabine, K. Lee, R. Wanninkhof, J. L. Bullister, R. A. Feely, F. J.
615 Millero, C. Mordy, and T. H. Peng (2004), A global ocean carbon climatology: Results from
616 Global Data Analysis Project (GLODAP), *Global Biogeochem. Cycles*, 18(4), 1–23,
617 doi:10.1029/2004GB002247.

- Kim, S., and J. O'Neil (1997), Equilibrium and nonequilibrium oxygen isotope effects in synthetic carbonates, *Geochimica et Cosmochimica Acta*.
- Labeyrie, L. D., J. C. Duplessy, and P. L. Blanc (1987), Variations in mode of formation and temperature of oceanic deep waters over the past 125,000 years, *Nature*, 327(6122), 477–482, doi:10.1038/327477a0.
- Lalicata, J. J., and D. W. Lea (2011), Pleistocene carbonate dissolution fluctuations in the eastern equatorial Pacific on glacial timescales: Evidence from ODP Hole 1241, *Marine Micropaleontology*, 79(1-2), 41–51, doi:10.1016/j.marmicro.2011.01.002.
- Lawrence, K., S. Sosdian, H. White, and Y. Rosenthal (2010), North Atlantic climate evolution through the Plio-Pleistocene climate transitions, *Earth and Planetary Science Letters*, 300(3-4), 329–342, doi:10.1016/j.epsl.2010.10.013.
- Lear, C. H., H. Elderfield, and P. Wilson (2000), Cenozoic Deep-Sea Temperatures and Global Ice Volumes from Mg/Ca in Benthic Foraminiferal Calcite, *Science*, 287(5451), 269–272, doi:10.1126/science.287.5451.269.
- Lear, C. H., Y. Rosenthal, and N. Slowey (2002), Benthic foraminiferal Mg/Ca-paleothermometry: A revised core-top calibration, *Geochimica et Cosmochimica Acta*, 66(19), 3375–3387.
- Lisiecki, L. E. (2014), Atlantic overturning responses to obliquity and precession over the last 3 Myr, *Paleoceanography*.
- Lisiecki, L. E., and M. E. Raymo (2005), A Pliocene-Pleistocene stack of 57 globally distributed benthic delta O-18 records, *Paleoceanography*, 20(1), PA1003, doi:10.1029/2004PA001071.
- Lisiecki, L. E., and M. E. Raymo (2007), Plio–Pleistocene climate evolution: trends and transitions in glacial cycle dynamics, *Quaternary Science Reviews*, 26(1-2), 56–69, doi:10.1016/j.quascirev.2006.09.005.
- Lisiecki, L. E., M. E. Raymo, and W. B. Curry (2008), Atlantic overturning responses to Late Pleistocene climate forcings, *Nature*, 456(7218), 85–88, doi:10.1038/nature07425.
- Liu, Z., M. Altabet, and T. Herbert (2005), Glacial-interglacial modulation of eastern tropical North Pacific denitrification over the last 1.8-Myr, *Geophysical Research Letters*, 32(23), L23607, doi:10.1029/2005GL024439.
- Lonsdale, P. (1977), Inflow of bottom water to the Panama Basin, *Deep-Sea Research*, 24(12), 1065–1101.
- Maasch, K. A. (1988), Statistical detection of the mid-Pleistocene transition, *Clim Dynam*, 2(3), 133–143.
- Martin, P. A., and D. W. Lea (2002), A simple evaluation of cleaning procedures on fossil benthic foraminiferal Mg/Ca, *Geochem. Geophys. Geosyst*, 3(10), 8,

653 doi:10.1029/2001GC000280.

654 Martinez-Garcia, A., A. Rosell-Mele, E. L. McClymont, R. Gersonde, and G. H. Haug (2010),
 655 Subpolar Link to the Emergence of the Modern Equatorial Pacific Cold Tongue, *Science*,
 656 328(5985), 1550–1553, doi:10.1126/science.1184480.

657 Martínez-Méndez, G., D. Hebbeln, M. Mohtadi, F. Lamy, R. De Pol-Holz, D. Reyes-Macaya,
 658 and T. Freudenthal (2013), Changes in the advection of Antarctic Intermediate Water to the
 659 northern Chilean coast during the last 970 kyr, *Paleoceanography*, 28(4), 607–618,
 660 doi:10.1002/palo.20047.

661 McClymont, E. L., A. Rosell-Mele, G. H. Haug, and J. M. Lloyd (2008), Expansion of subarctic
 662 water masses in the North Atlantic and Pacific oceans and implications for mid-Pleistocene
 663 ice sheet growth, *Paleoceanography*, 23(4), n–a–n–a, doi:10.1029/2008PA001622.

664 McClymont, E. L., and A. Rosell-Mele (2005), Links between the onset of modern Walker
 665 circulation and the mid-Pleistocene climate transition, *Geol*, 33(5), 389–392,
 666 doi:10.1130/G21292.1.

667 McClymont, E. L., S. M. Sosdian, A. Rosell-Mele, and Y. Rosenthal (2013), Pleistocene sea-
 668 surface temperature evolution: Early cooling, delayed glacial intensification, and
 669 implications for the mid-Pleistocene climate transition, *Earth Science Reviews*, 123(C), 173–
 670 193, doi:10.1016/j.earscirev.2013.04.006.

671 Medina-Elizalde, M., and D. W. Lea (2005), The Mid-Pleistocene Transition in the Tropical
 672 Pacific, *Science*, 310(5750), 1009–1012, doi:10.1126/science.1115933.

673 Mix, A. C., J. Le, and N. J. Shackleton (1995a), 43. Benthic foraminiferal stable isotope
 674 stratigraphy of site 846: 0-1.8 Ma, edited by N. Pisias, L. Mayer, T. Janecek, A. Palmer-
 675 Julson, and T. van Andel, *Proceedings of the Ocean Drilling Program, Scientific Results*,
 676 Vol. 138, 138, 839–854.

677 Mix, A. C., N. G. Pisias, R. Zahn, and W. Rugh (1991), Carbon 13 in Pacific Deep and
 678 Intermediate Waters, 0-370 ka: Implications for Ocean Circulation and Pleistocene CO₂,
 679 *Paleoceanography*, 6(2), 205–226.

680 Mix, A., N. Pisias, W. Rugh, J. Wilson, A. Morey, and T. Hagelberg (1995b), 17. Benthic
 681 Foraminifer Stable Isotope Record From Site 849 (0-5Ma): Local and Global Climate
 682 Changes, *Proceedings of the Ocean Drilling Program, Scientific Results*, 138, 371–412.

683 Mudelsee, M., and K. Stattegger (1997), Exploring the structure of the mid-Pleistocene
 684 revolution with advanced methods of time-series analysis, *Geologische Rundschau*, 86(2),
 685 499–511.

686 Mudelsee, M., and M. Schulz (1997), The Mid-Pleistocene climate transition: onset of 100 ka
 687 cycle lags ice volume build-up by 280 ka, *Earth and Planetary Science Letters*, 151(1), 117–
 688 123.

- 689 Murray, R. W., C. Knowlton, M. Leinen, A. C. Mix, and C. H. Polsky (2000), Export production
690 and carbonate dissolution in the central equatorial Pacific Ocean over the past 1 Myr,
691 *Paleoceanography*, 15(6), 570–592.
- 692 O'Neil, J. R., R. N. Clayton, and T. K. Mayeda (1969), Oxygen isotope fractionation in divalent
693 metal carbonates, *The Journal of Chemical Physics*, 51, 5547.
- 694 Oppo, D. W., and R. G. Fairbanks (1987), Variability in the deep and intermediate water
695 circulation of the Atlantic Ocean during the past 25,000 years: Northern Hemisphere
696 modulation of the Southern Ocean, *Earth and Planetary Science Letters*, 86(1), 1–15.
- 697 Pena, L. D., and S. L. Goldstein (2014), Thermohaline circulation crisis and impacts during the
698 mid-Pleistocene transition, *Science*, 345(6194), 318–322, doi:10.1126/science.1249770.
- 699 Pena, L. D., I. Cacho, P. Ferretti, and M. A. Hall (2008), El Nino-Southern Oscillation-like
700 variability during glacial terminations and interlatitudinal teleconnections,
701 *Paleoceanography*, 23(3), PA3101, doi:10.1029/2008PA001620.
- 702 Pisias, N. G., and T. C. Moore Jr (1981), The evolution of Pleistocene climate: a time series
703 approach, *Earth and Planetary Science Letters*, 52(2), 450–458.
- 704 Poirier, R. K., and K. Billups (2014), The intensification of northern component deepwater
705 formation during the mid-Pleistocene climate transition, *Paleoceanography*, 29(11), 1046–
706 1061, doi:10.1002/2014PA002661.
- 707 Prell, S. C. C. A. W. L. (2003), Data Report: Oxygen and Carbon Isotopes from Site 1146,
708 Northern South China Sea, edited by W. Prell, P. Wang, D. K. Rea, and S. C. Clemens,
709 *Proceedings of the Ocean Drilling Program, Scientific Results, Volume*, 184, 1–8.
- 710 Raymo, M. E., and K. H. Nisancioglu (2003), The 41 kyr world: Milankovitch's other unsolved
711 mystery, *Paleoceanography*, 18(1), 1011, doi:10.1029/2002PA000791.
- 712 Raymo, M. E., D. W. Oppo, and W. Curry (1997), The mid-Pleistocene climate transition: A
713 deep sea carbon isotopic perspective, *Paleoceanography*, 12(4), 546–559.
- 714 Raymo, M. E., D. W. Oppo, B. P. Flower, D. A. Hodell, J. F. McManus, K. A. Venz, K. F.
715 Kleiven, and K. McIntyre (2004), Stability of North Atlantic water masses in face of
716 pronounced climate variability during the Pleistocene, *Paleoceanography*, 19(2).
- 717 Raymo, M. E., L. E. Lisiecki, and K. H. Nisancioglu (2006), Plio-Pleistocene ice volume,
718 Antarctic climate, and the global delta O-18 record, *Science*, 313(5786), 492–495,
719 doi:10.1126/science.1123296.
- 720 Raymo, M. E., W. F. Ruddiman, N. J. Shackleton, and D. W. Oppo (1990), Evolution of
721 Atlantic-Pacific $\delta^{13}\text{C}$ gradients over the last 2.5 my, *Earth and Planetary Science Letters*,
722 97(3), 353–368.
- 723 Rosenthal, Y. et al. (2004), Interlaboratory comparison study of Mg/Ca and Sr/Ca measurements

- 724 in planktonic foraminifera for paleoceanographic research, *Geochem. Geophys. Geosyst.* 5,
725 Q04D09, doi:10.1029/2003GC000650.
- 726 Rosenthal, Y., M. P. Field, and R. M. Sherrell (1999), Precise Determination of
727 Element/Calcium Ratios in Calcareous Samples Using Sector Field Inductively Coupled
728 Plasma Mass Spectrometry, *Anal. Chem.*, 71(15), 3248–3253, doi:10.1021/ac981410x.
- 729 Ruddiman, W., M. E. Raymo, D. G. Martinson, B. M. Clement, and J. Backman (1989),
730 Pleistocene evolution: Northern hemisphere ice sheets and North Atlantic Ocean,
731 *Paleoceanography*, 4(4), 353–412.
- 732 Rutherford, S., and S. D'Hondt (2000), Early onset and tropical forcing of 100,000-year
733 Pleistocene glacial cycles, *Nature*, 408(6808), 72–75.
- 734 Saltzman, B., and K. A. Maasch (1991), A first-order global model of late Cenozoic climatic
735 change II. Further analysis based on a simplification of CO₂ dynamics, *Clim Dynam.*, 5(4),
736 201–210.
- 737 Sarinthein, M., K. Winn, S. J. Jung, J.-C. Duplessy, L. Labeyrie, H. Erlenkeuser, and G. Ganssen
738 (1994), Changes in east Atlantic deepwater circulation over the last 30,000 years: Eight time
739 slice reconstructions, *Paleoceanography*, 9(2), 209–267.
- 740 Schlitzer, R., Ocean Data View,
- 741 Schmieder, F., T. von Dobeneck, and U. Bleil (2000), The Mid-Pleistocene climate transition as
742 documented in the deep South Atlantic Ocean: initiation, interim state and terminal event,
743 *Earth and Planetary Science Letters*, 179(3), 539–549.
- 744 Sexton, P. F., and S. Barker (2012), Onset of “Pacific-style” deep-sea sedimentary carbonate
745 cycles at the mid-Pleistocene transition, *Earth and Planetary Science Letters*, 321–322(C),
746 81–94, doi:10.1016/j.epsl.2011.12.043.
- 747 Shackleton, N. (1967), Oxygen isotope analyses and Pleistocene temperatures re-assessed,
748 *Nature*, 215(5096), 15–17, doi:10.1038/215015a0.
- 749 Shackleton, N. J. (1974), Attainment of isotopic equilibrium between ocean water and the
750 benthonic foraminifera genus *Uvigerina*: Isotopic changes in the ocean during the last glacial,
751 *Colloques Internationaux du C.N.R.S.*, 203–209.
- 752 Shackleton, N. J., A. Berger, and W. R. Peltier (1990), An alternative astronomical calibration of
753 the lower Pleistocene timescale based on ODP Site 677,, 81(04), 251–261,
754 doi:10.1017/S0263593300020782.
- 755 Shackleton, N. J., and M. A. Hall (1984), Oxygen and carbon isotope stratigraphy of Deep-Sea
756 Drilling Project Hole-552A-Plio-Pleistocene glacial history, *Initial Reports of the Deep Sea*
757 *Drilling Project*, 81(DEC), 599–609.
- 758 Shackleton, N. J., and N. D. Opdyke (1976), Oxygen-isotope and paleomagnetic stratigraphy of

- 759 Pacific core V28-239 late Pliocene to latest Pleistocene, *Geological Society of America*
 760 *Memoir*, 145, 449–464.
- 761 Shakun, J. D., D. W. Lea, L. E. Lisiecki, and M. E. Raymo (2015), An 800-kyr record of global
 762 surface ocean $\delta^{18}\text{O}$ and implications for ice volume-temperature coupling, *Earth and*
 763 *Planetary Science Letters*, 426(C), 58–68, doi:10.1016/j.epsl.2015.05.042.
- 764 Sosdian, S., and Y. Rosenthal (2009), Deep-Sea Temperature and Ice Volume Changes Across
 765 the Pliocene-Pleistocene Climate Transitions, *Science*, 325(5938), 306–310,
 766 doi:10.1126/science.1169938.
- 767 Sosdian, S., and Y. Rosenthal (2010), Response to Comment on “Deep-sea temperature and ice
 768 volume changes across the Pliocene-Pleistocene climate transitions,” *Science*, 328(5985),
 769 1480–1480.
- 770 Tian, J., P. Wang, X. Cheng, and Q. Li (2002), Astronomically tuned Plio–Pleistocene benthic δ
 771 ^{18}O record from South China Sea and Atlantic–Pacific comparison, *Earth and Planetary*
 772 *Science Letters*, 203, 1015–1029.
- 773 Troupin, C. et al. (2012), Generation of analysis and consistent error fields using the Data
 774 Interpolating Variational Analysis (DIVA), *Ocean Modelling*, 52-53, 90–101,
 775 doi:10.1016/j.ocemod.2012.05.002.
- 776 Venuti, A., F. Florindo, E. Michel, and I. R. Hall (2007), Magnetic proxy for the deep (Pacific)
 777 western boundary current variability across the mid-Pleistocene climate transition, *Earth and*
 778 *Planetary Science Letters*, 259(1-2), 107–118, doi:10.1016/j.epsl.2007.04.032.
- 779 Venz, K. A., and D. A. Hodell (2002), New evidence for changes in Plio–Pleistocene deep water
 780 circulation from Southern Ocean ODP Leg 177 Site 1090, *Palaeogeogr Palaeocl*, 182(3),
 781 197–220.
- 782 Venz, K. A., D. A. Hodell, C. Stanton, and D. A. Warnke (1999), A 1.0 Myr record of Glacial
 783 North Atlantic Intermediate Water variability from ODP site 982 in the northeast Atlantic,
 784 *Paleoceanography*, 14(1), 42–52.
- 785 Winnick, M. J., and J. K. Caves (2015), Oxygen isotope mass-balance constraints on Pliocene
 786 sea level and East Antarctic Ice Sheet stability, *Geol*, 43(10), 879–882,
 787 doi:10.1130/G36999.1.
- 788 Yu, J., and H. Elderfield (2007), Benthic foraminiferal B/Ca ratios reflect deep water carbonate
 789 saturation state, *Earth and Planetary Science Letters*, 258(1), 73–86.
- 790 Yu, J., and H. Elderfield (2008), Mg/Ca in the benthic foraminifera *Cibicidoides wuellerstorfi*
 791 and *Cibicidoides mundulus*: Temperature versus carbonate ion saturation, *Earth and*
 792 *Planetary Science Letters*, 276, 129–139, doi:10.1016/j.epsl.2008.09.015.
- 793 Yu, J., and W. S. Broecker (2010), Comment on “Deep-Sea Temperature and Ice Volume
 794 Changes Across the Pliocene-Pleistocene Climate Transitions,” *Science*, 328(5985), 1480–

795 1480, doi:10.1126/science.1186544.

796

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8