Geochemical Response of the Mid-Depth Northeast Atlantic Ocean to Freshwater Input During Heinrich Events 1 to 4

Anya J. Crocker^{1,2*}, Thomas B. Chalk^{1,3}, Ian Bailey^{1,4}, Megan R. Spencer¹, Marcus Gutjahr^{1,5}, Gavin L. Foster¹, Paul A. Wilson¹

¹National Oceanography Centre Southampton, University of Southampton, Waterfront Campus, Southampton SO14 3ZH, UK

²Department of Animal and Plant Science, University of Sheffield, Western Bank, Sheffield, S10 2TN, UK

³Department of Physical Oceanography, Woods Hole Oceanographic Institution, Woods Hole, Massachusetts, USA

⁴Now at: Camborne School of Mines & Environment and Sustainability Institute, College of Engineering, Mathematics & Physical Sciences, University of Exeter, Penryn Campus, Treliever Road, Penryn Cornwall, TR10 9FE, UK

⁵Now at: GEOMAR Helmholtz Centre for Ocean Research Kiel, Wischofstraße 1 – 3, 24148 Kiel, Germany

*Corresponding author

Correspondence to: Anya J. Crocker, anya.crocker@noc.soton.ac.uk

- 1 **Heinrich events are intervals of rapid iceberg-sourced freshwater release to the high**
- 2 **latitude North Atlantic Ocean that punctuate late Pleistocene glacials. Delivery of**
- 3 **fresh water to the main North Atlantic sites of deep water formation during**
- 4 **Heinrich events may result in major disruption to the Atlantic Meridional**

Overturning Circulation (AMOC), however, the simple concept of an AMOC shutdown in response to each freshwater input has recently been shown to be overly simplistic. Here we present a new multi-proxy dataset spanning the last 40,000 years that resolves four Heinrich events at a classic mid-depth North Atlantic drill site, employing four independent geochemical tracers of water mass properties: boron/calcium, carbon and oxygen isotopes in foraminiferal calcite and neodymium isotopes in multiple substrates. We also report rare earth element distributions to investigate the fidelity by which neodymium isotopes record changes in water mass distribution in the northeast Atlantic. Our data reveal distinct geochemical signatures for each Heinrich event, suggesting that the sites of fresh water delivery and/or rates of input played at least as important a role as the stage of the glacial cycle in which the fresh water was released. At no time during the last 40 kyr was the mid-depth northeast North Atlantic dominantly ventilated by southern-sourced water. Instead, we document persistent ventilation by Glacial North Atlantic Intermediate Water (GNAIW), albeit with variable properties signifying changes in supply from multiple contributing northern sources.

Keywords: Heinrich events; North Atlantic palaeoceanography; Last glacial period; Neodymium isotopes; B/Ca; Carbon and oxygen isotopes; Ice-rafted debris; ODP Site 980.

1. Introduction

The climate of the last glacial [period](#) was punctuated by a number of pronounced events with near global impacts, known as Heinrich (or H-) events (e.g. Heinrich, 1988; Hemming, 2004). During these events, transient catastrophic collapses of the North American Laurentide Ice Sheet (LIS), centered over the Canadian Hudson Bay, produced armadas of icebergs that [travelled through the Hudson Strait into the North Atlantic Ocean, eventually depositing](#) large volumes of ice-rafted debris (IRD), including distinctive detrital limestone clasts, in a belt across the North Atlantic (Ruddiman, 1977). LIS surging, together with iceberg calving from other circum-North Atlantic and Arctic ice sheets during H-events added large volumes of fresh (low density) water to the (sub)polar oceans with [important](#) implications for northern-sourced deep water formation and the Atlantic Meridional Overturning Circulation (AMOC). The AMOC exerts a strong control on regional and global climate on orbital to suborbital timescales, influencing latitudinal heat budgets (e.g. Boyle and Keigwin, 1987; Broecker and Denton, 1989) and partitioning of carbon between the atmosphere and deep ocean (e.g. Adkins, 2013; Sigman and Boyle, 2000; Sigman et al., 2010). Understanding the response of AMOC to fresh water addition is therefore crucial, particularly in light of recently accelerating mass loss from the Greenland Ice Sheet and other circum-Atlantic ice masses (Gierz et al., 2015; Vaughan et al., 2013).

Previously, it was suggested that fresh water inputs at high latitudes during H-events caused a cessation of deepwater formation in the North Atlantic, resulting in a shoaling of the northern component water overturning cell by ~1 km (e.g. Alley et al., 1999; Sarnthein et al., 1994; Swingedouw et al., 2009). Support for this concept of a near complete shutdown of AMOC during H-events [came](#) from two main lines of evidence: (i)

51 a dramatic drop in circulation vigour (e.g. McCave et al., 1995a; McManus et al., 2004)
52 and (ii) water mass provenance reconstructions (including those based on $\delta^{13}\text{C}$, Cd/Ca
53 and ^{14}C) indicating an increased presence of nutrient-rich southern-source waters (SSW)
54 below ~2 – 2.5 km depth in the North Atlantic Ocean (e.g. Keigwin et al., 1991;
55 Robinson et al., 2005; Stern and Lisiecki, 2013; Vidal et al., 1997). More recently,
56 however, both of these arguments have been questioned. $^{231}\text{Pa}/^{230}\text{Th}_{\text{xs}}$ records from sites
57 across a wide range of water depths suggests that overturning persisted at shallower
58 depths during H1 (Bradtmeier et al., 2014; Gherardi et al., 2009; Lippold et al., 2016),
59 with values compatible with a near complete shutdown only identified in the SSW cell
60 and during H-events close to glacial maxima (Böhm et al., 2015; Lippold et al., 2009;
61 McManus et al., 2004). A further recent development is the documentation of a poorly
62 ventilated water mass in the Nordic Seas during the last glacial period that overflowed the
63 Greenland-Scotland Ridge (GSR) into the Atlantic basin during the deglaciation
64 (Thornalley et al., 2015). This discovery means that the presence of nutrient-rich waters
65 in the North Atlantic basin may no longer be simply attributed to the incursion of waters
66 from the south. Indeed, excursions to low oxygen and sometimes also low carbon
67 isotopic signatures of benthic foraminifera in the North Atlantic during H-events have
68 been interpreted to suggest that there may also have been overflow of waters from the
69 Nordic Seas at these times (e.g. Meland et al., 2008; Thornalley et al., 2010; Vidal et al.,
70 1998). In addition, bulk sediment leachate ϵ_{Nd} values from the northeast Atlantic (Crocket
71 et al., 2011) are argued to support a persistent presence of overflow waters from the
72 Nordic Seas in the North Atlantic throughout most, if not all, of the last glacial cycle,
73 providing a northern source of nutrient-rich waters to the northeast Atlantic Ocean.

Detailed palaeoceanographic reconstructions show that the classic concept of a simple repeated response of AMOC to freshwater addition during each H-event may be overly simplistic. Not all H-events show a clear perturbation in every oceanographic reconstruction, and there is no consensus on which H-events involve the largest disruption of the ocean-atmosphere system. There is also little agreement about the factors that control the amplitude of AMOC response to freshwater input and considerable debate over the importance of H-event timing relative to the last glacial cycle (Böhm et al., 2015; Lynch-Stieglitz et al., 2014). New high-resolution data sets spanning multiple H-events therefore are needed to help us to gain a better understanding of the range of associated circulation changes in the Atlantic. Records from sites proximal to the GSR are particularly valuable because this region is especially sensitive to changes in northern deep water formation and therefore can help to better constrain variations in the influence of Nordic Seas overflow waters (NSOW).

Bulk sediment leachate Nd isotope data from northeast Atlantic Ocean Drilling Program (ODP) Site 980 (55°29.1'N, 14°42.1'W; 2170 m water depth; location shown in Fig. 1) have been interpreted to suggest that overflow waters crossing the Wyville-Thomson Ridge (WTR) from the Nordic Seas were supplied to the Feni Basin at intermediate depths throughout the past 40 kyr (Crocket et al., 2011). This interpretation invokes either concentrated overflow waters without substantial entrainment of North Atlantic waters or SSW at this site during H-events 1 – 3. Yet, the potential for overprinting of water mass Nd isotope signatures in this region (Lacan and Jeandel, 2004a; Roberts and Piotrowski, 2015), concerns about the fidelity of bulk sediment leachate Nd isotope records (Elmore et al., 2011; Wilson et al., 2013) and the range of

97 water masses influencing the intermediate-depth northeast Atlantic (Lacan and Jeandel,
98 2005) are all sources of uncertainty demanding further careful assessment of the problem.

99 Additional independent proxies of water mass provenance are therefore required to help
100 reconstruct past vertical water mass structure from the Feni Drift.

101 To tackle these gaps in our knowledge, we report the results of a new multi-proxy
102 reconstruction of bottom water chemistry and inferred water mass distribution from ODP
103 Site 980. First, we present neodymium isotope reconstructions from three phases (fish
104 debris, planktonic foraminifera and bulk sediment leachates), together with rare earth
105 element distributions to better understand the controls on neodymium association with
106 foraminifera at Site 980. Then we combine our Nd isotope reconstructions with new, high
107 resolution records of three further proxies for water mass chemistry (B/Ca, $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$
108 in benthic foraminifera) at our study site to shed new light on the response of the AMOC
109 in the northeast Atlantic to freshwater addition during four H-events of the last glacial
110 period.

Fig. 1: (a) The North Atlantic region showing the location of the main study site (ODP Site 980, in black). Dotted lines mark the approximate maximum spatial extent of continental ice sheets during the last glacial maximum (Clark et al., 2012; Dyke et al., 2002; Funder and Hansen, 1996; Sejrup et al., 2005). Arrows represent modern major intermediate and deep current pathways with Nordic Sea overflows (NSOW, orange, also includes Wyville-Thomson Ridge overflow waters or WTROW), Labrador Sea Water (LSW, purple), Northeast Atlantic Deep Water (NEADW, light blue) and Antarctic Bottom Water (AABW, red), based upon McCartney (1992), Hansen and Østerhus (2000), New and Smythe-Wright (2001) and Lacan and Jeandel (2005). (b) Schematic cross-section of the major water masses in the modern Rockall Trough, based upon McGrath et

al. (2012) and Johnson et al. (2010). Colours as panel (a), with the addition of Subpolar Mode Water (SPMW, dark blue). Numbers give neodymium isotope signature of water masses today (Crocker, 2014; Crocket et al., 2011; Lacan and Jeandel, 2004c; Lacan and Jeandel, 2005; Lacan et al., 2012; Stichel et al., 2012). Note that the ε_{Nd} value of AABW shown the AABW endmember recorded in the Atlantic section of the Southern Ocean; this value is likely modified by mixing and boundary exchange as the water mass moves northwards (e.g. Lambelet et al., 2016). (c) North Atlantic region showing the location of the main study site (ODP Site 980, in red) and BOFS sites referred to in the text (black). Grey stripes indicate the main belt of ice-rafted debris deposition in the glacial north Atlantic during the Last Glacial Maximum (Ruddiman, 1977). Dark blue numbers: 1 – Rockall Plateau, 2 – Rockall Trough, 3 – Feni Drift. Last Glacial Maximum limit of perennial sea ice indicated by solid light blue line with dashed light blue line marking extreme limit of winter sea ice (de Vernal et al., 2005; Hillaire-Marcel and de Vernal, 2008).

2. Background

2.1 Site Location and Oceanography

ODP Site 980 is situated on the Feni sediment drift, and features high sedimentation rates across the studied interval (mean 20 cm kyr⁻¹, see details of age model in supplementary information). Its position on the northern fringe of the main Atlantic belt of North American-sourced detrital carbonate deposition (Hemming, 2004) results in lithologically distinct and hence unambiguously identifiable IRD layers. Site 980 is known for its benchmark archives of millennial-scale climate variability during the Quaternary (e.g. McManus et al., 1999; Oppo et al., 2003; 2006, Fig. 2). Existing records from this site cover an unusually long interval for suborbitally resolved records, extending back

147 approximately 500,000 years. Published data from Site 980 spanning the last glacial
148 period are, however, of insufficient resolution to resolve changes in ocean chemistry
149 across H-events clearly.

150 Today, Site 980 is bathed by a mixture of North East Atlantic Deep Water
151 (NEADW) and Labrador Sea Water (LSW), with only a minimal influence of Wyville
152 Thomson Ridge overflow waters (WTROW) from the Nordic Seas (Ellett and Martin,
153 1973; McGrath et al., 2012). During the last glacial period, Site 980 lay close to the
154 interpreted depth of the boundary between northern- and southern-sourced waters (e.g.
155 Boyle and Keigwin, 1987; Curry and Oppo, 2005; Oppo and Lehman, 1993), but this
156 depth is thought to have been bathed by Glacial North Atlantic Intermediate Water
157 (GNAIW) during background glacial conditions (Yu et al., 2008). If AMOC shutdown
158 occurred (as is classically suggested for last glacial H-events), model simulations suggest
159 that Site 980 would have been bathed by Glacial Antarctic Bottom Water (GAABW)
160 during these times (Flückiger et al., 2008; Singarayer and Valdes, 2010; Swingedouw et
161 al., 2009).

Fig. 2: Suborbital variability at ODP Site 980. (a) Data set of McManus et al. (1999), benthic foraminiferal oxygen isotopes in brown and IRD concentrations in pale blue. (b) New oxygen isotope record of benthic foraminiferal calcite (*C. wuellerstorfi*) in red (this study) with data of McManus et al. (1999) in brown (McManus data adjusted by -0.3 ‰ due to an inter-lab offset). (c) IRD concentrations, new data (this study) in dark blue (150 – 500 μm), data from McManus et al. (1999) in pale blue (grain size >150 μm). (d) Percentage of polar species *N. pachyderma* (s.)

as a proportion of the total number of planktonic foraminifera, with high values indicating cold sea surface temperatures (this study). (e) Oxygen isotope values of ice from NGRIP ice core, with less negative values indicating warmer temperatures (North Greenland Ice Core Project Members et al., 2004). [NGRIP data plotted on the GICC05 age model \(Andersen et al., 2006; Rasmussen et al., 2006; Svensson et al., 2008; Vinther et al., 2006\)](#). Grey arrows mark [Site 980](#) recalibrated radiocarbon ages from Oppo et al. (2003) and Benway et al. (2010). Data in panels (b) – (d) are plotted on our new age model (see methods), with intervals of high ice rafting shaded in yellow. [IRD peaks associated with H1-4 are labelled.](#)

2.2 Neodymium isotopes

Over the last few decades, neodymium isotopes have become an important tool for reconstructing the vertical structure of the water column and the provenance and circulation pathways of water masses. A major strength of the Nd isotope technique is that, unlike many commonly used proxies for water mass chemistry (e.g. $\delta^{13}\text{C}$), neodymium isotopes are not influenced by biological processes. The estimated residence time of Nd in the ocean (200 – 1000 years (Arsouze et al., 2009; Tachikawa et al., 2003; Tachikawa et al., 1999)) is shorter than the modern oceanic mixing time of 1000 – 1600 years (e.g. Broecker and Peng, 1982; Garrison, 2011; Sarmiento and Gruber, 2004), giving rise to spatial variation in the isotopic signature of different water masses. Rocks exhibit a wide range of neodymium isotopic compositions, depending upon both their age and initial Sm/Nd ratios. These Nd isotope signatures are typically expressed in epsilon notation or ϵ_{Nd} (representing the $^{143}\text{Nd}/^{144}\text{Nd}$ deviation of a sample relative to a chondrite uniform reservoir in parts per 10,000) with values ranging from $\epsilon_{\text{Nd}} = -56$ for old granitic cratons to +12 for young mid-ocean ridge basalts (Jeandel et al., 2007; Lacan et al., 2012;

Sarbas and Nohl, 2008, and references therein). Neodymium from this range of sources is transferred from the continents to the ocean via both riverine and aeolian inputs, resulting in different water masses acquiring distinct neodymium isotopic signatures (Goldstein et al., 1984; Grousset et al., 1992; Mearns, 1988; Tachikawa et al., 1999). Addition and exchange of neodymium at the sediment-bottom water interface along continental margins also modifies the neodymium isotopic signature of bottom waters through the process of boundary exchange (Lacan and Jeandel, 2004a; Rickli et al., 2014; Wilson et al., 2012).

Analysis of Fe-Mn oxyhydroxides within marine sediments extracted by a leaching procedure (e.g. Gutjahr et al., 2007; Rutberg et al., 2000) allows the production of relatively rapid and high resolution records of bottom water ϵ_{Nd} . This technique has now become widely used (e.g. Böhm et al., 2015; Jonkers et al., 2015; Wei et al., 2015; Wilson et al., 2015). A wide range of other phases have also been proposed to record and preserve bottom water ϵ_{Nd} , including corals (e.g. Copard et al., 2010; van de Flierdt et al., 2006), ferromanganese nodules and crusts (e.g. O'Nions et al., 1978; Piepgras et al., 1979), fish debris (e.g. Lang et al., 2016; Martin and Haley, 2000; Staudigel et al., 1985) and foraminifera both with and without authigenic Fe-Mn oxide coatings (e.g. Elmore et al., 2011; Palmer and Elderfield, 1985).

The ϵ_{Nd} signatures recorded by different substrates do not always agree with one another (e.g. Elmore et al., 2011). For example, a pronounced difference between the glacial and Holocene ϵ_{Nd} values of uncleaned foraminifera is seen along a depth transect of [Biogeochemical Ocean Flux Study](#) (BOFS) cores on and to the south of the Rockall Plateau (Fig. 1), but this difference is muted or absent in co-registered bulk sediment

leachate ϵ_{Nd} (Piotrowski et al., 2012). The leaching of bulk sediment can give a neodymium isotope signature influenced by reactive fine grained sediment components such as volcanic ash (which has a radiogenic signature) and detrital carbonate (which is unradiogenic) (Elmore et al., 2011; Roberts et al., 2010), and thus bulk sediment leachates may not accurately record bottom water chemistry at certain locations. The neodymium isotope signature extracted by bulk sediment leaching is also highly sensitive to the methodology used to separate the authigenic coating signal from the primary sediment (Wilson et al., 2013).

The mostly constant value of $\epsilon_{\text{Nd}} = -10$ recorded by bulk sediment leachates from Site 980 has been interpreted to demonstrate a continuous presence of WTROW in the Rockall Trough for most, if not all, of the last 40 [kyrs](#) due to the similarity of this value to the estimated signature of overflow waters (Crocket et al., 2011). [In contrast](#), modern oceanographic [observations indicate](#) only a minimal influence of WTROW at Site 980 today (Johnson et al., 2010; McGrath et al., 2012). Short-lived radiogenic Nd isotopic excursions identified in the Site 980 record with ages estimated as approximately concurrent to H-events 1 – 4 are inferred to record times of either increased influence of SSW during AMOC shutdowns or a more concentrated presence of overflow waters (Crocket et al., 2011). [However, the](#) limited resolution of the Site 980 bulk sediment leachate ϵ_{Nd} dataset and the absence of direct identification of H-layers in the site stratigraphy [make](#) these conclusions tentative (see also supplementary information). In addition, there are a number of other processes with the potential to alter bottom water ϵ_{Nd} (e.g. variable composition of NEADW or influence of LSW in the Rockall Trough, modification of bottom water isotopic signature by interaction with lithogenic material).

The circulation history of the northeast Atlantic during the last glacial period therefore requires further investigation.

A decoupling between ϵ_{Nd} values and other proxies used as water mass tracers ($\delta^{13}C$, B/Ca (Yu et al., 2008)) suggests that the glacial-interglacial variability in foraminiferal ϵ_{Nd} on and to the south of the Rockall Plateau may not be a result of changing bottom water provenance (Roberts and Piotrowski, 2015). Instead, reactive IRD grains may modify water mass ϵ_{Nd} in this region as they sink through the water column, with large inputs of volcanic glass ($\epsilon_{Nd} = +4.3 \pm 0.1$) to the Rockall Plateau and its southern flank appearing to modify bottom water towards more radiogenic values during the Last Glacial Maximum (Roberts and Piotrowski, 2015). Unradiogenic labelling of the water column by Hudson Bay-derived detrital limestone grains ($\epsilon_{Nd} = -18.6 \pm 0.2$) may also have occurred during Heinrich events (Roberts and Piotrowski, 2015). These observations raise questions over the temporal and spatial extent of water mass ϵ_{Nd} relabelling.

To better understand the meaning of bulk sediment ϵ_{Nd} data from Site 980 and to investigate whether issues of water column relabelling by IRD identified on the Rockall Plateau and its southern margin are applicable to the entire northeast Atlantic, we therefore need higher resolution ϵ_{Nd} records from multiple phases at Site 980 supplemented by a record of concentrations and lithologies of IRD. Furthermore, independent proxies of bottom water chemistry are also required to improve our understanding of water mass structure of the northeast Atlantic during the last 40,000 years at Site 980, therefore we also present co-registered benthic $\delta^{13}C$, $\delta^{18}O$, B/Ca records.

3. Materials and Methods

3.1 Stratigraphy

We sampled Holes 980B (0.05 – 4.12 meters below seafloor (mbsf) at 8 cm resolution) and 980A (0.32 – 4.08 mbsf at 2 cm resolution). We sampled the same core sections as Crocket et al. (2011) to allow direct comparison with their data. Our Hole A samples are not part of the original primary shipboard splice (Shipboard Scientific Party, 1996), therefore, to assign composite depths (metres composite depth, mcd) to these samples, we correlated Hole A mbsf to the primary splice using shipboard-derived whole core volume-specific magnetic susceptibility data measured shipboard (see supplementary information) (Shipboard Scientific Party, 1996). Our study interval corresponds to 0.06 – 8.12 mcd.

3.2 IRD and % *N. pachyderma* (s.)

Sediment samples were sieved at 150–500 μm , and then split until approximately 300 lithic grains remained, which were identified and counted using a binocular microscope, applying the categorizations of Hall et al. (2011). Pumice grains are assumed to have a volcanic ash origin, and so are not included in IRD counts. Reproducibility of IRD counts determined by repeat counts is $\pm 1\%$. IRD fluxes were estimated following Peck et al. (2007), with bulk density values corrected for a systematic error in the ODP Leg 162 shipboard gamma-ray attenuation porosity evaluator (GRAPE) measurements (Jansen et al., 2000). Samples prepared for IRD counts were then split again until approximately

300 foraminifera remained. Counts were performed of the number of specimens of *Neogloboquadrina pachyderma* (sinistral) in these splits and the value expressed as a percentage of the total number of planktonic foraminifera. Reproducibility of % *N. pachyderma* (s.) values determined by repeat counts is $\pm 3\%$.

3.3 Stable isotopes

To generate benthic stable oxygen and carbon isotope reconstructions for our target interval, specimens of *Cibicides wuellerstorfi* (typically 2 – 4) were picked from the >212 μm size fraction (total carbonate mass 60 – 100 μg). Oxygen and carbon isotopic compositions of this carbonate were analysed using a Europa Geo 20-20 stable isotope ratio mass spectrometer at Southampton. All sample values are expressed in delta notation, relative to the Vienna Pee Dee Belemnite standard (VPDB). External reproducibility is better than 0.053 ‰ for $\delta^{18}\text{O}$ and 0.027 ‰ for $\delta^{13}\text{C}$. Species-specific disequilibrium from seawater $\delta^{18}\text{O}$ values was removed by applying an adjustment of +0.64 ‰ (Shackleton and Opdyke, 1973). *C. wuellerstorfi* $\delta^{18}\text{O}$ values were adjusted for global sea level change ($\delta^{18}\text{O}_{\text{ivc}}$) to allow us to understand temporal changes in regional water mass chemistry, following the method of Meland et al. (2008) and using the sea level record shown in Grant et al. (2012) with a maximum glacial-interglacial sea level change of 110 m.

3.4 Neodymium isotope ratios, rare earth elements and trace elements

3.4.1 Sample preparation

To generate neodymium isotope and rare earth element profiles from foraminifera, a

large number (600 – 1600) of planktic foraminifera of mixed species were picked in the >212 μm size fraction from each sample. These were **broken open** then sonicated in ELGA water and methanol to remove clays, before being dissolved in 1.75 M HCl. Ferromanganese coatings were not removed from the foraminifera, as these are rich in pore water-derived Nd and have been shown to record a bottom water signature (Elmore et al., 2011; Roberts et al., 2012).

Fish debris ϵ_{Nd} is considered to be a faithful recorder of the neodymium isotopic composition of bottom waters (e.g. Grandjean et al., 1987; Martin and Haley, 2000; Staudigel et al., 1985; Stille, 1992; Stille and Fischer, 1990). To provide an additional check on the fidelity of our new foraminiferal ϵ_{Nd} record, we therefore picked 1 – 15 pieces of fish debris from the >125 μm size fraction of a subset of samples ($n = 8$). Adhering clays were removed by sonication in methanol and ELGA water. All samples were oxidatively cleaned based upon the methods of Boyle and Keigwin (1985), although a reductive cleaning step was not necessary (Martin and Haley, 2000). An aliquot of 10 μl was extracted from each of the dissolved foraminifera and fish debris samples for rare earth **element** (REE) and other trace element analysis, with the remainder of the sample used for neodymium isotope analysis.

3.4.2 Rare and trace elements

The mechanism by which Nd becomes associated with foraminifera strongly influences whether the isotopic signature preserved is representative of the bottom water signature (Roberts et al., 2012; Tachikawa et al., 2014). The extent of remobilization of Nd in sediment pore waters is of particular interest, and can be investigated using REE

concentrations and distributions (Roberts et al., 2012). Subtle differences in the behaviour of the various REEs result from a decrease in atomic radius as mass increases, leading to differences in their speciation in seawater (Elderfield and Greaves, 1982; Goldberg et al., 1963). The behaviour of cerium is also distinct from the other elements because it can be oxidized from a soluble (3+) to an insoluble (4+) state (de Baar et al., 1985; Elderfield et al., 1988). This distinction can be expressed by the cerium anomaly (Ce/Ce^*), which is the difference in the shale-normalised abundance of cerium compared to the expected value calculated from the nearby light rare earth (LREE) elements ($Ce/Ce^* = 3Ce_n / (2La_n + Nd_n)$, where $_n$ indicates concentrations relative to the Post-Archean Australian Shale, PAAS (McLennan, 1989; Taylor and McLennan, 1985)). Negative cerium anomalies (values <1) indicate a depletion of cerium relative to the other REEs (de Baar et al., 1988).

Rare earth element samples were diluted with 3% HNO₃ spiked with In, Re and Be. Calcium and strontium concentrations were measured with a Quadrupole ICP-MS: Thermo X-Series 2 at Southampton. The remaining sample solutions were analysed for REE and trace element concentrations by High Resolution ICP-MS: Thermo ELEMENT 2XR at Southampton. All samples were corrected for matrix effects and instrument drift using the In, Re and Be standard incorporated into each sample. A blank correction was then applied and rare earth element standards used to correct samples for oxide formation. External reproducibility is estimated as 4–5 %, with internal reproducibility much less than this for the majority of samples. Rare earth element concentrations of the samples (REE_{sample}) were expressed relative to the REE signature of the PAAS (REE_{PAAS}) (McLennan, 1989; Taylor and McLennan, 1985).

3.4.3 Neodymium isotope ratios

Neodymium was purified from the dissolved samples for isotope analysis using standard procedure column chemistry, based upon the methods of Cohen et al. (1988). Cation columns were used to strip iron and titanium from the samples. The remaining material was then run through LN SpecTM columns to **purify and concentrate** neodymium (Pin and Zalduegui, 1997). Samples were analysed by Multi-collector ICP-MS: Thermo NEPTUNE **at** Southampton. Instrumental mass bias ratios were corrected using the procedure of Vance and Thirlwall (2002), adjusting to a $^{146}\text{Nd}/^{144}\text{Nd}$ of 0.7219 and using cerium-doped standards to correct for interference of ^{142}Ce on ^{142}Nd . All values were normalised to the JNdi-1 Standard ($^{143}\text{Nd}/^{144}\text{Nd} = 0.512115 \pm 7$) (Tanaka et al., 2000), and isotopic signatures are expressed in epsilon notation, using a chondritic uniform reservoir value of 0.512638 (Jacobsen and Wasserburg, 1980). Replicate measurements of the JNdi-1 standard across the three analytical sessions gave an external reproducibility 2 s.d. = 0.2 ϵ_{Nd} units for replicate analyses of Nd standard solutions (n = 25). **Error bars plotted show either this external reproducibility or the sample internal reproducibility, whichever is larger.** We compared our Nd data to co-measured concentrations of lithophilic elements (e.g. Al, Ti, Zr, Pb) and volcanic grains, and **found** no evidence to suggest a strong **influence** of volcanic ash, clay contamination or other terrestrial material on foraminiferal ϵ_{Nd} values at Site 980 (see supplementary information).

3.5 Boron/calcium ratios

Between 8 and 12 *C. wuellerstorfi* tests (212 – 500 µm) were cleaned for trace metal analysis following Barker et al. (2003). Foraminiferal tests were cracked and ultrasonicated in Milli-Q 18.2 MΩcm water to remove clay materials. Samples were then oxidatively cleaned in a 1% hydrogen peroxide solution before a weak acid leach is applied to remove adsorbed cations. Finally, samples were dissolved in ~ 0.075M nitric acid. The above steps were all undertaken in a specialised boron free clean laboratory at the University of Southampton. Element ratios were measured using a ThermoFisher Scientific Element 2XR-ICPMS at Southampton, using the protocol described by Rae et al. (2011) employing matrix-matched in house standards and a variety of consistency standards to ensure reproducibility. Samples were screened for clay contamination using Al/Ca and other contaminant ratios (e.g. Ba/Ca, Fe/Ca), with samples Al/Ca > 100 µmol/mol excluded from this study (n = 2). No anomalous values of other elements were identified. Near complete clay removal is assumed for the remaining samples. Long-term reproducibility for B/Ca during this study was within 4 % at 2 s.d. (~ 8 µmol/mol). Carbonate ion concentrations were calculated using a sensitivity of $\Delta[\text{CO}_3^{2-}]$ on B/Ca of 1.14 ± 0.048 for *C. wuellerstorfi* (Yu and Elderfield, 2007), and a $[\text{CO}_3^{2-}]_{\text{sat}}$ calculated from modern local pressure, temperature and salinity.

3.6 Age model generation

A new age model was developed for Site 980 by combining previously published radiocarbon ages (Benway et al., 2010; Oppo et al., 2003) with a new record of the abundance of planktonic foraminifera *N. pachyderma* (s.) correlated to Greenland proxy records (Fig. 2 and supplementary information). An average sedimentation rate of ~ 0.2

mm/yr means that samples with 2 cm spacing give an age resolution of approximately 100 years.

Radiocarbon ages were updated using the Marine13 calibration curve (Reimer et al., 2013) in conjunction with Calib 7.1 software (Stuiver and Reimer, 1993; Stuiver et al., 2005). A constant reservoir age of 400 ± 100 years was assumed, with the exception of the Younger Dryas (800 ± 300 years) and H1 (1600 ± 1000) (e.g. Bard, 1988; Bard et al., 1994; Bondevik et al., 2006; Stanford et al., 2011; Stern and Lisiecki, 2013; Waelbroeck et al., 2001). Additional age constraints were provided by correlating the Site 980 % *N. pachyderma* (s.) record (a proxy for upper ocean temperature when sea surface temperatures are between 4°C and 10°C (Darling et al., 2006)) to North Greenland Ice Core Project (NGRIP) $\delta^{18}\text{O}_{\text{ice}}$ (a proxy for atmospheric temperatures) (North Greenland Ice Core Project Members et al., 2004) on the GICC05 age model (Andersen et al., 2006; Rasmussen et al., 2006; Svensson et al., 2008; Vinther et al., 2006). Visual correlation was carried out using the Analyseries software (Paillard et al., 1996), and is based on the assumption that temperature changes in the upper North Atlantic Ocean are synchronous with Greenland air temperatures. Distinctive peaks in detrital carbonate fluxes were used to confirm the position of H4, H2 and H1.

4. Results and Discussion

4.1 Assessing neodymium isotopes as a water mass tracer at ODP Site 980

4.1.1 Exploring inter-substrate differences in neodymium isotope signatures

New records of REE distributions and Nd isotope signatures of mixed planktonic foraminifera and fish debris from Site 980 are shown together with IRD concentrations in Fig. 3. Throughout our study interval, there is close agreement between new data from fish debris and planktonic foraminifera ϵ_{Nd} data. Glacial ϵ_{Nd} values recorded by fish and foraminifera (-9 to -10) are more radiogenic than Holocene values (~ -11.5), with suborbital unradiogenic excursions from baseline values during H4 (~ 38 ka) and the early Holocene ($\sim 5 - 8$ ka), and a strong radiogenic excursion across H2 ($\sim 23 - 26$ ka) (Fig. 3a). Previously published sediment leachate ϵ_{Nd} data from Site 980 (Crocket et al., 2011) generally show a systematic offset towards more radiogenic values than our new foraminifera and fish debris values by $1 - 2$ ϵ_{Nd} units over the last ~ 11.5 ka (Fig. 3a). This systematic offset is not present in the samples of peak glacial age, where agreement between the two substrates is often much closer, although there are some sporadic disagreements between the two substrates.

Modern oceanographic observations show that the mid-depth western Rockall Trough is predominantly bathed by NEADW ($\epsilon_{Nd} = -12.8 \pm 0.2$) and LSW (-13.9 ± 0.4), with a small influence of overflow waters from the Nordic Seas (-8.2 ± 0.6) (Ellett and Martin, 1973; Johnson et al., 2010; Lacan and Jeandel, 2005; McGrath et al., 2012; Olsen and Ninnemann, 2010). This suggests that the youngest foraminiferal ϵ_{Nd} values at Site 980 (-11.7 ± 0.3) capture bottom water chemistry more accurately than bulk sediment leachate ϵ_{Nd} (-10.2 ± 0.3). A stronger overflow signature than indicated by modern observations (Johnson et al., 2010) or local modification of bottom waters towards more radiogenic values is therefore required if the youngest leachate ϵ_{Nd} signatures at our study site accurately represent recent bottom water chemistry. Alternative explanations include

analytical issues offsetting the bulk sediment leachates to more radiogenic values than the fish debris and foraminifera, and the influence of an additional radiogenic phase is shifting bulk sediment leachates towards more radiogenic values in the Holocene.

Bulk sediment leachates can provide a relatively rapid way to reconstruct seawater chemistry, but a number of studies have highlighted the sensitivity of the technique to the precise method used. Wilson et al. (2013) and Blaser et al. (2016) have recently shown that carbonates play an important role as a buffer when extracting the ferromanganese coatings from sediments, preventing the acid-induced mobilization of Fe oxides and volcanogenic material (which has the potential to shift the measured ϵ_{Nd} towards more radiogenic values). Crocket et al. (2011) followed the earlier procedure of Gutjahr et al. (2007) which included a prior decarbonation step. Leaching time, sample size and pH have also been shown to influence the bulk sediment leachate ϵ_{Nd} signature (Blaser et al., 2016; Wilson et al., 2013)

Radiogenic offsets of bulk sediment leachate ϵ_{Nd} values from foraminiferal Nd signatures have also been documented in core top samples at other sites in the North Atlantic and attributed to the influence of volcanic ash that is easily leached during the extraction process (Elmore et al., 2011). There is no evidence, however, for increased or sustained volcanic activity throughout the Holocene that would increase the accumulation rate of ash in the sediment at Site 980 (see supplementary information). Instead, North Atlantic ash deposition is much more frequent during the last deglaciation (e.g. Abbott and Davies, 2012; Davies et al., 2012; Lowe et al., 2008), when very good agreement exists between foraminiferal and bulk sediment leachate ϵ_{Nd} at Site 980. Airborne volcanic ash therefore cannot easily be invoked to explain the observed offset between

substrates during the Holocene.

Variable transport of silt-sized titanomagnetite grains from the GSR by bottom currents has been documented at a number of sediment drift sites that lie in the path of overflow waters. Higher accumulation of titanomagnetite is documented associated with higher current velocities under interstadial conditions compared to stadials, with grains likely originating from the young basaltic (and radiogenic) rocks of the Nordic Basaltic Province (Ballini et al., 2006; Kissel, 2005; Kissel et al., 1999). Bottom currents were much stronger in the Holocene than the glacial in the North Atlantic (Innocent et al., 1997; Manighetti and McCave, 1995; McCave et al., 1995a; McIntyre and Howe, 2009; McManus et al., 2004; Thornalley et al., 2013), therefore, increased accumulation of leachable fine material at the Feni Drift during the Holocene represents a possible alternative explanation for the more radiogenic signature recorded by Site 980 sediment leachates. There is a possibility that certain other drift deposits in the region may be susceptible to a similar process because the spatial distribution of sites showing unreasonably radiogenic core top sediment leachate ϵ_{Nd} values (as identified by Elmore et al. (2011)) agrees well with the major pathways of modern Nordic Overflow waters (e.g. Hansen and Østerhus, 2000), and bulk sediment leachate ϵ_{Nd} records showing the largest radiogenic offset compared to other substrates during the Holocene have been reported at some other sites in the North Atlantic (Piotrowski et al., 2012; Roberts et al., 2010).

We conclude that the influence of bottom current transported, silt-sized radiogenic particles on sediment leachate ϵ_{Nd} is the most likely explanation for the offset between the Nd isotope substrates during the Holocene at our study site, with leachate methods also contributing to the variable offsets throughout the record. This implies that published

bulk sediment leachate values are not an accurate record of bottom water chemistry at Site 980, at least in the uppermost sedimentary column (above 2.9 mcd). As a consequence, we only consider the Nd isotopic signatures of foraminifera and fish debris as potential tracers of water mass provenance at Site 980.

Fig. 3: ODP Site 980 neodymium isotope and rare earth element chemistry. (a) Neodymium isotope composition (and 2σ error) of mixed planktonic foraminifera (blue), fish debris (red), sediment leachates (grey, (Crocket et al., 2011)) and the detrital fraction (black, (Crocket et al., 2011)). (b) and (c) Nd/Ca and Nd/Mn ratios of mixed planktonic foraminifera. (d), (e) and (f) Ratios of mixed planktonic foraminifera PAAS-normalised REEs (X_n) with (d) $Ce/Ce^* = 3Ce_n/(2La_n+Nd_n)$, (e) $HREE/LREE = (Er_n+Yb_n+Lu_n)/(La_n+Pr_n+Nd_n)$, (f) $MREE/MREE^* = Gd_n+Tb_n+Dy_n/0.5(La_n+Pr_n+Nd_n+Er_n+Yb_n+Lu_n)$. Fish Ce/Ce^* ratios are shown by crosses in panel (d). (g) Fluxes of ice-rafted debris (black) and detrital carbonate (red) grains from the 150 – 500 μm size fraction. Purple shading marks the timing of Heinrich-layer deposition (identified from the IRD and detrital carbonate records). Yellow shading highlights the upper section of the core in which Nd isotope ratios derived from sediment leachates are offset to more radiogenic values than the records of planktonic foraminifera and fish debris (named zone I). The deeper section of the core with higher Nd/Ca values and no systematic inter-substrate ϵ_{Nd} offset is named zone II. Note that depth within sediment core (in metres composite depth) is plotted on a non-linear scale.

4.1.2 Controls on downcore rare earth element distributions

The downcore REE signatures of planktonic foraminifera (shown in Fig. 3b–e) are characterized by upper (zone I) and lower (zone II) intervals with a distinct transition at 2.9 mcd (11.5 ka). Nd/Ca values (Fig. 3b) are consistently relatively low throughout zone I (~700 – 800 nmol/mol), with higher and more variable values in zone II (900 – 3000 nmol/mol). Nd/Mn values are also more variable in zone II, with the highest values typically associated with (or close to) H-events. Below the transition at 2.9 mcd there is less enrichment of heavy REEs over light REEs (Fig. 3d) and cerium anomaly values are closer to 1 (Fig. 3c). A systematic offset between the sediment leachate ϵ_{Nd} values and the other substrates is also recorded in zone I but is not clearly present in zone II (Fig. 3a).

This observation raises the possibility of a link between the processes controlling REE distributions across 2.9 mcd and inter-substrate offsets in ϵ_{Nd} values. We explore three potential explanations for the distinct difference in behavior of REEs in the upper (zone I) and lower (zone II) sections of the studied interval. These are (i) exchange between detrital and authigenic phases, (ii) an active redox front in the sediment and (iii) preserved differences between glacial and interglacial bottom water chemistry.

The REE signature of bulk sediment can co-vary with lithology (e.g. Sholkovitz, 1988). The detrital fraction, however, is typically unreactive, as evidenced by its extraction procedure (Bayon et al., 2002; Jones et al., 1994), and REE concentrations are much lower than in the Fe-Mn oxyhydroxide fraction (Gutjahr et al., 2007). The ϵ_{Nd} values of bulk sediment at Site 980 (Crocket et al., 2011) are consistently offset towards less radiogenic values than any of the substrates discussed here (Fig. 3a). There is also no clear relationship between the relative proportions of radiogenic and unradiogenic IRD and the co-registered foraminiferal ϵ_{Nd} (see supplementary information). These observations suggest that significant exchange between the detrital and authigenic phases is not the main control on foraminiferal REE distributions.

Many of the phases which have been shown to host REEs associated with foraminifera (Fe-Mn (hydr)oxides, organic matter and Mn carbonates) are sensitive to redox conditions (Tachikawa et al., 2014). Therefore, the presence of an active redox front at 2.9 m depth could result in a remobilisation and redistribution of foraminiferal REEs. The similarity of shift in REE distribution recorded by both foraminifera and fish debris, however, argues against this explanation at Site 980. The REE signal of fish debris is acquired at shallow depths in the sediment and is relatively resistant to

diagenesis, with substitution of rare earth elements into the crystal lattice occurring only under more extreme conditions and resulting in a distinctive bell-shaped REE profile (Martin and Scher, 2004; Reynard et al., 1999; Staudigel et al., 1985). This profile is not seen in any of the samples presented here (Fig. 4). In addition, strong negative Ce/Ce* values are preserved in Miocene fish teeth at nearby Site 982 (Martin et al., 2010), similar to that preserved by Holocene (but not glacial) samples at Site 980, suggesting that REE distributions are not reset below the top few metres of sediment. Therefore, it is unlikely that diagenetic redistribution explains the shift in REE distribution in either uncleaned foraminifera or fish debris, supporting our use of fish debris and foraminiferal REE chemistry at Site 980 to reconstruct past oceanographic conditions.

The cerium anomaly (Ce/Ce*) is used to track oxidation because cerium is readily converted from the insoluble Ce (IV) to the soluble Ce (III) phase when water column oxygenation decreases (Elderfield et al., 1988). It should, however, be noted that Ce/Ce* preserved by sediment components is not an infallible recorder of local bottom water oxidation, with questions raised regarding the incorporation of REEs into various marine substrates (e.g. German and Elderfield, 1990; Holser, 1997; MacLeod and Irving, 1996). In addition, lower Ce/Ce* ratios are recorded in the modern Pacific and Indian Oceans compared to the North Atlantic Ocean; a result which is attributed to a progressive removal of Ce as the age of a bottom water mass increases (German and Elderfield, 1990). The higher Ce/Ce* values at Site 980 during the glacial could therefore indicate a more vigorous overturning circulation than during the Holocene, with several studies supporting vigorous renewal of intermediate waters in the North Atlantic under glacial conditions (e.g. Gherardi et al., 2009; McCave et al., 1995a). An alternative explanation

569 is that the higher Ce/Ce* values in zone II (close to 1; Fig. 3c) in our record indicate that
570 glacial bottom waters at Site 980 had a relatively low oxygen content compared with
571 conditions in the Holocene. A similar increase in both Ce/Ce* and Nd/Ca of uncleaned
572 planktonic foraminifera during the transition into the Holocene has been recorded at
573 Bermuda Rise (Roberts et al., 2012). Glacial-interglacial differences in the degree of
574 oxygenation of shallowly buried sediments have been documented in a number of studies
575 around the world (Jaccard and Galbraith, 2012), with decreased oxygenation of glacial
576 bottom waters reported in the intermediate depth northeast Atlantic (Baas et al., 1998;
577 Schönfeld et al., 2003). The broad coincidence of the boundary between zone I and II
578 with the climatic transition into the Holocene therefore raises the possibility that the
579 differences in REE behaviour at Site 980 are the product of differences in the degree of
580 pore water oxidation between glacial and interglacial conditions.

Fig. 4: Comparison of REE profiles of ODP Site 980 fish debris (dotted lines) and mixed planktonic foraminifera (solid lines). Values are normalised to PAAS and La = 1, with samples from zone I in panel (a) and zone II in panel (b). Sample ages given in figure legend.

4.1.3 Assessing the influence of water mass labelling by IRD at Site 980

Foraminiferal ϵ_{Nd} values during the late Holocene (-11.7 ± 0.3) at Site 980 are consistent with the representative signatures of the water masses bathing this site today but determining whether older ϵ_{Nd} signatures are also representative of regional water mass chemistry requires further evaluation. The radiogenic signal (-7.9 to -4.2) in foraminiferal ϵ_{Nd} of the nearby BOFS cores (Fig. 1) during the Last Glacial Maximum has been interpreted to reflect labelling of the bottom waters bathing these sites by IRD raining through the water column (Roberts and Piotrowski, 2015). Therefore, before we can use foraminifera-based ϵ_{Nd} data from Site 980 to determine past variations in water mass sourcing, first we must explore the possibility that exchange with lithogenic material has influenced glacial ϵ_{Nd} values in our datasets.

Three lines of evidence suggest that there was a smaller influence of local modification of water mass chemistry by IRD at ODP Site 980 than the BOFS sites. First, the radiogenic ϵ_{Nd} values (≥ -8) reported by Roberts and Piotrowski (2015) during MIS 2 are not recorded at Site 980 (Fig. 5a). Second, we observe ϵ_{Nd} excursions of opposing signs during H2 and H4, despite lithologically similar IRD assemblages (Fig. 3a and supplementary information). Third, our new IRD records demonstrate that IRD inputs during the last glacial period at Site 980 (typically 0 – 650 grains/g, peaking at 1450 grains/g during H2) are lower than those recorded at the BOFS sites (400 – 4700 grains/g, (Roberts and Piotrowski, 2015)) (Fig. 5c and d). The same result is also true of the concentrations of volcanic grains, highlighting that the Feni Drift was likely subject to a different IRD depositional regime than the one influencing the BOFS sites during the last glacial period. Although we cannot discount the influence at Site 980 of water masses modified by IRD upstream of our site, and no direct correlation was found between

volcanic IRD concentration and ϵ_{Nd} values by Roberts and Piotrowski (2015), we suggest that the absence of large volumes of reactive material sinking through the water column at Site 980 during the last glacial period reduced the potential for local modification of water mass chemistry. We therefore suggest that IRD water mass relabelling is not the dominant control on the down core variability in our ϵ_{Nd} data. Hence, modification of the water column ϵ_{Nd} by IRD (Roberts and Piotrowski, 2015) may not be significantly problematic in all regions of IRD deposition.

Fig. 5: Comparison of palaeoclimate records from Site 980, Feni Drift (black, this study and Crocket et al. (2011)) and BOFS sites (colour) on and along the southern flank of the Rockall

Plateau (Roberts and Piotrowski, 2015) (water depths in legend, site locations shown in Fig. 1).
(a) Uncleaned foraminifera (solid circles and lines) and fish debris ϵ_{Nd} (crosses). (b) Bulk
sediment leachate ϵ_{Nd} values. (c) Concentration of IRD grains 150 – 500 μm (Site 980) and >150
 μm (BOFS cores (Roberts and Piotrowski, 2015)). (d) Concentration of volcanic IRD (including
basalt and mafic and felsic volcanic glasses) and pumice clasts with grain size 150 – 500 μm (Site
980) and >150 μm (BOFS sites, Roberts and Piotrowski (2015)). (e) Sedimentation rates (cm/kyr)
based on published age models (BOFS sites, Roberts and Piotrowski (2015)) and this study (Site
980).

4.2 Water mass changes

Having established that fish debris and uncleaned planktic foraminifera provide a reliable
record of bottom water Nd isotope composition at ODP Site 980, next we combine these
 ϵ_{Nd} data with three other proxies for water mass chemistry (B/Ca, $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$) to
reconstruct oceanographic variability in the mid-depth northeast Atlantic over the past
40,000 years.

Fig. 6: Bottom water chemistry shifts at ODP Site 980 during H-events. Shaded bands mark the position of the H-events identified by IRD abundance at Site 980. (a) Bottom water carbonate ion

concentrations reconstructed from B/Ca ratios of benthic foraminifera *C. wuellerstorfi*, with error bars indicating analytical uncertainty (calibration uncertainty $\pm 10\mu\text{mol/kg}$ (Yu and Elderfield, 2007)). (b) Neodymium isotope ratios of planktonic foraminifera with ferromanganese coatings not removed (in blue) and fish debris (red) (error bars $\pm 2\sigma$). (c) $\delta^{13}\text{C}$ of *C. wuellerstorfi*, with darker line colour marking the three-point running mean. (d) $\delta^{18}\text{O}$ of *C. wuellerstorfi*, adjusted for global ice volume changes, with darker line colour marking the three-point running mean. (e) Fluxes of ice-rafted debris (black) and detrital carbonate clasts (red) in grains $150 - 500\ \mu\text{m cm}^{-2}\ \text{kyr}^{-1}$.

4.2.1 Glacial-interglacial variability

In Fig. 6, we present Site 980 fish debris and foraminiferal ϵ_{Nd} alongside benthic $\delta^{13}\text{C}$ and B/Ca ratios for the past 41 **kyrs**. A distinct difference in the properties of the water mass bathing Site 980 can be seen between the glacial and Holocene, with glacial waters having a more radiogenic ϵ_{Nd} signature (Fig. 6b), slightly **lower** carbon isotope values (Fig. 6c) and higher $[\text{CO}_3^{2-}]$ (indicated by higher benthic foraminiferal B/Ca ratios, Fig. 6a). **These observations strongly suggest that the interpretation of Crocket et al. (2011) invoking the persistent dominance of WTROW throughout the last 40 kyr at this site cannot be correct without a significant shift in the properties of WTROW.**

Marked similarities exists between the long-term evolution of ϵ_{Nd} at Site 980 and deep water sites at the Bermuda Rise (Böhm et al., 2015; Gutjahr and Lippold, 2011; Roberts et al., 2010) and South Atlantic Cape Basin (Piotrowski et al., 2012; Piotrowski et al., 2005; Rutberg et al., 2000) (Fig. 7). The Bermuda Rise and Cape Basin records are primarily controlled by mixing between **unradiogenic northern- and radiogenic southern-sourced** waters. An increased contribution of radiogenic **southern-sourced** waters mixing

with northern-sourced waters in the glacial northeast North Atlantic cannot be ruled out on the basis of Nd isotopes alone, but B/Ca-based reconstructions of carbonate ion concentrations argue against this interpretation. Both modern and glacial SSW typically have low $[\text{CO}_3^{2-}]$ (75 – 95 $\mu\text{mol/mol}$), whereas modern North Atlantic Deep Water and Nordic Sea overflows are much richer in $[\text{CO}_3^{2-}]$ (>100 $\mu\text{mol/mol}$) (Key et al., 2004; Yu et al., 2014; Yu et al., 2008). Reconstructed glacial carbonate ion concentrations (~130 – 150 $\mu\text{mol/mol}$) at Site 980 are higher than those in the Holocene (~115 – 135 $\mu\text{mol/mol}$) (Fig. 6), the opposite direction of change as would be expected if an increased glacial presence of southern-sourced waters is the dominant driver of the difference in bottom water chemistry observed at Site 980 between the Holocene and last glacial period.

A reduction in the influence of unradiogenic LSW under glacial conditions (Cottet-Puinel et al., 2004; Hillaire-Marcel et al., 2001) could contribute to the more radiogenic glacial ϵ_{Nd} signature at Site 980, however, as modern LSW has high $[\text{CO}_3^{2-}]$ (Key et al., 2004; Yu et al., 2008), a decrease in glacial LSW influence alone cannot explain the observed bottom water chemistry changes at Site 980. A decreased export of Mediterranean outflow water (MOW) has been documented under glacial conditions (Zahn et al., 1997), but as modern MOW is moderately radiogenic (-9.4 ± 0.5 (Spivack and Wasserburg, 1988)), a reduction in MOW cannot explain the glacial shift towards more radiogenic bottom waters at Site 980. Alternatively, Rogerson et al. (2005) suggest that the core of MOW deepened which could increase its influence at Site 980, and as MOW has a high modern $[\text{CO}_3^{2-}]$, potentially contribute to the observed shift in glacial-interglacial bottom water chemistry. An increased influence of relatively radiogenic and high $[\text{CO}_3^{2-}]$ Nordic Sea overflow waters at Site 980 during the last glacial period as

suggested by Yu et al. (2008) can also explain the more radiogenic and $[\text{CO}_3^{2-}]$ -rich glacial bottom water signature at Site 980. It should be noted that an increased influence of overflow waters on bottom water chemistry at Site 980 does not require an increase in overflow current vigour, particularly in light of the decreased contribution of LSW (Cottet-Puinel et al., 2004; Hillaire-Marcel et al., 2001) to the main Atlantic basin during the last glacial period.

Fig. 7: Comparison of Site 980 uncleaned foraminiferal ϵ_{Nd} records (black) with deep sites at the Bermuda Rise (ODP 1063/OCE326-GGC6 in red/orange (Böhm et al., 2015; Gutjahr and Lippold, 2011; Roberts et al., 2010)) and South Atlantic (TNO57-21/RC11-83 in blue/purple (Piotrowski et al., 2012; Piotrowski et al., 2004, 2005)). Foraminiferal ϵ_{Nd} values are shown with solid lines and filled circles, while bulk sediment leachates are shown with dotted lines and

hollow circles. Note that GGC6 bulk sediment leachate values are not shown as these were deemed not to be representative of bottom water chemistry by Roberts et al. (2010).

4.2.2 Heinrich events

Our multi-proxy reconstruction of bottom water properties at Site 980 shows prominent excursions associated with increased concentrations of IRD during each of H-events 4 – 1 (Fig. 6). Large shifts in the ϵ_{Nd} of bottom waters are documented for both H4 ($\epsilon_{\text{Nd}} = -13.6$) and H2 (-7.1), but these excursions are of opposing signs. A small ($\sim 20 \mu\text{mol/kg}$) reduction in bottom water $[\text{CO}_3^{2-}]$ is recorded during H2 and just prior to H3, but there is no significant change during H4, and at no point in the record do the values get close to SSW values ($75 - 95 \mu\text{mol/kg}$ (Key et al., 2004; Yu et al., 2014; Yu et al., 2008)). Excursions towards low benthic $\delta^{13}\text{C}$ are recorded during each of the H-events, although typically lagging slightly behind the main IRD peak.

Arguably, the most notable feature is the unique expression of each event in our proxy records of bottom water chemistry, with differences in the sign of change, as well as the magnitude of excursions. This simple observation shows that circulation changes in the mid-depth glacial northeast Atlantic are much more complex than simple two component northern- and southern-sourced endmember mixing model. Instead, either three or more water masses must have influenced Site 980 during the last glacial period, or alternatively, shifts in surface ocean properties (particularly in deep water formation regions) or circulation vigour significantly modified the properties of water masses.

The surface of the North Atlantic Ocean is thought to have been a highly dynamic environment during the last glacial period, with temporally variable freshwater input

from both icebergs and continental sources (Heinrich, 1988; Hemming, 2004; Leksens et al., 2006; Stanford et al., 2011). Fluctuations in sea surface temperature, salinity, sea ice cover and productivity have all been documented in the high northern latitudes (e.g. Dokken et al., 2013; Maslin et al., 1995; van Kreveld, 1996). An increase in sea ice cover has the potential to reduce air-sea exchange in the deep water formation regions, with CO₂ invasion and evasion acting upon both the $\delta^{13}\text{C}$ and carbonate ion concentration of a water mass in a specific ratio (Fig. 8) (Yu et al., 2008). The variability in our data between H-events and non-Heinrich glacial intervals does not follow this trend, therefore we infer that variable sea-ice cover in deep water formation regions is not a major contributor to the variability in bottom water chemistry recorded at Site 980 during H-events. Instead, there is a closer fit to the “biology” slope, which represents the influence of biological regeneration on seawater chemistry and incorporates the effects of organic tissue degradation and CaCO₃ remineralisation (Yu et al., 2008). The influence of a water mass with increased organic matter remineralisation may therefore explain part of the H-event signals in our proxy records. This signature could reach the site either by increased surface productivity or increased age of the bottom waters reflecting a longer time interval since the water mass was exposed to the atmosphere, either as a result of a rearrangement of circulation patterns, a shift in water mass boundaries or a decrease in circulation vigour. Additional independent proxy records for current vigour such as sortable silt (McCave et al., 1995b) or $^{231}\text{Pa}/^{230}\text{Th}$ (Yu et al., 1996) could therefore prove valuable in better understanding the origin of the observed excursions of bottom water chemistry in the northeast Atlantic during H-events.

Fig. 8: Comparison of Site 980 bottom water chemistry to regional water mass signatures. (a) Cross-plot of co-measured oxygen and carbon isotopes of *C. wuellerstorfi* from ODP Site 980. All values are smoothed (3-point running mean), with $\delta^{18}\text{O}$ values adjusted for global ice volume changes. Estimated water mass compositions shown in grey, based upon Bertram et al. (1995), Voelker et al. (2006), Meland et al. (2008) and Thornalley et al. (2010). Water masses abbreviations: GAABW: Glacial Antarctic Bottom Water; GAAIW: Glacial Antarctic Intermediate Water; LSW: Labrador Sea Water (Holocene value); GNAIW: Glacial North Atlantic Intermediate Water; GNSDW: Glacial Norwegian Sea Deep Water. Note that the isotopic composition of brines has high uncertainty (e.g. Meland et al., 2008; Thornalley et al., 2010). (b) Cross-plot of $\delta^{13}\text{C} C. wuellerstorfi$ and estimated bottom water $[\text{CO}_3^{2-}]$ calculated from *C. wuellerstorfi* B/Ca ratios (see methods). Also shown are lines indicating the calculated slope of co-variance in $\delta^{13}\text{C}$ and $[\text{CO}_3^{2-}]$ generated by biological regeneration and CO_2 invasion/evasion (Yu et al., 2008). Reconstructed water mass signatures are shown in grey (data from Yu et al., 2014; Yu et al., 2008). NEAP 8K is thought to record a strong influence of overflow waters from the Nordic Seas at the [Last Glacial Maximum](#) (Yu et al., 2008).

4.2.2.1 Heinrich event 4

Fig. 9: Reconstruction of changes in surface and deep ocean during H4. (a) Bottom water carbonate ion concentrations reconstructed from B/Ca ratios of benthic foraminifera *C. wuellerstorfi* with error bars indicating analytical uncertainty (calibration uncertainty ± 10

769 $\mu\text{mol/kg}$ (Yu and Elderfield, 2007)). (b) Neodymium isotope ratios of planktonic foraminifera
770 with ferromanganese coatings not removed (in blue) and fish debris (red). Error bars indicate
771 $\pm 2\sigma$. (c) $\delta^{13}\text{C}$ of *C. wuellerstorfi*, with darker line colour marking the three-point running mean.
772 (d) $\delta^{18}\text{O}$ of *C. wuellerstorfi*, corrected for global ice volume changes, with darker line colour
773 marking the three-point running mean. (e) Percentage of polar species *N. pachyderma* (s.) as a
774 proportion of the total number of planktonic foraminifera. (f) Fluxes of ice-rafted debris (black)
775 and detrital carbonate (red) grains from the 150 – 500 μm size fraction of Site 980 samples.

776
777 Heinrich event 4 is distinct from the other H-events in our record because it is
778 characterised by an excursion towards unradiogenic ϵ_{Nd} values and the most significant
779 decrease in benthic foraminiferal $\delta^{18}\text{O}$ in our record (Figs. 6d and 9). Similar $\delta^{18}\text{O}$
780 excursions have been linked to episodes of brine formation in the Nordic Seas (Dokken
781 and Jansen, 1999; Meland et al., 2008; Vidal et al., 1998), although the origin of the low
782 $\delta^{18}\text{O}$ signature is debated (e.g. Bagniewski et al., 2015; Bauch and Bauch, 2001;
783 Rasmussen and Thomsen, 2009; Stanford et al., 2011). Regardless, a Nordic Sea origin
784 for the waters bathing Site 980 during H4 can likely be discounted because modern
785 Nordic Seas Overflows have radiogenic signatures of $\epsilon_{\text{Nd}} = -8.2 \pm 0.6$ (Lacan and
786 Jeandel, 2004b) while our data record a shift of ~ 2 ϵ_{Nd} units towards more unradiogenic
787 values. A very large shift in the neodymium isotope composition of the Nordic Seas (for
788 example, due to overprinting by unradiogenic IRD from Greenland) would be required
789 for the shift in bottom water chemistry recorded at Site 980 to be attributed to increased
790 presence of Nordic Seas Overflows at the site. The main source of unradiogenic waters in
791 the North Atlantic today is the Labrador Sea. LSW generation is thought to have been
792 weaker during the last glacial period, although with some evidence of sinking of brine-

rich waters during Heinrich events (Cottet-Puinel et al., 2004; Hillaire-Marcel and de Vernal, 2008; Hillaire-Marcel et al., 2001; Nuttin et al., 2015; Weber et al., 2001).

Alternatively, an increase in deep water formation south of the GSR could explain the ϵ_{Nd} signal at Site 980 during H4 (Duplessy et al., 1980; Labeyrie et al., 1992). Surface waters today in the Northeast Atlantic have ϵ_{Nd} values of -13 to -14.8 (Lacan and Jeandel, 2004c), although southward migration of the polar front during H-events (e.g. Eynaud et al., 2009) may have resulted in more radiogenic surface water signatures. Reconstructions of surface water properties in the open ocean suggest that they may have been insufficiently dense to sink (Maslin et al., 1995). Alternatively, brine formation, could have occurred on the European and/or Icelandic margins, Rockall Plateau or GSR (Meland et al., 2008; Thornalley et al., 2010), with a possible additional input of waters with low $\delta^{18}O$ and $\delta^{13}C$ signatures from the European continent (e.g. Eynaud et al., 2007; Toucanne et al., 2009).

4.2.2.2 Heinrich event 2

Heinrich event 2 shows a very different signal to H4 in our multi-proxy records. Unlike H4, there is no $\delta^{18}O$ excursion (Fig. 6d), and ϵ_{Nd} values shift ~ 2 units towards more radiogenic values (> -8 ; Fig. 6b). A combination of radiogenic ϵ_{Nd} and low $\delta^{13}C$ with no concurrent excursion in $\delta^{18}O$ in the Atlantic basin are commonly associated with SSW in the Atlantic (e.g. Jonkers et al., 2015; Piotrowski et al., 2008; Rutberg et al., 2000; Thornalley et al., 2010). We also see a small decrease in carbonate ion concentrations ($\sim 20 \mu\text{mol/kg}$), which supports an increase in the contribution of [southern-sourced](#) waters to the northeast North Atlantic. Our reconstructed $[\text{CO}_3^{2-}]$ of $> 120 \mu\text{mol/kg}$ at Site 980

during H2 is, however, well above modern Antarctic bottom and intermediate waters (~ 85 $\mu\text{mol/kg}$) (Key et al., 2004; Yu et al., 2008), and even more discrepant with values of ~ 80 $\mu\text{mol/kg}$ reported for deep SSW during the Last Glacial Maximum (Yu et al., 2014), so any southern-sourced water would need to be highly diluted by mixing with northern-sourced endmembers. A alternative source of radiogenic waters with high $[\text{CO}_3^{2-}]$ is the Nordic Seas (Lacan and Jeandel, 2004b; Lacan et al., 2012), a possibility consistent with proposed strengthening of GSR overflows into the North Atlantic during H-events (Crocket et al., 2011; Meland et al., 2008; Thornalley et al., 2010).

A clear temporal lag (~ 2 kyr, represented by 30 cm of core) exists between the onset of excursions in ϵ_{Nd} and $\delta^{13}\text{C}$ during H2 (Fig. 10). The initial shift in ϵ_{Nd} values is approximately concurrent with increasing % *N. pachyderma* (s.) values. There is no distinct offset between % *N. pachyderma* (s.) and IRD fluxes during either the onset or termination of Heinrich stadial 2, providing independent support of the suggestion that cooling preceded IRD deposition in the North Atlantic during H-events (Barker et al., 2015). Bottom water $[\text{CO}_3^{2-}]$ values only start to decrease when stadial conditions are well established. The delayed excursion towards low $\delta^{13}\text{C}$ values could be a result of a local or regional pulse in the flux of organic matter to the seafloor (Mackensen et al., 1993), or the flushing of previously poorly ventilated waters once ice-rafting has ceased and circulation vigour is increasing. Alternatively, the observed lag may indicate the presence of multiple phases of AMOC reorganization (Wilson et al., 2014), involving the existence of short-term circulation modes, rather than a simple switch between two distinct AMOC states ('glacial' and 'off') (Rahmstorf, 2002).

Fig. 10: Reconstruction of changes in surface and deep ocean during H2. Symbols as in Fig. 9.

840

841 4.2.2.3 Exploration of the differences between H-events

842 Comparison between multiple H-events allows greater insight into the processes
843 governing AMOC response to fresh water additions. The results of two recent studies
844 (Böhm et al., 2015; Lynch-Stieglitz et al., 2014) suggest that an important role in
845 determining the magnitude of oceanographic response to freshwater input is played by
846 the stage in the glacial cycle at which freshwater was released. One of these studies,
847 based on reconstruction of flow through the Florida Straits, suggests that AMOC was
848 least sensitive to freshwater forcing during full last glacial conditions when stratification
849 of the Atlantic Ocean was at its greatest (Lynch-Stieglitz et al., 2014). The other, a study
850 of water chemistry in the deep North Atlantic on Bermuda Rise, suggests that during the
851 last glacial cycle, AMOC shutdown only occurs when a threshold in continental ice sheet
852 size is crossed, with ice volumes close to their maximum (Böhm et al., 2015). While the
853 magnitude of AMOC perturbation cannot be simply gauged by the amplitude of the
854 excursion in our proxy records, the variability across our data sets strongly suggests a
855 discernible reorganization in ocean structure associated with both H2 (which occurs at
856 close to peak glacial conditions) and H4 (when ice sheets were much smaller (e.g. Siddall
857 et al., 2003)). Our results therefore offer a different perspective, suggesting that the stage
858 of the glacial cycle occupied by H-events may not be the principle factor governing the
859 response of the mid-depth North Atlantic to freshwater release.

860 The unique chemical signature of intermediate waters during each of the H-events
861 examined (H4 to H1) at Site 980 therefore requires an alternative explanation to changes
862 in global ice volume. Differences in the location, duration and magnitude of melt water

inputs have been shown to exert a clear influence over the response of AMOC in modelling experiments (Bigg et al., 2011; Otto-Bliesner and Brady, 2010; Smith and Gregory, 2009). Differences in the IRD assemblages have been documented between H-events (e.g. Grousset et al., 1993; Hall et al., 2011; Peck et al., 2007; Snoeckx et al., 1999), most notably a decreased flux of Hudson Bay-sourced detrital carbonates reaching the North Atlantic during H3 and H6 (e.g. Bond et al., 1992; Hemming, 2004; Hodell and Curtis, 2008), suggesting differences in the relative fresh water contributions from the circum-Atlantic ice sheets. Each H-event is therefore likely associated with different volumes, rates and/or delivery routes of freshwater to the North Atlantic Ocean. These differences are likely to have resulted in shifts in the relative strength of surface water subduction in a number of different potential deep water formation areas. This interpretation of events also offers a potential explanation as to why different events have a different prominence in different locations, and therefore why records from different study regions point to different relationships between global ice volume and AMOC stability (Böhm et al., 2015; Lynch-Stieglitz et al., 2014). AMOC strength has also been shown to exert a strong control over continental hydroclimate (e.g. Burckel et al., 2015; Mulitza et al., 2008), therefore, variations in meltwater inputs between H-events and the resulting AMOC perturbations may have important implications for terrestrial climate.

The unique response of intermediate water to fresh water input during each H-event examined (4 to 1) also raises the possibility of variations in the properties of GNAIW during H-events; a result that may have implications for downstream reconstructions of the strength of exported water from the high northern latitudes at these

times. Glacial bottom water chemistry at Site 980 however appears relatively stable between H-events.

Our records indicate that a dominance of northern-sourced waters in the mid-depth northeast North Atlantic persisted throughout the last glacial period, but the absence of a strong influence of SSW at our intermediate depth study site does not preclude a dramatic slowdown of the deep ocean during certain H-events (Burckel et al., 2015; Mulitza et al., 2008). In fact, a shoaling of the overturning circulation cell is a feature common to many simulations of freshwater input to the North Atlantic where AMOC shutdown is incomplete (Böhm et al., 2015).

5. Summary and Conclusions

We investigate the extent to which uncleaned foraminifera, fish debris and bulk sediment leachates can record regionally representative bottom water neodymium isotope compositions in the northeast North Atlantic by examining the isotopic signature of multiple substrates and associated rare earth element distributions. We find that neither local modification of bottom water chemistry by IRD labelling or redistribution of REE at depth within the sediment column exert a strong control over the REE and ϵ_{Nd} signatures preserved by planktonic foraminifera without their Fe-Mn oxide coatings removed at Site 980, making them suitable substrates for palaeoceanographic reconstructions. Foraminiferal and fish debris ϵ_{Nd} values show good agreement throughout the last 40,000 years, however, bulk sediment leachate values are offset towards more radiogenic values by 1–2 ϵ_{Nd} units throughout the Holocene, which we

908 attribute to the increased influence of fine-grained radiogenic material transported by
909 bottom currents on the sediment leachate values at this time. This result calls for careful
910 evaluation of extracted leachate neodymium isotope compositions for leaching artifacts at
911 North Atlantic drift sites.

912 By combining our new Nd isotope records with three additional proxies for water
913 mass chemistry (B/Ca, $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ signatures of *C. wuellerstorfi*), we demonstrate that
914 there is an increased influence of Nordic Seas overflow waters (and not southern-sourced
915 waters) at 2.2 km depth in the Rockall Trough during the late glacial, possibly related in
916 part to a reduction in LSW generation. A dominant presence of SSW at Site 980 at any
917 time within the past 40 kyr is ruled out by reconstructed $[\text{CO}_3^{2-}]$ greater than 110 $\mu\text{mol/kg}$
918 throughout our record. We find that all four H-events within our study interval have
919 different geochemical signatures, with H4 and H2 marked by ϵ_{Nd} excursions in opposing
920 directions. Unradiogenic ϵ_{Nd} values and light oxygen and carbon isotope values during
921 H4 could indicate an increased contribution of waters from either the Labrador Sea or
922 deep waters forming south of the Greenland-Scotland Ridge, while the radiogenic ϵ_{Nd}
923 signature recorded during H2 is more likely explained by an increased presence of
924 overflow waters from the Nordic Seas. We find that the stage of the glacial cycle
925 occupied by H-events may not be the principle factor governing the response of the mid-
926 depth North Atlantic to freshwater release. Instead, we suggest that this heterogeneity in
927 the intermediate depth northeast North Atlantic most likely arises due to differences in
928 fresh water input locations, magnitudes and fluxes amongst H-events. The balance of
929 contributions from different source regions to the northern intermediate/deep water
930 endmember does not appear to remain constant, with important implications for the

chemical signature of GNAIW that is transported downstream during millennial-scale climate events.

Acknowledgements

This research used samples provided by the Integrated Ocean Drilling (Discovery) Program IODP, which is sponsored by the US National Science Foundation and participating countries under management of Joint Oceanographic Institutions, Inc. We thank Walter Hale and Alex Wülbers for help with sampling and Matt Cooper, Andy Milton, Mike Bolshaw and Dave Spanner for analytical support. Heiko Pälike, David Thornalley and Rachel Mills are thanked for productive discussions and comments on earlier versions of this work. We also thank three anonymous reviewers for their constructive feedback, which greatly improved the manuscript. Funding for this project was provided by NERC studentships to A.J.C. (grant NE/D005728/2) and T.B.C. (NE/I528626/1), with additional funding support from a Royal Society Wolfson Research Merit Award and NERC grants NE/F00141X/1 and NE/I006168/1 to P.A.W. and NE/D00876X/2 to G.L.F.

References

953 Abbott, P.M., Davies, S.M., 2012. Volcanism and the Greenland ice-cores: the tephra
 954 record. *Earth-Science Reviews* 115, 173-191.
 955 Adkins, J.F., 2013. The role of deep ocean circulation in setting glacial climates.
 956 *Paleoceanography* 28, 539-561.
 957 Alley, R.B., Clark, P.U., Keigwin, L.D., Webb, R.S., 1999. Making sense of millennial-
 958 scale climate change. *Geophysical Monograph - American Geophysical Union* 112, 385-
 959 394.
 960 Andersen, K.K., Svensson, A., Johnsen, S.J., Rasmussen, S.O., Bigler, M., Röthlisberger,
 961 R., Ruth, U., Siggaard-Andersen, M.-L., Steffensen, J.P., Dahl-Jensen, D., Vinther, B.M.,
 962 Clausen, H.B., 2006. The Greenland Ice Core Chronology 2005, 15-42 ka. Part 1:
 963 constructing the time scale. *Quaternary Science Reviews* 25, 3246-3257.
 964 Arsouze, T., Dutay, J.C., Lacan, F., Jeandel, C., 2009. Reconstructing the Nd oceanic
 965 cycle using a coupled dynamical-biogeochemical model. *Biogeosciences* 6, 2829-2846.
 966 Baas, J.H., Schönfeld, J., Zahn, R., 1998. Mid-depth oxygen drawdown during Heinrich
 967 events: evidence from benthic foraminiferal community structure, trace-fossil tiering, and
 968 benthic $\delta^{13}\text{C}$ at the Portuguese Margin. *Marine Geology* 152, 25-55.
 969 Bagniewski, W., Meissner, K.J., Menviel, L., Brennan, C.E.C.P.A., 2015. Quantification
 970 of factors impacting seawater and calcite $\delta^{18}\text{O}$ during Heinrich Stadials 1 and 4.
 971 *Paleoceanography* 30, 895-911.
 972 Ballini, M., Kissel, C., Colin, C., Richter, T., 2006. Deep-water mass source and dynamic
 973 associated with rapid climatic variations during the last glacial stage in the North
 974 Atlantic: A multiproxy investigation of the detrital fraction of deep-sea sediments.
 975 *Geochem. Geophys. Geosyst.* 7, Q02N01.

976 Bard, E., 1988. Correction of Accelerator Mass Spectrometry ^{14}C Ages Measured in
 977 Planktonic Foraminifera: Paleoceanographic Implications. *Paleoceanography* 3, 635-645.
 978 Bard, E., Arnold, M., Mangerud, J., Paterne, M., Labeyrie, L., Duprat, J., Mélières, M.-
 979 A., Sønstegaard, E., Duplessy, J.-C., 1994. The North Atlantic atmosphere-sea surface
 980 ^{14}C gradient during the Younger Dryas climatic event. *Earth and Planetary Science*
 981 *Letters* 126, 275-287.
 982 Barker, S., Chen, J., Gong, X., Jonkers, L., Knorr, G., Thornalley, D., 2015. Icebergs not
 983 the trigger for North Atlantic cold events. *Nature* 520, 333-336.
 984 Barker, S., Greaves, M., Elderfield, H., 2003. A study of cleaning procedures used for
 985 foraminiferal Mg/Ca paleothermometry. *Geochemistry, Geophysics, Geosystems* 4, 8407.
 986 Bauch, D., Bauch, H.A., 2001. Last glacial benthic foraminiferal $\delta^{18}\text{O}$ anomalies in the
 987 polar North Atlantic: A modern analogue evaluation. *Journal of Geophysical Research:*
 988 *Oceans* 106, 9135-9143.
 989 Bayon, G., German, C.R., Boella, R.M., Milton, J.A., Taylor, R.N., Nesbitt, R.W., 2002.
 990 An improved method for extracting marine sediment fractions and its application to Sr
 991 and Nd isotopic analysis. *Chemical Geology* 187, 179-199.
 992 Benway, H.M., McManus, J.F., Oppo, D.W., Cullen, J.L., 2010. Hydrographic changes in
 993 the eastern subpolar North Atlantic during the last deglaciation. *Quaternary Science*
 994 *Reviews* 29, 3336-3345.
 995 Bertram, C.J., Elderfield, H., Shackleton, N.J., MacDonald, J.A., 1995.
 996 Cadmium/Calcium and Carbon Isotope Reconstructions of the Glacial Northeast Atlantic
 997 Ocean. *Paleoceanography* 10, 563-578.

998 Bigg, G.R., Levine, R.C., Green, C.L., 2011. Modelling abrupt glacial North Atlantic
 999 freshening: Rates of change and their implications for Heinrich events. *Global and*
 1000 *Planetary Change* 79, 176-192.

1001 Blaser, P., Lippold, J., Gutjahr, M., Frank, N., Link, J.M., Frank, M., 2016. Extracting
 1002 foraminiferal Nd isotope signatures from bulk deep sea sediment by chemical leaching.
 1003 *Chemical Geology*.

1004 Böhm, E., Lippold, J., Gutjahr, M., Frank, M., Blaser, P., Antz, B., Fohlmeister, J., Frank,
 1005 N., Andersen, M.B., Deininger, M., 2015. Strong and deep Atlantic meridional
 1006 overturning circulation during the last glacial cycle. *Nature* 517, 73-76.

1007 Bond, G., Heinrich, H., Broecker, W., Labeyrie, L., McManus, J., Andrews, J., Huon, S.,
 1008 Jantschik, R., Clasen, S., Simet, C., 1992. Evidence for massive discharges of icebergs
 1009 into the North Atlantic ocean during the last glacial period. *Nature* 360, 245-249.

1010 Bondevik, S., Mangerud, J., Birks, H.H., Gulliksen, S., Reimer, P., 2006. Changes in
 1011 North Atlantic Radiocarbon Reservoir Ages During the Allerod and Younger Dryas.
 1012 *Science* 312, 1514-1517.

1013 Boyle, E.A., Keigwin, L., 1987. North Atlantic thermohaline circulation during the past
 1014 20,000 years linked to high-latitude surface temperature. *Nature* 330, 35-40.

1015 Boyle, E.A., Keigwin, L.D., 1985. Comparison of Atlantic and Pacific paleochemical
 1016 records for the last 215,000 years: changes in deep ocean circulation and chemical
 1017 inventories. *Earth and Planetary Science Letters* 76, 135-150.

1018 Bradtmiller, L.I., McManus, J.F., Robinson, L.F., 2014. $^{231}\text{Pa}/^{230}\text{Th}$ evidence for a
 1019 weakened but persistent Atlantic meridional overturning circulation during Heinrich
 1020 Stadial 1. *Nat Commun* 5.

1021 Broecker, S.W., Peng, T.-H., 1982. Tracers in the Sea. Eldigo Press, Palisades, NY.

1022 Broecker, W.S., Denton, G.H., 1989. The role of ocean-atmosphere reorganizations in
1023 glacial cycles. *Geochimica et Cosmochimica Acta* 53, 2465-2501.

1024 Burckel, P., Waelbroeck, C., Gherardi, J.M., Pichat, S., Arz, H., Lippold, J., Dokken, T.,
1025 Thil, F., 2015. Atlantic Ocean circulation changes preceded millennial tropical South
1026 America rainfall events during the last glacial. *Geophysical Research Letters*,
1027 2014GL062512.

1028 Clark, C.D., Hughes, A.L.C., Greenwood, S.L., Jordan, C., Sejrup, H.P., 2012. Pattern
1029 and timing of retreat of the last British-Irish Ice Sheet. *Quaternary Science Reviews* 44,
1030 112-146.

1031 Cohen, A.S., O'Nions, R.K., Siegenthaler, R., Griffin, W.L., 1988. Chronology of the
1032 pressure-temperature history recorded by a granulite terrain. *Contributions to Mineralogy
1033 and Petrology* 98, 303-311.

1034 Copard, K., Colin, C., Douville, E., Freiwald, A., Gudmundsson, G., De Mol, B., Frank,
1035 N., 2010. Nd isotopes in deep-sea corals in the North-eastern Atlantic. *Quaternary
1036 Science Reviews* 29, 2499-2508.

1037 Cottet-Puinel, M., Weaver, A.J., Hillaire-Marcel, C., de Vernal, A., Clark, P.U., Eby, M.,
1038 2004. Variation of Labrador Sea Water formation over the Last Glacial cycle in a climate
1039 model of intermediate complexity. *Quaternary Science Reviews* 23, 449-465.

1040 Crocker, A., 2014. Coupling of the Cryosphere and Ocean During Intervals of Rapid
1041 Climate Change
1042 in the Palaeo Record: A Multi-Proxy Study of the Heinrich Events of the Last

1043 Glacial from the Northeast Atlantic Department of Ocean and Earth Science. University
 1044 of Southampton, p. 268.
 1045 Crocket, K.C., Vance, D., Gutjahr, M., Foster, G.L., Richards, D.A., 2011. Persistent
 1046 Nordic deep-water overflow to the glacial North Atlantic. *Geology* 39, 515-518.
 1047 Curry, W.B., Oppo, D.W., 2005. Glacial water mass geometry and the distribution of
 1048 $\delta^{13}\text{C}$ of ΣCO_2 in the western Atlantic Ocean. *Paleoceanography* 20, PA1017.
 1049 Darling, K.F., Kucera, M., Kroon, D., Wade, C.M., 2006. A resolution for the coiling
 1050 direction paradox in *Neoglobobulimina pachyderma*. *Paleoceanography* 21, PA2011.
 1051 Davies, S.M., Abbott, P.M., Pearce, N.J.G., Wastegård, S., Blockley, S.P.E., 2012.
 1052 Integrating the INTIMATE records using tephrochronology: rising to the challenge.
 1053 *Quaternary Science Reviews* 36, 11-27.
 1054 de Baar, H.J.W., Bacon, M.P., Brewer, P.G., Bruland, K.W., 1985. Rare earth elements in
 1055 the Pacific and Atlantic Oceans. *Geochimica et Cosmochimica Acta* 49, 1943-1959.
 1056 de Baar, H.J.W., German, C.R., Elderfield, H., van Gaans, P., 1988. Rare earth element
 1057 distributions in anoxic waters of the Cariaco Trench. *Geochimica et Cosmochimica Acta*
 1058 52, 1203-1219.
 1059 de Vernal, A., Eynaud, F., Henry, M., Hillaire-Marcel, C., Londeix, L., Mangin, S.,
 1060 Matthiessen, J., Marret, F., Radi, T., Rochon, A., Solignac, S., Turon, J.L., 2005.
 1061 Reconstruction of sea-surface conditions at middle to high latitudes of the Northern
 1062 Hemisphere during the Last Glacial Maximum (LGM) based on dinoflagellate cyst
 1063 assemblages. *Quaternary Science Reviews* 24, 897-924.
 1064 Dokken, T.M., Jansen, E., 1999. Rapid changes in the mechanism of ocean convection
 1065 during the last glacial period. *Nature* 401, 458-461.

1066 Dokken, T.M., Nisancioglu, K.H., Li, C., Battisti, D.S., Kissel, C., 2013. Dansgaard-
 1067 Oeschger cycles: Interactions between ocean and sea ice intrinsic to the Nordic seas.
 1068 *Paleoceanography* 28, 491-502.
 1069 Duplessy, J.-C., Moyes, J., Pujol, C., 1980. Deep water formation in the North Atlantic
 1070 Ocean during the last ice age. *Nature* 286, 479-482.
 1071 Dyke, A.S., Andrews, J.T., Clark, P.U., England, J.H., Miller, G.H., Shaw, J., Veillette,
 1072 J.J., 2002. The Laurentide and Innuitian ice sheets during the Last Glacial Maximum.
 1073 *Quaternary Science Reviews* 21, 9-31.
 1074 Elderfield, H., Greaves, M.J., 1982. The rare earth elements in seawater. *Nature* 296,
 1075 214-219.
 1076 Elderfield, H., Whitfield, M., Burton, J.D., Bacon, M.P., Liss, P.S., 1988. The Oceanic
 1077 Chemistry of the Rare-Earth Elements. *Philosophical Transactions of the Royal Society*
 1078 of London. Series A, Mathematical and Physical Sciences 325, 105-126.
 1079 Ellett, D.J., Martin, J.H.A., 1973. The physical and chemical oceanography of the
 1080 Rockall channel. *Deep Sea Research and Oceanographic Abstracts* 20, 585-625.
 1081 Elmore, A.C., Piotrowski, A.M., Wright, J.D., Scrivner, A.E., 2011. Testing the
 1082 extraction of past seawater Nd isotopic composition from North Atlantic deep sea
 1083 sediments and foraminifera. *Geochem. Geophys. Geosyst.* 12, Q09008.
 1084 Eynaud, F., de Abreu, L.c., Voelker, A., Sch^nfeld, J., Salgueiro, E., Turon, J.-L., Penaud,
 1085 A., Toucanne, S., Naughton, F., S-nchez GoÒi, M.F., MalaizÈ, B., Cacho, I., 2009.
 1086 Position of the Polar Front along the western Iberian margin during key cold episodes of
 1087 the last 45 ka. *Geochem. Geophys. Geosyst.* 10, Q07U05.

1088 Eynaud, F., Zaragosi, S., Scourse, J.D., Mojtahid, M., Bourillet, J.F., Hall, I.R., Penaud,
 1089 A., Locascio, M., Reijonen, A., 2007. Deglacial laminated facies on the NW European
 1090 continental margin: The hydrographic significance of British-Irish Ice Sheet deglaciation
 1091 and Fleuve Manche paleoriver discharges. *Geochem. Geophys. Geosyst.* 8, Q06019.
 1092 Flückiger, J., Knutti, R., White, J.C., Renssen, H., 2008. Modeled seasonality of glacial
 1093 abrupt climate events. *Climate Dynamics* 31, 633-645.
 1094 Funder, S., Hansen, L., 1996. The Greenland ice sheet - a model for its culmination and
 1095 decay during and after the last glacial maximum. *Bulletin of the Geological Society of*
 1096 *Denmark* 42, 137-152.
 1097 Garrison, T., 2011. *Essentials of oceanography*. Thomson Brooks/Cole, Cengage Learning,
 1098 Inc.
 1099 German, C.R., Elderfield, H., 1990. Application of the Ce anomaly as a paleoredox
 1100 indicator: The ground rules. *Paleoceanography* 5, 823-833.
 1101 Gherardi, J.-M., Labeyrie, L., Nave, S., Francois, R., McManus, J.F., Cortijo, E., 2009.
 1102 Glacial-interglacial circulation changes inferred from $^{231}\text{Pa}/^{230}\text{Th}$ sedimentary record in
 1103 the North Atlantic region. *Paleoceanography* 24, PA2204.
 1104 Gierz, P., Lohmann, G., Wei, W., 2015. Response of Atlantic overturning to future
 1105 warming in a coupled atmosphere-ocean-ice sheet model. *Geophysical Research Letters*
 1106 42, 6811-6818.
 1107 Goldberg, E.D., Koide, M., Schmitt, R.A., Smith, R.H., 1963. Rare-Earth distributions in
 1108 the marine environment. *Journal of Geophysical Research* 68, 4209-4217.

1109 Goldstein, S.L., O'Nions, R.K., Hamilton, P.J., 1984. A Sm-Nd isotopic study of
 1110 atmospheric dusts and particulates from major river systems. *Earth and Planetary Science*
 1111 *Letters* 70, 221-236.

1112 Grandjean, P., Cappetta, H., Michard, A., Albarède, F., 1987. The assessment of REE
 1113 patterns and $^{143}\text{Nd}/^{144}\text{Nd}$ ratios in fish remains. *Earth and Planetary Science Letters* 84,
 1114 181-196.

1115 Grant, K.M., Rohling, E.J., Bar-Matthews, M., Ayalon, A., Medina-Elizalde, M.,
 1116 Ramsey, C.B., Satow, C., Roberts, A.P., 2012. Rapid coupling between ice volume and
 1117 polar temperature over the past 150,000 years. *Nature* 491, 744-747.

1118 Grousset, F.E., Biscaye, P.E., Revel, M., Petit, J.-R., Pye, K., Joussaume, S., Jouzel, J.,
 1119 1992. Antarctic (Dome C) ice-core dust at 18 k.y. B.P.: Isotopic constraints on origins.
 1120 *Earth and Planetary Science Letters* 111, 175-182.

1121 Grousset, F.E., Labeyrie, L., Sinko, J.A., Cremer, M., Bond, G., Duprat, J., Cortijo, E.,
 1122 Huon, S., 1993. Patterns of Ice-Rafted Detritus in the Glacial North Atlantic (40-55°N).
 1123 *Paleoceanography* 8, 175-192.

1124 Gutjahr, M., Frank, M., Stirling, C.H., Klemm, V., van de Flierdt, T., Halliday, A.N.,
 1125 2007. Reliable extraction of a deepwater trace metal isotope signal from Fe-Mn
 1126 oxyhydroxide coatings of marine sediments. *Chemical Geology* 242, 351-370.

1127 Gutjahr, M., Lippold, J., 2011. Early arrival of Southern Source Water in the deep North
 1128 Atlantic prior to Heinrich event 2. *Paleoceanography* 26, PA2101.

1129 Hall, I.R., Colmenero-Hidalgo, E., Zahn, R., Peck, V.L., Hemming, S.R., 2011.
 1130 Centennial- to millennial-scale ice-ocean interactions in the subpolar northeast Atlantic
 1131 18-41 kyr ago. *Paleoceanography* 26, PA2224.

1132 Hansen, B., Østerhus, S., 2000. North Atlantic-Nordic Seas exchanges. Progress In
 1133 Oceanography 45, 109-208.
 1134 Heinrich, H., 1988. Origin and consequences of cyclic ice rafting in the Northeast
 1135 Atlantic Ocean during the past 130,000 years. Quaternary Research 29, 142-152.
 1136 Hemming, S.R., 2004. Heinrich events: Massive late Pleistocene detritus layers of the
 1137 North Atlantic and their global climate imprint. Rev. Geophys. 42, RG1005.
 1138 Hillaire-Marcel, C., de Vernal, A., 2008. Stable isotope clue to episodic sea ice formation
 1139 in the glacial North Atlantic. Earth and Planetary Science Letters 268, 143-150.
 1140 Hillaire-Marcel, C., de Vernal, A., Candon, L., Bilodeau, G., Stoner, J., 2001. Changes of
 1141 potential density gradients in the northwestern North Atlantic during the last climatic
 1142 cycle based on a multiproxy approach, The Oceans and Rapid Climate Change: Past,
 1143 Present, and Future. AGU, Washington, DC, pp. 83-100.
 1144 Hodell, D.A., Curtis, J.H., 2008. Oxygen and carbon isotopes of detrital carbonate in
 1145 North Atlantic Heinrich Events. Marine Geology 256, 30-35.
 1146 Holser, W.T., 1997. Evaluation of the application of rare-earth elements to
 1147 paleoceanography. Palaeogeography, Palaeoclimatology, Palaeoecology 132, 309-323.
 1148 Innocent, C., Fagel, N., Stevenson, R.K., Hillaire-Marcel, C., 1997. Sm–Nd signature of
 1149 modern and late Quaternary sediments from the northwest North Atlantic: Implications
 1150 for deep current changes since the Last Glacial Maximum. Earth and Planetary Science
 1151 Letters 146, 607-625.
 1152 Jaccard, S.L., Galbraith, E.D., 2012. Large climate-driven changes of oceanic oxygen
 1153 concentrations during the last deglaciation. Nature Geosci 5, 151-156.

1154 Jacobsen, S.B., Wasserburg, G.J., 1980. Sm-Nd isotopic evolution of chondrites. *Earth*
 1155 *and Planetary Science Letters* 50, 139-155.

1156 Jansen, E., Fronval, T., Rack, F., Channell, J.E.T., 2000. Pliocene-Pleistocene Ice Rafting
 1157 History and Cyclicity in the Nordic Seas During the Last 3.5 Myr. *Paleoceanography* 15,
 1158 709-721.

1159 Jeandel, C., Arsouze, T., Lacan, F., Téchiné, P., Dutay, J.C., 2007. Isotopic Nd
 1160 compositions and concentrations of the lithogenic inputs into the ocean: A compilation,
 1161 with an emphasis on the margins. *Chemical Geology* 239, 156-164.

1162 Johnson, C., Sherwin, T., Smythe-Wright, D., Shimmield, T., Turrell, W., 2010. Wyville
 1163 Thomson Ridge Overflow Water: Spatial and temporal distribution in the Rockall
 1164 Trough. *Deep Sea Research Part I: Oceanographic Research Papers* 57, 1153-1162.

1165 Jones, C.E., Halliday, A.N., Rea, D.K., Owen, R.M., 1994. Neodymium isotopic
 1166 variations in North Pacific modern silicate sediment and the insignificance of detrital
 1167 REE contributions to seawater. *Earth and Planetary Science Letters* 127, 55-66.

1168 Jonkers, L., Zahn, R., Thomas, A., Henderson, G., Abouchami, W., François, R.,
 1169 Masque, P., Hall, I.R., Bickert, T., 2015. Deep circulation changes in the central South
 1170 Atlantic during the past 145 kyrs reflected in a combined $^{231}\text{Pa}/^{230}\text{Th}$, Neodymium
 1171 isotope and benthic record. *Earth and Planetary Science Letters* 419, 14-21.

1172 Keigwin, L.D., Jones, G.A., Lehman, S.J., Boyle, E.A., 1991. Deglacial meltwater
 1173 discharge, North Atlantic Deep Circulation, and abrupt climate change. *Journal of*
 1174 *Geophysical Research: Oceans* 96, 16811-16826.

1175 Key, R.M., Kozyr, A., Sabine, C.L., Lee, K., Wanninkhof, R., Bullister, J.L., Feely, R.A.,
 1176 Millero, F.J., Mordy, C., Peng, T.H., 2004. A global ocean carbon climatology: Results

1177 from Global Data Analysis Project (GLODAP). *Global Biogeochemical Cycles* 18,
 1178 GB4031.

1179 Kissel, C., 2005. Magnetic signature of rapid climatic variations in glacial North Atlantic,
 1180 a review. *Comptes Rendus Geoscience* 337, 908-918.

1181 Kissel, C., Laj, C., Labeyrie, L., Dokken, T., Voelker, A., Blamart, D., 1999. Rapid
 1182 climatic variations during marine isotopic stage 3: magnetic analysis of sediments from
 1183 Nordic Seas and North Atlantic. *Earth and Planetary Science Letters* 171, 489-502.

1184 Labeyrie, L.D., Duplessy, J.-C., Duprat, J., Juillet-Leclerc, A., Moyes, J., Michel, E.,
 1185 Kallel, N., Shackleton, N.J., 1992. Changes in the vertical structure of the North Atlantic
 1186 Ocean between glacial and modern times. *Quaternary Science Reviews* 11, 401-413.

1187 Lacan, F., Jeandel, C., 2004a. Denmark Strait water circulation traced by heterogeneity in
 1188 neodymium isotopic compositions. *Deep Sea Research Part I: Oceanographic Research*
 1189 *Papers* 51, 71-82.

1190 Lacan, F., Jeandel, C., 2004b. Neodymium isotopic composition and rare earth element
 1191 concentrations in the deep and intermediate Nordic Seas: Constraints on the Iceland
 1192 Scotland Overflow Water signature. *Geochem. Geophys. Geosyst.* 5, Q11006.

1193 Lacan, F., Jeandel, C., 2004c. Subpolar Mode Water formation traced by neodymium
 1194 isotopic composition. *Geophys. Res. Lett.* 31, L14306.

1195 Lacan, F., Jeandel, C., 2005. Acquisition of the neodymium isotopic composition of the
 1196 North Atlantic Deep Water. *Geochem. Geophys. Geosyst.* 6, Q12008.

1197 Lacan, F., Tachikawa, K., Jeandel, C., 2012. Neodymium isotopic composition of the
 1198 oceans: a compilation of seawater data. *Chemical Geology* 300-301, 177-184.

1199 Lambelet, M., van de Flierdt, T., Crocket, K., Rehkämper, M., Kreissig, K., Coles, B.,
 1200 Rijkenberg, M.J.A., Gerringa, L.J.A., de Baar, H.J.W., Steinfeldt, R., 2016. Neodymium
 1201 isotopic composition and concentration in the western North Atlantic Ocean: results from
 1202 the GEOTRACES GA02 section. *Geochimica et Cosmochimica Acta*.
 1203 Lang, D.C., Bailey, I., Wilson, P.A., Chalk, T.B., Foster, G.L., Gutjahr, M., 2016.
 1204 Incursions of southern-sourced water into the deep North Atlantic during late Pliocene
 1205 glacial intensification. *Nature Geosci* 9, 375-379.
 1206 Lekens, W.A.H., Sejrup, H.P., Haflidason, H., Knies, J., Richter, T., 2006. Meltwater and
 1207 ice rafting in the southern Norwegian Sea between 20 and 40 calendar kyr B.P.:
 1208 Implications for Fennoscandian Heinrich events. *Paleoceanography* 21, PA3013.
 1209 Lippold, J., Grützner, J., Winter, D., Lahaye, Y., Mangini, A., Christl, M., 2009. Does
 1210 sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ from the Bermuda Rise monitor past Atlantic Meridional
 1211 Overturning Circulation? *Geophysical Research Letters* 36, L12601.
 1212 Lippold, J., Gutjahr, M., Blaser, P., Christner, E., de Carvalho Ferreira, M.L., Mulitza, S.,
 1213 Christl, M., Wombacher, F., Böhm, E., Antz, B., Cartapanis, O., Vogel, H., Jaccard, S.L.,
 1214 2016. Deep water provenance and dynamics of the (de)glacial Atlantic meridional
 1215 overturning circulation. *Earth and Planetary Science Letters* 445, 68-78.
 1216 Lowe, J.J., Rasmussen, S.O., Björck, S., Hoek, W.Z., Steffensen, J.P., Walker, M.J.C.,
 1217 Yu, Z.C., 2008. Synchronisation of palaeoenvironmental events in the North Atlantic
 1218 region during the Last Termination: a revised protocol recommended by the INTIMATE
 1219 group. *Quaternary Science Reviews* 27, 6-17.

1220 Lynch-Stieglitz, J., Schmidt, M.W., Gene Henry, L., Curry, W.B., Skinner, L.C., Mulitza,
 1221 S., Zhang, R., Chang, P., 2014. Muted change in Atlantic overturning circulation over
 1222 some glacial-aged Heinrich events. *Nature Geosci* 7, 144-150.

1223 Mackensen, A., Hubberten, H.W., Bickert, T., Fischer, G., Fütterer, D.K., 1993. The
 1224 $\delta^{13}\text{C}$ in benthic foraminiferal tests of *Fontbotia wuellerstorfi* (Schwager) Relative to the
 1225 $\delta^{13}\text{C}$ of dissolved inorganic carbon in Southern Ocean Deep Water: Implications for
 1226 glacial ocean circulation models. *Paleoceanography* 8, 587-610.

1227 MacLeod, K.G., Irving, A.J., 1996. Correlation of cerium anomalies with indicators of
 1228 paleoenvironment. *Journal of Sedimentary Research* 66, 948-955.

1229 Manighetti, B., McCave, I.N., 1995. Late Glacial and Holocene Palaeocurrents Around
 1230 Rockall Bank, NE Atlantic Ocean. *Paleoceanography* 10, 611-626.

1231 Martin, E.E., Blair, S.W., Kamenov, G.D., Scher, H.D., Bourbon, E., Basak, C., Newkirk,
 1232 D.N., 2010. Extraction of Nd isotopes from bulk deep sea sediments for
 1233 paleoceanographic studies on Cenozoic time scales. *Chemical Geology* 269, 414-431.

1234 Martin, E.E., Haley, B.A., 2000. Fossil fish teeth as proxies for seawater Sr and Nd
 1235 isotopes. *Geochimica et Cosmochimica Acta* 64, 835-847.

1236 Martin, E.E., Scher, H.D., 2004. Preservation of seawater Sr and Nd isotopes in fossil
 1237 fish teeth: bad news and good news. *Earth and Planetary Science Letters* 220, 25-39.

1238 Maslin, M.A., Shackleton, N.J., Pflaumann, U., 1995. Surface Water Temperature,
 1239 Salinity, and Density Changes in the Northeast Atlantic During the Last 45,000 Years:
 1240 Heinrich Events, Deep Water Formation, and Climatic Rebounds. *Paleoceanography* 10,
 1241 527-544.

1242 McCartney, M.S., 1992. Recirculating components to the deep boundary current of the
 1243 northern North Atlantic. *Progress In Oceanography* 29, 283-383.
 1244 McCave, I.N., Manighetti, B., Beveridge, N.A.S., 1995a. Circulation in the glacial North
 1245 Atlantic inferred from grain-size measurements. *Nature* 374, 149-152.
 1246 McCave, I.N., Manighetti, B., Robinson, S.G., 1995b. Sortable silt and fine sediment
 1247 size/composition slicing: Parameters for palaeocurrent speed and palaeoceanography.
 1248 *Paleoceanography* 10, 593-610.
 1249 McGrath, T., Nolan, G., McGovern, E., 2012. Chemical characteristics of water masses
 1250 in the Rockall Trough. *Deep Sea Research Part I: Oceanographic Research Papers*.
 1251 McIntyre, K.L., Howe, J.A., 2009. Bottom-current variability during the last glacial-
 1252 deglacial transition, Northern Rockall Trough and Faroe Bank Channel, NE Atlantic.
 1253 *Scottish Journal of Geology* 45, 43-57.
 1254 McLennan, S.M., 1989. Rare earth elements in sedimentary rocks; influence of
 1255 provenance and sedimentary processes. *Reviews in Mineralogy and Geochemistry* 21,
 1256 169-200.
 1257 McManus, J.F., Francois, R., Gherardi, J.M., Keigwin, L.D., Brown-Leger, S., 2004.
 1258 Collapse and rapid resumption of Atlantic meridional circulation linked to deglacial
 1259 climate changes. *Nature* 428, 834-837.
 1260 McManus, J.F., Oppo, D.W., Cullen, J.L., 1999. A 0.5-Million-Year Record of
 1261 Millennial-Scale Climate Variability in the North Atlantic. *Science* 283, 971-975.
 1262 Mearns, E.W., 1988. A samarium-neodymium isotopic survey of modern river sediments
 1263 from Northern Britain. *Chemical Geology: Isotope Geoscience section* 73, 1-13.

1264 Meland, M.Y., Dokken, T.M., Jansen, E., Hevrøy, K., 2008. Water mass properties and
 1265 exchange between the Nordic seas and the northern North Atlantic during the period 23-6
 1266 ka: Benthic oxygen isotopic evidence. *Paleoceanography* 23, PA1210.
 1267 Mulitza, S., Prange, M., Stuut, J.-B., Zabel, M., von Dobeneck, T., Itambi, A.C., Nizou,
 1268 J., Schulz, M., Wefer, G., 2008. Sahel megadroughts triggered by glacial slowdowns of
 1269 Atlantic meridional overturning. *Paleoceanography* 23, PA4206.
 1270 New, A.L., Smythe-Wright, D., 2001. Aspects of the circulation in the Rockall Trough.
 1271 *Continental Shelf Research* 21, 777-810.
 1272 North Greenland Ice Core Project Members, Andersen, K.K., Azuma, N., Barnola, J.-M.,
 1273 Bigler, M., Biscaye, P., Caillon, N., Chappellaz, J., Clausen, H.B., Dahl-Jensen, D.,
 1274 Fischer, H., Flückiger, J., Fritzsche, D., Fujii, Y., Goto-Azuma, K., Grønvold, K.,
 1275 Gundestrup, N.S., Hansson, M., Huber, C., Hvidberg, C.S., Johnsen, S.J., Jonsell, U.,
 1276 Jouzel, J., Kipfstuhl, S., Landais, A., Leuenberger, M., Lorrain, R., Masson-Delmotte, V.,
 1277 Miller, H., Motoyama, H., Narita, H., Popp, T., Rasmussen, S.O., Raynaud, D.,
 1278 Rothlisberger, R., Ruth, U., Samyn, D., Schwander, J., Shoji, H., Siggard-Andersen, M.-
 1279 L., Steffensen, J.P., Stocker, T., Sveinbjörnsdóttir, A.E., Svensson, A., Takata, M., Tison,
 1280 J.-L., Thorsteinsson, T., Watanabe, O., Wilhelms, F., White, J.W.C., 2004. High-
 1281 resolution record of Northern Hemisphere climate extending into the last interglacial
 1282 period. *Nature* 431, 147-151.
 1283 Nuttin, L., Maccali, J., Hillaire-Marcel, C., 2015. U, Th and Pa insights into
 1284 sedimentological and paleoceanographic changes off Hudson Strait (Labrador Sea)
 1285 during the last 37-ka with special attention to methodological issues. *Quaternary*
 1286 *Science Reviews* 115, 39-49.

1287 O'Nions, R.K., Carter, S.R., Cohen, R.S., Evensen, N.M., Hamilton, P.J., 1978. Pb, Nd
 1288 and Sr isotopes in oceanic ferromanganese deposits and ocean floor basalts. *Nature* 273,
 1289 435-438.

1290 Olsen, A., Ninnemann, U., 2010. Large $\delta^{13}\text{C}$ Gradients in the Preindustrial North
 1291 Atlantic Revealed. *Science* 330, 658-659.

1292 Oppo, D.W., Lehman, S.J., 1993. Mid-Depth Circulation of the Subpolar North Atlantic
 1293 During the Last Glacial Maximum. *Science* 259, 1148-1152.

1294 Oppo, D.W., McManus, J.F., Cullen, J.L., 2003. Palaeo-oceanography: Deepwater
 1295 variability in the Holocene epoch. *Nature* 422, 277-277.

1296 Oppo, D.W., McManus, J.F., Cullen, J.L., 2006. Evolution and demise of the Last
 1297 Interglacial warmth in the subpolar North Atlantic. *Quaternary Science Reviews* 25,
 1298 3268-3277.

1299 Otto-Bliesner, B.L., Brady, E.C., 2010. The sensitivity of the climate response to the
 1300 magnitude and location of freshwater forcing: last glacial maximum experiments.
 1301 *Quaternary Science Reviews* 29, 56-73.

1302 Paillard, D., Labeyrie, L., Yiou, P., 1996. Macintosh program performs time-series
 1303 analysis. *Eos. Trans. AGU* 77.

1304 Palmer, M.R., Elderfield, H., 1985. Variations in the Nd isotopic composition of
 1305 foraminifera from Atlantic Ocean sediments. *Earth and Planetary Science Letters* 73,
 1306 299-305.

1307 Peck, V.L., Hall, I.R., Zahn, R., Grousset, F., Hemming, S.R., Scourse, J.D., 2007. The
 1308 relationship of Heinrich events and their European precursors over the past 60 ka BP: a

1309 multi-proxy ice-rafted debris provenance study in the North East Atlantic. *Quaternary*
 1310 *Science Reviews* 26, 862-875.

1311 Piepgras, D.J., Wasserburg, G.J., Dasch, E.J., 1979. The isotopic composition of Nd in
 1312 different ocean masses. *Earth and Planetary Science Letters* 45, 223-236.

1313 Pin, C., Zalduegui, J.S., 1997. Sequential separation of light rare-earth elements, thorium
 1314 and uranium by miniaturized extraction chromatography: Application to isotopic analyses
 1315 of silicate rocks. *Analytica Chimica Acta* 339, 79-89.

1316 Piotrowski, A.M., Galy, A., Nicholl, J.A.L., Roberts, N., Wilson, D.J., Clegg, J.A., Yu,
 1317 J., 2012. Reconstructing deglacial North and South Atlantic deep water sourcing using
 1318 foraminiferal Nd isotopes. *Earth and Planetary Science Letters* 357-358, 289-297.

1319 Piotrowski, A.M., Goldstein, S.L., Hemming, S.R., Fairbanks, R.G., 2004. Intensification
 1320 and variability of ocean thermohaline circulation through the last deglaciation. *Earth and*
 1321 *Planetary Science Letters* 225, 205-220.

1322 Piotrowski, A.M., Goldstein, S.L., Hemming, S.R., Fairbanks, R.G., 2005. Temporal
 1323 Relationships of Carbon Cycling and Ocean Circulation at Glacial Boundaries. *Science*
 1324 307, 1933-1938.

1325 Piotrowski, A.M., Goldstein, S.L., Hemming, S.R., Fairbanks, R.G., Zylberberg, D.R.,
 1326 2008. Oscillating glacial northern and southern deep water formation from combined
 1327 neodymium and carbon isotopes. *Earth and Planetary Science Letters* 272, 394–405.

1328 Rae, J.W.B., Foster, G.L., Schmidt, D.N., Elliott, T., 2011. Boron isotopes and B/Ca in
 1329 benthic foraminifera: Proxies for the deep ocean carbonate system. *Earth and Planetary*
 1330 *Science Letters* 302, 403-413.

1331 Rahmstorf, S., 2002. Ocean circulation and climate during the past 120,000 years. *Nature*
 1332 419, 207-214.

1333 Rasmussen, S.O., Andersen, K.K., Svensson, A.M., Steffensen, J.P., Vinther, B.M.,
 1334 Clausen, H.B., Siggaard-Andersen, M.L., Johnsen, S.J., Larsen, L.B., Dahl-Jensen, D.,
 1335 Bigler, M., Röthlisberger, R., Fischer, H., Goto-Azuma, K., Hansson, M.E., Ruth, U.,
 1336 2006. A new Greenland ice core chronology for the last glacial termination. *J. Geophys.*
 1337 *Res.* 111, D06102.

1338 Rasmussen, T.L., Thomsen, E., 2009. Stable isotope signals from brines in the Barents
 1339 Sea: Implications for brine formation during the last glaciation. *Geology* 37, 903-906.

1340 Reimer, P.J., Bard, E., Bayliss, A., Beck, J.W., Blackwell, P.G., Bronk Ramsey, C.,
 1341 Buck, C.E., Cheng, H., Edwards, R.L., Friedrich, M., Grootes, P.M., Guilderson, T.P.,
 1342 Haflidason, H., Hajdas, I., Hatté, C., Heaton, T.J., Hoffmann, D.L., Hogg, A.G., Hughen,
 1343 K.A., Kaiser, K.F., Kromer, B., Manning, S.W., Niu, M., Reimer, R.W., Richards, D.A.,
 1344 Scott, E.M., Southon, J.R., Staff, R.A., Turney, C.S.M., van der Plicht, J., 2013. IntCal13
 1345 and Marine13 Radiocarbon Age Calibration Curves 0–50,000 Years cal BP.

1346 Reynard, B., Lécuyer, C., Grandjean, P., 1999. Crystal-chemical controls on rare-earth
 1347 element concentrations in fossil biogenic apatites and implications for
 1348 paleoenvironmental reconstructions. *Chemical Geology* 155, 233-241.

1349 Rickli, J.r., Gutjahr, M., Vance, D., Fischer-Gödde, M., Hillenbrand, C.-D., Kuhn, G.,
 1350 2014. Neodymium and hafnium boundary contributions to seawater along the West
 1351 Antarctic continental margin. *Earth and Planetary Science Letters* 394, 99-110.

1352 Roberts, N.L., Piotrowski, A.M., 2015. Radiogenic Nd isotope labeling of the northern
 1353 NE Atlantic during MIS 2. *Earth and Planetary Science Letters* 423, 125-133.

1354 Roberts, N.L., Piotrowski, A.M., Elderfield, H., Eglinton, T.I., Lomas, M.W., 2012. Rare
 1355 earth element association with foraminifera. *Geochimica et Cosmochimica Acta* 94, 57-
 1356 71.

1357 Roberts, N.L., Piotrowski, A.M., McManus, J.F., Keigwin, L.D., 2010. Synchronous
 1358 Deglacial Overturning and Water Mass Source Changes. *Science* 327, 75-78.

1359 Robinson, L.F., Adkins, J.F., Keigwin, L.D., Southon, J., Fernandez, D.P., Wang, S.L.,
 1360 Scheirer, D.S., 2005. Radiocarbon Variability in the Western North Atlantic During the
 1361 Last Deglaciation. *Science* 310, 1469-1473.

1362 Rogerson, M., Rohling, E.J., Weaver, P.P.E., Murray, J.W., 2005. Glacial to interglacial
 1363 changes in the settling depth of the Mediterranean Outflow plume. *Paleoceanography* 20,
 1364 n/a-n/a.

1365 Ruddiman, W.F., 1977. Late Quaternary deposition of ice-rafted sand in the subpolar
 1366 North Atlantic (lat 40° to 65°N). *Geological Society of America Bulletin* 88, 1813-1827.

1367 Rutberg, R.L., Hemming, S.R., Goldstein, S.L., 2000. Reduced North Atlantic Deep
 1368 Water flux to the glacial Southern Ocean inferred from neodymium isotope ratios. *Nature*
 1369 405, 935-938.

1370 Sarbas, B., Nohl, U., 2008. The GEOROC database as part of a growing geoinformatics
 1371 network, 2008 Geoinformatics conference.

1372 Sarmiento, J.L., Gruber, N., 2004. *Ocean Biogeochemical Dynamics*. Princeton
 1373 University Press, Princeton, NJ.

1374 Sarthein, M., Winn, K., Jung, S.J.A., Duplessy, J.-C., Labeyrie, L., Erlenkeuser, H.,
 1375 Ganssen, G., 1994. Changes in East Atlantic Deepwater Circulation Over the Last 30,000
 1376 years: Eight Time Slice Reconstructions. *Paleoceanography* 9, 209-267.

1377 Schönfeld, J., Zahn, R., de Abreu, L., 2003. Surface and deep water response to rapid
 1378 climate changes at the Western Iberian Margin. *Global and Planetary Change* 36, 237-
 1379 264.

1380 Sejrup, H.P., Hjelstuen, B.O., Torbjørn Dahlgren, K.I., Haflidason, H., Kuijpers, A.,
 1381 Nygård, A., Praeg, D., Stoker, M.S., Vorren, T.O., 2005. Pleistocene glacial history of the
 1382 NW European continental margin. *Marine and Petroleum Geology* 22, 1111-1129.

1383 Shackleton, N.J., Opdyke, N.D., 1973. Oxygen isotope and palaeomagnetic stratigraphy
 1384 of Equatorial Pacific core V28-238: Oxygen isotope temperatures and ice volumes on a
 1385 10^5 year and 10^6 year scale. *Quaternary Research* 3, 39-55.

1386 Shipboard Scientific Party, 1996. Sites 980/981, in: Jansen, E., Raymo, M.E., Blum P. et
 1387 al. (Eds.), *Proceedings of the Ocean Drilling Program, Initial Reports*. College station,
 1388 TX, pp. 49-90.

1389 Sholkovitz, E.R., 1988. Rare earth elements in the sediments of the North Atlantic Ocean,
 1390 Amazon Delta, and East China Sea; reinterpretation of terrigenous input patterns to the
 1391 oceans. *American Journal of Science* 288, 236-281.

1392 Siddall, M., Rohling, E.J., Almogi-Labin, A., Hemleben, C., Meischner, D., Schmelzer,
 1393 I., Smeed, D.A., 2003. Sea-level fluctuations during the last glacial cycle. *Nature* 423,
 1394 853-858.

1395 Sigman, D.M., Boyle, E.A., 2000. Glacial/interglacial variations in atmospheric carbon
 1396 dioxide. *Nature* 407, 859-869.

1397 Sigman, D.M., Hain, M.P., Haug, G.H., 2010. The polar ocean and glacial cycles in
 1398 atmospheric CO₂ concentration. *Nature* 466, 47-55.

1399 Singarayer, J.S., Valdes, P.J., 2010. High-latitude climate sensitivity to ice-sheet forcing
 1400 over the last 120 kyr. *Quaternary Science Reviews* 29, 43-55.
 1401 Smith, R.S., Gregory, J.M., 2009. A study of the sensitivity of ocean overturning
 1402 circulation and climate to freshwater input in different regions of the North Atlantic.
 1403 *Geophysical Research Letters* 36, L15701.
 1404 Snoeckx, H., Grousset, F., Revel, M., Boelaert, A., 1999. European contribution of ice-
 1405 rafted sand to Heinrich layers H3 and H4. *Marine Geology* 158, 197-208.
 1406 Spivack, A.J., Wasserburg, G.J., 1988. Neodymium isotopic composition of the
 1407 Mediterranean outflow and the eastern North Atlantic. *Geochimica et Cosmochimica*
 1408 *Acta* 52, 2767-2773.
 1409 Stanford, J.D., Rohling, E.J., Bacon, S., Roberts, A.P., Grousset, F.E., Bolshaw, M.,
 1410 2011. A new concept for the paleoceanographic evolution of Heinrich event 1 in the
 1411 North Atlantic. *Quaternary Science Reviews* 30, 1047-1066.
 1412 Staudigel, H., Doyle, P., Zindler, A., 1985. Sr and Nd isotope systematics in fish teeth.
 1413 *Earth and Planetary Science Letters* 76, 45-56.
 1414 Stern, J.V., Lisiecki, L.E., 2013. North Atlantic circulation and reservoir age changes
 1415 over the past 41,000 years. *Geophysical Research Letters* 40, 3693-3697.
 1416 Stichel, T., Frank, M., Rickli, J., Haley, B.A., 2012. The hafnium and neodymium isotope
 1417 composition of seawater in the Atlantic sector of the Southern Ocean. *Earth and Planetary*
 1418 *Science Letters* 317, 282-294.
 1419 Stille, P., 1992. Nd-Sr isotope evidence for dramatic changes of paleocurrents in the
 1420 Atlantic Ocean during the past 80 m.y. *Geology* 20, 387-390.

1421 Stille, P., Fischer, H., 1990. Secular variation in the isotopic composition of Nd in Tethys
 1422 seawater. *Geochimica et Cosmochimica Acta* 54, 3139-3145.
 1423 Stuiver, M., Reimer, P.J., 1993. Extended ^{14}C database and revised CALIB radiocarbon
 1424 calibration program. *Radiocarbon* 35, 215-230.
 1425 Stuiver, M., Reimer, P.J., Reimer, R.W., 2005. CALIB 5.0.
 1426 Svensson, A., Andersen, K.K., Bigler, M., Clausen, H.B., Dahl-Jensen, D., Davies, S.M.,
 1427 Johnsen, S.J., Muscheler, R., Parrenin, F., Rasmussen, S.O., Röthlisberger, R., Seierstad,
 1428 I., Steffensen, J.P., Vinther, B.M., 2008. A 60 000 year Greenland stratigraphic ice core
 1429 chronology. *Clim. Past* 4, 47-57.
 1430 Swingedouw, D., Mignot, J., Braconnot, P., Mosquet, E., Kageyama, M., Alkama, R.,
 1431 2009. Impact of Freshwater Release in the North Atlantic under Different Climate
 1432 Conditions in an OAGCM. *Journal of Climate* 22, 6377-6403.
 1433 Tachikawa, K., Athias, V., Jeandel, C., 2003. Neodymium budget in the modern ocean
 1434 and paleo-oceanographic implications. *J. Geophys. Res.* 108, 3254.
 1435 Tachikawa, K., Jeandel, C., Roy-Barman, M., 1999. A new approach to the Nd residence
 1436 time in the ocean: the role of atmospheric inputs. *Earth and Planetary Science Letters*
 1437 170, 433-446.
 1438 Tachikawa, K., Piotrowski, A.M., Bayon, G., 2014. Neodymium associated with
 1439 foraminiferal carbonate as a recorder of seawater isotopic signatures. *Quaternary Science*
 1440 *Reviews* 88, 1-13.
 1441 Tanaka, T., Togashi, S., Kamioka, H., Amakawa, H., Kagami, H., Hamamoto, T.,
 1442 Yuhara, M., Orihashi, Y., Yoneda, S., Shimizu, H., Kunimaru, T., Takahashi, K., Yanagi,
 1443 T., Nakano, T., Fujimaki, H., Shinjo, R., Asahara, Y., Tanimizu, M., Dragusanu, C.,

1444 2000. JNdi-1: a neodymium isotopic reference in consistency with LaJolla neodymium.
 1445 Chemical Geology 168, 279-281.
 1446 Taylor, S.R., McLennan, S.M., 1985. The continental crust: its composition and
 1447 evolution. Blackwell, Oxford.
 1448 Thornalley, D.J.R., Barker, S., Becker, J., Hall, I.R., Knorr, G., 2013. Abrupt changes in
 1449 deep Atlantic circulation during the transition to full glacial conditions.
 1450 Paleoceanography 28, 253-262.
 1451 Thornalley, D.J.R., Bauch, H.A., Gebbie, G., Guo, W., Ziegler, M., Bernasconi, S.M.,
 1452 Barker, S., Skinner, L.C., Yu, J., 2015. A warm and poorly ventilated deep Arctic
 1453 Mediterranean during the last glacial period. Science 349, 706-710.
 1454 Thornalley, D.J.R., Elderfield, H., McCave, I.N., 2010. Intermediate and deep water
 1455 paleoceanography of the northern North Atlantic over the past 21,000 years.
 1456 Paleoceanography 25, PA1211.
 1457 Toucanne, S., Zaragosi, S., Bourillet, J.F., Cremer, M., Eynaud, F., Van Vliet-Lanoë, B.,
 1458 Penaud, A., Fontanier, C., Turon, J.L., Cortijo, E., Gibbard, P.L., 2009. Timing of
 1459 massive 'Fleuve Manche' discharges over the last 350 kyr: insights into the European ice-
 1460 sheet oscillations and the European drainage network from MIS 10 to 2. Quaternary
 1461 Science Reviews 28, 1238-1256.
 1462 van de Flierdt, T., Robinson, L.F., Adkins, J.F., Hemming, S.R., Goldstein, S.L., 2006.
 1463 Temporal stability of the neodymium isotope signature of the Holocene to glacial North
 1464 Atlantic. Paleoceanography 21, PA4102.
 1465 van Kreveld, S.A., 1996. Northeast Atlantic Late Quaternary planktic Foraminifera as
 1466 primary productivity and water mass indicators. Scripta Geologica 113, 23-91.

1467 Vance, D., Thirlwall, M., 2002. An assessment of mass discrimination in MC-ICPMS
 1468 using Nd isotopes. *Chemical Geology* 185, 227-240.

1469 Vaughan, D.G., Comiso, J.C., Allison, I., Carrasco, J., Kaser, G., Kwok, R., Mote, P.,
 1470 Murray, T., Paul, F., Ren, J., Rignot, E., Solomina, O., Steffen, K., Zhang, T., 2013.
 1471 Observations: Cryosphere, in: Stocker, T.F., Qin, D., Plattner, G.-K., Tignor, M., Allen,
 1472 S.K., Boschung, J., Nauels, A., Xia, Y., Bex, V., Midgley, P.M. (Eds.), *Climate Change*
 1473 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth
 1474 Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge
 1475 University Press, Cambridge, United Kingdom and New York, NY, USA.

1476 Vidal, L., Labeyrie, L., Cortijo, E., Arnold, M., Duplessy, J.C., Michel, E., Becqué, S.,
 1477 van Weering, T.C.E., 1997. Evidence for changes in the North Atlantic Deep Water
 1478 linked to meltwater surges during the Heinrich events. *Earth and Planetary Science*
 1479 *Letters* 146, 13-27.

1480 Vidal, L., Labeyrie, L., van Weering, T.C.E., 1998. Benthic $\delta^{18}\text{O}$ Records in the North
 1481 Atlantic Over the Last Glacial Period (60-10 kyr): Evidence for Brine Formation.
 1482 *Paleoceanography* 13, 245-251.

1483 Vinther, B.M., Clausen, H.B., Johnsen, S.J., Rasmussen, S.O., Andersen, K.K., Buchardt,
 1484 S.L., Dahl-Jensen, D., Seierstad, I.K., Siggaard-Andersen, M.L., Steffensen, J.P.,
 1485 Svensson, A., Olsen, J., Heinemeier, J., 2006. A synchronized dating of three Greenland
 1486 ice cores throughout the Holocene. *J. Geophys. Res.* 111, D13102.

1487 Voelker, A.H.L., Lebreiro, S.M., Schönfeld, J., Cacho, I., Erlenkeuser, H., Abrantes, F.,
 1488 2006. Mediterranean outflow strengthening during northern hemisphere coolings: A salt
 1489 source for the glacial Atlantic? *Earth and Planetary Science Letters* 245, 39-55.

1490 Waelbroeck, C., Duplessy, J.-C., Michel, E., Labeyrie, L., Paillard, D., Duprat, J., 2001.
 1491 The timing of the last deglaciation in North Atlantic climate records. *Nature* 412, 724-
 1492 727.
 1493 Weber, M.E., Mayer, L.A., Hillaire-Marcel, C., Bilodeau, G., Rack, F., Hiscott, R.N.,
 1494 Aksu, A.E., 2001. Derivation of $\delta^{18}\text{O}$ from Sediment Core Log Data: Implications for
 1495 Millennial-Scale Climate Change in the Labrador Sea. *Paleoceanography* 16, 503-514.
 1496 Wei, R., Abouchami, W., Zahn, R., Masque, P., 2015. Deep circulation changes in the
 1497 South Atlantic since the Last Glacial Maximum from Nd isotope and multi-proxy
 1498 records. *Earth and Planetary Science Letters* 434, 18-29.
 1499 Wilson, D.J., Crocket, K.C., van de Flierdt, T., Robinson, L.F., Adkins, J.F., 2014.
 1500 Dynamic intermediate ocean circulation in the North Atlantic during Heinrich Stadial 1:
 1501 A radiocarbon and neodymium isotope perspective. *Paleoceanography*, 2014PA002674.
 1502 Wilson, D.J., Piotrowski, A.M., Galy, A., Banakar, V.K.C.P.A., 2015. Interhemispheric
 1503 controls on deep ocean circulation and carbon chemistry during the last two glacial
 1504 cycles. *Paleoceanography* 30, 621-641.
 1505 Wilson, D.J., Piotrowski, A.M., Galy, A., Clegg, J.A., 2013. Reactivity of neodymium
 1506 carriers in deep sea sediments: implications for boundary exchange and
 1507 paleoceanography. *Geochimica et Cosmochimica Acta* 109, 197-221.
 1508 Wilson, D.J., Piotrowski, A.M., Galy, A., McCave, I.N., 2012. A boundary exchange
 1509 influence on deglacial neodymium isotope records from the deep western Indian Ocean.
 1510 *Earth and Planetary Science Letters* 341–344, 35-47.
 1511 Yu, E.-F., Francois, R., Bacon, M.P., 1996. Similar rates of modern and last-glacial ocean
 1512 thermohaline circulation inferred from radiochemical data. *Nature* 379, 689-694.

1513 Yu, J., Anderson, R.F., Jin, Z., Menviel, L., Zhang, F., Ryerson, F.J., Rohling, E.J., 2014.
 1514 Deep South Atlantic carbonate chemistry and increased interocean deep water exchange
 1515 during last deglaciation. *Quaternary Science Reviews* 90, 80-89.
 1516 Yu, J., Elderfield, H., 2007. Benthic foraminiferal B/Ca ratios reflect deep water
 1517 carbonate saturation state. *Earth and Planetary Science Letters* 258, 73-86.
 1518 Yu, J., Elderfield, H., Piotrowski, A.M., 2008. Seawater carbonate ion- $\delta^{13}\text{C}$ systematics
 1519 and application to glacial-interglacial North Atlantic ocean circulation. *Earth and*
 1520 *Planetary Science Letters* 271, 209-220.
 1521 Zahn, R., Schönfeld, J., Kudrass, H.-R., Park, M.-H., Erlenkeuser, H., Grootes, P., 1997.
 1522 Thermohaline Instability in the North Atlantic During Meltwater Events: Stable Isotope
 1523 and Ice-Rafted Detritus Records from Core SO75-26KL, Portuguese Margin.
 1524 *Paleoceanography* 12, 696-710.
 1525