**The effect of acute alcohol on motor-related EEG asymmetries during preparation of
approach or avoid alcohol responses**

Dr. Ozlem Korucuoglu^{1,2}, Dr. Thomas E. Gladwin^{1,3}, & Prof. Dr. Reinout W. Wiers¹

1 Addiction, Development and Psychopathology (ADAPT)-lab, Department of Psychology,

University of Amsterdam, Weesperplein 4, 1018 XA Amsterdam, The Netherlands

2 Department of Psychological Sciences, University of Missouri, Columbia, USA

3 Research Centre – Military Mental Health, Ministry of Defense, The Netherlands

Correspondence concerning this article should be addressed to Ozlem Korucuoglu*, Department
of Developmental Psychology, University of Amsterdam, Weesperplein 4, 1018 XA Amsterdam,
The Netherlands

*Present Address: Department of Psychological Sciences, University of Missouri, Columbia, MO
65211, USA, korucuogluo@missouri.edu

Running head: LATERALIZED EEG COMPONENTS IN ALCOHOL APPROACH
TENDENCIES

Abstract

Alcohol-approach tendencies have been associated with heavy drinking and play a role in the transition to alcohol abuse. Such cognitive biases might predict future alcohol use better under a low dose of alcohol. The aim of this prospective study was to investigate both the magnitude and the predictive power of alcohol-induced changes on approach-avoidance bias and bias-related cortical asymmetries during response preparation across heavy and light drinking adolescents. In heavy drinking adolescents greater approach-related asymmetry index in the beta-band was observed for soft-drink cues compared to alcohol ones and this increase was associated with increase in difficulty to regulate alcohol intake. Earlier findings demonstrated that young heavy drinkers hold both positive and negative implicit alcohol associations, reflecting an ambiguity towards alcohol. The increase in approach related beta-lateralization for soft-drink cues measured in this study may represent a compensatory effort for the weaker S-R mapping (approaching soft drink). The MRAA findings in this study may highlight a mechanism related to overcompensation due to ambivalent attitudes towards drinking in our heavy drinking sample who had greater problems to limit their alcohol intake compared to light drinkers. Moreover, a relatively strong approach soft-drink and weak approach alcohol reaction-time bias after alcohol predicted decreasing drinking; suggesting that the capacity to control the bias under alcohol could be a protective factor.

Keywords: Alcohol approach bias, alpha, beta, asymmetry, acute alcohol, prospective predictions

1. Introduction

In recent years, researchers have shown an increasing interest in drug-related cognitive biases due to their value in predicting drug-related behaviours and clinical outcomes. Cognitive biases have been found in adolescents and young adults in attentional processes (e.g. Field, Christiansen, Cole, & Goudie, 2007), action tendencies (approach biases, Field, Kiernan, Eastwood, & Child, 2008; Wiers, Rinck, Dictus, & van den Wildenberg, 2009) and implicit memory associations (e.g. Thush, Wiers, Ames, Grenard, Sussman, & Stacy, 2007). In adolescents these biases have been found to be predictive of drinking (memory bias: Thush & Wiers, 2007; Thush, Wiers, Ames, Grenard, Sussman, & Stacey, 2008; approach bias: Peeters et al., 2013). Note that some of these studies involved high-risk groups, either defined by education (special education for adolescents with externalizing problems, Peeters et al., 2013) or by genotype (e.g., Wiers et al., 2009). Training varieties of these tasks have been found to change the bias and reduce relapse rates (Eberl et al., 2013; Schoenmakers & Wiers, 2010; Wiers, Eberl, Rinck, Becker, & Lindenmeyer, 2011). Such results have clinical implications but are also of theoretical interest. Studies in young samples may provide important insights for our understanding of the role of automatic motivational processes in the continuation of drug use later in life (i.e. Curtin, Barnett, Colby, Rohsenow, & Monti, 2005).

The approach avoidance task (AAT) assesses automatically activated action tendencies to approach or avoid a category of stimuli (Rinck & Becker, 2007; Wiers et al., 2009). Facilitations in response times when approaching alcohol-related stimuli compared to avoidance response indicates that alcohol stimuli are compatible with approach versus avoidance responses. These stimulus-response compatibility effects are thought to emerge when implicit action tendencies are in line with the instructed responses during congruent blocks and/or it is difficult to maintain a stimulus-response association during incongruent blocks. If indeed the motivational value of the alcohol cues drives the bias in the alcohol AAT, facilitation in approach alcohol responses might be related to subjects' drinking profile. This was exemplified by the finding of a stronger

approach bias in heavier drinkers (Field et al., 2008; especially in those with a g-allele in the OPRM1 gene, Wiers et al., 2009).

Stimulus-response compatibility effects on motor programs can be studied through the hand-related response preparation. Regarding hand-related neural activity, both during movement preparation and execution, the beta (14–30 Hz) and mu (8-12 Hz) amplitude, decrease in amplitude (event-related desynchronization, ERD) over the motor cortex contralateral to the movement limb (Doyle, Yarrow, & Brown, 2005; Gladwin, Lindsen, & de Jong, 2006; Gladwin, 't Hart, & de Jong, 2008; Pfurtscheller, Neuper, Pichler-Zalaudek, Edlinger, & Lopes da Silva, 2000; Poljac & Yeung, 2014; Stancák & Pfurtscheller, 1995). These movement-related amplitude asymmetries (MRAA) can be quantified according to the double subtraction rationale of the Lateralized Readiness Potentials in the time domain (LRP; Colebatch, 2007); by taking the difference in amplitude between contralateral and ipsilateral activity during preparation of right-hand response minus difference between ipsi- and contra-lateral activity during preparation of left-hand response (Gladwin et al., 2006). Given that the calculation of the MRAA eliminates motor-unrelated hemispheric lateralization, the remaining activity reflects motor-related preparatory lateralized activity. With repetitive alcohol use, cues associated with alcohol gain incentive salience and induce motivational responses (Robinson & Berridge, 1993; 2008). However, the process of preparing an action (approach/avoid) for conditioned cues with a specific affective connotation (alcohol vs. control cues, or positive vs. negative cues) requires mapping a motor response to a stimulus and in the case of approach tendencies this mapping is strengthened for approach responses on alcohol cues. Therefore both incentive motivations and pavlovian responses are likely to play a role in the approach bias. Compared to many other cognitive biases, stimulus-response associations are more central to approach tendencies, given that in this bias certain stimulus categories are inextricably linked with specific motor responses. To this end, studying MRAA indices allow us to differentiate tasks that require much skill and effort from automatically activated tasks; by comparing stimulus-response associations with different

strengths. The first aim of the present study was to investigate the motor preparation in alcohol approach–avoidance bias by means of motor-related asymmetries as a function of drinking profile (light and heavy drinkers). Thus, we used a modified version of the AAT task that resembles the one used in our previous study (Korucuoglu, Gladwin, & Wiers, 2014), extending it by focusing on lateralized spectral analysis. In our previous study, preparatory activity was measured by presenting a warning (or a preparatory) stimulus before the presentation of an imperative stimulus (S2) to which the subject had to give a motor response. Contrary to our previous study where right-hand joystick movement was required for response, the task used in this study consisted of trials requiring both left and right hand approach or avoid responses for alcohol-related and control cues to allow the study of motor-related lateralization.

In an earlier study, we showed that a low dose of alcohol administration increased the parietal beta-ERD during preparation for the alcohol-compatible trials (‘approach-alcohol/avoid-control picture trials’) following alcohol administration (Korucuoglu et al., 2014), similar to facilitating effects of alcohol on appetitive processes (Duka & Townshend, 2004; Hodgson, Rankin, & Stockwell, 1979). A second aim of the current study was to assess whether acute alcohol would enhance asymmetries associated with drug-related approach/avoidance motivations. Finally, we tested whether alcohol-induced effects on lateralized power spectra would be related to alcohol consumption, problems, and motivations; and would predict alcohol escalation in a young sample.

2. Methods

2.1. Participants

Forty adolescents (age range = 16-20 years) were recruited from local high schools in Amsterdam. Prior to the testing sessions, participants were informed about the study restrictions by email. Participants were required to be minimally 16 years-old (minimum drinking age in

Netherlands at the time of the study), with a minimum weight of 50 kg and to have had at least one full drink in their lifetime. Participants were requested not to drink any alcohol 24 hours before testing and eat a meal or drink caffeine 4 hours prior to testing. Participants' compliance with these restrictions was confirmed with self-report. Moreover, participants were instructed to abstain from any legal and illegal drugs for at least 1 week; their compliance with this restriction was confirmed with a urine test. Exclusion criteria were psychiatric disorders, diagnosed cases of drug use disorder, head trauma, seizures, severe physical illness, cardiovascular disease, chronic obstructive pulmonary disease, the presence of major medical conditions, and use of medication. Further exclusion criteria for female participants were pregnancy and breast-feeding; which were assessed with self-report.

Seven participants were excluded from data analysis (two due to a positive drug test for THC, two due to missing data in one session, one due to broken electrode, one due to being left handed, and one subject's AUDIT score was missing), analysis was conducted with the remaining 33 participants. In this study we examined participants with light and heavy drinking patterns, drinking groups were formed by using an inventory on alcohol use and problems (Alcohol Use Disorder Identification Test, AUDIT) using a median-split (heavy drinking: AUDIT > 8). All participants had normal or corrected-to-normal visual acuity. Prior to the experiment, written informed consent was obtained from all participants and from parents of participants under 18. The study was approved by the Ethical Committee of University of Amsterdam Psychology Department. Participants received financial compensation.

[Insert Table1 about here]

2.2. Alcohol Administration

All subjects participated in two sessions administered on two different days, between 2 to 7 days apart. Sessions started between 12:00 and 18:00 PM. Alcohol was administered in one

session and placebo in the other. Dose order was counterbalanced across subjects. Participants were told that they would receive a different dose of alcohol during both sessions, to keep expectancy effects similar across sessions.

To keep the participants as well as the experimenter oblivious to the condition, a double blind procedure was used. Over-age subjects (18 year-olds and above) received a mix of vodka and orange juice. Under-age subjects (16 and 17 year-olds) received a vodka-orange premixed drink (Eristoff & Orange Can, commercial ready-to-drink alcoholic beverages with a 7 % Vol). The alcohol content and the total volume of the liquid delivered to the participants under and over the age of 18 were the same (0.45g/kg with a maximum cut-off of 100 ml vodka). The mix was divided into three equal portions. Two of the drinks were served with 5 minutes apart, prior to commencing the task, and after electrode placement. Up to 3 minutes was allowed for drinking followed by 2 minutes of mouthwash to remove the residual alcohol in the mouth. In between the tasks 1/3 of the mix was administered as a booster drink in order to eliminate measurement during the descending limb of the BrAC. To enhance the alcohol taste, all the drinks had a lemon soaked in vodka and the glass in which drinks were served was sprayed with vodka beforehand. To mask the alcohol taste all drinks had three drops of tabasco sauce (McIlhenny Co., USA). The procedure was identical in each session, except alcohol was replaced with orange juice in the placebo condition.

Breath alcohol concentration (BrAC) was collected 5 minutes after the first two drinks, before and after the booster drink, and at the end of the experiment by using the Lion alcolmeter® SD-400 (Lion Laboratories Limited, South Glamorgan, Wales). Participants filled out the *Brief Biphaseic Alcohol Effects Scale* (B-BAES, Rueger, McNamara, & King, 2009) each time a breath sample was taken, except before the booster drink. Throughout the experiment the BrAC was measured three times during which subjects also filled the B-BAES questionnaire: after alcohol administration, before the booster drink and at the end of the experiment. Moreover, an additional

BrAC measurement was collected after the booster drink in order to monitor alcohol level following the top-up dose.

After completion of both sessions, a short manipulation check interview was conducted to determine whether the participants were aware of the alcohol contents of the drinks. Deception was not successful for one of the participants. Participants were debriefed about the true nature of the study and remained at the research site until their breath sample was 25mg/100ml or less.

2.3. Procedure

Upon arrival in the lab, participants filled out demographics, questionnaires related to personality and drinking habits. At the start of each session, participants completed the *Desire for Alcohol Questionnaire* (DAQ; Love, Es, & Willner, 1998) and the *Positive and Negative Affect Scale* (PANAS; Watson, Clark, & Tellegen, 1988) to measure differences in current mood and craving across sessions. Current alcohol use and problems were assessed with the AUDIT (Saunders, Aasland, Babor, de la Fuente, & Grant, 1993); we used both the standard past year version, and a version about the past 3 months. Motives to drink alcohol and drinking restraint were assessed with the *Drinking Motives Questionnaire-Revised* (DMQR-R; Cooper, 1994) and *Temptation and Restraint Inventory* (TRI; Collins & Lapp, 1992), respectively. Before and after alcohol administration, participants also performed other unrelated tasks (See Supplementary materials Figure S2, for timeline of events). Order of the tasks was counterbalanced across participants, but was kept same across sessions for each subject. The data of the other tasks are not reported in this paper.

Each session took approximately two and a half hours, including breaks and the application of electrodes, during one afternoon. Six months after these two assessments, participants were contacted with e-mail for an online assessment on recent alcohol and drug use. If no response was received within a week, participants were contacted by phone. During follow-up assessment, participants filled out the same alcohol-related scales as during pre-test.

2.3. Alcohol Approach Avoidance Task (A-AAT)

In the original A-AAT (Wiers et al., 2009) participants were instructed to pull (approach) or push (avoid) alcohol-related and control pictures by using a joystick. The EEG version of the AAT used in the current study was developed to compare the neural activity during preparation of alcohol approach and avoidance responses. Compared to the relevant-feature version used in our previous study (Korucuoglu et al., 2014), in this experiment the irrelevant-feature version of the task was used, where participants were presented with alcohol-related or soft-drink pictures in a portrait or landscape orientation with participants being instructed to approach or avoid pictures depending on the orientation of the picture (cf. Cousijn, Goudriaan, & Wiers, 2011).

The sequence of events in the trial was as follows (Figure 1): The trial started with a fixation period (500 ms or 700 ms), followed by a preparation period. During the preparation period, the word “Voorbereiden” (“prepare”) was presented on top of the stimuli. Participants were instructed to prepare their response depending on the orientation of the picture (portrait or landscape) and instructions (pull or push by pressing the left or right button, assigned per block; see below for more details), and to withhold their response until the word disappeared. The word “Voorbereiden” was displayed centrally for a randomly selected amount of time between 1000 ms and 1500 ms with 100-ms increments. The stimuli remained on the screen until the response was given. After the response there was a zoom effect with a fixed duration of 500 ms. During this zoom effect, pulled pictures became bigger and pushed pictures became smaller. Participants received feedback only if the response was incorrect. Each picture was presented equally often in the portrait and landscape orientations. Since each picture (alcohol-related or soft-drink pictures) was presented in both orientations (left and right-tilted), the task consisted of *four experimental conditions*: 1) approach alcohol-related pictures, 2) avoid soft-drink pictures, 3) approach soft-drink pictures and 4) avoid alcohol-related pictures.

The task contained four blocks in total. In order to disentangle left/right hand and push/pull responses and allow motor-related asymmetry analyses, the assignment of the buttons (left or right hand side) to each action type (approach or avoid) alternated across blocks. For this reason, the sequence of block type during 4 experimental blocks followed either ABBA for half of the participants or BAAB design for the other half. During the block type A, the left button was assigned to the approach action and the right button was assigned to the avoid action. During the block type B, the mapping of left-right response buttons on action type was reversed. The contingencies of orientation (portrait or landscape) and the target action (pull or push) were randomized across participants in such a way that half of the participants were instructed to pull the pictures in portrait orientation and push pictures in landscape orientation and the other half received opposite instructions.

Each block started with 16 practice trials and was followed by 48 experimental trials. Non-beverage images (grey rectangles) were used during practice trials. During the first 6 practice trials in each block, the correct response was presented on top of the rectangles. Participants repeated a trial during practice block if the response was incorrect. To control for the effect of picture familiarity across sessions, two sets of pictures with and without alcohol contents were matched (12 alcohol-related and 12 control pictures, each presented equally often), and each stimulus set was randomly assigned to a session.

[Insert Figure1 about here]

2.4. Behavioural Data Preparation

Practice trials and trials with incorrect response (i.e a pull response in a push trial) were excluded from the behavioral data for RT analysis. RT was calculated from the end of the preparation period until the motor response. Due to the preparation period, responses were fast

and no trials were excluded based on RT. Median RTs were analyzed as in previous AAT studies (e.g., Cousijn et al., 2011; Wiers et al., 2009).

2.5. Electroencephalogram (EEG) recording and data preparation

Electrophysiological data were recorded from the scalp using an Active-Two amplifier (Biosemi, Amsterdam, the Netherlands) from 32-scalp sites. Electrodes were placed at the standard positions of the 10-20 international system. Two electrodes were placed at the outer canthi of the eyes to measure horizontal eye movements. Two electrodes were placed at below and above the left eye to measure vertical eye movements. EEG was recorded at 2048 Hz sampling rate. The distance between the screen and the subject was kept at 75 cm.

EEG preprocessing was conducted using Brain Vision Analyzer (version 2.0, Brain Products GmbH, Munich, Germany). Data were down-sampled to 250Hz, re-referenced offline to the average of left and right mastoids, low pass filtered at 50Hz, and high pass filtered at 0.1 Hz. Ocular correction was applied using the algorithm of (Gratton, Coles, & Donchin, 1983). EEG data were segmented into 3 sec epochs starting 1 sec. before the cue presentation to 2 sec. afterwards. Trials were considered artefacts when the difference between consecutive data points was larger than 75 mV and the difference between the lowest and the highest voltage within a segment was higher than 200 mV. Epochs with an amplitude exceeding ± 100 mV were excluded. After the exclusion of artefacts and noisy data, in the placebo condition an average of 38.39, 38.03, 38.18, 38.63 trials remained for subsequent analysis for the avoid soft, avoid alcohol, approach soft, and approach alcohol conditions, respectively. In the alcohol condition the numbers of remaining trials were 39.85, 39.21, 40.39, and 39.42, respectively for the avoid soft, avoid alcohol, approach soft, and approach alcohol conditions.

The Fieldtrip toolbox for EEG/MEG analysis was used for the time-frequency analysis (Oostenveld, Fries, Maris, & Schoffelen, 2011) running under Matlab 2010b. Because of their sensitivity to muscle activity, the (most) peripheral electrodes from left to right earlobes (Fp1,

FP2, F7, F8, T7, and T8) were excluded from further data analysis. Time-frequency was performed by convolving the time series with a family of Morlet Wavelets with a family ratio of ($f_0/\sigma_f=7$), where f_0 represent the frequency of interest. Frequencies of interest were alpha (8-12 Hz with 1 Hz frequency steps) and beta (13-30 Hz with 2 Hz frequency steps) frequency ranges. An absolute baseline correction was applied to the power spectrum by using the time period of -600 to -200 ms preceding the presentation of the cue.

Condition specific grand averages (e.g. approach alcohol cue trials) were calculated separately for each response hand. Based on previous reports MRAA was calculated for the central beta and mu (de Jong, Gladwin, & 't Hart, 2006; Gladwin, Lindsen, & de Jong, 2006; Poljac and Yeung, 2014) and for the parietal alpha (Deiber, Sallard, Ludwig, Ghezzi, Barral, & Ibañez, 2012; note that for the parietal alpha we used P3-P4 channels in the same line with C3-C4, instead of CP3-CP4 used by the authors). To estimate lateralization, first the difference in power between two equal measuring points in the left and right hemispheres (C3-C4, P3-P4) was calculated for the left and right hand responses separately. Subsequently, a difference score was calculated between the left and the right hand responses (example for the central electrodes: $[(C3-C4)_{\text{Right hand response}} - (C3-C4)_{\text{Left hand response}}])$). These MRAA values are presented in supplementary materials FigureS1. Given that a decrease in power is expected for the hemisphere contralateral to the movement, more negative MRAA values would indicate greater motor-related lateralization due to increased ERD contralateral to the movement. The power estimates were averaged for three successive time points (T1: 0-350ms, T2: 350-700ms, T3: 700-1000ms) following the presentation of the cue to separate early, middle and late stages of motor preparation (Poljac and Yeung, 2014).

2.6. Procedure to calculate bias scores and contrasts

Calculation of bias scores and contrasts are summarized in Table2. First, cue-specific bias-scores were calculated separately for alcohol-related and soft-drink pictures by subtraction the median

RT in pull trials from the median RT in push trials. Positive scores represent an approach bias and negative scores represent an avoidance bias. Similar to the calculation of bias scores, cue-specific MRAA bias-scores were calculated separately for alcohol and soft-drink cues by subtracting the MRAA in push trials from the MRAA in pull trials (i.e. $\text{MRAA}_{\text{alcohol bias-scores in placebo}} = \text{MRAA}_{\text{pull}} - \text{MRAA}_{\text{push alcohol-stimuli}}$ at T1 after placebo). Given that MRAA is calculated based on ERDs (decrease in activity), negative MRAA bias-scores represent relatively higher ERDs for the approach compared to avoid responses, and positive MRAA bias-scores represent relatively higher ERDs for the avoid compared to approach responses.

Brain-behavior relationship between MRAA scores and questionnaires was investigated with correlational analysis. For the purpose of this analysis, overall bias score was calculated for the MRAA data. For the calculation of overall bias, a double subtraction procedure was followed where the difference between the bias-score for the soft-drink and for the alcohol cues were calculated both for the MRAA data separately in the placebo and alcohol conditions (i.e. $(\text{MRAA}_{\text{pull}} - \text{MRAA}_{\text{push}})_{\text{alcohol-stimuli}} - (\text{MRAA}_{\text{pull}} - \text{MRAA}_{\text{push}})_{\text{control-stimuli}}$ in the placebo dose). Overall bias scores are MRAA differences between push and pull trials across alcohol-related and soft drink stimuli, note that this controls for general response bias due to a specific action (approach/avoid) or due to a specific stimulus category (alcohol/control cues). Therefore, negative overall MRAA bias-scores represent relatively greater alcohol-cue MRAA bias than soft-drink-cue MRAA bias and vice versa for the positive overall MRAA bias-scores.

Lastly, to study the predictive power of alcohol-induced effects on future drinking, dose-contrast scores, representing the difference between the alcohol and placebo conditions, were calculated both for the overall RT and the MRAA bias and used as predictors in the regression analysis.

[Insert Table2 about here]

2.7.1 Statistical Analysis

All analyses were conducted using a repeated measures ANOVA (RM-ANOVA) in SPSS. The Stimulation and Sedation subscales of B-BAES scores were separately analysed with *Dose* (Placebo, Alcohol), and *Time* (pre-task and post-task) as within-subjects factors. Similarly, the estimated blood alcohol levels (BAL) were subjected to RM-ANOVA, with time (BAL pre-AAT and BAL post-AAT) as within subject variable.

For the analysis of accuracy, RM-ANOVA was conducted with *Dose* (placebo, alcohol), *Action* (approach, avoid) and *Stimulus Category* (alcohol-related, soft-drink cues) as within-subject factors and *Group* (Light, Heavy drinkers) as between-subjects factor. The cue-specific bias scores were analyzed with *Dose* (placebo, alcohol) and *Bias Type* (alcohol-cue RT bias, soft-drink-cue RT bias) as within-subject factors and *Group* (Light, Heavy drinkers) as between-subjects factor. Note that for all the analysis conducted with RM-ANOVA, the cue-specific bias scores (difference between push-pull trials) were used, except for the accuracy and RT data we also presented results with *Dose* (placebo, alcohol), *Action* (approach, avoid) and *Stimulus Category* (alcohol-related, soft-drink pictures) as within-subject factors and *Group* (Light, Heavy drinkers) as between-subjects factor.

For the lateralization, statistical analysis was conducted with *Dose* (Placebo, Alcohol), and *MRAA Bias-scores* (Alcohol-cue MRAA bias, Soft-drink-cue MRAA bias) as within-subject variables and *Group* (Light, Heavy drinkers) as between-subject variable, and for each time interval (T1:0-350ms, T2:350-700ms, T3:700-1000ms) separately. Note that following earlier studies, the MRAA bias scores were analysed separately for early (T1), middle (T2) and late (T3) preparation intervals, because it was shown that preparatory activity in the frequency domain unfold over time (De Jong et al., 2006; Polzac and Yeung, 2014). In the supplementary materials, we presented analysis showing an increase in motor preparation with the approaching response deadline (see section ‘desynchronization across timepoints’). The MRAA in different time intervals may represent quite different processes, and so even in the absence of significant

interactions with the *Time* factor, we provide tests for each time interval separately. However, we acknowledge and emphasize that such effects must be considered exploratory.

Post-hoc comparisons were conducted by using paired sample t-tests while comparing task conditions (reported under ‘task effects per group’) and independent sample t-tests when comparing heavy vs. light drinkers.

2.7.2 Relationships between MRAA and Individual Differences

Correlations between overall MRAA bias score and alcohol-related problems/drinking motives (TRI/DMQR) were assessed with Pearson correlations.

2.7.3. Prediction of future drinking

In order to assess whether differences in RT bias-scores and MRAA bias-scores (for all three frequency bands) across sessions predicted unique variance in the change in alcohol use during the six months after the experiment, a hierarchical multiple regression analysis was conducted. First, behavioural measures (AUDIT score for recent use at baseline from the version about the past 90 days -sum of scores of items on frequency of drinking, typical quantity and frequency of heavy drinking-) were entered to the regression model, followed by the RT and MRAA dose-contrast scores (alcohol-induced changes in the RT and the MRAA). We focused on 1) whether alcohol effects on behaviour predicted change in alcohol use, and 2) whether alcohol effects on brain responses predicted alcohol use beyond the predictive value of behavioural measures.

3. Results

3.1. Manipulation Checks

For the Stimulation and Sedation subscales of B-BAES scores, results revealed a significant main effect of *Dose* for the sedation subscale ($F(1, 32) = 5.016, p = .032, \eta^2_p = .15$).

Sedation scores were higher for the alcohol dose compared to the placebo dose, however post-hoc analysis with paired t-tests did not reveal any differences across conditions. All other main and interaction effects were not significant ($ps > .15$).

Three subjects' post-task BrAC data were lost, the analysis was completed with the remaining participants. Results revealed that subjects performed the task during the steady state of alcohol level ($p > .198$) (See Table 3).

[Insert Table3 about here]

3.2. Accuracy

Accuracy data revealed a two-way interaction effect of *Dose* by *Action Type* ($F(1, 31) = 5.874, p = .021$). Post-hoc analysis with paired sample t-test revealed that compared to placebo, after alcohol administration participants made more errors in trials in which they had to avoid alcohol-related stimuli ($t(32) = -2.292, p = .029$).

3.3. Bias Scores

The analysis of bias scores revealed a two-way interaction of *Dose* and *Bias Type* ($F(1, 31) = 6.602, p = .015, \eta^2_p = .176$). No interaction with the *Group* variable was observed. Participants demonstrated a non-significant positive alcohol-cue RT bias (approach bias for alcohol pictures) after a placebo dose and a positive soft-drink-cue RT bias (an approach bias for soft-drink pictures) after an alcohol dose. Post-hoc analysis revealed that after alcohol administration, soft-drink-cue RT bias scores tended to be higher compared to alcohol-cue RT bias scores, which did not reach significance ($t(32) = 1.816, p = .079$) (See Figure 2, upper panel).

3.4. Reaction times

Reaction time data comparing all task conditions revealed a three-way interaction of *Dose* by *Action Type* and *Stimulus Category* ($F(1, 31) = 6.579, p = .015, \eta_p^2 = .18$) and *Stimulus Category* by *Group* ($F(1, 31) = 4.48, p = .042, \eta_p^2 = .13$) interaction effect. To follow-up three-way interaction of *Dose* by *Action* by *Stimulus Category*, we performed analysis separately for the placebo and alcohol dose. Results revealed a faster approach alcohol response compared to avoid alcohol response in the placebo condition ($t(32) = -2.1, p = .044$). To follow-up *Stimulus Category* by *Group* interaction, we collapsed the data across levels of *Dose* and *Action* variables. T-tests showed that overall heavy drinkers were faster for alcohol cues compared to soft drink cues, $t(14) = -2.879, p = .012$. No effect was observed for light drinkers. (See Figure 2, lower panel).

[Insert Figure2 about here]

3.5. Beta-MRAA Bias

Beta MRAA bias scores are presented in Figure 2 (top panel) for light and heavy drinkers. As can be seen, in the placebo condition light drinkers had positive alcohol-cue and soft-drink-cue MRAA bias scores representing avoid-related lateralization and heavy drinkers had negative alcohol-cue and soft-drink-cue MRAA bias scores representing approach-related lateralization for both cue types. In the alcohol condition, this pattern was reversed. The differences across groups were most prominent for the soft-drink-cue MRAA bias, especially in the placebo condition. We tested whether the positive avoid-related MRAA scores in heavy drinkers and negative approach-related MRAA scores in light drinkers were statistically different from each other and also across conditions. Analysis of the beta-MRAA revealed a *Dose* by *Group* interaction effect at early ($F(1, 31) = 7.927, p = .008, \eta_p^2 = .204$), at middle ($F(1, 31) = 9.158, p = .005, \eta_p^2 = .113$) and late preparation periods ($F(1, 31) = 8.958, p = .005, \eta_p^2 = .224$). At late preparation period, a significant *Dose* by *Bias* by *Group* ($F(1, 31) = 6.988, p = .013, \eta_p^2 = .224$).

=.184) and marginally significant *Dose* by *Bias* ($F(1, 31) = 3.987, p = .055, \eta^2_p = .114$) interaction effects were observed. To understand the nature of this three-way interaction, we conducted paired sample t-tests per group separately and independent samples t-tests to compare light and heavy drinkers.

Task effect per Group: At 700-1000ms, after a placebo dose heavy drinkers' negative MRAA soft-drink-cue bias-scores were different from the positive alcohol-cue MRAA bias-scores ($t(14)=3.143, p=.007$). Also, for heavy drinkers at 700-1000ms, negative soft-drink-cue MRAA bias-scores after a placebo dose were different from positive soft-drink-cue MRAA bias-scores after an alcohol dose ($t(14)=-2.641, p=.019$). Light drinkers had negative soft-drink-cue MRAA bias-scores after an alcohol dose which was different than the positive soft-drink-cue MRAA bias-scores after a placebo dose at 0-350ms ($t(17)=2.742, p=.014$), at 350-700ms ($t(17)=2.447, p=.026$), and at 700-1000ms ($t(17)=2.608, p=.022$). Moreover, at 700-1000ms, light drinkers had negative alcohol-cue MRAA bias-scores after an alcohol dose which was different from the positive alcohol-cue MRAA bias-scores after a placebo dose ($t(17)=2.527, p=.018$).

Heavy vs. Light Drinkers: At 0-350ms, no differences were observed. At 350-700ms and 700-1000ms, after a placebo dose, heavy drinkers' negative soft-drink-cue MRAA bias was different from light drinkers' positive soft-drink-cue MRAA bias, (350-700ms: $t(31)=2.644, p=.013$; 700-1000ms: $t(31)=4.055, p<.001$). In the alcohol condition, light drinkers' negative alcohol-cue MRAA bias was different from heavy drinkers' positive alcohol-cue MRAA bias ($t(31)=-1.987, p=.056$) at 700-1000ms.

3.6. Mu-MRAA Bias

Mu MRAA bias scores are presented in Figure 2 (middle panel) for light and heavy drinkers. As can be seen, both for alcohol and soft-drink cues, light drinkers had negative MRAA bias scores representing approach-related lateralization and heavy drinkers had positive MRAA

bias scores representing avoid-related lateralization, except for the late preparatory period where this pattern was reversed. Next we tested if the differences across heavy and light drinkers were statistically significant differences.

During the early preparation period (0-350ms), analysis of the mu-MRAA bias revealed an interaction effect of *Dose* by *Bias* by *Group* ($F(1, 31) = 4.12, p = .051, \eta^2_p = .117$). During the middle preparation period (350-700 ms), *Dose* by *Bias* by *Group* was marginally significant ($F(1, 31) = 3.846, p = .059, \eta^2_p = .11$). During the late preparation period (700-1000 ms) a marginally significant *Dose* by *Group* interaction effect was observed ($F(1, 31) = 3.958, p = .056, \eta^2_p = .113$).

Task effect per Group: No significant differences were observed across task conditions with pairwise t-tests conducted separately in heavy and light drinkers.

Heavy vs. Light Drinkers: Independent sample t-tests comparing heavy and light drinkers revealed that after an alcohol dose, light drinkers' negative alcohol-cue MRAA bias-scores were different than heavy drinkers' positive alcohol-cue MRAA bias-scores at 0-350ms ($t(31) = -2.332, p = .026$) and 350-700ms ($t(31) = -2.08, p = .046$). At 700-1000ms, after an alcohol dose, light drinkers' negative soft-drink-cue MRAA bias-scores were different from the heavy drinkers' positive soft-drink-cue MRAA bias-scores ($t(31) = -2.178, p = .037$) (See Figure 3).

3.7. Parietal alpha-MRAA Bias

The parietal alpha revealed an interaction effect of *Bias* by *Group* at 0-350ms ($F(31) = 6.284, p = .018, \eta^2_p = .169$) and an interaction effect of *Dose* by *Group* at 700-1000ms ($F(31) = 4.374, p = .045, \eta^2_p = .124$).

Task effect per Group: At 0-350ms, after a placebo dose, light drinkers' negative alcohol-cue MRAA bias-scores was different than the positive soft-drink-cue MRAA bias-scores ($t(17) = -3.008, p = .008$). At 700-1000ms, positive soft-drink-cue MRAA bias-scores (representing greater

avoid-related lateralization) after a placebo dose and the negative soft-drink-cue MRAA bias-scores (representing greater approach-related lateralization) after an alcohol dose were significantly different ($t(17)=2.31, p=.034$).

Heavy vs. Light Drinkers: No differences were observed across groups.

[Insert Figure3 about here]

3.8. Correlations

The Govern subscale of the TRI questionnaire ('difficulty controlling alcohol intake') positively correlated with the overall central beta-MRAA bias scores in the alcohol condition at 350-700ms ($r=.34, p=.05$) and 700-1000ms ($r=.43, p=.012$) and with the overall MRAA bias scores in the placebo condition at 700-1000ms ($r=.37, p=.032$) (See Figure 4). Individuals with higher TRI scores had more positive overall MRAA bias scores, and individuals with lower TRI scores had more negative overall MRAA bias scores.

[Insert Figure4 about here]

3.9. Neural Predictors of Alcohol Use After Six Months

Six months after the baseline assessment (alcohol challenge session), 82.5% follow-up response rate was achieved in the full sample of 40 participants. RT dose-contrast scores (depicting alcohol-induced changes on the overall RT bias scores) and the parietal alpha-MRAA dose-contrast scores at 350-700ms (depicted alcohol-induced changes on the overall alpha-MRAA bias scores) predicted future alcohol use beyond the variance explained by baseline AUDIT scores. The total variance explained by the full model was 81.5% ($F\text{-change}_{1,24}=5.903, p=.023$). The baseline AUDIT scores explained 70.5% of the variance ($F\text{-change}_{1,26}=62.103, p<.001$). RT dose-contrast scores and the parietal alpha-MRAA dose-contrast scores explained an

additional 6.4 and 4.6% of the variance ($F\text{-change}_{1,25}=6.931, p=.014$; $F\text{-change}_{1,24}=5.903, p=.023$) (See Figure 5). To follow up, a correlation analysis between change in AUDIT scores ($\text{AUDIT}_{\text{follow-up}} - \text{AUDIT}_{\text{baseline}}$) and the predictors of the change in alcohol use was conducted (the RT and the parietal alpha-MRAA dose-contrast scores). Individuals who had relatively more negative overall bias scores for the RT after alcohol administration at baseline (due to a stronger approach soft-drink and weaker approach alcohol bias, as depicted in Figure5b), had lower Audit scores, 6 months later ($r=.384, p=.044$). Follow-up correlations for the parietal-MRAA dose-contrast scores did not reveal significant effects.

[Insert Figure5 about here]

4. Discussion

In the current EEG study, we focused on motor-related lateralization during preparation for approach and avoidance behaviors in the context of alcohol cues and investigated the effects of a prime dose of alcohol on these neurophysiological measures in heavy and light drinking adolescents. As in previous studies of motor preparation, preparation of a left/right hand response during the alcohol approach-avoidance task led to an MRAA following the presentation of the imperative stimulus (Supplementary materials, Figure S1). This preparatory MRAA was found to be related to experimental conditions and drinking behavior. Behavioural results revealed faster responses for approaching alcohol-related cues compared to avoiding alcohol ones in the placebo condition. After alcohol, a non-significant decrease was observed for approach alcohol bias, especially in heavy drinkers. A relatively strong approach soft-drink and weak approach alcohol reaction-time bias after alcohol predicted decreasing drinking; suggesting that the capacity to modulate an alcohol-approach bias while under the influence of an alcohol dose could be a protective factor. In heavy drinking adolescents increased approach-related asymmetry in the beta-band was observed for soft-drink cues compared to alcohol ones and this increase was

544 associated with an increase in difficulty in regulating alcohol intake. Possible implications of this
545 effect on the interpretation of the MRAA are discussed below.

546 In earlier studies, the mu- and beta-MRAA indices have been studied with switch task,
547 pre-cueing RT paradigm, and motor imagery task (de Jong, Gladwin, & 't Hart, 2006; Deiber et
548 al., 2012; Doyle et al., 2005; Gladwin et al., 2008; Gladwin et al., 2006; Nam, Jeon, Kim, Lee, &
549 Park, 2011; Poljac & Yeung, 2014). During task switching paradigms (subjects need to switch
550 their response hand when the current task switches), a reversal of lateralization of the mu and
551 beta-MRAA from previous to current task set has been observed (de Jong et al., 2006; Gladwin et
552 al., 2006; Poljac & Yeung, 2014), suggesting that MRAA reflects selection of motor goal and
553 advance task preparation. This interpretation is strengthened by the finding of higher beta-band
554 MRAA in 100% informative cues compared to 50% informative one (Doyle et al., 2005). In this
555 study visuospatial attention to the imperative cues was also measured and it was found to be
556 unrelated to the magnitude of the MRAA index. However, in another pre-cueing RT task, a cento-
557 parietal alpha-MRAA was found to be reflecting visuospatial attention (Deiber et al., 2012). This
558 study revealed a spectral pattern for weak lateralizers suggesting the recruitment of more
559 visuospatial attentional resources (alpha ERD) and for high lateralizers suppression of irrelevant
560 visual activity (alpha event-related synchronization, ERS).

561 Based on earlier findings, we expected that heavier drinkers would show an increased
562 (more negative) mu- and beta-MRAA index for the approach versus avoidance alcohol-related
563 cues compared to soft-drink cues, representing advance response preparation for these trial types.
564 In heavy drinkers, greater approach-related lateralization was observed for approach soft-drink
565 cues especially during the late preparation period, suggesting an increased asymmetry index for
566 the bias in the direction opposite to the one hypothesized. The effects for the mu- and alpha-
567 MRAA bias scores were found to be in the same direction, higher lateralization of the ERD for
568 soft-drink bias in heavy drinkers. For the alpha and the mu, differences across conditions were
569 moderate and did not lead to significant results. An exploratory analysis also performed on

behavior indicated that increased approach alcohol reaction (compared to avoidance reaction) was more pronounced in heavy drinkers ($p < .05$). Given that heavy drinkers had a greater approach bias for alcohol cues and also greater lateralization for approaching soft drink; the findings of the current study suggest that the asymmetry index measured with the AAT is likely to reflect an effortful response preparation process rather than an automatic advance response preparation. This could be due to two reasons: first, a lack of lateralization for approach alcohol response might represent the presence of an automatic response bias. However, another likely scenario is a possible relationship between behavior and lateralization, which resembles the “speed-accuracy” tradeoff for perceptual tasks. In the present case, our heavy drinking participants showed faster responses for approaching alcohol stimuli (compared to avoiding alcohol pictures) after placebo administration, however, they lacked approach-related lateralization for alcohol-related stimuli in the brain. In line with this, for the soft-drink stimuli an increased lateralization was observed for more effortful approach behavior. In sum, rather than a lack of lateralization possibly meaning an automatized process, the presence of lateralization could reflect effortful processing to overcome preexisting response preferences. Also our correlational analysis revealed that individuals with greater difficulty in regulating their drinking (note that heavy drinkers had greater difficulty), had greater approach-related lateralization for soft-drink cues and individuals with less problem with control over drinking had greater approach-related lateralization for alcohol cues. Using an alcohol implicit association test (IAT), it has been shown that young heavy drinkers hold both positive and negative alcohol associations (Houben & Wiers, 2006), most likely reflecting ambiguity towards alcohol. Therefore, a likely explanation for the MRAA pattern in heavy drinkers is that problems in controlling alcohol intake may have caused ambivalence in these individuals, and subjects may have compensated for this ambiguity by putting more effort in preparing their response for trials incongruent with their state of drinking profile (approaching soft-drink cues).

Based on earlier findings of alcohol's priming effects on cognitive biases in adult samples (Field, Schoenmakers, & Wiers, 2011), one may expect acute alcohol to increase this bias. However, earlier studies failed to show such an effect on RT with a relevant-feature version of the task with explicit instructions to approach/avoid alcohol-related cues (Korucuoglu et al., 2014; Schoenmakers, Wiers, & Field, 2008). Results of the current study employing an irrelevant-feature version of the task demonstrated that after alcohol, the bias for the alcohol cues decreased especially in heavy drinkers. With alcohol administration, while heavy drinkers slowed down their responding for approach alcohol and avoid soft-drink cue trials (this could be due to a decrease in inhibition or an increase in distraction), light drinkers showed a non-significant decrease in response time during approach soft-drink and avoid alcohol cue trial types. However, this difference between groups was not significant, and thus the direction of this effect can only be tentatively interpreted. Moreover, regression analysis revealed that individuals who had a relatively strong avoid alcohol bias after alcohol administration during the alcohol challenge (due to a stronger approach soft-drink and a weaker approach alcohol bias), had lower Audit scores, six months later. The evidence in this study suggests that the ability to respond adaptively under the influence of alcohol can be a protective factor for the development of addictive behaviours. Earlier studies showed that if alcohol is consumed in the presence of conditioned cues (drug-related environmental cues), individuals are able to counter the effects of alcohol on cognitive function (Birak, Terry, & Higgs, 2010; Birak, Higgs, & Terry, 2011), suggesting a cognitive tolerance to drugs in the presence of drug cues. It is important to note that these results might be specific to irrelevant version of the task used here, given that the implicit nature of the instructions probably gave more room for the top-down influence of task instructions on performance.

To conclude, results revealed greater preparatory approach-related lateralized activity for approach soft-drink cues in heavier drinkers in comparison to light drinkers and also in comparison to lateralization for the alcohol cues. The beta-lateralization measured in this study

may represent a compensatory effort for the weaker S-R mapping in heavy drinkers. The extent of alcohol-induced changes on the bias were related to changes in alcohol use, suggesting that the capacity to control the bias under alcohol could be a protective factor. It is important to note here, heavier drinkers in the present study also reported greater problems with controlling their drinking behavior. Studies with preselected samples can be considered to compare lateralization index in heavy drinkers with and without problems to control their drinking levels. Also future studies with a larger sample can focus on asymmetry differences between heavy drinking individuals who can and cannot overcome their approach alcohol bias.

Acknowledgements

The authors are supported by VICI award 453.08.01 from the Netherlands National Science Foundation (N.W.O.). Thomas E. Gladwin is supported by ERAB grant EA 1239.

635

References

- 636 Birak, K. S., Higgs, S., & Terry, P. (2011). Conditioned tolerance to the effects of alcohol on
 637 inhibitory control in humans. *Alcohol and Alcoholism (Oxford, Oxfordshire)*, 46(6), 686–93.
 638 doi:10.1093/alcalc/agr084
- 639 Birak, K. S., Terry, P., & Higgs, S. (2010). Effect of cues associated with an alcoholic beverage
 640 on executive function. *Journal of Studies on Alcohol and Drugs*, 71(4), 562–9.
- 641 Cooper, M. (1994). Motivations for alcohol use among adolescents: Development and validation
 642 of a four-factor model. *Psychological Assessment*, 6(2), 117–128.
- 643 Cousijn, J., Goudriaan, A. E., & Wiers, R. W. (2011). Reaching out towards cannabis: approach-
 644 bias in heavy cannabis users predicts changes in cannabis use. *Addiction (Abingdon,*
 645 *England)*, 106(9), 1667–74. doi:10.1111/j.1360-0443.2011.03475.x
- 646 Curtin, J. J., Barnett, N. P., Colby, S. M., Rohsenow, D. J., & Monti, P. M. (2005). Cue reactivity
 647 in adolescents: measurement of separate approach and avoidance reactions. *Journal of*
 648 *Studies on Alcohol*, 66(3), 332–43.
- 649 De Jong, R., Gladwin, T. E., & 't Hart, B. M. (2006). Movement-related EEG indices of
 650 preparation in task switching and motor control. *Brain Research*, 1105(1), 73–82.
 651 doi:10.1016/j.brainres.2006.03.030
- 652 Deiber, M.-P., Sallard, E., Ludwig, C., Ghezzi, C., Barral, J., & Ibañez, V. (2012). EEG alpha
 653 activity reflects motor preparation rather than the mode of action selection. *Frontiers in*
 654 *Integrative Neuroscience*, 6, 59. doi:10.3389/fnint.2012.00059
- 655 Doyle, L. M. F., Yarrow, K., & Brown, P. (2005). Lateralization of event-related beta
 656 desynchronization in the EEG during pre-cued reaction time tasks. *Clinical*
 657 *Neurophysiology : Official Journal of the International Federation of Clinical*
 658 *Neurophysiology*, 116(8), 1879–88. doi:10.1016/j.clinph.2005.03.017
- 659 Duka, T., & Townshend, J. M. (2004). The priming effect of alcohol pre-load on attentional bias
 660 to alcohol-related stimuli. *Psychopharmacology*, 176(3-4), 353–61. doi:10.1007/s00213-
 661 004-1906-7
- 662 Eberl, C., Wiers, R. W., Pawelczack, S., Rinck, M., Becker, E. S., & Lindenmeyer, J. (2013).
 663 Approach bias modification in alcohol dependence: do clinical effects replicate and for
 664 whom does it work best? *Developmental Cognitive Neuroscience*, 4, 38–51.
 665 doi:10.1016/j.dcn.2012.11.002
- 666 Field, M., Christiansen, P., Cole, J., & Goudie, A. (2007). Delay discounting and the alcohol
 667 Stroop in heavy drinking adolescents. *Addiction (Abingdon, England)*, 102(4), 579–86.
 668 doi:10.1111/j.1360-0443.2007.01743.x

- 669 Field, M., Kiernan, A., Eastwood, B., & Child, R. (2008). Rapid approach responses to alcohol
670 cues in heavy drinkers. *Journal of Behavior Therapy and Experimental Psychiatry*, 39(3),
671 209–18. doi:10.1016/j.jbtep.2007.06.001
- 672 Field, M., Schoenmakers, T., & Wiers, R. (2008). Cognitive processes in alcohol binges : A
673 review and research agenda. *Curr Drug Abuse Rev*, 1(3), 263–279.
- 674 Gladwin, T. E., 't Hart, B. M., & de Jong, R. (2008). Dissociations between motor-related EEG
675 measures in a cued movement sequence task. *Cortex*, 44(5), 521–36.
676 doi:10.1016/j.cortex.2007.10.005
- 677 Gladwin, T. E, Lindsen, J., & de Jong, R. (2006). Pre-stimulus EEG effects related to response
678 speed, task switching and upcoming response hand. *Biological Psychology*, 72(1), 15–34.
679 doi:10.1016/j.biopsycho.2005.05.005
- 680 Gratton, G., Coles, M. G. H., & Donchin, E. (1983). A new method for off-line removal of ocular
681 artifact. *Electroencephalography and Clinical Neurophysiology*, 55, 468–484.
- 682 Hodgson, R., Rankin, H., & Stockwell, T. (1979). Alcohol dependence and the priming effect.
683 *Behaviour Research and Therapy*, 17(4), 379–87.
- 684 Houben, K., & Wiers, R. W. (2006). Assessing implicit alcohol associations with the Implicit
685 Association Test: fact or artifact? *Addictive Behaviors*, 31(8), 1346–62.
686 doi:10.1016/j.addbeh.2005.10.009
- 687 Korucuoglu, O., Gladwin, T. E., & Wiers, R. W. (2014). Preparing to approach or avoid alcohol:
688 EEG correlates, and acute alcohol effects. *Neuroscience Letters*, 559, 199–204.
689 doi:10.1016/j.neulet.2013.12.003
- 690 Love, A., Es, D. J. A. M., & Willner, P. (1998). A comparison of two alcohol craving
691 questionnaires. *Addiction*, 93(7), 1091–1102.
- 692 Nam, C. S., Jeon, Y., Kim, Y.-J., Lee, I., & Park, K. (2011). Movement imagery-related
693 lateralization of event-related (de)synchronization (ERD/ERS): motor-imagery duration
694 effects. *Clinical Neurophysiology : Official Journal of the International Federation of*
695 *Clinical Neurophysiology*, 122(3), 567–77. doi:10.1016/j.clinph.2010.08.002
- 696 Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J.M. (2011). FieldTrip: Open source software
697 for advanced analysis of MEG, EEG, and invasive electrophysiological data. *Computational*
698 *Intelligence and Neuroscience*, 156869. doi:10.1155/2011/156869
- 699 Peeters, M., Monshouwer, K., van de Schoot, R. A., Janssen, T., Vollebergh, W. A., & Wiers, R.
700 W. (2013). Automatic processes and the drinking behavior in early adolescence: a
701 prospective study. *Alcoholism, Clinical and Experimental Research*, 37(10), 1737–1744.
702 doi:10.1111/acer.12156
- 703 Pfurtscheller, G., Neuper, C., Pichler-Zalaudek, K., Edlinger, G., & Lopes da Silva, F. H. (2000).
704 Do brain oscillations of different frequencies indicate interaction between cortical areas in
705 humans? *Neuroscience Letters*, 286(1), 66–8.

- 706 Poljac, E., & Yeung, N. (2014). Dissociable neural correlates of intention and action preparation
707 in voluntary task switching. *Cerebral Cortex*, 24(2), 465–478. doi:10.1093/cercor/bhs326
- 708 Robinson, T.E., & Berridge, K.C. (1993) The neural basis of drug craving: an incentive-
709 sensitization theory of addiction. *Brain Res Rev* 18(3):247-291.
710
- 711 Robinson, T.E., & Berridge, K.C. (2008) The incentive sensitization theory of addiction: some
712 current issues. *Philos Trans R Soc Lond Biol Sci* 363(1507): 3137-3146.
- 713 Saunders, J. B., Aasland, O. G., Babor, T. F., de la Fuente, J. R., & Grant, M. (1993).
714 Development of the Alcohol Use Disorders Identification Test (AUDIT): WHO
715 Collaborative Project on Early Detection of Persons with Harmful Alcohol Consumption--
716 II. *Addiction (Abingdon, England)*, 88(6), 791–804.
- 717 Schoenmakers, T. M., & Wiers, R. W. (2010). Craving and attentional bias respond differently to
718 alcohol priming: a field study in the pub. *European Addiction Research*, 16(1), 9–16.
719 doi:10.1159/000253859
- 720 Schoenmakers, T., Wiers, R. W., & Field, M. (2008). Effects of a low dose of alcohol on
721 cognitive biases and craving in heavy drinkers. *Psychopharmacology*, 197(1), 169–78.
722 doi:10.1007/s00213-007-1023-5
- 723 Stancák, A., & Pfurtscheller, G. (1995). Desynchronization and recovery of β rhythms during
724 brisk and slow self-paced finger movements in man. *Neuroscience Letters*, 196, 21–24.
- 725 Thush, C., & Wiers, R. (2007). Explicit and implicit alcohol-related cognitions and the prediction
726 of future drinking in adolescents. *Addictive Behaviors*, 32, 1367–83.
727 doi:10.1016/j.addbeh.2006.09.011
- 728 Thush, C., Wiers, R., Ames, S., Grenard, J., Sussman, S., & Stacey, A. (2008). Interactions
729 between implicit and explicit cognition and working memory capacity in the prediction of
730 alcohol use in at-risk adolescents. *Drug and Alcohol ...*, 94(1-3), 116–124.
731 doi:10.1016/j.drugalcdep.2007.10.019.
- 732 Thush, C., Wiers, R. W., Ames, S. L., Grenard, J. L., Sussman, S., & Stacy, A. W. (2007). Apples
733 and oranges? Comparing indirect measures of alcohol-related cognition predicting alcohol
734 use in at-risk adolescents. *Psychology of Addictive Behaviors : Journal of the Society of*
735 *Psychologists in Addictive Behaviors*, 21(4), 587–91. doi:10.1037/0893-164X.21.4.587
- 736 Watson, D., Clark, L. A., & Tellegen, A. (1988). Development and validation of brief measures
737 of positive and negative affect: the PANAS scales. *Journal of Personality and Social*
738 *Psychology*, 54(6), 1063–70.
- 739 Wiers, R., Rinck, M., Dictus, M., & van den Wildenberg, E. (2009). Relatively strong automatic
740 appetitive action-tendencies in male carriers of the OPRM1 G-allele. *Genes, Brain, and*
741 *Behavior*, 8(1), 101–6. doi:10.1111/j.1601-183X.2008.00454.x
- 742 Wiers, R. W., Eberl, C., Rinck, M., Becker, E. S., & Lindenmeyer, J. (2011). Retraining
743 automatic action tendencies changes alcoholic patients' approach bias for alcohol and

744 improves treatment outcome. *Psychological Science*, 22(4), 490–7.
 745 doi:10.1177/0956797611400615

746

747

748

Figure Captions

749 **Figure 1. Task illustration**

750 Schematic representation of the Approach Avoidance Task. S1 represents the warning stimulus
 751 and S2 represents the imperative stimulus to which motor response (MR) should be given.
 752 Following the motor response, stimuli becomes bigger or smaller during approach and avoid
 753 action (feedback period), respectively.

754 **Figure 2. Behavioral results**

755 Upper panel: Bias scores for the light and heavy drinkers after placebo and alcohol dose. Positive
 756 scores represent an approach bias and negative scores represent an avoidance bias. Error bars
 757 represent standard errors (SE). Lower panel: Mean reaction times (RT) per trial type for the light
 758 and heavy drinkers after placebo (left) and alcohol dose (right). AvoidSoft: Avoid soft-drink cue
 759 trials, ApproSoft: Approach alcohol cue trials, AvoidAlc: Avoid alcohol cue trials, ApproAlc:
 760 Approach alcohol cue trials.

761 **Figure 3. MRAA**

762 Beta-, mu- and alpha-MRAA bias scores for three successive time points (T1: 0-350ms, T2: 350-
 763 700ms, T3: 700-1000ms) following the presentation of the cue. ¥: indicates differences across
 764 placebo and alcohol conditions at $p \leq .05$. Note that bias scores are calculated, separately for
 765 alcohol and soft drink pictures, by subtracting MRAA scores for pull action from the MRAA
 766 scores for push action. Negative MRAA bias-scores represent relatively higher ERDs for the
 767 approach compared to avoid responses (similar to approach bias based on RT), and positive
 768 MRAA bias-scores represent relatively higher ERDs for avoid compared to approach responses

(similar to avoid bias based on RT). AlcBias: Alcohol-cue MRAA bias, SoftBias: soft-drink-cue MRAA bias.

Figure 4. Correlations

Scatterplots for the TRI Govern sub-scale and the overall beta-MRAA bias scores after an alcohol dose at T2 (350-700ms) and T3 (700-1000ms) and after a placebo dose at T3 (700-1000ms). Positive overall MRAA bias scores represent relatively higher ERDs for the soft-drink bias (and greater lateralization for the approach soft-drink bias relative to approach alcohol bias) and negative overall MRAA bias scores represent relatively higher ERDs for the alcohol bias (and greater lateralization for the approach alcohol bias relative to approach soft-drink bias).

Figure 5. Predictors of Alcohol Use

a) Hierarchical multiple regression analysis for variables predicting AUDIT at 6-month follow-up (n=28). **b)** Scatterplots between change in AUDIT scores ($AUDIT_{\text{follow-up, last 90 days}} - AUDIT_{\text{baseline, last 90 days}}$) and (from left to right) alcohol-induced changes on the overall RT bias (alcohol minus soft-drink cue RT bias) and alcohol-induced changes on alcohol-cue and the control-cue (soft-drink-cue) RT bias, separately.

Table 1. Demographic information for the light and heavy drinking groups.

Variable	Light (n=18)	Heavy (n=15)	Light vs. Heavy (p-values)
Age (mean, SD)	18(1.19)	17.4(1.24)	.167
Sex (M/F)	6/12	7/8	-
Current Education Level†			
High school (Level1/Level2/Level3)	1/1/7	-/4/1	-
Tertiary Education (Level1/Level2/Level3)	3/4/2	3/4/3	-
AUDIT(mean, SD)	5.33(2)	13.87(3.14)	<.001
Smoking? (lifetime) (Yes/No, frequency)*	10/7, 31-40 times	14/1, 61-70 times	-
Drug Use (lifetime)*			
Marijuana (Yes/No, frequency)	10/7, 11-20 times	13/2, 31-40 times	-
Ecstasy (Yes/No, frequency)	0/17	4/11, 1-10 times	-
Hallucinogens (Yes/No, frequency)	1/16, 1-10 times	2/13, 1-10 times	-
Hallucinogens (Yes/No, frequency)	0/17	3/12, 1-10 times	-
Stimulants (Yes/No, frequency)	1/16, 1-10 times	4/11, 1-10 times	-
Volatile Substances (Yes/No, frequency)			
RAPI (last 3 months)	1.72(1.52)	5(4.07)	.003
RAPI (lifetime)	6.17(6.01)	14.67(6.02)	<.001
TRI			
Govern	4.11(2.63)	9.53(4.19)	<.001
Restrict	8.83(4.96)	13.8(5.43)	.01
Emotion	4.78(3.3)	9.6(5.05)	.002
Concern	6.33(4.65)	6.73(4.43)	.803
Cognitive	3.39(1.65)	5.53(2.72)	.009
Total	27.44(14.08)	45.2(14.62)	.001
DMQR			
Social	15.44(3.96)	17.33(2.87)	.134
Coping	6.83(1.51)	10(4.49)	.001
Enhancement	12.44(4.38)	14.13(4.88)	.303
Conformity	6.56(2.09)	6.33(1.84)	.751
Total	41.28(9.18)	47.8(8.98)	.049

High School Level1: VMBO, Level2: HAVO, Level3: VWO, Tertiary Education Level1: MBO, Level2: HBO, Level3: WO. † Units of measurement: total number of subjects. * One light drinkers smoking and drug use information was missing.

Table 2. List of variables used in the statistical analysis, their calculations and interpretations.

	Interpretation		
	Calculation	Positive Score	Negative Score
Cue-specific biases *			
Alcohol-cue RT bias	RT in push alcohol cue trials – RT in pull alcohol cue trials	Approach bias for alcohol cues	Avoid bias for alcohol cues
Soft-drink-cue RT bias	RT in push soft-drink cue trials – RT in pull soft-drink cue trials	Approach bias for soft-drink cues	Avoid bias for soft-drink cues
Alcohol-cue MRAA bias	MRAA in pull alcohol cue trials – MRAA in push alcohol cue trials	Relatively greater ERD for avoid than approach alcohol response, i.e. avoid-related lateralization for alcohol cues	Relatively greater ERD for approach than avoid alcohol response, i.e. approach-related lateralization for alcohol cues
Soft-drink-cue MRAA bias	MRAA in pull soft-drink cue trials – MRAA in push soft-drink cue trials	Relatively greater ERD for avoid than approach soft-drink response; i.e. avoid-related lateralization for soft-drink cues	Relatively greater ERD for approach than avoid soft-drink response, i.e. approach-related lateralization for soft-drink cues
Overall biases †			
Overall MRAA bias	Alcohol-cue MRAA bias – Soft-drink-cue MRAA bias	Relatively greater soft-drink-cue MRAA bias than alcohol-cue MRAA bias; i.e. greater lateralization for soft-drink bias	Relatively greater alcohol-cue MRAA bias than soft-drink-cue MRAA bias; i.e. greater lateralization for alcohol bias
Contrasts capturing alcohol-induced effects ¥			
RT dose-contrast score	Overall RT bias in alcohol dose – Overall RT bias in placebo dose	Relatively stronger approach alcohol-cue RT bias score in alcohol dose than in placebo dose	Relatively stronger avoid alcohol-cue RT bias score in alcohol dose than in placebo dose
MRAA dose-contrast score	Overall MRAA bias in alcohol dose – Overall MRAA bias in placebo dose	Relatively stronger approach alcohol-cue MRAA bias score in alcohol dose than in placebo dose	Relatively stronger avoid alcohol-cue MRAA bias score in alcohol dose than in placebo dose

*: list of variables used in repeated measures ANOVA; †: list of variables used in correlational analysis; ¥: list of variables used to test alcohol-induced effects as predictor of future alcohol escalation. Variable in grey was not discussed in the paper due to lack of a statistical effect, but listed here only for the completeness. Movement-related amplitude asymmetry (MRAA) is quantified by taking the difference in amplitude between contralateral and ipsilateral activity during preparation of right-hand response minus difference between ipsi- and contra-lateral activity during preparation of left-hand response (Gladwin et al., 2006). Given that a decrease in power (ERD) is expected for the hemisphere contralateral to the movement, more negative MRAA values indicate greater motor-related lateralization due to increased ERD contralateral to the movement.

Table 3. Mean scores and standard deviations for the BrAC and the Brief Biphasic Alcohol Effects Scale (B-BAES) before (pre-task) and after (post-task) participants completed the alcohol-Approach-Avoidance Task in the placebo and in the alcohol condition (n=33).

	Pre-AAT	Post-AAT
<i>BAL (g/l, [Mean (SD)])</i>	.55(.4)	.46(.15)
<i>B-BAES Stimulation subscale</i>		
Placebo [Mean (SD)]	18.15(5.72)	17.09(5.8)
Alcohol [Mean (SD)]	17.39(4.87)	16.12(5.7)
<i>B-BAES Sedation subscale</i>		
Placebo [Mean (SD)]	11.64(5.32)	12.(4.43)
Alcohol [Mean (SD)]	13.24(5.6)	14.24(5.49)

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

	B	SE B	Beta	P – values
a) Step1: change R^2: .705, $p < .001$				
AUDIT baseline	.676	.086	.84	<.001
Step2: change R^2: .064, $p = .014$				
AUDIT baseline	.679	.077	.844	<.001
RT Contrast	.013	.005	.253	.014
Step3: change R^2: .046, $p = .023$				
AUDIT baseline	.643	.072	.798	<.001
RT dose-contrast	.016	.005	.305	.003
Parietal alpha-MRAA dose-contrast	-.001	.00026	.225	.023

RT dose-contrast: (Alcohol-cue RT bias minus Soft-drink-cue RT bias) in alcohol dose - (Alcohol-cue RT bias minus Soft-drink-cue RT bias) in placebo dose. MRAA dose-contrast: (Alcohol-cue MRAA bias minus Soft-drink-cue MRAA bias) in alcohol dose - (Alcohol-cue MRAA bias minus Soft-drink-cue MRAA bias) in placebo dose. Audit baseline refers to the alcohol challenge session.