1 **Weekend working: A retrospective cohort study of maternal and**
2 **neonatal outcomes in a large NHS delivery unit**

3
4 Catherine E Aiken¹ *, Abigail R Aiken², James G Scott³, Jeremy C Brockelsby¹,
5 James Trussell²

6
7 ¹Department of Obstetrics and Gynaecology, University of Cambridge; NIHR
8 Cambridge Comprehensive Biomedical Research Centre, CB2 2SW, UK

9 ² Office of Population Research, Princeton University; Princeton, NJ, USA, 08544

10 ³ McCombs School of Business, University of Texas at Austin, Texas, USA, 78712

11

12 ***Correspondence to:** Email: cema2@cam.ac.uk, Telephone: +44(0)1223 336871,

13 Address as above

14

15

16

17

18 **Condensation**

19 Adopting mandatory 7-day working contracts in the UK National Health Service is
20 unlikely to make any difference to consultant presence during the weekend or to
21 maternal or neonatal morbidity.

22

Weekend working: A retrospective cohort study of maternal and neonatal outcomes in a large NHS delivery unit

Catherine E Aiken, Abigail R Aiken, James G Scott, Jeremy C Brockelsby & James Trussell

Abstract (260)

Objectives: Mandatory weekend working for NHS consultants is currently the subject of intense political debate. The Secretary of State for Health's proposed 7-day contract policy is based on the claim that such working patterns will improve patient outcomes. We evaluate this claim by taking advantage of as-if-at-random presentation of women for non-elective deliveries throughout the week. We examine (i) whether consultants currently perform fewer deliveries during weekends *versus* weekdays, and (ii) whether adverse outcomes increase during weekends

Study Design: We conducted a retrospective cohort study using data on all non-elective deliveries from January 2008-December 2013 in a large UK obstetrics center (n=27,466). We used Pearson's chi-squared tests to make direct comparisons of adverse outcome rates during weekdays *versus* weekends. Outcomes included: estimated maternal blood loss ≥ 1.5 litres; severe perineal trauma; delayed neonatal respiration; umbilical arterial pH < 7.1 ; and critical incidents at delivery.

Results: Consultants currently perform the same proportion of non-elective deliveries on weekends and weekdays (2.3% *versus* 2.6%, $p = 0.25$). We found no increase in any adverse maternal or neonatal outcomes during weekends *versus* weekdays, despite high statistical power to detect such differences. Moreover, adverse outcomes

are no higher during periods of the weekend when consultants are not routinely present compared to equivalent periods during weekdays.

Conclusions: Under current working arrangements, women who would benefit from consultant-led delivery are equally likely to receive one on weekends compared to weekdays. Weekend delivery has no effect on maternal or neonatal morbidity. Adopting mandatory 7-day contracts is unlikely to make any difference to either consultant-led delivery during weekends or to patient outcomes.

Keywords: weekend working; consultants; delivery outcomes; health policy; maternity services

59 **Introduction**

60 Increased risk of adverse events during weekends compared to weekdays in the UK
61 National Health Service (NHS) has long been a concern of doctors, patients, and
62 policy-makers alike (1). This topic recently came into the public spotlight because of
63 remarks made by the Secretary of State for Health, Jeremy Hunt: “Around 6,000
64 people lose their lives every year because we do not have a proper 7-day service in
65 hospitals” (2). Mr Hunt further argued that requiring mandatory weekend-working
66 contracts for consultants would increase their presence in hospitals during weekends
67 and reduce these additional deaths. These remarks are echoed by current policy
68 recommendations to improve NHS services by reconfiguring consultants’ working
69 hours (1).

70
71 Yet the presumed causal link between consultant working patterns and higher rates of
72 adverse clinical outcomes is far from clear-cut. We aim to evaluate this link using
73 data on consultants working within maternity services, which are a touchstone for the
74 provision of safe and high-quality care across the NHS (3). Specifically, we examine
75 the risks of adverse outcomes arising from non-elective deliveries in a large UK
76 centre. We compare complication rates during weekdays and weekends to determine
77 (i) whether consultants perform fewer deliveries during weekends than during
78 weekdays, and (ii) whether rates of adverse outcomes increase during weekends.

79
80 Previous studies examining rates of neonatal deaths during weekends have
81 demonstrated higher rates outside 09.00-17.00 on weekdays than at other times (4).
82 However, studies specifically comparing weekends to weekdays suggest no
83 differences in neonatal death rates (5-7). Aside from neonatal mortality, there is little

evidence regarding rates of other serious adverse outcomes during weekends compared to weekdays, despite their potentially profound impacts on women and infants.

Our design takes advantage of several important features of obstetric data. First, delivery is a clearly defined, high-risk event at which the presence of a consultant could potentially reduce the risk of adverse outcomes (4). Second, by limiting our focus to non-elective deliveries, our sample is plausibly distributed as-if-at-random between weekend and weekdays, since these women have not chosen when to deliver. This strategy avoids possible selection bias, where the weekend patient population differs from the weekday population in ways that are likely related to the risk of adverse outcomes. Third, the obstetric consultants in our sample have a clear and consistent working pattern throughout the study period, allowing establishment of a reliable link between day and time of delivery and the presence of a consultant.

Methods

32,078 deliveries occurred during a 6-year period (January 2008 - December 2013) in a single large NHS maternity unit in the UK. Elective deliveries were excluded, as they are overwhelmingly more likely to occur during weekdays and carry a substantially lower risk of adverse outcomes. We identified a sub-cohort of 27,466 non-elective deliveries that occurred by spontaneous, instrumental delivery or non-elective Caesarean section for analysis. Inductions of labour were included, as initial analysis determined that these were no more likely to deliver during weekdays than at weekends. Spontaneous vaginal deliveries performed by midwives were also included since senior obstetricians may significantly influence decision-making and

management during these deliveries. We also present results for a second separate sub-cohort of operative deliveries (both instrumental vaginal deliveries and non-elective Caesarean sections, $n = 9,010$), as the outcomes of these deliveries are the most likely to be directly influenced by the presence of a consultant obstetrician.

In the study centre, 3 doctors are available for emergency work on the delivery unit at any given time. The difference in direct consultant presence on the delivery unit between weekends and weekdays is limited to the hours of 12.00 – 19.00. Outside of these times, the consultant is either present at the same times as during the weekdays (08.00 - 12.00) or is not present at either the weekends or weekdays (19.00 - 08.00).

We therefore identified a third sub-cohort of non-elective deliveries that occurred between 12.00 and 19.00 ($n = 7,361$) to allow separate analysis of outcomes during the time-period when no consultant is directly present during the weekends, but would have been on a weekday. No consultant opted out of weekend duty during the study period.

Study data were obtained from an electronic maternity data-recording system, which is updated by midwives shortly after delivery. The database is regularly validated by a rolling program of audits, where the original case notes are checked against the information recorded. No patient-identifiable data were accessed in the course of this research, which was performed as part of a provision-of-service study for the obstetrics centre. Individual medical records were not accessed at any stage, and the study was therefore deemed exempt from full institutional review board approval.

Data obtained on delivery characteristics included maternal age in years (at time of delivery), BMI (measured at first trimester prenatal booking), parity (prior to delivery), and the birth-weight of the infant (recorded to the nearest gram). Gestational age was determined from first trimester ultrasound and recorded to the nearest week. Deliveries were classified as either spontaneous onset or induced. The healthcare professional delivering the baby was either a midwife or a doctor classified by years of specific obstetric training at the time of the delivery. Categories of experience were: ≤ 2 years (including those in the second year of foundation training, vocational general practitioner training, or the first 2 years of specialty training); 3-5 years (including both doctors in years 3-5 inclusive of their specialty training and those of equivalent experience not enrolled in a specialty training programme); > 5 years (doctors in years 6/7 of the specialty training programme or those of equivalent or greater experience not employed as NHS consultants); and consultants (all of whom must have a minimum of 7 years obstetric training). Delivery type was classified as elective Caesarean section, emergency Caesarean section, instrumental delivery (sub-classified as forceps or ventouse) and vaginal deliveries (sub-classified as either breech or cephalic). Elective Caesarean deliveries were excluded from the analysis.

Outcome data on maternal and neonatal complications were obtained from the same database. Delay in neonatal respiration was defined as no spontaneous neonatal respiration within 1 minute of delivery. Where the healthcare professional performing delivery deemed it necessary (typically all non-elective operative deliveries and those involving concern about neonatal well-being before delivery or at birth), the pH of umbilical arterial blood was tested immediately following delivery. Umbilical arterial

pH was categorized as ≥ 7.1 or < 7.1 (8). A critical-incident form was generated at delivery in the case of any obstetric or neonatal emergency, including maternal death, full neonatal resuscitation, shoulder dystocia, maternal visceral injury or any other event triggering an obstetric emergency call. Maternal blood loss was estimated as soon as possible after delivery. Estimated blood loss was categorized as < 1.5 litres or ≥ 1.5 litres. Severe maternal perineal trauma was defined as any third or fourth degree tear.

Standard significance tests were used to assess whether patients delivering at the weekend *versus* weekdays exhibited any imbalances in risk factors for adverse neonatal and maternal outcomes. A two-sided, two-sample t-test with unequal sample sizes was used for each continuous numerical risk factor (maternal age, maternal BMI, gestational age, and birth weight). A Pearson chi-squared test was used for each categorical risk factor (parity, race of the mother, delivery type, induction of labor, and the delivering healthcare professional).

All five adverse outcomes analysed are binary events. Complication rates on weekends versus weekdays were compared using two-sample tests of proportions with unequal sample sizes. For each outcome, a one-sided test was conducted, in which the alternative hypothesis is that the adverse-outcome rate is higher on the weekend than on the weekday. Compared with a two-sided test, this allowed greater power to detect excess complications for weekend deliveries.

Power calculations were performed for all comparisons of adverse-outcome rates.

For each test, the minimum detectable effect size was calculated: that is, the smallest

effect size (Δ) that could be detected at a significance level of 0.05 with power of at least 80%. These effect sizes are expressed as an absolute difference in rates (e.g. 4.9% on weekends versus 4.8% on weekdays is a $\Delta = 0.1\%$ effect size). These power calculations were initially performed using the standard Gaussian approximation to the binomial test but were also verified using Monte Carlo simulation. The Monte Carlo simulations showed slightly lower power than the Gaussian approximation. In our results, we therefore quote the more conservative numbers from the Monte Carlo simulations. Based on our findings of no statistically significant differences in any adverse outcomes between deliveries during weekends and weekdays, no corrections for multiplicity in our assessments of statistical significance were required. Correcting these p-values for multiplicity could only make them appear less significant, meaning that would be impossible for such a correction to materially change our findings.

All data analyses were conducted using the R statistical software package version 3.2.0 (9). Findings were considered statistically significant at an alpha level of 0.05. An R script containing code for all adverse-outcome comparisons and power calculations is available as a supplemental file.

Results

There were no significant differences in the maternal, neonatal or delivery-related characteristics for non-elective deliveries occurring on weekdays ($n = 19,626$) compared to those occurring at weekends ($n = 7,840$) (Table 1) and no difference in the total number of non-elective deliveries that occurred on any day. This finding suggests that cases of broadly similar clinical difficulty present during weekdays and

weekends and that comparisons of complication rates are not prone to any obvious source of confounding. There were no differences in the rates of any adverse outcomes for non-elective deliveries that occurred during the weekdays compared to the weekends (Table 2). Our power calculations demonstrate that for all non-elective deliveries, the minimum detectable effect sizes range from 0.5% (for estimated blood loss) to 1.2% (for arterial umbilical pH < 7.1). These minimum detectable effect sizes can be interpreted as a likely upper bound on the magnitude of any discrepancy between the weekend and weekday rates. When non-elective operative deliveries performed by doctors only were considered (n=9,010), none of the rates of adverse outcomes at weekends were significantly different from those occurring during the weekdays (Table 2). In this sub-cohort, the minimum detectable effect sizes range from 1.1% (for estimated blood loss) to 1.8% (for delayed neonatal respiration).

Deliveries were equally likely to be performed by consultants at weekends as during the weekdays: 508/19,626 (2.3%) v. 184/7,840 (2.6%), p=0.25. The characteristics of mothers, neonates and deliveries were not significantly different during afternoons during weekdays (when consultants were routinely present) compared to weekends (when consultants were not routinely present) (Table 3). There was no increase in the rates of any adverse outcomes during the afternoon period at weekends, compared to during weekdays (Table 4). The minimum detectable effect sizes for this analysis range from 1.0% (for estimated blood loss) to 2.5% (for low arterial umbilical pH).

Discussion

We present evidence that serious adverse delivery events within NHS maternity services are not increased at the weekend compared to weekdays. The study cohort is

well powered for all outcomes examined and allows direct comparison of outcomes during consultant presence with times when no consultant was present. No differences were found in any of the adverse outcomes studied, either in all non-elective deliveries or in those undergoing non-elective operative delivery. Moreover, despite consultants being routinely present on the delivery unit for fewer hours at weekends, the proportion of babies delivered by consultants did not decrease at the weekends. Specific examination of the period when consultants would be additionally present if their working patterns were identical during weekdays and weekends (12.00-19.00 on Saturday and Sunday) revealed no increase in the rates of any adverse outcomes.

Our results accord with those from a large, recent North American cohort, which reported no increase in rates of pelvic morbidity (including perineal trauma as defined here) or other severe maternal morbidity on weekends when compared to weekdays (10). By contrast, a recent UK study found that there was an increase in perinatal mortality and maternal infection on weekend days (11). This study took the unusual step of comparing weekend days to those deliveries occurring on Tuesdays only, rather than a comparison over all weekdays. Furthermore, all stillbirths, including those where the death occurred antepartum, were attributed to the day of delivery rather than the day of death. These methodological steps may account for the differences in the detection of a 'weekend effect' in the previous study, although we do not find a similar effect in our cohort. Direct comparisons on all outcomes between studies are precluded by the differences in outcomes assessed.

Our study has several important limitations. The study was not powered to detect an increase in maternal mortality or to consider neonatal mortality, except within the

composite outcome of ‘critical incidents’. Other studies, however, suggest that neonatal deaths may not be significantly increased during weekends (5-7), although they are higher outside of 09.00-17.00 weekdays than at other times (4). Maternal death rates in the UK are as low as 1/10,000 (12). Although neonatal deaths are more common, with a perinatal extended death rate at 6/1,000, this figure includes antepartum stillbirths, which would not be affected by weekend consultant working patterns (13). While our conclusions are applicable only to non-elective deliveries, elective deliveries account for only a small proportion of all deliveries in the UK and are not scheduled over weekends, thus precluding weekday versus weekend comparisons.

A further limitation of our study is that data from a single site may not be generalizable to other dissimilar populations. However, the characteristics of our population (including maternal age, birth weight and mode of delivery) are similar to those of maternity service users elsewhere in England (14), implying that our results are likely applicable to a high proportion of maternity services. Indeed, one advantage of a single site design is that it ensures detailed reporting of serious adverse events other than mortality, which are less well captured in other cohorts. While our study is focused exclusively on outcomes from maternity services and thus is of particular relevance to obstetricians, we suggest that delivery outcome data represent a good model for the multitude of high-risk emergency services that the NHS provides to the general population on a short-term basis.

Consultant involvement in care is certainly important in reducing adverse outcomes both within obstetric services across the NHS overall (15). However, we find no

evidence in support of a causal link between consultant contractual obligations and higher rates of adverse clinical outcomes. Across the NHS as a whole, mortality rates for patients admitted during the weekend are higher than for those admitted during the week (16, 17), and similar trends have been observed among patients presenting for acute emergency care (18). But, the weekend patient population in most specialties differs from the weekday population in ways that are likely related to the risk of adverse outcomes—for example, major trauma is most likely to occur on Saturday night (19). While some evidence suggests that increased consultant presence could reduce the weekend fatality rate for acute medical inpatients (20), other evidence bears out the selection-bias hypothesis: among non-acute-emergency patients, the association between weekend admission and increased mortality does not hold true for all conditions (21, 22), and it is stronger for conditions with higher baseline mortality rates (22, 23). Our study also highlights the importance of considering adverse outcomes other than mortality. Although overall mortality within 30 days after emergency admission for high-risk conditions in England is 5.59% (24), a high proportion of patients seeking non-elective weekend care in the NHS do so for conditions with very low baseline rates of mortality, including the users of maternity services.

In contrast to the current policy change proposed by the Secretary of State for Health, our findings imply that mandatory ‘7-day working’ by consultants is unlikely to have an impact on the rates of common adverse outcomes in maternity care. Within the current system, consultants appear to be readily available for deliveries if and when required. In light of multiple competing demands on NHS finances (25), restructuring

working patterns to mandate continuous consultant presence at weekends is unlikely to be either an effective or efficient use of resources to improve patient care.

Acknowledgements

ARA and JT are supported by grant P2CHD047879, awarded to the Office of Population Research at Princeton University by The Eunice Kennedy Shriver National Institute of Child Health and Human Development of the National Institutes of Health. JGS is supported by a CAREER grant from the U.S. National Science Foundation (DMS-1255187). None of the sponsors played any role in the collection, analysis or interpretation of the data, or and in the writing of the manuscript.

Disclosures of Interest

The authors have no conflicts of interest to declare.

References

- (1) NHS Services, seven days a week forum: Evidence base and clinical standards for the care and onward transfer of acute inpatients. Gateway reference : 00889.
- (2) Department of Health and the Rt Hon Jeremy Hunt MP. Making healthcare more human-centred and not system-centred. 16 July 2015, King's Fund, London, UK <https://www.gov.uk/government/speeches/making-healthcare-more-human-centred-and-not-system-centred>.
- (3) Department of Health, 2009, The Year: NHS Chief Executive's Annual Report. London: Department of Health http://www.healthcare-today.co.uk/doclibrary/documents/pdf/215_the_year_2008-09.pdf.

- 331 (4) Pasupathy D, Wood AM, Pell JP, Fleming M, Smith GC. Time of birth and
332 risk of neonatal death at term: retrospective cohort study. *BMJ*. 2010;341:c3498.
- 333 (5) Stephansson O, Dickman PW, Johansson AL, Kieler H, Cnattingius S. Time
334 of birth and risk of intrapartum and early neonatal death. *Epidemiology*. 2003
335 Mar;14(2):218-22.
- 336 (6) Gijzen R, Hukkelhoven CW, Schipper CM, Ogbu UC, de Bruin-Kooistra M,
337 Westert GP. Effects of hospital delivery during off-hours on perinatal outcome in
338 several subgroups: a retrospective cohort study. *BMC Pregnancy Childbirth*.
339 2012;12:92.
- 340 (7) Dowding VM, Duignan NM, Henry GR, MacDonald DW. Induction of
341 labour, birthweight and perinatal mortality by day of the week. *Br J Obstet Gynaecol*.
342 1987 May;94(5):413-9.
- 343 (8) Yeh P, Emary K, Impey L. The relationship between umbilical cord arterial
344 pH and serious adverse neonatal outcome: analysis of 51,519 consecutive validated
345 samples. *BJOG*. 2012 Jun;119(7):824-31.
- 346 (9) R Core Team (2014). R: A language and environment for statistical
347 computing. R Foundation for Statistical Computing, Vienna, Austria. URL
348 <http://www.R-project.org/>.
- 349 (10) Lyndon A, Lee HC, Gay C, Gilbert WM, Gould JB, Lee KA. Effect of time of
350 birth on maternal morbidity during childbirth hospitalization in California. *Am J*
351 *Obstet Gynecol*. 2015 Nov;213(5):705 e1- e11.
- 352 (11) Palmer WL, Bottle A, Aylin P. Association between day of delivery and
353 obstetric outcomes: observational study. *BMJ*. 2015;351:h5774.
- 354 (12) Knight M, Kenyon S, Brocklehurst P, Neilson J, Shakespeare J, Kurinczuk JJ.
355 Saving Lives, Improving Mothers' Care: Lessons learned to inform future maternity

- 356 care from the UK and Ireland Confidential Enquiries into Maternal Deaths and
 357 Morbidity 2009-2012. MBRRACE-UK, London, UK. December 2014. 2014.
- 358 (13) Manktelow BN, Smith LK, Evans TA, et al. Perinatal Mortality Surveillance
 359 Report: UK perinatal deaths for births from January to December 2013. MBRRACE-
 360 UK, Maternal, Newborn and Infant Clinical Outcome Review Programme. June 2015
 361 2015.
- 362 (14) Hospital Episode Statistics NHS Maternity Statistics 2012-13, Health and
 363 Social Care Information Centre 2013
 364 <http://www.hscic.gov.uk/catalogue/PUB12744/nhs-mate-eng-2012-13-summ-repo->
 365 [rep.pdf](http://www.hscic.gov.uk/catalogue/PUB12744/nhs-mate-eng-2012-13-summ-repo-rep.pdf).
- 366 (15) Academy of Royal Medical Colleges, The Benefits of Consultant Delivered
 367 Care. January 2012.
- 368 (16) Freemantle N, Richardson M, Wood J, et al. Weekend hospitalization and
 369 additional risk of death: an analysis of inpatient data. J R Soc Med. 2012
 370 Feb;105(2):74-84.
- 371 (17) Aylin P, Yunus A, Bottle A, Majeed A, Bell D. Weekend mortality for
 372 emergency admissions. A large, multicentre study. Qual Saf Health Care. 2010
 373 Jun;19(3):213-7.
- 374 (18) Barba R, Losa JE, Velasco M, Guijarro C, Garcia de Casasola G, Zapatero A.
 375 Mortality among adult patients admitted to the hospital on weekends. Eur J Intern
 376 Med. 2006 Aug;17(5):322-4.
- 377 (19) The National Confidential Enquiry into Patient Outcome and Death (2007).
 378 Trauma:
 379 Who cares?, London: UK.

- (20) Bell D, Lambourne A, Percival F, Lavery AA, Ward DK. Consultant input in acute medical admissions and patient outcomes in hospitals in England: a multivariate analysis. PLoS One. 2013;8(4):e61476.
- (21) Ricciardi R, Roberts PL, Read TE, Baxter NN, Marcello PW, Schoetz DJ. Mortality rate after nonelective hospital admission. Arch Surg. 2011 May;146(5):545-51.
- (22) Bell CM, Redelmeier DA. Mortality among patients admitted to hospitals on weekends as compared with weekdays. N Engl J Med. 2001 Aug 30;345(9):663-8.
- (23) Concha OP, Gallego B, Hillman K, Delaney GP, Coiera E. Do variations in hospital mortality patterns after weekend admission reflect reduced quality of care or different patient cohorts? A population-based study. BMJ Qual Saf. 2014 Mar;23(3):215-22.
- (24) Roberts SE, Thorne K, Akbari A, Samuel DG, Williams JG. Weekend emergency admissions and mortality in England and Wales. Lancet. 2015 May 9;385(9980):1829.
- (25) Five Year Forward View. NHS, October 2014.
<http://www.england.nhs.uk/wp-content/uploads/2014/10/5yfv-web.pdf>.

Table legends

Table 1: Characteristics of non-elective deliveries occurring in the study centre (January 2008 - December 2013), by delivery during weekdays or weekends. P values represent the results of comparison of means via Student's t-test (2-tailed, unequal sample size) for continuous variables, and Pearson's chi-squared (2-tailed, unequal sample size) for categorical variables.

405

406 Table 2: Adverse outcomes by delivery during weekdays or weekends. Percentages
407 represent the percentage experiencing the adverse outcome from all deliveries where
408 outcome data were available. P values are calculated using one-tailed Pearson's chi-
409 squared. Δ is the smallest effect size that could be detected at a significance level of
410 0.05 with power of at least 80%.

411

412 Table 3: Characteristics of non-elective deliveries occurring during 12.00 – 19.00 in
413 the study centre (January 2008 - December 2013), by delivery during weekdays or
414 weekends. P values represent the results of comparison of means via Student's t-test
415 (2-tailed, unequal sample size) for continuous variables, and Pearson's chi-squared
416 (2-tailed, unequal sample size) for categorical variables.

417

418 Table 4: Adverse outcomes by delivery during weekdays or weekends, for non-
419 elective deliveries occurring between 12.00 and 19.00. Percentages represent the
420 percentage experiencing the adverse outcome from all deliveries where outcome data
421 were available. P values are calculated using one-tailed Pearson's chi-squared. Δ is
422 the smallest effect size that could be detected at a significance level of 0.05 with
423 power of at least 80%.

Table 1

Characteristic		Weekday deliveries (n= 19,626) Mean $\pm$ SD	Weekend deliveries (n= 7,840) Mean $\pm$ SD	p value
Maternal age (years)		30.3 $\pm$ 5.6	30.4 $\pm$ 5.6	0.15
Maternal BMI		24.7 $\pm$ 5.1	24.8 $\pm$ 5.2	0.11
Gestational age (weeks)		39.2 $\pm$ 2.3	39.2 $\pm$ 2.2	0.99
Birth weight (grams)		3372 $\pm$ 612.6	3370 $\pm$ 616.8	0.78
		Frequency (%)	Frequency (%)	
Parity	0	6226 (32%)	2528 (32%)	0.62
	1	8800 (45%)	3476 (44%)	
	≥ 2	4342 (22%)	1712 (22%)	
	Unknown	258 (1%)	124 (2%)	
Race	Caucasian	17089 (87%)	6819 (87%)	0.42
	South Asian	995 (5%)	387 (5%)	
	Black	344 (2%)	129 (2%)	
	East Asian	298 (2%)	129 (2%)	
Delivery type	SVD	13,309 (68%)	5331 (68%)	0.37
	Caesarean	3317 (17%)	1358 (17%)	
	Instrumental	3000 (15%)	1151 (15%)	
Induction of labour	Yes	5577 (28%)	2202 (28%)	0.25
	No	12910 (66%)	5168 (66%)	
	Unknown	1139 (6%)	470 (6%)	
Delivering healthcare professional	Doctor ≤ 2 years obstetric training	217 (1%)	83 (1%)	0.75
	Doctor 3-5 years obstetric training	4621 (24%)	1881 (24%)	
	Doctor >5 years obstetric training	1400 (7%)	559 (7%)	
	Consultant	508 (3%)	184 (2%)	
	Midwife	12880 (65%)	5133 (66%)	

Table 2

Outcome	All non-elective deliveries (n=27,466)				All non-elective operative deliveries by doctors (n=9010)			
	Weekday	Weekend	p	Δ	Weekday	Weekend	p	Δ
Estimated blood loss ≥1.5 litres	326/1929 8 (1.7%)	123/7678 (1.6%)	0.70	0.5%	194/6204 (3.1%)	78/2460 (3.2%)	0.49	1.1%
Severe maternal perineal trauma	689/1784 2 (3.9%)	286/7122 (4%)	0.29	0.7%	228/4931 (4.6%)	85/1970 (4.3%)	0.71	1.5%
Delay to neonatal respiration	999/1926 2 (5.1%)	409/7840 (5.2%)	0.33	0.8%	614/6440 (9.5%)	224/2570 (8.7%)	0.89	1.8%
Umbilical Arterial pH <7.1	371/7501 (4.9%)	145/2999 (4.8%)	0.59	1.2%	261/5623 (4.6%)	97/2224 (4.4%)	0.71	1.4%
Any critical incident at delivery	742/1877 7 (4%)	289/7451 (3.9%)	0.61	0.7%	423/5948 (7.1%)	166/2342 (7.1%)	0.52	1.7%

Table 3

Characteristic		Weekday deliveries (n= 5,303) Mean $\pm$ SD	Weekend deliveries (n= 2,058) Mean $\pm$ SD	p value
Maternal age (years)		30.3 $\pm$ 5.6	30.3 $\pm$ 5.6	0.74
Maternal BMI		24.7 $\pm$ 5.1	24.7 $\pm$ 5.6	0.99
Gestational age (weeks)		39.1 $\pm$ 2.3	39.1 $\pm$ 2.5	0.38
Birth weight (grams)		3366 $\pm$ 619.7	3345 $\pm$ 666.6	0.22
		Frequency (%)	Frequency (%)	
Parity	0	1815 (35%)	686 (33%)	0.53
	1	2346 (44%)	938 (45%)	
	≥ 2	1066 (20%)	400 (20%)	
	Unknown	76 (1%)	34 (2%)	
Race	Caucasian	4629 (87%)	1773 (86%)	0.56
	South Asian	258 (5%)	119 (5%)	
	Black	83 (2%)	32 (2%)	
	East Asian	79 (1%)	32 (2%)	
Delivery type	SVD	3422 (64%)	1321 (64%)	0.44
	Caesarean	983 (19%)	405 (20%)	
	Instrumental	898 (17%)	332 (16%)	
Induction of labour	Yes	1406 (27%)	513 (25%)	0.34
	No	3529 (67%)	1391 (68%)	
	Unknown	368 (6%)	154 (7%)	
Delivering healthcare professional	Doctor ≤ 2 years obstetric training	78 (2%)	26 (1%)	0.90
	Doctor 3-5 years obstetric training	1384 (26%)	548 (27%)	
	Doctor > 5 years obstetric training	280 (5%)	105 (5%)	
	Consultant	278 (5%)	115 (6%)	
	Midwife	3283 (62%)	1264 (61%)	

Table 4

Outcome	All non-elective deliveries occurring between 12.00 and 19.00 (n=7,361)			
	Weekday	Weekend	p	Δ
Estimated blood loss ≥ 1.5 litres	101/5208 (1.9%)	28/2007 (1.4%)	0.95	1.0%
Severe maternal perineal trauma	188/4756 (3.9%)	75/1823 (4.1%)	0.38	1.5%
Delay to neonatal respiration	312/5303 (5.8%)	126/2058 (6.1%)	0.35	1.6%
Umbilical Arterial pH < 7.1	104/2097 (5%)	33/825 (4%)	0.88	2.5%
Any critical incident at delivery	212/5049 (4%)	70/1948 (3.9%)	0.88	1.5%