Impact of Targeted PPAR γ Disruption on Bone Remodeling

Jay Cao¹, Guomin Ou^{2,#}, Kehong Ding², Nianlan Yang², Barbara E. Kream³, Mark W. Hamrick⁴, Carlos M. Isales^{2,5}, Xing-Ming Shi^{2,5*}

¹USDA ARS Grand Forks Human Nutrition Research Center, Grand Forks, ND, USA

²Department of Neuroscience & Regenerative Medicine, Georgia Regents University, Augusta, GA, USA

³Departments of Medicine, University of Connecticut Health Center, Farmington, CT, USA

Departments of ⁴Cell Biology, ⁵Orthopaedic Surgery, Georgia Regents University, Augusta, GA, USA

[#]Current address: West China College of Stomatology, Sichuan University, Chengdu, China

Address correspondence to: Xing-Ming Shi, Departments of Neuroscience & Regenerative Medicine and Orthopedic Surgery, Georgia Regents University, 1120 15th Street, CA-2003, Augusta, GA 30912, USA. Phone: (706) 721-1097; Fax: (706) 434-7440; Email: xshi@gru.edu

This work was supported by grants from The National Institute on Aging (AG046248 and AG036675) and The National Institute of Diabetes and Digestive and Kidney Diseases (DK76045), National Institutes of Health (NIH).

ABSTRACT

Peroxisome proliferator-activated receptor gamma (PPAR γ), known as the master regulator of adipogenesis, has been regarded as a promising target for new anti-osteoporosis therapy due to its role in regulating bone marrow mesenchymal stem/progenitor cell (BMSC) lineage commitment. However, the precise mechanism underlying PPAR γ regulation of bone is not clear as a bone-specific PPAR γ conditional knockout (cKO) study has not been conducted and evidence showed that deletion of PPAR γ in other tissues also have profound effect on bone. In this study, we show that mice deficiency of PPAR γ in cells expressing a 3.6kb type I collagen promoter fragment (PPAR^{fl/fl}:Col3.6-Cre) exhibit a moderate, site-dependent bone mass phenotype. *In vitro* studies showed that adipogenesis is abolished completely and osteoblastogenesis increased significantly in both primary bone marrow culture and the BMSCs isolated from PPAR γ cKO mice. Histology and histomorphometry studies revealed significant increases in the numbers of osteoblasts and surface in the PPAR γ cKO mice. Finally, we found that neither the differentiation nor the function of osteoclasts was affected in the PPAR γ cKO mice. Together, our studies indicate that PPAR γ plays an important role in bone remodeling by increasing the abundance of osteoblasts for repair, but not during skeletal development.

Keywords: PPAR γ , MSC, aging, osteoblasts, bone loss.

1. INTRODUCTION

Peroxisome proliferator-activated receptor gamma (PPAR γ) is a ligand-activated nuclear receptor and is indispensable for adipocyte differentiation both *in vitro* and *in vivo* (1, 2). Bone marrow mesenchymal stem cell (BMSC) is a common precursor for bone-forming osteoblasts and marrow adipocytes. With advancing age, bone mass decreases and marrow fat increases (fatty marrow) (3-5), indicating that an imbalance develops between these two pathways (6-8). Since the PPAR γ is a key adipogenic regulator and the osteoblast and adipocyte, which are the two dominant pathways for BMSC to differentiate, have a reciprocal relationship (9-11), the PPAR γ has been viewed as a prominent target for new anti-osteoporosis therapies that could increase bone formation or prevent bone loss. Indeed, studies have shown that PPAR γ heterozygous (PPAR $\gamma^{+/-}$) mice exhibited an extraordinary high bone mass phenotype (12). However, global PPAR γ insufficiency reduced body fat mass, therefore it is not clear whether the high bone mass observed in these PPAR γ insufficient mice was due to a direct effect of PPAR γ on BMSC lineage determination or an indirect effect via modulating the functions of adipose tissue or both, since disruption of PPAR γ in adipose tissue (lipodystrophic PPAR $\gamma^{\text{hyp/hyp}}$ mice) also enhanced osteoblastic activity and increased bone formation (13). In this study, we generated mice in which PPAR γ gene is deleted in cells that support the activity of a 3.6kb type I collagen promoter fragment (PPAR $\gamma^{\text{fl/fl}}$:Col3.6Cre) and report here the bone phenotype of these PPAR γ conditional knockout (PPAR γ cKO) mice.

2. MATERIALS AND METHODS

2.1. Animals

PPAR γ cKO mice were created by breeding Col3.6-Cre transgenic mice (14) with PPAR γ floxed (PPAR $\gamma^{fl/fl}$) mice (15) using standard breeding procedures.

2.2. Ethics Statement

Animals were maintained in a centralized barrier facility and all experimental procedures were approved by the Institutional Animal Care and Use Committee (IACUC) at the Georgia Regents University.

2.3. Genotyping

Genotype was analyzed using primers and PCR conditions described previously (14, 16). In brief, tail genomic DNA was extracted and amplified with one forward primer (F1: 5'-CTCCAATGTTCTCAAACCTTAC-3') and two reverse primers (R1: 5'-GATGAGTCATGTAA GTTGACC-3'; R2: 5'-GTATTCTATGGCTTCCAGTGC-3'). The expected sizes of the PCR products are wild-type allele ~250bp, floxed allele ~285bp, and null allele ~450bp. PCR analysis of Cre was performed using primers 5'-GCATTTCTGGGGATTGCTTA-3' (forward) and 5'-GTCATCCTTAGCGCCGTAAA-3' (reverse). The expected size of PCR product for Cre is ~350 bp.

2.4. DXA and Micro Computed Tomography (μ CT) analysis

Bone mineral density (BMD) and bone mineral content (BMC) were measured at 6, 12, and 30 weeks of age by dual-energy X-ray absorptiometry (DXA) (GE Lunar PIXImus system, software version 1.4x) as previously described (17). In brief, mice were anesthetized with a cocktail of ketamine hydrochloride and xylazine (30 mg/ml and 4 mg/ml, respectively, 0.1 ml/25 g body weight, IP) and were placed on the blocks supplied with the instrument according to the manufacturer's recommendations. All mice were analyzed at three distinct regions: (1) the total region, the whole body excluding the head; (2) a femoral region of interest that was defined as the maximal box fitted within a portion of the right femur; (3) a lumbar region of interest that was defined as the biggest box fitted within a portion of the lumbar vertebrae. The stability of the measurement was controlled by performing a quality control procedure before acquiring images. μ CT scan (μ CT-40; Scanco Medical AG, Bassersdorf, Switzerland) was used to evaluate the bone structural parameters as previously described (17).

2.5. Histology and histomorphometry assays

Bone tissues were collected, processed, and analyzed following our procedures described previously (18, 19). Standard bone histomorphometrical nomenclatures, symbols, and units were used as recommended by Parfitt et al (20).

2.6. Primary bone marrow cell culture, isolation and differentiation of BMSCs. Bone marrow cells were flushed from long bones of PPAR γ cKO, heterozygous, and PPAR $\gamma^{\text{fl/fl}}$ mice. Colony-forming unit (CFU) assays, BMSC isolation and differentiation were performed as previously described (21, 22).

2.6. In vitro osteoclast differentiation and bone resorption assay

Bone marrow monocyte and macrophage precursor cells (BMMs) were harvested from long bones and induced for osteoclast differentiation as described (17). In brief, BMMs were seeded in 96-well plates at a density of 1×10^4 cells/well in triplicates and cultured in α -MEM containing FBS (10%), M-CSF (50 ng/ml) and RANKL (50 ng/ml) (R&D Systems, Inc. Minneapolis, MN) for 5-6 days. Differentiated cells were fixed, stained for tartrate-resistant acid phosphatase (TRAP) (#387A-1KT, Sigma Aldrich). For Pit assay, BMMs were seeded in coated quartz slides (Osteologic bonecell culture system, BD Biosciences) and induced for osteoclast differentiation as above. The slides were then stained with 5% silver nitrate solution (von Kossa staining) and the bone resorption was visualized as white spots (pits) on the slides.

2.7. Western blot analysis

Western blot analyses were performed as previously described (21, 23). Protein levels of PPAR γ were detected using a monoclonal antibody raised against a sequence mapping at the C-terminus of PPAR γ (Cat #: sc-7273, Santa Cruz Biotechnology, Inc.).

2.8. Immunofluorescence labeling and Imaging. This experiment was performed as previously described (24) except that a PPAR γ 2 monoclonal antibody was used. Images were acquired using a Nikon TE2000 fluorescence microscope equipped with COOLSNAP Monochrome Camera and processed with Metamorph Imaging System.

2.9. Real-time qRT-PCR analysis

BMSC culture, RNA isolation, and quantitative analysis of levels of mRNA expression were performed as described previously using TaqMan Reverse Transcription Reagents (Applied Biosystems) and a Chromo-4 real-time RT-PCR instrument (MJ Research) (22). The mRNA levels were normalized to β -actin (internal control) and gene expression was presented as fold changes ($\Delta\Delta$ Ct method). The primer sequences used in the PCR reactions are: Runx2: 5'-CCACCACTCACTACCACACG-3' (forward) and 5'-TCAGCGTCAACACCATCATT-3' (reverse); Colla1: 5'-CACCTCAAGAGCCTGAGTC-3' (forward) and 5'-CGGGCTGATGTACCAGTTCT-3' (reverse); ALP: 5'-AACCCAGACACAAGCATTCC-3' (forward) and 5'-CCAGCAAGAAGAAGCCTTTG-3' (reverse); Ocn: 5'-TTCTGCTCACTCTGCTGACC-3' (forward) and 5'-TTTGTAGGCGGTCTTCAAGC-3' (reverse); PPAR γ 2: 5'-TTTTCCGAAGAACCATCCGAT-3' (forward) and 5'-ACAAATGGTGATTTGTCCGTT-3' (reverse); β -actin: 5'-CTGGCACCACACCTTCTACA-3' (forward) and 5'-GGTACGACCAGAGGCATACA-3' (reverse).

2.10. Statistical analysis

Results are expressed as mean $\pm$ SD. All in vitro experiments (qRT-PCR, cell differentiation etc.,) were performed in triplicates except where noted. Data were analyzed using either ANOVA with Bonferroni post hoc test or unpaired *t*-test, using a commercial statistical package (Instat, Graphad Inc.). A *p*-value less than 0.05 was considered significant.

3. RESULTS

3.1. Bone phenotype of PPAR γ cKO mice

Bone mineral density (BMD) and bone mineral content (BMC): DXA scan was performed to detect dynamic changes of BMD and BMC. The measurement was started with mice at 6 weeks of age. This time point was based on studies by Akune et al (12), which showed that PPAR γ haploinsufficient (PPAR $\gamma^{+/-}$) mice exhibit high bone mass at 8 weeks of age. We predicted that the PPAR $\gamma^{fl/fl}$:Col3.6-Cre mice would have a greater bone mass increase at an earlier time point than the PPAR $\gamma^{+/-}$ mice. However, we found no difference in BMD and BMC between cKO and floxed littermate control mice at 6 weeks of age either at the whole body or at the regional levels (data not shown). At 3 months of age, a moderate, yet statistically significant, increase in BMD and BMC (7.9% and 7.7%, respectively, cKO vs. flox) was detected in the lumbar spine of the male cKO mice. The magnitude of increase became more evident at 7.5 months (BMD 12%, BMC 10%, cKO vs. flox) (Table 1).

Table 1. **BMD and BMC of male PPAR γ cKO and fl/fl mice at 3 and 7.5 months of age.**

Age Genotype	3 month			7.5 month		
	cKO (n=12)	fl/fl (n=8)	<i>p</i>	cKO (n=13)	fl/fl (n=13)	<i>p</i>
Whole body						
BMD (g/cm ²)	0.0538±0.0021	0.0526±0.0014	0.2820	0.0593±0.0031	0.0556±0.0021	0.0017
BMC (g)	0.534±0.047	0.526±0.038	0.6749	0.639±0.063	0.574±0.045	0.0053
Spine						
BMD (g/cm ²)	0.0818±0.0031	0.0573±0.0035	0.0067	0.0864±0.0054	0.0594±0.0052	0.0026
BMC (g)	0.042±0.002	0.039±0.003	0.0423	0.044±0.005	0.040±0.006	0.0407
Femur						
BMD (g/cm ²)	0.0772±0.0060	0.0775±0.0023	0.9019	0.0855±0.0077	0.0809±0.0053	0.0880
BMC (g)	0.034±0.003	0.033±0.002	0.7735	0.038±0.005	0.036±0.003	0.1053

μ CT analysis of lumbar spine (L4, 6.5 months of age) showed that the bone volume density (BV/TV) and trabecular number (Tb. N) were increased (56% and 15%, respectively; $p < 0.05$, $n = 5$) and trabecular separation (Tb. Sp) decreased (-12%, $p < 0.05$, $n = 5$) significantly in $\text{PPAR}\gamma$ cKO mice compared to that in $\text{PPAR}\gamma^{\text{fl/fl}}$ control mice (Fig. 1A-C). The trabecular thickness (Tb. Th) showed an increase in cKO mice (+15%, $p = 0.270$, $n = 5$), although this increase is not statistically significant (Fig. 1D). Representative re-constructed 3D images are shown in Fig. 1E.

Figure 1. (A-D) μ CT analysis of lumbar spine (L4) showing BV/TV, Tb. N, Tb. Sp and Tb. Th of 6.5-month-old male $\text{PPAR}\gamma$ cKO (solid bar) and double-floxed control (empty bar) mice. (E) Representative three-dimensional reconstructed images of L4 lumbar spine. Values are given as mean $\pm$ SD.

Histology and histomorphometric analyses of lumbar spine (L2, 6.5 months of age) showed that both the number and the surface of osteoblasts were increased significantly in the PPAR γ cKO mice compared to that in floxed control mice (+98% and +72%, respectively) (Fig. 2).

Figure 2. **Histology and histomorphometry analyses.** (A, B) Representative H&E stained lumbar vertebrae (L4) from 6.5-month-old male mice. (C, D) Enlarged images showing the boxed areas in A and B. (E, F) Quantitative results showing the numbers of osteoblasts and the osteoblast covered bone surfaces.

3.2. PPAR γ gene is ablated successfully in BMSCs

Since an expected high bone mass phenotype was not observed in the PPAR γ cKO mice, we suspected that the Col3.6kb promoter fragment was probably not active in BMSCs but only in a small fraction of committed osteoblasts. To clarify this conjecture, we isolated BMSCs using a negative-immuno-depletion and positive-immuno-selection procedure established in our

laboratory (22) and examined the expression of PPAR γ mRNA. RT-PCR analysis showed that the PPAR γ mRNA is, indeed, absent in the BMSCs of cKO mice while its expression detected in cells of PPAR γ -flox and heterozygous mice (Fig. 3A). The Cre mRNA, which is present in cells of heterozygous and cKO mice, is absent in cells of double floxed mice. These results are consistent with the genotyping results obtained from PCR amplification of the tail genomic DNA (Fig. 3B). To confirm the deletion of PPAR γ at the protein level, we performed Western blot and immunocytochemistry studies using a monoclonal antibody specific to PPAR γ . Results of both experiments revealed, again, that the PPAR γ protein is absent in cells isolated from cKO mice, but expressed in cells of heterozygous and PPAR γ -flox mice (Fig. 3C, D). These results demonstrated that the Col3.6kb promoter fragment is active, robustly drives Cre expression, and ablates PPAR γ allele in BMSCs.

Figure 3. **Characterization of BMSCs isolated from the PPAR γ cKO mice.** (A) RT-PCR analysis showing the expression of PPAR γ and Cre mRNA in BMSCs isolated from PPAR γ cKO, heterozygous (Het), and Floxed (Flox) mice. (B) Genotyping results showing distinct PCR products amplified from tail genomic DNA. KO and floxed PPAR γ alleles are indicated. WT: tail DNA from a wild-type mouse was used as template in PCR reaction to show the wild type allele. (C, D) Western blot (C) and immunofluorescence labeling (D) showing PPAR γ protein expression in BMSCs.

3.3. Adipogenic differentiation is abolished in PPAR γ cKO cell cultures *in vitro*.

To demonstrate, at the functional level, that PPAR γ gene is deleted in BMSCs, we subjected BMSCs to adipogenic and osteogenic differentiation program. Fig. 4 shows that while the PPAR γ -flox BMSCs differentiated into adipocytes normally, as determined by Oil Red O staining of the intracellular lipid droplets, the PPAR γ KO cells completely lost their ability for adipocyte differentiation (Fig. 4A). As expected, *in vitro* osteogenic differentiation of PPAR γ KO BMSC is enhanced significantly as assessed by Alizarin red S (ARS) staining of the mineralized bone nodules (Fig. 4A). This result is replicated in colony forming unit (CFU) assays using freshly prepared whole bone marrow cells, i.e., CFU-adipocyte (CFU-ad) is completely abolished and CFU-osteoblast (CFU-ob) enhanced significantly in marrow cell cultures of PPAR γ cKO mice (Fig. 4B). Quantitative results of these assays are shown in Fig. 4C and D. These results are in line with the RT-PCR and western blot and immunocytochemistry studies. Together, these data demonstrated that the PPAR γ gene is deleted by Col3.6kb promoter-driven Cre in BMSCs or progenitor cells that are capable of differentiating into at least two cell lineages, the adipocytes and osteoblasts.

Figure 4. Adipogenic and osteogenic differentiation assays. (A) BMSCs were subjected to standard adipocyte or osteoblast differentiation program. Cells were either stained with Oil red O for detection of adipocytes 9 days after induction (A, left panel), or stained with Alizarin red S for detection of mineralized bone nodules 21 days after induction (A, right panel). (B) Assays for the number of colony forming units (CFUs). Equal numbers of bone marrow cells (5×10^6 cells/ 25cm^2 flask) from cKO and double-floxed mice were cultured in adipogenic or osteogenic induction media and stained with Oil Red O at day 11 for CFU-ad or with Alizarin red S at day 21 for CFU-ob. Colonies (>50 cells in size) were counted visually. These experiments were repeated two times in triplicates. Only one representative flask is shown for each group. (C, D) Quantitative results of B.

3.4. Expression of osteoblast lineage-associated genes in PPAR γ KO BMSCs. To determine if the observed increase in osteogenesis of PPAR γ KO cells correlates with the expression pattern of osteoblast lineage genes, we performed real-time qRT-PCR analysis. BMSCs from PPAR γ cKO and floxed mice were treated with osteogenic induction media and analyzed for the expression of mRNAs. Results showed that the mRNA levels of Runx2 (Fig. 5A), type I collagen

(5B), alkaline phosphatase (5C), and osteocalcin (5D) were all increased significantly in PPAR γ KO cells compared to that in PPAR γ -floxed cells. Levels of PPAR γ mRNA in these cells are also shown (5E). These results demonstrated that BMSCs lacking PPAR γ do have an enhanced *in vitro* osteogenic differentiation capability.

Figure 5. Real-time qRT-PCR analysis of mRNA expression during BMSC osteogenic differentiation. BMSCs were cultured in osteogenic induction media and harvested at days 3, 7, and 14 for total RNA isolation. Equal amounts of RNA were reverse transcribed and the mRNA levels of the indicated genes were analyzed. Data were normalized to β -actin and expressed as fold changes. Levels of mRNA in double-floxed BMSCs at day 3 were arbitrarily set as 1. Experiments were repeated two times. PCR reactions were performed in triplicates. Error bars indicate SD. * $p < 0.05$.

3.5. KO of PPAR γ in BMSCs does not affect osteoclast differentiation and function. Lastly, we examined osteoclastogenesis both *in vivo*, using decalcified tibia samples, and *in vitro*, using bone marrow monocyte/macrophage precursor cells (BMMs). TRAP stain experiment showed that neither the numbers of osteoclasts (N.Oc/B.Pm) nor the osteoclast surfaces (N.Oc/BS) were different between PPAR γ cKO and floxed control mice (Fig. 6A). *In vitro* osteoclast induction experiment showed that BMMs from PPAR γ cKO and floxed mice differentiated into multinucleated TRAP-positive osteoclasts equally well (Fig. 6B), demonstrating that the osteoclast differentiation is not affected. To determine whether these osteoclasts are functionally different, we performed a bone resorption assay. BMMs were seeded in coated quartz slides and induced for osteoclast differentiation. The slides were then fixed and stained with 5% silver nitrate solution (von Kossa staining) to assess the bone resorption activity. Again, the results showed that the osteoclasts differentiated from BMMs of cKO and floxed control mice had similar bone resorption activity (Fig. 6C). These results suggested that deletion of PPAR γ in osteoblastic lineage cells or BMSCs does not have an effect on osteoclast differentiation or function. This conclusion is consistent to and supported, in some degree, by studies of Akune et al, which showed that the osteoclastogenesis is not affected in PPAR γ -haploinsufficient mice (12).

Figure 6. **Effect of PPAR γ on osteoclast differentiation and function.** (A) TRAP stain of decalcified tibiae samples showing the TRAP-positive osteoclasts in PPAR γ cKO and double-floxed control mice at 7 months of age. (B) BMMs isolated from PPAR γ cKO and floxed mice were induced for osteoclast differentiation in the presence of M-CSF (50ng/ml) and RANKL (50ng/ml) for 6 days and stained for TRAP activity. (C) Pit assay showing bone resorption activity. BMMs were induced as above in quartz chamber slides and stained with 5% silver nitrate. White spots indicate the area where bone resorption occurred. Insets: representative high power images of osteoclasts and pits shown in A and B. Original magnification of the images including insets is indicated. These experiments were performed at least 3 times and with different concentrations of RANKL, same results were obtained.

4. DISCUSSION

In this study we showed that deletion of PPAR γ gene in 3.6kb type I collagen promoter-expressing cells (PPAR^{fl/fl}:Col3.6-Cre) resulted in a moderately enhanced bone mineral density preferentially in the vertebrae of the elder mice (Table 1). Multiple experiments, from DNA to protein expression showed clearly that the PPAR γ gene is deleted in cells supporting the activities of Col3.6kb promoter fragment (Fig. 3). Giving the high expectations for the impact of PPAR γ knockout may have on bone, and the fact that the adipogenesis of both primary bone marrow cells and the BMSCs isolated from the PPAR γ cKO mice is abolished completely (Fig. 4), it is a little disappointing that the bone density increase was not dramatic in the PPAR^{fl/fl}:Col3.6-Cre mice. Interestingly, however, a similar bone phenotype was also observed in another conditional PPAR γ knockout mouse model, the PPAR γ ^{fl/fl}:Sox2-Cre mouse (25). This mouse, which carries total deletion of PPAR γ , showed an increased BMD only in vertebrae but not in long bones (25). More recently, Sun et al (26) reported that osteoblast-specific PPAR γ KO (at ~3 weeks of age) using an inducible Osx promoter-driven Cre (PPAR γ ^{fl/fl}:Sp7-tTA,tetO-EGFP/Cre) increased the Tb. N and decreased Tb. Sp in 6-month-old mice, but had no effect on BMD of the femur (long bone) and the authors concluded that suppression of PPAR γ increases osteogenesis through activation of mTOR signaling. These results clearly indicated that other mechanisms by which PPAR γ regulates bone exist. A plausible one, as proposed by Gimble and colleagues (27) and Ferrari and colleagues (25), in addition to mTOR pathway, is that PPAR γ regulates bone turnover/remodeling by both a direct effect on cell differentiation and an indirect effect on cell fate determination through adipocyte and adipokine secretion (25). Supporting this notion, studies of lipodystrophic PPAR γ ^{hyp/hyp} mouse, which lacks PPAR γ only in adipose tissues, also showed increased osteogenic activity and enhanced bone formation (13). The adipose tissue

is an endocrine organ and secretes large amounts of factors including adipokines and inflammatory cytokines that are known to regulate bone formation and metabolism (28-40). Currently, it is not clear, and controversial, whether PPAR γ plays a role in osteoclast differentiation or function. Our studies showed that the osteoclast differentiation was not affected in PPAR^{fl/fl}:Col3.6-Cre cKO mice (Fig. 6). Studies by Akune et al showed that osteoclast differentiation and function are not affected in PPAR γ -haploinsufficient mice (12). Interestingly, Wan et al showed that mice lacking PPAR γ in osteoclast lineage cells (Tie2-Cre:PPAR^{fl/fl}) develop severe osteopetrosis, suggesting that PPAR γ is a strong pro-osteoclast factor (41). In contrast, several lines of *in vitro* evidence showed that activation of PPAR γ by its ligands inhibits osteoclast differentiation (42-49). However, it is not clear whether this ligand-mediated inhibition is PPAR γ dependent or independent.

Bone is a dynamic enterprise; it is constantly destroyed by osteoclasts and rebuilt by osteoblasts in a process called bone remodeling. Bone remodeling occurs constantly throughout the developed skeleton and is a key process maintaining bone homeostasis. While the activity of resorption and formation may keep in balance when bone mass is at its peak level or shortly after, during the phase of plateau, this balance is broken in old human or animals with more bones destroyed than new bones formed, thus generates a deficit in each remodeling cycle due to the lack of osteoblasts for repair (33, 50). Bone loss with aging is accompanied by an increase in marrow fat content (3-5), and this coincide with the increase of PPAR γ (51-53). Our data that knockout PPAR γ in BMSC/progenitor cells increased the numbers of osteoblasts suggests that PPAR γ may play an important role in aging-induced bone loss, i.e., as the bone marrow microenvironment deteriorates with advancing age, the levels of PPAR γ expression and/or

activity increase. These changes lead to an increased marrow adipogenesis and the production of inflammatory cytokines and adipokines, which impact negatively on the production of osteoblasts and consequently, the bone repair during bone remodeling.

ACKNOWLEDGEMENTS

We thank Dr. Frank J. Gonzalez, National Cancer Institute, NIH, for providing floxed PPAR γ mouse. We thank Byung Lee, Michael (Konglun) Yu and Hongfan Yu for their involvement of this study. This work was supported by grants from The National Institute on Aging (AG046248 and AG036675) and The National Institute of Diabetes and Digestive and Kidney Diseases (DK76045), National Institutes of Health (NIH).

REFERENCES

1. Barak, Y., Nelson, M.C., Ong, E.S., Jones, Y.Z., Ruiz-Lozano, P., Chien, K.R., Koder, A., and Evans, R.M. 1999. PPAR gamma is required for placental, cardiac, and adipose tissue development. *Mol.Cell* 4:585-595.
2. Rosen, E.D., Sarraf, P., Troy, A.E., Bradwin, G., Moore, K., Milstone, D.S., Spiegelman, B.M., and Mortensen, R.M. 1999. PPAR gamma is required for the differentiation of adipose tissue in vivo and in vitro. *Mol.Cell* 4:611-617.
3. Justesen, J., Stenderup, K., Ebbesen, E.N., Mosekilde, L., Steiniche, T., and Kassem, M. 2001. Adipocyte tissue volume in bone marrow is increased with aging and in patients with osteoporosis. *Biogerontology*. 2:165-171.
4. Verma, S., Rajaratnam, J.H., Denton, J., Hoyland, J.A., and Byers, R.J. 2002. Adipocytic proportion of bone marrow is inversely related to bone formation in osteoporosis. *J.Clin.Pathol.* 55:693-698.
5. Meunier, P., Aaron, J., Edouard, C., and Vignon, G. 1971. Osteoporosis and the replacement of cell populations of the marrow by adipose tissue. A quantitative study of 84 iliac bone biopsies. *Clin.Orthop.* 80:147-154.
6. Ahdjoudj, S., Lasmoles, F., Oyajobi, B.O., Lomri, A., Delannoy, P., and Marie, P.J. 2001. Reciprocal control of osteoblast/chondroblast and osteoblast/adipocyte differentiation of multipotential clonal human marrow stromal F/STRO-1(+) cells. *J Cell Biochem.* 81:23-38.

7. Beresford, J.N., Bennett, J.H., Devlin, C., Leboy, P.S., and Owen, M.E. 1992. Evidence for an inverse relationship between the differentiation of adipocytic and osteogenic cells in rat marrow stromal cell cultures. *J.Cell Sci.* 102 (Pt 2):341-351.
8. Jaiswal, R.K., Jaiswal, N., Bruder, S.P., Mbalaviele, G., Marshak, D.R., and Pittenger, M.F. 2000. Adult human mesenchymal stem cell differentiation to the osteogenic or adipogenic lineage is regulated by mitogen-activated protein kinase. *Journal of Biological Chemistry* 275:9645-9652.
9. Minguell, J.J., Erices, A., and Conget, P. 2001. Mesenchymal stem cells. *Exp.Biol.Med.(Maywood.)* 226:507-520.
10. Pittenger, M.F., Mackay, A.M., Beck, S.C., Jaiswal, R.K., Douglas, R., Mosca, J.D., Moorman, M.A., Simonetti, D.W., Craig, S., and Marshak, D.R. 1999. Multilineage Potential of Adult Human Mesenchymal Stem Cells. *Science* 284:143-147.
11. Prockop, D.J. 1997. Marrow stromal cells as stem cells for nonhematopoietic tissues. *Science* 276:71-74.
12. Akune, T., Ohba, S., Kamekura, S., Yamaguchi, M., Chung, U.I., Kubota, N., Terauchi, Y., Harada, Y., Azuma, Y., Nakamura, K., et al. 2004. PPARgamma insufficiency enhances osteogenesis through osteoblast formation from bone marrow progenitors. *J.Clin.Invest* 113:846-855.
13. Cock, T.A., Houten, S.M., and Auwerx, J. 2004. Peroxisome proliferator-activated receptor-gamma: too much of a good thing causes harm. *EMBO Rep.* 5:142-147.
14. Liu, F., Woitge, H.W., Braut, A., Kronenberg, M.S., Lichtler, A.C., Mina, M., and Kream, B.E. 2004. Expression and activity of osteoblast-targeted Cre recombinase transgenes in murine skeletal tissues. *Int J Dev.Biol* 48:645-653.
15. Akiyama, T.E., Sakai, S., Lambert, G., Nicol, C.J., Matsusue, K., Pimprale, S., Lee, Y.H., Ricote, M., Glass, C.K., Brewer, H.B., Jr., et al. 2002. Conditional disruption of the peroxisome proliferator-activated receptor gamma gene in mice results in lowered expression of ABCA1, ABCG1, and apoE in macrophages and reduced cholesterol efflux. *Mol Cell Biol* 22:2607-2619.
16. Zhang, H., Zhang, A., Kohan, D.E., Nelson, R.D., Gonzalez, F.J., and Yang, T. 2005. Collecting duct-specific deletion of peroxisome proliferator-activated receptor {gamma} blocks thiazolidinedione-induced fluid retention. *Proceedings of the National Academy of Sciences* 102:9406-9411.
17. Pan, G., Cao, J., Yang, N., Ding, K., Fan, C., Xiong, W.-C., Hamrick, M., Isales, C.M., and Shi, X.-M. 2014. Role of Glucocorticoid-Induced Leucine Zipper (GILZ) in Bone Acquisition. *Journal of Biological Chemistry.*
18. Hamrick, M.W., Della-Fera, M.A., Choi, Y.H., Pennington, C., Hartzell, D., and Baile, C.A. 2005. Leptin Treatment Induces Loss of Bone Marrow Adipocytes and Increases Bone Formation in Leptin-Deficient ob/ob Mice. *Journal of Bone and Mineral Research* 20:994-1001.
19. Hamrick, M.W., Shi, X., Zhang, W., Pennington, C., Thakore, H., Haque, M., Kang, B., Isales, C.M., Fulzele, S., and Wenger, K.H. 2007. Loss of myostatin (GDF8) function increases osteogenic differentiation of bone marrow-derived mesenchymal stem cells but the osteogenic effect is ablated with unloading. *Bone* 40:1544-1553.
20. Parfitt, A.M., Drezner, M.K., Glorieux, F.H., Kanis, J.A., Malluche, H., Meunier, P.J., Ott, S.M., and Recker, R.R. 1987. Bone histomorphometry: standardization of

- nomenclature, symbols, and units. Report of the ASBMR Histomorphometry Nomenclature Committee. *J.Bone Miner.Res.* 2:595-610.
21. Zhang, W., Yang, N., and Shi, X.M. 2008. Regulation of Mesenchymal Stem Cell Osteogenic Differentiation by Glucocorticoid-induced Leucine Zipper (GILZ). *Journal of Biological Chemistry* 283:4723-4729.
 22. Zhang, W., Ou, G., Hamrick, M., Hill, W., Borke, J., Wenger, K., Chutkan, N., Yu, J., Mi, Q.S., Isales, C.M., et al. 2008. Age-Related Changes in the Osteogenic Differentiation Potential of Mouse Bone Marrow Stromal Cells. *Journal of Bone and Mineral Research* 23:1118-1128.
 23. Shi, X., Shi, W., Li, Q., Song, B., Wan, M., Bai, S., and Cao, X. 2003. A glucocorticoid-induced leucine-zipper protein, GILZ, inhibits adipogenesis of mesenchymal cells. *EMBO Rep.* 4:374-380.
 24. Yang, N., Zhang, W., and Shi, X.M. 2007. Glucocorticoid-induced leucine zipper (GILZ) mediates glucocorticoid action and inhibits inflammatory cytokine-induced COX-2 expression. *J Cell Biochem* 103:1760-1771.
 25. He, F., Dominique, P., Anne, W., Serge, F., and Beatrice, D. 2011. Bone Marrow Adiposity Affects Osteoclastogenesis by Modulating the Bone Marrow Niche *ASBMR Annual Meeting* 2011 Annual Meeting:S63 - S63
 26. Sun, H., Kim, J.K., Mortensen, R., Mutyaba, L.P., Hankenson, K.D., and Krebsbach, P.H. 2013. Osteoblast-targeted suppression of PPARgamma increases osteogenesis through activation of mTOR signaling. *Stem Cells* 31:2183-2192.
 27. Nuttall, M., Shah, F., Singh, V., Thomas-Porch, C., Frazier, T., and Gimble, J. 2014. Adipocytes and the Regulation of Bone Remodeling: A Balancing Act. *Calcified Tissue International* 94:78-87.
 28. Trujillo, M.E., and Scherer, P.E. 2006. Adipose Tissue-Derived Factors: Impact on Health and Disease. *Endocrine Reviews* 27:762-778.
 29. Abrahamsen, B.O., Bonnevie-Nielsen, V.A.G.N., Ebbesen, E.N., Gram, J.E.P.P., and Beck-Nielsen, H.E.N.N. 2000. Cytokines and Bone Loss in a 5-Year Longitudinal Study-Hormone Replacement Therapy Suppresses Serum Soluble Interleukin-6 Receptor and Increases Interleukin-1-Receptor Antagonist: The Danish Osteoporosis Prevention Study. *Journal of Bone and Mineral Research* 15:1545-1554.
 30. Khosla, S., Peterson, J.M., Egan, K., Jones, J.D., and Riggs, B.L. 1994. Circulating cytokine levels in osteoporotic and normal women. *Journal of Clinical Endocrinology Metabolism* 79:707-711.
 31. Jilka, R.L., Hangoc, G., Girasole, G., Passeri, G., Williams, D.C., Abrams, J.S., Boyce, B., Broxmeyer, H., and Manolagas, S.C. 1992. Increased osteoclast development after estrogen loss: mediation by interleukin-6. *Science* 257:88-91.
 32. Nanes, M.S. 2003. Tumor necrosis factor-[alpha]: molecular and cellular mechanisms in skeletal pathology. *Gene* 321:1-15.
 33. Manolagas, S.C., and Jilka, R.L. 1995. Bone Marrow, Cytokines, and Bone Remodeling - Emerging Insights into the Pathophysiology of Osteoporosis. *The New England Journal of Medicine* 332:305-311.
 34. Ding, C., Parameswaran, V., Udayan, R., Burgess, J., and Jones, G. 2008. Circulating Levels of Inflammatory Markers Predict Change in Bone Mineral Density and Resorption in Older Adults: A Longitudinal Study. *Journal of Clinical Endocrinology Metabolism* 93:1952-1958.

35. Maurin, A.C., Chavassieux, P.M., Frappart, L., Delmas, P.D., Serre, C.M., and Meunier, P.J. 2000. Influence of mature adipocytes on osteoblast proliferation in human primary cocultures. *Bone* 26:485-489.
36. Fried, S.K., Bunkin, D.A., and Greenberg, A.S. 1998. Omental and subcutaneous adipose tissues of obese subjects release interleukin-6: depot difference and regulation by glucocorticoid. *J.Clin.Endocrinol.Metab* 83:847-850.
37. Peng, X.D., Xie, H., Zhao, Q., Wu, X.P., Sun, Z.Q., and Liao, E.Y. 2008. Relationships between serum adiponectin, leptin, resistin, visfatin levels and bone mineral density, and bone biochemical markers in Chinese men. *Clinica Chimica Acta* 387:31-35.
38. Sayers, A., Timpson, N.J., Sattar, N., Deanfield, J., Hingorani, A.D., Davey-Smith, G., and Tobias, J.H. 2010. Adiponectin and its association with bone mass accrual in childhood. *Journal of Bone and Mineral Research* 25:2212-2220.
39. Filková, M., Haluzík, M., Gay, S., and Senolt, L. 2009. The role of resistin as a regulator of inflammation: Implications for various human pathologies. *Clinical Immunology* 133:157-170.
40. Richards, J.B., Valdes, A.M., Burling, K., Perks, U.C., and Spector, T.D. 2007. Serum Adiponectin and Bone Mineral Density in Women. *Journal of Clinical Endocrinology Metabolism* 92:1517-1523.
41. Wan, Y., Chong, L.W., and Evans, R.M. 2007. PPAR-gamma regulates osteoclastogenesis in mice. *Nat Med* 13:1496-1503.
42. Mbalaviele, G., Abu-Amer, Y., Meng, A., Jaiswal, R., Beck, S., Pittenger, M.F., Thiede, M.A., and Marshak, D.R. 2000. Activation of Peroxisome Proliferator-activated Receptor-gamma Pathway Inhibits Osteoclast Differentiation. *Journal of Biological Chemistry* 275:14388-14393.
43. Bendixen, A.C., Shevde, N.K., Dienger, K.M., Willson, T.M., Funk, C.D., and Pike, J.W. 2001. IL-4 inhibits osteoclast formation through a direct action on osteoclast precursors via peroxisome proliferator-activated receptor gamma 1. *Proc Natl Acad Sci U.S.A* 98:2443-2448.
44. Hounoki, H., Sugiyama, E., Mohamed, S.G., Shinoda, K., Taki, H., Abdel-Aziz, H.O., Maruyama, M., Kobayashi, M., and Miyahara, T. 2008. Activation of peroxisome proliferator-activated receptor gamma inhibits TNF-alpha-mediated osteoclast differentiation in human peripheral monocytes in part via suppression of monocyte chemoattractant protein-1 expression. *Bone* 42:765-774.
45. Abbas, S., Zhang, Y.H., Clohisy, J.C., and Abu-Amer, Y. 2003. Tumor necrosis factor-alpha inhibits pre-osteoblast differentiation through its type-1 receptor. *Cytokine* 22:33-41.
46. Kaneki, H., Guo, R., Chen, D., Yao, Z., Schwarz, E.M., Zhang, Y.E., Boyce, B.F., and Xing, L. 2006. Tumor Necrosis Factor Promotes Runx2 Degradation through Up-regulation of Smurf1 and Smurf2 in Osteoblasts. *Journal of Biological Chemistry* 281:4326-4333.
47. Yamazaki, M., Fukushima, H., Shin, M., Katagiri, T., Doi, T., Takahashi, T., and Jimi, E. 2009. Tumor necrosis factor alpha represses bone morphogenetic protein (BMP) signaling by interfering with the DNA binding of Smads through the activation of NF-kappaB. *Journal of Biological Chemistry* 284:35987-35995.

48. Bertolini, D.R., Nedwin, G.E., Bringman, T.S., Smith, D.D., and Mundy, G.R. 1986. Stimulation of bone resorption and inhibition of bone formation in vitro by human tumour necrosis factors. *Nature* 319:516-518.
49. Kobayashi, K., Takahashi, N., Jimi, E., Udagawa, N., Takami, M., Kotake, S., Nakagawa, N., Kinosaki, M., Yamaguchi, K., Shima, N., et al. 2000. Tumor Necrosis Factor {alpha} Stimulates Osteoclast Differentiation by a Mechanism Independent of the ODF/RANKL-RANK Interaction. *The Journal of Experimental Medicine* 191:275-286.
50. Manolagas, S.C. 2000. Birth and Death of Bone Cells: Basic Regulatory Mechanisms and Implications for the Pathogenesis and Treatment of Osteoporosis. *Endocrine Reviews* 21:115-137.
51. Jiang, Y., Mishima, H., Sakai, S., Liu, Y.k., Ohyabu, Y., and Uemura, T. 2008. Gene expression analysis of major lineage-defining factors in human bone marrow cells: Effect of aging, gender, and age-related disorders. *Journal of Orthopaedic Research* 26:910-917.
52. Almeida, M., Ambrogini, E., Han, L., Manolagas, S.C., and Jilka, R.L. 2009. Increased Lipid Oxidation Causes Oxidative Stress, Increased Peroxisome Proliferator-activated Receptor-__ Expression, and Diminished Pro-osteogenic Wnt Signaling in the Skeleton. *Journal of Biological Chemistry* 284:27438-27448.
53. Imai, T., Takakuwa, R., Marchand, S., Dentz, E., Bornert, J.M., Messaddeq, N., Wendling, O., Mark, M., Desvergne, B., Wahli, W., et al. 2004. Peroxisome proliferator-activated receptor gamma is required in mature white and brown adipocytes for their survival in the mouse. *Proc.Natl.Acad.Sci.U.S.A* 101:4543-4547.